



Applicable Analysis An International Journal

ISSN: 0003-6811 (Print) 1563-504X (Online) Journal homepage: https://www.tandfonline.com/loi/gapa20

A hyperbolic-type azimuthal velocity model for equatorial currents

Kateryna Marynets

To cite this article: Kateryna Marynets (2020): A hyperbolic-type azimuthal velocity model for equatorial currents, Applicable Analysis, DOI: 10.1080/00036811.2020.1774054

To link to this article: https://doi.org/10.1080/00036811.2020.1774054

Ω	
0	

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 04 Jun 2020.

	Submit	our	article to	o this	journal	
		/				

Article views: 182



View related articles



View Crossmark data 🗹



∂ OPEN ACCESS

Check for updates

A hyperbolic-type azimuthal velocity model for equatorial currents

Kateryna Marynets 🕩

Delft Institute of Applied Mathematics, Faculty of Electrical Engineering, Mathematics and Computer Science Delft University of Technology, Delft, Netherlands

ABSTRACT

We discuss a mathematical model for the equatorial current across the Pacific Ocean, obtained as a leading-order solution to the Navier-Stokes governing equations for geophysical flows in a rotating frame.

ARTICLE HISTORY Received 4 March 2020 Accepted 11 May 2020

COMMUNICATED BY Y. S. Xu

KEYWORDS

Geophysical flow; boundary-value problem; vorticity; viscosity

2010 MATHEMATICS SUBJECT CLASSIFICATIONS

Primary 34B15; 35J15; Secondary37N10

1. Introduction

Equatorial currents are among the most significant currents in the oceans, being an essential part of the flow. Classical theories ignore their effect, but in recent years their significance was recognized and various theoretical investigations were devoted to this important aspect of geophysical flows.

The main features of the equatorial current are that the current is mainly azimuthal and presents a strong depth-variation in the upper 500 m of the ocean (of about 4 km mean depth), while at depths larger than 500 m there is practically no motion. The most significant features of the depth-variation consist of a westward near-surface current (induced by the trade winds that blow towards the west) and a stronger eastward flow – the Equatorial Undercurrent (EUC) that is a major feature of equatorial circulation. It occurs in Pacific (Cromwell Current), Atlantic (Lomonosov Current) and Indian oceans, but only in the first two of them has a permanent character. In the equatorial Indian Ocean region it is temporary and appears in its western part in spring towards the end of the northeast monsoon season (see [1]).

A lot of attention is paid to study and mathematical modeling of the ocean flows in the equatorial regions and their underlying currents, using the *f* - and β -plain approximations and dealing with wave solutions for the governing equations in Lagrangian coordinates (see discussions in [2–8], [14–35]). In addition, a recent research (see [9]) proposed an approach to explain how such a current with flow-reversal might be induced by the action of the wind. Moreover the velocity profile derived in [10], also allowed to include the westward Equatorial Intermediate Current (EIC), found under the EUC, and which was not taken into account in [9]. The present paper is a natural continuation of [10], where

CONTACT Kateryna Marynets 🖾 K.Marynets@tudelft.nl

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (http://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

2 🐱 K. MARYNETS

we prove that not only polynomial profiles of the velocity components, but also the hyperbolic ones allow us to model the aforementioned flow behavior in the equatorial regions.

2. Preliminaries

Consider the Navier–Stokes equation and the equation of mass conservation in rotating spherical coordinates (see discussion in [9])

$$\begin{pmatrix} \frac{\partial}{\partial t'} + \frac{u'}{r'\cos\theta} \frac{\partial}{\partial\varphi} + \frac{v'}{r'} \frac{\partial}{\partial\theta} + w' \frac{\partial}{\partial r'} \end{pmatrix} (u', v', w') + \frac{1}{r'} \left(-u'v' \tan\theta + u'w', u'^2 \tan\theta + v'w', -u'^2 - v'^2 \right) + 2\Omega'(-v'\sin\theta + w'\cos\theta, u'\sin\theta, -u'\cos\theta) + r'\Omega'^2(0, \sin\theta\cos\theta, -\cos^2\theta) = -\frac{1}{\rho'} \left(\frac{1}{r'\cos\theta} \frac{\partial p'}{\partial\varphi}, \frac{1}{r'} \frac{\partial p'}{\partial\theta}, \frac{\partial p'}{\partial r'} \right) + (0, 0, -g') + v'_1 \left(\frac{\partial^2}{\partial r'^2} + \frac{2}{r'} \frac{\partial}{\partial r'} \right) (u', v', w') + \frac{v'_2}{r'^2} \left(\frac{1}{\cos^2\theta} \frac{\partial^2}{\partial\varphi^2} - \tan\theta \frac{\partial}{\partial\theta} + \frac{\partial^2}{\partial\theta^2} \right) (u', v', w'),$$
(1)

and

$$\frac{1}{r'\cos\theta}\frac{\partial u'}{\partial\varphi} + \frac{1}{r'\cos\theta}\frac{\partial}{\partial\theta}(v'\cos\theta) + \frac{1}{r'^2}\frac{\partial}{\partial r'}(r'^2w') = 0,$$
(2)

respectively, where

- (φ, θ, r') are the near the Equator spherical coordinates with r' is the radius;
- $\theta \in (-\pi/2, \pi/2)$ is the angle of latitude and $\varphi \in [0, 2\pi)$ is the angle of longitude;
- (e_φ, e_θ, e_r) is the system of unit vectors, with e_φ pointing from West to East, e_θ from South to North and e_r upwards;
- (u', v', w') are the ocean flow velocity components;
- $p'(\varphi, \theta, r', t')$ is the pressure in the fluid;
- ρ' the constant density;
- $\Omega' \approx 7.29 \times 10^{-5}$ rad s⁻¹ is the constant rate of rotation of the Earth;
- $g' = \text{constant} \approx 9.81 \text{ m s}^{-2}$ is the gravitational acceleration.

Here the notations with primes mean the physical (dimensional) variables. These will be removed after the non-dimensionalization of Equations (1) and (2).

In the Navier–Stokes equation (1) the coefficients of the viscous terms are taken to be constant, where ν'_1 represents the vertical and ν'_2 the horizontal kinematic eddy viscosities.

At the ocean's surface the dynamic and kinematic boundary constraints and the wind-stress are expressed by

$$p' = P'_s(\varphi, \theta, t') \quad \text{on} \quad r' = R' + h'(\varphi, \theta, t'), \tag{3}$$

$$w' = \frac{\partial h'}{\partial t'} + \frac{u'}{r'\cos\theta} \frac{\partial h'}{\partial \varphi} + \frac{v'}{r'} \frac{\partial h'}{\partial \theta} \quad \text{on} \quad r' = R' + h'(\varphi, \theta, t'), \tag{4}$$

$$\begin{cases} \tau_1'(\varphi,\theta,t') = \rho' \nu_1' \frac{\partial u'}{\partial r'}, \\ \tau_2'(\varphi,\theta,t') = \rho' \nu_1' \frac{\partial \nu'}{\partial r'}, \end{cases} \quad \text{on} \quad r' = R' + h'(\varphi,\theta,t'), \tag{5}$$

respectively, where

- $R' \approx 6378$ km is the radius of the spherical Earth;
- P'_{s} is the surface pressure;
- (τ'_1, τ'_2) is the surface wind stress, related to the vertical eddy viscosity by $\nu'_1 = \sigma' |(\tau'_1, \tau'_2)|$ on the surface;
- σ' is a (dimensional) constant;
- (τ'₁, τ'₂) = c_Dρ'_{air}U'_{wind}|U'_{wind}|, ρ'_{air} ≈ 1.2 kg/m³ is the density of air;
 c_D ≈ 0.0013 is a (dimensionless) drag coefficient.

The boundary condition at the bottom of the ocean $r' = R' + d'(\varphi, \theta)$ for the viscous flow is given by relation

$$u' = v' = w' = 0$$
 on $r' = R' + d'(\varphi, \theta)$. (6)

By setting r' = R' + z' and

$$p' = \rho' \left(-g'r' + \frac{1}{2}r'^2 \Omega'^2 \cos^2 \theta \right) + P'(\varphi, \theta, r', t'),$$
(7)

the problem can be non-dimensionalized via following substitutions (see the discussion in [9]):

$$z' = D'z, \quad (u', v', w') = U'(u, v, kw), \quad P' = \rho' U'^2 P,$$
(8)

where

- $D' \approx 200$ m is the average depth of the near-surface layer;
- U' ≈ 0.1 m s⁻¹ is the typical speed of mid-latitude ocean currents at the surface;
 k < 10⁻⁴ is the scaling factor for the vertical velocity.

In terms of the the shallow-water parameter

$$\varepsilon = D'/R',$$

with a typical value of the order 10^{-5} , we rewrite the governing equation (1) and the equation of mass conservation (2) for steady flow in the non-dimensional form

$$\left(\frac{u}{(1+\varepsilon z)\cos\theta}\frac{\partial}{\partial\varphi} + \frac{v}{1+\varepsilon z}\frac{\partial}{\partial\theta} + \frac{k}{\varepsilon}w\frac{\partial}{\partial z}\right)(u, v, kw) + \frac{1}{1+\varepsilon z}\left(-uv\tan\theta + kuw, u^{2}\tan\theta + kvw, -u^{2} - v^{2}\right) + 2\omega(-v\sin\theta + kw\cos\theta, u\sin\theta, -u\cos\theta) = -\left(\frac{1}{(1+\varepsilon z)\cos\theta}\frac{\partial P}{\partial\varphi}, \frac{1}{1+\varepsilon z}\frac{\partial P}{\partial\theta}, \frac{1}{\varepsilon}\frac{\partial P}{\partial z}\right) + \frac{1}{R_{\varepsilon 1}}\left(\frac{1}{\varepsilon^{2}}\frac{\partial^{2}}{\partial z^{2}} + \frac{2}{(1+\varepsilon z)}\frac{1}{\varepsilon}\frac{\partial}{\partial z}\right)(u, v, kw) + \frac{1}{R_{\varepsilon 2}(1+\varepsilon z)^{2}}\left(\frac{1}{\cos^{2}\theta}\frac{\partial^{2}}{\partial\varphi^{2}} - \tan\theta\frac{\partial}{\partial\theta} + \frac{\partial^{2}}{\partial\theta^{2}}\right)(u, v, kw),$$
(9)

4 👄 K. MARYNETS

and

$$\frac{1}{(1+\varepsilon z)\cos\theta} \left\{ \frac{\partial u}{\partial\varphi} + \frac{\partial}{\partial\theta} (v\cos\theta) \right\} + \frac{k/\varepsilon}{(1+\varepsilon z)^2} \frac{\partial}{\partial z} \left\{ (1+\varepsilon z)^2 w \right\} = 0, \tag{10}$$

respectively, where $\omega = \Omega' R' / U' = O(1)$ and $R_{ei} = U' R' / \nu'_j$ (j = 1, 2) are the inverse Rossby number and the pair of Reynolds numbers, respectively.

In addition, for (h', d') = D'(h, d), the non-dimensional boundary restrictions are

$$\begin{cases} P = \overline{P}_{s}(\varphi, \theta), \\ \frac{\partial u}{\partial z} = \tau_{1}(\varphi, \theta), & \text{on } z = h(\varphi, \theta), \\ \frac{\partial v}{\partial z} = \tau_{2}(\varphi, \theta), \end{cases}$$
(11)

$$\frac{k}{\varepsilon}w = \frac{u}{(1+\varepsilon h)\cos\theta}\frac{\partial h}{\partial\varphi} + \frac{v}{1+\varepsilon h}\frac{\partial h}{\partial\theta} \quad \text{on} \quad z = h(\varphi,\theta),$$
(12)

(u, v) decays rapidly below $z = h(\varphi, \theta)$. (13)

In [9] Constantin, Johnson show that for the equatorial Pacific flows it is adequate to consider

$$\omega = \mathcal{O}(1), \quad \frac{1}{R_{e2}} = \varepsilon^2 \mu \quad \text{with } \mu = \frac{\nu'_2}{\nu'_1}.$$

By multiplication of the third component in the governing equation (9) by ε , we obtain the Navier-Stokes system with Coriolis effects in the non-dimensional form:

$$\left(\frac{u}{(1+\varepsilon z)\cos\theta}\frac{\partial}{\partial\varphi} + \frac{v}{1+\varepsilon z}\frac{\partial}{\partial\theta} + \frac{k}{\varepsilon}w\frac{\partial}{\partial z}\right)(u, v, \varepsilon kw) + \frac{1}{1+\varepsilon z}\left(-uv\tan\theta + kuw, u^{2}\tan\theta + kvw, -\varepsilon u^{2} - \varepsilon v^{2}\right) + 2\omega(-v\sin\theta + kw\cos\theta, u\sin\theta, -\varepsilon u\cos\theta) = -\left(\frac{1}{(1+\varepsilon z)\cos\theta}\frac{\partial P}{\partial\varphi}, \frac{1}{1+\varepsilon z}\frac{\partial P}{\partial\theta}, \frac{\partial P}{\partial z}\right) + \left(\frac{\partial^{2}}{\partial z^{2}} + \frac{2\varepsilon}{(1+\varepsilon z)}\frac{\partial}{\partial z}\right)(u, v, \varepsilon kw) + \frac{\varepsilon^{2}\mu}{(1+\varepsilon z)^{2}}\left(\frac{1}{\cos^{2}\theta}\frac{\partial^{2}}{\partial\varphi^{2}} - \tan\theta\frac{\partial}{\partial\theta} + \frac{\partial^{2}}{\partial\theta^{2}}\right)(u, v, \varepsilon kw).$$
(14)

In the non-dimensional system of Equations (14) and (10), the parameters ($\omega, \mu, k, \varepsilon$) are held fixed.

In fact the shallow-water model of the equatorial current may be obtained by passing to the limit for $\varepsilon \to 0$ and $k/\varepsilon \to 0$ (see the discussion in [9]). This regime ignores the vertical velocity component and the flow dynamics, governed by the horizontal flow components *u* and *v*, subject to the nonlinear system

$$\left(\frac{u}{\cos\theta}\frac{\partial}{\partial\varphi} + v\frac{\partial}{\partial\theta}\right)u - uv\tan\theta - 2\omega v\sin\theta = -\frac{1}{\cos\theta}\frac{\partial P}{\partial\varphi} + \frac{\partial^2 u}{\partial z^2},\tag{15}$$

$$\left(\frac{u}{\cos\theta}\frac{\partial}{\partial\varphi} + v\frac{\partial}{\partial\theta}\right)v + u^2\tan\theta + 2\omega u\sin\theta = -\frac{\partial P}{\partial\theta} + \frac{\partial^2 v}{\partial z^2},\tag{16}$$

$$\frac{\partial u}{\partial \varphi} + \frac{\partial}{\partial \theta} (v \cos \theta) = 0, \tag{17}$$

which features Coriolis terms, viscous terms and the horizontal pressure gradients. On the other hand, the boundary conditions (11)–(13) are simplified to

$$\begin{cases} P = \overline{P}_{s}(\varphi, \theta), \\ \frac{\partial u}{\partial z} = \tau_{1}(\varphi, \theta), & \text{on } z = 0, \\ \frac{\partial v}{\partial z} = \tau_{2}(\varphi, \theta), \end{cases}$$
(18)

(u, v) decays rapidly below z = 0. (19)

since setting w = 0 and h = 0 makes (12) irrelevant.

3. Main results

Relation (17) ensures (see the discussion in [11]) the existence of a stream function in spherical coordinates, $\psi(\varphi, \theta, z)$, such that

$$u = -\psi_{\theta}, \quad v = \frac{1}{\cos\theta} \psi_{\varphi}.$$
 (20)

Then we can exclude the pressure terms from Equations (15) and (16) to derive (see [9]) the vorticity equation

$$\left(\psi_{\varphi}\frac{\partial}{\partial\theta} - \psi_{\theta}\frac{\partial}{\partial\varphi}\right) \left(\frac{1}{\cos^{2}\theta}\psi_{\varphi\varphi} - \psi_{\theta}\tan\theta + \psi_{\theta\theta} + 2\omega\sin\theta\right)$$
$$= \cos\theta \left(\frac{1}{\cos^{2}\theta}\psi_{\varphi\varphi} - \psi_{\theta}\tan\theta + \psi_{\theta\theta}\right)_{zz}.$$
(21)

We restrict our attention to the equatorial zonal band located in the Pacific between between 160° E and 80° W. Its characteristic feature is the symmetry about the Equator and it corresponds to the angle of latitude θ close to 0 and the angle of longitude φ near $\varphi_0 \in (8\pi/9, 14\pi/9)$. For the twice differentiable functions $\alpha(z)$ and $\beta(z)$, we look for longitude-independent meridional velocity components under a linear dependence of the azimuthal velocity component on the longitude. As showed in [9], this leads to the stream function

$$\psi(\varphi,\theta,z) = \{\varphi\alpha(z) + \beta(z)\}\ln\left[\frac{\cos\theta}{1-\sin\theta}\right] - \omega\left\{\sin\theta + \ln\left[\frac{\cos\theta}{1-\sin\theta}\right]\right\},\tag{22}$$

satisfying the governing equation (21), valid for $|\varphi - \varphi_0| < \hat{\varphi}$ and $|\theta| < \hat{\theta}$ with some fixed (and small) values $\hat{\varphi} > 0$ and $\hat{\theta} > 0$.

Taking into account relations (20), we obtain the horizontal velocity field and the boundary constraints given by

$$u(\varphi, \theta, z) = -\frac{\varphi \alpha(z) + \beta(z)}{\cos \theta} + \omega \frac{\sin^2 \theta}{\cos \theta},$$

$$v(\varphi, \theta, z) = \frac{\alpha(z)}{\cos \theta} \ln \left[\frac{\cos \theta}{1 - \sin \theta} \right],$$
(23)

6 🕢 K. MARYNETS

and

$$\tau_1(\theta,\varphi) = -\frac{\varphi\alpha'(0) + \beta'(0)}{\cos\theta},$$

$$\tau_2(\theta,\varphi) = \frac{\alpha'(0)}{\cos\theta} \ln\left[\frac{\cos\theta}{1 - \sin\theta}\right]$$
(24)

respectively.

We will interpret the solution (23) as defined in the near-surface ocean region above the thermocline z = -T, where we require the no-stress boundary condition

$$u_z = v_z = 0$$
 on $z = -T$. (25)

The fundamental characteristic of the trade winds in the equatorial Pacific is that, in each hemisphere, they are oriented towards the Equator, blowing westwards with a more stronger westward direction as the Equator ($\theta = 0$) is approached. This property holds for the wind stress specified in (24) if

$$\alpha'(0) < 0 < \varphi \alpha'(0) + \beta'(0) \tag{26}$$

for all relevant values of the longitude φ , that is, for $-\hat{\varphi} < \varphi - \varphi_0 < \hat{\varphi}$. Note that this orientation of the wind, in combination with the fact that the vanishing of the meridional component of the Coriolis force, $\omega \sin \theta$, at the Equator prevents it from inducing a deflection from the wind direction (as is typical in Ekman theory at mid-latitudes), leads to a near-surface current that moving westward. This is ensured for the solution (23) if

$$\varphi\alpha(0) + \beta(0) > 0. \tag{27}$$

Moreover, if

$$\alpha(0) > 0, \tag{28}$$

then the meridional flow described by (23) is poleward near the surface. Note that the main features of the near-surface flow in the equatorial Pacific are a westward motion at the surface and a poleward meridional flow (see the discussion in [12]). One can see that a linear or quadratic dependence on the *z*-variable of the azimuthal velocity profile $u(\varphi, \theta, z)$ of type (23) can not accommodate the constraints (26)–(28) and (25).

In [9] Constantin, Johnson showed that at any fixed longitude $\varphi \in (\varphi_0, \frac{13}{12}\varphi_0)$ the constraints (25)–(28) are accommodated by the cubic azimuthal profile that represents the flow which feathtures the westward drift at the surface and an eastward jet along the termocline.

In addition, a recent result [see discussion in [10]] proved that it is possible to capture not only the presence of the eastward EUC but also the presence of the weaker westward EIC below the EUC, which wasn't taken into account in [9]. In particular a suitable quintic polynomial expressions for the azimuthal velocity profile

$$\begin{aligned} \alpha(z) &= \frac{144}{2125} \frac{(195T^2 - 2264)z^5}{T^5} + \frac{36}{245} \frac{(195T^2 - 2264)z^4}{T^4} - \frac{1}{2}z^2 - Tz + 1, \\ \beta(z) &= \varphi_0 \left\{ \frac{18}{2125} \frac{(2285T^2 - 24982)z^4}{T^4} + \frac{1}{4250} \frac{141180T^2 - 1487411}{T^3} z^3 + \left(\frac{3}{340} \frac{1260T^2 - 11531}{T^2} + 1 \right) z^2 + 2Tz \right\} \end{aligned}$$

can accommodate this regime.



Figure 1. Depiction of the vertical profile of the current in the upper 200–400 m of the Pacific Ocean along the Equator: the westward wind-drift current is near the surface z = 0, below it is the eastward EUC that dominates the subsurface flows, with the weaker westward EIC found directly below the EUC (see the data provided in [13]).

Moreover the following theorem proves that also a hyperbolic azimuthal velocity profile coupled with the third-order polynomial is suitable in modeling of the aforementioned currents.

Theorem 3.1. An equatorial current profile featuring an eastward jet at mid-depth of the near-surface layer above the thermocline z = -T, a weak eastward jet just above the thermocline and a strong westward jet near the surface z = 0 is accommodated by hyperbolic functions coupled with the third-order polynomial

$$\alpha(z) = \left(z^3 + \frac{21T}{40}z^2 - \frac{17T^2}{40}z + \frac{T^3}{20}\right)\sinh(z+T),$$

$$\beta(z) = \varphi_0\left(\left(z^2 + Tz\right)T\sinh(z+T) - \frac{1}{100}\right).$$

Proof: It is easy to show that the azimuthal velocity component $u(\varphi, \theta, z)$ defined in the first relation of system (23), vanishes at z = -T/10, z = -T/2 and z = -4T/5, giving the profile depicted in Figure 1.

Simple calculations prove that

$$\begin{aligned} \alpha(0) &= \frac{T^3}{20} \sinh(T) > 0, \\ \phi_0 \alpha(0) + \beta(0) &= \phi_0 \left(\frac{T^3}{20} \sinh(T) - \frac{1}{100} \right) > 0 \\ \alpha'(0) &= -\frac{17T^2}{40} \sinh(T) + \frac{T^3}{20} \cosh(T) < 0, \end{aligned}$$

$$\begin{split} \phi_0 \alpha'(0) &+ \beta'(0) = \frac{T^2}{40} \phi_0(2T \cosh(T) + 23 \sinh(T)) > 0, \\ \alpha'(-T) &= 0, \\ \beta'(-T) &= 0. \end{split}$$

i.e. all the constraints (25)-(28) hold.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Kateryna Marynets D http://orcid.org/0000-0002-0043-6336

References

- [1] Philander SGH. The equatorial undercurrent revised. Ann Rev Earth Planetary Sci. 1980;8:191–204.
- Basu B. On a three-dimensional nonlinear model of Pacific equatorial ocean dynamics: velocities and flow paths. Oceanography. 2018;31:51–58.
- [3] Basu B. Some numerical investigations into a nonlinear three-dimensional model of the Pacific equatorial ocean flows. Deep Sea Res Part II: Top Stud Oceanogr. 2019;160:7–15.
- [4] Constantin A. An exact solution for equatorially trapped waves. J Geophys Res: Oceans. 2012;117:C05029.
- [5] Constantin A, Johnson RS. An exact, steady, purely azimuthal equatorial flow with a free surface. J Phys Oceanogr. 2016;46:1935–1945.
- [6] Constantin A, Johnson RS. A nonlinear, three-dimensional model for ocean flows, motivated by some observations of the Pacific equatorial undercurrent and thermocline. Phys Fluids. 2017;29:056604.
- [7] Henry D. An exact solution for equatorial geophysical water waves with an underlying current. European J Mech B/Fluids. 2013;38:18–21.
- [8] Henry D. Equatorially trapped nonlinear water waves in a β-plane approximation with centripetal forces. J Fluid Mech. 2016;804:R1.
- [9] Constantin A, Johnson RS. Ekman-type solutions for shallow-water flows on a rotating sphere: a new perspective on a classical problem. Phys Fluids. 2019;31:021401.
- [10] Marynets K. The modeling of the equatorial undercurrent using the Navier-Stokes equations in rotating spherical coordinates. Appl Anal. 2019; doi:10.1080/00036811.2019.1673375.
- [11] Constantin A, Johnson RS. Steady large-scale ocean flows in spherical coordinates. Oceanography. 2018;31:42-50.
- [12] Boyd JP. Dynamics of the equatorial ocean. Berlin: Springer; 2018.
- [13] Lukas R. Pacific ocean equatorial currents. In: Thorpe SA, Turekian KK, editors. Encyclopedia of ocean sciences. San Diego: Academic Press; 2001. p. 2069–2076.
- [14] Constantin A. Some nonlinear, equatorially trapped, nonhydrostatic internal geophysical waves. J Phys Oceanogr. 2014;44:781–789.
- [15] Constantin A, Germain P. Instability of some equatorially trapped waves. J Geophys Res: Oceans. 2013;118:2802–2810.
- [16] Constantin A, Ivanov RI. Equatorial wave-current interactions. Comm Math Phys. 2019;370:1-48.
- [17] Constantin A, Johnson RS. The dynamics of waves interacting with the equatorial undercurrent. Geophys Astrophs Fluid Dyn. 2015;109:311–358.
- [18] Constantin A, Johnson RS. An exact, steady, purely azimuthal flow as a model for the antarctic circumpolar current. J Phys Oceanogr. 2016;46:3585–3594.
- [19] Constantin A, Johnson RS. Large gyres as a shallow-water asymptotic solution of Euler's equation in spherical coordinates. Proc Roy Soc London A. 2017;473:Art. No. 20170063.
- [20] Constantin A, Johnson RS. Large-scale oceanic currents as shallow-water asymptotic solutions of the Navier–Stokes equation in rotating spherical coordinates. Deep Sea Res Part II: Top Stud Oceanogr. 2019;160:32–40.
- [21] Constantin A, Johnson RS. On the nonlinear, three-dimensional structure of equatorial oceanic flows. J Phys Oceanogr. 2019;49:2029–2042.
- [22] Haziot S. Study of an elliptic partial differential equation modelling the antarctic circumpolar current. Discr Cont Dyn Syst. 2019;39:4415–4427.
- [23] Haziot S, Marynets K. Applying the stereographic projection to modeling of the flow of the antarctic circumpolar current. Oceanography. 2018;31:68–75.

- [24] Henry D. Nonlinear features of equatorial ocean flows. Oceanography. 2018;31:22-27.
- [25] Henry D, Martin CI. Exact, free-surface equatorial flows with general stratification in spherical coordinates. Arch Ration Mech Anal. 2019;233:497–512.
- [26] Ionescu-Kruse D. A three-dimensional autonomous nonlinear dynamical system modelling equatorial ocean flows. J. Differ Equations. 2018;264:4650–4668.
- [27] Ionescu-Kruse D, Martin CI. Local stability for an exact steady purely azimuthal equatorial flow. J Math Fluid Mech. 2018;20:27–34.
- [28] Marshall J, Plumb RA. Atmosphere, ocean and climate dynamics: an introductory text. London: Academic Press; 2016.
- [29] Martin CI. On the existence of free-surface azimuthal equatorial flows. Appl Anal. 2017;96:1207–1214.
- [30] Marynets K. Two-point boundary problem for modeling the jet flow of the antarctic circumpolar current. Electron J Differ Equ. 2018;56:12.
- [31] Marynets K. A nonlinear two-point boundary-value problem in geophysics. Monatsh Math. 2019;188:287–295.
- [32] Marynets K. On a two-point boundary-value problem in geophysics. Appl Anal. 2019;98:553–560.
- [33] Marynets K. On the modeling of the flow of the antarctic circumpolar current. Monatsh Math. 2019;188:561-565.
- [34] Marynets K. The antarctic circumpolar current as a shallow-water asymptotic solution of Euler's equation in spherical coordinates. Deep Sea Res Part II: Top Stud Oceanogr. 2019;160:58–62.
- [35] Vallis GK. Atmosphere and ocean fluid dynamics. Cambridge: Cambridge University Press; 2006.