

Mitigation of greenhouse gas emissions from beef cattle production systems

Stine Samsonstuen , Bente A. Åby , Paul Crosson , Karen A. Beauchemin & Laila Aass

To cite this article: Stine Samsonstuen , Bente A. Åby , Paul Crosson , Karen A. Beauchemin & Laila Aass (2020): Mitigation of greenhouse gas emissions from beef cattle production systems, Acta Agriculturae Scandinavica, Section A — Animal Science, DOI: [10.1080/09064702.2020.1806349](https://doi.org/10.1080/09064702.2020.1806349)

To link to this article: <https://doi.org/10.1080/09064702.2020.1806349>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



[View supplementary material](#)



Published online: 27 Aug 2020.



[Submit your article to this journal](#)



Article views: 438







[View related articles](#)



[View Crossmark data](#)

Mitigation of greenhouse gas emissions from beef cattle production systems

Stine Samsonstuen ^a, Bente A. Åby ^a, Paul Crosson ^b, Karen A. Beauchemin ^c and Laila Aass^a

^aDepartment of Animal Sciences, Norwegian University of Life Sciences, Ås, Norway; ^bAnimal Grassland Research and Innovation Centre, Teagasc, Meath, Ireland; ^cAgriculture and Agri-Food Canada, Lethbridge Research and Development Centre, Lethbridge, Canada

ABSTRACT

The whole-farm model HolosNorBeef was used to estimate the efficiency of GHG emission mitigation strategies in Norwegian beef cattle herds. Various mitigation scenarios, involving female reproductive performance (i.e. calf mortality rate and the number of calves produced per cow per year), production efficiency of young bulls for slaughter (i.e. age at slaughter and carcass weight), and supplementation of an inhibitor currently reported as promising for enteric methane (CH₄) inhibition (3-nitrooxypropanol; 3-NOP) was investigated in herds of British and Continental breeds. Reducing calf mortality and increasing the number of produced calves per cow per year both reduced emission intensities by 3% across breeds. Continental breeds showed greater potential of reducing emission intensities due to increased carcass production. Combining mitigation options in a best case scenario reduced the total emissions by 11.7% across breeds. The emission intensities could be further reduced by 8.3% with the use of 3-NOP.

ARTICLE HISTORY

Received 24 March 2020
Accepted 17 July 2020

KEYWORDS

Beef production efficiency; farm scale model; methane inhibitor; mitigation options

Introduction


Beef consumption is expected to increase in both developed and developing countries as a consequence of global population growth (OECD/FAO, 2018). Thus, greenhouse gas (GHG) emissions from beef production are expected to increase. Beef has a large GHG emission intensity albeit with considerable variation among continents, countries (Gerber et al., 2013) and farms within a country (Bonesmo et al., 2013). The emission intensity of beef production depends upon breed (Hyslop, 2008), geographical location (White et al., 2010; Samsonstuen et al., 2019), farming system (Nguyen et al., 2010), and management practices (Alemu et al., 2017; Stanley et al., 2018). Hristov et al. (2013) showed potential long-term mitigation effects from ruminant production through improved reproductive performance, increased beef production and various management practices such as diet formulation, feed supplements, and manure management. Thus, the potential to reduce emission intensities is significant.

Animal productivity is important for beef farm profitability and is positively related to reductions in GHG emissions (Åby et al., 2014). The environmental impact of improved carcass production has been investigated by a number of studies (Thornton & Herrero, 2010; Desjardins et al., 2012; Legesse et al., 2016; Murphy et al., 2017; Legesse et al., 2018). Murphy et al. (2017)

showed decreased emission intensity when reducing age at slaughter, while increased average daily gain (ADG) reduced the emission intensities of Irish beef production systems (Casey & Holden, 2006; Crosson et al., 2010). The emission intensities from Canadian beef production have decreased from 1981 to 2011 due to improved reproduction efficiency, increased ADG, increased slaughter weight, reduced age at slaughter, and use of high grain diets that enabled slaughtering at a younger age (Legesse et al., 2016, 2018).

The environmental impact of female fertility and calf survival is inadequate or absent in most studies, as research mainly focuses on carcass production efficiency. Poor fertility and low calf survival increases the number of animals to maintain production levels and a stable herd size, hence a greater proportion of the GHG emissions is produced by herd replacements (Garnsworthy, 2004; Wall et al., 2010; Bell et al., 2011). Calf survival is of great importance in beef production systems, as the calf is the main product from the enterprise. Improvements in calf survival and cow fertility are known to reduce the overall emissions from beef production, as well as improving animal welfare (Wall et al., 2010). Beauchemin et al. (2011) reported a 4% reduction in GHG emissions following practices that improved calf survival to weaning, and Navajas et al.

CONTACT Stine Samsonstuen  stine.samsonstuen@nmbu.no

 Supplemental data for this article can be accessed at <https://doi.org/10.1080/09064702.2020.1806349>

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

(2010) reported reduction in emission intensities due to genetic improvement of fertility and calf survival.

Enteric CH₄ emissions account for approximately half the emissions from beef cattle production (Foley et al., 2011; Mogensen et al., 2015; Samsonstuen et al., 2019), hence various feed additives have been examined for their anti-methanogenic properties. These include various phyto-compounds (essential oils, oregano, garlic, green tea extract, condensed and hydrolysable tannins), microbials (live yeast, bacterial direct-fed probiotics), ionophores, dietary lipids, and chemical inhibitors (Hristov et al., 2013; Bayat et al., 2015, 2017; Kolling et al., 2018). However, many inhibitors have negative effects on feed intake (Hristov et al., 2013), organic matter fermentation in the rumen, digestibility (Johnson & Johnson, 1995), animal health and production (Hristov et al., 2013). However, the inhibitor 3-nitrooxypropanol (3-NOP, DSM Nutritional Products Ltd., Kaiseraugst, Switzerland) has shown promising long-term mitigation effects on enteric CH₄ emissions with no compromising effect on diet digestibility (Romero-Perez et al., 2014) or animal performance (Vyas et al., 2018).

The emission intensities from typical herds of British and Continental breeds in two geographically different regions in Norway were estimated by Samsonstuen et al. (2019). However, that study did not include GHG mitigation options such as improved cow efficiency, beef production efficiency or the effect of inhibitors. Furthermore, previous studies of calf survival (e.g. Beauchemin et al., 2011) and fertility (e.g. Bell et al., 2011) did not use nationally representative statistics or combinations of mitigation options. Additionally, pastoral systems have the potential to remove carbon (C) from the atmosphere through sequestration (Soussana et al., 2007), yet few studies have accounted for C sequestration when investigating mitigation options.

Thus, the aim of the study was to estimate the net mitigation potential in Norwegian beef cattle production systems using a whole-farm approach by investigating various scenarios, including variable cow and young bull beef production efficiency scenarios, as well as the mitigating effect of inhibitor (3-NOP) for enteric CH₄ reduction. The inhibitor evaluated was 3-NOP (DSM Nutritional Products Ltd., Kaiseraugst, Switzerland).

Materials and methods

This study was based on a previous study of GHG emissions from typical herds of British and Continental breeds in Norway (Samsonstuen et al., 2019). Fourteen mitigation scenarios were designed to reflect the variation in production efficiency among Norwegian beef cattle herds. The variable herd performances were

compared to the typical herds to investigate GHG mitigation potentials. For each scenario, the amount of beef carcass (kg) produced was based on the number of animals sent to slaughter, body weights and dressing percentages for the specific breed and animal class. Production enterprises on the farm not related to the cattle operation, such as the use of farm inputs (i.e. area, fertilizer, and pesticides) for grain production, were excluded from the analysis as the grain crops are sold from the farm and not used as feed. All cattle manure was applied on the included grasslands.

Baseline scenarios

Baseline (BL) scenarios were developed to represent each typical herd; British (Angus and Hereford) and Continental (Limousin, Simmental, and Charolais) breeds with associated geographical location, management, and production levels as described by Samsonstuen et al. (2019). For both breeds the farms were located in the flatlands (average altitude 246 m above sea level) of Norway, with a land size ranging between 45.4 and 50.1 ha.

The Norwegian beef cattle production system is semi-intensive with the cow-calf enterprise and finishing of bulls at the same farm. The BL farms were stocked with 28 spring-calving cows with the replacement rate set at 36% to keep the herd size constant (NIBIO, 2015). All progeny were retained for slaughter with males finished as bulls at 17.5 and 16.8 months, and surplus heifers not required to replace culled cows finished at 18.2 and 17.5 months for British and Continental breeds, respectively (Åby et al., 2012). Amount of beef carcass (kg) produced was calculated based on the number of cattle slaughtered to remain a stable herd size, slaughter weights and dressing percentages. Estimates of proportion of energy from concentrates and forage in the diet were from Åby et al. (2012). From birth to age at slaughter female progeny was fed 22% and 38% concentrates, and male progeny was fed 53% and 50% concentrates for British and Continental breeds, respectively. Cows were fed 25% (British) and 17% (Continental) concentrates throughout the year. Time spent on pasture for heifers were 19 and 13%, and for cows the time on pasture were 36 and 38% for British and Continental breeds, respectively. Manure was assumed to be deposited on pasture during the grazing period (June 1 to Sept 15) and handled as deep bedding during the housing period (Sept 16 to May 31). Manure was applied on ley area during spring. Silage yield (3350 kg dry matter (DM) ha⁻¹), pesticide (1.1 MJ ha⁻¹), and silage additive (21 kg Formic acid (CH₂O₂) ha⁻¹) use for a typical farm in the flatlands were

Table 1. Animal performance, land use and farm inputs for cow efficiency scenarios used to estimate GHG emission intensities from beef cattle operations.

Scenario	Cow efficiency scenarios									
	British					Continental				
	BL	1	2	3	4	BL	1	2	3	4
Animal system										
Still born calves (%)	3.5 ¹	0.0 ²	8.3 ²	3.5 ¹	3.5 ¹	3.9 ¹	0.0 ²	6.2 ²	3.9 ¹	3.9 ¹
Dead calves < 180 days (%)	3.6 ¹	0.0 ¹	10.8 ²	3.6 ¹	3.6 ¹	4.1 ¹	0.0 ²	10.9 ²	4.1 ²	4.1 ²
Calves cow ⁻¹ per year	1.0 ¹	1.0 ¹	1.0 ¹	0.9 ³	1.1 ³	1.0 ¹	1.0 ¹	1.0 ¹	0.9 ³	1.1 ³
Replacement heifers (year ⁻¹)	10	10	10	10	10	10	10	10	10	10
Heifers slaughtered (year ⁻¹)*	4	4	2	2	5	4	4	2	2	5
Young bulls slaughtered (year ⁻¹)*	13	14	12	12	15	13	14	12	12	15
ADG** heifers (g day ⁻¹)	645	645	645	645	645	784	784	784	784	784
ADG** young bulls (g day ⁻¹)	974	974	974	974	974	1212	1212	1212	1212	1212
Beef produced (kg carcass) ¹⁴	7699	8190	6841	7004	8303	9635	10311	8862	8815	10362
Land use										
Farm size (ha) ^{5***}	45.4	47.5	45.3	45.5	47.8	50.1	52.5	50.0	50.2	52.8
Of which: Ley area (ha) ^{5***}	39.7	41.8	39.7	39.8	42.1	44.4	46.8	44.4	44.5	47.1
Input use										
Concentrates (kg DM year ⁻¹)	44300	46178	41266	41141	47037	61244	63527	58937	57938	64184
Fuel (L year ⁻¹) ^{5***}	3931	4138	3930	3947	4171	4394	4641	4394	4406	4668
Silage additive (kg CH ₂ O ₂ year ⁻¹) ^{5***}	819	863	819	823	869	916	967	916	918	973

Note: BL = Baseline, typical beef cattle herd; CML = Calf mortality low; CMH = Calf mortality high; CYL = Calves cow⁻¹ per year low; CYH = Calves cow⁻¹ per year high; ADG = average daily gain.

¹Animalia (2017).

²Animalia (2019).

³Animalia (2018).

⁴Norwegian Institute of Bioeconomy research (NIBIO, 2015).

⁵Norwegian Institute of Bioeconomy research (NIBIO, 2016).

*Heifers and young bulls available for slaughter varies across scenarios dependent on number of produced calves cow per year and calf mortality

**Average daily gain from birth to age at slaughter.

***Silage additives for conservation of grass by ensiling. Corresponds to the ley area required to cover the forage requirement.

obtained from Norwegian Institute of Bioeconomy Research (NIBIO, 2015). The ley area (ha) corresponded to the calculated forage requirements plus an additional 10% (DM basis) to account for losses due to ensiling according to Bonesmo et al. (2013). N-fertilizer application for conserved feed (13 kg N ha⁻¹) followed advisory based recommendations for forage production (NIBIO, 2016). Dry matter content and nutritive value (0.87 FU kg DM⁻¹) of forage was estimated for the flatlands based on laboratory analysis information (Eurofins, Moss, Norway). Use of electricity (26,300 kWh year⁻¹) and fuel (99 L ha⁻¹) for a typical farm in the flatlands was from operational farm data (NIBIO, 2015). Seasonal soil and weather data were available through Skjelvåg et al. (2012; supplemental Table S1).

Alternative scenarios

For each alternative scenario, the herd size and structure (number of cows and replacement heifers) were kept constant corresponding to the BL scenario. The ley area (ha) varied across scenarios and corresponded to the calculated forage requirements plus 10% loss (Tables 1 and 2). Forage yields (kg ha⁻¹), use of silage additives for conservation of grass through ensiling (kg CH₂O₂ ha⁻¹),

fertilizers (kg N ha⁻¹), and fuel (L ha⁻¹) were kept constant per ha, yielding different total amounts for each scenario (Tables 1 and 2).

Cow efficiency scenarios (Table 1) were based on the observed variation in calf mortality and the number of calves born per cow per year from the Norwegian Beef Herd Recording System (NBS; Animalia, 2018, 2019). The calf mortality among Norwegian beef cattle production herds ranges from 0% to 20% with a positively skewed distribution. Scenarios were based on the observed proportion of stillborn and dead calves prior to 180 days among 95% of the herds with British and Continental breeds in the NBS (Scenario 1 (CML) and 2 (CMH)). The number of calves produced per cow per year was based on the observed production among the worst 1/3 and best 1/3 of Norwegian herds (Scenario 3 (CYL) and 4 (CYH)).

Young bull beef production efficiency scenarios (Table 2) were based on age at slaughter and carcass weight for young bulls among the worst and best 1/3 of the Norwegian herds from the annual report of NBS (Animalia, 2018; Scenario 5 (BPL) and 6 (BPH)). The proportion of concentrates in the diet and days on pasture were kept constant across scenarios, influencing the required ley areas (ha) to cover the animal requirements.

Table 2. Animal performance, land use and farm inputs for young bull beef production efficiency, best case (BC) and worst case (WC) scenarios used to estimate GHG emission intensities from beef cattle operations.

Scenario	Young bull beef production efficiency scenarios						Best case/worst case scenarios			
	British			Continental			British		Continental	
	BL	5 BPL	6 BPH	BL	5 BPL	6 BPH	7 WC	8 BC	7 WC	8 BC
Animal system										
Still born calves (%)	3.5 ¹	3.5 ¹	3.5 ¹	3.9 ¹	3.9 ¹	3.9 ¹	8.3 ²	0.0 ²	6.2 ²	0.0 ²
Dead calves <180 days (%)	3.6 ¹	3.6 ¹	3.6 ¹	4.1 ¹	4.1 ¹	4.1 ¹	10.8 ²	0.0 ²	10.9 ²	0.0 ²
Calves cow ⁻¹ per year	1.0 ¹	1.0 ¹	1.0 ¹	1.0 ¹	1.0 ¹	1.0 ¹	0.9 ³	1.1 ³	0.9 ³	1.1 ³
Young bulls, age at slaughter (month)	17.5 ⁴	18.7 ²	16.1 ²	16.8 ⁴	18.1 ²	15.4 ²	18.7 ²	16.1 ²	18.1 ²	15.4 ²
Young bulls, carcass weight (kg)	291 ¹	256 ³	334 ³	353 ¹	317 ³	392 ³	256 ³	334 ³	317 ³	392 ³
Replacement heifers (year ⁻¹)	10	10	10	10	10	10	10	10	10	10
Heifers slaughtered (year ⁻¹)*	4	4	4	4	4	4	1	6	1	6
Young bulls slaughtered (year ⁻¹)*	13	13	13	13	13	13	10	16	10	16
ADG** heifers (g day ⁻¹)	645	645	645	784	784	784	645	645	784	784
ADG** young bulls (g day ⁻¹)	974	794	1227	1212	1002	1479	794	1227	1002	1479
Beef produced (kg carcass) ¹⁴	7699	7232	8272	9635	9157	10159	5868	9509	7721	11700
Land use										
Farm size (ha) ^{5***}	45.4	45.6	47.9	50.1	51.5	50.8	43.6	44.9	49.1	53.0
Of which: Ley area (ha) ^{5***}	39.7	39.9	42.2	44.4	45.8	45.1	37.9	44.2	43.4	47.3
Input use										
Concentrates (kg DM year ⁻¹)	44300	40941	48299	61244	61554	58725	35688	53736	55987	63604
Fuel (L year ⁻¹) ^{5***}	3931	3951	4177	4394	4533	4472	3751	4375	4304	4683
Silage additive (kg CH ₂ O ₂ year ⁻¹) ^{5***}	819	823	871	916	945	932	782	912	897	976

Note: BL = baseline, typical beef cattle herd; BPL = Young bull beef production efficiency, low; BPH = Young bull beef production efficiency, high; WC = Worst case, worst performing 1/3 of Norwegian beef cattle farms; BC = Best case, best performing 1/3 of Norwegian beef cattle farms; ADG = average daily gain.

¹Animalia (2017).

²Animalia (2019).

³Animalia (2018).

⁴Norwegian Institute of Bioeconomy research (NIBIO, 2015).

⁵Norwegian Institute of Bioeconomy research (NIBIO, 2016).

* Heifers and young bulls available for slaughter varies across scenarios dependent on number of produced calves cow per year and calf mortality

** Average daily gain from birth to age at slaughter

*** Silage additives for conservation of grass by ensiling. Corresponds to the ley area required to cover the forage requirements.

Unfavorable cow efficiency and young bull beef production efficiency scenarios were combined in a worst case (WC; Scenario 7) scenario. The corresponding best case (BC; Scenario 8) scenario was a combination of favorable scenarios (Table 2). The effect of feeding a low level of the inhibitor 3-NOP (100 mg/kg DM) on enteric CH₄ emissions was included in the BL scenario (BLinL; Scenario 9), the WC scenario (WCinL; Scenario 10), and BC scenario (BCinL; Scenario 11) for the two typical herds of British and Continental breeds (Table 2). The effect of feeding a high level of the inhibitor (237 mg/kg DM) was included in the BL scenario (BLinH; Scenario 12), the WC scenario (WCinH; Scenario 13), and corresponding BC scenario (BCinH; Scenario 14). Dietary supplementation of the inhibitor 3-NOP was based on the findings by Romero-Perez et al. (2014), Vyas et al. (2016) and Vyas et al. (2018). The inhibitor was fed during the housing period (8.5 months) for cows and backgrounding and finishing stock (6–24 months in age), respectively. It was assumed that on the days the inhibitor was fed, enteric CH₄ emissions expressed as a percentage of dry matter intake (DMI) were decreased by 7 (low) and 33% (high) with no negative effects on DMI or ADG.

Modeling GHG emissions

The GHG emissions were estimated using HolosNorBeef developed by Samsonstuen et al. (2019). HolosNorBeef is an empirical model specifically developed for beef cattle production systems under Norwegian conditions, using Tier 2 methodology of the Intergovernmental Panel on Climate Change (IPCC, 2006). The model estimated the GHG emissions on an annual time step for the land use and management changes and on a monthly time step for animal production, accounting for differences in diet, housing, and climate. HolosNorBeef estimated whole-farm GHG emissions by considering direct emissions of CH₄ from enteric fermentation and manure, nitrous oxide (N₂O) and carbon dioxide (CO₂) from on-farm livestock production including soil C changes, and indirect N₂O and CO₂ emissions associated with run-off, nitrate leaching, ammonia volatilization and from inputs used on the farm. The sources of emissions, emission factors, and equations used by the HolosNorBeef model are given in supplemental Table S2. All emissions were expressed as CO₂ eq to account for the global warming potential (GWP) of the respective gases for a time horizon of 100 years: CH₄

Table 3. Total greenhouse gas (GHG) emissions (kg CO₂ eq) and emission intensities (kg CO₂ eq kg⁻¹ carcass) by source for cow efficiency scenarios from beef cattle operations.

Scenario	British					Continental				
	BL	1 CML	2 CMH	3 CYL	4 CYH	BL	1 CML	2 CMH	3 CYL	4 CYH
Total GHG emissions	236,984	244,780	229,886	229,503	247,387	281,879	291,482	275,796	273,465	293,554
Emission intensities by source										
Enteric CH ₄	14.03	13.50	15.20	14.82	13.45	13.24	12.70	14.03	13.95	12.73
Manure CH ₄	3.22	3.11	3.47	3.40	3.09	3.17	3.05	3.34	3.33	3.05
Manure N ₂ O	3.01	2.89	3.28	3.20	2.87	2.78	2.66	2.96	2.95	2.66
Soil N ₂ O	3.03	2.94	3.33	3.25	2.93	2.86	2.77	3.05	3.06	2.77
Soil C*	-1.72	-1.62	-1.77	-1.74	-1.61	-1.85	-1.74	-1.90	-1.89	-1.74
Off-farm barley	1.94	1.90	2.04	1.98	1.91	2.15	2.08	2.25	2.22	2.09
Off-farm soy	1.89	1.85	1.98	1.93	1.86	2.09	2.02	2.18	2.16	2.03
Indirect energy	3.03	3.00	3.43	3.35	2.98	2.70	2.67	2.94	2.97	2.67
Direct energy	2.34	2.31	2.65	2.58	2.30	2.09	2.06	2.27	2.29	2.06
Total emission intensities	30.78	29.89	33.61	32.77	29.80	29.23	28.27	31.12	31.02	28.33
Total emission intensities excluding soil C	32.49	31.50	35.37	34.51	31.40	31.08	30.01	33.02	32.92	30.07

Note: BL: Baseline, typical beef cattle herd; CML: Calf mortality low; CMH: Calf mortality high; CYL: Calves cow⁻¹ per year low; CYH: Calves cow⁻¹ per year high. *Negative values indicate carbon sequestration.

(kg) × 28 + N₂O (kg) × 265 + CO₂ (kg) (Myhre et al., 2013). Emission intensities were expressed as kg CO₂ eq (kg beef carcass)⁻¹.

Results

Total emissions

The total emissions per year for the BL scenario representing typical herds were 237 t CO₂ eq for British and 282 t CO₂ eq for Continental breeds (Tables 3–5). For both breeds, cow efficiency scenarios (Scenario 1–4) resulted in decreased total emissions when reducing the calf survival (CMH) and number of calves per year (CYL). Increased calf survival (CML) and number of calves per cow (CYH) increased the total emissions, compared with BL scenarios. The young bull carcass

production scenarios (Scenario 5–6) differed across breeds. High age at slaughter and low carcass weight (BPL) reduced the total emission by 3.3% for the British breeds, whereas the scenario with low age at slaughter and high carcass weight (BPH) had a 1.6% lower total emissions for Continental breeds. In the combined scenarios (Scenario 7–8), high calf survival, high number of calves produced per cow and high young bull carcass production (BC) increased total emissions by 12.1 and 4.1% for British and Continental breeds, respectively. By including the effect of 3-NOP in the combined scenarios (Scenario 9–14), the total emissions were decreased with both supplementation rates for baseline (BLinL, BLinH) and worst case (WCinL, WCinH) scenarios compared with the BL scenarios. In the best case scenarios, a low supplementation rate (BCinL) increased the total emissions by 9.6 and 1.7%

Table 4. Total greenhouse gas (GHG) emissions (kg CO₂ eq) and emission intensities by source (kg CO₂ eq kg⁻¹ carcass) for young bull beef production efficiency, worst case (WC) and best case (BC) scenarios from beef cattle operations.

Scenario	Young bull beef production efficiency scenarios						Best case/worst case scenarios			
	British			Continental			British		Continental	
	BL	5 BPL	6 BPH	BL	5 BPL	6 BPH	7 WC	8 BC	7 WC	8 BC
Total GHG emissions	236,984	229,264	249,453	281,879	283,990	277,415	213,280	265,652	266,249	293,415
Emission intensities by source										
Enteric CH ₄	14.03	14.34	13.60	13.24	13.94	12.28	16.47	12.58	15.51	11.27
Manure CH ₄	3.22	3.29	3.14	3.17	3.36	2.91	3.76	2.92	3.72	2.68
Manure N ₂ O	3.01	3.08	2.91	2.78	2.91	2.59	3.58	2.67	3.27	2.36
Soil N ₂ O	3.03	3.14	2.97	2.86	3.05	2.68	3.64	2.73	3.41	2.45
Soil C*	-1.72	-1.67	-1.67	-1.85	-1.94	-1.63	-1.88	-1.57	-2.12	-1.51
Off-farm barley	1.94	1.91	1.97	2.15	2.27	1.95	2.05	1.91	2.45	1.84
Off-farm soy	1.89	1.86	1.92	2.09	2.21	1.90	2.00	1.86	2.38	1.79
Indirect energy	3.03	3.25	3.00	2.70	2.94	2.61	3.80	2.73	3.31	2.38
Direct energy	2.34	2.50	2.31	2.09	2.27	2.02	2.93	2.11	2.55	1.83
Total emission intensities	30.78	31.70	30.16	29.23	31.01	27.31	36.34	27.94	34.48	25.08
Total emission intensities excluding soil C	32.49	33.37	31.82	31.08	32.96	28.94	38.22	29.51	36.61	26.59

Note: BL: Baseline, typical beef cattle herd; BPL: Young bull beef production efficiency, low; BPH: Young bull beef production efficiency, high; WC: Worst case, poorest performing 1/3 of Norwegian beef cattle farms; BC: Best case, best performing 1/3 of Norwegian beef cattle farms.

*Negative values indicate carbon sequestration.

Table 5. Total greenhouse gas (GHG) emissions (kg CO₂ eq) and emission intensities by source (kg CO₂ eq kg⁻¹ carcass) for worst case (WC) and best case (BC) scenarios with and without the reduction of enteric methane (CH₄) from low and high levels of the inhibitor 3-nitrooxypropanol (3-NOP).

Application level 3-NOP Scenario	British							Continental						
	100 mg (kg DM) ⁻¹			237 mg (kg DM) ⁻¹				100 mg (kg DM) ⁻¹			237 mg (kg DM) ⁻¹			
	BL	9 BLinL	10 WCinL	11 BCinL	12 BLinH	13 WCinH	14 BCinH	BL	9 BLinL	10 WCinL	11 BCinL	12 BLinH	13 WCinH	14 BCinH
Total GHG emissions	236,984	232,262	209,525	260,619	214,186	193,386	240,975	281,879	262,247	258,580	287,795	254,829	234,798	265,594
Emission intensities by source														
Enteric CH ₄	14.03	13.29	15.67	11.95	10.77	12.71	9.74	13.24	11.11	14.40	10.70	10.15	11.85	8.66
Manure CH ₄	3.22	3.22	3.76	2.92	3.22	3.76	2.92	3.17	3.17	3.72	2.68	3.17	3.72	2.68
Manure N ₂ O	3.01	3.01	3.58	2.67	3.01	3.58	2.67	2.78	2.78	3.27	2.36	2.78	3.27	2.36
Soil N ₂ O	3.03	3.03	3.64	2.73	3.03	3.64	2.73	2.86	2.86	3.41	2.45	2.86	3.41	2.45
Soil C*	-1.72	-1.72	-1.88	-1.57	-1.72	-1.88	-1.57	-1.85	-1.85	-2.12	-1.51	-1.85	-2.12	-1.51
Off-farm barley	1.94	1.94	2.05	1.91	1.94	2.05	1.91	2.15	2.15	2.45	1.84	2.15	2.45	1.84
Off-farm soy	1.89	1.89	2.00	1.86	1.89	2.00	1.86	2.09	2.09	2.38	1.79	2.09	2.38	1.79
Indirect energy	3.03	3.03	3.80	2.73	3.03	3.80	2.73	2.70	2.70	3.31	2.38	2.70	3.31	2.38
3-NOP**	0.00	0.12	0.16	0.10	0.30	0.38	0.25	0.00	0.10	0.12	0.09	0.29	0.36	0.22
Direct energy	2.34	2.34	2.93	2.11	2.34	2.93	2.11	2.09	2.09	2.55	1.83	2.09	2.55	1.83
Total emission intensities	30.78	30.16	35.70	27.41	27.82	32.95	25.34	29.23	27.19	33.49	24.60	26.42	31.19	22.70
Total emission intensities excluding soil C	32.49	31.88	37.58	28.98	29.53	34.83	26.91	31.08	29.04	35.61	26.11	28.27	33.31	24.21

Note: BL: baseline, typical beef cattle herd; BLinL: baseline with 7% reduction of enteric CH₄ emissions from the inhibitor; WCinL: Worst case farms with 7% reduction of enteric CH₄ emissions from the inhibitor; BCinL: Best case farms with 7% reduction of enteric CH₄ emissions from the inhibitor; BLinH: baseline with 33% reduction of enteric CH₄ emissions from the inhibitor; WCinH: Worst case farms with 33% reduction of enteric CH₄ emissions from the inhibitor; BCinH: Best case farms with 33% reduction of enteric CH₄ emissions from the inhibitor.

*Negative values indicate carbon sequestration.

**Emissions related to production and transport of 3-NOP.




Figure 1. Emission intensities (CO₂ eq kg⁻¹ carcass) for each scenario including and excluding soil carbon (C) balance. In all scenarios, the farm was located in the flatlands in Norway and stocked with 28 cows. BL: baseline; CML: calf mortality low; CMH: calf mortality high; CYL: calves per cow per year low; CYH: calves per cow per year high; BPL: young bull beef production efficiency low; BPH: young bull beef production efficiency high; WC: worst case; BC: best case; BLinL: baseline with inhibitor low; WCinL: worst case with inhibitor low; BCinL: best case with inhibitor low; BLinH: baseline with inhibitor high; WCinH: worst case with inhibitor high; BCinH: best case with inhibitor high.

for British and Continental breeds, respectively. At high application level (BCinH) the total emissions increased 0.7% for the British breeds and decreased 5.8% for the Continental breeds.

Emission intensities

The emission intensities (Figure 1) for the BL scenario were greater for the British breeds (30.8 kg CO₂ eq (kg carcass)⁻¹) compared with the Continental breeds (29.2 kg CO₂ eq (kg carcass)⁻¹; Table 4). Enteric CH₄ contributed most to the GHG emissions, accounting for 45–46% of the total emissions. Nitrous oxide from manure and soil were the second largest source, accounting for

20–21% of the total emissions. Manure CH₄ accounted for 10–11% and soil C balance was negative for both breed types, indicating C sequestration. Emission intensities for the cow efficiency scenarios varied from 28.3 kg CO₂ eq (kg carcass)⁻¹ for the CML and CYH scenarios for Continental breeds to 33.6 kg CO₂ eq (kg carcass)⁻¹ for the CMH scenario for British breeds (Table 3). Across breeds, reduced calf mortality and increased number of calves per cow per year each reduced the emission intensities by 3.1% compared with the BL scenario, whereas CMH and CYL increased the emission intensities by 7.8 and 6.3%, respectively.

The Continental breeds demonstrated greater reduction in emission intensities with increased carcass

Table 6. Net reduction potential (t CO₂ eq) from implementing mitigation options by Norwegian from beef cattle operations assuming a constant production of 28,516 t carcass year⁻¹.

Scenario	Net reduction potential (t CO ₂ eq)		
	Improving the 1/3 poorest performing herds to BL level	Improving the BL level to the best performing level	
Calf mortality	CML + CMH	-8,792	-26,377
Calves per cow year ⁻¹	CYL + CYH	-8,935	-26,805
Young bull beef production	BPL + BPH	-12,072	-36,215
Combined*	WC + BC	-33,221	-99,662
BL inhibitor, low rate	BLinL	-12,642	-37,926
BL inhibitor, high rate	BLinH	-27,423	-82,268
Combined* inhibitor, low rate	WCinL + BCinL	-38,021	-114,063
Combined* inhibitor, high rate	WCinH + BCinH	-56,699	-170,096

Note: BL: baseline; (855,611 t CO₂ eq); CML: calf mortality low; CMH: calf mortality high; CYL: calves per cow per year low; CYH: calves per cow per year high; BPL: young bull beef production efficiency low; BPH: young bull beef production efficiency high; WC: worst case; BC: best case; BLinL: baseline with inhibitor low; BLinH: baseline with inhibitor high; WCinL: worst case with inhibitor low; BCinL: best case with inhibitor low; WCinH: worst case with inhibitor high; BCinH: best case with inhibitor high.

*Combination of calf mortality, calves per cow year⁻¹ and young bull beef production.

production compared to the British breeds (Table 4). Reduced carcass production and increased age at slaughter in the BPL scenario, increased the emission intensities by 3.0% and 6.1% for British and Continental breeds, respectively. Increased carcass weight (BPH) and reduced age at slaughter reduced the emission intensities by 2.0% for British and 6.6% for Continental breeds.

In the combined scenarios, larger effects on GHG intensities were observed. For the BC and WC scenarios, emission intensities varied from 27.9 to 36.3 kg CO₂ eq (kg carcass)⁻¹ for British breeds and from 25.1 to 34.5 kg CO₂ eq (kg carcass)⁻¹ for Continental breeds (Table 4). The BC scenario reduced the emission intensities by 11.7% on average. When the inhibitor 3-NOP was included during the housing period, the emission intensities were reduced by 4.5 and 9.6% across breeds for the BLinL and BLinH scenarios, respectively (Table 5). For the British breeds, the inhibitor reduced the emission intensities by 2.0 (BLinL) and 9.6% (BLinH) compared with the BL scenario, whereas the Continental breeds had 7.0 (BLinL) and 9.6% (BLinH) reductions. High supplementation rate of the inhibitor in the WCinH scenario offset more than half the increase in emission intensities in the WC scenario, resulting in only 6.9% greater emission intensity across breeds compared to the BL scenario (Table 5).

Improving cow efficiency and young bull carcass production among Norwegian beef cattle production herds reduced net emissions by 26,377 and 26,805 t CO₂ eq, respectively (Table 6). The largest net mitigation potential was obtained by combining both improved performance and feeding of the inhibitor 3-NOP at a high supplementation rate (Table 6).

Discussion

Our study investigated GHG mitigation options including cow efficiency, young bull beef production efficiency, combination of cow and bull efficiency, and use of CH₄ inhibitor (3-NOP) for the typical beef herds of British and Continental breeds in Norway. The study adopted the unique approach of combining mitigation strategies because, individually, most mitigation strategies have low to moderate impact on decreasing emissions, whereas our study shows that combining strategies may help achieve the decreases in GHG emissions needed from the beef industry. 3-Nitrooxypropanol has been identified as a highly promising mitigation strategy for enteric CH₄ reduction. However the inhibitor 3-NOP is currently not approved by the Norwegian authorities, so the reduction potential from applying the inhibitor is theoretical at this point. Should the approval status of

3-NOP change in the near future (i.e. 3-NOP is undergoing review by authorities in the E.U.), the study demonstrates the possible implications for beef production in Norway. Unlike many previous farm scale studies for beef production (e.g. Foley et al., 2011), our analysis included changes in soil C in the model and demonstrates that forage-based beef production systems can offset some of the CH₄ emissions by preserving or enhancing soil C reserves. Differences in soil C cause differences across farms. However, as the yields and use of fertilizer are kept constant within a farm, the different scenarios causes a proportional change in soil C due to changes in animal manure.

HolosNorBeef estimated emission intensities from the BL scenarios of 29.2–30.8 kg CO₂ eq (kg carcass)⁻¹ for typical herds of British and Continental breeds. This range of emission intensities is similar to both other Nordic countries; Denmark 23.1–29.7 kg CO₂ eq (kg carcass)⁻¹ and Sweden 25.4 kg CO₂ eq (kg carcass)⁻¹ (Mogensen et al., 2015), and the typical herds of British and Continental breeds considered by Samsonstuen et al. (2019) (range: 27.5–32.01 kg CO₂ eq (kg carcass)⁻¹). The present study found that Norwegian beef production systems have potential to reduce emission intensities without substantial changes in the enterprise by adopting practices that improve female fertility and calf survival and increase carcass production. The risk of stillbirth could be reduced by supervision during calving, whereas ensuring colostrum and good hygiene could reduce mortality after calving. Good management and feeding of heifers to ensure optimal growth prior to mating help reduce calving difficulties and stillbirths. In addition to herd size and housing conditions, the on-farm calving management and workload is dependent on the length of the calving period (Murray et al., 2015). Seasonal calving could also ease the hygiene management and thus be beneficial to improving calf health and survival. Increased number of calves produced per cow per year through improved culling management, higher pregnancy rates, and fewer abortions and empty cycles is highly dependent on the individual farmers professional knowledge and experience and do not require additional resource input in the enterprise. Carcass production could also be increased by selecting breeding candidates with high genetic merit for feed efficiency that produce larger offspring (Arthur & Herd, 2005), or by dietary improvements such as higher forage quality from improved agronomic practices (Randby et al., 2010). Differences in GHG emissions were demonstrated between typical farm conditions (BL scenario). Across breeds, the alternative scenarios CML, CYH, BC, and BCinL resulted in greater total emissions, whereas scenarios CMH, CYL, BPL, WC, BLinL,

WCinL, BLinH, and WCinH resulted in lower total emissions compared to BL. Scenarios BPH and BCinH were dependent upon breed. Higher levels of production were associated with higher levels of inputs (total use of pesticides, fertilizer, and fuel) resulting in greater total on-farm emissions compared with the BL scenarios. However, when expressed per kg carcass, the scenarios with increased cow and beef production efficiency reduced emission intensities (by 2.0–14.2%) compared with the BL scenarios.

In all scenarios, C sequestration had a mitigating effect on GHG emissions. Emission intensities vary due to location, resources, and climatic conditions (White et al., 2010; Samsonstuen et al., 2019). While most farm-level modeling studies assume that soil carbon is at equilibrium, Soussana et al. (2007) has shown that European grasslands act as atmospheric carbon sinks. In Denmark, Sweden and Norway, studies have also shown forage lands to be sequestering carbon (Mogensen et al., 2015; Samsonstuen et al., 2019). Those estimates of carbon sequestration are in the range of the level of C sequestration estimated in the present study. Bonesmo et al. (2013) reported variability in emission intensities from soil N₂O and soil C among Norwegian dairy farms. In the current study, a single location was considered with the initial SOC, temperature, and moisture held constant across scenarios. Forage production and application of N-fertilizer were also held constant per ha. Hence, differences in C sequestration were dependent upon the application of manure and the ley area (ha). As the ley area was a function of animal requirements and DMI, these relationships resulted in lower C sequestration (kg CO₂ eq kg⁻¹ carcass) for scenarios where the production efficiency was increased (CML, CYH, BPH, BC) (Soil C; Tables 3 and 4).

Due to low reproductive rate, the impact of offspring survival is larger for cattle compared to pigs. Hence, offspring survival is of great importance for both economics (Azzam et al., 1993) and GHG emissions from beef cattle production (Wall et al., 2010). Calf mortality may be reduced by improving calving and maternal traits both through breeding (i.e. breeding for moderate birth weights) and improved management, such as providing colostrum, good hygiene at calving and navel dipping to reduce infections (Wall et al., 2010; Murray et al., 2015). The CML scenario had low calf mortality, which increased total forage requirements, area needed for forage production, and the total use of inputs (i.e. N-fertilizer and fuel). A larger number of heifer and bull calves were sent to slaughter, which increased the total beef production from the farm. Hence, the low calf mortality scenario (CML) lowered the emission intensities by 3.1% compared with the BL

scenarios, which corresponded to the reported reduction in emission intensity (4%) from improved calf survival reported by Beauchemin et al. (2011). Improved female fertility may reduce both management costs and emissions (Wall et al., 2010). The best and worst 1/3 of the Norwegian beef cattle farms produce on average 1.1 and 0.9 calves per cow per year, respectively (Animalia, 2018). An increased number of calves produced per cow may be obtained by improved culling management, higher pregnancy rates, and fewer abortions and empty cycles, which may all be achieved through good management, health, and nutrition.

Production efficiency is essential for reducing the emission intensities from beef production systems (Hyslop, 2008). At low production levels, the number of cattle required to produce the same amount increases. Historically, the number of dairy cattle in Norway has decreased approximately 35% since 1990 as a consequence of increased milk yield (Statistics Norway, 2019). Thus, the number of beef cattle with a greater carcass production has increased to meet the increasing demand for domestic beef. Greater animal productivity through increased carcass production increases the gross efficiency by diluting the maintenance costs of the production animals (Wall et al., 2010). Intensive concentrate-based systems produce lowest emissions per kg beef (Hyslop, 2008) as such diets increase ADG and shorten the finishing period, thereby reducing enteric CH₄ emissions (Lovett et al., 2010). In the present study, the carcass output from the farms varied across scenarios (Tables 1 and 2) with a constant number of cows due to differences in female fertility, calf survival and animal productivity. In accordance with Veysset et al. (2014), the emission intensities decreased with greater animal productivity, due to reduced age at slaughter and increased carcass weights. Higher young bull efficiency (BPH) resulted in a larger reduction in emission intensities for the Continental breeds (6.6%) than the British breeds (2.0%) compared to BL, which reflects greater unexploited potential for increased carcass production for Continental breeds.

Enteric CH₄ accounts for 43.9–55.7% of total GHG emissions from beef cattle production (Foley et al., 2011; Mogensen et al., 2015; Samsonstuen et al., 2019) and is mainly related to variation in feed quality (Ominski et al., 2011) and DMI (Herd et al., 2014). Alemu et al. (2017) reported substantial variation in enteric CH₄ emissions among Canadian farms due to variation in diet composition and diet quality. Hence, reduced enteric CH₄ emissions through nutrition is often seen to be an ideal mitigation strategy. In the present study, differences in enteric CH₄ emissions were related to the number of animals, ADG, and age

at slaughter, as the forage quality and proportion of concentrates/pasture were kept constant within breed across scenarios. The reduction in enteric CH₄ emissions from beef cattle by feeding 3-NOP in backgrounding diets varies from 4 to 59% dependent on diet composition and level of application (Romero-Perez et al., 2014; Vyas et al., 2016, 2018). The scenarios investigating the effect of 3-NOP assumed 7 or 33% reduction of enteric CH₄ emissions with no negative effects on performance or DMI. The effect of the inhibitor was only considered during the housing period as feeding supplements to cattle on pasture is challenging. At high supplementation rates, Vyas et al. (2016) reported reduced DMI ($P < 0.01$) during the backgrounding phase and a tendency ($P = 0.06$) for reduced DMI during the finishing phase, whereas Romero-Perez et al. (2014) showed no significant reduction of DMI. Hence, the emissions in the present study might be over-estimated as the inhibitor was assumed to have no effect on DMI. The reduction in emission intensities could potentially be greater if performance was improved or higher if DMI decreased with no influence on ADG. With 3-NOP, the reduction in enteric CH₄ emissions more than offset the increase in indirect energy emissions from manufacturing the inhibitor, regardless of level of supplementation. At high supplementation levels, 3-NOP offset more than half the increase in emission intensities of low production efficiency and poor management, as the WCin scenario had 6.9% greater CO₂ eq emissions across breeds compared with the BL scenario. Currently, the inhibitor (3-NOP) is only available for research purposes as the long-term effect of feeding the supplement needs further investigation for the inhibitor to be approved for use on commercial farms. Hence, the scenarios investigating the mitigation potential by feeding 3-NOP are highly theoretical. Additionally, 3-NOP might influence other emission sources, such as cattle manure and corresponding soil C balance, which warrants further investigation of the inhibitor as a mitigation option.

The market demand for beef is a prerequisite for domestic production, and mitigation options to reduce the national GHG emissions from beef production need to be investigated in relation to the production level. Increased animal production efficiency could contribute to reduced national emissions both by reducing the total number of beef cows and by reducing emission intensities at the farm-level. The Norwegian beef cattle population produces approximately 28,516 t carcass year⁻¹ (Nortura, 2019), corresponding to approximately 856,000 t CO₂ eq at BL level. By improving calf mortality, increasing the number of calves per cow per year, and increasing young bull carcass production of the

poorest performing 1/3 herds to BL level, total emissions could be reduced by 8,792, 8,935 and 12,072 t CO₂ eq, respectively (Table 6). The results from this study indicate that the total potential for reducing GHG emissions from Norwegian beef cattle production ranges from 8,000 to 57,000 t CO₂ eq year⁻¹ depending upon mitigation option. Over a 10 year period, the scenario combining improved performance and high 3-NOP supplementation rates (BCinH) exceeds the 5 mill ton reduction of GHG emissions required from the agricultural sector according to the agreement between the Norwegian Farmers Union and the Norwegian Government (Norwegian Farmers Union et al., 2019) while maintaining the same level of production.

Genetic improvement of livestock is cost effective and produces permanent and cumulative changes in performance, and can improve farm profitability and reduce emissions through improved animal productivity and efficiency, reduced wastage (i.e. reduced involuntary culling and empty reproductive cycles) and direct selection for low-emission animals (Wall et al., 2010; Åby et al., 2014). Other measures, such as the use of inhibitors are highly effective at reducing emissions, but increase input costs. A premise for farmers to implement on-farm mitigation options is that the extra efforts are considered profitable. Thus, adoption may require subsidy financing to encourage implementation unless a gain in production efficiency is also realized.

Conclusions

The baseline scenario estimated a farm gate GHG emission intensity of 30.8 and 29.2 kg CO₂ eq (kg carcass)⁻¹ for British and Continental breeds, respectively. Mitigation strategies that improve cow efficiency by reducing calf mortality and increasing the number of calves born per cow per year each reduced emission intensities by 3.1% across breeds. Improving young bull beef production efficiency had greater mitigation potential for Continental breeds (-6.6%) compared with British breeds (-2.0%). When mitigation options were combined, the emission intensities were reduced by 11.7% across breeds. Assuming no negative effect on performance or DMI, the inhibitor 3-NOP reduced the net GHG emissions from beef cattle production dependent on application level. At a constant national level of beef production, total national emissions can be reduced by implementing one or a combination of mitigation options aimed at improving female fertility, increasing carcass production or reducing enteric CH₄ production using an inhibitor. However, despite a decrease in emission intensity, the total emissions might increase as a consequence of increased production followed by

annual population growth and/or increased demand for beef.

Acknowledgements

The authors would like to thank the Norwegian University of Life Sciences and Department of Animal and Aquacultural Sciences for funding the Ph.D. project, the Animalia Meat and Poultry Research Center for data from the Norwegian Beef Cattle Recording System, and Eurofins Food & Agro Testing Norway AS for forage quality data.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCID

Stine Samsonstuen  <http://orcid.org/0000-0001-8309-3457>
Bente A. Åby  <http://orcid.org/0000-0003-0139-5377>
Paul Crosson  <http://orcid.org/0000-0001-7724-8571>
Karen A. Beauchemin  <http://orcid.org/0000-0002-5070-4554>

References

- Åby, B. A., Aass, L., Sehested, E. & Vangen, O. (2012). A bio-economic model for calculating economic values of traits for intensive and extensive beef cattle breeds. *Livestock Science*, doi:10.1016/j.livsci.2011.10.003
- Åby, B. A., Kantanen, J., Aass, L. & Meuwissen, T. (2014). Current status of livestock production in the Nordic countries and future challenges with a changing climate and human population growth. *Acta Agriculturae Scandinavica, Section A — Animal Science*, doi:10.1080/09064702.2014.950321
- Alemu, A. W., Amiro, B. D., Bittman, S., MacDonald, D. & Ominski, K. H. (2017). Greenhouse gas emission of Canadian cow-calf operations: A whole-farm assessment of 295 farms. *Agricultural Systems*, doi:10.1016/j.agsy.2016.11.013
- Animalia. (2017). Slaughter statistics, available at: <https://www.animalia.no/contentassets/3dce35cde68a47b091097fa8c6c2dd5/animalia-kjottetets-tilstand-2017.pdf>.
- Animalia. (2018). Norwegian Beef Cattle Recording System – Annual Report 2017. Norwegian Meat and Poultry Research Centre, Oslo, available at: <https://www.animalia.no/globalassets/storfekjottkontrollen-dokumenter/178153-animalia-arsmelding-storfe-web.pdf>.
- Animalia. (2019). Norwegian Beef Cattle Recording System 2018, Available at: <https://www.animalia.no/no/Dyr/husdyrkontrollene/storfekjottkontrollen/>.
- Arthur, P. F. & Herd, R. M. (2005). Efficiency of feed utilisation by livestock — implications and benefits of genetic improvement. *Canadian Journal of Animal Science*, 85, 281–290.
- Azzam, S. M., Kinder, J. E., Nielsen, M. K., Werth, L. A., Gregory, K. E., Cundiff, L. V. & Koch, R. M. (1993). Environmental effects on neonatal mortality of beef calves. *Journal of Animal Science*, doi:10.2527/1993.712282x
- Bayat, A. R., Kairenius, P., Stefański, T., Leskinen, H., Comtet-Marre, S., Forano, E., Chaucheyras-Durand, F. & Shingfield, K. J. (2015). Effect of camelina oil or live yeasts (*Saccharomyces cerevisiae*) on ruminal methane production, rumen fermentation, and milk fatty acid composition in lactating cows fed grass silage diets. *Journal of Dairy Science*, doi:10.3168/jds.2014-7976
- Bayat, A. R., Tapio, I., Vilkki, J., Shingfield, K. J. & Leskinen, H. (2017). Plant oil supplements reduce methane emissions and improve milk fatty acid composition in dairy cows fed grass silage-based diets without affecting milk yield. *Journal of Dairy Science*, doi:10.3168/jds.2017-13545
- Beauchemin, K. A., Janzen, H. H., Little, S. M., McAllister, T. A. & McGinn, S. M. (2011). Mitigation of greenhouse gas emissions from beef production in western Canada – evaluation using farm-based life cycle assessment. *Animal Feed Science and Technology*, doi:10.1016/j.anifeedsci.2011.04.047
- Bell, M. J., Wall, E., Russell, G., Simm, G. & Stott, A. W. (2011). The effect of improving cow productivity, fertility, and longevity on the global warming potential of dairy systems. *Journal of Dairy Science*, doi:10.3168/jds.2010-4023
- Bonesmo, H., Beauchemin, K. A., Harstad, O. M. & Skjelvåg, A. O. (2013). Greenhouse gas emission intensities of grass silage based dairy and beef production: A systems analysis of Norwegian farms. *Livestock Science*, doi:10.1016/j.livsci.2012.12.016
- Casey, J. W. & Holden, N. M. (2006). Quantification of GHG emissions from sucker-beef production in Ireland. *Agricultural Systems*, doi:10.1016/j.agsy.2005.11.008
- Crosson, P., Foley, P. A., Shalloo, L., O'Brien, D. & Kenny, D. A. (2010). Greenhouse gas emissions from Irish beef and dairy production systems. *Advances in Animal Biosciences*, doi:10.1017/s2040470010004930
- Desjardins, R. L., Worth, D. E., Vergé, X. P. C., Maxime, D., Dyer, J. & Cerkowniak, D. (2012). Carbon footprint of beef cattle. *Sustainability*, doi:10.3390/su4123279
- Foley, P. A., Crosson, P., Lovett, D. K., Boland, T. M., O'Mara, F. P. & Kenny, D. A. (2011). Whole-farm systems modelling of greenhouse gas emissions from pastoral suckler beef cow production systems. *Agriculture Ecosystems & Environment*, doi:10.1016/j.agee.2011.05.010
- Garnsworthy, P. C. (2004). The environmental impact of fertility in dairy cows: A modelling approach to predict methane and ammonia emissions. *Animal Feed Science and Technology*, doi:10.1016/j.anifeedsci.2003.10.011
- Gerber, P. J., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Falucci, A. & Tempio, G. (2013). Tackling Climate Change Through Livestock: A global Assessment of Emissions and Mitigation Opportunities, Most. doi:10.1016/j.anifeedsci.2011.04.074
- Herd, R. M., Arthur, P. F., Donoghue, K. A., Bird, S. H., Bird-Gardiner, T. & Hegarty, R. S. (2014). Measures of methane production and their phenotypic relationships with dry matter intake, growth, and body composition traits in beef cattle. *Journal of Animal Science*, doi:10.2527/jas.2014-8273
- Hristov, A. N., Oh, J., Lee, C., Meinen, R., Montes, F., Ott, T., Firkins, J., Rotz, A., Dell, C., Adesogan, A., Yang, W., Tricarico, J., Kebreab, E., Waghorn, G., Dijkstra, J. & Oosting, S. (2013). Mitigation of greenhouse gas emissions in livestock production – A review of technical options for non-CO2 emissions. P. J. Gerber, B. Henderson and H. P. S. Makkar (eds.). FAO Animal Production and Health Paper No. 177. FAO, Rome, Italy.
- Hyslop, P.P. (2008). Simulated global warming potential and ammonia emission figures for a range of suckler herd breed-

- ing strategies and beef cattle finishing systems, in: livestock and global climate change. Proceedings of a BSAS Conference, May 17–20 2008, Hammamet, Tunisia, p 127.
- IPCC (2006). Guidelines for National Greenhouse Gas Inventories. Agric. For. other L. use. doi:10.1016/j.phrs.2011.03.002
- IPCC (2019). Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P. and Federici, S. (eds). (Switzerland: IPCC).
- Johnson, K. A. & Johnson, D. E. (1995). Methane emissions from cattle. *Journal of Animal Science*, doi:10.2527/1995.7382483x
- Kolling, G. J., Stivanin, S. C. B., Gabbi, A. M., Machado, F. S., Ferreira, A. L., Campos, M. M., Tomich, T. R., Cunha, C. S., Dill, S. W., Pereira, L. G. R. & Fischer, V. (2018). Performance and methane emissions in dairy cows fed oregano and green tea extracts as feed additives. *Journal of Dairy Science*, doi:10.3168/jds.2017-13841
- Legesse, G., Beauchemin, K. A., Ominski, K. H., McGeough, E. J., Kroebel, R., MacDonald, D., Little, S. M. & McAllister, T. A. (2016). Greenhouse gas emissions of Canadian beef production in 1981 as compared with 2011. In *Animal Production Science*. doi:10.1071/AN15386
- Legesse, G., Kroebel, R., Alemu, A. W., Ominski, K. H., McGeough, E. J., Beauchemin, K. A., Chai, L., Bittman, S. & McAllister, T. A. (2018). Effect of changes in management practices and animal performance on ammonia emissions from Canadian beef production in 1981 as compared with 2011. *Canadian Journal of Animal Science*, doi:10.1139/cjas-2017-0184
- Lovett, D. K., Stack, L. J., Lovell, S., Callan, J., Flynn, B., Hawkins, M. & O'Mara, F. P. (2010). Manipulating enteric methane emissions and animal performance of late-lactation dairy cows through concentrate supplementation at pasture. *Journal of Dairy Science*, doi:10.3168/jds.s0022-0302(05)72964-7
- Mogensen, L., Kristensen, T., Nielsen, N. I., Spleth, P., Henriksson, M., Swensson, C., Hesse, A. & Vestergaard, M. (2015). Greenhouse gas emissions from beef production systems in Denmark and Sweden. *Livestock Science*, doi:10.1016/j.livsci.2015.01.021
- Murphy, B., Crosson, P., Kelly, A. K. & Prendiville, R. (2017). An economic and greenhouse gas emissions evaluation of pasture-based dairy calf-to-beef production systems. *Agricultural Systems*, doi:10.1016/j.agsy.2017.03.007
- Murray, C. F., Fick, L. J., Pajor, E. A., Barkema, H. W., Jelinski, M. D. & Windeyer, M. C. (2015). Calf management practices and associations with herd-level morbidity and mortality on beef cow-calf operations. *Animal*, doi:10.1017/S1751731115002062
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T. & Zhang, H. (2013). IPCC 2013 AR5 – Chapter8: Anthropogenic and Natural Radiative Forcing. *Clim. Chang. 2013 Phys. Sci. Basis. Contrib. Work. Gr. I to Fifth Assess. Rep. Intergov. Panel Clim. Chang.* doi:10.1017/CBO9781107415324.018
- Navajas, E. A., Duthie, C. A., Amer, P. R., Sawalha, R. M., Roehe, R. & Roughsedge, T. (2010). Impact of Genomic selection for Residual feed intake and calf survival in beef cattle on profit and greenhouse Gas mitigation, In *Proceedings of the 9th World Congress of Genetics Applied to Livestock Production*.
- Nguyen, T. L. T., Hermansen, J. E. & Mogensen, L. (2010). Environmental consequences of different beef production systems in the EU. *Journal of Cleaner Production*, doi:10.1016/j.jclepro.2009.12.023
- NIBIO (2015). *Account Results in Agriculture and Forestry 2015* (Ås: Norsk institutt for bioøkonomi). 247 pp, available at: <https://www.nibio.no/tema/landbruksokonomi/driftsgranski-nyar-i-jordbruket>
- NIBIO (2016). Totalkalkylen for jordbruket. Jordbrukets totalregnskap 2014 og 2015. Budsjett 2016. In *Budsjettmemda for jordbruket* (Ed.).(in Norwegian). available at: <https://www.nibio.no/tema/landbruksokonomi/totalkalkylen>.
- Nortura (2019). Grunnlagsdokument storfe, sau/lam og egg – 2. halvår 2019.
- Norwegian Farmers Union, Norwegian Farmers and Smallholders Union, Norwegian Government (2019). Intensjonsavtale mellom jordbruket og regjeringen om reduserte klimagassutslipp og økt opptak av karbon fra jordbruket for perioden 2021-2030, available at: <https://www.bondelaget.no/getfile.php/13915420-1561096794/MMA/Bilder%20NB/Mat/Mat-%20og%20landbrukspolitikk/Milj%C3%B8%2C%20energi%20og%20klima/Klimaforhandlinger/Forhandlingsdokument%206.2/avtale%20210619.pdf> (in Norwegian).
- OECD/FAO (2018). OECD-FAO Agricultural Outlook 2018-2027, Agricultural Outlook 2016-2025. doi:10.1787/agr_outlook-2018-en
- Öko-Institut (2010). E.V., Tankstelle/Diesel-DE-2010, available at: <http://www.probas.umweltbundesamt.de/php/volltextsuche.php?&prozessid=%7b9F010C0D-A18D-4163-B86D-22E8656276F7%7d&id=1&step=1&search=Tankstelle/Diesel-DE&b=1#in-output>. (in German).
- Ominski, K. H., Boadi, D. A. & Wittenberg, K. M. (2011). Enteric methane emissions from backgrounded cattle consuming all-forage diets. *Canadian Journal of Animal Science*, doi:10.4141/a05-051
- Randby, ÅT, Nørgaard, P. & Weisbjerg, M. R. (2010). Effect of increasing plant maturity in timothy-dominated grass silage on the performance of growing/finishing Norwegian Red bulls. *Grass and Forage Science* 65, 273–286. doi:10.1111/j.1365-2494.2010.00745.x
- Romero-Perez, A., Okine, E. K., McGinn, S. M., Guan, L. L., Oba, M., Duval, S. M., Kindermann, M. & Beauchemin, K. A. (2014). The potential of 3-nitrooxypropanol to lower enteric methane emissions from beef cattle. *Journal of Animal Science*, doi:10.2527/jas.2014-7573
- Samsonstuen, S., Åby, B. A., Crosson, P., Beauchemin, K. A., Bonesmo, H. & Aass, L. (2019). Farm scale modelling of greenhouse gas emissions from semi-intensive suckler cow beef production. *Agricultural Systems* 176, doi:10.1016/J.AGSY.2019.102670
- Skjelvåg, A. O., Arnoldussen, A. H., Klakegg, O. & Tveito, O. E. (2012). Farm specific natural resource base data for estimating greenhouse gas emissions. *Acta Agriculturae Scandinavica, Section A — Animal Science*, doi:10.1080/09064702.2013.777092
- Soussana, J. F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E., Clifton-Brown, J., Czobel, S., Domingues, R., Flechard, C., Fuhrer, J., Hensen, A., Horvath, L., Jones, M., Kasper, G., Martin, C., Nagy, Z., Neftel, A.,

- Raschi, A., Baronti, S., Rees, R. M., Skiba, U., Stefani, P., Manca, G., Sutton, M., Tuba, Z. & Valentini, R. (2007). Full accounting of the greenhouse gas (CO₂, N₂O, CH₄) budget of nine European grassland sites. *Agriculture Ecosystems & Environment*, doi:10.1016/j.agee.2006.12.022
- Stanley, P. L., Rowntree, J. E., Beede, D. K., DeLonge, M. S. & Hamm, M. W. (2018). Impacts of soil carbon sequestration on life cycle greenhouse gas emissions in Midwestern USA beef finishing systems. *Agricultural Systems* doi:10.1016/j.agsy.2018.02.003
- Statistics Norway (2019). Livestock husbandry. Accessed 15 October, 2019, available at: <http://www.ssb.no/en/jord-skog-jaktog-fiskeri/statistikker/jordhus>: Statistics Norway.
- Thornton, P. K. & Herrero, M. (2010). Potential for reduced methane and carbon dioxide emissions from livestock and pasture management in the tropics. *Proceedings of the National Academy of Sciences of the United States of America*, doi:10.1073/pnas.0912890107
- Veysset, P., Lherm, M., Bébin, D., Roulenc, M. & Benoit, M. (2014). Variability in greenhouse gas emissions, fossil energy consumption and farm economics in suckler beef production in 59 French farms. *Agriculture Ecosystems & Environment*, doi:10.1016/j.agee.2014.03.003
- Vyas, D., McGinn, S. M., Duval, S. M., Kindermann, M. & Beauchemin, K. A. (2016). Effects of sustained reduction of enteric methane emissions with dietary supplementation of 3-nitrooxypropanol on growth performance of growing and finishing beef cattle. *Journal of Animal Science*, doi:10.2527/jas.2015-0268
- Vyas, D., Alemu, A. W., McGinn, S. M., Duval, S. M., Kindermann, M. & Beauchemin, K. A. (2018). The combined effects of supplementing monensin and 3-nitrooxypropanol on methane emissions, growth rate, and feed conversion efficiency in beef cattle fed high-forage and high-grain diets. *Journal of Animal Science*, doi:10.1093/jas/sky174
- Wall, E., Simm, G. & Moran, D. (2010). Developing breeding schemes to assist mitigation of greenhouse gas emissions. *Animal*. doi:10.1017/s175173110999070x
- White, T. A., Snow, V. O. & King, W. M. G. (2010). Intensification of New Zealand beef farming systems. *Agricultural Systems*, doi:10.1016/j.agsy.2009.08.003