

RESEARCH PAPER

 OPEN ACCESS



COVID-19 coronavirus vaccine T cell epitope prediction analysis based on distributions of HLA class I loci (HLA-A, -B, -C) across global populations

Yina Cun^{a*}, Chuanyin Li^{a*}, Lei Shi^a, Ming Sun^a, Shuying Dai^b, Le Sun^b, Li Shi^{id}^a, and Yufeng Yao^a

^aInstitute of Medical Biology, Chinese Academy of Medical Sciences & Peking Union Medical College, Yunnan Key Laboratory of Vaccine Research & Development on Severe Infectious Diseases, Yunnan Engineering Research Centre of Vaccine Research & Development on Severe Infectious Diseases, Kunming, China; ^bSchool of Basic Medical Science, Kunming Medical University, Kunming, China

ABSTRACT

T cell immunity, such as CD4 and/or CD8 T cell responses, plays a vital role in controlling the virus infection and pathological damage. Several studies have reported SARS-CoV-2 proteins could serve as ideal vaccine candidates against SARS-CoV-2 infection by activating the T cell responses. In the current study, based on the SARS-CoV-2 sequence and distribution of host human leukocyte antigen (HLA), we predicted the possible epitopes for the vaccine against SARS-CoV-2 infections. Firstly, the current study retrieved the SARS-CoV-2 S and N protein sequences from the NCBI Database. Then, using the Immune Epitope Database Analysis Resource, we predicted the CTL epitopes of the SARS-CoV-2 S and N proteins according to worldwide frequency distributions of HLA-A, -B, and -C alleles (>1%). Our results predicted 90 and 106 epitopes of N and S proteins, respectively. Epitope cluster analysis showed 16 and 34 respective clusters of SARS-CoV-2 N and S proteins, which covered 95.91% and 96.14% of the global population, respectively. After epitope conservancy analysis, 8 N protein epitopes and 6 S protein epitopes showed conservancy within two SARS-CoV-2 types. Of these 14 epitopes, 13 could cover SARS coronavirus and Bat SARS-like coronavirus. The remaining epitope (KWPWYIWLGF₁₂₁₁₋₁₂₂₀) could cover MERS coronavirus. Finally, the 14-epitope combination could vaccinate 89.60% of all individuals worldwide. Our results propose single or combined CTL epitopes predicted in the current study as candidates for vaccines to effectively control SARS-CoV-2 infection and development.

ARTICLE HISTORY

Received 28 April 2020
Revised 23 August 2020
Accepted 9 September 2020

KEYWORDS

SARS-CoV-2; Human Leukocyte Antigen (HLA); T cell epitopes; population coverage; vaccine

Introduction

Beginning December 2019, a cluster of acute respiratory disease, known as novel coronavirus-infected pneumonia (COVID-19), occurred in Wuhan, Hubei Province, China.¹⁻³

In January 2020, SARS-CoV-2 was identified and confirmed as the cause of COVID-19.⁴ Then, the full genome sequences of SARS-CoV-2 (NC_045512.2) were published in the National Center for Biotechnology Information (NCBI) website. The full-genome sequence and phylogenetic analysis indicated that SARS-CoV-2 forms a clade that is distinct from the beta coronaviruses associated with human severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS).^{3,4} Phylogenetic analysis also revealed that the gene sequence of SARS-CoV-2 is about 88% identical to that of bat SARS-like coronavirus ZXC21 (bat-SL-CoVZXC21, accession no. MG772934.1) and ZC45 (MG772933.1),⁵ which indicated SARS-CoV-2 has closer homology to bat SARS-like coronavirus and bats could be the primary source of SARS-CoV-2.^{6,7}

The antigenicity of coronaviruses, such as SARS-CoV, seems to be largely dependent upon two viral proteins that comprise the nucleocapsid protein (N) and the spike protein (S). The N protein, which is a nucleocapsid phosphoprotein, has been demonstrated to be highly immunogenic and seems to be an important component of the humoral response to SARS-CoV.^{8,9}

The S protein, which is a surface glycoprotein, is a large type-I transmembrane glycoprotein that is not only responsible for receptor binding and membrane fusion, but also serves as a potent immunogen that induces neutralizing antibodies.^{8,10} Whole-sequence alignment showed about 79% sequence identity between SARS-CoV-2 and SARS-CoV.⁵ Thus, the SARS-CoV-2 S and N proteins could serve as target antigens for immune intervention protocols against coronaviruses.

T cell immunity, such as CD4 and/or CD8 T cell responses, plays a vital role in controlling the SARS-CoV infection and/or pathological damage after infection with MERS-CoV.^{11,12} T cell responses have been shown to provide long-term protection.¹³⁻¹⁵ Several studies reported that T cells had the strongest immunogenicity to structural proteins in peripheral blood mononuclear cells of convalescent SARS-CoV patients.^{16,17} Based on the high avidity of cytotoxic T lymphocytes (CTLs), CTLs are considered the major eradicators of viral infections through adaptive immune response. Several studies showed that the presented CTL epitopes of S and N proteins in the context of human leukocyte antigen (HLA) alleles will aid in characterizing the virus control mechanisms and immunopathology of SARS-CoV infection and could provide a new strategy to develop an epitope-based vaccine for SARS.¹⁸⁻²² For example, in 2007 and 2008, Cheung et al. predicted the SARS N protein peptide

CONTACT Li Shi  shili.imb@gmail.com; Yufeng Yao  yufeng_yao@imbcams.com.cn  Institute of Medical Biology, Chinese Academy of Medical Sciences & Peking Union Medical College, Kunming 650118, China

*These authors contributed equally to this work.

© 2020 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

sequence for human MHC class I binding and screened for potential CTL epitopes to control the SARS-associated coronavirus infection *in vitro* and *in vivo*.^{18,23} In 2020, Ibrahim and Kafi used a computational approach for vaccine design to search for candidate epitopes to control MERS-CoV infections.²⁴ But the epitope vaccine method is only in the research stage, and there is no relevant clinical trial report for coronavirus. However, for other viruses, the peptides vaccines have entered clinical trial, such as HPV peptide vaccines.²⁵

When a virus infects cells, the viral antigens are presented to the host immune system through the antigen processing machine (APM). The APM is composed of a proteasome (where the antigens are degraded into peptides), transporters associated with antigen presentation (TAPs, which are responsible for translocating peptide precursors), endoplasmic reticulum aminopeptidases (ERAPs, which trim peptides to fit HLA molecules), and the major histocompatibility complex (HLA, which presents antigen peptides to the cell surface).^{26,27} The CTL epitopes bind to the cleft of various HLA-I molecules through features embedded in the peptide sequence and, more specifically, in anchor residues of HLA-I molecules.²⁸ Then, the HLA-I antigen processing system plays important roles in eliminating the infected cells. However, epitope-based vaccines are limited by HLA specificity, as HLA molecules are highly polymorphic. Therefore, it may be difficult to produce an epitope-based vaccine that is effective in patients with different HLA molecules, thus making it impractical for large-scale vaccination programs.²⁸

In the current study, based on the distribution characteristics of HLA alleles across all populations, we predicted putative CTL epitopes of the novel coronavirus (SARS-CoV-2) N and S proteins using immunoinformatic methods, which combine predictors of proteasomal processing, TAP transport, and MHC binding to produce an overall score indicating the intrinsic potential of each peptide as a T cell epitope. Our results provide likely candidate CTL epitopes or combinations thereof for vaccine development to effectively control SARS-CoV-2.

Methods

Sequence retrieval of SARS-CoV-2 N and S proteins

The ID of SARS-CoV-2 (NC_045512.2) was retrieved from NCBI (https://www.ncbi.nlm.nih.gov/nuccore/NC_045512.2).

Epitope prediction

T cell CTL epitopes of the SARS-CoV-2 N and S proteins were predicted using the Proteasomal cleavage/TAP transport/MHC class I combined predictor, which combines predictors of proteasomal processing, TAP transport, and MHC binding to produce a total score indicating the intrinsic potential of each peptide as a T cell epitope. Based on HLA allele data from the Immune Epitope Database (IEDB) Analysis Resource (<http://tools.iedb.org/processing/>), only HLA alleles occurring in at least 1% of the human population or with an allele frequency of 1% or more were selected. This prediction method is recommended by IEDB. The total score combines the proteasomal cleavage, TAP transport, and MHC binding predictions, which

predicts a quantity proportional to the amount of peptide presented by MHC molecules on the cell surface. As the prediction method recommended, high score equal to the high efficiency of the epitope presented by MHC molecules. The total score was used as the cut off value for epitope selection in the present study.

Epitope cluster analysis

Based on sequence identity, the predicted epitopes were grouped into clusters using Epitope Cluster Analysis (<http://tools.iedb.org/cluster/>). A cluster is defined as a group of sequences having a sequence similarity greater than the specified minimum sequence identity threshold. An identity set percentage means that any member of the cluster will be at least the set percentage identical to at least one member of the cluster.²⁹ In the current study, a cluster was defined as a group of sequences that have a sequence similarity >70% minimum sequence identity threshold, which is the default of the cluster analysis (<http://tools.iedb.org/cluster/help/>)

Population coverage

The population coverage method calculates the fraction of individuals predicted to respond to a given set of epitopes with known MHC restrictions (<http://tools.iedb.org/population/>). This calculation is based on HLA genotypic frequencies assuming non-linkage disequilibrium between HLA loci. According to the results, we selected >95% population coverage as the threshold value.

The blast analysis of the 13 coronavirus N and S proteins

Blast method was used to analyze identity proportion of the N and S protein sequences between NC_045512.2 (Wuhan seafood market pneumonia virus isolate Wuhan-Hu-1) and other 12 coronaviruses which are MT007544.1 (Severe acute respiratory syndrome coronavirus 2 isolate Australia/VIC01/2020), FJ882944.1 (SARS coronavirus ExoN1 isolate P3pp23), NC_004718.3 (SARS coronavirus Tor2), MG987420.1 (Middle East respiratory syndrome-related coronavirus isolate NL13892) and MG021451.1 (Middle East respiratory syndrome-related coronavirus isolate NL13845), NC_006213.1 (Human coronavirus OC43 strain ATCC VR-759), NC_006577.2 (Human coronavirus HKU1), KY983587.1 (Human coronavirus 229E strain HCoV_229E/Seattle/USA/SC3112/2015), NC_005831.2 (Human Coronavirus NL63) which contain all 7 known human coronaviruses, and MG772934.1 (Bat SARS-like coronavirus isolate bat-SL-CoVZXC21), KY417144.1 (Bat SARS-like coronavirus isolate Rs4084), KT444582.1 (SARS-like coronavirus WIV16) which contain none human coronaviruses.

Epitope conservancy analysis

Epitope conservancy analysis was used to calculate the degree of conservancy of an epitope within N and S protein sequence of 13 coronaviruses set at different degrees of sequence identity (<http://tools.iedb.org/conservancy/>). The degree of conservancy is defined as the fraction of protein sequences containing

the epitope at a given identity level that the selected epitopes completely matched at least two human coronaviruses above.

Results

The flow chart of epitope prediction analysis and results is illustrated in Figure 1.

HLA allele analysis

The average frequency of HLA alleles (>1%) across all population samples was selected in the current study (<http://tools.iedb.org/processing/help/>). In total, 70 HLA alleles were selected for analysis according to the IEDB Analysis Resource (Table 1). There were 18 HLA-A alleles, 32 HLA-B alleles, and 20 HLA-C alleles.




Figure 1. The flow chart of epitope prediction analysis and results.

Table 1. List of 70 HLA alleles (18 HLA-A alleles, 32 HLA-B alleles, 20 HLA-C alleles) selected occurring in at least 1% of the human population.

HLA Loci	Alleles
HLA-A	A*02:01, A*01:01, A*02:06, A*03:01, A*11:01, A*23:01, A*24:02, A*25:01, A*26:01, A*29:02, A*30:01, A*30:02, A*31:01, A*32:01, A*33:03, A*68:01, A*74:01, A*68:02
HLA-B	B*07:02, B*08:01, B*13:01, B*13:02, B*14:02, B*15:01, B*15:02, B*15:25, B*18:01, B*27:02, B*27:05, B*35:01, B*35:03, B*37:01, B*38:01, B*39:01, B*40:01, B*40:02, B*44:02, B*44:03, B*46:01, B*48:01, B*49:01, B*50:01, B*51:01, B*52:01, B*53:01, B*55:01, B*56:01, B*57:01, B*58:01, B*58:02
HLA-C	C*01:02, C*02:02, C*02:09, C*03:02, C*03:03, C*03:04, C*04:01, C*05:01, C*06:02, C*07:01, C*07:02, C*07:04, C*08:01, C*08:02, C*12:02, C*12:03, C*14:02, C*15:02, C*16:01, C*17:01

Epitope prediction

A total of 200,410 epitopes of SARS-CoV-2 N protein were predicted against the 70 alleles of the HLA-A, -B, and -C loci. We selected 0.60, 0.55, 0.50, 0.47, 0.45, 0.40, and 0.30 as the cut off values of total scores for epitope selection, and 71, 82, 90, 99, 103, 110, and 130 epitopes were found within these cut offs. According to the epitope cluster analysis and population coverage results, we chose 0.50 as the cut off value for epitope selection (90 epitopes), as this value identified few epitopes while reaching more than 95% population coverage (Table 2).

For the SARS-CoV-2 S protein, 618,871 total epitopes were predicted against the 70 alleles of the HLA-A, -B and -C loci. We selected 0.70, 0.80, 0.90, 0.95, 1.00, 1.05, 1.07, 1.10, and 1.15 as cut off values for epitope selection, and 80, 95, 106, 113, 133, 158, 179, 247, and 313 epitopes were found. According to the epitope cluster analysis and population coverage results, 1.07 was chosen as the cut off value for epitope selection (106 epitopes), as this value indicated few epitopes reaching more than 95% population coverage (Table 2).

Epitope cluster analysis

Using the selected epitopes (90 epitopes in N protein and 106 epitopes in S protein), the predicted epitopes were grouped into

Table 2. The epitopes number, cluster numbers, and population coverage results of the predicted SARS-CoV-2 N and S proteins.

Protein	Epitopes Number	Total Score	Population Coverage			Clustering
			coverage ^a	Average hit ^b	PC90 ^c	
N protein	130	0.30	0.9910	8.18	2.44	21
	110	0.40	0.9903	7.25	2.28	17
	103	0.45	0.9903	6.57	2.13	17
	99	0.47	0.9665	5.99	1.64	17
	90	0.50	0.9591	5.57	1.51	16
	82	0.55	0.9458	4.97	1.32	15
	71	0.60	0.9228	4.39	1.14	15
	S protein	313	0.70	0.9901	19.77	4.78
247		0.80	0.9864	16.12	4.33	49
179		0.90	0.9810	12.09	3.31	45
158		0.95	0.9788	10.65	3.23	40
133		1.00	0.9632	8.65	1.60	35
113		1.05	0.9614	7.35	1.56	33
106		1.07	0.9614	7.08	1.56	34
95		1.10	0.9319	6.17	1.28	33
80		1.15	0.9242	5.25	1.17	30

^aprojected population coverage

^baverage number of epitope hits/HLA combinations recognized by the population

^cminimum number of epitope hits/HLA combinations recognized by 90% of the population

16 and 34 clusters, respectively. Among the 16 N protein clusters, cluster 1 contained the most epitopes (11 epitopes), which was represented by HLA-A*30:02 (8/11), A*29:02 (8/11), A*30:01 (1/11), A*31:01 (1/11), B*15:25 (1/11), B*15:01 (1/11), A*03:01 (1/11), B*07:02 (1/11), and B*08:01 (1/11). Clusters 9 to 16 contained only one epitope each (Table 3). Among the 34 S protein clusters, clusters 1 and 2 contained the most epitopes (5 epitopes each). Cluster 1 was represented by HLA-B*15:25 (2/5), B*15:01 (1/5), A*03:01 (1/5), A*23:01 (1/5), A*24:02 (1/5), and A*30:02 (1/5). Cluster 2 included HLA-A*23:01 (2/5), A*29:02 (1/5) and A*02:01 (1/5). Clusters 16 to 34 contained only one epitope each (Table 4).

Population coverage

The HLA allele frequencies and associated data for different individual populations from worldwide studies were provided by the allelefrequencies.net database (<http://www.allelefrequencies.net/>), which covered 10,656,469 individuals from 1,270 populations in 16 different geographical areas including 115 countries and 21 different ethnicity groups.

The worldwide population coverage of the 90 and 106 epitopes of N and S proteins were 95.91% and 96.14%, respectively (Table 2). For the 90 N protein epitopes, the average number of epitope hits/HLA combinations recognized by the population was 5.57, and the minimum number recognized by 90% of the population was 1.51. For the 106 S protein epitopes, the average number of epitope hits/HLA combinations recognized by the population was 7.08, and the minimum number recognized by 90% of the population was 1.56.

The blast analysis of the 13 coronavirus N and S proteins

The BLAST results of N and S proteins between NC_045512.2 and 12 other sequences (MT007544.1, MG772934.1, KY417144.1, KT444582.1, FJ882944.1, NC_004718.3, MG987420.1, MG021451.1, NC_006213.1, NC_006577.2, KY983587.1, and NC_005831.2) selected in the current study are shown in Table 5. The percent identities of other human coronaviruses except SARS-CoV-2 and SARS were relatively low (29.43%~50.97% for N protein; 30.78%~37.63% for S protein).

Epitope conservancy analysis

The identity proportion results of 16 epitopes within 13 N protein sequences above are shown in Table 6. The identity is 80%~100% for 16 epitopes within 7 N protein sequences belong to the same subgenus (Sarbecovirus, including SARS-CoV-2). The identity is 23.53%~80% for 16 epitopes within other 6 N protein sequences belong to the different genus (Alphacoronavirus) or subgenus (Merbecovirus). The identity proportion results of 34 epitopes within 13 S protein sequences above are shown in Table 7. The identity is higher (60%~100%) for NP13 (KRSFIEDLLF₈₁₄₋₈₂₃), NP17 (KWPWYIWLGF₁₂₁₁₋₁₂₂₀) and NP28 (YEQYIKWPWY₁₂₁₄₋₁₂₂₃) within 13 S protein sequences. For the remaining 31 epitopes, the identity is 33.33%~100% for the epitopes within 7 S protein sequences belong to the same subgenus (Sarbecovirus, including SARS-CoV-2); the identity is 28.57%~55.56% for the epitopes within other 6 N protein sequences

Table 3. The epitopes sequences, amino acid position and its presented HLA alleles of SARS-CoV-2 N protein by cluster analysis.

Cluster Number	Epitope Number	Alignment	Epitope	Amino-acid Position	HLA Alleles
1.1	Consensus	DGKMKDLSRWYFYLL	-	98-113	
	1	DGKMKDLSRWYFY-	DGKMKDLSRWYFY	98-111	A*30:02
	2	-GKMKDLSRWYFY-	GKMKDLSRWYFY	99-111	A*30:02, A*29:02
	3	-KMKDLSRWYFY-	KMKDLSRWYFY	100-112	A*30:02, A*29:02
	4	-KMKDLSRWYFY-	KMKDLSRWYFY	100-111	A*30:02, A*29:02, A*30:01, A*31:01
	5	-KMKDLSRWY - -	KMKDLSRWY	100-109	A*30:02, B*15:25, B*15:01
	6	- MKDLSRWYFY-	MKDLSRWYFY	101-111	A*29:02, A*30:02
	7	- -KDLSRWYFY-	KDLSRWYFY	102-111	A*30:02, A*03:01, A*29:02
	8	- -DLSRWYFY-	DLSRWYFY	103-112	A*29:02
	9	- -DLSRWYFY-	DLSRWYFY	103-111	A*29:02
	10	- - LSPRWYFY-	LSPRWYFY	104-112	A*29:02, A*30:02
11	- - -SPRWYFYLL	SPRWYFYLL	105-113	B*07:02, B*08:01	
2.1	Consensus	IAQFAPSASAFF	-	304-315	
	1	IAQFAPSASAF-	IAQFAPSASAF	304-314	B*15:25
	2	-AQFAPSASAFF	AQFAPSASAFF	305-315	B*15:01, B*15:25
	3	-AQFAPSASAF-	AQFAPSASAF	305-314	B*15:25, B*15:01, B*15:02
	4	-QFAPSASAFF	QFAPSASAFF	306-315	A*23:01
	5	-QFAPSASAF-	QFAPSASAF	306-314	C*14:02
6	- FAPSASAFF	FAPSASAFF	307-315	C*12:03, B*35:01, C*03:03, C*03:04, C*03:02, C*14:02, C*16:01	
3.1	Consensus	MSRIGMEVTPSGTWLTY	-	317-333	
	1	MSRIGMEVTPSGTW -	MSRIGMEVTPSGTW	317-330	B*58:01
	2	- -MEVTPSGTWLTY	MEVTPSGTWLTY	322-333	B*35:01, B*18:01, B*44:03
	3	- -MEVTPSGTW -	MEVTPSGTW	322-330	B*44:02
	4	- - -VTPSGTWLTY	VTPSGTWLTY	324-333	A*29:02, B*35:01, A*30:02
5	- - -TPSGTWLTY	TPSGTWLTY	325-333	B*35:01, B*53:01	
4.1	Consensus	NTNSSPDDQIGYY	-	75-87	
	1	NTNSSPDDQIGYY	NTNSSPDDQIGYY	75-87	A*01:01
	2	-NSSPDDQIGYY	NSSPDDQIGYY	77-87	A*01:01
	3	- SSPDDQIGYY	SSPDDQIGYY	78-87	A*01:01
4	- SPDDQIGYY	SPDDQIGYY	79-87	B*35:01	
5.1	Consensus	FYYLGTGPEAGLPY	-	110-123	
	1	FYYLGTGPEAGLPY	FYYLGTGPEAGLPY	110-123	A*29:02, C*14:02
2	-YYLGTGPEAGLPY	YYLGTGPEAGLPY	111-123	A*29:02	
6.1	Consensus	LPQGTTLPKGFY	-	161-172	
	1	LPQGTTLPKGFY	LPQGTTLPKGFY	161-172	B*35:01
2	- GTTLPKGFY	GTTLPKGFY	164-172	A*30:02	
7.1	Consensus	ILLNKHIDAY	-	351-360	
	1	ILLNKHIDAY	ILLNKHIDAY	351-360	B*15:25, B*15:01
2	-LLNKHIDAY	LLNKHIDAY	352-360	B*15:25, B*15:01, B*15:02	
8.1	Consensus	KFPRGQGVPI	-	65-74	
	1	KFPRGQGVPI	KFPRGQGVPI	65-74	B*07:02
2	-FPRGQGVPI	FPRGQGVPI	66-74	B*07:02	
9.1	Singleton	NTASWFTAL	NTASWFTAL	48-56	A*68:02
10.1	Singleton	RQKRTATKAY	RQKRTATKAY	259-268	B*15:01, B*15:25, A*30:02
11.1	Singleton	KAYNVTQAF	KAYNVTQAF	266-274	C*03:02, B*15:25, C*12:03, A*32:01, C*16:01, C*03:03, C*03:04, B*15:01, C*14:02, B*58:01, C*12:02, B*15:02, B*35:01, C*02:09, C*02:02
12.1	Singleton	LPAADLDDF	LPAADLDDF	395-403	B*35:01
13.1	Singleton	LPNNTASWF	LPNNTASWF	45-53	B*35:01, B*53:01
14.1	Singleton	LLLDRLNQL	LLLDRLNQL	222-230	A*02:01, A*02:06
15.1	Singleton	IGYRRATR	IGYRRATR	84-92	A*31:01
16.1	Singleton	NQRNAPRITF	NQRNAPRITF	8-17	B*15:01

belong to different genus (Alphacoronavirus) or subgenus (Merbecovirus).

Among the 16 SARS-CoV-2 N protein epitopes, 7 epitopes showed 53.85% (7/13) match between 13 protein sequences (Table 6). For NP 3 (MSRIGMEVTPSGTWLTY₃₁₇₋₃₃₃), this match was 46.15% (6/13). The above 8 epitopes had 23 alleles (Table 8). HLA-B*35:01 (3/8) was most frequent in the epitopes, followed by B*15:01 (2/8), B*15:02 (2/8), B*15:25 (2/8), A*30:02 (2/8), and B*53:01 (2/8). The remaining 17 HLA alleles were present in only one epitope. No epitope was found with the N protein of other human coronaviruses except SARS-CoV-2 and SARS. Among the 34 SARS-CoV-2 S protein

epitopes, 3 epitopes showed 53.85% (7/13) match between 13 protein sequences (Table 7). Two epitopes had a 46.15% (6/13) match. The NP17 epitope (KWPWYIWLGF₁₂₁₁₋₁₂₂₀) was found in the S protein of two SARS-CoV-2 and two MERS-related coronaviruses. The above 6 epitopes had 13 alleles (Table 8). HLA-A*23:01, B*35:01, B*58:01, C*03:04, and C*03:03 were present in 2 epitopes. The remaining 8 HLA alleles were present in only one epitope.

Based on the epitope conservancy analysis results, we selected epitopes with 30.77%~53.85% match percentage as epitope candidates, which means the selected epitopes completely matched at least two human coronaviruses. Finally, 14 total epitopes (8

Table 4. The epitopes sequences, amino acid position and its presented HLA alleles of SARS-CoV-2 S protein by cluster analysis.

Cluster Number	Epitope Number	Alignment	Epitope	Amino-acid Position	HLA Alleles
1.1	Consensus	YSVLYNSASFSTFKCY	-	365–380	
	1	YSVLYNSASF --	YSVLYNSASF	365–374	B*15:01
	2	-SVLYNSASF --	SVLYNSASF	366–374	B*15:25
	3	-VLYNSASFSTFKCY	VLYNSASFSTFKCY	367–380	A*03:01
	4	-LYNSASFSTF --	LYNSASFSTF	368–377	A*23:01, A*24:02
2.1	Consensus	AYYVGYLQPRTFLLKY	-	264–279	B*15:25, A*30:02
	1	AYYVGYLQPRTF --	AYYVGYLQPRTF	264–275	A*23:01
	2	-YYVGYLQPRTF --	YYVGYLQPRTF	265–275	A*23:01
	3	--GYLQPRTFLLKY	GYLQPRTFLLKY	268–279	A*29:02
	4	--YLQPRTFLLKY	YLQPRTFLLKY	269–279	A*29:02
3.1	Consensus	RISNCVADYSVLY	-	357–369	A*02:01
	1	RISNCVADYSVLY	RISNCVADYSVLY	357–369	A*30:02
	2	RISNCVADY --	RISNCVADY	357–365	A*30:02
	3	-NCVADYSVLY	NCVADYSVLY	360–369	A*29:02
	4	--CVADYSVLY	CVADYSVLY	361–369	A*29:02, B*35:01, A*26:01, A*01:01
4.1	Consensus	LQIPFAMQMAYRF	-	894–906	
	1	LQIPFAMQMAY--	LQIPFAMQMAY	894–904	B*35:01, B*15:25, B*15:01
	2	-QIPFAMQMAY--	QIPFAMQMAY	895–904	B*35:01, A*29:02
	3	-IPFAMQMAY--	IPFAMQMAY	896–904	B*35:01, B*53:01
5.1	Consensus	RVYSSANNCTFEY	-	158–170	B*35:01, C*03:02, B*53:01, C*03:04, C*03:03, B*58:01, A*23:01
	1	RVYSSANNCTFEY	RVYSSANNCTFEY	158–170	A*30:02
	2	-VYSSANNCTF--	VYSSANNCTF	159–168	A*24:02
	3	-YSSANNCTF--	YSSANNCTF	160–168	C*03:02, C*03:04, C*03:03, B*35:01, C*16:01, C*12:03
6.1	Consensus	AYTMSLGAENSVAY	-	694–707	B*35:01, A*29:02
	1	AYTMSLGAENSVAY	AYTMSLGAENSVAY	694–707	A*29:02
	2	-YTMSLGAENSVAY	YTMSLGAENSVAY	695–707	A*29:02, B*15:25
	3	--SLGAENSVAY	SLGAENSVAY	698–707	B*15:25
7.1	Consensus	SWMESEFRVY	-	151–160	B*35:01
	1	SWMESEFRVY	SWMESEFRVY	151–160	A*29:02
	2	-WMESEFRVY	WMESEFRVY	152–160	B*15:25, B*15:02
8.1	Consensus	YTNSFTRGVVY	-	28–38	B*18:01
	1	YTNSFTRGVVY	YTNSFTRGVVY	28–38	A*01:01, A*29:02
	2	YTNSFTRGVY-	YTNSFTRGVY	28–37	A*30:02, A*01:01
9.1	Consensus	HWFVTQRNFY	-	1101–1110	C*12:03, A*29:02
	1	HWFVTQRNFY	HWFVTQRNFY	1101–1110	A*29:02
10.1	Consensus	FQFCNDPFLGVY	-	133–144	A*29:02
	1	FQFCNDPFLGVY	FQFCNDPFLGVY	133–144	B*15:25, B*15:01
11.1	Consensus	SVASQSIAY	-	686–695	A*02:06
	1	SVASQSIAY	SVASQSIAY	686–695	B*15:25
12.1	Consensus	FLPFFSNVTW	-	55–64	B*35:01, B*15:25, C*03:02
	1	FLPFFSNVTW	FLPFFSNVTW	55–64	B*53:01
13.1	Consensus	KRSFIEDLLF	-	814–823	B*53:01
	1	KRSFIEDLLF	KRSFIEDLLF	814–823	B*58:01
14.1	Consensus	LLTDEMIAQY	-	864–873	B*58:01, B*57:01
	1	LLTDEMIAQY	LLTDEMIAQY	864–873	A*01:01
15.1	Consensus	RVYSTGSNVF	-	634–643	A*01:01
	1	RVYSTGSNVF	RVYSTGSNVF	634–643	B*15:25, B*15:01, A*32:01
16.1	Consensus	WTAGAAAYY	-	258–266	C*14:02
	1	WTAGAAAYY	WTAGAAAYY	258–266	A*29:02, A*26:01, A*68:01
17.1	Consensus	KWPWYIWLGF	-	1211–1220	A*23:01, A*24:02
	1	KWPWYIWLGF	KWPWYIWLGF	1211–1220	A*23:01, A*24:02
18.1	Consensus	KSNIIRGWIF	-	97–106	B*58:01
	1	KSNIIRGWIF	KSNIIRGWIF	97–106	B*58:01
19.1	Consensus	CYFPLQSYGF	-	488–497	A*23:01, A*24:02
	1	CYFPLQSYGF	CYFPLQSYGF	488–497	A*23:01, A*24:02
20.1	Consensus	FEYVSQPFL	-	168–176	B*40:01
	1	FEYVSQPFL	FEYVSQPFL	168–176	B*40:01
21.1	Consensus	FVFNKIDGY	-	192–200	B*35:01, A*29:02
	1	FVFNKIDGY	FVFNKIDGY	192–200	B*35:01, A*29:02
22.1	Consensus	LMSFPQSAPHGVVF	-	1057–1070	B*15:01, B*15:25
	1	LMSFPQSAPHGVVF	LMSFPQSAPHGVVF	1057–1070	B*15:01, B*15:25
23.1	Consensus	NATRFASVY	-	151–159	B*35:01
	1	NATRFASVY	NATRFASVY	151–159	B*35:01
24.1	Consensus	KSFTVEKGIY	-	312–321	A*30:02
	1	KSFTVEKGIY	KSFTVEKGIY	312–321	A*30:02
25.1	Consensus	LPFNDGVYF	-	84–92	B*35:01, B*53:01
	1	LPFNDGVYF	LPFNDGVYF	84–92	B*35:01, B*53:01
26.1	Consensus	TLLALHRSY	-	248–256	B*15:25
	1	TLLALHRSY	TLLALHRSY	248–256	B*15:25
27.1	Consensus	NDLCFTNVY	-	396–404	B*18:01
	1	NDLCFTNVY	NDLCFTNVY	396–404	B*18:01
28.1	Consensus	YEQYIKWPWY	-	1214–1223	B*18:01
	1	YEQYIKWPWY	YEQYIKWPWY	1214–1223	B*18:01

(Continued)

Table 4. (Continued).

Cluster Number	Epitope Number	Alignment	Epitope	Amino-acid Position	HLA Alleles
29.1	Singleton	KRFDNPVLPF	KRFDNPVLPF	77–86	B*27:05
30.1	Singleton	FPNITNLCPF	FPNITNLCPF	337–346	B*35:01
31.1	Singleton	NVYADSFVIR	NVYADSFVIR	402–411	A*68:01
32.1	Singleton	QLTPTWRVY	QLTPTWRVY	636–644	B*15:25, B*15:02
33.1	Singleton	KVGGNYNYLY	KVGGNYNYLY	452–461	A*30:02
34.1	Singleton	WTFGAGAAL	WTFGAGAAL	894–902	C*03:04, C*03:03

epitopes for N protein, 6 epitopes for S protein) were selected, which cover 89.60% of the worldwide population (Table 9). The average number of epitope hits/HLA combinations recognized by the population was 3.22, and the minimum number recognized by 90% of the population was 0.96. For SARS-CoV (FJ882944.1) and SARS-like coronavirus (KT444582.1), 13 epitopes (8 clusters for N protein, 5 clusters for S protein) were covered, which occurred 84.94% of the worldwide population. For Bat SARS-like coronavirus (KY417144.1), 12 epitopes (7 clusters for N protein, 5 clusters for S protein) were covered, which occurred 81.91% of the worldwide population. For Bat SARS-like coronavirus (MG772934.1), 11 epitopes (8 clusters for N protein, 3 clusters for S protein) were covered, which occurred 84.94% of the worldwide population. For MERS (MG987420.1 and MG021451.1), only one epitope (S protein) was covered, which occurred 26.18% of the worldwide population (Table 9).

Discussion

Modern immunoinformatic methodologies provide new strategies for the design and synthesis of antigen-specific epitope-based vaccines against viral or pathogenic infections. In the current study, according to the SARS-CoV-2 S and N protein sequences, we predicted putative HLA-restricted CTL epitopes using immunoinformatic methods. We found 14-epitope combinations that have 30.77%–53.85% match percentage among the coronavirus sequences covering SARS-CoV-2, other 6 human coronaviruses and other coronavirus species (NC_045512.2, MT007544.1, MG772934.1, KY417144.1, KT444582.1, FJ882944.1, NC_004718.3, MG987420.1, MG021451.1, NC_006213.1, NC_006577.2, KY983587.1, and NC_005831.2). The worldwide population coverage is 89.60%, which indicates that these epitopes could serve as candidate epitopes for vaccines of SARS-CoV-2 among most of the global population.

Based on the antigenicity of the SARS-CoV-2 S and N proteins, they could be major targets for preventing and treating SARS-CoV-2 infection. In 2007 and 2008, Cheung *et al.* predicted the DNA vaccines encoding the N-protein peptides LLLDRLNQL₂₂₃₋₂₃₁ and LALLLLDRL₂₂₀₋₂₂₈ presented by HLA-A*02:01 could trigger the highest T-cell cytotoxicity toward N protein-expressing cells, which indicated that these two N-protein peptides could be valuable peptide candidates for SARS vaccine.^{18,23} In the current study, we also predicted that LLLDRLNQL₂₂₃₋₂₃₁ could be presented by HLA-A*02:01 and HLA-A*02:06, which covered 39.08% and 1.95% individuals worldwide, respectively. Moreover, the sequence of the epitope is identical to those of SARS coronavirus (KT444582.1 and FJ882944.1) and nucleocapsid protein Bat SARS-like coronavirus (MG772934.1 and KY417144.1), but not MERS (MG987420.1 and MG021451.1), which indicated that

LLLDRLNQL₂₂₃₋₂₃₁ could also be a valuable vaccine candidate peptide for SARS and Bat SARS-like coronavirus. In 2020, Ahmed *et al.* revisited previously tested and functional HLA-restricted SARS coronavirus epitopes and found many epitopes were also conserved in SARS-CoV-2, such as epitope LLLDRLNQL₂₂₃₋₂₃₁ presented by HLA-A*02:01 allele.³⁰ Thus, epitope LLLDRLNQL₂₂₃₋₂₃₁ could potentially offer protection against these two viruses. In addition, in 2020, Austin Nguyen *et al.* analyzed viral peptide-MHC class I binding affinity across HLA-A, -B, and -C genotypes for all SARS-CoV-2 peptides in silico, and found that different HLA alleles showed various capacities to present highly conserved SARS-CoV-2 peptides.³¹ Among the conserved sequences, PRWYFYLTGP₁₀₆₋₁₁₇ in N protein was highly conserved. In the current study, we also predicted that FYYLGTGPEAGLPY₁₁₀₋₁₂₃ was presented by HLA-A*29:02 and HLA-C*14:02 alleles. In 2006, Zhou *et al.* predicted the SARS S protein epitopes and identified KLPDDFMGCV₄₁₁₋₄₂₀ as a novel HLA-A*02:01-restricted S protein epitope.¹¹ Their results indicated that the epitope could be a novel SARS-associated coronavirus-specific CTL epitope and a potential target for characterizing virus control mechanisms and evaluating candidate SARS vaccines. Then, Liu *et al.* also predicted a SARS-CoV N protein-derived CTL epitope and identified QFKDENVILL₃₄₆₋₃₅₄, which was restricted by HLA-A*24:02, by a series of in vitro studies.¹⁹ In the current study, we did not predict these two epitopes in the SARS-CoV-2 N and S proteins. After aligning SARS-CoV-2 and SARS-CoV, we found only one mutation in the S protein epitope KLPDDFMGCV₄₁₁₋₄₂₀ and two different mutations in QFKDENVILL₃₄₆₋₃₅₄ between SARS-CoV-2 and SARS-CoV. Then, we replaced these mutations in SARS-CoV-2 and re-conducted the predictions; KLPDDFMGCV₄₁₁₋₄₂₀ and QFKDENVILL₃₄₆₋₃₅₄ could be predicted, which indicated that these mutations could change the affinity between epitopes and HLA molecules.

Evasion of the host CTL response through the mutation of key epitopes is a major challenge to achieving natural or therapeutic vaccine-induced immune control of SARS-CoV-2. Therefore, we used two SARS-CoV-2 sequences (NC_045512.2 and MT007544.1) retrieved from NCBI to predict the epitopes and found that 14 epitopes shared identity with these two SARS-CoV-2 sequences, which include two major types of SARS-CoV-2 (Designated L and S). Thus, the 14-epitope combination could be feasible in vaccines for SARS-CoV-2, including SARS-CoV-2 types L and S.

Moreover, 13 of the 14 epitopes we selected in the current study could also serve as candidate epitopes for the SARS coronavirus and Bat SARS-like coronavirus (MG772934.1, KT444582.1, KT444582.1, and FJ882944.1) due to identical sequences between the epitopes and coronaviruses; additionally, these 13 epitopes

Table 5. The results of comparing N and S protein of SARS-CoV-2 with Bat coronavirus and other human coronaviruses (SARS, MERS, HCoV-OC43, HCoV-HKU1, HCoV-229E and HCoV-NL63).

Serial number	complete genome ^a	Genus	Subgenus	Description	Max Score	Total Score	Query Cover ^b	E value ^c	Per. Ident ^d	Accession
1	NC_045512.2	Betacoronavirus	Sarbecovirus	nucleocapsid phosphoprotein	854	854	100%	0	100.00%	YP_009724397.2
				surface glycoprotein	2637	2637	100%	0	100.00%	YP_009724390.1
2	MT007544.1	Betacoronavirus	Sarbecovirus	nucleocapsid phosphoprotein	854	854	100%	0	100.00%	QHR84456.1
				surface glycoprotein	2634	2634	100%	0	99.92%	QHR84449.1
3	MG772934.1	Betacoronavirus	Sarbecovirus	nucleocapsid protein	715	715	100%	0	94.27%	AVP78049.1
				spike protein	2105	2105	99%	0	80.32%	AVP78042.1
4	KT444582.1	Betacoronavirus	Sarbecovirus	nucleocapsid protein	669	669	100%	0	90.28%	ALK02467.1
				spike protein	2065	2065	100%	0	77.07%	ALK02457.1
5	KY417144.1	Betacoronavirus	Sarbecovirus	nucleocapsid protein	666	666	100%	0	90.05%	ATO98142.1
				spike protein	2049	2049	99%	0	77.23%	ATO98132.1
6	FJ882944.1	Betacoronavirus	Sarbecovirus	nucleocapsid protein	672	672	100%	0	90.52%	ACZ72030.1
				spike glycoprotein precursor	2040	2040	100%	0	76.12%	ACZ72020.1
7	NC_004718.3	Betacoronavirus	Sarbecovirus	nucleocapsid protein	672	672	100%	0	90.52%	YP_009825061.1
				spike glycoprotein precursor	2038	2038	100%	0	75.96%	YP_009825051.1
8	MG987420.1	Betacoronavirus	Merbecovirus	N protein	288	288	88%	6E-90	50.26%	AWH65950.1
				S protein	561	636	85%	0	35.56%	AWH65943.1
9	MG021451.1	Betacoronavirus	Merbecovirus	N protein	284	284	82%	3E-88	50.97%	AVV62533.1
				S protein	556	618	83%	1E-179	34.70%	AVV62526.1
10	NC_006213.1	Betacoronavirus	Embecovirus	nucleocapsid protein	176	207	74%	6.00E-55	38.35%	YP_009555245.1
				spike surface glycoprotein	467	602	74%	2.00E-146	37.63%	YP_009555241.1
11	NC_006577.2	Betacoronavirus	Embecovirus	nucleocapsid phosphoprotein	197	212	82%	9.00E-63	36.74%	YP_173242.1
				spike glycoprotein	452	576	71%	3.00E-141	35.43%	YP_173238.1
12	KY983587.1	Alphacoronavirus	Duvinacovirus	nucleocapsid protein	89.4	89.4	65%	6.00E-24	29.43%	ARU07605.1
				spike protein	366	481	67%	2.00E-111	31.32%	ARU07601.1
13	NC_005831.2	Alphacoronavirus	Setracovirus	nucleocapsid protein	61.2	61.2	13%	9.00E-15	48.28%	YP_003771.1
				Spike protein	349	404	65%	3.00E-104	30.78%	YP_003767.1

^aNC_045512.2 is Wuhan seafood market pneumonia virus isolate Wuhan-Hu-1; MT007544.1 is Severe acute respiratory syndrome coronavirus 2 isolate Australia/VIC01/2020; MG772934.1 is Bat SARS-like coronavirus isolate bat-SL-CoVZXC21; KY417144.1 is Bat SARS-like coronavirus isolate Rs4084; KT444582.1 is SARS-like coronavirus WIV16; FJ882944.1 is SARS coronavirus ExoN1 isolate P3pp23; NC_004718.3 SARS coronavirus Tor2; MG987420.1 is Middle East respiratory syndrome-related coronavirus isolate NL13892; MG021451.1 is Middle East respiratory syndrome-related coronavirus isolate NL13845; NC_006213.1 Human coronavirus OC43 strain ATCC VR-759; NC_006577.2 Human coronavirus HKU1; KY983587.1 Human coronavirus 229E strain HCoV_229E/Seattle/USA/SC3112/2015; NC_005831.2 Human Coronavirus NL63;

^bQuery Coverage: coverage of the compared sequences

^cexpect value: the possibility of random matching. When the value of E is close to zero or zero, it is essentially a perfect match.

^dpercentage identity: the percentage of base number in the total sequence of the compared sequences

could cover 84.94% of individuals worldwide. However, the 13 combination epitopes did not cover MERS (MG987420.1 and MG021451.1) due to the different subgenus (MERS belongs to Merbecovirus and SARS-CoV-2 belongs to Sarbecovirus) with significant differences in the sequences between SARS-CoV-2 (NC_045512.2) and MERS (MG987420.1 and MG021451.1). The identity of the N and S proteins was 50.0% and 35.0%, respectively, between SARS-CoV-2 (NC_045512.2) and MERS (MG987420.1 and MG021451.1). However, it was interesting that one epitope (KWPWYIWLGF₁₂₁₁₋₁₂₂₀) had the same sequences between SARS-CoV-2 and MERS (NC_045512.2, MT007544.1, and MG987420.1, MG021451.1). This epitope (KWPWYIWLGF₁₂₁₁₋₁₂₂₀) presented by HLA-A*23:01 and HLA-A*24:02 could cover 26.18% of individuals globally. In 2020, Ibrahim and Kafi also predicted the epitope KWPWYIWLGF₁₂₁₁₋₁₂₂₀ because of the high scores that indicate high efficiency due to the prediction of a quantity proportional to the amount of peptide presented by MHC molecules on the cell surface.²⁴ No epitope completely matched with other four human coronaviruses (HCoV-OC43, HCoV-HKU1, HCoV-229E, and HCoV-NL63), because they belong to the different subgenus or different genus from SARS-

CoV-2. HCoV-OC43 (NC_006213.1) and HCoV-HKU1 (NC_006577.2) belong to Embecovirus. HCoV-229E (KY983587.1) and HCoV-NL63 (NC_005831.2) belong to Alphacoronavirus. Thus, the 14-epitope combination could serve as vaccine candidate epitopes for SARS-CoV-2, Bat SARS-like coronavirus, SARS-like coronavirus, and MERS.

Our results indicated the possibility of using candidate CTL epitopes to produce vaccines to effectively control SARS-CoV-2 infections and development. In the current study, we obtained the 14-epitope combination based on the distribution of HLA-A, -B, and -C could cover 89.60% of individuals globally and overcome the limitation of HLA specificity. However, the epitopes we selected were only predicted with respect to binding ability between epitopes and specific MHC molecules based on the total scores as the cut off value for epitope selection. Thus, there were epitopes being ignored because the total scores were below the cut off value (0.5 for N protein, 1.07 for S protein). For example, there are no HLA-A*30:01 predicted epitopes in the entire sequence of the SARS-CoV-2 S protein in our results because all total scores were ≤ 0.6 with all 8,841 peptides for HLA-A*30:01. Moreover, in 2004, Wang *et al.* demonstrated that SARS-

Table 7. The epitope conservancy analysis results for S protein among the 13 coronaviruses.

Name ^a	Epitope sequence	Length ^b	Percent of protein sequence matches ^c	Y_P_009724390.1	QHR84449.1	AVP7804.2.1	ALK02457.1	ATO9813.2.1	ACZ72020.1	Y_P_00982505.1.1	AWH65943.1	AVV62526.1	Y_P_009555241.1	Y_P_173238.1	ARU07601.1	Y_P_003767.1
NP 1	YSVLVNSASFSFKCY ³⁶⁵⁻³⁸⁰	16	15.38% (2/13)	100.00	100.00	81.25	93.75	87.50	87.50	87.50	31.25	31.25	31.25	37.50	31.25	31.25
NP 2	AYYVGLQPRTLKLY ²⁶⁴⁻²⁷⁹	16	15.38% (2/13)	100.00	100.00	81.25	75.00	75.00	75.00	75.00	37.50	37.50	37.50	43.75	31.25	37.50
NP 3	RISNCVADYSLVLY ³⁵⁷⁻³⁶⁹	13	23.08% (3/13)	100.00	100.00	61.54	100.00	100.00	92.31	92.31	30.77	38.46	38.46	46.15	38.46	38.46
NP 4	LOIPFAMQMAVRF ⁸⁹⁴⁻⁹⁰⁶	13	53.85% (7/13)	100.00	100.00	100.00	100.00	100.00	100.00	100.00	46.15	46.15	30.77	38.46	30.77	30.77
NP 5	RYVSSANNCTFEY ¹⁵⁸⁻¹⁷⁰	13	15.38% (2/13)	100.00	100.00	69.23	53.85	53.85	53.85	53.85	38.46	38.46	30.77	38.46	30.77	38.46
NP 6	AYTMSLGAENSVAY ⁶⁹⁴⁻⁷⁰⁷	14	15.38% (2/13)	100.00	100.00	92.86	78.57	78.57	78.57	78.57	42.86	42.86	35.71	28.57	35.71	35.71
NP 7	SWMESEFRVY ¹⁵¹⁻¹⁶⁰	10	15.38% (2/13)	100.00	100.00	81.82	54.55	40.00	40.00	40.00	30.00	40.00	40.00	40.00	40.00	30.00
NP 8	YTNSTFRGVY ²⁸⁻³⁸	11	15.38% (2/13)	100.00	100.00	100.00	40.00	40.00	40.00	40.00	36.36	36.36	36.36	36.36	36.36	36.36
NP 9	HWVFORNFY ¹¹⁰¹⁻¹¹¹⁰	10	23.08% (3/13)	100.00	100.00	100.00	70.00	70.00	70.00	70.00	30.00	30.00	30.00	40.00	40.00	40.00
NP 10	FQFCNDPFLGVY ¹³³⁻¹⁴⁴	12	15.38% (2/13)	100.00	100.00	58.33	41.67	41.67	41.67	41.67	33.33	33.33	33.33	33.33	33.33	33.33
NP 11	SVASQSIAY ⁶⁸⁶⁻⁶⁹⁵	10	15.38% (2/13)	100.00	100.00	40.00	50.00	50.00	50.00	50.00	40.00	40.00	40.00	40.00	50.00	50.00
NP 12	FLPFFSNVTW ⁵⁵⁻⁶⁴	10	15.38% (2/13)	100.00	100.00	80.00	80.00	80.00	80.00	80.00	40.00	40.00	40.00	40.00	40.00	40.00
NP 13	KRSFEDLLF ¹⁸¹⁴⁻⁸²³	10	53.85% (7/13)	100.00	100.00	100.00	100.00	100.00	100.00	100.00	80.00	80.00	80.00	80.00	70.00	70.00
NP 14	LLTDEMIAGY ⁸⁶⁴⁻⁸⁷³	10	15.38% (2/13)	100.00	100.00	90.00	80.00	80.00	80.00	80.00	40.00	50.00	40.00	40.00	40.00	40.00
NP 15	RYVSTGSNVYF ⁶³⁴⁻⁶⁴³	10	15.38% (2/13)	100.00	100.00	50.00	90.00	90.00	80.00	80.00	40.00	50.00	30.00	40.00	40.00	40.00
NP 16	WVTAAGAAAYY ²⁵⁸⁻²⁶⁶	9	15.38% (2/13)	100.00	100.00	66.67	55.56	55.56	55.56	55.56	44.44	44.44	44.44	33.33	44.44	33.33
NP 17	KWPVYWLGLF ¹²¹¹⁻¹²²⁰	10	30.77% (4/13)	100.00	100.00	90.00	90.00	90.00	90.00	90.00	100.00	100.00	70.00	70.00	60.00	60.00
NP 18	KSNIIRGWTF ⁹⁷⁻¹⁰⁶	10	15.38% (2/13)	100.00	100.00	80.00	70.00	70.00	70.00	70.00	40.00	30.00	40.00	40.00	40.00	40.00
NP 19	CYFPLQSYGF ⁴⁸⁸⁻⁴⁹⁷	10	15.38% (2/13)	100.00	100.00	40.00	70.00	70.00	70.00	70.00	40.00	50.00	40.00	40.00	40.00	40.00
NP 20	FVWSQPLF ¹⁶⁸⁻¹⁷⁶	9	15.38% (2/13)	100.00	100.00	66.67	55.56	66.67	55.56	55.56	44.44	44.44	44.44	44.44	44.44	44.44
NP 21	FVKNIIDGY ¹⁹²⁻²⁰⁰	9	15.38% (2/13)	100.00	100.00	66.67	77.78	66.67	77.78	77.78	44.44	44.44	44.44	33.33	44.44	44.44
NP 22	LMSFQSPHGWVF ¹⁰⁵⁷⁻¹⁰⁷⁰	14	23.08% (3/13)	100.00	100.00	100.00	92.86	92.86	92.86	92.86	42.86	42.86	42.86	42.86	42.86	42.86
NP 23	NATRFASWY ¹⁵¹⁻¹⁵⁹	9	15.38% (2/13)	100.00	100.00	88.89	77.78	77.78	77.78	77.78	55.56	55.56	44.44	44.44	44.44	44.44
NP 24	KFTVEKGVY ¹¹²⁻¹²¹	10	15.38% (2/13)	100.00	100.00	70.00	70.00	70.00	70.00	70.00	50.00	50.00	50.00	40.00	50.00	40.00
NP 25	LPNDGVVF ⁸⁴⁻⁹²	9	15.38% (2/13)	100.00	100.00	66.67	77.78	66.67	66.67	66.67	44.44	44.44	44.44	55.56	33.33	44.44
NP 26	TLALHRSY ²⁴⁸⁻²⁵⁶	9	7.69% (1/13)	100.00	88.89	55.56	33.33	33.33	33.33	33.33	44.44	44.44	44.44	44.44	33.33	44.44
NP 27	NDLCTFNWY ³⁹⁶⁻⁴⁰⁴	9	15.38% (2/13)	100.00	100.00	77.78	88.89	88.89	88.89	88.89	44.44	55.56	44.44	44.44	44.44	55.56
NP 28	YEQYKWPWY ¹²¹⁴⁻¹²²³	10	46.15% (6/13)	100.00	100.00	90.00	100.00	100.00	100.00	100.00	70.00	70.00	80.00	80.00	70.00	70.00
NP 29	KRFNDPVLPPF ⁷⁷⁻⁸⁶	10	15.38% (2/13)	100.00	100.00	70.00	80.00	60.00	60.00	60.00	40.00	40.00	30.00	30.00	40.00	40.00
NP 30	FPNITLCPF ³⁷⁻³⁴⁶	10	46.15% (6/13)	100.00	100.00	90.00	100.00	100.00	100.00	100.00	40.00	40.00	50.00	40.00	40.00	40.00
NP 31	NYVADSPVIR ⁴⁰²⁻⁴¹¹	10	15.38% (2/13)	100.00	100.00	70.00	80.00	80.00	80.00	80.00	40.00	40.00	40.00	40.00	50.00	40.00
NP 32	QLPTWRVY ⁶³⁶⁻⁶⁴⁴	9	15.38% (2/13)	100.00	100.00	77.78	88.89	88.89	77.78	77.78	44.44	33.33	44.44	44.44	44.44	44.44
NP 33	KVGGANVLY ⁴⁵²⁻⁴⁶¹	10	15.38% (2/13)	100.00	100.00	40.00	60.00	70.00	60.00	60.00	50.00	40.00	40.00	40.00	30.00	40.00
NP 34	WTFGGAAL ⁸⁹⁴⁻⁹⁰²	9	53.85% (7/13)	100.00	100.00	100.00	100.00	100.00	100.00	100.00	44.44	44.44	44.44	44.44	44.44	44.44

^aEpitope name

^bEpitope length

^cPercent of protein sequence matches at identity 100%

^dThe identity percentage results of each 34 epitopes with 13 S protein sequences

^eAmino-acid position

Table 8. The epitope sequences and its presented HLA alleles of the 14 combination epitopes.

Protein	Epitope sequence	HLA Alleles	
N protein	IAQFAPSASAFF ₃₀₄₋₃₁₅ ^a	A*23:01, B*15:01, B*15:02, B*15:25, B*35:01, C*03:02, C*03:03, C*03:04, C*12:03, C*14:02, C*16:01	
	MSRIGMEVTPSGTWLTY ₃₁₇₋₃₃₃	A*29:02, A*30:02, B*18:01, B*35:01, B*44:02, B*44:03, B*53:01, B*58:01	
	LPQGTTLPKGFY ₁₆₁₋₁₇₂	A*30:02, B*35:01	
	ILLNKHIDAY ₃₅₁₋₃₆₀	B*15:25, B*15:01, B*15:02	
	NTASWFTAL ₄₈₋₅₆	A*68:02	
	LPNNTASWF ₄₅₋₅₃	B*35:01, B*53:01	
	LLLDRLNQL ₂₂₂₋₂₃₀	A*02:01, A*02:06	
	IGYYRRATR ₈₄₋₉₂	A*31:01	
	S protein	LQIPFAMQMAYRF ₈₉₄₋₉₀₆	A*23:01, A*29:02, B*15:01, B*15:25, B*35:01, B*53:01, B*58:01, C*03:02, C*03:03, C*03:04
		KRSFIEDLLF ₈₁₄₋₈₂₃	B*58:01, B*57:01
YEQYIKWPWY ₁₂₁₄₋₁₂₂₃		B*18:01	
FPNITNLCPF ₃₃₇₋₃₄₆		B*35:01	
WTFGAGAAL ₈₉₄₋₉₀₂		C*03:04, C*03:03	
KWPWYIWLGF ₁₂₁₁₋₁₂₂₀		A*23:01, A*24:02	

^aAmino-acid position**Table 9.** The population coverage results (%) of the 14 combination epitopes.

Description	Total Score	Population Coverage		
		Coverage ^a	Average hit ^b	PC90 ^c
8 epitopes in N protein	0.50	0.8413	1.93	0.63
5 epitopes in S protein	1.07	0.5312	1.02	0.21
8 epitopes in N protein and 5 clusters in S protein	-	0.8494	2.95	0.66
14 epitopes for SARS-CoV-2 (NC_045512.2 and MT007544.1)	-	0.8960	3.22	0.96
13 epitopes for SARS-CoV (FJ882944.1 and NC_004718.3)	-	0.8494	2.95	0.66
13 epitopes for SARS-like coronavirus (KT444582.1)	-	0.8494	2.95	0.66
12 epitopes for Bat SARS-like coronavirus (KY417144.1)	-	0.8191	2.53	0.55
11 epitopes for Bat SARS-like coronavirus (MG772934.1)	-	0.8494	2.79	0.66
1 epitope (NP17 in S protein) for MERS (MG987420.1 and MG021451.1)	-	0.2618	0.27	0.14

^aprojected population coverage^baverage number of epitope hits/HLA combinations recognized by the population^cminimum number of epitope hits/HLA combinations recognized by 90% of the population

CoV protein-derived peptide-1 (RLNEVAKNL₁₁₆₇₋₁₁₇₅) could induce peptide-specific CTLs both in vivo and in vitro.²¹ In the current study, we have predicted this epitope using epitope prediction analysis; however, its total score was ≤ -0.54 . So, this epitope was not selected in the current study. Thus, the candidate epitopes which were predicted to reduce infection effects in the current study should also be demonstrated using peptide-sensitized peripheral blood mononuclear cells or isolated CD8⁺ CTL responses in vivo or in vitro level and animal model in the future. Our results will aid in exploring the possible use of these epitopes for the vaccine against SARS-CoV-2 infections.

Author contributions

Conceived and designed the experiments: Li Shi and Yufeng Yao. Performed the HLA data analysis: Sun Ming, Shuying Dai, Le Sun. Performed the immunoinformatic analysis: Yina Cun, Chuanyin Li, Lei Shi. Wrote the paper: Li Shi and Yufeng Yao.

Funding

This work was supported by grant from Special Funds for high-level health talents of Yunnan Province (D-201669 and L-201615) and the Yunnan Provincial Science and Technology Department (2019HC0060). The funders had no role in the design of the study, data collection and analysis, decision to publish, or preparation of the manuscript.

ORCID

Li Shi  <http://orcid.org/0000-0001-9508-7863>

References

- Hui DS, Madani EIA, Ntoumi TA, Kock F, Dar RO, et al. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health - The latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis.* 2020;91:264–66.
- Lu H, Stratton CW, Tang YW. Outbreak of pneumonia of unknown etiology in Wuhan, China: the mystery and the miracle. *J Med Virol.* 2020;92(4):401–02. doi:10.1002/jmv.25678.
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, et al. Clinical Characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA.* 2020;323(11):1061. doi:10.1001/jama.2020.1585.
- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727–33. doi:10.1056/NEJMoa2001017.
- Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, Wang W, Song H, Huang B, Zhu N, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet.* 2020;395(10224):565–74. doi:10.1016/S0140-6736(20)30251-8.
- Chan JF, Kok KH, Zhu Z, Chu H, To KK, Yuan S, Yuen K-Y. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect.* 2020;9(1):221–36. doi:10.1080/22221751.2020.1719902.
- Jiang S, Du L, Shi Z. An emerging coronavirus causing pneumonia outbreak in Wuhan, China: calling for developing therapeutic and prophylactic strategies. *Emerg Microbes Infect.* 2020;9(1):275–77. doi:10.1080/22221751.2020.1723441.
- Leung DT, Tam FC, Ma CH, Chan PK, Cheung JL, Niu H, et al. Antibody response of patients with severe acute respiratory syndrome (SARS) targets the viral nucleocapsid. *J Infect Dis.* 2004;190:379–86.

9. Shi Y, Wan Z, Li L, Li P, Li C, Ma Q, et al. Antibody responses against SARS-coronavirus and its nucleocapsid in SARS patients. *J Clin Virol.* 2004;31(1):66–68. doi:10.1016/j.jcv.2004.05.006.
10. He Y, Zhou Y, Wu H, Luo B, Chen J, Li W, et al. Identification of immunodominant sites on the spike protein of severe acute respiratory syndrome (SARS) coronavirus: implication for developing SARS diagnostics and vaccines. *J Immunol.* 2004;173:4050–57.
11. Zhou M, Xu D, Li X, Li H, Shan M, Tang J, Wang M, Wang F-S, Zhu X, Tao H, et al. Screening and identification of severe acute respiratory syndrome-associated coronavirus-specific CTL epitopes. *J Immunol.* 2006;177(4):2138–45. doi:10.4049/jimmunol.177.4.2138.
12. Coleman CM, Sisk JM, Halasz G, Zhong J, Beck SE, Matthews KL, et al. CD8+ T cells and macrophages regulate pathogenesis in a mouse model of middle east respiratory syndrome. *J Virol.* 2017;91.
13. Tang F, Quan Y, Xin ZT, Wrarmmert J, Ma MJ, Lv H, Wang T-B, Yang H, Richardus JH, Liu W, et al. Lack of peripheral memory B cell responses in recovered patients with severe acute respiratory syndrome: a six-year follow-up study. *J Immunol.* 2011;186(12):7264–68. doi:10.4049/jimmunol.0903490.
14. Peng H, Yang LT, Wang LY, Li J, Huang J, Lu ZQ, Koup RA, Bailer RT, Wu C-Y. Long-lived memory T lymphocyte responses against SARS coronavirus nucleocapsid protein in SARS-recovered patients. *Virology.* 2006;351(2):466–75. doi:10.1016/j.virol.2006.03.036.
15. Fan -Y-Y, Huang Z-T, Li L, Wu M-H, Yu T, Koup RA, Bailer RT, Wu C-Y. Characterization of SARS-CoV-specific memory T cells from recovered individuals 4 years after infection. *Arch Virol.* 2009;154(7):1093–99. doi:10.1007/s00705-009-0409-6.
16. Li CK, Wu H, Yan H, Ma S, Wang L, Zhang M, Tang X, Temperton NJ, Weiss RA, Brenchley JM, et al. T cell responses to whole SARS coronavirus in humans. *J Immunol.* 2008;181(8):5490–500. doi:10.4049/jimmunol.181.8.5490.
17. Ng O-W, Chia A, Tan AT, Jadi RS, Leong HN, Bertoletti A, Tan Y-J. Memory T cell responses targeting the SARS coronavirus persist up to 11 years post-infection. *Vaccine.* 2016;34(17):2008–14. doi:10.1016/j.vaccine.2016.02.063.
18. Cheung YK, Cheng SC, Sin FW, Chan KT, Xie Y. Investigation of immunogenic T-cell epitopes in SARS virus nucleocapsid protein and their role in the prevention and treatment of SARS infection. *Hong Kong Med J.* 2008;14:27–30.
19. Liu J, Wu P, Gao F, Qi J, Kawana-Tachikawa A, Xie J, Vavricka CJ, Iwamoto A, Li T, Gao GF, et al. Novel immunodominant peptide presentation strategy: a featured HLA-A*2402-restricted cytotoxic T-lymphocyte epitope stabilized by intrachain hydrogen bonds from severe acute respiratory syndrome coronavirus nucleocapsid protein. *J Virol.* 2010;84(22):11849–57. doi:10.1128/JVI.01464-10.
20. Tsao YP, Lin JY, Jan JT, Leng CH, Chu CC, Yang YC, et al. HLA-A*0201 T-cell epitopes in severe acute respiratory syndrome (SARS) coronavirus nucleocapsid and spike proteins. *Biochem Biophys Res Commun.* 2006;344:63–71.
21. Wang B, Chen H, Jiang X, Zhang M, Wan T, Li N, Zhou X, Wu Y, Yang F, Yu Y, et al. Identification of an HLA-A*0201-restricted CD8+ T-cell epitope SSp-1 of SARS-CoV spike protein. *Blood.* 2004;104(1):200–06. doi:10.1182/blood-2003-11-4072.
22. Wang YD, Sin WY, Xu GB, Yang HH, Wong TY, Pang XW, He X-Y, Zhang H-G, Ng JNL, Cheng CSS, et al. T-cell epitopes in severe acute respiratory syndrome (SARS) coronavirus spike protein elicit a specific T-cell immune response in patients who recover from SARS. *J Virol.* 2004;78(11):5612–18. doi:10.1128/JVI.78.11.5612-5618.2004.
23. Cheung YK, Cheng SC, Sin FW, Chan KT, Xie Y. Induction of T-cell response by a DNA vaccine encoding a novel HLA-A*0201 severe acute respiratory syndrome coronavirus epitope. *Vaccine.* 2007;25(32):6070–77. doi:10.1016/j.vaccine.2007.05.025.
24. Ibrahim HS, Kafi SK. A computational vaccine designing approach for MERS-CoV infections. *Methods Mol Biol.* 2020;2131:39–145.
25. Bezu L, Kepp O, Cerrato G, Pol J, Fucikova J, Spisek R, Zitvogel L, Kroemer G, Galluzzi L. Trial watch: peptide-based vaccines in anticancer therapy. *Oncoimmunology.* 2018;7(12):e1511506. doi:10.1080/2162402X.2018.1511506.
26. Germain RN, Margulies DH. The biochemistry and cell biology of antigen processing and presentation. *Annu Rev Immunol.* 1993;11(1):403–50. doi:10.1146/annurev.iy.11.040193.002155.
27. Heemels MT, Ploegh H. Generation, translocation, and presentation of MHC class I-restricted peptides. *Annu Rev Biochem.* 1995;64:463–91.
28. Hung CF, Ma B, Monie A, Tsen SW, Wu TC. Therapeutic human papillomavirus vaccines: current clinical trials and future directions. *Expert Opin Biol Ther.* 2008;8(4):421–39. doi:10.1517/14712598.8.4.421.
29. Dhanda SK, Vaughan K, Schulten V, Grifoni A, Weiskopf D, Sidney J, et al. Development of a novel clustering tool for linear peptide sequences. *Immunology.* 2018;155(3):331–45. doi:10.1111/imm.12984.
30. Ahmed SF, Quadeer AA, McKay MR. Preliminary identification of potential vaccine targets for the COVID-19 coronavirus (SARS-CoV-2) based on SARS-CoV immunological studies. *Viruses.* 2020;12(3):254. doi:10.3390/v12030254.
31. Nguyen A, David JK, Maden SK, Wood MA, Weeder BR, Nellore A, et al. Human leukocyte antigen susceptibility map for severe acute respiratory syndrome coronavirus 2. *J Virol.* 2020;94.