



Steiner Wiener index of block graphs

Matjaž Kovše, V A. Rasila & Ambat Vijayakumar

To cite this article: Matjaž Kovše, V A. Rasila & Ambat Vijayakumar (2020) Steiner Wiener index of block graphs, AKCE International Journal of Graphs and Combinatorics, 17:3, 833-840, DOI: [10.1016/j.akcej.2019.11.001](https://doi.org/10.1016/j.akcej.2019.11.001)

To link to this article: <https://doi.org/10.1016/j.akcej.2019.11.001>



© 2020 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 27 Apr 2020.



Submit your article to this journal [↗](#)



Article views: 1386



View related articles [↗](#)



View Crossmark data [↗](#)

Steiner Wiener index of block graphs

Matjaž Kovše^a , V. A. Rasila^b, and Ambat Vijayakumar^b

^aSchool of Basic Sciences, IIT Bhubaneswar, Bhubaneswar, India; ^bDepartment of Mathematics, Cochin University of Science and Technology, Kochi, India

ABSTRACT

Let S be a set of vertices of a connected graph G . The Steiner distance of S is the minimum size of a connected subgraph of G containing all the vertices of S . The Steiner k -Wiener index is the sum of all Steiner distances on sets of k vertices of G . Different simple methods for calculating the Steiner k -Wiener index of block graphs are presented.

KEYWORDS

Distance in graphs; Steiner distance; Wiener index; Steiner k -Wiener index; block graphs; poset of block graphs

MR SUBJECT CLASSIFICATIONS

05C12

1. Introduction

All graphs in this paper are simple, finite and undirected. Unless stated otherwise let $n = |V(G)|$ and $m = |E(G)|$, hence n denotes the order and m the size of a graph G . If G is a connected graph and $u, v \in V(G)$, then the (geodetic) distance $d_G(u, v)$ (or simply $d(u, v)$ if there is no confusion about G) between u and v is the number of edges on a shortest path connecting u and v . The Wiener index $W(G)$ of a connected graph G is defined as

$$W(G) = \sum_{\{u, v\} \in V(G)} d(u, v).$$

The first investigation of this distance-based graph invariant was done by Wiener in 1947, who realized in [21] that there exist correlations between the boiling points of paraffins and their molecular structure and noted that in the case of a tree it can be easily calculated from the edge contributions by the following formula:

$$W(T) = \sum_{e \in E(T)} n(T_1)n(T_2), \quad (1)$$

where $n(T_1)$ and $n(T_2)$ denote the number of vertices in connected components T_1 and T_2 formed by removing an edge e from the tree T .

The Steiner distance of a graph has been introduced in [6] by Chartrand et al., as a natural generalization of the geodetic graph distance. For a connected graph G and $S \subseteq V(G)$, the Steiner distance $d_G(S)$ (or simply $d(S)$) among the vertices of S is the minimum size among all connected subgraphs whose vertex sets contain S . Note that any such subgraph H is a tree, called a Steiner tree connecting vertices from S . Vertices of S are called terminal vertices of tree H ,

while the rest of the vertices of H are called inner vertices of the Steiner tree H . If $S = \{u, v\}$, then $d(S) = d(u, v)$ coincides with the geodetic distance between u and v . In [8] Dankelmann et al. followed by studying the average k -Steiner distance $\mu_k(G)$. In [15], Li et al. introduced a generalization of the Wiener index by using the Steiner distance. The Steiner k -Wiener index $SW_k(G)$ of a connected graph G is defined as

$$SW_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S| = k}} d(S).$$

For $k=2$, the Steiner k -Wiener index coincides with the Wiener index. The average k -Steiner distance $\mu_k(G)$ is related to the Steiner k -Wiener index via the equality $\mu_k(G) = SW_k(G) / \binom{n}{k}$. In [15] the exact values of the Steiner k -Wiener index of the path, star, complete graph, and complete bipartite graph and sharp lower and upper bounds for $SW_k(G)$ for connected graphs and for trees have been obtained. In [11] an application of Steiner k -Wiener index in mathematical chemistry is reported, and it is shown that the term $W(G) + \lambda SW_k(G)$ provides a better approximation for the boiling points of alkanes than $W(G)$ itself, and that the best such approximation is obtained for $k=7$. For a survey on Steiner distance see [16].

The problem of deciding whether for a given subset of vertices in a graph G there exist a Steiner tree of size at most t belongs to the classical NP-complete problems from [10]. Hence for $k > 2$ it is not very likely to find an efficient way to compute the Steiner k -Wiener index for general graphs. Therefore it becomes interesting to either find efficient procedures to compute the Steiner k -Wiener index or bounds for particular classes of graphs.

A vertex v is a cut vertex of graph G if deleting v and all edges incident to it increases the number of connected components G . A block of a graph is a maximal connected vertex induced subgraph that has no cut vertices. A block graph is a graph in which every block is a clique. Block graphs are a natural generalization of trees, and they arise in areas such as metric graph theory, [1], molecular graphs [3] and phylogenetics [9]. They have been characterized in various ways, for example, as certain intersection graphs [12], or in terms of distance conditions [4]. In [3] it has been shown that the determinant of the distance matrix of a block graph depends only on types of blocks, and not on how they are connected. Steiner distance, Steiner centers and Steiner medians of block graphs have been studied in [19].

A vertex of a block graph G that appears in only one block is called a pendant vertex. Hence there are exactly two types of vertices in a block graph: cut vertices and pendant vertices. A block B is called a pendant block if B has a non-empty intersection with a unique block in G . Any block graph different from a complete graph has at least two pendant blocks. Pendant vertex from a pendant block of G is called a leaf of G . For a block graph G with blocks B_1, B_2, \dots, B_t , let $b_i = |V(B_i)|$, for $i \in \{1, \dots, t\}$. We call a sequence $b_1 \geq b_2 \geq \dots \geq b_t$ the block order sequence of G .

The line graph $L(G)$ of a simple graph G is obtained by associating a vertex with each edge of the graph and connecting two vertices with an edge if and only if the corresponding edges of G have a vertex in common. A graph is a line graph of a tree if it is a connected block graph in which each cut vertex is in exactly two blocks, hence they are claw-free block graphs: no induced subgraph is a claw - a complete bipartite graph $K_{1,3}$. A caterpillar is a tree with the property that a path remains if all leaves are deleted. This path is called the backbone of the caterpillar. Line graphs of caterpillars are called path-like block graphs. These are all block graphs with precisely two pendant blocks. The windmill graph $Wd(r, t)$ is a block graph constructed by joining t copies of K_r at a shared vertex, where $r, t \geq 2$. If $v \in V(G)$ is adjacent to all other vertices of G , it is called a universal vertex. A block graph with a universal vertex is called star-like block graph.

For $v \in V(G)$ let $N(v)$ denote the set of all neighbours. The degree of vertex $v \in V(G)$ in graph G is defined as the number of neighbours and denoted with $deg(v)$, i.e., $deg(v) = |N(v)|$. Graph G has degree sequence $\lambda = (d_1, \dots, d_n)$, $d_1 \geq d_2 \geq \dots \geq d_k$, if vertices of G can be indexed from v_1 to v_n such that $deg(v_i) = d_i$, $i = 1, \dots, n$. Let $\mathcal{T}(\lambda)$ be the set of trees with degree sequence λ . It is known that $\mathcal{T}(\lambda)$ is not empty, if and only if $d_1 + \dots + d_n = 2(n - 1)$.

The Cartesian product $G \square H$ of two graphs G and H is the graph with vertex set $V(G) \times V(H)$ and $(a, x)(b, y) \in E(G \square H)$ whenever either $ab \in E(G)$ and $x = y$, or $a = b$ and $xy \in E(H)$. Cartesian products of complete graphs are called Hamming graphs. They can be alternatively described as follows. For $i = 1, 2, \dots, t$ let $r_i \geq 2$ be given integers. Let G be the graph whose vertices are the t -tuples a_1, a_2, \dots, a_t with $a_i \in \{0, 1, \dots, r_i - 1\}$. Two vertices are adjacent if the corresponding tuples differ in precisely one place. Then it is

straightforward to see that G is isomorphic to $K_{r_1} \square K_{r_2} \square \dots \square K_{r_t}$. A subgraph H of G is called isometric (or distance preserving) if $d_H(u, v) = d_G(u, v)$ for all $u, v \in V(H)$. Isometric subgraphs of Hamming graphs are called partial Hamming graphs. The Hamming distance between two t -tuples is defined as the number of positions in which these t -tuples differ, in a partial Hamming graph it coincides with the geodetic distance.

In this paper we obtain several simple methods for calculating the Steiner k -Wiener index of block graphs. In Section 3 we present the block decomposition formula of the Steiner k -Wiener index of block graphs. In Section 4 we present the edge decomposition formula of the Steiner 3-Wiener index of block graphs. In Section 5 we present the vertex decomposition formula of the Steiner k -Wiener index of block graphs and relate it to the k -Steiner betweenness centrality. In Section 6 we study the graphs which minimize or maximize Steiner k -Wiener index among all block graphs with the same set of blocks, and obtain the sharp lower bound. To describe block graphs that maximize the Steiner k -Wiener index we introduce a special graph transformation called generalized block shift, which generalizes the generalized tree shift transformation introduced in [7] by Csikvári.

2. Block decomposition formula of Steiner k -Wiener index of block graphs

Let $n(G)$ denote the number of vertices of a graph G . For a graph G with p , $p > 1$, connected components G_1, G_2, \dots, G_p we denote by $N_k(G)$ the sum over all partitions of k into at least two nonzero parts of products of combinations distributed among the p components of F :

$$N_k(G) = \sum_{\substack{l_1 + l_2 + \dots + l_p = k \\ 0 \leq l_1, l_2, \dots, l_p < k}} \binom{n(G_1)}{l_1} \binom{n(G_2)}{l_2} \dots \binom{n(G_p)}{l_p}$$

For a graph G and $e \in E(G)$, let $G - e$ denote a graph obtained by removing edge e from G . Then the following formula for a tree T has been shown in [11, 14, 15]

$$SW_k(T) = \sum_{e \in E(T)} N_k(T - e).$$

For a given partition $l_1 + l_2 + \dots + l_p = k$, let $\alpha(l_1, l_2, \dots, l_p)$ denote the number of nonzero summands minus 1. For a graph G with p , $p > 1$, connected components G_1, G_2, \dots, G_p , we define $N'_k(G)$ to be the sum over all partitions of k into at least two nonzero parts of products of combinations distributed among the p components of G multiplied by $\alpha(l_1, l_2, \dots, l_p)$:

$$N'_k(G) = \sum_{\substack{l_1 + l_2 + \dots + l_p = k \\ 0 \leq l_1, l_2, \dots, l_p < k}} \binom{n(G_1)}{l_1} \binom{n(G_2)}{l_2} \dots \binom{n(G_p)}{l_p} \cdot \alpha(l_1, l_2, \dots, l_p).$$

For a connected graph G , we define $N'_k(G) = 0$. Note that by definition $\binom{n}{0} = 1$, and $\binom{n}{k} = 0$ whenever $n < k$.




Figure 1. A block graph G with vertices $v_1, v_2, v_3, v_4, v_5, v_6, v_7$ and blocks of orders $b_1 = 2, b_2 = 3$ and $b_3 = 4$.

Let $G \setminus B_i$ denote the graph obtained from G by deleting all edges from block B_i .

Theorem 1. Let G be a connected block graph with blocks B_1, B_2, \dots, B_t . Then

$$SW_k(G) = \sum_{i=1}^t N'_k(G \setminus B_i).$$

Proof. Let $G_1, \dots, G_p, p > 1$ be the components of $G \setminus B_i$ for some $i \in \{1, \dots, t\}$. Let l_1, \dots, l_p vertices are chosen respectively from G_1, \dots, G_p such that $l_1 + l_2 + \dots + l_p = k$ and $\alpha(l_1, l_2, \dots, l_p)$ denote the number of nonzero summands minus one.

For a given set S of vertices with l_1, \dots, l_p of them chosen respectively from G_1, \dots, G_p , it is easy to see that the contribution of block B_i to its Steiner distance $d_G(S)$ is $\alpha(l_1, l_2, \dots, l_p)$, since G is a block graph. Then, $N'_k(G \setminus B_i)$ gives the contribution of block B_i to $SW_k(G)$. Counting the contribution of all blocks, the result follows. \square

Isometric embedding of a graph into Cartesian product of graphs is well known [13]. If G is a block graph with blocks B_1, B_2, \dots, B_t of orders b_1, b_2, \dots, b_t , then it can be isometrically embedded into the Hamming graph $K_{b_1} \square K_{b_2} \square \dots \square K_{b_t}$ as follows.

Let $C_i(1), \dots, C_i(b_i)$ be the connected components of $G \setminus B_i$. For $v \in C_i(j)$, denote $\alpha_i : V \rightarrow \{0, 1, \dots, b_i - 1\}$ the natural contraction of components of $G \setminus B_i$ to the vertices $\{0, 1, \dots, b_i - 1\}$ of K_{b_i} such that each component get different value. Now define an embedding $\alpha : G \rightarrow K_{b_1} \square K_{b_2} \square \dots \square K_{b_t}$ by $\alpha(v) = (\alpha_1(v), \dots, \alpha_t(v))$. We can show that the corresponding Hamming labelling of vertices of G can be further used to calculate the Steiner distance on a subset of more than two vertices.

Theorem 2. Let G be a connected block graph on n vertices with blocks B_1, B_2, \dots, B_t , of orders $b_1 \leq b_2 \leq \dots \leq b_t$. Let $S \subseteq V(G), |S| = k, 2 \leq k \leq n$.

Then

$$d(S) = \sum_{i=1}^t \ell_i(S) - t,$$

where $\ell_i(S)$ denotes the number of different values assigned to the i -th coordinates of vertices from S .

Proof. If all vertices from S belong to the same connected component of $G \setminus B_i$ then the contribution of the block B_i to the Steiner distance between vertices from S is 0 in all

other cases it is equal to $\ell_i(S) - 1$, and hence the formula follows. \square

Corollary 3. Let G be a connected block graph on n vertices with blocks B_1, B_2, \dots, B_t , of orders $b_1 \leq b_2 \leq \dots \leq b_t$. Then, $SW_k(G) = \sum_{S \subseteq V(G), |S|=k} \sum_{i=1}^t \ell_i(S) - \binom{n}{k} t$

Illustration 1. Consider the block graph G in Figure 1. Let $\ell(v)$ denote the Hamming labelling induced by an isometric embedding into $K_2 \square K_3 \square K_4$. Then one possible labeling of vertices of G reads as follows: $\ell(v_1) = (0, 0, 0), \ell(v_2) = (0, 1, 0), \ell(v_3) = (0, 2, 0), \ell(v_4) = (1, 2, 0), \ell(v_5) = (1, 2, 1), \ell(v_6) = (1, 2, 2)$ and $\ell(v_7) = (1, 2, 3)$. Thus for $S = \{v_1, v_2, v_5, v_6\}$, we have $\ell_1(S) = 2, \ell_2(S) = \ell_3(S) = 3$, and hence $d(S) = 5$.

3. Edge decomposition of the Steiner k -Wiener index of block graphs

For an edge $ab \in E(G)$, we denote $W_{ab} = \{v \in V(G) \mid d_G(a, v) < d_G(b, v)\}$, $W_{ba} = \{v \in V(G) \mid d_G(b, v) < d_G(a, v)\}$, and ${}_aW_b = \{v \in V(G) \mid d_G(a, v) = d_G(b, v)\}$. Hence for any edge $ab \in E(G)$, vertices of G are partitioned into three sets: W_{ab} - a set of vertices that are closer to a than to b , W_{ba} - a set of vertices that are closer to b than to a and ${}_aW_b$ - a set of vertices that are at the same distance to a and b . If G is a bipartite graph, then ${}_aW_b = \emptyset$. If ab is a cut edge then also ${}_aW_b = \emptyset$. Additionally we denote $N_{ab} = |W_{ab}|, N_{ba} = |W_{ba}|$ and ${}_aN_b = |{}_aW_b|$. Note that for a pendant vertex a and its neighbour b in a block graph $G, N_{ab} = 1$, and ${}_aN_b = |V(G)| - 2$. For a cut vertex a and its neighbour b in a block graph $G, N_{ab} > 1$.

Theorem 4. Let G be a connected block graph on n vertices. Then

$$W(G) = \sum_{ab \in E(G)} N_{ab} \cdot N_{ba}.$$

Proof. Let $u, v \in V(G)$. Note that any pair of vertices in a block graph is connected by a unique shortest path. Edge ab appears on a shortest path between u and v if and only if W_{ab} contains one of the vertices u, v and W_{ba} the other vertex. The expression on the right side of the formula thus counts the number of times edge ab appears as an edge in a shortest path between a pair of vertices u and v , hence contributing 1 to $d(u, v)$. \square

Theorem 5. Let G be a connected block graph on n vertices. Then

$$SW_3(G) = \sum_{ab \in E(G)} N_{ab} \cdot N_{ba} + \frac{2}{3} \sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot {}_aN_b.$$

Proof. Let $\{u, v, w\} = S \subseteq V(G)$. Edge ab appears in some Steiner tree connecting vertices of S if and only if $S \cap W_{ab} \neq \emptyset$

\emptyset and $S \cap W_{ba} \neq \emptyset$. The expression on the right side counts the contribution of an edge ab to Steiner 3-Wiener index of G . We distinguish two cases.

Case 1. $S \cap W_{ab} \neq \emptyset$, $S \cap W_{ba} \neq \emptyset$ and $S \cap_a W_b = \emptyset$.

In this case ab appears in any Steiner tree connecting vertices of S , therefore it contributes 1 to $d(S)$, altogether this happens $N_{ab} \cdot N_{ba}$ times.

Case 2. $S \cap W_{ab} \neq \emptyset$, $S \cap W_{ba} \neq \emptyset$ and $S \cap_a W_b \neq \emptyset$.

In this case each of W_{ab} , W_{ba} and ${}_a W_b$ includes one of the vertices from S . Without loss of generality, let $u \in W_{ab}$, $v \in W_{ba}$ and $w \in {}_a W_b$. Let c be a common neighbour of a and b that lies on a shortest path between a and w (as well on a shortest path between b and w). Any Steiner tree on S must include a , b and c . Note that $d(\{a, b, c\}) = 2$ and there are three different Steiner trees connecting a , b and c , each of them induced by a pair of edges among ab , ac and bc . Therefore the contribution of each edge among ab , ac and bc to the Steiner distance on S is $\frac{2}{3}$. There are exactly $N_{ab} \cdot N_{ba} \cdot N_b$ number of $\{u, v, w\}$ sets with the property that any Steiner tree connecting $\{u, v, w\}$ includes a and b and some common neighbour c of a and b . Since the contribution to the Steiner distance $d(\{u, v, w\})$ of each edge among ab , ac and bc is defined to be $\frac{2}{3}$, the common contribution of ab , ac and bc in the sum $\frac{2}{3} \sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot N_b$ equals $\frac{2}{3} + \frac{2}{3} + \frac{2}{3} = 2$, which is exactly the contribution of $\{a, b, c\}$ to the Steiner distance $d(\{u, v, w\})$. \square

Theorem 6. Let G be a connected block graph on n vertices. Then

$$SW_3(G) = \left(\sum_{ab \in E(G)} \left[\binom{n}{3} - \binom{n - N_{ab}}{3} - \binom{n - N_{ba}}{3} - \binom{{}_a N_b}{3} \right] \right) - \frac{1}{3} \sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot N_b.$$

Proof. For $ab \in E(G)$, let $n_3(ab)$ denote the number of subsets $S \subseteq V(G)$, $|S| = 3$, for which ab appears as an edge in a Steiner tree connecting vertices of S . Note that an edge ab will not be contained in any Steiner tree connecting vertices of S if and only if one of the cases holds: Case 1: $S \subseteq W_{ab}$, Case 2: $S \subseteq W_{ba}$, Case 3: $S \subseteq {}_a W_b$, Case 4: $S \cap W_{ab} \neq \emptyset$, $S \cap {}_a W_b \neq \emptyset$ and $S \cap W_{ba} = \emptyset$, Case 5: $S \cap W_{ba} \neq \emptyset$, $S \cap {}_a W_b \neq \emptyset$ and $S \cap W_{ab} = \emptyset$. Therefore it follows

$$\begin{aligned} n_3(ab) &= \binom{n}{3} - \binom{N_{ab}}{3} - \binom{N_{ba}}{3} - \binom{{}_a N_b}{3} \\ &\quad - \left[\binom{N_{ab} + {}_a N_b}{3} - \binom{N_{ab}}{3} - \binom{{}_a N_b}{3} \right] \\ &\quad - \left[\binom{N_{ba} + {}_a N_b}{3} - \binom{N_{ba}}{3} - \binom{{}_a N_b}{3} \right] \\ &= \binom{n}{3} - \binom{n - N_{ba}}{3} - \binom{n - N_{ab}}{3} - \binom{{}_a N_b}{3}. \end{aligned}$$

Note that $\sum_{ab \in E(G)} n_3(ab)$ has to be adjusted because of the double counting of the common edge contributions of ab , ac and bc to the Steiner distances of three vertices u , v and w as in the Case 2 in the proof of Theorem 5. For an edge ab , the number of such sets equals $N_{ab} \cdot N_{ba} \cdot N_b$. The common contribution of ab , ac and bc to the Steiner 3-Wiener index in $\sum_{ab \in E(G)} n_3(ab)$ is three instead of two. In $\sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot N_b$ each such triple a , b and c is counted 3 times, hence if we multiply the sum by $-\frac{1}{3}$ we get the common contribution of ab , ac and bc in $\sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot N_b$ is -1 and therefore the altogether contribution of a , b and c in $\sum_{ab \in E(G)} n_3(ab) - \frac{1}{3} \sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot N_b$ is 2, hence:

$$SW_3(G) = \sum_{ab \in E(G)} n_3(ab) - \frac{1}{3} \sum_{ab \in E(G)} N_{ab} \cdot N_{ba} \cdot N_b.$$

\square

Note that for $k \geq 4$ no simple formula for the edge decomposition of the Steiner k -Wiener index of block graph G seems to exist. The reason is that there are more and more different cases which needed to be considered when more than one common neighbour of a and b is included in some Steiner tree for a set of k vertices.

4. Vertex decomposition of the Steiner k -Wiener index of block graphs

For a graph G and $v \in V(G)$, let $G \setminus v$ denote a graph obtained by removing v from G .

Theorem 7. Let G be a connected block graph on n vertices with the set of cut vertices $V_c(G)$ and k an integer such that $2 \leq k \leq n$. Then

$$SW_k(G) = \sum_{v \in V_c(G)} N_k(G \setminus v) + (k - 1) \binom{n}{k}.$$

Proof. $N_k(G \setminus v)$ counts the number of times a cut vertex v is an inner vertex of Steiner tree. Since each such vertex adds 1 to the Steiner distance of a set of k vertices, Steiner distance between k vertices is by $k - 1$ greater than the number of inner vertices in the corresponding Steiner tree, adding $k - 1$ for each set of k vertices, we get the sum of Steiner distances between all k sets of vertices, and the equality in formula holds. \square

Special case of the Theorem 7 on trees has been proved in [14] by Kovše, who also introduced the k -Steiner betweenness centrality $B_k(v)$ of a vertex $v \in V(G)$ as the sum of the fraction of all Steiner trees connecting k terminal vertices, that include v as its inner vertex, across all Steiner trees connecting the same set of k terminal vertices:

$$B_k(v) = \sum_{\substack{A \subseteq V(G) \setminus \{v\} \\ |A| = k}} \frac{\sigma_A(v)}{\sigma_A},$$

where σ_A denotes the number of all Steiner trees between vertices of A in a graph G and $\sigma_A(v)$ denotes the number of

all Steiner trees between vertices of A in a graph G that include v as an inner vertex.

As pendant vertices do not appear as inner vertices of any Steiner tree of a block graph G , it follows that $B_k(v) = 0$ for any pendant vertex v and $B_k(v) > 0$ for all other vertices, if $k < n$. In fact $B_k(v) = N_k(G \setminus v)$ and we can now restate **Theorem 7** as follows.

Theorem 8. *Let G be a connected block graph on n vertices with the set of cut vertices $V_c(G)$ and k an integer such that $2 \leq k \leq n$. Then*

$$SW_k(G) = \sum_{v \in V_c(G)} B_k(v) + (k-1) \binom{n}{k}.$$

Note that in [14] it has been shown that the equality $SW_k(G) = \sum_{v \in V(G)} B_k(v) + (k-1) \binom{n}{k}$ holds for any graph G .

Corollary 9. *Let G be a star-like block graph on n vertices and blocks B_1, B_2, \dots, B_t of orders b_1, b_2, \dots, b_t and k an integer such that $2 \leq k \leq n$. Then*

$$SW_k(G) = (n-1) \binom{n-1}{k-1} - \sum_{i=1}^t \binom{b_i-1}{k}.$$

Proof. Let v be the universal vertex of G . Hence v is the only cut vertex of G . Then v is not an inner vertex of a Steiner tree connecting vertices of S if and only if $S \subseteq B_i \setminus \{v\}$ for some $i \in \{1, \dots, t\}$, hence $B_k(v) = \binom{n-1}{k} - \sum_{i=1}^t \binom{b_i-1}{k}$ and by Theorem

$$\begin{aligned} SW_k(G) &= B_k(v) + (k-1) \binom{n}{k} \\ &= \binom{n-1}{k} - \sum_{i=1}^t \binom{b_i-1}{k} + (k-1) \binom{n}{k}. \end{aligned}$$

From basic binomial identity $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$ it follows that $\binom{n-1}{k} + (k-1) \binom{n}{k} = \binom{n}{k} - \binom{n-1}{k-1} + (k-1) \binom{n}{k} = k \binom{n}{k} - \binom{n-1}{k-1} = n \binom{n-1}{k-1} - \binom{n-1}{k-1} = (n-1) \binom{n-1}{k-1}$, hence the formula follows. \square

Corollary 10. *Let k be an integer such that $2 \leq k \leq n$. For the windmill graph $Wd(r, t)$,*

$$SW_k(Wd(r, t)) = (n-1) \binom{n-1}{k-1} - t \binom{r-1}{k}.$$

Note that for $r=2$ the formula from **Corollary 10** gives formula $SW_k(S_n) = (n-1) \binom{n-1}{k-1}$ for a star S_n on n vertices, already noted in [15].

Corollary 11. *Let G be a path-like block graph on n vertices, with blocks B_1, B_2, \dots, B_t of orders b_1, b_2, \dots, b_t ordered from one pendant block to the other, and k an integer such that $2 \leq k \leq n-1$. Then*

$$\begin{aligned} SW_k(G) &= (k-1) \binom{n}{k} + (t-1) \binom{n-1}{k} \\ &\quad - \left(\sum_{i=1}^{t-1} \binom{\left(\sum_{j=1}^i b_j\right) - i}{k} \right) \\ &\quad + \sum_{i=1}^{t-1} \binom{n+i - \sum_{j=1}^i b_j}{k}. \end{aligned}$$

Proof. Let v_1, v_2, \dots, v_{t-1} denote cut vertices of G , where $v_i \in B_i \cap B_{i+1}$ for $i \in \{1, \dots, t-1\}$. Then v_i is not an inner vertex of a Steiner tree connecting vertices of S if and only if $S \subseteq B_1 \cup \dots \cup B_i \setminus \{v_i\}$ or $S \subseteq B_{i+1} \cup \dots \cup B_t \setminus \{v_i\}$. Hence

$$\begin{aligned} B_k(v_i) &= \binom{n-1}{k} - \binom{\left(\sum_{j=1}^i b_j\right) - i}{k} \\ &\quad - \binom{n+i - \sum_{j=1}^i b_j}{k}, \end{aligned}$$

and the formula follows. \square

Clearly $SW_n(G) = n-1$ for any graph of order n . For $k = n-1$ we can extend the Theorem on trees from [15] as follows.

Theorem 12. *Let G be a block graph of order n with p pendant vertices. Then*

$$SW_{n-1}(G) = n^2 - n - p.$$

Proof. If $k = n-1$, then subset S contains all but one vertex from G . If the vertex missing from S is pendant, then the vertices contained in S form a connected subgraph and its spanning tree of order $n-1$ provides the Steiner tree for S . Therefore $d(S) = n-2$. There are p such subsets, contributing to SW_{n-1} by $p(n-2)$.

If the vertex not present in S is a cut vertex of G , then the respective Steiner tree must contain all vertices of G . Therefore a spanning tree of order n provides the Steiner tree for S and $d(S) = n-1$. There are $n-p$ such subsets, contributing to SW_{n-1} by $(n-p)(n-1)$. Thus $SW_{n-1}(G) = p(n-2) + (n-p)(n-1)$, which proves the theorem. \square

5. Extremal values of Steiner k -Wiener index and GBS poset of block graphs

Among all trees on n vertices star has the minimal Wiener index and path has the largest Wiener index. More generally in extremal problems concerning trees on n vertices it turns out that the maximal (minimal) value of the examined parameter is attained at the star and the minimal (maximal) value is attained at the path. While often it is not hard to prove the extremality of the star, it turns out that to prove the extremality of the path usually requires some effort.

Let $\mathcal{G}(b_1, b_2, \dots, b_t)$ denote the set of all connected block graphs with block order sequence $b_1 \geq b_2 \geq \dots \geq b_t$. We will demonstrate that similar situation, as in the case of the




Figure 2. Proper generalized block shift transforming block graph G_2 on the left to the block graph G_1 on the right. Parallelograms denote blocks of G_1 and G_2 , vertices $x = c_1, c_2, \dots, c_q$ denote cut vertices.

extremality of the Wiener index over trees, happens also for the problem of the extremal values of the Steiner k -Wiener over $\mathcal{G}(b_1, b_2, \dots, b_t)$. If $t \leq 2$ then $\mathcal{G}(b_1, b_2, \dots, b_t)$ consist of a unique block graph, hence in what follows we always assume that $t \geq 3$.

Theorem 13. *Let $G \in \mathcal{G}(b_1, b_2, \dots, b_t)$ and $2 \leq k < n - 1$. Then*

$$(n-1) \binom{n-1}{k-1} - \sum_{i=1}^t \binom{b_i-1}{k} \leq SW_k(G). \quad (2)$$

The equality is attained for the star-like block graph.

Proof. Note that $d(S) \geq k - 1$ for any subset S of k vertices, with equality if and only if Steiner tree connecting S has no inner vertices. Note that the only vertices that can appear as inner vertices of some Steiner trees in a block graph are cut vertices. Hence $SW_k(G)$ will attain the minimum if and only if G has only one cut vertex, which is then a universal vertex of G . Therefore we can use Corollary 9 and the lower bound is proved. \square

In [7] Csikvári introduced a graph transformation called generalized tree shift and defined a partially ordered set (poset) on the set of unlabelled trees with n vertices. The minimal element of this level poset is the path, and the maximal element is the star. Among other results, he showed that going up on this poset decreases the Wiener index.

Next we introduce a graph transformation called a generalized block shift which defines a poset on $\mathcal{G}(b_1, b_2, \dots, b_t)$. We will show that going up on this poset does not increase the Steiner k -Wiener index, which will help us to describe block graphs with minimal and maximal Steiner k -Wiener index among all block graphs from $\mathcal{G}(b_1, b_2, \dots, b_t)$.

Let $G_2 \in \mathcal{G}(b_1, b_2, \dots, b_t)$ and $x, y \in V(G_2)$, such that all blocks having nonempty intersection with the path between x and y (if they exist) have exactly two cut vertices. The generalized block shift (GBS) of G_2 is the block graph G_1 obtained from G_2 as follows: let z be the neighbor of y lying on the path between x and y , we erase all the edges between y and $N(y) \setminus \{z\}$ and add the edges between x and $N(y) \setminus \{z\}$, see Figure 2. Note that $G_1 \in \mathcal{G}(b_1, b_2, \dots, b_t)$.

We denote the cut vertices on the path between x and y of length q by c_1, c_2, \dots, c_q , where $x = c_1$ and $y = c_q$. The set $A \subseteq V(G_2)$ consists of the vertices which can be reached with a path from c_q only through c_1 , and similarly the set $B \subseteq V(G_2)$ consists of those vertices which can be reached with a path from c_1 only through c_q . For the sake of simplicity we denote the corresponding sets in G_1 also with A and B . Furthermore let $A_0 = N(x) \cap A$ in both G_1 and G_2 , and let $B_0 = N(x) \cap B$ in G_1 and $B_0 = N(y) \cap B$ in G_2 .

As in [7] we call x the beneficiary and y the candidate (for being a leaf) of the generalized block shift. Note that if x or y is a leaf in G_2 , then G_1 and G_2 have the same number of pendant blocks, otherwise the number of pendant blocks in G_1 equals the number of pendant blocks in G_2 plus 1. In the latter case we call the generalized block shift proper.

Let $G_1, G_2 \in \mathcal{G}(b_1, b_2, \dots, b_t)$. We denote by $G_1 > G_2$ if G_1 can be obtained from G_2 by some proper generalized block shift. The relation $>$ induces a poset on $\mathcal{G}(b_1, b_2, \dots, b_t)$, which we call GBS poset.

We can always apply a proper generalized block shift to any block graph which has at least one non pendant block. Therefore the only maximal element of GBS poset is the (unique) star-like block graph from $\mathcal{G}(b_1, b_2, \dots, b_t)$. The following theorem shows that the minimal elements of the induced poset are path-like block graphs.

Theorem 14. *Every block graph from $\mathcal{G}(b_1, b_2, \dots, b_t)$ that is not a path-like block graph is the image of some proper generalized block shift.*

Proof. Let G be a block graph from $\mathcal{G}(b_1, b_2, \dots, b_t)$ that is not path-like, i. e., it has at least one cut vertex belonging to at least 3 blocks. Let v be a leaf of G and w the closest cut vertex to v which belongs to at least 3 blocks. Then the cut vertices (if they exist) on the path between v and w belong to exactly 2 blocks. Vertex w belongs to at least two blocks different from the one which has a nonempty intersection with the path between v and w , so we can split $N(w)$ into two nonempty sets A_0 and B_0 , where vertices of A_0 consist of those that form a block which has nonempty intersection with the path between v and w . Let G' be the block graph




Figure 3. A greedy tree T and its line graph $L(T)$.

obtained by erasing the edges between w and B_0 and adding the edges between v and B_0 . Then G can be obtained from G' by a generalized block shift, where w is the beneficiary and v is the candidate. Since both A_0 and B_0 are nonempty this is a proper generalized block shift. \square

Corollary 15. *The star-like block graph is the unique maximal element of the GBS poset on $\mathcal{G}(b_1, b_2, \dots, b_t)$. The set of minimal elements consist of all path-like block graphs.*

Theorem 16. *A proper generalized block shift decreases the value of the Steiner k -Wiener index if and only if $|A \cup B| \geq k$, otherwise the value remains the same.*

Proof. Let G_2 be a block graph and G_1 its image obtained by a proper generalized block shift, moreover let $2 \leq k \leq n - 2$, where $n = |V(G_1)| = |V(G_2)|$. Let d^1 and d^2 denote the Steiner distance in G_1 and G_2 respectively. For $i \in \{1, \dots, q\}$ it follows that $d^1(\{c_i\} \cup A_1) + d^1(\{c_{q+1-i}\} \cup A_1) = d^2(\{c_i\} \cup A_1) + d^2(\{c_{q+1-i}\} \cup A_1)$ for all $A_1 \subseteq A$ and $d^1(\{c_i\} \cup B_1) + d^1(\{c_{q+1-i}\} \cup B_1) = d^2(\{c_i\} \cup B_1) + d^2(\{c_{q+1-i}\} \cup B_1)$ for all $B_1 \subseteq B$. Moreover $d^1(A_1) = d^2(A_1)$ for $A_1 \subseteq A$, and $d_1(B_1) = d_2(B_1)$ for $B_1 \subseteq B$ and $d^2(A_1 \cup B_1) = d^1(A_1 \cup B_1) + (q - 1)$ for $A_1 \subseteq A$ and $B_1 \subseteq B$. Altogether we have

$$\begin{aligned} SW_k(G_2) &= \sum_{\substack{S \subseteq V(G_2) \\ |S|=k}} d^2(S) = \sum_{\substack{S \subseteq V(G_1) \\ |S|=k}} d^1(S) \\ &+ (q - 1) \cdot \sum_{\substack{l_1 + l_2 = k \\ 0 < l_1, l_2 < k}} \binom{|A|}{l_1} \binom{|B|}{l_2} \\ &= SW_k(G_1) + (q - 1) \cdot \sum_{\substack{l_1 + l_2 = k \\ 0 < l_1, l_2 < k}} \binom{|A|}{l_1} \binom{|B|}{l_2}. \end{aligned}$$

If $|A \cup B| \geq k$ then the generalized block shift decreases Steiner k -Wiener index, otherwise the value remains the same. \square

Hence going up the GBS poset never increases the Steiner k -Wiener index. Moreover from the Theorem 16 and the fact that the path-like block graphs are minimal elements, and the star-like block graph is the only maximal

element of the poset induced by the generalized block shift on $\mathcal{G}(b_1, b_2, \dots, b_t)$, we get the following corollary.

Corollary 17. *Let $2 \leq k \leq n$. Among all block graphs from $\mathcal{G}(b_1, b_2, \dots, b_t)$ the minimum value of the Steiner k -Wiener index is attained by the star-like block graph, and the maximum value of the Steiner k -Wiener index is attained among path-like block graphs.*

Hence Corollary 17 implies Theorem 13. Finding the upper bound of the Steiner k -Wiener index over $\mathcal{G}(b_1, b_2, \dots, b_t)$ turns out to be more challenging problem. Let $\mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$ denote the set of all connected claw-free block graphs with block order sequence (b_1, b_2, \dots, b_t) , where $b_1 \geq b_2 \geq \dots \geq b_t$. Recall that claw-free block graphs are precisely the line graphs of trees. In [4] Buckley has shown the following result.

Theorem 18. *Let T be a tree on n vertices. Then $W(L(T)) = W(T) - \binom{n}{2}$.*

Note that for any $G \in \mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$, any cut vertex of G is incident to exactly two blocks, therefore G has maximal possible number of cut vertices among all block graphs from $\mathcal{G}(b_1, b_2, \dots, b_t)$. Hence all graphs from $\mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$ have the same number of cut vertices, which is equal to $t - 1$, and therefore also the same number of pendant vertices, which equals $n - t + 1$.

By Theorem 18 finding minimum or maximum value of the Wiener index over the set $\mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$ is equivalent to finding a minimum or maximum value of the Wiener index on the set of trees with the degree sequence $(b_1, \dots, b_t, 1, 1, \dots, 1)$, where 1's corresponds to pendant vertices in block graphs from $\mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$, hence 1 appears $n - t + 1$ times in the sequence.

Let (d_1, \dots, d_n) be a degree sequence of the tree T with $d_1 \geq d_2 \geq \dots \geq d_k \geq 2 > 1 = d_{k+1} = \dots = d_n$. Then T is called greedy tree if it can be embedded in the plane as follows:

1. take the vertex v with degree d_1 as the root,
2. each vertex u lies on some line i where i is the distance between the root v and u ,
3. each line is filled up with vertices in decreasing degree order from left to right.

Next we provide an example of a greedy tree and its line graph.

Illustration 2. Let $(4, 4, 4, 3, 3, 3, 2, 2, 2, 1, 1, 1, 1, 1, 1, 1, 1, 1)$ be the given degree sequence. Then its corresponding greedy tree using the plane embedding from the definition is shown in Figure 3.

Wang [20] and Zhang et al. [22] have shown independently the following theorem.

Theorem 19. Let (d_1, \dots, d_n) be an integer sequence with $d_1 \geq d_2 \geq \dots \geq d_s \geq 2 > 1 = d_{s+1} = \dots = d_n$ and $\sum_{i=1}^n d_i = 2(n-1)$. Then the greedy tree with degree sequence (d_1, \dots, d_n) minimizes the Wiener index among all trees with same degree sequence.

From Theorem 18 and Theorem 19 we immediately get the following corollary.

Corollary 20. Then line graph of the greedy tree with the degree sequence $(b_t, b_{t-1}, \dots, b_2, b_1, 1, 1, \dots, 1)$ minimizes the Wiener index over the set $\mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$.

The maximization problem is somewhat more complicated. It has been proven by Shi in [17] that the problem can be reduced to the study of caterpillars. Schmuck et al. [18] have shown that an optimal tree is a caterpillar with vertex degrees non-increasing from the ends of the caterpillar towards its central part. In [5] Schmuck et al. obtained a polynomial time algorithm for finding a caterpillar T that maximizes the Wiener index among all trees with a prescribed degree sequence $(b_t, b_{t-1}, \dots, b_2, b_1, 1, 1, \dots, 1)$. We call T the optimal caterpillar for the degree sequence $(b_t, b_{t-1}, \dots, b_2, b_1, 1, 1, \dots, 1)$.

Corollary 21. The line graph of the optimal caterpillar for the degree sequence $(b_t, b_{t-1}, \dots, b_2, b_1, 1, 1, \dots, 1)$ maximizes the Wiener index over the set $\mathcal{G}_{CF}(b_1, b_2, \dots, b_t)$.

Although we are not aware of a possible generalization of Buckley's theorem which would relate the Steiner k -Wiener index of a tree to the Steiner k -Wiener index of its line graph, due to the fact that Steiner distance behaves nicely on block graphs, as demonstrated throughout this paper, we propose also the following problem.

Problem 1. Let k be an integer with $3 \leq k \leq n$. Let (d_1, \dots, d_n) be an integer sequence with $d_1 \geq d_2 \geq \dots \geq d_s \geq 2 > 1 = d_{s+1} = \dots = d_n$ and $\sum_{i=1}^n d_i = 2(n-1)$. Is it true that the line graph of the greedy tree with the degree sequence (d_1, \dots, d_n) minimizes the Steiner k -Wiener index over the set $\mathcal{G}(d_1, \dots, d_n)$, and that the line graph of the optimal caterpillar for the degree sequence (d_1, \dots, d_n) maximizes the Steiner k -Wiener index over the set $\mathcal{G}(d_1, \dots, d_n)$?

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

The second author acknowledges the support by KSCSTE research fellowship with reference No.001/FSHP-MAIN/2015/KSCSTE.

ORCID

Matjaž Kovše  <http://orcid.org/0000-0001-9473-7545>

References

- [1] Bandelt, H.-J., Mulder, H. M. (1986). Distance-hereditary graphs. *J. Comb. Theory, Ser. B* 41(2):182–208.
- [2] Bapat, R. B., Sivasubramanian, S. (2011). Inverse of the distance matrix of a block graph. *Linear Multilinear Algebra* 59(12): 1393–1397.
- [3] Behtoei, A., Mohsen, J., Bijan, T. (2010). A characterization of block graphs. *Discrete Appl. Math.* 158(3):219–221.
- [4] Buckley, F. (1981). Mean distance in line graphs. *Congr. Numer.* 32:153–162.
- [5] Čela, E., Schmuck, N. S., Wimer, S., Woeginger, G. J. (2011). The Wiener maximum quadratic assignment problem. *Discrete Optim.* 8(3):411–416.
- [6] Chartrand, G., Oellermann, O. R., Tian, S. L., Zou, H. B. (1989). Steiner distance in graphs. *Časopis pro pěstování matematiky* 114:399–410.
- [7] Csikvári, P. (2010). On a poset of trees. *Combinatorica* 30(2): 125–137.
- [8] Dankelmann, P., Oellermann, O. R., Swart, H. C. (1996). The average Steiner distance of a graph. *J. Graph Theory* 22(1):15–22.
- [9] Dress, A., Huber, K. T., Koolen, J., Moulton, V., Spillner, A. (2016). Characterizing block graphs in terms of their vertex-induced partitions. *Australas. J. Comb.* 66:1–9.
- [10] Garey, M. R., Johnson, D. S. (1979). *Computers and Intractability – A Guide to the Theory of NP-Completeness*, Freeman. San Francisco, CA: WH Freeman.
- [11] Gutman, I., Furtula, B., Li, X. (2015). Multicenter Wiener indices and their applications. *J. Serb. Chem. Soc.* 80(8):1009–1017.
- [12] Harary, F. (1963). A characterization of block-graphs. *Can. Math. Bull.* 6(1):1–6.
- [13] Imrich, W., Klavzar, S., Rall, D. F. (2008). *Topics in Graph Theory: Graphs and Their Cartesian Product*. Wellesley, MA: AK Peters/CRC Press.
- [14] Kovše, M. (2016). Vertex decomposition of Steiner Wiener index and Steiner betweenness centrality. arXiv:1605.00260 [math.CO].
- [15] Li, X., Mao, Y., Gutman, I. (2016). The Steiner Wiener index of a graph. *Discuss. Math. Graph Theory* 36(2):455–465.
- [16] Mao, Y. (2017). Steiner distance in graphs—A survey. Preprint, arXiv:1708.05779 [math.CO].
- [17] Shi, R. (1993). The average distance of trees. *Sys. Sci. Math. Sci.* 6:18–24.
- [18] Schmuck, N. S., Wagner, S. G., Wang, H. (2012). Greedy trees, caterpillars, and Wiener-type graph invariants. *MATCH Commun. Math. Comput. Chem.* 68:273–292.
- [19] Yeh, H.-G., Chiang, C.-Y., Peng, S.-H. (2008). Steiner centers and Steiner medians of graphs. *Discrete Math.* 308(22): 5298–5307.
- [20] Wang, H. (2008). The extremal values of the Wiener index of a tree with given degree sequence. *Discrete App. Math.* 156(14): 2647–2654.
- [21] Wiener, H. (1947). Structural determination of paraffin boiling points. *J. Am. Chem. Soc.* 69(1):17–20.
- [22] Zhang, X.-D., Xiang, Q.-Y., Xu, L.-Q., Pan, R.-Y. (2008). The Wiener index of trees with given degree sequences. *MATCH Commun. Math. Comput. Chem.* 60(2):623–644.