



Upper dimension and bases of zero-divisor graphs of commutative rings

S. Pirzada, M. Aijaz & S.P. Redmond

To cite this article: S. Pirzada, M. Aijaz & S.P. Redmond (2020) Upper dimension and bases of zero-divisor graphs of commutative rings, AKCE International Journal of Graphs and Combinatorics, 17:1, 168-173, DOI: [10.1016/j.akcej.2018.12.001](https://doi.org/10.1016/j.akcej.2018.12.001)

To link to this article: <https://doi.org/10.1016/j.akcej.2018.12.001>



© 2018 Kalasalingam University. Published with license by Taylor & Francis Group, LLC.



Published online: 01 Jun 2020.



Submit your article to this journal [↗](#)



Article views: 133



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 2 View citing articles [↗](#)



Upper dimension and bases of zero-divisor graphs of commutative rings

S. Pirzada^{a,*}, M. Aijaz^a, S.P. Redmond^b

^a Department of Mathematics, University of Kashmir, Srinagar, India

^b Department of Mathematics and Statistics, Eastern Kentucky University, Richmond, KY, USA

Received 20 July 2018; received in revised form 24 December 2018; accepted 27 December 2018

Abstract

For a commutative ring R with non-zero zero divisor set $Z^*(R)$, the zero divisor graph of R is $\Gamma(R)$ with vertex set $Z^*(R)$, where two distinct vertices x and y are adjacent if and only if $xy = 0$. The upper dimension and the resolving number of a zero divisor graph $\Gamma(R)$ of some rings are determined. We provide certain classes of rings which have the same upper dimension and metric dimension and give an example of a ring for which these values do not coincide. Further, we obtain some bounds for the upper dimension in zero divisor graphs of commutative rings and provide a subset of vertices which cannot be excluded from any resolving set.

Keywords: Ring; Zero-divisor graph; Upper dimension; Resolving number

1. Introduction

Throughout this article, we will consider only commutative rings R with unity $1 \neq 0$, and we will let $Z(R)$ be the set of zero divisors of R . The zero divisor graph of R denoted by $\Gamma(R)$ is the undirected graph having vertex set $V(\Gamma(R)) = Z^*(R) = Z(R) \setminus \{0\}$, with distinct vertices x and y being adjacent if and only if $xy = 0$. This definition of the zero divisor graph is due to D.F. Anderson and Livingston [1], who extended the earlier work of Beck [2] and D.D. Anderson and Naseer [3] which used all zero-divisors of the ring as vertices. The zero divisor graph translates the algebraic properties of a ring to graph theoretical tools, and therefore it can help in exploring interesting results in both graph theory and abstract algebra. The zero-divisor graph of a commutative ring has been studied by many authors and has been extended to several other algebraic structures.

This paper is organized as follows. In Section 2, we analyze the relation between the upper dimension and the graph structure of $\Gamma(R)$ for a commutative ring R . Rings with finite metric dimension and those for which upper dimension and resolving number are same are characterized. In Section 3, it is shown that the upper dimension and metric dimension (lower dimension) are the same in zero divisor graphs for all finite commutative rings of odd characteristic

Peer review under responsibility of Kalasalingam University.

* Corresponding author.

E-mail addresses: pirzadasd@kashmiruniversity.ac.in (S. Pirzada), ahajjaz99@gmail.com (M. Aijaz), shane.redmond@eku.edu (S.P. Redmond).

<https://doi.org/10.1016/j.akcej.2018.12.001>

© 2018 Kalasalingam University. Published with license by Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

and for rings of order $2k$, where k is an odd integer. Finally, several examples are discussed, with methods to compute the upper dimension.

2. Upper dimension and bases of $\Gamma(R)$

The concept of metric dimension of a graph was introduced in 1970s by Slater [4] and independently by Harary and Melter [5]. The concept of *upper dimension* of graphs was introduced by Chartrand et al. [6].

Definition 2.1. Consider a connected graph G on n vertices. For a given vertex $v \in V(G)$, the representation $r(v | W)$ for v with respect to an ordered set $W = \{w_1, w_2, \dots, w_k\}$ of vertices of G is a k -tuple defined as

$$r(v | W) = (d(v, w_1), d(v, w_2), \dots, d(v, w_k)),$$

where $d(x, y)$ represents the distance between the two vertices x and y of G . Clearly, the representation for the i th vertex in W has 0 in the i th coordinate and all other coordinates are non-zero. So the vertices of W necessarily have distinct representations. Thus the representations of only those vertices that are not in W need to be examined to check if these representations are distinct. The set W is called a *resolving set* if all vertices of G have different representations with respect to W . A resolving set W is called a *minimal resolving set* if no proper subset of W is a resolving set of G . A minimal resolving set containing the minimum number of vertices is called a *metric basis* for G and the cardinality of a metric basis is called the *metric dimension* of G , denoted by $\dim(G)$. A minimal resolving set with the largest number of vertices is called an *upper basis* of G and its cardinality is called the *upper dimension* which is denoted by $\dim^+(G)$. It is obvious that for a graph on n vertices, every subset of $(n - 1)$ vertices is a resolving set. Thus, for any connected graph G , $\dim^+(G) \leq n - 1$.

The *resolving number*, denoted by $\text{res}(G)$, of a connected graph G is the smallest positive integer k such that every set of k vertices of G is a resolving set of G . Since the order of an upper basis is the largest minimal resolving set and resolving number is the order of a resolving set (whether minimal or not) we have the following inequality on metric dimension, upper dimension and the resolving number for a connected graph G of order $n \geq 2$,

$$1 \leq \dim(G) \leq \dim^+(G) \leq \text{res}(G) \leq n - 1$$

We begin by summarizing some results on metric dimension and upper dimension of graphs which will be used in throughout this section. For undefined notations and terminology from graph theory, the readers are referred to [7].

Lemma 2.1 (Lemma 2.2 [8]). For a connected graph G of order $n \geq 1$, $\dim^+(G) = 1$ if and only if $G \cong P_2$ or P_3 and, for $n \geq 4$, $\dim^+(P_n) = 2$, where P_n denotes the path on n vertices.

Lemma 2.2 (Lemma 2.4 [8]). A connected graph G of order n has upper dimension equal to $n - 1$ if and only if $G \cong K_n$.

Lemma 2.3 (Lemma 2.5 [8]). The upper dimension of a cycle C_n is 2, where $n \geq 3$ is a positive integer.

Theorem 2.4 (Theorem 4.2 [8]). For a positive integer $n \geq 3$, $\dim^+(K_{1,n} \times K_2) = \dim^+(K_{1,n}) + 1 = n$.

In this and later sections, we denote the ring of integers by \mathbb{Z} , the ring of integers modulo n by \mathbb{Z}_n , and the field with q elements by \mathbb{F}_q . As we now begin to discuss zero-divisor graphs of commutative rings, we remind the reader of the most fundamental characteristics of the structure of such graphs.

Theorem 2.5 (Theorem 2.3 [1]). Let R be a commutative ring. Then $\Gamma(R)$ is connected and $\text{diam}(\Gamma(R)) \leq 3$.

Lemma 2.6. If R is a commutative ring with unity such that $\Gamma(R)$ is a path, then $|Z^*(R)| \leq 3$.

Proof. By [Theorem 2.3 [1]], $\Gamma(R)$ is connected and $\text{diam}(\Gamma(R)) \leq 3$. Thus, $\Gamma(R)$ cannot be a path of length greater than 4.

If possible, let $|Z^*(R)| = 4$ for some ring R , where $\Gamma(R)$ is a path, say $a - b - c - d$ such that $ab = bc = cd = 0$ are the only zero divisor relations. Note that $a + c \in Z(R)$ since $b(a + c) = 0$. Clearly, $a + c \neq a$ and $a + c \neq c$.

Also, $a + c \neq d$ since $bd \neq 0$ and $a + c \neq 0$ since $d(a + c) = da + dc = da \neq 0$. Therefore, $a + c = b$. A similar argument shows $c = b + d$. Hence, $c - d = b = a + c$. Thus $a = -d$. However, this is a contradiction, since $-dc = 0$ but $ac \neq 0$. Thus, $|Z^*(R)| = 4$ is not possible. \square

Theorem 2.7. *Let R be a commutative ring with unity. Then $\dim^+(\Gamma(R)) = 1$ if and only if R is one of the following rings.*

- (i) $\frac{\mathbb{Z}_3[x]}{(x^2)}, \mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_9$.
- (ii) $\mathbb{Z}_6, \mathbb{Z}_8, \frac{\mathbb{Z}_2[x]}{(x^3)}, \frac{\mathbb{Z}_4[x]}{(2x, x^2 - 2)}$.

Proof. The lists here give the only rings (up to isomorphism) whose zero-divisor graph is isomorphic to (i) P_2 or (ii) P_3 . Hence, the result follows by Lemmas 2.6 and 2.1. \square

Notice that $\dim^+(\Gamma(R)) = 1$ if and only if $\Gamma(R)$ is a path by Lemma 2.1. However, the same is not true for a graph G in general, since $\dim^+(G) \geq 2$ if $G \neq P_2, P_3$. Further, if $\Gamma(R)$ is a path, then $\Gamma(R)$ has exactly two upper basis sets, since only the end vertex forms a resolving set.

Theorem 2.8. *Let R be a commutative ring with unity. Then $\dim^+(\Gamma(R))$ is finite if and only if R is finite (and not a domain).*

Proof. If R is finite, then $|Z^*(R)|$ is finite and therefore $\dim^+(\Gamma(R))$ is finite. Now, suppose $\dim^+(\Gamma(R))$ is finite. Let W be the upper basis set with $|W| = k$, where k is some positive integer. For any two vertices x and y of $\Gamma(R)$, $d(x, y) \in \{0, 1, 2, 3\}$ by Theorem 2.5. Now, for each vertex $x \in Z^*(R)$, the representation $r(x | W)$ is a k -coordinate vector (d_1, d_2, \dots, d_k) , where each $d_i \in \{0, 1, 2, 3\}$, $1 \leq i \leq k$. As each d_i has four possibilities, therefore the total number of possibilities for $r(x | W)$ is 4^k . Since W is a resolving set, therefore $r(x | W)$ is unique for each vertex $x \in Z^*(R)$ so that $|Z^*(R)| \leq 4^k$. This implies that $Z^*(R)$ is finite and hence R is finite. \square

Note that $\dim^+(\Gamma(R))$ is finite if and only if R is finite, let $\dim^+(\Gamma(R)) = k$ and let $W = \{w_1, w_2, \dots, w_k\}$ be the upper basis. Since each coordinate of $r(x|W)$ is non-zero whenever $x \in V(\Gamma(R)) - W$, we see that every coordinate in $r(x|W)$ belongs to the set $\{1, 2, 3\}$, as $\text{diam}(\Gamma(R)) \leq 3$. Therefore $|Z^*(R)| \leq 3^k + k$. We note that this is a better bound than given in proof of Theorem 2.8 as $3^k + k < 4^k$ for all $k \geq 2$.

Corollary 2.9. *Let R be a commutative ring with unity 1 (and not a domain) such that $\dim^+(\Gamma(R)) = k$, where k is any positive integer. Then $|Z(R)| \leq 4^k + 1$.*

If R is a commutative ring (and not a domain), we also notice that $|Z(R)| \leq 4^k + 1$ gives $k \geq \lceil \log_4(|Z(R)| - 1) \rceil = \lceil \log(|Z^*(R)|) \rceil$, thus we have a lower bound for the upper dimension of $\Gamma(R)$. The equality holds when $|\Gamma(R)| = 1, 2$. For a ring R , $\dim^+(\Gamma(R)) = \text{res}(\Gamma(R)) = 1$ if and only if $\Gamma(R) \cong P_2$, that is, if and only if $R \cong \frac{\mathbb{Z}_3[x]}{(x^2)}, \mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_9$.

Theorem 2.10. *Let R be a commutative ring with unity.*

- (i) $\dim^+(\Gamma(R)) = \text{res}(\Gamma(R)) = 2$ if and only if $R \cong \frac{\mathbb{F}_4[x]}{(x^2)}, \frac{\mathbb{Z}_4[x]}{(x^2+x+1)}, \frac{\mathbb{Z}_4[x]}{(2,x)^2}, \frac{\mathbb{Z}_2[x,y]}{(x,y)^2}$.
- (ii) $\dim^+(\Gamma(R)) = \text{res}(\Gamma(R)) = |Z^*(R)| - 1$ if and only if $\Gamma(R)$ is complete

Proof. (i) To prove the result, we show if $\dim^+(\Gamma(R)) = \text{res}(\Gamma(R)) = 2$, then $\Gamma(R)$ is either a path or a cycle. For this, we first show that $\Delta(\Gamma(R)) = 2$, where $\Delta(\Gamma(R)) = 2$ denotes the largest degree of a vertex in $\Gamma(R)$.

We claim that there does not exist a subset of vertices $D = \{x, a, b, c\}$ in $\Gamma(R)$ with the property $ax = bx = cx = 0$ and the restriction $\text{res}(\Gamma(R)) = 2$, for otherwise, we have the following cases as given in Fig. 1. In each of the four graphs in Fig. 1, the bold vertices represent the set of two vertices that do not form a resolving set. Thus, the graphs (a), (b), (c), (d) in Fig. 1 completely describe the situation that a graph having a vertex of degree 3 or more cannot have resolving number equal to 2. Therefore, we must have $\Delta(\Gamma(R)) \leq 2$. Also, if $\Delta(\Gamma(R)) = 1$, then $\Gamma(R) = P_2$ and $\text{res}(P_2) = 1$. Thus, we have $\Delta(\Gamma(R)) = 2$. Hence $\Gamma(R)$ is either a path P_3 or a cycle C_3 or C_4 . Therefore, we must have $\Gamma(R) \cong C_3$ or C_4 . Since the two non-adjacent vertices of C_4 do not form a resolving set, therefore $\Gamma(R) \cong C_3$ and so $R \cong \frac{\mathbb{F}_4[x]}{(x^2)}, \frac{\mathbb{Z}_4[x]}{(x^2+x+1)}, \frac{\mathbb{Z}_4[x]}{(2,x)^2}, \frac{\mathbb{Z}_2[x,y]}{(x,y)^2}$.




Fig. 1. Graphs with resolving number 2

(ii). Clearly $\dim^+(\Gamma(R)) \leq \text{res}(\Gamma(R)) \leq n - 1$ and any subset of $n - 1$ vertices of complete graph forms a resolving set. Hence, the result follows by Lemma 2.2. \square

Example 2.11. It is easily verified that if $R \cong \mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}, \mathbb{Z}_2 \times \mathbb{F}_4, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_3 \times \mathbb{Z}_3$, then $\dim^+(\Gamma(R)) = 2$. However, in each case, one can find a set of two distinct vertices of $\Gamma(R)$ that do not form a resolving set. Thus $\text{res}(\Gamma(R)) > 2$.

Recall that an element x in a ring R is said to be nilpotent if there exists a positive integer k such that $x^k = 0$.

Theorem 2.12. Let R be a ring where every zero-divisor is nilpotent.

- (i) If $|Z^*(R)| = 1$ or 2 , then $\dim^+(\Gamma(R)) = 1$.
- (ii) If $|Z^*(R)| \geq 3$, and $Z(R)^2 = \{0\}$, then $\dim^+(\Gamma(R)) = |Z^*(R)| - 1$.
- (iii) If $|Z^*(R)| \geq 3$, and $Z(R)^2 \neq \{0\}$, then $\dim^+(\Gamma(R)) \leq |Z^*(R)| - 2$.

Proof. (i) If $|Z^*(R)| = 1$, then $R \cong \mathbb{Z}_4$ or $\frac{\mathbb{Z}_2}{(x^2)}$ and so $\Gamma(R) = K_1$. If $|Z^*(R)| = 2$, then $R \cong \mathbb{Z}_9$ or $\mathbb{Z}_3[x]/(x^2)$ or $\mathbb{Z}_2 \times \mathbb{Z}_2$ and so $\Gamma(R) = K_2$. Therefore, in each case $\dim^+(\Gamma(R)) = 1$.

(ii) If $|Z^*(R)| \geq 3$, and $Z(R)^2 = \{0\}$, then $xy = 0$ for all $x, y \in Z^*(R)$, by Theorem 2.8 [1]. Thus, $\Gamma(R)$ is a complete graph and so by Lemma 2.2, $\dim^+(\Gamma(R)) = |Z^*(R)| - 1$.

(iii) If $|Z^*(R)| \geq 3$ and $Z(R)^2 \neq \{0\}$, there exists some $x \in Z^*(R)$ such that $x^2 \neq 0$. So there exists $y \in Z^*(R)$ such that $d(x, y) \geq 2$. Therefore, $\dim^+(\Gamma(R)) \leq |Z^*(R)| - 2$ as $Z^*(R) - \{z, y\}$ is a resolving set for any vertex z adjacent to x . \square

3. Characteristic of a Ring and Star Subsets

Resolving sets for zero-divisor graphs have previously been studied in [9] and [10]. In these articles, it was noted that distance similarity was a key factor in determining resolving sets. Vertices x and y of a graph G are called *distance similar* if $d(x, a) = d(y, a)$ for all $a \in V(G) - \{x, y\}$. The following results illustrate this connection between concepts.

Theorem 3.1 (Theorem 2.1 [9]). Let G be a connected graph. Suppose G is partitioned into k distinct distance similar classes V_1, V_2, \dots, V_k (that is, $x, y \in V_i$ if and only if $d(x, a) = d(y, a)$ for all $a \in V(G) - \{x, y\}$).

- (i) Any resolving set W for G contains all but at most one vertex from each V_i .
- (ii) Each V_i induces a complete subgraph or a graph with no edges.
- (iii) $\dim(G) \geq |V(G)| - k$.
- (iv) There exists a minimal resolving set W for G such that if $|V_i| > 1$, at most $|V_i| - 1$ vertices of v_i are elements of W .
- (v) If m is the number of distance similar classes that consist of a single vertex, then $|V(G)| - k \leq \dim(G) \leq |V(G)| - k + m$.

The characteristic of a ring R is the least positive integer n such that $nx = 0$ for all $x \in R$, where 0 is the zero element of R . If no such integer exists, we say that R has characteristic 0 .

Theorem 3.2. Let R be a finite commutative ring that is not a field such that R has odd characteristic. Then $\dim^+(\Gamma(R)) = \dim(\Gamma(R))$.

Proof. Since the characteristic of R is odd, x and $-x$ are distance similar for any vertex x and $x \neq -x$. Thus, by [Theorem 3.1](#), $\dim^+(\Gamma(R)) = \dim(\Gamma(R))$. \square

Theorem 3.3. *Let S be a finite commutative ring of order $2k$, where k is an odd integer. Then $\dim^+(\Gamma(S)) = \dim(\Gamma(S))$.*

Proof. It can be shown that $S \simeq \mathbb{Z}_2 \times R$ for some finite ring R with odd characteristic. If R is a domain, then $\Gamma(S)$ is a star-graph and the result follows from [Theorem 2.4](#). (It is also trivial to prove that $\dim^+(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_p)) = \dim(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_p))$, where p is prime). Hence, we assume that R is not a domain for the rest of the proof.

The elements of $v(\Gamma(S))$ can be partitioned into the sets $A = \{(0, x) | x \in Z(R)^*\}$, $B = \{(0, y) | y \in R - Z(R)\}$, $C = \{(1, x) | x \in Z(R)^*\}$ and $D = \{(1, 0)\}$. Note that $|B| > 1$ since $R - Z(R) \neq \emptyset$ and $z \in R - Z(R)$ implies $-z \in R - Z(R)$ with $z \neq -z$. Also, all elements of B are distance similar as any element of B is only adjacent to $(1, 0)$. If x and y are distance similar vertices of $\Gamma(R)$, then both $(0, x)$ and $(0, y)$ are distance similar in $\Gamma(S)$ and $(1, x)$ and $(1, y)$ are distance similar in $\Gamma(S)$. (For example, $d((a, b), (0, x)) = 1$ if and only if $(a, b)(0, x) = (0, 0)$ if and only if $bx = 0$ if and only if $by = 0$ if and only if $(a, b)(0, y) = (0, 0)$ if and only if $d((a, b), (0, y)) = 1$). Thus, when $v(\Gamma(S))$ is partitioned into distance similar classes (as in [Theorem 3.1](#)), the only class that will have one element is $D = \{(1, 0)\}$.

Next, we show $(1, 0)$ cannot be an element of any minimal resolving set for $\Gamma(S)$. Suppose W is a resolving set with $(1, 0) \in W$. Let $W^* = W - \{(1, 0)\}$. We show that W^* is also a resolving set. Note that $W^* \cap B \neq \emptyset$. So, let $c \in W^* \cap B$. Then, for all $s \in v(\Gamma(S)) - W^*$, $r(s, W^*) \neq r((1, 0), W^*)$ because $(1, 0)$ is the only vertex of $\Gamma(S)$ whose distance to c is 1. So, suppose $x, y \in v(\Gamma(S)) - W$ with $x \neq y$ but $r(x, W^*) = r(y, W^*)$. However, since W is a resolving set, this implies $d((1, 0), x) \neq d((1, 0), y)$ and $d(z, x) = d(z, y)$ for all $z \in W^*$. This is impossible if both x and y are in $A \cup B$, since all elements of $A \cup B$ are distance 1 from $(1, 0)$. If $x, y \in C$, then $x = (1, x_2)$ and $y = (1, y_2)$, for some $x_2, y_2 \in Z(R)^*$. Clearly, $d(x, (1, 0)) > 1$ and $d(y, (1, 0)) > 1$. There must exist some $x_3, y_3 \in Z(R)^*$ such that $x_2x_3 = 0$ and $y_2y_3 = 0$. However, this implies $d(x, (1, 0)) = d(y, (1, 0)) = 2$ via the paths $x - (0, x_3) - (1, 0)$ and $y - (0, y_3) - (1, 0)$. Hence, it must be the case that (without loss of generality) $x \in A \cup B$ and $y \in C$. Thus $y = (1, y_2)$ for some $y_2 \in Z(R)^*$. Suppose $x \in A$ with $x = (0, x_2)$ for some $x_2 \in Z(R)^*$. Then there is some $t \in Z(R)^*$ with $x_2t = 0$. Therefore, there must be some $t^* \in Z(R)^*$ such that t and t^* are distance similar with $(1, t^*) \in W^*$. However, this implies $d(x, (1, t^*)) = 1$ but $d(y, (1, t^*)) \neq 1$. If $x \in B$ with $x = (0, u)$ for some $u \in Z(R)^*$, then there is some $v \in Z(R)^*$ with $uv = 0$. Therefore, there must be some $v^* \in Z(R)^*$ such that v and v^* are distance similar and $(0, v^*) \in W^*$. Then $d(y, (0, v^*)) = 1$ and $d(x, (0, v^*)) \neq 1$. Hence, in all cases, $r(x|W^*) \neq r(y|W^*)$.

Finally, we show that every minimal resolving set of $\Gamma(S)$ must contain all but one element of each distance similar class. Let K_1, K_2, \dots, K_n be the partition of $\Gamma(S)$ into distance similar classes (as in [Theorem 3.1](#)). By [Theorem 3.1](#), $|W \cap K_i| \geq |K_i| - 1$ for any minimal resolving set W . Therefore, assume $|W \cap K_i| = |K_i|$ (that is, $K_i \subseteq W_1$) for some minimal resolving set W_1 and some $1 \leq i \leq n$ with $K_i \neq \{(1, 0)\}$.

Let $x_1 \in K_i$ and let $W^* = W_1 - \{x_1\}$. As in [Theorem 3.1](#), we will show that W_1 is not a minimal resolving set by showing that W^* is a resolving set. Let $a, b \in V(\Gamma(S)) - W_1$. Then $r(a|W_1) \neq r(b|W_1)$, implying there is some $c \in W_1$ with $d(a, c) \neq d(b, c)$. If $c \neq x_1$, then $c \in W^*$ and $r(a|W^*) \neq r(b|W^*)$. If $c = x_1$, then let $v \in K_i$ with $v \neq x_1$. Therefore, v and x_1 are distance similar and $d(a, v) = d(a, x_1) \neq d(b, x_1) = d(b, v)$. Hence, $r(a|W^*) \neq r(b|W^*)$. Finally, if $t \in V(\Gamma(S)) - W^*$ with $t \neq x_1$, then t is not distance similar to x_1 . Thus, there is some vertex $z \in V(\Gamma(S)) - \{t, x_1\}$ such that $d(t, z) \neq d(x_1, z)$. If $z \neq (1, 0)$, there is some $z^* \in W^*$ such that $z = z^*$ or z is distance similar to z^* . Thus, $d(t, z^*) = d(t, z) \neq d(x_1, z) = d(x_1, z^*)$.

However, suppose $z = (1, 0)$. Choose $u \in R - Z(R)$ such that $(0, u) \in W^*$. Then, since the only vertex adjacent to $(0, u)$ is $(1, 0)$, we have $d((0, u), t) = d((0, u), (1, 0)) + d((1, 0), t) = 1 + d((1, 0), t) \neq 1 + d((1, 0), x_1) = d((0, u), (1, 0)) + d((1, 0), x_1) = d((0, u), x_1)$. Therefore, $r(t|W^*) \neq r(x_1|W^*)$. \square

Definition 3.1. A vertex of degree one (that is, a vertex adjacent to only one other vertex) is called a *pendant vertex*. Call the vertices which are adjacent with at least one pendent vertex *star vertices* and the subset of all such vertices a *star subset*. Also, the number of pendent edges incident on v is called the *star degree* of v , denoted by $sdeg(v)$. Clearly, if $d(v)$ denotes the degree of a vertex v , then $sdeg(v) \leq d(v)$ and the equality holds in star graphs. Also, a tree that is not a star graph has at least two star vertices.

Theorem 3.4. Let G be a graph of order n with vertex set $V(G)$ and a star subset $X = \{x_1, x_2, \dots, x_p\}$ such that the star degree of x_i is $k_i \geq 2$ for all $1 \leq i \leq p$. Then $\dim^+(G) \geq k - p$, where $k = k_1 + k_2 + \dots + k_p$.

Proof. For $1 \leq i \leq p$, choose x_i , and let v_1, v_2, \dots, v_{k_i} , $k_i > 1$ be pendent vertices incident on x_i . Then v_m is distance similar to v_j for each $1 \leq m \leq k_i$ and $1 \leq j \leq k_i$. Therefore, by Theorem 3.1, a subset of at least $k_i - 1$ of the vertices $\{v_1, v_2, \dots, v_{k_i}\}$ must be contained in any minimal resolving set. Thus any resolving set has cardinality greater than or equal to $k_1 + k_2 + \dots + k_p - p = k - p$. \square

Corollary 3.5. Let G be a graph as in Theorem 3.4. If, in addition, G is the zero divisor graph of some commutative ring, then $k - p \leq \dim^+(G) \leq n - p$, where $k = k_1 + k_2 + \dots + k_p$.

Example 3.6. By using the results in this article and examining the graphs found in [11], one can determine the upper dimension of all zero divisor graphs of a commutative ring with up to 14 vertices. Out of these examples, there is only one ring R such that $\dim(\Gamma(R)) \neq \dim^+(\Gamma(R))$. For $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, $\dim(\Gamma(R)) = 3$ with $W = \{(1, 1, 1, 0), (1, 1, 0, 1), (1, 0, 1, 1)\}$ an example of a minimal resolving set. However, it is straightforward to verify that $V = \{(1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1)\}$ also defines a resolving set for $\Gamma(R)$. We can see that V is minimal, as removing $(1,0,0,0)$ would give $r((1, 1, 1, 0)|V) = r((0, 1, 1, 0)|V)$, removing $(0,1,0,0)$ will give $r((1, 1, 1, 0)|V) = r((1, 0, 1, 0)|V)$, removing $(0,0,1,0)$ will give $r((1, 1, 1, 0)|V) = r((1, 1, 0, 0)|V)$, and removing $(0,0,0,1)$ will give $r((1, 1, 0, 1)|V) = r((1, 1, 0, 0)|V)$.

It is also interesting to note that for most of the zero divisor graphs on 14 or fewer vertices, a minimal resolving set can be determined by looking at classes of distance similar vertices — in particular, if K_1, K_2, \dots, K_p is a partition of $v(\Gamma(R))$ into distance similar sets of vertices, then a minimal resolving set is formed by taking all but one element from each K_i . The only exceptions are as follows.

- (i) for $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ or \mathbb{Z}_9 or $\mathbb{Z}_3[x]/(x^2)$, (where $\Gamma(R) \cong K_2$ and vacuously no distinct vertices of $\Gamma(R)$ are distance similar).
- (ii) for $R \cong \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$ or $\mathbb{Z}_2 \times \mathbb{Z}_4$, when $\Gamma(R)$ has only one pair of distance similar vertices but $\dim(\Gamma(R)) = \dim^+(\Gamma(R)) = 2$.
- (iii) for $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, when no two distinct vertices of $\Gamma(R)$ are distance similar.
- (iv) for $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, when no two distinct vertices of $\Gamma(R)$ are distance similar.

Acknowledgments

The authors are grateful to the anonymous referee for his useful suggestions which improved the presentation of the paper. This research is supported by JRF financial assistance (second author) by Council of Scientific and Industrial Research (CSIR), New Delhi India and the University Grants Commission, New Delhi with research project number MRP-MAJOR-MATH-2013-8034 (first author).

References

- [1] D.F. Anderson, P.S. Livingston, The zero-divisor graph of a commutative ring, *J. Algebra* 217 (1999) 434–447.
- [2] I. Beck, Coloring of commutative rings, *J. Algebra* 116 (1988) 208–226.
- [3] D.D. Anderson, M. Naseer, Beck's coloring of a commutative ring, *J. Algebra* 159 (1993) 500–517.
- [4] P.J. Slater, Leaves of trees, *Congr. Number.* 14 (1975) 549–559.
- [5] F. Harary, R.A. Melter, On the metric dimension of a graph, *Ars Combin.* 2 (1976) 191–195.
- [6] G. Chartrand, C. Poisson, P. Zhang, Resolvability and the upper dimension of graphs, *Int. J. Comput. Math. Appl.* 39 (2000) 19–28.
- [7] S. Pirzada, An Introduction to Graph Theory, Universities Press, Orient Blackswan, Hyderabad, India, 2012.
- [8] S. Pirzada, M. Aijaz, S.P. Redmond, On the metric and upper dimension of some graphs (submitted for publication).
- [9] S. Pirzada, Rameez Raja, S.P. Redmond, Locating sets and numbers of graphs associated to commutative rings, *J. Algebra Appl.* 13 (7) (2014) 1450047 (18 pp).
- [10] Rameez Raja, S. Pirzada, S.P. Redmond, On locating numbers and codes of zero-divisor graphs associated with commutative rings, *J. Algebra Appl.* 15 (1) (2016) 1650014 (22 pp).
- [11] S.P. Redmond, On zero divisor graphs of small finite commutative rings, *Discrete Math.* 307 (2007) 1155–1166.