



The upper domatic number of a graph

Teresa W. Haynes, Jason T. Hedetniemi, Stephen T. Hedetniemi, Alice McRae & Nicholas Phillips

To cite this article: Teresa W. Haynes, Jason T. Hedetniemi, Stephen T. Hedetniemi, Alice McRae & Nicholas Phillips (2020) The upper domatic number of a graph, AKCE International Journal of Graphs and Combinatorics, 17:1, 139-148, DOI: [10.1016/j.akcej.2018.09.003](https://doi.org/10.1016/j.akcej.2018.09.003)

To link to this article: <https://doi.org/10.1016/j.akcej.2018.09.003>



© 2018 Kalasalingam University. Published with license by Taylor & Francis Group, LLC.



Published online: 01 Jun 2020.



Submit your article to this journal [↗](#)



Article views: 92



View related articles [↗](#)



View Crossmark data [↗](#)



The upper domatic number of a graph

Teresa W. Haynes^{a,b,*}, Jason T. Hedetniemi^c, Stephen T. Hedetniemi^d, Alice McRae^e,
Nicholas Phillips^e

^a Department of Mathematics and Statistics, East Tennessee State University, Johnson City, TN 37614-0002, USA

^b Department of Mathematics, University of Johannesburg, Auckland Park, South Africa

^c Department of Mathematics, Wingate University, Wingate, NC 28174, USA

^d Professor Emeritus, School of Computing, Clemson University, Clemson, SC 29634, USA

^e Department of Computer Science, Appalachian State University, Boone, NC 28608, USA

Received 8 June 2018; received in revised form 13 September 2018; accepted 14 September 2018

Abstract

Let $G = (V, E)$ be a graph. For two disjoint sets of vertices R and S , set R dominates set S if every vertex in S is adjacent to at least one vertex in R . In this paper we introduce the *upper domatic number* $D(G)$, which equals the maximum order k of a vertex partition $\pi = \{V_1, V_2, \dots, V_k\}$ such that for every $i, j, 1 \leq i < j \leq k$, either V_i dominates V_j or V_j dominates V_i , or both. We study properties of the upper domatic number of a graph, determine bounds on $D(G)$, and compare $D(G)$ to a related parameter, the transitivity $Tr(G)$ of G .

Keywords: Domination; Domatic number; Upper domatic number; Transitivity

1. Introduction

Partitioning the vertex set of a graph into subsets with a specified property is a popular subject in graph theory. Arguably the most famous problem of graph partitioning is graph coloring, where the vertex set is partitioned into independent sets. The minimum order of such a partition is the well-studied *chromatic number* of a graph. Considering subsets that are dominating sets, Cockayne and Hedetniemi [1] in 1977 defined the *domatic number* to be the maximum order of a partition of the vertices of a graph into dominating sets.

More formally, a set $S \subseteq V(G)$ is a *dominating set* of a graph $G = (V, E)$ if every vertex in $V(G) - S$ is adjacent to a vertex in S . The *domination number* of G , denoted $\gamma(G)$, is the minimum cardinality of a dominating set of G . A dominating set with cardinality $\gamma(G)$ is called a γ -set of G . Given two disjoint sets of vertices $R, S \subset V$, we

Peer review under responsibility of Kalasalingam University.

* Corresponding author at: Department of Mathematics and Statistics, East Tennessee State University, Johnson City, TN 37614-0002, USA.

E-mail addresses: haynes@etsu.edu (T.W. Haynes), j.hedetniemi@wingate.edu (J.T. Hedetniemi), hedet@clemson.edu (S.T. Hedetniemi), mcræaa@appstate.edu (A. McRae), nickphillips@gmail.com (N. Phillips).

<https://doi.org/10.1016/j.akcej.2018.09.003>

© 2018 Kalasalingam University. Published with license by Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

say that R dominates S , denoted $R \rightarrow S$, if every vertex in S is adjacent to, or dominated by, at least one vertex in R . As previously mentioned, the *domatic number* of a graph G , denoted $d(G)$, is the maximum order k of a vertex partition $\pi = \{V_1, V_2, \dots, V_k\}$ such that for every i , $1 \leq i \leq k$, V_i is a dominating set of G . We call a partition π into dominating sets a *domatic partition* and a domatic partition of order $d(G)$ is called a *d-partition* of G . Well over 100 papers have been published on this concept and its variations. The bibliography contains a sampling of these papers. For examples, cf. [2–9].

An equivalent definition of the domatic number is that it equals the maximum order k of a vertex partition $\pi = \{V_1, V_2, \dots, V_k\}$ such that for all i, j , $1 \leq i < j \leq k$, $V_i \rightarrow V_j$ and $V_j \rightarrow V_i$. This definition of the domatic number suggests the following straightforward generalization. The *upper domatic number*, denoted $D(G)$, equals the maximum order k of a vertex partition $\pi = \{V_1, V_2, \dots, V_k\}$ such that for all i, j , $1 \leq i < j \leq k$, either $V_i \rightarrow V_j$ or $V_j \rightarrow V_i$, or both. A vertex partition π meeting this condition is called an *upper domatic partition* and an upper domatic partition of order $D(G)$ is called a *D-partition* of G .

In 2017, J. T. Hedetniemi and S. T. Hedetniemi [10] gave a different generalization of the domatic number as follows. The *transitivity of a graph* $Tr(G)$ is the maximum order k of a vertex partition $\pi = \{V_1, V_2, \dots, V_k\}$ such that for all i, j , $1 \leq i < j \leq k$, $V_i \rightarrow V_j$. A vertex partition of V meeting this condition is called a *transitive partition* and a transitive partition of G having order $Tr(G)$ is called a *Tr-partition* of G .

By definition, therefore, every domatic partition is both a transitive partition and an upper domatic partition, while every transitive partition is an upper domatic partition. Thus, we have the following basic inequalities for any graph G .

Proposition 1. For any graph G of order n , $1 \leq d(G) \leq Tr(G) \leq D(G) \leq n$.

Our aim in this paper is to introduce and study the upper domatic number of a graph. We present properties and bounds on $D(G)$, and compare the upper domatic number and transitivity of graphs G .

1.1. Notation and terminology

In general, we follow the notation and terminology in [11]. Let $G = (V, E)$ be a graph. The *open neighborhood* of a vertex $v \in V$ is the set $N(v) = \{u \mid uv \in E\}$. Each vertex $u \in N(v)$ is called a *neighbor* of v , and $|N(v)|$ is called the *degree* of v , denoted $\deg(v)$. The minimum and maximum degrees of vertices in a graph G are denoted $\delta(G)$ and $\Delta(G)$, respectively. The *closed neighborhood* of a vertex $v \in V$ is the set $N[v] = N(v) \cup \{v\}$. The *open neighborhood of a set* $S \subseteq V$ of vertices is $N(S) = \bigcup_{v \in S} N(v)$, while the *closed neighborhood of a set* S is the set $N[S] = \bigcup_{v \in S} N[v]$. Let $G[S]$ denote the subgraph induced by S in G .

A *cycle* and *path* on n vertices are denoted by C_n and P_n , respectively. The *complete graph* on n vertices is denoted by K_n , and its *complement*, the empty graph on n vertices, is denoted \overline{K}_n . A *clique* is a maximal complete subgraph of G , and the *clique number* $\omega(G)$ is the maximum cardinality of a clique of G .

Let $\pi = \{V_1, V_2, \dots, V_k\}$ be a vertex partition of a graph $G = (V, E)$. A subset V_i with $|V_i| = 1$ is called a *singleton*, and a subset $|V_i| \geq 2$ is called a *non-singleton*.

1.2. Domination digraphs

An equivalent definition of domatic partitions can be given using domination digraphs. Let $\pi = \{V_1, V_2, \dots, V_k\}$ be a vertex partition of a graph $G = (V, E)$. From this partition one can construct a directed graph, or digraph, $DG(\pi) = (\pi, E(\pi))$, the vertices of which correspond one-to-one with the sets V_1, V_2, \dots, V_k of π , with a directed arc $V_i \rightarrow V_j$ in $DG(\pi)$ if V_i dominates V_j in G , that is, $V_j \subseteq N(V_i)$. We call $DG(\pi)$ the *domination digraph* of π . Domination digraphs were introduced in 2012 by Goddard et al. [12]. A digraph $DG = (V, E)$ is called *complete* if for any two vertices $v_i, v_j \in V$, either $v_i \rightarrow v_j$ or $v_j \rightarrow v_i$, or both. A complete digraph is called *bi-directed* if for any two vertices v_i and v_j , both $v_i \rightarrow v_j$ and $v_j \rightarrow v_i$. A complete digraph $DG = (V, E)$ is called a *tournament* if for any two vertices v_i and v_j , either $v_i \rightarrow v_j$ or $v_j \rightarrow v_i$, but not both. A tournament $DG = (V, E)$ is called *transitive* if the vertices can be ordered v_1, v_2, \dots, v_n in such a way that for every $1 \leq i < j \leq n$, $v_i \rightarrow v_j$ if and only if $i < j$. A (complete) digraph DG of order n is said to be *transitive* if it contains a transitive tournament of order n as a sub-digraph, that is, DG contains a spanning transitive tournament.

A set V_i in a partition $\pi = \{V_1, V_2, \dots, V_k\}$ is called a *source set* if the out-degree of vertex V_i in its domination digraph $DG(\pi)$ is $k - 1$. And the set V_i is a *sink set* if the in-degree of vertex V_i in $DG(\pi)$ is $k - 1$. Note that if $\pi = \{V_1, V_2, \dots, V_k\}$ is a transitive partition, then V_1 is a source set and V_k is a sink set.

Using domination digraphs, we can give equivalent definitions of transitive, domatic, and upper domatic partitions. A vertex partition π is *transitive* if its domination digraph is transitive, and is an *upper domatic partition* if its domination digraph is complete. The *domatic number* of a graph is the maximum integer k such that G has a vertex partition $\pi = \{V_1, V_2, \dots, V_k\}$ whose domination digraph is a bi-directed complete graph.

2. The upper domatic number

2.1. Basic properties

We begin with some basic properties that will prove useful in the results that follow.

Proposition 2. *If $\pi = \{V_1, V_2, \dots, V_r\}$ is an upper domatic partition of a graph G , then the partition $\pi' = \{V_1, V_2, \dots, V_{i-1}, V_i \cup V_j, V_{i+1}, \dots, V_{j-1}, V_{j+1}, \dots, V_r\}$ obtained from π by deleting V_i and V_j and adding to π' the union $V_i \cup V_j$, for any $1 \leq i < j \leq r$, is also an upper domatic partition.*

Proof. In the vertex partition π' , we only need to show that for any $V_k, V_k \neq V_i \cup V_j$, either $V_k \rightarrow (V_i \cup V_j)$ or $(V_i \cup V_j) \rightarrow V_k$. There are essentially two cases: (i) $V_k \rightarrow V_i$ and $V_k \rightarrow V_j$, or (ii) either $V_i \rightarrow V_k$ or $V_j \rightarrow V_k$ or both. For (i), $V_k \rightarrow (V_i \cup V_j)$, and for any of the three possibilities of (ii), $(V_i \cup V_j) \rightarrow V_k$. \square

Corollary 3. *If a graph G has an upper domatic partition of order k , then it has an upper domatic partition of order j , for any $1 \leq j \leq k$.*

The following observations are straightforward.

Observation 4. *Let $\pi = \{V_1, V_2, \dots, V_k\}$ be an upper domatic partition of a graph G .*

1. *For each $V_i, V_j, i \neq j$, if $V_i \rightarrow V_j$, there are at least $|V_j|$ edges in G between vertices in V_i and V_j .*
2. *For each $V_i, V_j, i \neq j$, there are at least $\min\{|V_i|, |V_j|\}$ edges in G between vertices in V_i and V_j .*
3. *If vertices v_1, v_2, \dots, v_k are each in singleton sets in π , then the vertices v_1, v_2, \dots, v_k form a complete subgraph in G .*
4. *If a vertex $u \in V_i$ is not adjacent to any vertex in V_j , then $V_i \rightarrow V_j$.*
5. *If $u \in V_i$ with $\deg(u) = x < k$, then $\text{outdeg}(V_i) \geq k - x - 1$ in $DG(\pi)$. Further, if $V_i \rightarrow V_j$ and u is adjacent to a vertex in V_j for some j , then $\text{outdeg}(V_i) \geq k - x$ in $DG(\pi)$.*

It was shown in [10] that the transitivity of any graph G is at least as large as the transitivity of any subgraph of G .

Proposition 5 ([10]). *If H is any subgraph of a graph G , then $Tr(H) \leq Tr(G)$.*

This is not necessarily true for the upper domatic number. To see this, consider the bipartite graph G in Fig. 1. Let H be the induced subgraph $G - v_{11}$, and note that $\pi = \{V_1 = \{v_1, v_2, v_3\}, V_2 = \{v_4, v_5, v_6\}, V_3 = \{v_7, v_8, v_9\}, V_4 = \{v_{10}\}\}$ is an upper domatic partition of H in which $V_1 \rightarrow V_2 \rightarrow V_3 \rightarrow V_1$ and V_4 is a sink set. Thus, $D(H) \geq 4$, and it is straightforward to show that $D(H) = 4$.

The graph G , on the other hand, satisfies $D(G) < 4$. This follows by observing that since G has an isolated vertex v_{11} , any upper domatic partition of G of order 4, $\pi = \{V_1, V_2, V_3, V_4\}$, must have a source set V_1 containing vertex v_{11} . This means that the set $V_1 - \{v_{11}\}$ is a dominating set of H . However, it can be seen that the domination number of H is 3, and that the vertices of any dominating set are incident to at least 8 of the 12 edges. Thus, one can check that there is no upper domatic partition of G of order 4 containing such a source set.

The following relaxed form of Proposition 5, however, does hold for the upper domatic number.

Proposition 6. *If H is any subgraph of a graph G and H has a D -partition with a source set, then $D(H) \leq D(G)$.*




Fig. 1. $D(H) = 4 > D(G) = 3$, where $H = G - v_{11}$.

Proof. To get an upper domatic partition for G , use the D -partition for H and add the vertices in $V(G) - V(H)$ to the source set in the D -partition for H . \square

Corollary 7. For any graph G , $D(G) \geq \omega(G)$.

We note that the bound of Corollary 7 is sharp for complete graphs K_n , for which $\omega(K_n) = n = D(K_n)$. On the other hand, the difference $D(G) - \omega(G)$ can be arbitrarily large; for complete bipartite graphs $K_{k,k}$, $\omega(K_{k,k}) = 2 < D(K_{k,k}) = k + 1$. Note that an upper domatic partition of order $k + 1$ can be formed by placing one vertex from each partite set into singleton sets and adding $k - 1$ dominating sets containing one vertex from each partite set.

2.2. Small values of the upper domatic number

In this section we characterize the families of graphs G having upper domatic number equal to 1, 2, or at least 3. A *star* is a tree having at most one vertex of degree greater than one. A *galaxy* is a disjoint union of stars.

Theorem 8. For any graph G ,

- (1) $D(G) \geq 3$ if and only if G contains either a K_3 or a P_4 ,
- (2) $D(G) = 2$ if and only if G is a galaxy with at least one edge, and
- (3) $D(G) = 1$ if and only if $G = \overline{K}_n$.

Proof. (1) Assume that G contains a triangle or a P_4 subgraph. Each of these subgraphs has upper domatic number 3 and a D -partition with a source set. Thus, Proposition 6 implies that $D(G) \geq 3$.

Conversely, assume that $D(G) = k \geq 3$. Let $\pi = \{V_1, V_2, \dots, V_k\}$ be an upper domatic partition of G . Consider only V_1, V_2 , and V_3 . Since π is an upper domatic partition, it follows that either $V_i \rightarrow V_j$ or $V_j \rightarrow V_i$, or both, for every $1 \leq i < j \leq 3$. From this we can assume either (i) the existence of a *cyclic triple*, where, without loss of generality, $V_1 \rightarrow V_2, V_2 \rightarrow V_3$ and $V_3 \rightarrow V_1$, or (ii) the existence of a *transitive triple*, where $V_1 \rightarrow V_2, V_1 \rightarrow V_3$, and $V_2 \rightarrow V_3$.

If (i) holds, let v_1 be an arbitrary vertex in V_1 . Since $V_3 \rightarrow V_1$, there must be a vertex $v_3 \in V_3$, which is adjacent to v_1 . Similarly, since $V_2 \rightarrow V_3$, there must be a vertex $v_2 \in V_2$ which is adjacent to vertex v_3 . Finally, since $V_1 \rightarrow V_2$, there must be a vertex $v'_1 \in V_1$ which is adjacent to v_2 . If $v_1 = v'_1$, then G has a triangle $\{v_1, v_2, v_3\}$, but if $v_1 \neq v'_1$, then G has a P_4 subgraph (v_1, v_3, v_2, v'_1) .

If (ii) holds, let $v_3 \in V_3$. Since $V_2 \rightarrow V_3$, there must exist a vertex $v_2 \in V_2$ which is adjacent to v_3 . Since $V_1 \rightarrow V_2$, there is a vertex $v_1 \in V_1$ which is adjacent to v_2 . Finally, since $V_1 \rightarrow V_3$, there is a vertex $v'_1 \in V_1$, which is adjacent to v_3 . If $v'_1 = v_1$, then $\{v_1, v_2, v_3\}$ forms a triangle in G , but if $v'_1 \neq v_1$, then (v_1, v_2, v_3, v'_1) is a P_4 subgraph of G .

(2) Let $G = (V, E)$ be a galaxy with at least one edge, say uv . Then the partition defined by $\{V - \{u\}, \{u\}$ is an upper domatic partition, and so $D(G) \geq 2$. But since G is a galaxy, it does not contain a K_3 or a P_4 . Hence, from (1) we know that $D(G) < 3$. Thus, $D(G) = 2$.

Conversely, assume that $D(G) = 2$. Then by definition, G has at least one edge. By (1) we know that G has no triangle and no P_4 subgraph. Any trivial component of G is the star K_1 . Since G has at least one edge, G has at least one nontrivial component. Consider any nontrivial component H of G (if G is connected $H = G$), and let u be a vertex of H having maximum degree $\Delta(H)$. Since G has no triangles, $N(u)$ is an independent set. If u has degree 1, then since u has maximum degree, H is the star K_2 . Assume that u has degree 2 or more. If a neighbor of u , say v , has a neighbor $z \notin N[u]$, then (w, u, v, z) , where $w \in N(u)$ and $w \neq v$ is a P_4 subgraph of G , a contradiction. Hence, every neighbor of u has degree 1, so H is a star, implying that G is a galaxy.

(3) Trivially, $D(\overline{K}_n) = 1$. Assume that $D(G) = 1$. If $G \neq \overline{K}_n$, then G has at least one edge, say uv . Thus, the partition defined by $\{\{u\}, V - \{u\}\}$ is an upper domatic partition, and so, $D(G) \geq 2$, a contradiction. Thus, $G = \overline{K}_n$. \square

3. Bounds on the upper domatic number

For a set of vertices S , let $G - S$ be the graph formed from G by removing the set S from G .

Proposition 9. *If G is a graph with order n and S is a γ -set of G , then*

$$D(G) \geq D(G - S) + 1.$$

Proof. Since S along with the sets of a D -partition of $G - S$ is an upper domatic partition of G , $D(G) \geq D(G - S) + 1$. \square

By Proposition 1, we have that $d(G) \leq D(G)$. Proposition 9 suggests a way of improving this bound for graphs with a d -partition containing a set that is not independent.

Proposition 10. *If $\pi = \{V_1, V_2, \dots, V_k\}$ is a d -partition of G , with $G_i = G[V_i]$ for $1 \leq i \leq k$, then $D(G) \geq d(G) - 1 + \max\{D(G_i) : 1 \leq i \leq k\}$.*

Proof. Let $\pi = \{V_1, V_2, \dots, V_k\}$ be a d -partition of G , and let $G_i = G[V_i]$. Let $D(G_j) = \max\{D(G_i) : 1 \leq i \leq k\}$ and let π' be a D -partition of G_j . We note that $(\pi - \{V_j\}) \cup \pi'$ is an upper domatic partition of G . Thus, $D(G) \geq d(G) - 1 + D(G[V_j])$. \square

Corollary 11. *If G is a graph with $d(G) = D(G)$, then every d -partition of G is a partition of the vertices of G into independent dominating sets.*

Proof. Let π be a d -partition of G , and let V_j be a set in π that is not independent. Using the notation of Proposition 10, $D(G) \geq d(G) - 1 + D(G_j)$. Since V_j is not an independent set, Theorem 8 implies that $D(G_j) \geq 2$. Thus, $D(G) \geq d(G) - 1 + D(G_j) \geq d(G) - 1 + 2 = d(G) + 1$. \square

The following upper bound on the domatic number is well known.

Proposition 12 ([1]). *For any graph G , $d(G) \leq \delta(G) + 1$.*

This bound does not hold for $D(G)$. It is immediately obvious that for any complete graph K_n , the domatic number $d(K_n) = D(K_n) = n = \delta(G) + 1$. Consider, however, the graph G formed by adding a new vertex u adjacent to exactly one vertex, say v , of a complete graph K_n . It is easy to see that $d(G) = 2 = \delta(G) + 1$, while $D(G) = n$, since a set containing $\{u, v\}$ along with $n - 1$ singleton sets, each containing a vertex of $K_n - \{u, v\}$, is a D -partition of G . Thus, for $n \geq 3$, $D(G) > 2 = \delta(G) + 1$, and we see that a significant difference can exist between the domatic number and the upper domatic number. With this example, we have proved the following.

Proposition 13. *For any positive integer k , there is a graph G such that $D(G) - d(G) \geq k$, and such that $D(G) - (\delta(G) + 1) \geq k$.*

Although the bound involving minimum degree in Proposition 12 does not hold for the upper domatic number, we show that replacing minimum degree with maximum degree is an upper bound for $D(G)$.

Theorem 14. For any graph G , $D(G) \leq \Delta(G) + 1$.

Proof. Let $\pi = \{V_1, V_2, \dots, V_{k+1}\}$ be a D -partition of G ordered in such a way, that $|V_i| \geq |V_j|$ for $i < j$. We claim there is at least one vertex $v \in V_{k+1}$ with $\deg(v) \geq k$. By [Observation 4](#), it follows there must be at least $|V_{k+1}|$ edges between each of the sets V_i and V_{k+1} , for $1 \leq i \leq k$. Hence, the sum of the degrees of the vertices in V_{k+1} is at least $k|V_{k+1}|$, and therefore, the average degree of the vertices in this set is at least k . Hence, $D(G) = k+1 \leq \Delta(G)+1$. \square

It is shown in [\[1\]](#) that $d(G) + d(\overline{G}) \leq n + 1$. Our Nordhaus–Gaddum type bounds for the upper domatic number follow from [Theorem 14](#).

Corollary 15. Let G be a graph of order n . Then

$$D(G) + D(\overline{G}) \leq n + \Delta(G) - \delta(G) + 1.$$

Proof. [Theorem 14](#) implies that $D(G) + D(\overline{G}) \leq \Delta(G) + 1 + \Delta(\overline{G}) + 1$. The result follows since $\Delta(\overline{G}) = n - \delta(G) - 1$. \square

We note that the bound of [Corollary 15](#) is obtained by the self-complementary graph $G = P_4$ for which $D(G) + D(\overline{G}) = 3 + 3 = 4 + 2 - 1 + 1 = n + \Delta(G) - \delta(G) + 1$. It is also obtained the complete graph.

Corollary 16. If G is a regular graph, then $D(G) + D(\overline{G}) \leq n + 1$.

An upper bound on the sum of the upper domatic number and the domination number of a graph also follows from [Theorem 14](#). It is a well-known result (see [\[11\]](#)) that for graphs with order n , $\gamma(G) \leq n - \Delta(G)$.

Corollary 17. If G is a graph with order n , then $D(G) + \gamma(G) \leq n + 1$.

The following proposition was given in [\[10\]](#).

Proposition 18 ([\[10\]](#)). For any connected graph G with $\delta(G) = 3$, $Tr(G) \geq 4$.

Since $Tr(G) \leq D(G)$, we have the following corollary.

Corollary 19. For any connected graph G with $\delta(G) = 3$, $D(G) \geq 4$.

Based on [Corollary 19](#), one might think that for any graph G , $D(G) \geq \delta(G) + 1$. However, this is not the case. The following examples show that $D(G) < \delta(G)$ is possible. Indeed, the difference $\delta(G) - D(G)$ can be arbitrarily large.

For integers $t \geq 2$ and $k \geq 2$, let $G_{t,k}$ be the complete multipartite graph with t partite sets each having cardinality k . That is, $G_{t,k} = K_{\{k,k,\dots,k\}}$ with t partite sets.

Proposition 20. For the graph $G_{t,k}$ with t even, $D(G_{t,k}) = \frac{kt+t}{2}$.

Proof. Let the partite sets be U_1, U_2, \dots, U_t and let $U_i = \{u_{i,1}, u_{i,2}, \dots, u_{i,k}\}$ for $1 \leq i \leq t$. Note that for any $1 \leq i < j \leq t$, $\{u_i\} \rightarrow \{u_j\}$ and conversely, $\{u_j\} \rightarrow \{u_i\}$. Moreover, every two-element subset containing elements from different partite sets is a dominating set of $G_{t,k}$. That is, the partition $\pi = \{\{u_{1,1}\}, \{u_{2,1}\}, \dots, \{u_{t,1}\}, \{u_{1,2}, u_{2,2}\}, \{u_{3,2}, u_{4,2}\}, \dots, \{u_{t-1,2}, u_{t,2}\}, \dots, \{u_{1,k}, u_{2,k}\}, \{u_{3,k}, u_{4,k}\}, \dots, \{u_{t-1,k}, u_{t,k}\}\}$ is an upper domatic partition. Thus, it follows that for this graph, $D(G) \geq t + \frac{(k-1)t}{2} = \frac{kt+t}{2}$.

To see that $D(G) \leq \frac{kt+t}{2}$, observe that any upper domatic partition of order larger than $\frac{kt+t}{2}$ has at least $t + 1$ singleton sets. But then at least two of these singleton sets are vertices in the same partite set and are not adjacent, contradicting [Observation 4](#). \square

It follows from [Proposition 20](#) that there exists a graph G for which $D(G) = \delta(G)$, and there is a graph G for which $D(G) < \delta(G)$. For examples, the graph $G_{4,2}$ has $D(G_{4,2}) = \delta(G_{4,2}) = 6$, and the graph $G_{6,2}$ has $D(G_{6,2}) = 9 < \delta(G_{6,2}) = 10$.

4. Upper domatic number versus transitivity

In this section, we first consider graphs G having equal transitivity and upper domatic number. We then turn our attention to the differences in these two parameters.

By considering [Theorem 8](#) together with the following two results proven in [10], we see that graphs G satisfying $D(G) \leq 3$ have equal upper domatic and transitivity numbers.

Theorem 21 ([10]). *Let G be a graph.*

1. $Tr(G) = 1$ if and only if $G = \overline{K}_n$.
2. $Tr(G) = 2$ if and only if G is a galaxy with at least one edge.

Proposition 22 ([10]). *For any graph G , $Tr(G) \geq 3$ if and only if G contains either a triangle or a P_4 subgraph (not necessarily induced).*

We now consider other examples of graph families having this property. Recall that P_n and C_n denote the path and cycle, of order n , respectively. The Cartesian product $G \square H$ of two graphs G and H is the graph with vertex set $V(G) \times V(H)$ and edges such that two vertices (u, v) and (w, x) are adjacent in $E(G \square H)$ if and only if either $u = w$ and v is adjacent to x in H , or u is adjacent to w in G and $v = x$. The class of graphs called n -cubes is defined recursively as follows. The 0-cube, denoted Q_0 , is the graph K_1 . The n -cube Q_n is the graph $Q_n = Q_{n-1} \square P_2$.

Using the fact that $Tr(G) \leq D(G) \leq \Delta(G) + 1$ ([Theorem 14](#)) and proofs similar to those in [13,10], we can easily prove the following.

Theorem 23.

1. For the path P_n , $n \geq 4$, $Tr(P_n) = D(P_n) = \Delta(P_n) + 1 = 3$.
2. For the cycle C_n , $n \geq 3$, $Tr(C_n) = D(C_n) = \Delta(C_n) + 1 = 3$.
3. For any cubic graph G , $Tr(G) = D(G) = \Delta(G) + 1 = 4$.
4. For $m, n \geq 4$, $Tr(P_m \square P_n) = D(P_m \square P_n) = \Delta(P_m \square P_n) + 1 = 5$.
5. For $m, n \geq 3$, $Tr(C_m \square C_n) = D(C_m \square C_n) = \Delta(C_m \square C_n) + 1 = 5$.
6. For any positive integer n , $Tr(Q_n) = D(Q_n) = \Delta(Q_n) + 1 = n + 1$.

[Theorem 23](#) gives several families of graphs having equal transitivity and upper domatic numbers. Acyclic graphs are another family with this property.

Theorem 24. *If G is acyclic, then $D(G) = Tr(G)$.*

Proof. Let G be any acyclic graph, and let $\pi = \{V_1, V_2, \dots, V_k\}$ be an upper domatic partition of cardinality k for G . We show that the sets in π can be reordered V'_1, V'_2, \dots, V'_k , so that for $i < j$, V'_i dominates V'_j . We first show that π contains a source set. Let T_r be a tree in G rooted at r . We consider two cases:

Case (i). Let $r \in V_i$. For Case (i), if c is a child of r , then either $c \in V_i$ or V_i dominates the set of π containing c . Thus, V_i dominates all the sets of π containing vertices in $N[r]$. Suppose there is some set $V_j \in \pi$ that is not dominated by V_i . Then no child of r is in V_j , implying that V_j does not dominate V_i , contradicting that π is an upper domatic partition of G . Hence, V_i is a source set in π , and we can let $V'_1 = V_i$.

Case (ii). If Case (i) does not hold, let u be a vertex of maximum depth in T_r , such that $\text{parent}(u)$'s set, say V_i , is different from the set V_j containing u and V_i does not dominate V_j . Since Case (i) does not hold, at least one child of r has this property, so the vertex u exists. Moreover, since π is an upper domatic partition, $V_j \rightarrow V_i$. Also note by the choice of u , if u has a child, it is either in V_j or V_j dominates the child's set. Again, if there is a set say V_a such that V_j does not dominate V_a , then V_a has no vertex in $N(u)$. But then V_j does not dominate V_a and V_a does not dominate V_j , contradicting that π is an upper domatic partition. Thus, V_j is a source set in π .

Therefore, there exists a source set in π . We let this set be the first set V'_1 in a reordered π . We can create a new acyclic graph $G' = G - V'_1$. The partition $\pi' = \pi - V'_1$ will be an upper domatic partition for G' . By repeating the above process, we can find the next set in the ordering of π , and after $k - 1$ iterations the entire ordering. \square




Fig. 2. $Tr(G_i) < D(G_i)$ for $i = 1, 2$.

We have seen several examples where the inequality $Tr(G) \leq D(G)$ is tight. On the other hand, strict inequality is possible. For example, the graph G_1 in Fig. 2 has $4 = Tr(G_1) < D(G_1) = 5$. To see that $D(G_1) = 5$, we note that $\{\{u_1, u_2\}, \{u_3, u_4\}, \{u_5, u_6\}, \{u_7\}, \{u_8\}\}$ is an upper domatic partition, so $D(G_1) \geq 5$, and Theorem 14 implies that $D(G_1) \leq 5$. It can be verified by a straightforward, but detailed proof or by a computer search that G_1 is a graph with the fewest vertices having transitivity strictly less than its upper domatic number. Similarly, it can be proved that the bipartite graph G_2 in Fig. 2, for which $3 = Tr(G_2) < D(G_2) = 4$, is a graph with the fewest edges having this property. We note that Propositions 22 and 25 imply that $Tr(G_2) = 3$. From Theorem 14, $D(G_2) \leq 4$. The partition $\{\{v_1, v_2, v_3\}, \{v_4, v_5, v_6\}, \{v_7, v_8, v_9\}, \{v_{10}\}\}$ is an upper domatic partition, and so, $D(G_2) = 4$.

We next show that the difference between $D(G)$ and $Tr(G)$ can be made arbitrarily large. We will need the following result from [10].

Proposition 25 ([10]). *If G is a graph with $Tr(G) = k$, then there exist two adjacent vertices in G , each having degree $k - 1$ or more.*

Theorem 26. *There exists a graph G for which $D(G) - Tr(G)$ differ by at least p , for any positive integer p .*

Proof. Let $n = 2k + 1$ for some positive integer $k > 1$. We construct a bipartite graph G with $n(k + 1)$ vertices and $nk(k + 1)$ edges as follows.

Let $V(G) = \{s_0, s_1, s_2, \dots, s_{n-1}\} \cup \{x_{i,j} \mid 0 \leq i \leq n - 1, 1 \leq j \leq k\}$. For $0 \leq i \leq n - 1$, let $X_i = \{x_{i,j} \mid 1 \leq j \leq k\}$ and $V_i = \{s_i\} \cup X_i$.

For $0 \leq i \leq n - 1$, add edges such that s_i is adjacent to every vertex in X_p where p is the sum $i + j$ modulo n for $1 \leq j \leq k$. Finally, add the edges $s_i x_{a,b}$ such that $a + b \equiv i \pmod{n}$. Now G is a bipartite graph with partite sets $S = \{s_i \mid 0 \leq i \leq n - 1\}$ and $X = \bigcup X_i$ for $1 \leq i \leq n - 1$, where each vertex in X has degree $k + 1$ and each vertex in S has degree $k(k + 1)$.

We note that the partition $\pi = \{V_0, V_1, \dots, V_{n-1}\}$ is an upper domatic partition, so $D(G) \geq n = 2k + 1$. On the other hand, since every edge of G is incident to a vertex in X with degree $k + 1$, Proposition 25 implies that $Tr(G) \leq k + 2$, and so, $D(G) - Tr(G) \geq k - 1$. \square

5. Concluding remarks

We conclude by noting another difference in the transitivity number and the upper domatic number of a graph that leads us to making a conjecture. As we saw in Section 1.2, every transitive partition π has at least one source set and at least one sink set. In fact, it was observed in [10] that a graph G will always have a Tr -partition with at least two sink sets. Upper domatic partitions, on the other hand, do not necessarily behave similarly.

For example, consider the graph G_2 from Fig. 2. This graph has no D -partition with a source set. If G_2 had a D -partition with a source set V_1 , then as in the proof of Proposition 29, in the graph $G - V_1$, $\{V_2, V_3, V_4\}$ would be an upper domatic partition. But if $D(G - V_1) = 3$, then $Tr(G - V_1) = 3$ as well. This would imply that a transitive partition of G of order four could be constructed, a contradiction. Additionally, the graph G_2 in Fig. 2 has no D -partition with two sink sets. To see this, note that the graph G_2 has $Tr(G) < D(G) = 4$. Thus, if G_2 had a D -partition with two sink sets, then the partition would also be transitive, again a contradiction.

The question arises whether there are graphs with no D -partitions with a sink set. Unfortunately, we have neither found such a graph nor proven that one does not exist.

Conjecture 1. *There exists a graph G such that no D -partition of G contains a sink set.*

Our next results show that if Conjecture 1 is true for a graph G , then $D(G) \geq 5$.

Proposition 27. *If G has a D -partition π of order k that contains a set that is dominated by $k - 2$ sets, then G has a D -partition π' containing a sink set.*

Proof. Let $\pi = \{V_1, V_2, \dots, V_k\}$ be an upper domatic partition that contains a set V_j that is dominated by $k - 2$ sets. If π does not have a sink set, consider the set V_i that is dominated by V_j . Let u be a vertex in V_j that dominates a vertex in V_i . Let $X = V_i \cup V_j - \{u\}$. Then $\pi' = \pi - \{V_i, V_j\} \cup \{X, \{u\}\}$ is a D -partition of G with the sink set $\{u\}$ (cf. Proposition 2). \square

Corollary 28. *If $D(G) \leq 4$, then there exists a D -partition that contains a sink set.*

Proof. It follows from previous results that if $D(G) \leq 3$, then $Tr(G) = D(G)$, and there is a Tr -partition π that is a D -partition. Thus, π has a sink set. Hence, assume that $D(G) = 4$. Let π be a D -partition for G , and consider the domination digraph for π . There are four vertices and at least six arcs in this digraph, so some vertex must have in-degree at least $2 = D(G) - 2$. By Proposition 27, the result holds. \square

We also note that if Conjecture 1 holds, the following is true.

Proposition 29. *If there exist graphs that have no D -partitions containing a sink set, then there exist graphs that have no D -partitions containing either a sink set or a source set.*

Proof. Let G be a graph for which $D(G) = k$, and G has no D -partition containing a sink set. Assume that k is the minimum integer for which a graph G exists for which $D(G) = k$ and G has no D -partitions containing a sink set. Let $\pi = \{V_1, V_2, \dots, V_k\}$ be a D -partition of G , where, by assumption, V_k is not a sink set, and assume that V_1 is a source set. This means that V_1 is a dominating set of G . Then $\pi' = \{V_2, V_3, \dots, V_k\}$ is an upper domatic partition of order $k - 1$ of the graph $G' = G - V_1$.

First, we claim that $D(G') = k - 1$. From the upper domatic partition π' , we know that $D(G') \geq k - 1$. Suppose $D(G') > k - 1$. Let $\{W_1, W_2, \dots, W_r\}$ be a D -partition of G' , where $r \geq k$. Then it follows that $\{V_1, W_1, W_2, \dots, W_r\}$ is an upper domatic partition of G of order greater than k ; a contradiction. Thus, $D(G') = k - 1$.

Now suppose that G' has a D -partition $\{U_1, U_2, \dots, U_{k-1}\}$ in which U_{k-1} is a sink. Then $\{V_1, U_1, U_2, \dots, U_{k-1}\}$ is a D -partition of G containing a sink; a contradiction.

Thus, G' has no D -partition containing a sink, but $D(G') = k - 1 < k$, which contradicts our assumption that k is the minimum integer for which a graph G exists having $D(G) = k$ and no D -partitions containing a sink set.

Thus, it follows that G has no D -partitions containing either a sink set or a source set. \square

References

- [1] E.J. Cockayne, S.T. Hedetniemi, Towards a theory of domination in graphs, *Networks* 7 (1977) 247–261.
- [2] G.J. Chang, The domatic number problem, *Discrete Math.* 125 (1994) 115–122.
- [3] T.L. Lu, P.H. Ho, G.J. Chang, The domatic number problem in interval graphs, *SIAM J. Discrete Math.* 3 (1990) 531–536.
- [4] G.K. Manacher, T.A. Mankus, Finding a domatic partition of an interval graph in time $O(n)$, *SIAM J. Discrete Math.* 9 (1996) 167–172.
- [5] S.L. Peng, M.S. Chang, A simple linear time algorithm for the domatic partition problem on strongly chordal graphs, *Inform. Process. Lett.* 43 (1992) 297–300.

- [6] A.S. Rao, C. Pandu Rangan, Linear algorithm for domatic number problem on interval graphs, *Inform. Process. Lett.* 33 (1989-1990) 29–33.
- [7] B. Zelinka, On k -domatic numbers of a graph, *Czechoslovak Math. J.* 33 (1983) 309–313.
- [8] B. Zelinka, Domatically critical graphs, *Czechoslovak Math. J.* 30 (1980) 486–489.
- [9] B. Zelinka, Domatic number and linear arboricity of cacti, *Math. Slovaca* 36 (1986) 41–54.
- [10] J.T. Hedetniemi, S.T. Hedetniemi, The transitivity of a graph, *J. Combin. Math. Combin. Comput.* 104 (2018) 75–91.
- [11] T.W. Haynes, S.T. Hedetniemi, P.J. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, New York, 1998.
- [12] W. Goddard, S.M. Hedetniemi, S.T. Hedetniemi, A.A. McRae, The algorithmic complexity of domination digraphs, *J. Combin. Math. Combin. Comput.* 80 (2012) 367–384.
- [13] T.W. Haynes, J.T. Hedetniemi, S.T. Hedetniemi, A.A. McRae, N. Phillips, The transitivity of special graph classes, *J. Combin. Math. Combin. Comput.* (2018) (in press).