



## A semi-strong perfect digraph theorem

Stephan Dominique Andres, Helena Bergold, Winfried Hochstättler & Johanna Wiehe

To cite this article: Stephan Dominique Andres, Helena Bergold, Winfried Hochstättler & Johanna Wiehe (2020) A semi-strong perfect digraph theorem, AKCE International Journal of Graphs and Combinatorics, 17:3, 992-994, DOI: [10.1016/j.akcej.2019.12.018](https://doi.org/10.1016/j.akcej.2019.12.018)

To link to this article: <https://doi.org/10.1016/j.akcej.2019.12.018>



© 2020 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 23 Apr 2020.



Submit your article to this journal [↗](#)



Article views: 145



View related articles [↗](#)



View Crossmark data [↗](#)

## A semi-strong perfect digraph theorem

Stephan Dominique Andres, Helena Bergold, Winfried Hochstättler, and Johanna Wiehe

Fakultät für Mathematik und Informatik, IZ, FernUniversität in Hagen, Hagen, Germany

### ABSTRACT

Reed (1987) showed that, if two graphs are  $P_4$ -isomorphic, then either both are perfect or none of them is. In this note, we will derive an analogous result for perfect digraphs.

### KEYWORDS

Dichromatic number;  
perfect graph;  
perfect digraph

### 2000 MSC

05C17; 05C20; 05C15

### 1. Introduction and notation

Perfect digraphs have been introduced by Andres and Hochstättler [1] as the class of digraphs where the clique number equals the dichromatic number for every induced subdigraph. Reed [8] showed that, if two graphs are  $P_4$ -isomorphic, then either both are perfect or none of them is, which was conjectured by Chvátal [3]. In this note we will derive an analogous result for perfect digraphs.

We start with some definitions. For basic terminology we refer to Bang-Jensen and Gutin [2]. For the rest of the paper, we only consider digraphs without loops. Let  $D$  be a digraph. The *symmetric part*  $S(D)$  of  $D = (V, A)$  is the digraph  $(V, A_2)$  where  $A_2$  is the union of all pairs of anti-parallel arcs of  $D$ , the *oriented part*  $O(D)$  of  $D$  is the digraph  $(V, A_1)$  where  $A_1 = A \setminus A_2$ .

A proper  $k$ -coloring of  $D$  is an assignment  $c : V \rightarrow \{1, \dots, k\}$  such that for all  $1 \leq i \leq k$  the digraph induced by  $c^{-1}(\{i\})$  is acyclic. The *dichromatic number*  $\chi(D)$  of  $D$  is the smallest non-negative integer  $k$  such that  $D$  admits a proper  $k$ -coloring (see Neumann-Lara [7]). A *clique* in a digraph  $D$  is a subdigraph in which for any two distinct vertices  $v$  and  $w$  both arcs  $(v, w)$  and  $(w, v)$  exist. The *clique number*  $\omega(D)$  of  $D$  is the size of the largest clique in  $S(D)$ . The clique number is an obvious lower bound for the dichromatic number.  $D$  is called *perfect* if, for any induced subdigraph  $H$  of  $D$ ,  $\chi(H) = \omega(H)$ .

An (undirected) graph  $G = (V, E)$  can be considered as the symmetric digraph  $D_G = (V, A)$  with  $A = \{(v, w), (w, v) \mid vw \in E\}$ . In the following, we will not distinguish between  $G$  and  $D_G$ . In this way, the dichromatic number of a graph  $G$  is its chromatic number  $\chi(G)$ , the clique number of  $G$  is its usual clique number  $\omega(G)$ , and  $G$  is perfect as a digraph if and only if  $G$  is perfect as a graph.

A main result of Andres and Hochstättler [1] is the following:

**Theorem 1** (Andres and Hochstättler [1]). *A digraph  $D = (V, A)$  is perfect if and only if  $S(D)$  is perfect and  $D$  does not contain any directed cycle  $\vec{C}_n$  with  $n \geq 3$  as induced subdigraph.*

Together with the Strong Perfect Graph Theorem (see e.g. Golumbic [6]) this yields a characterization of perfect digraphs by forbidden induced minors. The Weak Perfect Graph Theorem (see Golumbic [6]), though, does not generalize. The directed 4-cycle  $\vec{C}_4$  is not perfect but its complement is perfect, thus perfection is in general not maintained under taking complements.

Two graphs  $G = (V, E_1)$  and  $H = (V, E_2)$  are  $P_4$ -isomorphic, if any set  $\{a, b, c, d\} \subseteq V$  induces a chordless path, i.e. a  $P_4$ , in  $G$  if and only if it induces a  $P_4$  in  $H$ .

**Theorem 2** (Semi-strong Perfect Graph Theorem (Reed [8])). *If  $G$  and  $H$  are  $P_4$ -isomorphic, then*

$$G \text{ is perfect} \iff H \text{ is perfect.}$$

The graphs without an induced  $P_4$  are the cographs, see Corniel et al. [4]. Thus any pair of cographs with the same number of vertices is  $P_4$ -isomorphic. In order to generalize Theorem 2 to digraphs we consider the class of directed cographs investigated by Crespelle and Paul [5], which are characterized by a set  $\mathcal{F}$  of eight forbidden induced minors. Since the class of directed cographs is invariant under taking complements and perfect digraphs are not, it is clear that isomorphism with respect to  $\mathcal{F}$  will not yield the right notion of isomorphism for our purposes. It turns out that restricting to five of these minors yields the desired result.

## 2. $P^4C$ -isomorphic digraphs

The five forbidden induced minors from the characterization by Crespelle and Paul [5] that we need are the symmetric path  $P_4$ , the directed 3-cycle  $\vec{C}_3$ , the directed path  $\vec{P}_3$  and the two possible augmentations  $\vec{P}_3^+$  and  $\vec{P}_3^-$  of the  $\vec{P}_3$  with one antiparallel edge (see Figure 1).

**Definition 3.** Let  $D = (V, A)$  and  $D' = (V, A')$  be two digraphs on the same vertex set. Then  $D$  and  $D'$  are said to be  $P^4C$ -isomorphic if and only if

1. any set  $\{a, b, c, d\} \subseteq V$  induces a  $P_4$  in  $S(D)$  if and only if it induces a  $P_4$  in  $S(D')$ ,
2. any set  $\{a, b, c\} \subseteq V$  induces a  $\vec{C}_3$  in  $D$  if and only if it induces a  $\vec{C}_3$  in  $D'$ ,
3. any set  $\{a, b, c\} \subseteq V$  induces a  $\vec{P}_3$  with midpoint  $b$  in  $D$  if and only if it induces a  $\vec{P}_3$  with midpoint  $b$  in  $D'$  and
4. any set  $\{a, b, c\} \subseteq V$  induces a  $\vec{P}_3^+$  or a  $\vec{P}_3^-$  in either case with midpoint  $b$  in  $D$  if and only if it induces one of them with midpoint  $b$  in  $D'$ .

Note that the  $P_4$  in case 1 is not necessarily induced in  $D$ , resp. in  $D'$ .

**Lemma 4.** If  $D$  and  $D'$  are  $P^4C$ -isomorphic, then  $D$  contains an induced directed cycle of length  $k \geq 3$  if and only if the same is true for  $D'$ .

*Proof.* By symmetry it suffices to prove that, if  $\{v_0, \dots, v_{k-1}\}$  induces a directed cycle  $\vec{C}_k$  in  $D$ , then the same holds for  $D'$ . The assertion is clear if  $k=3$ , thus assume  $k \geq 4$ . We may, furthermore, assume that the vertices are traversed in consecutive order in  $D$ . Since  $D$  and  $D'$  are  $P^4C$ -isomorphic, each set  $\{v_i, v_{i+1}, v_{i+2}\}$  induces a  $\vec{P}_3$  with midpoint  $v_{i+1}$  in  $D'$ , where indices are taken modulo  $k$ . This yields a directed cycle  $C$  on  $v_0, \dots, v_{k-1}$ , possibly with opposite orientation w.r.t.  $D$ . In that case we relabel the vertices such that the label coincides with the direction of traversal. We claim the cycle is induced in  $D'$ , too.

Assume it is not, i.e.  $C$  has a chord  $(v_i, v_j)$ ,  $j \neq i - 1$  in  $D'$ . We choose  $j$  such that the directed path from  $v_j$  to  $v_i$  on  $C$  is shortest possible. If  $(v_i, v_j)$  is an asymmetric arc, then, since  $\{v_i, v_j, v_{j+1}\}$  does not induce a  $\vec{C}_3$ , it must induce a  $\vec{P}_3$  with midpoint  $v_j$  in  $D'$  and hence the same must hold in  $D$ , contradicting  $\vec{C}_k$  being induced. If we have a pair of antiparallel edges between  $v_i$  and  $v_j$ , then, similarly,  $\{v_i, v_j, v_{j+1}\}$  induces a  $\vec{P}_3^+$  or a  $\vec{P}_3^-$  with midpoint  $v_j$ , also leading to a contradiction.  $\square$

**Theorem 5.** If  $D$  and  $D'$  are  $P^4C$ -isomorphic then

$$D \text{ is perfect} \iff D' \text{ is perfect.}$$




Figure 1. The five induced subdigraphs considered.

*Proof.* By assumption  $S(D)$  and  $S(D')$  are  $P_4$ -isomorphic, hence using Theorem 2 we find that  $S(D)$  is perfect if and only if  $S(D')$  is perfect. By Lemma 4,  $D$  contains an induced directed cycle of length at least three if and only if the same holds for  $D'$ . The assertion thus follows from Theorem 1.  $\square$

## 3. Transitive extensions of cographs

In this section we will analyse the class of digraphs without any of the five subgraphs, which thus are trivially pairwise  $P^4C$ -isomorphic.

Since the symmetric part of such a graph is a cograph, we may consider its cotree (cf. Corneil et al. [4]) in canonical form, where the labels alternate between 0 and 1. Since the 1-labeled tree vertices correspond to complete joins, there is no additional room for asymmetric arcs. The 0-labeled vertices correspond to disjoint unions. Assume the connected components in  $S(G)$  are  $G_1, \dots, G_k$ .

**Lemma 6.** If there exists an asymmetric arc connecting a vertex  $v_i$  in  $G_i$  to a vertex  $v_j$  in  $G_j$ , then  $G_i$  and  $G_j$  are connected by an orientation of the complete bipartite graph  $K_{V(G_i), V(G_j)}$ .

*Proof.* Since  $S(G_i)$  and  $S(G_j)$  are connected and by symmetry, it suffices to show that  $v_i$  must be connected by an asymmetric arc to all symmetric neighbors of  $v_j$ . Let  $w$  be such a neighbor. Since there is no symmetric arc from  $v_i$  to  $w$  and  $\{v_i, v_j, w\}$  must neither induce a  $\vec{P}_3^-$  nor a  $\vec{P}_3^+$ , we must have an asymmetric arc between  $v_i$  and  $w$ .  $\square$

Hence, the asymmetric arcs between the components  $G_1, \dots, G_k$  constitute an orientation of a complete  $\ell$ -partite graph for  $1 \leq \ell \leq k$ . The situation is further complicated by the fact that we must neither create a  $\vec{C}_3$  nor a  $\vec{P}_3$ , where we have to take into account that there may also be asymmetric arcs within the  $G_i$ .

We wonder whether this structure is strict enough to make some problems tractable that are  $\mathcal{NP}$ -complete in general. In particular we would be interested in the complexity of the problem to cover all vertices with a minimum number of vertex disjoint directed paths.

### Disclosure statement

No potential conflict of interest was reported by the author(s).

### References

- [1] Andres, S. D, Hochstättler, W. (2015). Perfect digraphs. *J. Graph Theory* 79(1): 21–29.
- [2] Bang-Jensen, J., Gutin, G. (2009). *Digraphs. Theory, Algorithms and Applications*. London: Springer-Verlag London Ltd.
- [3] Chvátal, V. (1984). A semi-strong perfect graph conjecture. *Ann. Discrete Math.* 21: 279–280.
- [4] Corneil, D. G., Lerchs, H, Burlingham, L. S. (1981). Complement reducible graphs. *Discrete Appl. Math.* 3(3): 163–174.
- [5] Crespelle, C, Paul, C. (2006). Fully dynamic recognition algorithm and certificate for directed cographs. *Discrete Appl. Math.* 154(12): 1722–1741.
- [6] Golubic, M. C. (2004). *Algorithmic Graph Theory and Perfect Graphs*. Ann. of Discrete Math., Vol. 57. 2nd ed. Amsterdam: Elsevier.

- [7] Neumann-Lara, V. (1982). The dichromatic number of a digraph. *J. Combin. Theory Ser. B* 33(3): 265–270.
- [8] Reed, B. (1987). A semi-strong perfect graph theorem. *J. Combin. Theory Ser. B* 43(2): 223–240.