



On strong Skolem starters for \mathbb{Z}_{pq}

Adrián Vázquez-Ávila

To cite this article: Adrián Vázquez-Ávila (2020) On strong Skolem starters for \mathbb{Z}_{pq} , AKCE International Journal of Graphs and Combinatorics, 17:3, 988-991, DOI: [10.1016/j.akcej.2019.12.017](https://doi.org/10.1016/j.akcej.2019.12.017)

To link to this article: <https://doi.org/10.1016/j.akcej.2019.12.017>



© 2020 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 23 Apr 2020.



Submit your article to this journal [↗](#)



Article views: 186




View related articles [↗](#)



View Crossmark data [↗](#)

On strong Skolem starters for \mathbb{Z}_{pq}

Adrián Vázquez-Ávila 

Subdirección de Ingeniería y Posgrado, Universidad Aeronáutica en Querétaro, Querétaro, México

ABSTRACT

In 1991, N. Shalaby conjectured that any additive group \mathbb{Z}_n , where $n \equiv 1$ or $3 \pmod{8}$ and $n \geq 11$, admits a strong Skolem starter and constructed these starters of all admissible orders $11 \leq n \leq 57$. Shalaby et al. [24] was proved if $n = \prod_{i=1}^k p_i^{\alpha_i}$, where p_i is a prime number such that $\text{ord}(2)_{p_i} \equiv 2 \pmod{4}$ and α_i is a non-negative integer, for all $i = 1, \dots, k$, then \mathbb{Z}_n admits a strong Skolem starter. On the other hand, the author [30] gives different families of strong Skolem starters for \mathbb{Z}_p than Shalaby et al, where $p \equiv 3 \pmod{8}$ is an odd prime. Recently, the author [31] gives different families of strong Skolem starters of \mathbb{Z}_{p^n} than Shalaby et al, where $p \equiv 3 \pmod{8}$ and n is an integer greater than 1. In this paper, we give some different families of strong Skolem starters of \mathbb{Z}_{pq} , where $p, q \equiv 3 \pmod{8}$ are prime numbers such that $p < q$ and $(p-1) \nmid (q-1)$.

KEYWORDS

Strong starters; Skolem starters; quadratic residues

1. Introduction

Let G be a finite additive abelian group of odd order $n = 2k + 1$, and let $G^* = G \setminus \{0\}$ be the set of non-zero elements of G . A *starter* for G is a set $S = \{\{x_i, y_i\}, i = 1, \dots, k\}$ such that $\{\{x_i\} \cup \{y_i\} : i = 1, \dots, k\} = G^*$ and $\{\pm(x_i - y_i) : i = 1, \dots, k\} = G^*$. Moreover, if all elements $\{x_i + y_i : i = 1, \dots, k\} \subseteq G^*$ are different, then S is called *strong starter* for G . The reader may consult [8, 9, 13, 14, 18, 20, 22, 23] for works related with.

Strong starters were first introduced by Mullin and Stanton in [28] in constructing Room squares. Starters and strong starters have been useful to construct many combinatorial designs such as Room cubes [12], Howell designs [3, 20], Kirkman triple systems [20, 25], Kirkman squares and cubes [26, 29], and factorizations of complete graphs [2, 4, 10, 11, 13, 16, 17, 21].

Let $n = 2k + 1$ and $1 < 2 < \dots < 2k$ be the order of \mathbb{Z}_n^* . A starter for \mathbb{Z}_n is *Skolem* if it can be written as $S = \{\{x_i, y_i\} : i = 1, \dots, k\}$ such that $y_i > x_i$ and $y_i - x_i = i \pmod{n}$, for $i = 1, \dots, k$. In [27], it was proved the Skolem starter for \mathbb{Z}_n exists if and only if $n \equiv 1, 3 \pmod{8}$. A starter which is both Skolem and strong is called *strong Skolem starter*.

Shalaby in [27] proposed the following:

Conjecture 1.1. *If $n \equiv 1, 3 \pmod{8}$ and $n \geq 11$, then \mathbb{Z}_n admits a strong Skolem starter.*

In [24], it was proved if $n = \prod_{i=1}^k p_i^{\alpha_i}$, where p_i is a prime such that $\text{ord}(2)_{p_i} \equiv 2 \pmod{4}$ and α_i is a non-negative integer, for all $i = 1, \dots, k$, then \mathbb{Z}_n admits a strong Skolem starter, where $\text{ord}(2)_{p_i}$ is the order of the element 2 in \mathbb{Z}_{p_i} . In [30], it was given different families of strong Skolem

starters of \mathbb{Z}_p , where $p \equiv 3 \pmod{8}$ is an odd prime, using a different method than in [24]. Recently in [31], it was given different families of strong Skolem starters of \mathbb{Z}_{p^n} , where $p \equiv 3 \pmod{8}$ and n is an integer greater than 1, than in [24]. In this paper, we use the method of Vázquez-Ávila to give different families of strong Skolem starters for \mathbb{Z}_{pq} than in [24], where $p, q \equiv 3 \pmod{8}$ are prime numbers such that $p < q$ and $(p-1) \nmid (q-1)$.

This paper is organized as follows. In Section 2, we recall some basic properties about quadratic residues and we present the strong Skolem starters of \mathbb{Z}_p given in [30]; this idea is used in the main result of this paper, Theorem 3.3. Finally, in Section 3, we give the main result of this paper, and we present one example. The main theorem states the following:

Theorem 1.2. *Let $p, q \equiv 3 \pmod{8}$ be odd prime numbers such that $p < q$ and $(p-1) \nmid (q-1)$. If $r \in \mathbb{Z}_p^*$ is a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* , then \mathbb{Z}_{pq} admits a strong Skolem starter.*

2. A family of strong Skolem starters for \mathbb{Z}_p

The following definitions and notations are obtained from [30] and [31]: Let p be an odd prime power. An element $x \in \mathbb{Z}_p^*$ is called a *quadratic residue* if there exists an element $y \in \mathbb{Z}_p^*$ such that $y^2 = x$. If there is no such y , then x is called a *non-quadratic residue*. The set of quadratic residues of \mathbb{Z}_p^* is denoted by $QR(p)$ and the set of non-quadratic residues is denoted by $NQR(p)$. It is well known that $QR(p)$ is a (cyclic) subgroup of \mathbb{Z}_p^* of cardinality $\frac{p-1}{2}$ (see for example [15]); also, if either $x, y \in$

$QR(p)$ or $x, y \in NQR(p)$, then $xy \in QR(p)$, and if $x \in QR(p)$ and $y \in NQR(p)$, then $xy \in NQR(p)$.

The following theorems are well known results on quadratic residues. For more details of this kind of results the reader may consult [5, 15].

Theorem 2.1 (Euler’s criterion). *If p is an odd prime and $x \in \mathbb{Z}_p^*$, then*

1. $x \in QR(p)$ if and only if $x^{\frac{p-1}{2}} = 1$.
2. $x \in NQR(p)$ if and only if $x^{\frac{p-1}{2}} = -1$.

Theorem 2.2. *Let p be an odd prime power, then*

1. $-1 \in QR(p)$ if and only if $p \equiv 1 \pmod{4}$.
2. $-1 \in NQR(p)$ if and only if $p \equiv 3 \pmod{4}$.

Theorem 2.3. *Let p be an odd prime. If $p \equiv 3 \pmod{4}$, then*

1. $x \in QR(p)$ if and only if $-x \in NQR(p)$.
2. $x \in NQR(p)$ if and only if $-x \in QR(p)$.

In [1], it was proved the following (see also [22]):

Lemma 2.4. [1] *If $p \equiv 3 \pmod{4}$ is an odd prime with $p \neq 3$, then the following set*

$$S_\beta = \{ \{x, \beta x\} : x \in QR(p) \},$$

is a strong starter for \mathbb{Z}_p , for all $\beta \in NQR(p) \setminus \{-1\}$.

In [30], it was proved the following:

Theorem 2.5. [30] *Let $p \equiv 3 \pmod{8}$ be an odd prime, then the following strong starter*

$$S_\beta = \{ \{x, \beta x\} : x \in QR(p) \}$$

for \mathbb{Z}_p is Skolem, if $\beta = 2$ and $\beta = \frac{1}{2}$.

We will use the strong Skolem starter of Theorem 2.5 for the main result of this paper.

3. A family of strong Skolem starters for \mathbb{Z}_{pq}

In this section, we give different families of strong Skolem starters that given in [24]. The following definitions and notations are obtained from [24]: Let G_n be the group of units of the ring \mathbb{Z}_n (elements invertible with respect to multiplication). It is denoted by $\langle x \rangle_n$ the cyclic subgroup of G_n generated by $x \in G_n$. Also, we will use the notation $aB = \{ab : b \in B\}$, where $a \in \mathbb{Z}$ and $B \subseteq \mathbb{Z}$. On the other hand, it is denoted by $ord(x)_n$ the order of the element $x \in G_n$; hence, $ord(x)_n = |\langle x \rangle_n|$. Whenever the group operation is irrelevant, it will consider G_n and its cyclic multiplicative subgroups $\langle x \rangle_n$ in the set-theoretical sense and denote them by \underline{G}_n and $\langle \underline{x} \rangle_n$, respectively.

Let $p, q \equiv 3 \pmod{8}$ be odd prime numbers such that $p < q$ and $(p-1) \nmid (q-1)$. We have $\underline{G}_{pq} = \{x \in \mathbb{Z}_{pq}^* : gcd(x, pq) = 1\}$, with $|\underline{G}_{pq}| = (p-1)(q-1)$, see for example [6]. Hence, $p\mathbb{Z}_q^*, q\mathbb{Z}_p^*$ and \underline{G}_{pq} forms a partition of \mathbb{Z}_{pq}^* , since every element $x \in \mathbb{Z}_{pq}^*$ lies in one and only one of these sets. Moreover, it is not difficult to prove that if $r \in \mathbb{Z}_p^*$ (or $r \in \mathbb{Z}_q^*$) is a primitive root, then $|\langle r \rangle_{pq}| = lcm(p-1, q-1) = \frac{(p-1)(q-1)}{2}$, since $(p-1) \nmid (q-1)$.

Lemma 3.1. *If $p, q \equiv 3 \pmod{8}$ are odd prime numbers with $p < q$, and $r \in \mathbb{Z}_p^*$ is such that $r \in NQR(p)$ and $r \in NQR(q)$, then $r^{\frac{(p-1)(q-1)}{4}} \equiv -1 \pmod{pq}$. Moreover, if r is a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* , then $-1 \in \langle r^2 \rangle_{pq}$.*

Proof. Recall that G_m is the group of units of \mathbb{Z}_m . It is well known that the map $\Psi : G_{pq} \rightarrow G_p \times G_q$ defined by $\Psi(k_{pq}) = (k_p, k_q)$, is an isomorphism between G_{pq} and $G_p \times G_q$, see for example [7]. Since p and q are prime numbers then $G_p = \mathbb{Z}_p^*$ and $G_q = \mathbb{Z}_q^*$. Let $r \in \mathbb{Z}_p^*$ such that $r \in NQR(p)$ and $r \in NQR(q)$. Hence $r^{\frac{p-1}{2}} = -1 \pmod{p}$ and $r^{\frac{q-1}{2}} = -1 \pmod{q}$, by Theorem 2.1. Then $\Psi(r_{pq}^{\frac{(p-1)(q-1)}{4}}) = (r_p^{\frac{(p-1)(q-1)}{4}}, r_q^{\frac{(p-1)(q-1)}{4}}) = (r_p^{\frac{p-1}{2} \cdot \frac{q-1}{2}}, r_q^{\frac{q-1}{2} \cdot \frac{p-1}{2}}) = (-1_p, -1_q) = \Psi(-1_{pq})$, since $\frac{p-1}{2}$ and $\frac{q-1}{2}$ are odd integers. On the other hand, if r is a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* , then by Theorem 2.2, it follow that $-1_p \in \langle r^2 \rangle_p$ and $-1_q \in \langle r^2 \rangle_q$, which implies that $-1_{pq} \in \langle r^2 \rangle_{pq}$. \square

Lemma 3.2. *Let $p, q \equiv 3 \pmod{8}$ be odd prime numbers such that $p < q$. If $r \in \mathbb{Z}_p^*$ is a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* , then $2 \notin \langle r^2 \rangle_{pq}$.*

Proof. Let $\Psi : G_{pq} \rightarrow G_p \times G_q$ given by $\Psi(k_{pq}) = (k_p, k_q)$, the isomorphism between G_{pq} and $G_p \times G_q$, and let $r \in \mathbb{Z}_p^*$ be a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* . It is well known that $2 \in NQR(p)$ and $2 \in NQR(q)$, where $NQR(p) = \langle r^2 \rangle_p$ and $NQR(q) = \langle r^2 \rangle_q$, see for example [19]. Hence, if $2 \in \langle x^2 \rangle_{pq}$, then there exists $j \in \{0, \dots, \frac{(p-1)(q-1)}{4}\}$ such that $x^{2j} = 2 \pmod{pq}$, which implies that $\Psi(2) = \Psi(x_{pq}^{2j}) = (x_p^{2j}, x_q^{2j}) \neq (2, 2)$, a contradiction. Hence $2 \notin \langle r^2 \rangle_{pq}$. \square

Theorem 3.3. *Let $p, q \equiv 3 \pmod{8}$ be odd prime numbers such that $p < q$ and $(p-1) \nmid (q-1)$. If $r \in \mathbb{Z}_p^*$ is a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* , then \mathbb{Z}_{pq} admits a strong Skolem starter.*

Proof. Let $r \in \mathbb{Z}_p^*$ be a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* . Hence, r^2 is a generator of $QR(p)$ and $QR(q)$. Since $p\mathbb{Z}_q^*, q\mathbb{Z}_p^*$ and \underline{G}_{pq} forms a partition of \mathbb{Z}_{pq}^* , define

$$\begin{aligned} pS_q &= \{ \{px, 2px\} : x \in QR(q) \} \\ qS_p &= \{ \{qx, 2qx\} : x \in QR(p) \} \\ S_{pq} &= \{ \{x, 2x\} : x \in \langle r^2 \rangle_{pq} \} \cup \{ \{ \lambda x, 2\lambda x \} : x \in \langle r^2 \rangle_{pq} \}, \end{aligned}$$

where $\lambda \notin \langle r^2 \rangle_{pq} \cup 2\langle r^2 \rangle_{pq}$. It is easy to see that $\{ \pm px : x \in QR(q) \} = p\mathbb{Z}_q^*, \{ \pm qx : x \in QR(p) \} = q\mathbb{Z}_p^*$ and $\{ \pm x : x \in \langle r^2 \rangle_{pq} \} \cup \{ \pm \lambda x : x \in \langle r^2 \rangle_{pq} \} = \underline{G}_{pq}$, by Lemmas 3.1 and 3.2. Hence, the set $S = pS_q \cup qS_p \cup S_{pq}$ is a starter.

Let define

$$\begin{aligned} qS_p^+ &= \{ 3px : x \in QR(p) \} \\ pS_q^+ &= \{ 3qx : x \in QR(q) \} \\ S_{pq}^+ &= \{ 3x : x \in \langle r^2 \rangle_{pq} \} \cup \{ 3\lambda x : x \in \langle r^2 \rangle_{pq} \} \end{aligned}$$

Since $l_1x + l_2y \neq 0$, for all different $x, y \in QR(p) \cup QR(q) \cup \langle r^2 \rangle_{pq}$, with $l_1, l_2 \in \{1, p, q, \lambda\}$. Then $|qS_p^+| = |QR(p)|, |pS_q^+| = |QR(q)|$ and $|S_{pq}^+| = |\langle r^2 \rangle_{pq}|$. By Lemma 3.1, we have $|qS_p^+ \cup pS_q^+ \cup S_{pq}^+| = |qS_p^+| + |pS_q^+| + |S_{pq}^+| = \frac{p-1}{2} + \frac{q-1}{2} + \frac{(p-1)(q-1)}{2} = \frac{pq-1}{2}$. Hence, the set S is strong.

Finally, we give a proof analogous to the case (i) of the proof of Theorem 2.5 given in [30] to prove S is Skolem. Let $pq = 2t + 1$ and $1 < 2 < \dots < 2t$ be the order of the non-zero elements of \mathbb{Z}_{pq}^* . Define $Q_{\frac{1}{2}} = \{1, 2, \dots, t\}$. To prove that S is Skolem, it is sufficient to prove that, if $2lx > lx$ then $lx \in Q_{\frac{1}{2}}$, and if $lx > 2lx$ then $-lx \in Q_{\frac{1}{2}}$, where $l \in \{1, p, q, \lambda\}$. Suppose that $lx \in Q_{\frac{1}{2}}$, for $l \in \{1, p, q, \lambda\}$, then $2lx > lx$, which implies that $2lx - lx = lx \in Q_{\frac{1}{2}}$. On the other hand, if $lx \notin Q_{\frac{1}{2}}$, for $l \in \{1, p, q, \lambda\}$, then $-lx \in Q_{\frac{1}{2}}$. Hence, $2(-lx) > -lx$, which implies that $-2lx + lx = -lx \in Q_{\frac{1}{2}}$. Hence, the set S is Skolem. \square

Corollary 3.4. *Let $p, q \equiv 3 \pmod{8}$ be odd prime numbers such that $p < q$ and $(p-1) \nmid (q-1)$, and let $r \in \mathbb{Z}_p^*$ be a primitive root of \mathbb{Z}_p^* and \mathbb{Z}_q^* . If $pS_q^- = \{px, 2^{-1}px\} : x \in QR(q)\}$, $S_p^- = \{qx, 2^{-1}qx\} : x \in QR(p)\}$ and $S_{pq}^- = \{x, 2^{-1}x\} : x \in \langle r^2 \rangle_{pq} \cup \{\lambda x, 2^{-1}\lambda x\} : x \in \langle r^2 \rangle_{pq}\}$, where $\lambda \notin \langle r^2 \rangle_{pq} \cup 2\langle r^2 \rangle_{pq}$, then set $S^- = pS_q^- \cup qS_p^- \cup S_{pq}^-$ is a strong Skolem starter for \mathbb{Z}_{pq} .*

Proof. The proof is analogous to the proof of Theorem 3.3, using the case (ii) of Theorem 2.5 given in [30]. \square

Example 1. Consider $\mathbb{Z}_{11 \cdot 19}$. We have $\mathbb{Z}_{11 \cdot 19}^* = 19\underline{G}_{11} \cup 11\underline{G}_{19} \cup \underline{G}_{11 \cdot 19}$, and $r=2$ is a primitive root of \mathbb{Z}_{11}^* and \mathbb{Z}_{19}^* . Then, taking $\lambda=3$, the pairs from $19\underline{G}_{11}$, $11\underline{G}_{19}$ and $\underline{G}_{11 \cdot 19}$ are:

$$\begin{aligned} 19\underline{G}_{11} &: \{19, 38\}, \{76, 152\}, \{95, 190\}, \{171, 133\}, \{57, 114\}. \\ 11\underline{G}_{19} &: \{11, 22\}, \{44, 88\}, \{176, 143\}, \{77, 154\}, \{99, 198\}, \\ &\quad \{187, 165\}, \{121, 33\}, \{66, 132\}, \{55, 110\}. \\ \underline{G}_{11 \cdot 19} &: \{1, 2\}, \{4, 8\}, \{16, 32\}, \{64, 128\}, \{47, 94\}, \\ &\quad \{188, 167\}, \{125, 41\}, \{82, 164\}, \{119, 29\}, \{58, 116\}, \\ &\quad \{23, 46\}, \{92, 184\}, \{159, 109\}, \{9, 18\}, \{36, 72\}, \\ &\quad \{144, 79\}, \{158, 107\}, \{5, 10\}, \{20, 40\}, \{80, 160\}, \\ &\quad \{111, 13\}, \{26, 52\}, \{104, 208\}, \{207, 205\}, \{201, 193\}, \\ &\quad \{177, 145\}, \{81, 162\}, \{115, 21\}, \{42, 84\}, \\ &\quad \{168, 127\}, \{45, 90\}, \{180, 151\}, \{93, 186\}, \{163, 117\}, \\ &\quad \{25, 50\}, \{100, 200\}, \{191, 173\}, \{137, 65\}, \\ &\quad \{130, 51\}, \{102, 204\}, \{199, 189\}, \{169, 129\}, \{49, 98\}, \\ &\quad \{196, 183\}, \{157, 105\}, \{3, 6\} \\ &\quad \cup \{12, 24\}, \{48, 96\}, \{192, 175\}, \{141, 73\}, \\ &\quad \{146, 83\}, \{166, 123\}, \{37, 74\}, \{148, 87\}, \{174, 139\}, \\ &\quad \{69, 138\}, \{67, 134\}, \{59, 118\}, \{27, 54\}, \\ &\quad \{108, 7\}, \{14, 28\}, \{56, 112\}, \{15, 30\}, \{60, 120\}, \{31, 62\}, \\ &\quad \{124, 39\}, \{78, 156\}, \{103, 206\}, \{203, 197\}, \\ &\quad \{185, 161\}, \{113, 17\}, \{34, 68\}, \{136, 63\}, \{126, 43\}, \\ &\quad \{86, 172\}, \{135, 61\}, \{122, 35\}, \{70, 140\}, \\ &\quad \{71, 142\}, \{75, 150\}, \{91, 182\}, \{155, 101\}, \{202, 195\}, \\ &\quad \{181, 153\}, \{97, 194\}, \{179, 149\}, \{89, 178\}, \\ &\quad \{147, 85\}, \{170, 131\}, \{53, 106\}. \end{aligned}$$

This strong Skolem starter of $\mathbb{Z}_{11 \cdot 19}$ is the same from the Example 4.12 given in [24]. Now, by Corollary 3.4, we have

a different strong Skolem starter of $\mathbb{Z}_{11 \cdot 19}$, using the same parameters:

$$\begin{aligned} 19\underline{G}_{11} &: \{19, 114\}, \{76, 38\}, \{95, 152\}, \{171, 190\}, \{57, 133\}. \\ 11\underline{G}_{19} &: \{11, 110\}, \{44, 22\}, \{176, 88\}, \{77, 143\}, \{99, 154\}, \\ &\quad \{187, 198\}, \{121, 165\}, \{66, 33\}, \{55, 132\}. \\ \underline{G}_{11 \cdot 19} &: \{1, 105\}, \{4, 2\}, \{16, 8\}, \{64, 32\}, \{47, 128\}, \\ &\quad \{188, 94\}, \{125, 167\}, \{82, 41\}, \{119, 164\}, \{58, 29\}, \\ &\quad \{23, 116\}, \{92, 46\}, \{159, 84\}, \{9, 109\}, \{36, 18\}, \\ &\quad \{144, 72\}, \{158, 79\}, \{5, 107\}, \{20, 10\}, \{80, 40\}, \\ &\quad \{111, 160\}, \{26, 13\}, \{104, 52\}, \{207, 208\}, \\ &\quad \{201, 205\}, \{177, 193\}, \{81, 145\}, \{115, 162\}, \{42, 21\}, \\ &\quad \{168, 84\}, \{45, 127\}, \{180, 90\}, \{93, 151\}, \\ &\quad \{163, 186\}, \{25, 117\}, \{100, 50\}, \{191, 200\}, \{137, 173\}, \\ &\quad \{130, 65\}, \{102, 51\}, \{199, 204\}, \{169, 189\}, \\ &\quad \{49, 129\}, \{196, 98\}, \{157, 183\}, \{3, 106\} \\ &\quad \cup \{12, 6\}, \{48, 24\}, \{192, 96\}, \{141, 175\}, \{146, 73\}, \\ &\quad \{166, 83\}, \{37, 123\}, \{148, 74\}, \{174, 87\}, \{69, 139\}, \\ &\quad \{67, 138\}, \{59, 134\}, \{27, 118\}, \{108, 54\}, \\ &\quad \{14, 7\}, \{56, 28\}, \{15, 112\}, \{60, 30\}, \{31, 120\}, \{124, 62\}, \\ &\quad \{78, 39\}, \{103, 156\}, \{203, 206\}, \{185, 197\}, \\ &\quad \{113, 161\}, \{34, 17\}, \{136, 68\}, \{126, 63\}, \{86, 43\}, \\ &\quad \{135, 172\}, \{122, 61\}, \{70, 35\}, \{71, 140\}, \\ &\quad \{75, 142\}, \{91, 150\}, \{155, 182\}, \{202, 101\}, \{181, 195\}, \\ &\quad \{97, 153\}, \{179, 194\}, \{89, 149\}, \\ &\quad \{147, 178\}, \{170, 85\}, \{53, 131\}. \end{aligned}$$

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

Research was partially supported by SNI and CONACyT.

ORCID

Adrián Vázquez-Ávila  <http://orcid.org/0000-0003-1602-2872>

References

- [1] Alfaro, C. A., Rubio-Montiel, C., Vázquez-Ávila, A. (2020). On two-quotient strong starters for \mathbb{F}_q . *Util. Math.* In press.
- [2] Anderson, B. A. (1974). A Class of Starter Induced 1-Factorizations. In: Bari R. A., Harary F. (eds). *Graphs and Combinatorics*. Lecture Notes in Math., Vol. 406. Berlin: Springer, pp. 180–185.
- [3] Anderson, B. A., Schellenberg, P. J., Stinson, D. R. (1984). The existence of Howell designs of even side. *J. Combin. Theory Ser. A* 36(1): 23–55.
- [4] Bao, J, Ji, L. (2019). Two orthogonal 4-cycle-free one-factorizations of complete graphs. *Graphs Combin.* 35(2): 373–392.

- [5] Burton, D. M. (2007). *Elementary Number Theory*. International series in pure and applied mathematics. 6th ed. McGraw-Hill.
- [6] Cohen, H. (1993). *A Course in Computational Algebraic Number Theory*. Heidelberg: Springer-Verlag.
- [7] Childs, L. N. (2009). *A Concrete Introduction to Higher Algebra*. 3rd ed. New York: Springer-Verlag.
- [8] Dillon, J. F, Morris, R. A. (1973). A skew Room square of side 257. *Utilitas Math.* 4: 187–192.
- [9] Dinitz, J. H. (1984). Room n -cubes of low order. *J. Aust. Math. Soc. A* 36(2): 237–252.
- [10] Dinitz, J. H, Dukes, P. (2006). On the structure of uniform one-factorizations from starters in finite fields. *Finite Fields Appl.* 12(2): 283–300.
- [11] Dinitz, J. H., Dukes, P, Stinson, D. R. (2005). Sequentially perfect and uniform one-factorizations of the complete graph. *Electron. J. Combin.* 12(1).
- [12] Dinitz, J. H, Stinson, D. R. (1981). The spectrum of room cubes. *Eur. J. Combin.* 2(3): 221–230.
- [13] Dinitz, J. H, Stinson, D. R. (1989). Some new perfect one-factorizations from starters in finite fields. *J. Graph Theory* 13(4): 405–415.
- [14] Gross, K. B. (1975). Some new classes of strong starters. *Discrete Math.* 12(3): 225–243.
- [15] Hardy, G. H, Wright, E. M. (2008). *An Introduction to the Theory of Numbers*. 6th ed. Oxford: Oxford University Press. Revised by D. R. Heath-Brown and J. H. Silverman, With a foreword by Andrew Wiles.
- [16] Horton, J. D. (1981). Room designs and one-factorizations. *Aeq. Math.* 22(1): 56–63.
- [17] Horton, J. D. (1983). The construction of Kotzig factorizations. *Discrete Math* 43(2-3): 199–206.
- [18] Horton, J. D. (1990). Orthogonal starters in finite abelian groups. *Discrete Math.* 79(3): 265–278.
- [19] Ireland, K, Rosen, M. A. (1990). *A Classical Introduction to Modern Number Theory*. 2nd ed. New York: Springer-Verlag.
- [20] Kocay, W. L., Stinson, D. R, Vanstone, S. A. (1985). On strong starters in cyclic groups. *Discrete Math.* 56(1): 45–60.
- [21] Meszka, M. (2009). k -cycle free one-factorizations of complete graphs. *Electron. J. Combin.* 16(1): 255–258.
- [22] Mullin, R. C, Nemeth, E. (1969). An existence theorem for room squares. *Can. Math. Bull.* 12(4): 493–497.
- [23] Mullin, R. C, Nemeth, E. (1969). On furnishing room squares. *J. Combin. Theory* 7(3): 266–272.
- [24] Ogandzhanyants, O., Kondratieva, M, Shalaby, N. (2019). Strong Skolem starters. *J. Combin. Des.* 27(1): 5–21.
- [25] Ray-Chaudhuri, D. K, Wilson, R. M. (1971). Solution of Kirkman’s Schoolgirl Problem. In *Combinatorics (Proc. Sympos. Pure Math., Vol. XIX, Univ. California, Los Angeles, Calif., 1968)*. Providence, RI: Amer. Math. Soc., pp. 187–203.
- [26] Rosa, A, Vanstone, S. A. (1985). On the existence of strong Kirkman cubes of order 39 and block size 3. In *Algorithms in Combinatorial Design Theory, North-Holland Math. Stud., Vol. 114*. Amsterdam: North-Holland, pp. 309–319.
- [27] Shalaby, N. (1991). Skolem sequences: generalizations and applications. Thesis (PhD). McMaster University, Canada.
- [28] Stanton, R. G, Mullin, R. C. (1968). Construction of Room squares. *Ann. Math. Stat.* 39(5): 1540–1548.
- [29] Stinson, D. R, Vanstone, S. A. (1985). A Kirkman square of order 51 and block size 3. *Discrete Math.* 55(1): 107–111.
- [30] Vázquez-Ávila, A. (2020). A note on strong Skolem starters. *Discrete Math.* 343(2): 111672.
- [31] Vázquez-Ávila, A. (in press). On new families of strong Skolem starters for \mathbb{Z}_p^n . <https://arxiv.org/abs/1907.05266>