



## Decomposition of complete bipartite graphs into cycles and stars with four edges

M. Ilayaraja & A. Muthusamy

To cite this article: M. Ilayaraja & A. Muthusamy (2020) Decomposition of complete bipartite graphs into cycles and stars with four edges, AKCE International Journal of Graphs and Combinatorics, 17:3, 697-702, DOI: [10.1016/j.akcej.2019.12.006](https://doi.org/10.1016/j.akcej.2019.12.006)

To link to this article: <https://doi.org/10.1016/j.akcej.2019.12.006>



© 2020 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 22 Apr 2020.



Submit your article to this journal [↗](#)



Article views: 496



View related articles [↗](#)



View Crossmark data [↗](#)

# Decomposition of complete bipartite graphs into cycles and stars with four edges

M. Ilayaraja and A. Muthusamy

Department of Mathematics, Periyar University, Salem, Tamil Nadu, India

## ABSTRACT

Let  $C_k$ ,  $S_k$  denote a cycle, star with  $k$  edges and let  $K_{m,n}$  denotes a complete bipartite graph with  $m$  and  $n$  vertices in the parts. In this paper, we obtain necessary and sufficient conditions for the existence of a decomposition of complete bipartite graphs into cycles and stars with four edges.

## KEYWORDS

Cycles; star; graph decomposition

## 2010 MSC

05B30; 05C38;

## 1. Introduction

All graphs considered here are finite. For the standard graph-theoretic terminology the reader is referred to [4]. Let  $C_k$ ,  $S_k$  denote a cycle, star with  $k$  edges and let  $K_{m,n}$  denotes a complete bipartite graph with  $m$  and  $n$  vertices in the parts. Also we denote the cycle  $C_k$  with vertices  $x_0, x_1, \dots, x_{k-1}$  and edges  $x_0x_1, x_1x_2, \dots, x_{k-2}x_{k-1}, x_{k-1}x_0$  as  $(x_0, x_1, \dots, x_{k-1}, x_0)$  and a star  $S_k$  consists of a centre vertex  $x_0$  of degree  $k$  (i.e.,  $d(x_0) = k$ ) and  $k$  end vertices  $x_1, x_2, \dots, x_k$  as  $(x_0; x_1, \dots, x_k)$ . If there are  $t$  stars with same end vertices  $x_1, x_2, \dots, x_k$  and different centres  $y_1, y_2, \dots, y_t$ , we denote it by  $(y_1, y_2, \dots, y_t; x_1, x_2, \dots, x_k)$ . Note that  $S_k$  is isomorphic to  $K_{1,k}$ . By a *decomposition* of  $G$ , we mean a list of edge-disjoint subgraphs of  $G$  whose union is  $G$  (ignoring isolated vertices). For the graph  $G$ , if  $E(G)$  can be partitioned into  $E_1, \dots, E_k$  such that the subgraph induced by  $E_i$  is  $H_i$ , for all  $i$ ,  $1 \leq i \leq k$ , then we say that  $H_1, \dots, H_k$  decompose  $G$  and we write  $G = H_1 \oplus \dots \oplus H_k$ . For  $1 \leq i \leq k$ , if  $H_i \cong H$ , we say that  $G$  has a *H-decomposition* and it is denoted by  $H|G$ . If  $G$  can be decomposed into  $p$  copies of  $H_1$  and  $q$  copies of  $H_2$ , then we say that  $G$  has a  $\{pH_1, qH_2\}$ -decomposition or  $(H_1, H_2)$ -multidecomposition. If such a decomposition exists for all  $p$  and  $q$  satisfying trivial necessary conditions, then we say that  $G$  has a  $\{H_1, H_2\}_{\{p, q\}}$ -decomposition or complete  $\{H_1, H_2\}$ -decomposition. We denote the number of edges of  $G$  by  $e(G)$ .

Cycle decomposition of graphs and star decomposition of graphs are popular topic of research in graph theory; see [3, 12–14]. The study of  $(K, H)$ -multidecomposition has been introduced by Atif Abueida and M. Daven [1]. Moreover, Atif Abueida and Theresa O’Neil [2] have settled the existence of  $(K, H)$ -multidecomposition of  $K_m(\lambda)$  when  $(K, H) = (K_{1, n-1}, C_n)$  for  $n = 3, 4, 5$ . Priyadharsini and Muthusamy [9] established necessary and sufficient condition for the existence

of  $(G_n, H_n)$ -multidecomposition of  $\lambda K_n$  where  $G_n, H_n \in \{C_n, P_{n-1}, S_{n-1}\}$ . Lee [7], gave necessary and sufficient condition for the multidecomposition of  $K_{m,n}$  into at least one copy of  $C_k$  and  $S_k$ . Lee and J.J. Lin [8], have obtained necessary and sufficient condition for the decomposition of complete bipartite graph minus a one factor into cycles and stars. Shyu [10] considered the existence of a decomposition of  $K_{m,n}$  into paths and stars with  $k$  edges, giving a necessary and sufficient condition for  $k = 3$ . Jeevadoss and Muthusamy [5] have obtained some necessary and sufficient condition for the existence of a decomposition of complete bipartite graphs into paths and cycles. Recently, Lee [6] established necessary and sufficient conditions for the existence of a decomposition of complete bipartite multigraph into cycles and stars with at least one copy of each. In this paper, we study about the existence of a decomposition of complete bipartite graphs into  $p$  copies of  $C_4$  and  $q$  copies of  $S_4$  for all possible values of  $p$  and  $q$ . We abbreviate the notation for such a decomposition as  $\{pC_4, qS_4\}$ -decomposition. In fact, we establish necessary and sufficient conditions for the existence of  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ .

To prove our main results, we state the following:

**Theorem 1.1.** [11] *Let  $m, n$  and  $l \in \mathbb{Z}_+$ . There exists an  $C_{2l}$ -decomposition of  $K_{m,n}$  if and only if  $m$  and  $n$  are even,  $m, n \geq l \geq 2$  and  $mn \equiv 0 \pmod{2l}$ .*

**Theorem 1.2.** [14] *Let  $k, m$  and  $n \in \mathbb{Z}_+$  with  $m \leq n$ . There exists an  $S_k$ -decomposition of  $K_{m,n}$  if and only if one of the following holds:*

- (i)  $k \leq m$  and  $mn \equiv 0 \pmod{k}$ ;
- (ii)  $m < k \leq n$  and  $n \equiv 0 \pmod{k}$ .

## Remarks.

1. If  $G_1$  and  $G_2$  have a  $\{pC_4, qS_4\}$ -decomposition, then  $G_1 \oplus G_2$  has a such decomposition.

2. Let  $X + Y = \{(x_1 + y_1, x_2 + y_2) \mid (x_1, x_2) \in X, (y_1, y_2) \in Y\}$  and  $rX$  is the sum of  $r$  copies of  $X$ .

**2. Necessary conditions**

The following Lemmas gives necessary conditions for the existence of a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ .

**Lemma 2.1.** *Let  $p, q \geq 0$  and even  $n \geq 2$ . If  $K_{2,n}$  has a  $\{pC_4, qS_4\}$ -decomposition, then  $q$  must be even.*

*Proof.* Let  $D$  be an arbitrary  $\{pC_4, qS_4\}$ -decomposition of  $K_{2,n}$ . Then  $4(p + q) = e(K_{2,n})$ . Let  $V(K_{2,n}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, x_{12}\}$  and  $X_2 = \{x_{21}, \dots, x_{2n}\}$ . Since  $x_{11}$  and  $x_{12}$  must be a centre vertex of each  $S_4$ 's in  $D$  and each  $C_4$ 's in  $D$  must contains both  $x_{11}$  and  $x_{12}$ . Therefore the number of copies of  $S_4$  centered in both  $x_{11}$  and  $x_{12}$  are the same. □

**Lemma 2.2.** *Let  $p, q \geq 0$  and even  $n \geq 4$ . If  $K_{4,n}$  has a  $\{pC_4, qS_4\}$ -decomposition, then  $p, q \neq 1$ .*

*Proof.* Let  $D$  be an arbitrary  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ . Then  $4(p + q) = e(K_{m,n})$ . On the contrary, suppose that  $q = 1$ . Let  $S_4^1$  denote the only star in  $D$ . It follows that each end vertex of  $S_4^1$  has odd degree in  $K_{4,n} \setminus E(S_4^1)$ , which cannot have a cycle decomposition and hence a contradiction. On the other hand, let  $V(K_{4,n}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, \dots, x_{14}\}$  and  $X_2 = \{x_{21}, \dots, x_{2n}\}$ . Assume that there exist a  $(4; 1, n - 1)$ -decomposition of  $K_{4,n}$ . Without loss of generality, let  $C_4^1 = (x_{11}, x_{21}, x_{12}, x_{22}, x_{11})$  be the only  $C_4$  in  $D$ . Then by assumption  $K_{4,n} \setminus E(C_4^1) = G$  has an  $S_4$ -decomposition. In  $G$ , the vertices  $x_{21}$  and  $x_{22}$  have exactly degree 2. So it could not be a centre vertex of any stars in  $D$ . Therefore we have two stars  $S_4^1$  and  $S_4^2$  whose centre vertex is  $x_{13}$  and  $x_{14}$  respectively, which consist of  $x_{21}$  and  $x_{22}$  as end vertices. That is,  $S_4^1 = (x_{13}; x_{21}, x_{22}, x_{2i}, x_{2j})$  and  $S_4^2 = (x_{14}; x_{21}, x_{22}, x_{2i'}, x_{2j'})$  for some  $i \neq j$  and  $i' \neq j' \in \mathbb{Z}_+$ . We define  $D' = \{S_4 \in D \mid \text{Centre vertex of } S_4 \text{ is } x_{14}\}$ ;  $|D'| = r$ , where  $r \in \mathbb{Z}_+$  and let  $X_2' = \{y \in X_2 \mid y \text{ is an end vertex of } S_4 \in D'\}$ . Then clearly  $S_4^2 \in D'$  and  $|X_2'| = 4r$ . Every vertex in  $X_2 \setminus X_2'$  must be centre vertex of some stars in  $D \setminus D'$ . Otherwise we cannot use the edges between  $x_{14} \in X_1$  and  $X_2 \setminus X_2'$  in any star in  $D$ , whose centre vertex is not  $x_{14}$ .

Now we collect all the stars whose centre in  $X_2 \setminus X_2'$  and denote it  $D''$ . That is,  $D'' = \{S_4 \in D \mid \text{Centre vertex of } S_4 \text{ in } X_2 \setminus X_2'\}$ . Then the new graph  $G' = K_{4,n} \setminus \{E(C_4^1) \cup E(D') \cup E(D'') \cup E(S_4^1)\} \cong K_{3,4r-2} \setminus \{x_{13}x_{2i}\} \cup \{x_{13}x_{2j}\}$ . Let  $V(K_{3,4r-2}) = (Y_1, Y_2)$ , where  $Y_1 = X_1 \setminus \{x_{14}\}$  and  $Y_2 = X_2 \setminus (X_2 \setminus X_2' \cup \{x_{21}, x_{22}\})$ . By our assumption  $G'$  has an  $S_4$ -decomposition. In  $G'$ ,  $d(x_{11}) = d(x_{12}) = 4r - 2$  and  $d(x_{13}) = 4r - 4$  also  $d(x_{2i}) = d(x_{2j}) = 2$  and  $d(x_{2k}) = 3$ , where  $x_{2k} \neq x_{2i}, x_{2j}$  and  $x_{2k} \in X_2$ . It follows that no vertex of  $Y_2$  can be a centre vertex of any stars, so the centre vertices of stars must be in  $Y_1$ . Since  $d(x_{11}) = d(x_{12}) = 4r - 2 \not\equiv 0 \pmod{4}$ , by **Theorem 1.2**, the

graph  $G'$  cannot have  $S_4$ -decomposition, which is contradiction to our assumption. □

**Lemma 2.3.** *Let  $p, q \geq 0$  and even  $m, n \geq 6$  with  $m \leq n$ . If  $K_{m,n}$  has a  $\{pC_4, qS_4\}$ -decomposition, then  $q \neq 1$ .*

*Proof.* Let  $D$  be an arbitrary  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ . On the contrary, suppose  $q = 1$ , we obtain a contradiction as in **Lemma 2.2**. □

**Lemma 2.4.** *Let  $p, q$  be nonnegative integers,  $m$  is odd (resp.,  $n$  is odd) and  $n \equiv 0 \pmod{4}$  (resp.,  $m \equiv 0 \pmod{4}$ ) such that  $m < n$ . If  $K_{m,n}$  has a  $\{pC_4, qS_4\}$ -decomposition, then  $q \geq \frac{n}{4}$  (resp.,  $q \geq \frac{m}{4}$ ).*

*Proof.* Let  $D$  be an arbitrary  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ . Let the  $q$  copies of stars be  $S_4^i \in D, 1 \leq i \leq q$  and  $G = K_{m,n} \setminus E(\sum_{i=1}^q S_4^i)$ . By hypothesis,  $G$  has a  $C_4$ -decomposition. It follows that every vertex of  $G$  must be of even degree. Note that when  $n$  is even (resp.,  $m$ ), each vertex of  $X_2$  (resp.,  $X_1$ ) must be either an end vertex or the centre of some  $S_4^i, 1 \leq i \leq q$ . It implies that  $4q \geq n$  (resp.,  $4q \geq m$ ). □

**3. Sufficient conditions**

The following sequence of lemmas we show that the above necessary conditions are also sufficient.

**Lemma 3.1.** *If  $m, n \in 2\mathbb{Z}_+$  with  $2 \leq m \leq n \leq 8$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ .*

*Proof. Case 1.* For  $m = 2$  and  $n = 2$ , trivially one  $C_4$ . For  $n = 4$ , let  $V(K_{2,4}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, x_{12}\}$  and  $X_2 = \{x_{21}, \dots, x_{24}\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

1.  $p = 2$  and  $q = 0$ .  
By **Theorem 1.1**, we get the required  $2C_4$ 's.
2.  $p = 0$  and  $q = 2$ .  
By **Theorem 1.2**, we get the required  $2S_4$ 's.

For  $n = 6$ , let  $V(K_{2,6}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, x_{12}\}$  and  $X_2 = \{x_{21}, \dots, x_{26}\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

1.  $p = 3$  and  $q = 0$ .  
By **Theorem 1.1**, we get the required  $3C_4$ 's.
2.  $p = 1$  and  $q = 2$ .  
 $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11})$  and  $(x_{11}, x_{12}; x_{23}, x_{24}, x_{25}, x_{26})$ .

For  $n = 8$ , we can write,  $K_{2,8} = K_{2,4} \oplus K_{2,4}$ . Then the graph  $K_{2,4}$  has a  $\{pC_4, qS_4\}$ -decomposition, by the starting of the proof. Hence, by the remark, the graph  $K_{2,8}$  has a desired decomposition.

**Case 2.** For  $m = 4$  and  $n = 4$ , let  $V(K_{4,4}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, \dots, x_{14}\}$  and  $X_2 = \{x_{21}, \dots, x_{24}\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

- $p = 4$  and  $q = 0$ .  
 $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13})$  and  $(x_{13}, x_{23}, x_{14}, x_{24}, x_{13})$ .
- $p = 2$  and  $q = 2$ .  
 The first two  $C_4$ 's in (1) and the  $2S_4$ 's  $(x_{13}, x_{14}; x_{21}, x_{22}, x_{23}, x_{24})$  gives the required decomposition.
- $p = 0$  and  $q = 4$ .  
 By [Theorem 1.2](#), we get the required  $4S_4$ 's.

For  $m = 4$  and  $n = 6$ , let  $V(K_{4,6}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, \dots, x_{14}\}$  and  $X_2 = \{x_{21}, \dots, x_{26}\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

- $p = 6$  and  $q = 0$ .  
 $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11}), (x_{11}, x_{25}, x_{12}, x_{26}, x_{11}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13}), (x_{13}, x_{23}, x_{14}, x_{24}, x_{13})$  and  $(x_{13}, x_{25}, x_{14}, x_{26}, x_{13})$ .
- $p = 4$  and  $q = 2$ .  
 The first four  $C_4$ 's in (1) and the  $2S_4$ 's  $(x_{13}, x_{14}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 3$  and  $q = 3$ .  
 The first three  $C_4$ 's in (1) and the  $3S_4$ 's  $(x_{11}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{12}; x_{21}, x_{22}, x_{25}, x_{26}), (x_{13}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 2$  and  $q = 4$ .  
 The  $2C_4$ 's  $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13})$  and the  $4S_4$ 's  $(x_{11}, x_{12}, x_{13}, x_{14}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 0$  and  $q = 6$ .  
 By [Theorem 1.2](#), we get the required  $6S_4$ 's.

For  $n = 8$ , we can write,  $K_{4,8} = K_{4,6} \oplus K_{4,2}$ . Both the graphs  $K_{4,6}$  and  $K_{4,2}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{4,8}$  has a desired decomposition.

**Case 3.** For  $m = 6$  and  $n = 6$ , let  $V(K_{6,6}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, \dots, x_{16}\}$  and  $X_2 = \{x_{21}, \dots, x_{26}\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

- $p = 9$  and  $q = 0$ .  
 $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11}), (x_{11}, x_{25}, x_{12}, x_{26}, x_{11}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13}), (x_{13}, x_{23}, x_{14}, x_{24}, x_{13}), (x_{13}, x_{25}, x_{14}, x_{26}, x_{13}), (x_{15}, x_{21}, x_{16}, x_{22}, x_{15}), (x_{15}, x_{23}, x_{16}, x_{24}, x_{15}), (x_{15}, x_{25}, x_{16}, x_{26}, x_{15})$ .
- $p = 7$  and  $q = 2$ .  
 The first four and last three  $C_4$ 's in (1) and the  $2S_4$ 's  $(x_{13}, x_{14}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 6$  and  $q = 3$ .  
 The first three and last three  $C_4$ 's in (1) and the  $3S_4$ 's  $(x_{11}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{12}; x_{21}, x_{22}, x_{25}, x_{26}), (x_{13}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 5$  and  $q = 4$ .  
 The last three  $C_4$ 's in (1) along with  $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13})$  and the  $4S_4$ 's  $(x_{11}, x_{12}, x_{13}, x_{14}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.

- $p = 4$  and  $q = 5$ .  
 The  $4C_4$ 's  $(x_{11}, x_{25}, x_{13}, x_{26}, x_{11}), (x_{12}, x_{23}, x_{14}, x_{24}, x_{12}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13}), (x_{15}, x_{21}, x_{16}, x_{22}, x_{15})$  and the  $5S_4$ 's  $(x_{11}, x_{12}; x_{21}, x_{22}, x_{25}, x_{26}), (x_{13}, x_{15}, x_{16}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 3$  and  $q = 6$ .  
 The  $3C_4$ 's  $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{13}, x_{21}, x_{14}, x_{22}, x_{13}), (x_{15}, x_{21}, x_{16}, x_{22}, x_{15})$  and the  $6S_4$ 's  $(x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}; x_{23}, x_{24}, x_{25}, x_{26})$  gives the required decomposition.
- $p = 2$  and  $q = 7$ .  
 The  $2C_4$ 's  $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11})$  and the  $7S_4$ 's  $(x_{13}; x_{21}, x_{22}, x_{23}, x_{26}), (x_{14}; x_{21}, x_{22}, x_{23}, x_{26}), (x_{15}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{16}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{24}; x_{11}, x_{12}, x_{13}, x_{14}), (x_{25}; x_{13}, x_{14}, x_{15}, x_{16}), (x_{26}; x_{11}, x_{12}, x_{15}, x_{16})$  gives the required decomposition.
- $p = 1$  and  $q = 8$ .  
 $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11})$  and the  $8S_4$ 's  $(x_{15}, x_{16}; x_{23}, x_{24}, x_{25}, x_{26}), (x_{21}, x_{22}, x_{23}, x_{24}, x_{25}, x_{26}; x_{11}, x_{12}, x_{13}, x_{14})$  gives the required decomposition.
- $p = 0$  and  $q = 9$ .  
 By [Theorem 1.2](#), we get the required  $9S_4$ 's.

For  $n = 8$ , we can write,  $K_{6,8} = K_{6,4} \oplus K_{6,4}$ . Then we obtained  $(p, q) \in \{(12, 0), (10, 2), \dots, (2, 10), (0, 12)\}$ . The case  $(1, 11)$  can be obtain by taking,  $K_{6,8} = K_{6,2} \oplus K_{6,6}$ , we have  $(1, 11) = (1, 2) + (0, 9)$ , by the Cases 1 and 3 above procedure. Hence, the graph  $K_{6,8}$  has a desired decomposition.

**Case 4.** For  $m = 8$  and  $n = 8$ , we can write,  $K_{8,8} = 2K_{4,6} \oplus 2K_{2,4}$ . By Case 1 above, we obtained  $(p, q) \in \{(16, 0), (14, 2), \dots, (2, 14), (0, 16)\}$ . The case  $(1, 15)$  can be obtain by taking,  $K_{8,8} = K_{6,8} \oplus 2K_{2,4}$ , we have  $(1, 15) = (1, 11) + (0, 4)$ , by the Case 1 and 3 above procedure. Hence, the graph  $K_{8,8}$  has a desired decomposition.  $\square$

**Lemma 3.2.** If  $m, n \in 2\mathbb{Z}_+$  with  $2 \leq m \leq 8$  and  $n \geq 10$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ .

*Proof.* **Case 1.** For  $m = 2$ , we distinguish two subcases.

**Subcase 1.**  $n \equiv 2 \pmod{4}$ , we have  $n = 4x + 2$ , where  $x \geq 2$ .

For  $x = 2$  we have,  $K_{2,10} = K_{2,8} \oplus K_{2,2}$ . Then the graph  $K_{2,10}$  has a  $(4; p, q)$ -decomposition, by [Lemma 3.1](#). For  $x \geq 3$ , we can write,  $K_{2,4x+2} = K_{2,10} \oplus (x-2)K_{2,4}$ . By the above procedure and [Lemma 3.1](#), both the graphs  $K_{2,10}$  and  $K_{2,4}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{2,4x+2}$  has a desired decomposition.

**Subcase 2.**  $n \equiv 0 \pmod{4}$ , we have  $n = 4x$ , where  $x \geq 3$ . We can write,  $K_{2,4x} = xK_{2,4}$ . Hence, the graph  $K_{2,4x}$  has a  $\{pC_4, qS_4\}$ -decomposition, by [Lemma 3.1](#).

For  $m = 4, 6$  and  $n = 10$ , we can write,  $K_{m,10} = K_{m,6} \oplus K_{m,4}$ . Hence, the graph  $K_{m,10}$  has a  $\{pC_4, qS_4\}$ -decomposition, by [Lemma 3.1](#).

Let  $n > 10$ , we have  $n = 4x + y$ , where  $3 \leq x \in \mathbb{Z}_+$  and  $y = 0, 2$ . We can write,  $K_{m,n} = K_{m,4x+y}$ .

**Case 2.** For  $m = 4$ , we can write,  $K_{4,4x+y} = K_{4,10} \oplus \left(\frac{4x+y}{2} - 5\right)K_{4,2}$ . By the above procedure and [Lemma 3.1](#),

both the graphs  $K_{4,10}$  and  $K_{4,2}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{4,4x+y}$  has the desired decomposition.

**Case 3.** For  $m = 6$ . We distinguish two subcases.

**Subcase 1.**  $n \equiv 2 \pmod{4}$ , we have  $n = 4x + 2$ , where  $3 \leq x \in \mathbb{Z}_+$ , we can write,  $K_{6,4x+2} = (x - 1)K_{6,4} \oplus K_{6,6}$ . Hence, the graph  $K_{6,4x+2}$  has a  $\{pC_4, qS_4\}$ -decomposition, by Lemma 3.1.

**Subcase 2.**  $n \equiv 0 \pmod{4}$ , we have  $n = 4x$ , where  $3 \leq x \in \mathbb{Z}_+$ , we can write,  $K_{6,4x} = (x - 1)K_{6,4} \oplus K_{6,4}$ . Hence, the graph  $K_{6,4x}$  has a  $\{pC_4, qS_4\}$ -decomposition, by Lemma 3.1.

**Case 4.** For  $m = 8$  and  $n = 2x$ , where  $x \geq 5$ , we can write,  $K_{8,2x} = K_{8,8} \oplus (x - 4)K_{8,2}$ . Note that  $K_{8,2} \cong K_{2,8}$ . Hence, the graph  $K_{8,2x}$  has a  $\{pC_4, qS_4\}$ -decomposition, by Lemma 3.1.  $\square$

**Lemma 3.3.** *If  $m \in 2\mathbb{Z}_+$  with  $m \equiv 0 \pmod{4} \geq 12$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{\frac{m}{2},m}$ , where  $q \neq 1$ .*

*Proof.* We distinguish two cases.

**Case 1.**  $m = 12$  and  $q \neq 1$ .

We can write,  $K_{6,12} = K_{6,6} \oplus K_{6,6}$ . By Lemma 3.1, the graph  $K_{6,6}$  has a  $\{pC_4, qS_4\}$ -decomposition. Hence, the graph  $K_{6,12}$  has the desired decomposition

**Case 2.**  $m > 12$  and  $q \neq 1$ . We can write,

$$K_{\frac{m}{2},m} = K_{6,12} \oplus K_{6,m-12} \oplus \left(\frac{m-12}{4}\right)(K_{2,12} \oplus K_{2,m-12}).$$

By the Case 1 above and Lemma 3.2, the graphs  $K_{6,12}, K_{2,12}$  and  $K_{2,m-12}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{\frac{m}{2},m}$  has the desired decomposition.  $\square$

**Lemma 3.4.** *If  $m \in 2\mathbb{Z}_+$  with  $m \equiv 0 \pmod{4} \geq 12$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,m}$ , where  $q \neq 1$ .*

*Proof.* For  $m \geq 12$  and  $q \neq 1$ . We can write,

$$K_{m,m} = K_{8,8} \oplus \left(\frac{m-12}{4}\right)K_{4,4} \oplus \left\{ \bigoplus_i^{m-4} K_{i,4} \right\} \oplus \left\{ \bigoplus_j^{m-4} K_{4,j} \right\},$$

where  $i \equiv 0 \pmod{4} \geq 4$  and  $j \equiv 0 \pmod{4} \geq 8$ . Note that  $K_{i,4} \cong K_{4,i}$ . By Lemmas 3.1 and 3.2, the graphs  $K_{4,4}, K_{8,8}, K_{i,4}$  and  $K_{4,j}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{m,m}$  has the desired decomposition.  $\square$

**Lemma 3.5.** *If  $m \in 2\mathbb{Z}_+$  with  $m \equiv 2 \pmod{4} \geq 10$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,m}$ , where  $q \neq 1$ .*

*Proof.* We can write,

$$K_{m,m} = K_{6,6} \oplus \left(\frac{m-6}{4}\right)K_{4,4} \oplus \left\{ \bigoplus_i^{m-4} K_{4,i} \right\} \oplus \left\{ \bigoplus_j^{m-4} K_{j,4} \right\},$$

where  $i, j \equiv 2 \pmod{4} \geq 6$ . Note that  $K_{j,4} \cong K_{4,j}$ . By Lemmas 3.1 and 3.2, the graphs  $K_{4,4}, K_{6,6}, K_{4,i}$  and  $K_{j,4}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{m,m}$  has the desired decomposition.  $\square$

**Lemma 3.6.** *If  $m, n \in 2\mathbb{Z}_+$  with  $n \geq m \geq 12$  and  $m, n \equiv 0 \pmod{4}$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \neq 1$ .*

*Proof.* We can write,

$$K_{m,n} = K_{m,m} \oplus \left(\frac{n-m}{4}\right)K_{m,4}.$$

Note that  $K_{m,4} \cong K_{4,m}$ . By Lemmas 3.2 and 3.4, the graphs  $K_{m,4}$  and  $K_{m,m}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{m,n}$  has the desired decomposition.  $\square$

**Lemma 3.7.** *If  $m, n \in 2\mathbb{Z}_+$  with  $n \geq m \geq 12$ ;  $m \equiv 0 \pmod{4}$  and  $n \equiv 2 \pmod{4}$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \neq 1$ .*

*Proof.* Let  $m = 4x$  and  $n = 4y + 2$ , where  $x, y \in \mathbb{Z}_+$  and  $y \geq x \geq 3$ . Hence  $K_{m,n} = K_{4x,4y+2}$ . We can write,  $K_{4x,4y+2} = K_{4x,4(y-1)} \oplus K_{4x,4+2}$ . Since the graph  $K_{4x,4(y-1)}$  can be viewed as  $(y - 1)$  copies of  $K_{4x,4}$ . Note that  $K_{4x,6} \cong K_{6,4x}$ . By Lemma 3.2, both the graphs  $K_{4x,4}$  and  $K_{4x,6}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{m,n}$  has the desired decomposition.  $\square$

**Lemma 3.8.** *If  $m, n \in 2\mathbb{Z}_+$  with  $n \geq m \geq 10$ ;  $m \equiv 2 \pmod{4}$  and  $n \equiv 0 \pmod{4}$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \neq 1$ .*

*Proof.* By the similar argument as in Lemma 3.7, we get a required decomposition.  $\square$

**Lemma 3.9.** *If  $m, n \in 2\mathbb{Z}_+$  with  $n \geq m \geq 10$ ;  $m, n \equiv 2 \pmod{4}$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \neq 1$ .*

*Proof.* We can write,

$$K_{m,n} = K_{m,m} \oplus \left(\frac{n-m}{4}\right)K_{m,4}.$$

Note that  $K_{m,4} \cong K_{4,m}$ . By Lemma 3.2 and 3.5, the graphs  $K_{m,4}$  and  $K_{m,m}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{m,n}$  has the desired decomposition.  $\square$

**Lemma 3.10.** *If  $m \in \{3, 5, 7\}$  and  $n = 4$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$ .*

*Proof.* We distinguish three cases.

**Case 1.** For  $m = 3$  and  $n = 4$ . Let  $V(K_{3,4}) = (X_1, X_2)$ , where  $X_1 = \{x_{11}, x_{12}, x_{13}\}$  and  $X_2 = \{x_{21}, \dots, x_{24}\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

1.  $p = 0$  and  $q = 3$ .  
 $(x_{11}, x_{12}, x_{13}; x_{21}, x_{22}, x_{23}, x_{24}).$



2.  $p = 2$  and  $q = 1$ .

$(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11})$  and  $(x_{13}; x_{21}, x_{22}, x_{23}, x_{24})$ .

**Case 2.** For  $m = 5$  and  $n = 4$ . We can write,  $K_{5,4} = K_{4,4} \oplus K_{1,4}$ . Note that  $K_{5,4} \cong K_{4,5}$ . By Lemma 3.1, the graph  $K_{4,4}$  has a  $\{pC_4, qS_4\}$ -decomposition and trivially the graph  $K_{1,4}$  is  $S_4$ . Hence, the graph  $K_{5,4}$  has the desired decomposition.

**Case 3.** For  $m = 7$  and  $n = 4$ . We can write,  $K_{7,4} = K_{6,4} \oplus K_{1,4}$ . By Lemma 3.1 the graph  $K_{6,4}$  has a  $\{pC_4, qS_4\}$ -decomposition and trivially the graph  $K_{1,4}$  is  $S_4$ . Hence, the graph  $K_{7,4}$  has the desired decomposition.  $\square$

**Lemma 3.11.** *If  $m = 3$  and  $n \in 2\mathbb{Z}_+$  with  $n \equiv 0 \pmod{4} \geq 8$ , then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \geq \frac{n}{4}$ .*

*Proof.* We distinguish two cases.

**Case 1.** For  $m = 3$  and  $n = 8$ , let  $V(K_{3,8}) = V(X_1, X_2)$ , where  $X_1 = \{x_{11}, x_{12}, x_{13}\}$ ,  $X_2 = \{x_{21}, \dots, x_{28}\}$  and  $E(K_{3,8}) = \{x_i x_{2j} \mid i = 1, 2, 3 \text{ and } j = 1, \dots, 8\}$ . Then the required  $\{pC_4, qS_4\}$ -decompositions are as given below:

1.  $p = 4$  and  $q = 2$ .

The  $4C_4$ 's  $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11}), (x_{11}, x_{25}, x_{12}, x_{26}, x_{11}), (x_{11}, x_{27}, x_{12}, x_{28}, x_{11})$  and the  $2S_4$ 's  $(x_{13}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{13}; x_{25}, x_{26}, x_{27}, x_{28})$  gives the required decomposition.

2.  $p = 3$  and  $q = 3$ .

The  $3C_4$ 's  $(x_{11}, x_{21}, x_{23}, x_{22}, x_{11}), (x_{11}, x_{24}, x_{13}, x_{25}, x_{11}), (x_{12}, x_{26}, x_{13}, x_{28}, x_{12})$  and the  $3S_4$ 's  $(x_{11}; x_{22}, x_{26}, x_{27}, x_{28}), (x_{12}; x_{22}, x_{24}, x_{25}, x_{27}), (x_{13}; x_{21}, x_{22}, x_{23}, x_{27})$  gives the required decomposition.

3.  $p = 2$  and  $q = 4$ .

The  $2C_4$ 's  $(x_{11}, x_{21}, x_{12}, x_{22}, x_{11}), (x_{11}, x_{23}, x_{12}, x_{24}, x_{11})$  and the  $4S_4$ 's  $(x_{13}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{11}, x_{12}, x_{13}; x_{25}, x_{26}, x_{27}, x_{28})$  gives the required decomposition.

4.  $p = 0$  and  $q = 6$ .

The  $6S_4$ 's  $(x_{11}, x_{12}, x_{13}; x_{21}, x_{22}, x_{23}, x_{24}), (x_{11}, x_{12}, x_{13}; x_{25}, x_{26}, x_{27}, x_{28})$ .

**Case 2.** For  $m = 3, n > 8$  and  $q \geq \frac{n}{4}$ . We can write,

$$K_{3,n} = K_{3,8} \oplus \left(\frac{n-8}{4}\right)K_{3,4}.$$

By Lemma 3.10 and the Case 1 above, the graphs  $K_{3,4}$  and  $K_{3,8}$  have a  $\{pC_4, qS_4\}$ -decomposition. Hence, by the remark, the graph  $K_{3,n}$  has the desired decomposition.  $\square$

**Lemma 3.12.** *Let  $m$  be an odd integer and  $n \in 2\mathbb{Z}_+$  with  $2 < m < n$  and  $n \equiv 0 \pmod{4} \geq 4$ . Then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \geq \frac{n}{4}$ .*

*Proof.* Let  $m = 4x + s$  and  $n = 4y$ , where  $x, y \in \mathbb{Z}_+$  with  $y \geq x \geq 1$  and  $s = 1, 3$ . We can write,  $K_{m,n} = K_{m-1,n} \oplus K_{1,n}$ , we have  $K_{4x+s,4y} = K_{4x+s-1,4y} \oplus K_{1,4y}$ . Since  $K_{4x+s-1,4y}$  can

be viewed as  $x$  copies of  $K_{4+s-1,4y}$ . Then the graph  $K_{4+s-1,4y}$  has a  $\{pC_4, qS_4\}$ -decomposition, by Lemma 3.2 and trivially the graph  $K_{1,4y}$  has an  $S_4$ -decomposition. Hence, the graph  $K_{4x+s,4y}$  has the desired decomposition.  $\square$

**Lemma 3.13.** *Let  $n$  be an odd integer and  $m \equiv 0 \pmod{4}, n > m \geq 4$ . Then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  where  $q \geq \frac{m}{4}$ .*

*Proof.* By the similar argument as in Lemma 3.12, we get a required decomposition.  $\square$

## 4. Conclusion

As a consequence of Lemmas 2.1–2.4 and 3.1–3.13, our main result immediately follows.

**Theorem 4.1.** *Let  $p$  and  $q$  be nonnegative integers, and let  $m$  and  $n$  be positive integers such that  $m \leq n$ . Then there exists a  $\{pC_4, qS_4\}$ -decomposition of  $K_{m,n}$  if and only if one of the following holds:*

1.  $q$  is even, when  $m = 2$  and even  $n \geq 2$ ;
2.  $p, q \neq 1$ , when  $m = 4$  and even  $n \geq 4$ ;
3.  $q \neq 1$  when even  $m, n \geq 6$ ;
4.  $q \geq \frac{n}{4}$  (resp.,  $q \geq \frac{m}{4}$ ), when  $m$  (resp.,  $n$ ) is an odd integer and  $n \equiv 0 \pmod{4}$  (resp.,  $m \equiv 0 \pmod{4}$ ).

## Acknowledgments

The authors thank the University Grant Commission, Government of India, New Delhi for its support through the Grant No.F.510/7/DRS-I/2016(SAP-I). Also would like to thank Dr. A. Shanmuga Vadivu (NBHM-PDF) whose contribution was helpful in developing ideas for this paper.

## Disclosure statement

The following authors have affiliations with organizations with direct or indirect partial financial support in the subject matter discussed in the manuscript: M. Ilayaraja, Periyar University, Salem, Tamil Nadu, India.

## References

- [1] Abueida, A. A, Daven, M. (2003). Multidesigns for graph-pairs of order 4 and 5. *Graphs Combin* 19(4):433–447.
- [2] Abueida, A. A, O'Neil, T. (2007). Multidecomposition of  $\lambda K_m$  into small cycles and claws. *Bull. Inst. Combin. Appl.* 49:32–40.
- [3] Alspach, B, Gavlas, H. (2001). Cycle decompositions of  $K_n$  and  $K_n - I$ . *J. Combin. Theory Ser. B* 81(1):77–99.
- [4] Bondy, J. A, Murty, U. R. S. (1976). *Graph Theory with Applications*. New York: The Macmillan Press Ltd.,
- [5] Jeevadoss, S, Muthusamy, A. (2014). Decomposition of complete bipartite graphs into paths and cycles. *Discrete Math.* 331: 98–108.
- [6] Lee, H. C. (2015). Decomposition of the complete bipartite multigraph into cycles and stars. *Discrete Math.* 338(8): 1362–1369.
- [7] Lee, H.-C, Chu, Y.-P. (2013). Multidecompositions of complete bipartite graphs into cycles and stars. *Ars Combin.* 2013:1–364.

- [8] Lee, H. C, Lin, J.-J. (2013). Decomposition of the complete bipartite graph with a 1-factor removed into cycles and stars. *Discret Math.* 313(20):2354–2358.
- [9] Priyadharsini, H. M, Muthusamy, A. (2012).  $(G_m, H_m)$ -multidecomposition of  $K_{m,m}(\lambda)$ . *Bull. Inst. Combin. Appl.* 66: 42–48.
- [10] Shyu, T.-W. (2013). Decomposition of complete bipartite graphs into paths and stars with same number of edges. *Discret. Math.* 313(7):865–871.
- [11] Sotteau, D. (1981). Decomposition of  $K_{m,n}(K_{m,n}^*)$  into cycles (circuits) of length  $2k$ . *J. Combin. Theory Ser. B* 30(1):75–81.
- [12] Sajna, M. (2002). Cycle decompositions III; complete graphs and fixed length cycles. *J. Combin. Des.* 10:27–78.
- [13] Ushio, K, Tazawa, S, Yamamoto, S. (1978). On claw-decomposition of complete multipartite graphs. *Hiroshima Math. J.* 8(1):207–210.
- [14] Yamamoto, S., Ikeda, H., Shige-Eda, S., Ushio, K, Hamada, N. (1975). On claw decomposition of complete graphs and complete bipartite graphs. *Hiroshima Math. J.* 5(1):33–42.