

SENSMG: First-Order Sensitivities of Neutron Reaction Rates, Reaction-Rate Ratios, Leakage, k_{eff} , and α Using PARTISN

Jeffrey A. Favorite

To cite this article: Jeffrey A. Favorite (2018) SENSMG: First-Order Sensitivities of Neutron Reaction Rates, Reaction-Rate Ratios, Leakage, k_{eff} , and α Using PARTISN, Nuclear Science and Engineering, 192:1, 80-114, DOI: [10.1080/00295639.2018.1471296](https://doi.org/10.1080/00295639.2018.1471296)

To link to this article: <https://doi.org/10.1080/00295639.2018.1471296>



© J. A. Favorite. Published with license by Taylor & Francis.



Published online: 23 Jul 2018.



Submit your article to this journal [↗](#)



Article views: 665



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 3 View citing articles [↗](#)

SENSMG: First-Order Sensitivities of Neutron Reaction Rates, Reaction-Rate Ratios, Leakage, k_{eff} , and α Using PARTISN

Jeffrey A. Favorite*

Los Alamos National Laboratory, Computational Physics (X-CP) Division, MS F663, Los Alamos, New Mexico 87545

Received March 26, 2018

Accepted for Publication April 26, 2018

Abstract — *SENSMG* is a tool for calculating the first-order sensitivities of reaction-rate ratios, k_{eff} , and α in critical problems and reaction-rate ratios, reaction rates, and leakage in fixed-source problems to multigroup cross sections, isotope densities, material mass densities, and interface locations using the PARTISN multigroup discrete-ordinates code by implementing Generalized Perturbation Theory. *SENSMG* can be used for one-dimensional spherical and slab (r) and two-dimensional cylindrical (r - z) geometries. For fixed-source (leakage) problems, *SENSMG* relies on the MISC and/or SOURCES4C codes to compute neutron source rate densities from spontaneous fission and (α , n) sources. *SENSMG* is a combination of Python and Fortran and was developed under Linux. This computer code abstract describes all user inputs, the input file, and output files. This computer code abstract describes how *SENSMG* can be modified to support different computer platforms, PARTISN versions, or cross-section availability. Several verification problems are presented in which *SENSMG* results are compared with MCNP6, SCALE6.2, and direct perturbations (central differences). *SENSMG* is available at <https://github.com/jafavorite/SENSMG>. *SENSMG* can be modified to accommodate other deterministic transport codes that have an adjoint capability.

Keywords — Multigroup, discrete ordinates, adjoint, first-order sensitivity analysis.

Note — Some figures may be in color only in the electronic version.

I. INTRODUCTION

SENSMG is a tool for computing first-order sensitivities of neutron reaction rates, reaction-rate ratios, leakage, k_{eff} , and α using the PARTISN multigroup discrete-ordinates code.¹ Both codes are developed at Los Alamos National Laboratory (LANL). SENS MG computes sensitivities to all of the transport cross sections and data (total, fission,^a nu, chi, and all scattering moments), two edit cross sections

(absorption and capture), and the density for every isotope and energy group. The tool can be used for one-dimensional spherical and slab (r) and two-dimensional cylindrical (r - z) geometries. The tool can be used for fixed-source and eigenvalue problems. The tool implements Generalized Perturbation Theory (GPT) as discussed by Williams² and Stacey.³ The tool is thus limited to computing sensitivities only for GPT-allowable responses.⁴

Neutron leakage is computed only for fixed-source problems, for which an additional code, MISC (Ref. 5) and/or SOURCES4C (Ref. 6), is used to compute the neutron source rate density from spontaneous fission and (α , n) reactions based on the materials present in the geometry. MISC and SOURCES4C are both developed at LANL.

SENSMG is a combination of Python and Fortran and was developed under Linux. SENS MG is available at <https://github.com/jafavorite/SENSMG>.

*E-mail: fave@lanl.gov

^a Technically, fission is an edit cross section in PARTISN because only the product with nu is used in the transport.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Section II briefly describes the theory behind adjoint-based sensitivities, gives the equations that SENSIMG solves, and defines the sensitivities that are output. Section III describes the user interface, including the input file and command line options. Section IV describes the output. Section V gives some notes about the coding that may be of interest. Section VI discusses verification, which is ongoing. Section VII lists needs and ideas for future work. Appendix A lists all of the input files whose results are presented in Sec. VI. Appendix B provides some useful details on one of the cross-section libraries that SENSIMG supports.

II. THEORY AND DEFINITIONS

II.A. Reaction Rates, Reaction-Rate Ratios, and Sensitivities

Define two reaction rates as

$$R_1 \equiv \int_V dV \sum_{g=1}^G \Sigma_1^g(r) \phi^g(r) \quad (1)$$

and

$$R_2 \equiv \int_V dV \sum_{g=1}^G \Sigma_2^g(r) \phi^g(r) , \quad (2)$$

where

- V = problem spatial domain
- r = a point within the volume V
- g = energy group index
- G = number of energy groups

$\Sigma_1^g(r), \Sigma_2^g(r)$ = arbitrary macroscopic cross sections in group g at point r .

Also in Eqs. (1) and (2), $\phi^g(r)$ is the scalar flux in group g at point r , defined as

$$\phi^g(r) \equiv \int_{4\pi} d\hat{\Omega} \psi^g(r, \hat{\Omega}) , \quad (3)$$

where $\psi^g(r, \hat{\Omega})$ is the angular flux in group g at point r in direction $\hat{\Omega}$. The reaction cross sections $\Sigma_1^g(r)$ and $\Sigma_2^g(r)$ are for arbitrary nuclides that need not be present in the problem. They are defined to be zero outside of a subset of V within which the reaction rates are assumed to be measured.

The ratio of the reaction rates is

$$R \equiv \frac{R_1}{R_2} = \frac{\int_V dV \sum_{g=1}^G \Sigma_1^g(r) \phi^g(r)}{\int_V dV \sum_{g=1}^G \Sigma_2^g(r) \phi^g(r)} . \quad (4)$$

The relative sensitivity $S_{R,x}^g$ of R to a transport cross section σ_x^g for reaction x in group g is defined to be

$$S_{R,x}^g \equiv \left. \frac{\sigma_{x,0}^g}{R_0} \frac{\partial R}{\partial \sigma_x^g} \right|_{\sigma_x^g = \sigma_{x,0}^g} , \quad (5)$$

where subscript 0 indicates the initial base case. The transport cross section σ_x^g is for a specific nuclide that is present in the problem.

The derivative of R with respect to σ_x^g is found by differentiating Eq. (4) with respect to σ_x^g , noting that g denotes a specific energy group. The result is

$$\frac{\partial R}{\partial \sigma_x^g} = \frac{1}{R_2} \left(\int_V dV \frac{\partial \Sigma_1^g(r)}{\partial \sigma_x^g} \phi^g(r) + \int_V dV \Sigma_1^g(r) \frac{\partial \phi^g(r)}{\partial \sigma_x^g} \right) - \frac{R_1}{R_2^2} \left(\int_V dV \frac{\partial \Sigma_2^g(r)}{\partial \sigma_x^g} \phi^g(r) + \int_V dV \Sigma_2^g(r) \frac{\partial \phi^g(r)}{\partial \sigma_x^g} \right) . \quad (6)$$

Rearranging Eq. (6) and using the result in Eq. (5) yields

$$S_{R,x}^g = \left(\frac{\sigma_x^g}{R_1} \int_V dV \frac{\partial \Sigma_1^g(r)}{\partial \sigma_x^g} \phi^g(r) - \frac{\sigma_x^g}{R_2} \int_V dV \frac{\partial \Sigma_2^g(r)}{\partial \sigma_x^g} \phi^g(r) \right) + \left(\frac{\sigma_x^g}{R_1} \int_V dV \Sigma_1^g(r) \frac{\partial \phi^g(r)}{\partial \sigma_x^g} - \frac{\sigma_x^g}{R_2} \int_V dV \Sigma_2^g(r) \frac{\partial \phi^g(r)}{\partial \sigma_x^g} \right) . \quad (7)$$

In Eqs. (6) and (7) and the rest of this computer code abstract, we assume that all quantities are evaluated for the base configuration.

The total first-order sensitivity of R to reaction x is the sum over groups:

$$S_{R,x} = \sum_{g=1}^G S_{R,x}^g . \quad (8)$$

As shown in Eq. (7), the relative sensitivity is the sum of two components. The first term in parentheses on the right side is called the *direct effect* of a perturbation in σ_x^g and the second term in parentheses is called the *indirect effect*.

The derivatives in the direct effect are nonzero only when $\Sigma_1^g = N_1 \sigma_1^g = N_1 \sigma_x^g$ and/or $\Sigma_2^g = N_2 \sigma_2^g = N_2 \sigma_x^g$, where N_1 and N_2 are the atom densities for nuclides 1 and 2. The direct effect can therefore be written

$$\begin{aligned} (S_{R,x}^g)_{direct} &= \frac{\int_V dV \Sigma_1^g(r) \phi^g(r)}{R_1} \delta_{x1} \\ &- \frac{\int_V dV \Sigma_2^g(r) \phi^g(r)}{R_2} \delta_{x2} , \end{aligned} \quad (9)$$

where δ_{xj} ($j = 1, 2$) is the Kronecker delta that is equal to one when $x = j$ and 0 otherwise. Equation (9) can readily be evaluated given the forward scalar flux $\phi^g(r)$.

Rearranging the second term in parentheses in Eq. (7), the indirect effect, yields

$$(S_{R,x}^g)_{indirect} = \sigma_x^g \int_V dV \left(\frac{\Sigma_1^g(r)}{R_1} - \frac{\Sigma_2^g(r)}{R_2} \right) \frac{\partial \phi^g(r)}{\partial \sigma_x^g} . \quad (10)$$

This term uses the derivative of the scalar flux with respect to σ_x^g . GPT provides an economical means of computing the indirect effect.

II.B. k_{eff} -Eigenvalue Problems

Details of applying GPT to λ -mode (k_{eff}) eigenvalue problems are given in Ref. 2; a good summary is given in Ref. 7. Very briefly, for k_{eff} calculations, the forward and adjoint neutron fluxes $\psi^g(r, \hat{\Omega})$ and $\psi^{*g}(r, \hat{\Omega})$ satisfy

$$(A - \lambda F)\psi = 0 \quad (11)$$

and

$$(A^* - \lambda F^*)\psi^* = 0 , \quad (12)$$

respectively, where

F = fission neutron production operator

A = neutron loss operator

$\lambda = 1/k_{eff}$

$*$ = adjoint operators.

Vacuum boundary conditions apply to Eqs. (11) and (12). The indirect effect [Eq. (10)] is

$$\begin{aligned} (S_{R,x}^g)_{indirect} &= \sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \\ &\times \left(\lambda \frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega}) , \end{aligned} \quad (13)$$

where $\Gamma^{*g}(r, \hat{\Omega})$ is the generalized adjoint function that satisfies

$$(A^* - \lambda F^*)\Gamma^* = S^* , \quad (14)$$

with

$$S^{*g}(r, \hat{\Omega}) = \frac{1}{R} \frac{\partial R}{\partial \phi} = \frac{\Sigma_1^g(r)}{R_1} - \frac{\Sigma_2^g(r)}{R_2} , \quad (15)$$

which is an isotropic volume source. A vacuum boundary condition applies to Eq. (14). Because the operator acting on Γ^* in Eq. (14) is singular [see Eq. (12)], special techniques are required to solve Eq. (14). Another way of stating this is that Eq. (14) represents a source inserted into a critical system. One technique is the method of successive approximations,^{3,8} which calculates $\Gamma^{*g}(r, \hat{\Omega})$ using

$$\Gamma^{*g}(r, \hat{\Omega}) = \sum_{n=0}^{\infty} \Gamma_n^{*g}(r, \hat{\Omega}) , \quad (16)$$

where

$$\begin{aligned} \Gamma_n^{*g}(r, \hat{\Omega}) &= \xi_n^{*g}(r, \hat{\Omega}) \\ &- \frac{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \xi_n^{*g}(r, \hat{\Omega}) F \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) F \psi^g(r, \hat{\Omega})} \\ &\times \psi^{*g}(r, \hat{\Omega}) . \end{aligned} \quad (17)$$

The $\xi_n^{*g}(r, \hat{\Omega})$ are generated recursively using

$$\begin{cases} A^* \xi_0^* = S^*, \\ A^* \xi_n^* = \lambda F^* \Gamma_{n-1}^*, n > 0. \end{cases} \quad (18)$$

The second term on the right side of Eq. (17) is included to remove any fundamental mode contamination. Williams² has a thorough discussion of this issue.

The generalized adjoint for k_{eff} -eigenvalue problems must satisfy²

$$\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) F \psi^g(r, \hat{\Omega}) = 0. \quad (19)$$

In fact, as Cacuci has shown,⁴ Eq. (19) represents a limitation of GPT and thus of SENSIMG because only those responses whose generalized adjoints satisfy Eq. (19) are GPT-allowable responses that can be treated with GPT and thus SENSIMG. In Ref. 4, Cacuci has also proposed an innovative framework that alleviates the shortcomings of GPT, but that framework has not been implemented in PARTISN or in SENSIMG.

In practice, of course, the sum in Eq. (16) goes until some user-defined convergence criterion is achieved. Cacuci's framework⁴ avoids the use of successive approximations altogether.

A more efficient method of solving Eq. (14) was recently implemented into Oak Ridge National Laboratory's (ORNL's) Denovo code.⁹ The method

takes advantage of Denovo's generalized minimal residual method (GMRES) and Krylov solvers.

It may also be of interest to compute the sensitivity of an "absolute" reaction rate, such as R_1 of Eq. (1). There is no absolute reaction rate in a critical system unless the amplitude of the flux is constrained. The power level is often used as a constraint on the flux, but for critical experiments the power level is generally unknown. In Williams's development,² the constraint is

$$P \equiv \int_V dV \sum_{g=1}^G \Sigma_p^g(r) \phi^g(r), \quad (20)$$

and P and Σ_p^g essentially replace R_2 and Σ_2^g in the equations in this section, particularly Eqs. (4), (9), and (15). In the SENSIMG sensitivity tool, we use 1 for Σ_p^g so that P becomes the flux in the reaction-rate region. (Williams² uses N where we use P .)

The λ -mode forward angular flux $\psi^g(r, \hat{\Omega})$ is computed to obtain the baseline ratio R from Eq. (4). It is also used in the solution for $\Gamma^{*g}(r, \hat{\Omega})$ through R_1 and R_2 in Eqs. (15) and through Eq. (17). In addition, Eq. (17) requires the λ -mode adjoint angular flux $\psi^{*g}(r, \hat{\Omega})$. Because the forward and adjoint fluxes are both computed, k_{eff} sensitivities can always be computed with λ -mode reaction-rate ratio sensitivities.

The relative sensitivity $S_{k,x}^g$ of k_{eff} to a transport cross section σ_x^g for reaction x in group g is defined to be

$$S_{k,x}^g \equiv \left. \frac{\sigma_{x,0}^g \partial k_{eff}}{k_{eff,0} \partial \sigma_x^g} \right|_{\sigma_x^g = \sigma_{x,0}^g}. \quad (21)$$

The adjoint-based formulation for $S_{k,x}^g$ is

$$S_{k,x}^g = \frac{\sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \left(\lambda \frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \lambda F \psi^g(r, \hat{\Omega})}. \quad (22)$$

Again, the total sensitivity $S_{k,x}$ of k_{eff} to reaction x is the sum of $S_{k,x}^g$ over groups.

Table I lists the specific equations that are used for the indirect effect of the sensitivities in the SENSIMG sensitivity tool with the label that appears in the output (Sec. IV). In Table I, subscript i indicates an isotope. In the scattering sensitivities of Table I, $\phi_l^g(r)$ and $\Gamma_l^{*g}(r)$ are the l 'th moments of the forward flux and generalized adjoint flux, defined as

$$\phi_l^g(r) \equiv \int_{4\pi} d\hat{\Omega} P_l(\hat{\Omega}) \psi^g(r, \hat{\Omega}) \quad (23)$$

TABLE I

Reaction-Rate Ratio Sensitivities for λ-Mode Flux, Indirect Effect*

Sensitivity	Label	Equation
Total cross section	total	$(S_{R,\sigma_{t,i}}^g)_{indirect} = - \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) N_i \sigma_{t,i}^g \psi^g(r, \hat{\Omega})$
Absorption cross section	abs	$(S_{R,\sigma_{a,i}}^g)_{indirect} = - \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) N_i \sigma_{a,i}^g \psi^g(r, \hat{\Omega})$
Capture cross section	(n,g)	$(S_{R,\sigma_{c,i}}^g)_{indirect} = - \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) N_i \sigma_{c,i}^g \psi^g(r, \hat{\Omega})$
Chi (not normalized)	chi_(nn)	$(S_{R,\chi_i}^g)_{indirect} = \lambda \int_V dV \sum_{g'=1}^G \Gamma_0^{*g'}(r) \chi_i^{g' \rightarrow g} N_i v_i^{g'} \sigma_{f,i}^{g'} \phi_0^{g'}(r)$
Nu	nu	$(S_{R,\nu_i}^g)_{indirect} = \lambda \int_V dV \sum_{g'=1}^G \Gamma_0^{*g'}(r) \chi_i^{g \rightarrow g'} N_i v_i^g \sigma_{f,i}^g \phi_0^g(r)$
Fission cross section	fiss	$(S_{R,\sigma_{f,i}}^g)_{indirect} = \lambda \int_V dV \sum_{g'=1}^G \Gamma_0^{*g'}(r) \chi_i^{g \rightarrow g'} N_i v_i^g \sigma_{f,i}^g \phi_0^g(r) - \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) N_i \sigma_{f,i}^g \psi^g(r, \hat{\Omega})$
In-scattering, P^0	in-scat-0	$(S_{R,\sigma_{s,0,i}}^{g' \rightarrow g})_{indirect} = \int_V dV \sum_{g' \neq g, g'=1}^G \Gamma_0^{*g'}(r) N_i \sigma_{s,0,i}^{g' \rightarrow g} \phi_0^{g'}(r) - \int_V dV \int_{4\pi} d\hat{\Omega} \sum_{g' \neq g, g'=1}^G \Gamma^{*g'}(r, \hat{\Omega}) \times N_i \sigma_{s,0,i}^{g' \rightarrow g} \psi^{g'}(r, \hat{\Omega})$
Self-scattering, P_0	self-scat-0	$(S_{R,\sigma_{s,0,i}}^{g \rightarrow g})_{indirect} = \int_V dV \Gamma_0^{*g}(r) N_i \sigma_{s,0,i}^{g \rightarrow g} \phi_0^g(r) - \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) N_i \sigma_{s,0,i}^{g \rightarrow g} \psi^g(r, \hat{\Omega})$
Out-scattering, P^0	out-scat-0	$(S_{R,\sigma_{s,0,i}}^{g \rightarrow g'})_{indirect} = \int_V dV \sum_{g' \neq g, g'=1}^G \Gamma_0^{*g'}(r) N_i \sigma_{s,0,i}^{g \rightarrow g'} \phi_0^{g'}(r) - \int_V dV \int_{4\pi} d\hat{\Omega} \sum_{g' \neq g, g'=1}^G \Gamma^{*g'}(r, \hat{\Omega}) \times N_i \sigma_{s,0,i}^{g \rightarrow g'} \psi^{g'}(r, \hat{\Omega})$
In-scattering, $P_l, 1 \leq l \leq L$	in-scat-[l]	$(S_{R,\sigma_{s,l,i}}^{g' \rightarrow g})_{indirect} = \int_V dV \sum_{\substack{g'=1 \\ g' \neq g}}^G \Gamma_l^{*g'}(r) N_i (2l+1) \sigma_{s,l,i}^{g' \rightarrow g} \phi_l^{g'}(r)$
Self-scattering, $P_l, 1 \leq l \leq L$	self-scat-[l]	$(S_{R,\sigma_{s,l,i}}^{g \rightarrow g})_{indirect} = \int_V dV \Gamma_l^{*g}(r) N_i (2l+1) \sigma_{s,l,i}^{g \rightarrow g} \phi_l^g(r)$

(Continued)

TABLE I (Continued)

Sensitivity	Label	Equation
Out-scattering, $P_l, 1 \leq l \leq L$	out-scat-[l]	$(S_{R,\sigma_{s,i}}^{g \rightarrow g'})_{indirect} = \int_V dV \sum_{g' \neq g, g'=1}^G \Gamma_0^{*g'}(r) N_i (2l + 1) \sigma_{s,0,i}^{g \rightarrow g'} \phi_l^g(r)$
Source ^a	source	$S_{R,Q_{N_i}}^g = \int_V dV \Gamma_0^{*g}(r) \left(Q_{s,f,i}^g + N_i \frac{\partial Q_{(a,i)}^g}{\partial N_i} \right)$
Total scattering	ssctt	$(S_{R,\sigma_{s,i}}^g)_{indirect} = \sum_{l=0}^L \int_V dV \sum_{g'=1}^G \Gamma_l^{*g'}(r) N_i (2l + 1) \sigma_{s,l,i}^{g \rightarrow g'} \phi_l^g(r)$
Nuclide density ^b	density	$(S_{R,N_i}^g)_{indirect} = (S_{R,N_i}^g)_{indirect} + (S_{R,\sigma_{t,i}}^g)_{indirect} + (S_{R,\sigma_{s,i}}^g)_{indirect}$

*Except for density (see note b), these equations are also used for α -mode and fixed-source flux, but $\lambda = 1/k_{eff}$ is set to 1 and a divisor is used where needed.

^aFor leakage problems only. See Eq. (61) and the text that follows it.

^bFor leakage, the nuclide density sensitivity includes $S_{R,Q_{N_i}}^g$ as a summand.

and

$$\Gamma_l^{*g}(r) \equiv \int_{4\pi} d\hat{\Omega} P_l(-\hat{\Omega}) \Gamma^{*g}(r, \hat{\Omega}), \quad (24)$$

where $P_l(\hat{\Omega})$ are Legendre and associated Legendre polynomials appropriate for the geometry.¹ Note that the generalized adjoint moments are inner product moments defined using the Legendre polynomial for the direction opposite the direction of the angular flux.¹⁰ When λ -mode, α -mode, or fixed-source adjoint moments are needed, their definition is analogous to Eq. (24):

$$\phi_l^{*g}(r) \equiv \int_{4\pi} d\hat{\Omega} P_l(-\hat{\Omega}) \phi^{*g}(r, \hat{\Omega}). \quad (25)$$

Also note that the zeroth forward moment $\phi_0^g(r)$ is just $\phi^g(r)$ of Eq. (3). The number of moments, L in Table I, depends on the scattering order and the number of dimensions (it is equal to the scattering order in one dimension).¹

The sensitivity to the total cross section and all contributors to the total cross section have an angle integral of the product of angular fluxes. For α calculations, the integrals in Eq. (34) do as well. If discrete-ordinates angular fluxes are available, then these integrals are computed using

$$\int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \psi^g(r, \hat{\Omega}) \approx \sum_{m=1}^M \omega_m \Gamma_{-m}^{*g}(r) \psi_m^g(r), \quad (26)$$

where

M = number of discrete ordinates

ω_m = weight of ordinate m

subscript $-m$ = ordinate that is in the opposite direction of ordinate m .

If discrete-ordinates angular fluxes are not available, then angle integrals are estimated using a moments expansion:¹⁰

$$\int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \psi^g(r, \hat{\Omega}) \approx \sum_{l=0}^L (2l + 1) \Gamma_l^{*g}(r) \phi_l^g(r). \quad (27)$$

Note that the second term in the fission cross section and P_0 scattering equations appears because these cross sections are addends in the total cross section.

The numerator of Eq. (22) is the same as Eq. (13) except that the latter has the generalized adjoint $\Gamma^{*g}(r, \hat{\Omega})$ where the former has the λ -mode adjoint $\psi^{*g}(r, \hat{\Omega})$. Thus, the sensitivities of k_{eff} are computed using the equations in Table I, with the following modifications: (1) k is substituted for R in the subscript on S , (2) λ -mode $\phi^{*g}(r, \hat{\Omega})$ and $\phi_l^{*g}(r)$ [Eq. (25)], the solution to Eq. (12), are used instead of $\Gamma^{*g}(r, \hat{\Omega})$ and $\Gamma_l^{*g}(r)$, and (3) the integrals are divided by the denominator of Eq. (22).

SENSMG also computes the relative sensitivity S_{k, ρ_n}^g of k_{eff} with respect to the mass density of each material defined in the SENSMG input file (Sec. III.B). The adjoint-based formulation for S_{k, ρ_n}^g is

$$S_{k, \rho_n}^g = \frac{\int_{V_n} dV \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) (\lambda F_n - A_n) \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \lambda F \psi^g(r, \hat{\Omega})} . \quad (28)$$

The total sensitivity S_{k, ρ_n} of k_{eff} to the mass density of coarse mesh n is the sum of S_{k, ρ_n}^g over groups. It is important to note that the derivative in S_{k, ρ_n}^g is a constant-volume partial derivative.

SENSMG also computes the derivative of k_{eff} with respect to the location of each material interface location and the location of the outer boundary. The derivative, computed for each energy group, is denoted $\partial k_{eff}^g / \partial r_n$, where r_n is the location of the n 'th interface, surface S_n . The adjoint-based formulation for $\partial k_{eff}^g / \partial r_n$ is¹¹

$$\frac{\partial k_{eff}^g}{\partial r_n} = \frac{k_{eff} \int_{S_n} dS \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) (\lambda \Delta F_n - \Delta A_n) \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \lambda F \psi^g(r, \hat{\Omega})} , \quad (29)$$

where ΔF_n is the difference in the fission operator across surface S_n , defined as the operator on the negative side (left, inside, or bottom) minus the operator on the positive side (right, outside, or top), and ΔA_n is defined analogously for the transport operator. The total derivative of k_{eff} to the location of surface S_n , $\partial k_{eff} / \partial r_n$, is the sum of $\partial k_{eff}^g / \partial r_n$ over groups. It is important to note that $\partial k_{eff}^g / \partial r_n$ is a constant-density partial derivative.

For all of the problems in this computer code abstract, the constant-volume and constant-density partial derivatives can be combined (with cell volumes and interface areas) to compute constant-mass partial derivatives.¹²

II.C. α -Eigenvalue Problems

GPT may also be used for α -eigenvalue calculations. The α -mode forward and adjoint neutron fluxes $\psi^g(r, \hat{\Omega})$ and $\psi^{*g}(r, \hat{\Omega})$ satisfy

$$\left(\frac{\alpha}{v} + A - F \right) \psi = 0 \quad (30)$$

and

$$\left(\frac{\alpha}{v} + A^* - F^* \right) \psi^* = 0 , \quad (31)$$

respectively, where v is the neutron speed. Vacuum boundary conditions apply to Eqs. (30) and (31). The indirect effect is

$$\left(S_{R,x}^g \right)_{indirect} = \sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \left(\frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega}) , \quad (32)$$

where now $\Gamma^{*g}(r, \hat{\Omega})$ satisfies

$$\left(\frac{\alpha}{v} + A^* - F^* \right) \Gamma^* = S^* , \quad (33)$$

with S^* given by Eq. (15). A vacuum boundary condition applies to Eq. (33). The method of successive approximations can be used to solve Eq. (33) using Eq. (16) with

$$\Gamma_n^{*g}(r, \hat{\Omega}) = \xi_n^{*g}(r, \hat{\Omega}) - \frac{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \xi_n^{*g}(r, \hat{\Omega}) \frac{1}{v^g} \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \frac{1}{v^g} \psi^g(r, \hat{\Omega})} \psi^{*g}(r, \hat{\Omega}), \quad (34)$$

where v^g is the neutron speed associated with energy group g . The $\xi_n^{*g}(r, \hat{\Omega})$ are generated recursively using

$$\begin{cases} \left(\frac{\alpha}{v} + A^*\right) \xi_0^* = S^*, \\ \left(\frac{\alpha}{v} + A^*\right) \xi_n^* = F^* \Gamma_{n-1}^*, \quad n > 0. \end{cases} \quad (35)$$

The second term on the right side of Eq. (34) is included to remove any fundamental mode contamination.

It can be shown that the generalized adjoint for α -eigenvalue problems must satisfy

$$\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \frac{1}{v^g} \psi^g(r, \hat{\Omega}) = 0. \quad (36)$$

Like k_{eff} calculations, in α calculations, the α -mode forward and adjoint angular fluxes are computed to apply GPT for reaction-rate ratios, so α sensitivities can always be computed with α -mode reaction-rate ratio sensitivities. The relative sensitivity $S_{\alpha,x}^g$ of the α eigenvalue to a transport cross section σ_x^g for reaction x in group g is defined to be

$$S_{\alpha,x}^g \equiv \frac{\sigma_{x,0}^g}{\alpha_0} \frac{\partial \alpha}{\partial \sigma_x^g} \Big|_{\sigma_x^g = \sigma_{x,0}^g}. \quad (37)$$

The adjoint-based formulation for $S_{\alpha,x}^g$ is

$$S_{\alpha,x}^g = \frac{1}{\alpha} \frac{\sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \left(\frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \frac{1}{v^g} \psi^g(r, \hat{\Omega})}. \quad (38)$$

Again, the total sensitivity $S_{\alpha,x}$ of α to reaction x is the sum of $S_{\alpha,x}^g$ over groups.

The equations in Table I are also used for reaction-rate ratios of the α -mode flux, except that k_{eff} is set to 1 (and the appropriate fluxes are used).

The numerator of Eq. (38) is the same as Eq. (13) except that the latter has the generalized adjoint $\Gamma^{*g}(r, \hat{\Omega})$ where the former has the α -mode adjoint $\psi^{*g}(r, \hat{\Omega})$. Thus, the sensitivities of α are computed using the equations in Table I, with the following modifications: (1) λ is set to 1, (2) α is substituted for R in the subscript on S , (3) α -mode $\phi^{*g}(r, \hat{\Omega})$ and $\phi_l^{*g}(r)$ [Eq. (25)], the solution to Eq. (31), are used instead of $\Gamma^{*g}(r, \hat{\Omega})$ and $\Gamma_l^{*g}(r)$, and (4) the integrals are divided by the denominator of Eq. (38) (including the factor of α).

SENSMG also computes the relative sensitivity S_{α,ρ_n}^g of α with respect to the mass density of each material defined in the SENSMG input file (Sec. III.B). The adjoint-based formulation for S_{α,ρ_n}^g is

$$S_{\alpha, \rho_n}^g = \frac{1}{\alpha} \frac{\int_V dV \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) (F_n - A_n) \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \frac{1}{v^g} \psi^g(r, \hat{\Omega})} . \quad (39)$$

The total sensitivity S_{α, ρ_n} of α to the mass density of coarse mesh n is the sum of S_{α, ρ_n}^g over groups. It is important to note that the derivative in S_{α, ρ_n}^g is a constant-volume partial derivative.

SENSMG also computes the derivative of α with respect to the location of each material interface location and the location of the outer boundary. The derivative, computed for each energy group, is denoted $\partial\alpha^g/\partial r_n$. The adjoint-based formulation for $\partial\alpha^g/\partial r_n$, whose derivation is similar to that of $\partial\lambda/\partial r_n$ in Ref. 11, is

$$\frac{\partial\alpha^g}{\partial r_n} = \frac{\int_{S_n} dS \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) (\Delta F_n - \Delta A_n) \psi^g(r, \hat{\Omega})}{\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \frac{1}{v^g} \psi^g(r, \hat{\Omega})} , \quad (40)$$

where again the Δ terms are differences across surface S_n . The total derivative of α to the location of surface S_n , $\partial\alpha/\partial r_n$, is the sum of $\partial\alpha^g/\partial r_n$ over groups. It is important to note that $\partial\alpha^g/\partial r_n$ is a constant-density partial derivative.

II.D. Fixed-Source Problems

For fixed-source calculations, the forward and adjoint neutron fluxes $\psi^g(r, \hat{\Omega})$ and $\psi^{*g}(r, \hat{\Omega})$ satisfy

$$(A - F)\psi = Q \quad (41)$$

and

$$S_{L,x}^g = \frac{1}{L} \sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \left\{ \psi^{*g}(r, \hat{\Omega}) \frac{\partial Q}{\partial \sigma_x^g} + \psi^{*g}(r, \hat{\Omega}) \left(\frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega}) \right\} . \quad (45)$$

The intrinsic neutron source rate density Q depends on isotopic (α, n) cross sections. However, SENSMG presently considers only neutron reaction cross sections. Therefore, the derivative $\partial Q/\partial \sigma_x^g$ is zero, and SENSMG uses

$$S_{L,x}^g = \frac{1}{L} \sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) \left(\frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega}) . \quad (46)$$

Again, the total sensitivity $S_{L,x}$ of L to reaction x is the sum of $S_{L,x}^g$ over groups.

$$(A^* - F^*)\psi^* = Q^* , \quad (42)$$

respectively. In SENSMG, the neutron source $Q^g(r)$ is an energy- and material-dependent intrinsic source of spontaneous fission neutrons and neutrons from (α, n) reactions occurring in a homogeneous material. It is assumed to be isotropic. The adjoint source Q^* is defined by the quantity of interest. In SENSMG, the quantity of interest for the adjoint flux $\psi^{*g}(r, \hat{\Omega})$ is, for spheres and cylinders, the total neutron leakage L from the system, which is

$$L \equiv \int_{S_b} dS \sum_{g=1}^G \int_{\hat{\Omega} \cdot \hat{n} > 0} d\hat{\Omega} \hat{\Omega} \cdot \hat{n} \psi^g(r, \hat{\Omega}) , \quad (43)$$

where S_b represents the outer boundary and \hat{n} is the outward unit normal vector at each point on S_b . For slabs, the quantity of interest is the leakage from the right boundary, computed using Eq. (43) with S_b representing just the right boundary. Therefore, the adjoint source Q^* is an isotropic incoming flux distribution on the outer boundary (for spheres/cylinders) or the right boundary (for slabs). (An isotropic incoming distribution in the flux is equivalent, within the limits of the discrete-ordinates approximation, to a cosine incoming distribution in the neutron density.) Other than that incoming flux condition for Eq. (42), vacuum boundary conditions apply to Eqs. (41) and (42).

The relative sensitivity $S_{L,x}^g$ of the leakage to a transport cross section σ_x^g for reaction x in group g is defined to be

$$S_{L,x}^g \equiv \frac{\sigma_{x,0}^g}{L_0} \frac{\partial L}{\partial \sigma_x^g} \Big|_{\sigma_x^g = \sigma_{x,0}^g} . \quad (44)$$

The adjoint-based formulation for $S_{L,x}^g$ is

For reaction-rate ratios, the indirect effect of Eq. (10) is

$$\left(S_{R,x}^g\right)_{indirect} = \sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \left\{ \Gamma^{*g}(r, \hat{\Omega}) \frac{\partial Q}{\partial \sigma_x^g} + \Gamma^{*g}(r, \hat{\Omega}) \left(\frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega}) \right\}. \quad (47)$$

Again, SENSMSG considers only neutron reaction cross sections, so $\partial Q/\partial \sigma_x^g = 0$ and the indirect effect is

$$\left(S_{R,x}^g\right)_{indirect} = \sigma_x^g \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \left(\frac{\delta F}{\delta \sigma_x^g} - \frac{\delta A}{\delta \sigma_x^g} \right) \psi^g(r, \hat{\Omega}). \quad (48)$$

In Eqs. (47) and (48), $\Gamma^{*g}(r, \hat{\Omega})$ satisfies

$$(A^* - F^*)\Gamma^* = S^*, \quad (49)$$

with S^* given by Eq. (15). A vacuum boundary condition applies to Eq. (49). For fixed-source problems, the operator on $\Gamma^{*g}(r, \hat{\Omega})$ in Eq. (49) is not singular, so Eq. (49) is solved directly, without the need for successive approximations.

For fixed-source problems, Eq. (49) is the same form as Eq. (42) (the sources are different). For reaction-rate ratios, we are following the convention of Stacey³ in referring to the adjoint function as a generalized adjoint and using the symbol Γ^* . We use ψ^* for the neutron leakage adjoint so that the discussion in this section parallels somewhat the discussion of eigenvalue problems in Secs. II.B and II.C. However, for reaction-rate ratios in fixed-source problems, only one adjoint is needed, Γ^* . Therefore, in this case, the neutron leakage sensitivity does require an extra adjoint transport calculation beyond what the reaction-rate ratios sensitivities require, unlike in the eigenvalue case.

For fixed-source problems, the magnitude of the neutron flux is fixed, and an absolute reaction rate R_1 of Eq. (1) can be computed. In this case, S^* is

$$S^{*g}(r, \hat{\Omega}) = \frac{1}{R_1} \frac{\partial R_1}{\partial \phi} = \frac{\Sigma_1^g(r)}{R_1}. \quad (50)$$

For a reaction-rate ratio in fixed-source problems, the generalized adjoint must satisfy

$$\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) Q^g(r) = 0. \quad (51)$$

For an absolute reaction rate in fixed-source problems, the generalized adjoint must satisfy

$$\int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) Q^g(r) = 1. \quad (52)$$

With one exception, the equations in Table I are also used for reaction-rate ratios of a fixed-source flux, except that k_{eff} is set to 1 (and the appropriate fluxes are used). The exception is that the sensitivity of the reaction-rate ratio to the density of nuclide i , N_i , in a fixed-source problem requires the derivative $\partial Q^g/\partial N_i$ (but $\partial Q/\partial \sigma_x^g$ is still zero). The contribution of the source sensitivity to the sensitivity of R to the density of isotope i is

$$S_{R,Q_{N_i}}^g = N_i \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \frac{\partial Q^g}{\partial N_i}. \quad (53)$$

The sensitivity $S_{R,Q_{N_i}}^g$ of Eq. (53) is included as another addend on the right side of the equation for S_{R,N_i}^g in Table I. The source Q^g is the sum of the spontaneous fission source in group g and the (α,n) source in group g :

$$Q^g = Q_{s.f.}^g + Q_{(\alpha,n)}^g. \quad (54)$$

For spontaneous fission sources, the source rate density in group g for isotope i is

$$Q_{s.f.,i}^g = \lambda_i N_i \chi_{s.f.,i}^g, \quad (55)$$

where λ_i is the decay constant for isotope i and $\chi_{s.f.,i}^g$ includes a spectrum function and the spontaneous fission branching ratio,⁶ neither of which depend on any isotopic densities. The total spontaneous fission source rate density in group g is the sum over fissioning isotopes:

$$Q_{s.f.}^g = \sum_{j=1}^J Q_{s.f.,j}^g = \sum_{j=1}^J \lambda_j N_j \chi_{s.f.,j}^g. \quad (56)$$

The derivative of the spontaneous fission neutron source rate density in group g with respect to the density of spontaneous-fissioning isotope i is

$$\frac{\partial Q_{s.f.}^g}{\partial N_i} = \frac{\partial Q_{s.f.,i}^g}{\partial N_i} = \frac{Q_{s.f.,i}^g}{N_i} . \quad (57)$$

Whether MISC or SOURCES4C is used to compute spontaneous fission sources, the overall spontaneous-fission neutron spectrum is applied for each spontaneous-fissioning isotope i .

For (α,n) sources, the source rate density in group g for α -emitting isotope j due to target isotope k is

$$Q_{(\alpha,n),j,k}^g = \lambda_j N_j \chi_{(\alpha,n),j,k}^g , \quad (58)$$

where $\chi_{(\alpha,n),j,k}^g$ depends on the density of the target isotope but also on the density of every isotope in the material because they all contribute to the α -stopping power of the material. The total (α,n) source rate density in group g is the sum over α -emitting isotopes and target isotopes:

$$\begin{aligned} Q_{(\alpha,n)}^g &= \sum_{j=1}^J \sum_{k=1}^K Q_{(\alpha,n),j,k}^g \\ &= \sum_{j=1}^J \lambda_j N_j \sum_{k=1}^K \chi_{(\alpha,n),j,k}^g . \end{aligned} \quad (59)$$

The derivative of the (α,n) neutron source rate density in group g with respect to the density of isotope i is

$$\begin{aligned} \frac{\partial Q_{(\alpha,n)}^g}{\partial N_i} &= \delta_{ij} \lambda_j \sum_{k=1}^K \chi_{(\alpha,n),j,k}^g \\ &\quad + \sum_{j=1}^J \lambda_j N_j \sum_{k=1}^K \frac{\partial \chi_{(\alpha,n),j,k}^g}{\partial N_i} . \end{aligned} \quad (60)$$

The derivative of $\chi_{(\alpha,n),j,k}^g$ is too complicated to present here, but it has been worked out¹³ and implemented in SENSMG.

Using Eqs. (54) and (57) and the left side of Eq. (60), Eq. (53) becomes

$$S_{R,Q_{N_i}}^g = \int_V dV \int_{4\pi} d\hat{\Omega} \Gamma^{*g}(r, \hat{\Omega}) \left(Q_{s.f.,i}^g + N_i \frac{\partial Q_{(\alpha,n)}^g}{\partial N_i} \right) . \quad (61)$$

The neutron sources are isotropic, so only the zeroth moment of $\Gamma^{*g}(r, \hat{\Omega})$ is needed.

SENSMG makes an approximation in evaluating the Q^g terms in Eq. (61). The total spontaneous fission source rate density for each isotope $Q_{s.f.,i}^g$ is

known, but the spontaneous fission spectrum calculated for the material (the mixture of isotopes) is applied to approximate $Q_{s.f.,i}^g$:

$$Q_{s.f.,i}^g \approx Q_{s.f.,i} \frac{\sum_{j=1}^J Q_{s.f.,j}^g}{\sum_{g=1}^G \sum_{j=1}^J Q_{s.f.,j}^g} , \quad (62)$$

where j indexes fissioning isotopes. The derivative of the total (α,n) source rate density (for the material) with respect to each isotope is known,¹³ but the (α,n) neutron spectrum is computed and applied for each α -emitting isotope and each target isotope:

$$\frac{\partial Q_{(\alpha,n)}^g}{\partial N_i} \approx \frac{\partial Q_{(\alpha,n)}}{\partial N_i} \frac{\sum_{j=1}^J Q_{(\alpha,n),i,j}^g}{\sum_{g=1}^G \sum_{j=1}^J Q_{(\alpha,n),i,j}^g} , \quad (63)$$

where j indexes α emitters if i is a target isotope and target isotopes if i is an α emitter. If i is neither an α emitter nor a target, the overall (α,n) neutron spectrum is applied:

$$\frac{\partial Q_{(\alpha,n)}^g}{\partial N_i} \approx \frac{\partial Q_{(\alpha,n)}}{\partial N_i} \frac{\sum_{j=1}^J \sum_{k=1}^K Q_{(\alpha,n),k,j}^g}{\sum_{g=1}^G \sum_{j=1}^J \sum_{k=1}^K Q_{(\alpha,n),k,j}^g} . \quad (64)$$

In some cases, a weak α emitter is more important due to its contribution to the material's α -particle stopping power than to its contribution to the α -particle source rate density.¹³ Then Eq. (64) is used rather than Eq. (63).

The numerator of Eq. (45) is the same as Eq. (13) except that the latter has the generalized adjoint $\Gamma^{*g}(r, \hat{\Omega})$ where the former has the fixed-source adjoint $\psi^{*g}(r, \hat{\Omega})$. Thus, the sensitivities of the leakage are computed using the equations in Table I, with the following modifications: (1) λ is set to 1, (2) L is substituted for R in the subscript on S , (3) the fixed-source $\phi^{*g}(r, \hat{\Omega})$ and $\phi_l^{*g}(r)$ [Eq. (25)], the solution to Eq. (42), are used instead of $\Gamma^{*g}(r, \hat{\Omega})$ and $\Gamma_l^{*g}(r)$, (4) $S_{R,Q_{N_i}}^g$ of Eq. (61) is included (see 3), and (5) the integrals are divided by L of Eq. (43).

SENSMG also computes the relative sensitivity S_{L,ρ_n}^g of L with respect to the mass density of each material defined in the SENSMG input file (Sec. III.B). The adjoint-based formulation for S_{L,ρ_n}^g is

$$S_{L,\rho_n}^g = \frac{1}{L} \int_{V_n} dV \int_{4\pi} d\hat{\Omega} \left\{ \psi^{*g}(r, \hat{\Omega}) Q_n + \psi^{*g}(r, \hat{\Omega}) (F_n - A_n) \psi^g(r, \hat{\Omega}) \right\}. \quad (65)$$

The total sensitivity S_{L,ρ_n} of L to the mass density of coarse mesh n is the sum of S_{L,ρ_n}^g over groups. The derivative of $\chi_{(a,n),j}^g$ with respect to the material mass density is zero.¹³ It is important to note that the derivative in S_{L,ρ_n}^g is a constant-volume partial derivative.

SENSMG also computes the derivative of L with respect to the location of each material interface location and the location of the outer boundary. The derivative, computed for each energy group, is denoted $\partial L^g / \partial r_n$. The adjoint-based formulation for $\partial L^g / \partial r_n$ is¹⁴

$$\frac{\partial L^g}{\partial r_n} = \int_{S_n} dS \int_{4\pi} d\hat{\Omega} \left\{ \psi^{*g}(r, \hat{\Omega}) \Delta Q_n + \psi^{*g}(r, \hat{\Omega}) (\Delta F_n - \Delta A_n) \psi^g(r, \hat{\Omega}) \right\}, \quad (66)$$

where again the Δ terms are differences across surface S_n . The total derivative of L to the location of surface S_n , $\partial L / \partial r_n$, is the sum of $\partial L^g / \partial r_n$ over groups. It is important to note that $\partial L^g / \partial r_n$ is a constant-density partial derivative.

III. USER INTERFACE

The SENSMSG sensitivity tool is run by invoking the name of a Python control script with options. On the LANL high-performance computing (HPC) machines, the control script is called `/usr/projects/data/nuclear/working/sensitivities/bin/sensmg.py`.

There are four ways to input information into SENSMSG: (1) through preset environment variables, (2) through an input file, (3) through the command line, and (4) in special cases, through the Python script itself. These are discussed in Secs. III.A through III.D.

III.A. Environment Variables

SENSMSG checks the environment variable `SENS_PARTISN` for the specific executable version of PARTISN to run. If that variable is not set, SENSMSG will use (in C-shell notation) `/usr/projects/lin-det/rel8_27/8_27_15/${HN}-intel-17.0.4-openmpi-2.1.2/partisn`, where `${HN}` is derived from the `HOSTNAME` describing the computer that the user is on. See the PARTISN documentation for available versions and systems. SENSMSG loads the appropriate modules for the supported PARTISN versions.

When using the Nuclear Data Interface (NDI) at LANL, the user should have an `NDI_GENDIR_PATH` set. If there is none, SENSMSG will use `/usr/projects/data/nuclear/ndi/2.1.3/share/gendir.all`.

III.B. Input File

The default input file name is `sensmg_inp`, but this can be changed with the `-i` parameter on the command line (Sec. III.C). The input file tells what type of calculation to run, fully specifies the geometry and materials, and lists the reaction-rate ratios of interest.

A forward slash (/) designates that the rest of the line is a comment. Forward slashes are required to end material composition lines and reaction-rate ratio lines. Each line of the input file is free format, but the lines must be in a certain order. In some cases, the entries on a line must also be in a certain order. This section describes the order.

SENSMSG computes one-dimensional (r) spheres or slabs or two-dimensional (r - z) cylinders. The input file for a spherical geometry is described in Table II. The first card is a title line. The second card has `sph` or `sphere` for a sphere and it also tells the type of calculation to run, either `keff` (`keff`), `α` (`alpha`), or neutron leakage (`lkg`). The order of these words is not important.

The third card tells the LIBNAME used to specify cross sections to PARTISN.

The fourth and fifth cards describe materials. The number of materials (NM) is given, and then the composition for each material is specified (each material on a single line) by ZAIID (the nuclide identification number) and weight fraction. Numeric ZAIIDs, not the full cross-section-specification ZAIIDs, are used to identify nuclides. After the material specifications, the material densities are listed in order on a single line. The same material can appear at more than one density but only if it is listed each time (once for each density) in the material list.

It is at this point that sphere/slab and cylinder input files differ. For a spherical or slab input, the sixth card tells the number of radial shells or slabs (NR) and the seventh card specifies the location (not the thickness) of

TABLE II
Spherical or Slab Input File Description

Card	Description	Number of Entries per Line
1	Title	Maximum 120 characters
2	[sphere sph slab] [keff alpha lkg]	2 (words)
3	LIBNAME	1
4.0	NM	1
4.1 to 4.NM	Specification for each material. Each line has an index followed by a list of pairs of ZAID and weight fraction. End each line with “/”.	Variable
5	Mass density of each material (g/cm ³)	NM
6	NR	1
7	Radius of each shell (cm) ^a	NR or NR + 1 ^a
8	Material assignments to radial shells (0 means void)	NR
9	Index of radial shell to use for reaction rates	1
10.0	NRRR ^b	1
10.1 to 10.NRRR	Reaction-rate ratios; end each line with “/”.	2 or 4

^aFor spheres, the inner-most surface is assumed to be 0. For slabs, the left-most surface is entered.

^bNRRR = Number of reaction-rate ratios.

each shell or slab, i.e., the outer radius or surface location. The eighth card assigns materials to spherical shells or slab layers in order.

The rest of the spherical input file tells where the reaction-rate ratios are to be calculated and what they are. The ninth card tells the index of the radial shell in which the ratios are calculated. All reaction rates and fluxes are computed in the same location. The tenth card tells the number of reaction-rate ratios, and the next set of lines gives the reaction-rate ratios in ZAID-reaction pairs; see later in this section for details. If the number of reaction-rate ratios is zero, SENSMG will only compute k_{eff} , α , or leakage sensitivities.

The input file for a cylindrical geometry is described in Table III. The first five cards are the same as described above and in Table II for a sphere/slab geometry. The sixth card tells not just NR but also the number of axial layers (NZ). The outer radius of each shell is given on the seventh card. The eighth card gives the location of each axial layer, starting at the bottom. The number of axial surfaces to specify is thus one plus the number of layers.

Next, materials are assigned to two-dimensional regions. The layout here is somewhat graphical in that each line has an

entry for each radial shell (in order), and there is a line for each axial layer in order from top to bottom.

The tenth card needs to specify both the radial shell and the axial layer in which reaction-rate ratios are to be computed. The rest of the input file is the same as in the sphere/slab case.

Cross sections are specified using the LIBNAME given in the input file. As when PARTISN is run on its own, the LIBNAME library must exist in the `gendir` file specified in the environment variable `NDI_GENDIR_PATH` (Sec. III.A). SENSMG searches for LIBNAME in the `gendir` file to set the cross sections. For each isotope in the problem, including edit isotopes (those only used in reaction rates), SENSMG sets the full ZAID (and therefore the cross sections) using the first instance of the isotope found with the specified LIBNAME in `NDI_GENDIR_PATH`. As when PARTISN is run on its own, the isotopes in the input file must match those available in the specified LIBNAME and `NDI_GENDIR_PATH`.

In addition to the libraries available through the NDI at LANL, SENSMG can use the Kynea3 cross sections developed by Varley and Mattingly.¹⁵ The Kynea3 cross

TABLE III
Cylindrical Input File Description

Card	Description	Number of Entries per Line
1	Title	Maximum 120 characters
2	[cyl cylinder] [keff alpha lkg]	2 (words)
3	LIBNAME	1
4.0	NM	1
4.1 to 4.NM	Specification for each material. Each line has an index followed by a list of pairs of ZAID and weight fraction. End each line with “/”.	Variable
5	Mass density of each material (g/cm ³)	NM
6	NR and NZ	2
7	Radius of each shell (cm)	NR
8	Location of each axial layer starting with the bottom of the first layer (cm)	NZ + 1
9.1 to 9.NZ	Material assignments to radial zones for each axial layer (0 means void); axial layers are listed top to bottom.	NR
10	Index of radial zone and index of axial zone to use for reaction rates	2
11.0	NRRR ^a	1
11.1 to 11.NRRR	Reaction-rate ratios; end each line with “/”.	2 or 4

^aNRRR = Number of reaction-rate ratios.

sections are in 79 groups and have upscattering in the lower 35 groups. Use `kynea3` for LIBNAME in the input file. The code will automatically set the number of energy groups to 79. Because the Kynea3 cross sections do not have a full set of edit reactions, this capability is generally applicable only to k_{eff} , α , and leakage sensitivities. (The Kynea3 set does include the first four edit positions of Table IV, discussed below.) The isotopes available in the Kynea3 set are listed with their ZAIDs in Appendix B.

SENSMG can support users who do not have access to the NDI or Kynea3. If the word `special` is used for LIBNAME in the input file, SENSMG will look in directory `SENS_DATA` (see Sec. III.D) for four files from a forward run (`for_macrxs`, `for_snxedt`, `for_ndxsrf`, and `for_znatdn`), four files from a cross-section run (`xs1_macrxs`, `xs1_snxedt`, `xs1_ndxsrf`, and `xs1_znatdn`), and a file called `spc_gendir`, copied from the `sensmg.log` file (see Sec. IV; below “diagnostics from `rdgendir`” or “diagnostics from `rdbxslib`”), that gives the

ZAIDs and atomic weights used in the problem. These nine files must correspond exactly to the end user’s input file. The eight cross-section files must be created with the end user’s PARTISN version. SENSMG is presently hard-wired to accept MENDF71X cross sections using this capability.

Using $\Sigma_1 = N_1 \sigma_{x1}$ and $\Sigma_2 = N_2 \sigma_{x2}$ in Eqs. (1), (2), and (4), reaction-rate ratios are specified as

$$\text{ZAID1 } x1 \text{ ZAID2 } x2 /$$

Note that the forward slash at the end of the line is required. For k_{eff} and α problems, if ZAID2 and $x2$ are not present, it means to use 1 for Σ_2 , as discussed in Sec. II.B. Numeric ZAIDS are used to specify nuclides (i.e., the full cross-section specification is not used here). Reactions $x1$ and $x2$ are given as integers indicating the position of the desired cross section in the PARTISN edit tables. Available edit positions when the NDI is used are given in Table IV. Note from Table IV that the location of certain edit reactions changed with PARTISN version 8.

TABLE IV
Edit Reactions Available from the NDI

Position	PARTISN 7 and Earlier		PARTISN 8 and Later	
	Description	Name	Description	Name
1	fission spectrum	chi	fission spectrum	chi
2	effective nusigf	nusigf	effective nusigf	nusigf
3	total	total	total	total
4	absorption	abs	absorption	abs
5	chi	chi	chi	chi
6	elastic	mend1	elastic	(n,n)
7	inelastic	mend2	inelastic	(n,n')
8	(n,2n)	mend3	(n,2n)	(n,2n)
9	(n,3n)	mend4	(n,3n)	(n,3n)
10	(n,gamma)	mend5	(n,gamma)	(n,g)
11	(n,alpha)	mend6	(n,proton)	(n,p)
12	(n,proton)	mend7	(n,alpha)	(n,a)
13	(n,deuteron)	mend8	1st-chance fission	(n,f)
14	(n,triton)	mend9	2nd-chance fission	(n,n')f
15	(n,3He)	mend10	3rd-chance fission	(n,2n)f
16	sum of all fission	n-fiss	sum of all fission	(n,F)
17	prompt fission spectrum	mend12	prompt fission spectrum	chi_pr
18	total fission spectrum	mend13	total fission spectrum	chi_tot
19	—	—	(n,deuteron)	(n,d)
20	—	—	(n,triton)	(n,t)

An example of a spherical input file follows (note that though the composition of material 1 wraps here, it should be on a single line):

```
Pu-Flattop (PU-MET-FAST-006)
keff sph
mt71x / libname
2 / no of materials
1 94239 9.38001E-01 94240 4.79988E-02 94241 2.99996E-03 31069 6.53652E-03 31071
4.46355E-03 / Pu-alloy
2 92234 5.40778E-05 92235 7.10966E-03 92238 9.92836E-01 / Flattop natural U
15.53 19.00 / densities
3 / no of shells
0.155 4.5332 24.142 / outer radii
1 1 2 / material nos
1 / index of coarse mesh to use for reaction rates
3 / number of reaction-rate ratios
92238 16 92235 16 /numerator and denominator
93237 16 92235 16 /numerator and denominator
92235 10 /numerator and flux
```

An example of a cylindrical input file follows:

```
HEU-MET-FAST-001-02 Godiva-II
keff cyl
mt71x / libname
1 / no of materials
```



```

1 92234 1.01999E-02 92235 9.37100E-01 92238, 5.27004E-02 / Godiva
18.74 / densities
2 3 / no of r layers, no of z layers
0.635 7.62 / outer radii
0. 7.86 9.13 16.99 / heights
1 1 / material nos, top
1 1 / material nos
1 1 / material nos, bottom
1 2 / r and z indices of coarse mesh to use for reaction rates
1 /number of reaction-rate ratios
92238 16 92235 16 /numerator and denominator

```

More examples are given in [Appendix A](#) and discussed in [Sec. VI](#).

III.C. Command Line

The available command line options are listed and described in [Table V](#).

If PARTISN version 7 or earlier is used, then spherical and slab problems should be run using a single processor (`-np 1`). The reason is that [Eq. \(26\)](#) is more accurate than [Eq. \(27\)](#), and PARTISN’s angular fluxes are readily available for one-dimensional problems, except that message passing interface (MPI) runs of PARTISN 7 (and earlier) cannot print the angular flux file. If PARTISN version 8 is used, then spherical and slab problems can be run with as many processors as are available.

Because of memory limitations, cylindrical problems approximate the volume inner product of angular fluxes with a moment expansion [[Eq. \(27\)](#)]. Therefore, cylindrical problems can always be run with as many processors as are available. (This approximation is too crude for surface integrals, so the `-rplane` and `-zplane` options discussed below were developed.)

The number of energy groups “`-ngroup [G]`” available when using the NDI at LANL is $G = 618, 301, 250, 141, 133, 130, 91, 75, 70, 49, 47, 30, 21, 16, 14, 13, 12, 10, 9, 8, 7, 6, 5, 4, 3, 2$, or 1. Not every value is available for all possible values of LIBNAME ([Sec. III.B](#)). (The 130-group library¹⁶ is particularly limited.) The 79-group Kynea3 set¹⁵ is also available ([Sec. III.B](#)).

For spheres and slabs, the traditional S_N constants are used (`iquad = 1` in PARTISN lingo¹). For cylinders, the square Chebychev-Legendre built-in set is used (`iquad = 6`). For both geometries, any even S_N quadrature order can be used¹; set with the `-isn` parameter.

The `-epsig` parameter is the convergence criterion for the sum in [Eq. \(16\)](#). Before the latest $\Gamma_n^{*g}(r, \hat{\Omega})$ is added to the current sum up to $n - 1$, each element of $\Gamma_n^{*g}(r, \hat{\Omega})$ (radial fine mesh, axial fine mesh, energy group, and either

moment or discrete ordinate) is divided by the corresponding element in the current sum. If the absolute value of the largest ratio is smaller than `epsig`, the sum is considered to be converged. The latest $\Gamma_n^{*g}(r, \hat{\Omega})$ is then added to the current sum (whether the sum is converged or not).

The `-chinorm` parameter controls constrained normalization for chi. Constrained normalization is described in [Ref. 17](#) for material compositions, but SENSIMG uses the same equations for the normalization of chi. The `-chinorm` parameter is similar to the `CONSTRAIN` keyword on MCNP6’s `KSEN` card.¹⁸ In MCNP6, “`CONSTRAIN = yes`” gives full normalization¹⁹ and is the default, as in SENSIMG. “`CONSTRAIN = no`” is the same as “`-chinorm none`”. MCNP6 does not have a partial normalization option.

For fixed-source neutron leakage problems (keyword `lkg` in the input file; [Section III.B](#)), SENSIMG uses the `MISC` code⁵ to compute spontaneous fission sources and the `SOURCES4C` code⁶ to compute (α, n) sources. “`-misc no`” can be used to have SENSIMG use `SOURCES4C` rather than `MISC` for spontaneous fission sources. “`-alpha_n no`” can be used to ignore (α, n) sources altogether.

The `-srcacc_no` parameter is useful if the PARTISN iterations for the forward or adjoint files are observed to “flip flop” between two states or otherwise have trouble converging. If that happens, turn off source acceleration for the difficult calculation. Source acceleration is always turned off for the generalized adjoint functions.

The `-use_existing` option is a quick way to see the effect of a change in the chi normalization. However, it was mainly implemented for development and debugging of the SENSIMG code.

The `-rplane` and `-zplane` options activate PARTISN’s `ASLEFT/ASRITE` and `ASBOTT/ASTOP`

TABLE V
Command Line Options

Flag	Description	Default
-i	Input file name	sensmg_inp
-np	Number of MPI processes to use for PARTISN	1
-ngroup	Number of energy groups	30
-isn	S_N order	32
-isct	Legendre scattering expansion order	3
-epsi	Convergence criterion for PARTISN calculations (PARTISN's <code>epsi</code>) (Ref. 1)	1E-06
-epsig	Convergence criterion for GPT adjoint calculations in the method of successive approximations	1E-05
-chinorm	full/partial/none constrained normalization for the chi matrix	full
-fissdata	0/1/2 use fission matrix/chi matrix/chi vector (PARTISN's <code>fissdata</code>) (Ref. 1)	0
-aflxfrm	0/1 no/yes use an α -eigenvalue search method that may help for subcritical problems [PARTISN's <code>aflxfrm</code> (Ref. 1); ignored for versions before 7.72]	0
-alpha_n	yes/no use/ignore (α ,n) neutron sources	yes
-misc	yes/no use MISC/use SOURCES4C for spontaneous fission sources	yes
-nag	Number of α -particle energy groups in SOURCES4C	100
-srcacc_no	none/for/adj/for+adj turn off source acceleration for neither/forward/adjoint/both	none
-use_existing	yes/no use/don't use PARTISN output files that have already been calculated.	no
-my_modules	yes/no use/don't use user's own loaded module files. "yes" skips automatic module file loading, used to override defaults or if the user's system does not use modules.	no
-rplane	Radial surface on which to compute angular fluxes	-1 (i.e., do not use)
-zplane	Axial surface on which to compute angular fluxes	-1 (i.e., do not use)
-h	Show options (i.e., help)	None

keywords¹ to write surface angular flux files on the specified surfaces. The radial surfaces are indexed from 1 to NR, where NR is the number of radial regions, and the axial surfaces are indexed from 0 to NZ (bottom to

top), where NZ is the number of axial regions. Without this option, in cylindrical problems, the surface fluxes appearing in Eqs. (29), (40), and (66) are approximated using mesh-centered moments. This feature is intended

for PARTISN versions before 8.27. Those versions allow only one surface in each coordinate direction to have all angular fluxes written. The default value for both directions is -1 , which deactivates this feature.

Beginning with version 8.27, PARTISN is able to write surface angular fluxes for a user-input list of surfaces using the AFLUXX and AFLUXY keywords.¹ For cylindrical problems, SENSMG automatically generates that list.

III.D. Python Control Script

The full path specifying the executable code for SENSMG is set in the Python control script as a local variable. The paths to the SOURCES4C code⁶ and its data files, used in fixed-source problems, are also set in the script (as local variables). The path to the MISC code⁵ and its data files, also used in fixed-source problems, are also set in the script (as local variable for the code and environment variable ISCDATA for the data).

The path to group-structure files, Kynea3 cross sections, and special cross-section files (LIBNAME = special; see Sec. III.B) is set in the script as environment variable SENS_DATA.

IV. OUTPUT

The main output of SENSMG consists of four files. The files sens_rr_x and one of sens_k_x, sens_a_x, or sens_l_x are the sensitivities of the specified reaction rates and k_{eff} or α or neutron leakage, respectively, to all cross sections, as shown in Table I. In these files, sensitivities are given in units of %/%. The sensitivity to the total cross section (denoted “total” in the output tables) is the relative change in the ratio or the eigenvalue if only the total cross section for each isotope changes. This notion of the sensitivity to the total cross section is different from the way it is understood in MCNP6 (Ref. 18) and SCALE 6.2 (Ref. 20). There, the total cross section is the sum of all other cross sections and the sensitivity to the total cross section is the same as the sensitivity to the isotopic density.¹⁷ In PARTISN, it is possible for the total cross section to change independently of all other cross sections. SENSMG uses the label “density” to denote the sensitivity to the total cross section or density as it is understood in MCNP and SCALE (Table I). This explains why it is possible to have an energy-dependent sensitivity to the density in PARTISN (see the equation for the sensitivity to the density in Table I).

The file sens_rr_x includes a block of data for each reaction-rate ratio. Each ratio block includes the indirect effect as a block for each isotope in each material, the direct effect as a single block that includes the numerator and the denominator, and the total (indirect plus direct) as a block for each unique isotope in the problem. If an isotope appears in more than one material, the total will include all of its appearances and include indirect and direct effects, if any.

The file sens_k_x or sens_a_x or sens_l_x includes a single k_{eff} block or α block or leakage block, respectively, that includes the result of Eqs. (22), (38), or (45), respectively, as a block for each isotope in each material and the total as a block for each unique isotope in the problem. If each isotope only appears in one material, then the total isotope block will be the same as the isotope by material block.

The isotope blocks begin with an isotope summary, which is just the sum over groups of each sensitivity of Table I. The group-dependent sensitivities follow. The isotope blocks include the full matrix of sensitivities to chi, each element of which is, for example,

$$S_{R,\chi_i}^{g' \rightarrow g} = \frac{1}{k_{eff}} \int_V dV \Gamma_0^{*g}(r) \chi_i^{g' \rightarrow g} N_i v_i^{g'} \sigma_{f,i}^{g'} \phi_0^{g'}(r), \quad (67)$$

and the full matrix of sensitivities to scattering, each element of which is, for example,

$$S_{R,\sigma_{s,i}}^{g' \rightarrow g} = \int_V dV \Gamma_l^{*g}(r) N_i (2l + 1) \sigma_{s,i}^{g' \rightarrow g} \phi_l^{g'}(r) \quad (68)$$

for $l > 0$. [For α and leakage calculations, k_{eff} is set to 1 in Eq. (67).] The zeroth-moment scattering elements include the effect on the total cross section as shown in the entries on Table I. In the chi scattering matrices that are printed, columns are *from* groups [g' in Eqs. (67) and (68)] and rows are *to* groups (g).

The normalization of chi is indicated in the label for chi: chi_(nn) is used for no normalization, chi_(fn) for full normalization, and chi_(pn) for partial normalization. In full normalization, the sum over groups is analytically zero, but the output may have very small nonzero values due to numerical imprecision.

Using Eq. (9), it can be shown that the sum over groups of the direct effect should be one.

The third main output file is one of sens_k_r, sens_a_r, or sens_l_r. These are the sensitivities of k_{eff} or α or neutron leakage, respectively, to the mass densities, as given by Eqs. (28), (39), or (65),

respectively, and the derivatives with respect to the interface locations, as given by Eqs. (29), (40), or (66), respectively. The sensitivities have units of %/% and the derivatives have units of (units of quantity of interest)/centimeter. This output file has a block for the density sensitivities, followed by a block for the radial surface derivatives, followed by a block for the axial surface derivatives (for cylinders).

For cylinders, the derivative for each surface of each coarse mesh is output individually. The block for radial surface derivatives has an entry for each of the NR radial coarse mesh surfaces and each of the NZ axial zones (in the notation of Table III). The block for axial surface derivatives has an entry for each of the NR radial zones and each of the NZ + 1 axial surfaces.

If the material is the same on both sides of the coarse mesh, the derivatives will be zero, but they will still be output.

The fourth main output file is `sensmg.log`, which has all of the input parameters, any warnings that were generated, and information from SENSMG's reading of `NDI_GENDIR_PATH`. Diagnostic information about the forward and adjoint transport calculations and the iterations for the generalized adjoint functions is printed as the calculations are run. For example, the normalization of Eqs. (19), (36), (51), or (52) is checked at each iteration. It is difficult to monitor the progress of the iterations in the foreground window because the PARTISN screen output is constantly scrolling. The user can `cat` the contents of the log file in another window in order to monitor progress. A convergence monitor can be extracted with a command like "`grep convergence sensmg.log`".

For leakage calculations, this file also includes $\partial Q_{(\alpha,n)}/\partial N_i$, the derivative of the total (α,n) source rate density with respect to each isotope, labeled as "dQ/dN for (α,n) sources", separated into contributions from each nuclide as a target, stopping element, and α emitter.

SENSMG also keeps all input files, all text output files from transport and source calculations, and all binary cross-section and flux files.

V. SOME NOTES ON THE CODING

SENSMG is a combination of a Python script and a Fortran code. The Python script runs MISC and/or SOURCES4C (for fixed-source problems). It runs PARTISN and controls the iterations in the solution of Eqs. (14) and (33). The Fortran code writes MISC, SOURCES4C, and PARTISN input files and reads

MISC, SOURCES4C, and PARTISN output files, including cross-section files. The Fortran code computes and writes the final sensitivities. The Fortran code can probably be replaced with Python.

For fixed-source calculations, SENSMG first computes the intrinsic neutron source using the MISC code⁵ and/or the SOURCES4C code.⁶ The energy group structure is needed for this input before it is available from the PARTISN output files. Thus, the available energy group structures are listed in data files in the `sensmg/data` directory (set with the `SENS_DATA` environment variable in the `sensmg.py` script; see Sec. III.D). These are only used in fixed-source problems. The available energy-group structures are listed with the `ngroup` keyword in Sec. III.C.

To add a new group structure, follow the example in source file `set_ebins.F`. Increase the number of available group-structure files and add the file name to the `DATA` statement. Add logic to select the correct file name. Finally, put the group structure data file in the `sensmg/data` directory. If it is to be available through the NDI, the data file name is `[G]_lan1`; otherwise, it is `[G]_[LIBNAME]`.

SENSMG gets cross sections from PARTISN outputs. It writes a special PARTISN input file with one problem or edit isotope per zone, with atom density equal to 1 atom/(b·cm), and runs it with "`nosolv = 1,`" a PARTISN block 1 input entry¹ that turns off the solver module but not before PARTISN writes a `macrxs` macroscopic cross-section file and an `sxnedt` microscopic edit cross-section file. The `macrxs` file now has microscopic transport data for every isotope. The fission and capture cross sections for every isotope are read from the `sxnedt` file.

It has been suggested to use existing tools to use the NDI to read cross-section files directly. However, the present scheme allows SENSMG to compute sensitivities for any cross-section library, not just those available through the NDI. The ability to use the Kynea3 library has been very useful.

After computing the neutron source rate density (if necessary), SENSMG runs PARTISN for the forward input file and the special cross-section input file. In subsequent adjoint and generalized adjoint calculations, SENSMG uses the cross-section files from the forward run with PARTISN's "`lib = macrxs`" option.¹ This saves a lot of time.

The adjoint k_{eff} or α from Eq. (12) or (31) is read from the adjoint output file and compared with the forward value; the comparison is printed in the log file. For fixed-source calculations, the adjoint-based leakage is computed using

$$\begin{aligned}
 L &= \int dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) Q^g(r) \\
 &= \int dV \sum_{g=1}^G \phi^{*g}(r) Q^g(r) \\
 &= \sum_{n=1}^{N_z} \sum_{g=1}^G \phi_n^{*g}(r) Q_n^g(r), \quad (69)
 \end{aligned}$$

where n is an index representing coarse meshes and N_z is the number of coarse meshes (zones). Equation (69) is evaluated using PARTISN's edit module for the adjoint calculation, with the forward source $Q_n^g(r)$ in coarse mesh n used as the response function in coarse mesh n .

SENSMG supports older versions of PARTISN (5.69 and 7.72) as well as newer (8, up to 8.27). One place in the Fortran code that depends on the PARTISN version is where the code writes the forward input file. Because it might be of interest to the user, all available reaction-rate edits are specified for the edit isotopes. There are 18 available in PARTISN version 7 (and earlier) but 20 in version 8 (see Table IV). For now, it appears to be an easy thing to support this difference, but if it becomes a burden, then the code will be changed to print only the reaction rates specified in the input file. If an unsupported PARTISN version is used, SENSMG will simply not specify edits in the forward input file.

SENSMG automatically loads the correct modules for the specified PARTISN version and the detected host. As modules, PARTISN versions, and computers get updated and deprecated, this logic will have to be maintained. The “`-use_existing yes`” option is intended for use when the user knows that his/her modules are the correct ones.

New cross-section libraries that become available through the NDI should work automatically. Stand-alone libraries such as Kynea3 can be accommodated with a small amount of effort, using the coding for Kynea3 as an example.

SENSMG is generally LANL-HPC-centric, but it should be straightforward to generalize it to other platforms. In the Python script, modify the default value of `SENS_PARTISN` and the construction of `PARTISN_EXE`, which uses the variable `MACH`, derived from the environment variable `HOSTNAME`. If your system does not use modules, set the default `MY_MODULES` to `yes` (for convenience). Modify the location of your cross-section data files. The Fortran code does not need to be modified.

PARTISN solves Eqs. (18) and (35) using “`nosigf = 1`” to inhibit fission multiplication. PARTISN solves Eq. (35) using “`anorm = [α]`” to put the α/v term on the left side. For Eqs. (18), (35), and (49), the sources are input to PARTISN using the `fixsrc` file rather than the standard input file.

SOURCES4C can be used to compute the intrinsic neutron source for fixed-source problems. [It is presently required for (α, n) sources.] A special version of SOURCES4C was made that does not stop on certain errors (namely, “Target nuclide not found in tape3”, “Target nuclide not found in tape4”, “Source nuclide not found in tape5”, “S.F. source nuclide not found on tape5”, and “D.N. source nuclide not found on tape5”). Also, more digits are written to the SOURCES4C output files in certain places. These digits are not improving the accuracy of the simulation, but they are needed so that sources are balanced correctly, particularly for the central-difference calculations used for verification. In addition, SOURCES4C was modified to print the quantities that are needed to compute Eq. (60) (Ref. 13).

VI. VERIFICATION

This section provides a few results for a few problems. A more thorough verification by comparison with results from MCNP6 (Ref. 18) has been performed for k_{eff} problems for fast spherical systems.^{21,22}

VI.A. Comparison with MCNP6 and SCALE 6.2 for k_{eff} Sensitivities for a Cylinder

A test problem was documented in Ref. 17 in which the sensitivities of k_{eff} to isotopic densities in a homogeneous cylinder were computed using MCNP6 (Ref. 18), SCALE 6.2 (Ref. 20), and PARTISN. Results are repeated here. The fuel is similar to that used in the Transient Reactor Test Facility²³ (TREAT) at Idaho National Laboratory. The height and radius of the reactor are 240 and 60 cm, respectively. The atom density and the mass density of the fuel are 0.113705 atoms/(b·cm) and 2.27 g/cm³, respectively. The composition of the fuel is given in Table VI. The enrichment is 92.9168 wt% ²³⁵U. The input file for SENSMG is listed in Appendix A. The calculations used S_{64} angular quadrature and a P_7 scattering expansion.

In MCNP6, the $S(\alpha, \beta)$ table `grph.20t` was associated with the fuel. Otherwise, ENDF/B-VII.1 cross sections

TABLE VI
Composition of TREAT Fuel

Nuclide	Atom Density [atoms/(b·cm)]	Weight Fraction
¹ H	1.13694E-4	8.38215E-5
¹⁰ B	5.68468E-7	4.16390E-6
C (natural)	1.13579E-1	9.97955E-1
²³⁵ U	1.05735E-5	1.81803E-3
²³⁸ U	7.95855E-7	1.38591E-4

were used. In SCALE, the cross-section table for carbon in graphite was associated with natural carbon, and ENDF/B-VII.1 cross sections were used. SENSMG used Kynea3 with the “C(gph)” $S(\alpha,\beta)$ data for carbon and it used S_{64} angular quadrature and a P_7 scattering expansion. Also it used 16 processors.

Results are shown in Table VII. The deterministic sensitivities are in excellent agreement with the Monte Carlo sensitivities. They are within 1.4% of the MCNP6 results, except for the sensitivity to ¹H, for which the deterministic sensitivity is 11% smaller. It is likely that the deterministic neutron flux spectrum, with only 79 groups, is not quite as thermal as the continuous-energy Monte Carlo spectrum.

VI.B. Comparison with MCNP6 KSEN Results for a Sphere

Results from SENSMG were compared with results from the KSEN card¹⁸ of MCNP6 for a one-dimensional spherical Flattop-Pu model. The SENSMG input file and the MCNP6 input file are listed in Appendix A. The PARTISN sensitivities used 30 groups, S_{32} angular quadrature, and a P_3 scattering expansion.

The sensitivity profiles of k_{eff} to the density of ²³⁵U in the reflector are compared in Fig. 1. The agreement is excellent.

Group-dependent results for the unconstrained (unnormalized) sensitivity of k_{eff} to the chi distribution of ²³⁵U in the reflector are compared in Fig. 2, and results using full normalization are compared in Fig. 3. In both cases, the agreement is excellent. The full-normalization results are in the same output file as the density results of Fig. 1. The unconstrained results were obtained using the same flux files with “-use_existing yes -chinorm none” (Sec. III.C).

VI.C. Comparison with TSUNAMI-1D for a Reaction-Rate Ratio with λ -Mode Flux

As part of a verification for a continuous-energy Monte Carlo GPT capability for SCALE, Perfetti recently published TSUNAMI-1D results.⁷ The multigroup sensitivity tool SENSMG of this paper is more similar to the deterministic TSUNAMI-1D capability than to the Monte Carlo capability, so TSUNAMI-1D results were used for comparison. Reference 7 shows graphs of sensitivity profiles.²⁴

The problem is the one-dimensional spherical Flattop-Pu model and the ratio of interest is the ²³⁷Np fission rate (F37) to the ²³⁵U fission rate (F25) in a central sphere of radius 0.155 cm (whose volume matches that of the cylindrical foil described in Ref. 7). The TSUNAMI-1D calculation used 238 energy groups and S_{16} angular quadrature.⁷ It used ENDF/B-VII.0 cross sections with self-shielding and a P_5 scattering expansion.²⁴ SENSMG used ENDF/B-VII.1 cross sections in 70 energy groups with no self-shielding, S_{32} quadrature, and a P_3 scattering expansion. The SENSMG input file is listed in Appendix A.

The sensitivity profiles of the ratio F37/F25 to the ²³⁹Pu density and fission cross section are shown in Fig. 4, where SENSMG results are compared with

TABLE VII
Sensitivity S_{k,N_j} to Constituents of TREAT Fuel

Nuclide	MCNP	SCALE	PARTISN
¹ H	4.0059E-03 ± 4.37%	4.0194E-03 ± 1.63%	3.5619E-03
¹⁰ B	-2.0248E-01 ± 0.02%	-2.0235E-01 ± 0.00%	-2.0292E-01
C (natural)	5.9836E-01 ± 0.46% ^a	5.8209E-01 ± 1.58%	5.9970E-01
²³⁵ U	3.5071E-01 ± 0.04%	3.5032E-01 ± 0.01%	3.4584E-01
²³⁸ U	-2.5392E-03 ± 0.50%	-2.5494E-03 ± 0.14%	-2.5535E-03
$S(\alpha,\beta)$	2.3217E-01 ± 0.82%	N/A ^b	N/A ^b

^aIncludes the $S(\alpha,\beta)$ sensitivity for the grph table.

^bNot applicable— $S(\alpha,\beta)$ scattering is not calculated separately in SCALE or PARTISN.




Fig. 1. Sensitivity of k_{eff} to ^{235}U density in Flattop-Pu reflector.




Fig. 3. Sensitivity (full normalization) of k_{eff} to ^{235}U chi in Flattop-Pu reflector.




Fig. 2. Sensitivity (unconstrained) of k_{eff} to ^{235}U chi in Flattop-Pu reflector.




Fig. 4. Sensitivity of the ^{237}Np fission to ^{235}U fission ratio to the ^{239}Pu density and fission cross section in Flattop-Pu.

TSUNAMI-1D results (compare with Fig. 9 of Ref. 7, where “total” is “density” of this report; see Sec. IV). The two codes generally agree very well. The peaks in the SENSMG profiles are about 5% smaller than the peaks in the TSUNAMI-1D profiles.

VI.D. Comparison with Direct Perturbations for a Reaction Rate with λ -Mode Flux

A spherical Jezebel model was run. The reaction rate of interest, calculated in a central sphere with a radius of 0.5 cm, was capture in ^{239}Pu . The SENSMG input file is listed in Appendix A. As discussed in Sec. II.B, the reaction rate is put in a ratio with the flux in the reaction-rate region in the denominator. The sensitivity of this reaction rate to the ^{239}Pu capture cross section (indirect effect only, sum of groups) was computed by SENSMG

to be $-3.1228\text{E}-02\%/%$. (The direct effect, of course, is $1\%/%$, which was correctly calculated by SENSMG.) A chi vector was used (`-fissdata 2`) to facilitate the comparisons with direct perturbations.

A direct calculation of this reaction rate is shown in Fig. 5. The reaction rate of interest, called R1, and the flux are plotted as a function of the relative change in the ^{239}Pu capture cross section in Fig. 5a. Figure 5b plots the ratio.

The plots show that the reaction rate, the flux, and the ratio are all very linear in this range of perturbations ($\pm 10\%$). The sensitivity estimated from the slopes of the reaction rate and the flux is $-3.1253\text{E}-02\%/%$, and the sensitivity estimated from the slope of the ratio is $-3.1251\text{E}-02\%/%$. Both are in excellent agreement with the SENSMG result.

To run the direct perturbation cases, the cross sections for each perturbed case were written directly into a PARTISN

input file. A chi matrix cannot be entered this way, which is why a chi vector was used for the sensitivity calculations.

Twenty PARTISN calculations in addition to the base case were run to produce the plots of Fig. 5. This seems like overkill, when only two calculations are required for a central difference. However, in verifying SENSMG, many cases were encountered in which central differences gave unreliable results. Indeed, the difficulty computing central differences is another argument for the use of GPT.

VI.E. Comparison with Direct Perturbations for a Reaction-Rate Ratio with λ -Mode Flux

A spherical Godiva model was run. The reaction-rate ratio of interest, calculated in a central sphere with a radius of 0.574 cm, was fission in ^{238}U to fission in ^{235}U . The SENSMG input file is listed in Appendix A. The number of energy groups was 70. The sensitivity of this reaction rate to the ^{234}U fission cross section in group 30 (indirect effect only; the direct effect is zero) was computed by SENSMG to be $2.1113\text{E-}04\%/%$. A chi vector was used (`-fissdata 2`) to facilitate the comparisons with direct perturbations.




Fig. 5. Sensitivity of the ratio of R1 = ^{239}Pu capture reaction rate to the flux to the ^{239}Pu capture cross section (indirect effect) in a one-dimensional Jezebel model: (a) R1 and flux plotted separately and (b) R1/flux.

The sensitivity of the reciprocal of this reaction rate was also calculated. It was (correctly) the negative of the original sensitivity.

A direct calculation of this reaction-rate ratio is shown in Fig. 6. The numerator, fission in ^{238}U , is R1 and the denominator, fission in ^{235}U , is R2. R1 and R2 are plotted as a function of the relative change in the ^{234}U fission cross section in Fig. 6a. Figure 6b plots the ratio.

Note from the y-axis scales that the changes plotted are extremely small, even though the range of perturbations is $\pm 100\%$. The reason that R1 is jagged is that the reaction rates were obtained from PARTISN's edit output, which prints only six significant figures. A smoother curve would be obtained by reading the binary flux file and the binary edit cross-section file and constructing the reaction rates in postprocessing, as SENSMG does. However, the slope would be close to the least-squares slope shown in Fig. 6 and the conclusions of this section would be the same.

The plots show that R1 is fairly linear and R2 and the ratio are very linear in this range of perturbations ($\pm 100\%$). The sensitivity estimated from the slopes of the reaction rates is $2.3777\text{E-}04\%/%$, and the sensitivity estimated from the slope of the ratio is $2.3778\text{E-}04\%/%$. The difference between these and the SENSMG result is $\sim 13\%$.




Fig. 6. Sensitivity of the ratio of R1 = ^{238}U fission to R2 = ^{235}U fission to the ^{234}U fission cross section in group 30 in a one-dimensional Godiva model: (a) R1 and R2 plotted separately and (b) R1/R2.

The sensitivity for each of the 70 groups was computed directly as in Fig. 6 (20 PARTISN calculations in addition to the base case and a least-squares fit for the slope of the ratio). The sensitivities from SENSMG are compared with the direct sensitivities in Fig. 7. The average energy of group 30, whose result is shown in Fig. 6, is 0.732 MeV; the sensitivity there is the largest in both curves.

Figure 7 shows the difference discussed previously in this section for group 30 and also differences for other energies. The shapes of the curves are similar, but the adjoint-based sensitivities are more negative by ~10% to 15%.

Since the energy-dependent sensitivities are both positive and negative, these differences strongly affect the total (sum over groups). The total adjoint-based sensitivity is $-2.6119\text{E}-04\%/%$ and the total direct sensitivity is $3.3178\text{E}-05\%/%$. The direct sensitivity obtained by perturbing the ^{234}U fission cross section in all 70 groups is $2.6441\text{E}-05$, indicating either nonlinear behavior or inaccuracies in the methodology used to compute direct sensitivities.

It should be noted that the sensitivities computed and discussed in this section are extremely small.

VI.F. Comparison with Direct Perturbations for α and a Reaction-Rate Ratio with α -Mode Flux

This is an α -eigenvalue problem. The problem is the one-dimensional Jezebel model with the benchmark radius (6.39157 cm) but the realistic Pu-alloy density (15.82 g/cm^3) rather than the benchmark density (15.61 g/cm^3). The reaction-rate ratio of interest, fission in ^{238}U to fission in ^{235}U , was computed in a central sphere of radius 0.5 cm. The SENSMG input file is listed in Appendix A. The 30-group MENDF71X cross sections, S_{32} angular quadrature, and P_3 scattering were used. A chi vector was used (`-fissdata 2`) to facilitate




Fig. 7. Sensitivity profile of the ratio of $R1 = ^{238}\text{U}$ fission to $R2 = ^{235}\text{U}$ fission to the ^{234}U fission cross section in a one-dimensional Godiva model.

comparisons with direct perturbations. The α eigenvalue for this model was 0.0373818/sh.

The sensitivities of the α eigenvalue and the reaction-rate ratio to the P_0 and P_1 (separately) ^{239}Pu self scattering cross sections in every group (simultaneously) are compared with central differences (computed using cross-section perturbations of $\pm 1\%$) in Table VIII. The agreement is excellent.

VI.G. Comparison with Direct Perturbations for Leakage and a Reaction-Rate Ratio in a Fixed-Source Problem

This is a fixed-source problem. The problem is a simplified model of the BeRP ball reflected by 3.81 cm of polyethylene.^{25,26} The simplified model has only α -phase plutonium and polyethylene, no steel can or gas fill. The ratio of fission in ^{238}U to fission in ^{235}U was computed in a central sphere of radius 0.25 cm (the actual BeRP ball does not have a cavity for samples). The absolute fission rate in ^{238}U was also computed. The SENSMG input file is listed in Appendix A. The 30-group MENDF71X cross sections, S_{32} and S_{128} angular quadrature, and P_3 scattering were used. A chi vector was used (`-fissdata 2`). The neutron leakage for this model [using MISC for spontaneous fission and SOURCES4C for (α, n)] was $1.81174 \times 10^6/\text{s}$ using S_{32} and $1.80663 \times 10^6/\text{s}$ using S_{128} . A slightly modified version of this problem is discussed in detail in Ref. 13, but without the reaction rates.

The sensitivities of the leakage to the two material densities and to the ^9Be and ^{240}Pu atom densities in the plutonium are shown in Table IX. They are compared with central differences. The results of Table IX used S_{32} quadrature. The agreement is excellent.

The sensitivities of the ratio of fission in ^{238}U to fission in ^{235}U and the absolute fission rate in ^{238}U to the ^{240}Pu density are shown in Table X, where they are compared with central differences. These calculations were done with S_{32} and S_{128} quadrature, as shown in Table X. When going from S_{32} to S_{128} quadrature, the relative differences are reduced by a factor of about 10.

The results of Table X suggest that this problem is not converged with respect to angular quadrature order at S_{32} . Table XI compares several adjoint-weighted quantities as a function of S_N order. In Table XI, L^* is the leakage computed using the solution of Eq. (42):

$$L^* \equiv \int_V dV \sum_{g=1}^G \int_{4\pi} d\hat{\Omega} \psi^{*g}(r, \hat{\Omega}) Q^g(r), \quad (70)$$

TABLE VIII
Sensitivity to ²³⁹Pu Self-Scattering for the High-Density Jezebel Model

Scattering Order	α			Reaction-Rate Ratio		
	SENSMG	Central Difference	Difference	SENSMG	Central Difference	Difference
0	1.63687E+01	1.6338E+01	0.186%	7.96962E-02	7.9501E-02	0.246%
1	-9.49096E+00	-9.4800E+00	0.116%	-9.73175E-02	-9.7199E-02	0.122%

TABLE IX
Sensitivity of the Leakage to Density for the BeRP Ball Model

Density	Adjoint	Central Difference	Difference
Pu bulk mass density	7.20992E+00	7.21712E+00	-0.100%
Poly bulk mass density	1.56066E+00	1.56032E+00	0.022%
⁹ Be atom density	1.15770E-02	1.15836E-02	-0.057%
²⁴⁰ Pu atom density	1.18230E+00	1.18305E+00	-0.064%

TABLE X
Sensitivity of Reaction Rates to ²⁴⁰Pu Density for the BeRP Ball Model

Reaction	S_N	Adjoint	Central Difference	Difference
²³⁸ U(n, f)/ ²³⁵ U(n, f)	32	-1.11391E-02	-1.14502E-02	-2.717%
	128	-1.14238E-02	-1.14488E-02	-0.218%
²³⁸ U(n, f)	32	1.20874E+00	1.20539E+00	0.277%
	128	1.20497E+00	1.20461E+00	0.030%

TABLE XI
Adjoint-Weighted Quantities as a Function of S_N Order for the BeRP Ball Model

S_N	Fine Mesh Width (cm)	$(L^* - L)/L$ (Should Be 0)	Equation (51) (Should Be 0)	Equation (52) (Should Be 1)
16	0.005	-2.3660E-03	9.0572E-04	1.009210
32	0.005	-6.0765E-04	2.7421E-04	1.002783
64	0.005	-1.5487E-04	7.7350E-05	1.000797
128	0.005	-4.5194E-05	2.1796E-05	1.000219
256	0.005	-1.6662E-05	7.8144E-06	1.000070
512	0.005	-2.4125E-06	4.3070E-06	1.000038
1024	0.005	-3.2741E-06	5.4009E-06	1.000025
1024	0.0005	3.1054E-06	1.0501E-06	1.000007

which should be equal to L of Eq. (43). Table XI also evaluates Eq. (51) for the reaction-rate ratio and Eq. (52) for the absolute reaction rate. These quantities are all quite different from their exact values at low S_N orders

but they all approach their exact values as the S_N order increases, particularly when the spatial discretization is refined as well. The PARTISN convergence criterion for all cases was 10^{-6} .

The results of this section suggest that adjoint-based sensitivities for leakage problems require larger angular quadrature orders than might be expected.

VI.H. Comparison with Direct Perturbations for Leakage from a Slab in a Fixed-Source Problem

This is a fixed-source problem. The problem is a three-region slab. The left region is 1.8 cm of polyethylene, the middle region is 2 cm of a simplified version of the BeRP plutonium, and the right region is 1.8 cm of polyethylene. The only quantity of interest is the leakage from the right side of the slab. The SENSMSG input file is listed in [Appendix A](#). The 30-group MENDF71X cross sections, S_{32} angular quadrature, and P_3 scattering were used. A chi vector was used (`-fissdata 2`). The neutron leakage for this model [using MISC for spontaneous fission; there was no (α, n)] was $1.08877 \times 10^4/s$.

The sensitivities of the leakage to the four boundaries and internal interfaces are shown in [Table XII](#). They are compared with central differences. The agreement is excellent.

The derivatives of [Table XII](#) are not symmetric about the slab's center because the quantity of interest is the leakage from the right side, not the total leakage.

VI.I. Comparison with Direct Perturbations for k_{eff} and Total Neutron Leakage from a Cylinder

Constant-density adjoint-based derivatives of k_{eff} and total neutron leakage with respect to interface and outer boundary locations for a cylinder are compared in detail with direct perturbations in [Ref. 12](#). These calculations used the `rplane/zplane` capability ([Sec. III.C](#)). General comparisons for constant-volume adjoint-based derivatives with respect to material mass density are also given in [Ref. 12](#). The main point of [Ref. 12](#) is to derive an equation for constant-mass derivatives with respect to interface and outer boundary locations and to make comparisons with direct perturbations. The SENSMSG

input files used for the neutron problems in [Ref. 12](#) are listed in [Appendix A](#).

For these problems, particularly the k_{eff} problem, the moments expansion approximation of the inner product of angular fluxes, [Eq. \(27\)](#), was extremely inaccurate. Using it lead to errors of 11%, 13%, and 15% in the sensitivity of k_{eff} to the mass density of lead, aluminum, and ^{238}U , respectively (and 0.3% in the sensitivity of k_{eff} to the mass density of ^{235}U). A seventeenth-order expansion was required to reduce all of the errors below 0.1%.

These results suggest that caution should be applied when using the moments expansion approximation. Better memory management in SENSMSG and the ability for PARTISN to print mesh-centered angular flux files from MPI runs of cylindrical problems would obviate this problem.

VII. SUMMARY AND FUTURE WORK

SENSMSG is a tool for calculating the first-order sensitivities of reaction-rate ratios, k_{eff} , and α in critical problems and reaction-rate ratios, reaction rates, and leakage in fixed-source problems to multigroup cross sections, isotope densities, material mass densities, and interface locations using PARTISN. SENSMSG treats only GPT-allowable responses.⁴ SENSMSG is a combination of Python and Fortran and was developed under Linux.

For all of the problems that SENSMSG can treat, the constant-volume and constant-density partial derivatives can be combined (with cell volumes and interface areas) to compute constant-mass partial derivatives.¹

SENSMSG continues to undergo verification. Future work for SENSMSG includes:

1. implement a more efficient strategy for memory management so that larger cylindrical problems can be run and angular fluxes used
2. develop and implement a better method for an absolute reaction rate for critical problems
3. implement a more efficient convergence scheme for the generalized adjoint calculations

TABLE XII

Sensitivity of the Leakage to Boundaries and Internal Interfaces for the BeRP Slab

Surface Location (cm)	Adjoint	Central Difference	Difference
-2.8	-9.58828E+03	-9.58820E+03	0.0008%
-1.0	-3.25842E+04	-3.25967E+04	-0.0384%
1.0	3.39701E+04	3.39837E+04	-0.0398%
2.8	8.20235E+03	8.20220E+03	0.0018%

4. compute the derivatives of reaction-rate ratios to material densities
5. compute the derivatives of reaction-rate ratios to internal interface locations
6. compute higher-order sensitivities
7. implement a continue-run capability.

It would be worth making some modifications to PARTISN that would allow for more efficient and accurate sensitivity calculations. Two specific suggestions are

1. have PARTISN print more flux moments than the number used in the transport
2. implement a solver such as described in Ref. 4 so that the successive approximation scheme is unnecessary. (This would obviate items 2 and 3 in the SENSMG to-do list.)

SENSMG can be modified to accommodate other deterministic transport codes that have an adjoint capability.

APPENDIX A SENSMG AND MCNP6 INPUT FILES

A.I. SENSMG Input File for Sec. VI.A

(Because of memory management issues, this file will only run if it has the title shown [line 1] and if it is named “bck01”.)

```
brian's problem, inspired by TREAT
keff 2d
kynea3
1 / no of materials
1 1001 8.38215E-05 5010 4.16390E-06 106012 9.97955E-01 92235 1.81803E-03 92238 1.38591E-04 / TREAT fuel
2.27 / densities
1 1 / no of r layers, no of z layers
    60. / outer radii
    0.240. / heights
1 / material nos
0 0 / index of coarse mesh to use for reaction rates
0 / number of reaction-rate ratios
```

A.II. SENSMG Input File for Sec. VI.B

```
Pu-Flattop (PU-MET-FAST-006)
keff 1d
mendf71x
2 / no of materials
1 94239 9.38001E-01 94240 4.79988E-02 94241 2.99996E-03 31069 6.53652E-03 31071 4.46355E-03 / Pu-alloy
2 92234 5.40778E-05 92235 7.10966E-03 92238 9.92836E-01 / Flattop natural U
15.53 19.00 / densities
3 / no of shells
0.5 4.5332 24.142 / outer radii
1 1 2 / material nos
1 / index of coarse mesh to use for reaction rates
0 / number of reaction-rate ratios
```

A.III. MCNP6 Input File for Sec. VI.B

```
Pu-Flattop (PU-MET-FAST-006)
```

```

1  1  -15.53  -1      imp:n = 1
2  1  -15.53  -2    1  imp:n = 1
3  2  -19.00  -3    2  imp:n = 1
9  0                      3  imp:n = 0

1  so  0.5
2  so  4.5332
3  so  24.142

mode n
totnu
rand gen = 2 seed = 100001
kcode 1200000 1.0 50 2050
prdmp j 500
sdef rad = d1 erg = d2
si1  0. 24.142
sp1  -21 2
sp2  -3 0.988 2.249
m1   94239-9.38001E-01
      94240-4.79988E-02
      94241-2.99996E-03
      31069-6.53652E-03
      31071-4.46355E-03
m2   92234-5.40778E-05
      92235-7.10966E-03
      92238-9.92836E-01
m3   77191 1.
m4   92235 1.
m5   77193 1.
m6   77193 1.
e0   1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
      8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
      1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
      1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
      1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
      7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
      1.7000E+01t

f04:n 1
fm04 (1. 3-1) $ total
sd04 1.
f14:n 1
fm14 (1. 4-6) $ fission
sd14 1.
f24:n 1
fm24 (1. 5-1) $ total
sd24 1.
f34:n 1
fm34 (1. 6-2:-6) $ absorption (capture+fission)
sd34 1.
print -30
kopts  blocksize = 5
c total

```

```

ksen01  xs    cell = (1 2)
ksen02  xs    cell = (1 2)
        erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
           8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
           1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
           1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
           1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
           7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
           1.7000E+01

c chi
ksen03  xs    cell = (1 2) rxn = -1018 constrain = yes
        erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
           8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
           1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
           1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
           1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
           7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
           1.7000E+01

c chi
ksen04  xs    cell = (1 2) rxn = -1018 constrain = no
        erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
           8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
           1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
           1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
           1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
           7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
           1.7000E+01

c nu
ksen05  xs    cell = (1 2) rxn = -7
        erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
           8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
           1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
           1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
           1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
           7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
           1.7000E+01

c total
ksen11  xs    cell = 3
ksen12  xs    cell = 3
        erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
           8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
           1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
           1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
           1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
           7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
           1.7000E+01

c chi
ksen13  xs    cell = 3 rxn = -1018 constrain = yes
        erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
           8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
           1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02

```

```

1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
1.7000E+01

```

c chi

```

ksen14  xs    cell = 3 rxn = -1018 constrain = no
erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
      8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
      1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
      1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
      1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
      7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
      1.7000E+01

```

c nu

```

ksen15  xs    cell = 3 rxn = -7
erg = 1.3900E-10 1.5200E-07 4.1400E-07 1.1300E-06 3.0600E-06
      8.3200E-06 2.2600E-05 6.1400E-05 1.6700E-04 4.5400E-04
      1.2350E-03 3.3500E-03 9.1200E-03 2.4800E-02 6.7600E-02
      1.8400E-01 3.0300E-01 5.0000E-01 8.2300E-01 1.3530E+00
      1.7380E+00 2.2320E+00 2.8650E+00 3.6800E+00 6.0700E+00
      7.7900E+00 1.0000E+01 1.2000E+01 1.3500E+01 1.5000E+01
      1.7000E+01

```

A.IV. SENSMG Input File for Sec. VI.C

```

Pu-Flattop (PU-MET-FAST-006)
keff
mendf71x
2 / no of materials
1 94239 9.38001E-01 94240 4.79988E-02 94241 2.99996E-03 31069 6.53652E-03 31071 4.46355E-03 / Pu-alloy
2 92234 5.40778E-05 92235 7.10966E-03 92238 9.92836E-01 / Flattop natural U
15.53 19.00 / densities
3 / no of shells
0.155 4.5332 24.142 / outer radii
  1 1 2 / material nos
  1 / index of coarse mesh to use for reaction rates
  2 / number of reaction-rate ratios
  92238 16 92235 16/numerator and denominator
  93237 16 92235 16/numerator and denominator

```

A.V. SENSMG Input File for Sec. VI.D

```

Pu-239 Jezebel (PU-MET-FAST-001)
1d keff
mendf71x / libname
1/no of materials
1 94239 9.42090E-01 94240 4.47189E-02 94241 2.99354E-03 31069 6.12984E-03 31071 4.06790E-03 / Pu-alloy
15.61 / densities
2 / no of shells

```

```

0.5 6.39157 / outer radii
  1  1      /material nos
  1      / index of coarse mesh to use for reaction rates
  1      / number of reaction-rate ratios
94239 10    / denominator is flux

```

A.VI. SENSMG Input File for Sec. VI.E

```

HEU-MET-FAST-001-02 Godiva
1d keff
mendf71x / libname
1 / no of materials
1 92234 1.01999E-02 92235 9.37100E-01 92238 5.27004E-02/Godiva
18.74 / densities
2 / no of shells
0.5738912155190165 8.7407 / outer radii
  1  1      / material nos
  1      / index of coarse mesh to use for reaction rates
  2      / number of reaction-rate ratios
  92238 16  92235 16 / numerator and denominator
  92235 16  92238 16 / numerator and denominator

```

A.VII. SENSMG Input File for Sec. VI.F

```

Pu-239 Jezebel (benchmark radius, realistic density)
alpha 1d
mendf71x / libname
1 / no of materials
1 94239 9.42090E-01 94240 4.47189E-02 94241 2.99354E-03 31069 6.12984E-03 31071 4.06790E-03 / Pu-alloy
15.82 / densities
2/no of shells
0.5 6.39157 / outer radii
  1  1      / material nos
  1      / index of coarse mesh to use for reaction rates
  1      / number of reaction-rate ratios
  92238 16  92235 16 / numerator and denominator

```

A.VIII. SENSMG Input File for Sec. VI.G

(Though the composition of material 1 wraps here, it should be on a single line.)

```

BeRP ball, 38.1 mm reflector, no steel or NO
1d lkg
mendf71x
2/no of materials
1 31069 1.99071E-04 31071 1.35938E-04 6000 2.29800E-04 26054 5.65054E-07 26056 9.19000E-06 26057
2.16145E-07 26058 2.90477E-08 20040 2.89993E-06 20042 2.03213E-08 20043 4.34122E-09 20044 6.86363E-08
20046 1.37596E-10 20048 6.71243E-09 42092 1.27960E-06 42094 8.14938E-07 42095 1.41752E-06 42096
1.50082E-06 42097 8.68256E-07 42098 2.21645E-06 42100 9.02647E-07 40090 5.07075E-05 40091 1.11812E-05
40092 1.72786E-05 40094 1.78916E-05 40096 2.94387E-06 11023 5.00013E-05 24050 2.08690E-07 24052
4.18508E-06 24053 4.83692E-07 24054 1.22671E-07 28058 3.36013E-06 28060 1.33868E-06 28061 5.91747E-08

```



```

28062 1.91509E-07 28064 5.06485E-08 4009 1.00003E-06 5010 1.84314E-07 5011 8.15712E-07 12024 7.79520E-07
12025 1.02804E-07 12026 1.17702E-07 94238 1.59585E-04 92234 3.94427E-05 94239 9.35361E-01 92235
7.51202E-04 94240 5.92435E-02 92236 1.75085E-04 94241 6.77410E-04 95241 2.48513E-03 93237 7.84266E-05
94242 2.79593E-04/BeRP Pu from John Mattingly
2 6000 8.56299E-01 1001 1.43701E-01/HDPE, C2H4
19.6 0.95/densities
3/no of shells
    0.25 3.794 7.604/radii
    1 1 2 / material nos
    1 / index of coarse mesh to use for reaction rates
    2 / number of reaction-rate ratios
    92238 16 92235 16 / numerator and denominator
    92238 16 / numerator and denominator

```

A.IX. SENSMG Input File for Sec. VI.H

```

BeRP ball slab
slab lkg
mendf71x
2/no of materials
1 94239 9.38039E-01 94240 5.94113E-02 31069 1.51516E-03 31071 1.03465E-03/simple BeRP
2 6000 8.56299E-01 1001 1.43701E-01/HDPE, C2H4
19.6 0.95/densities
3/no of shells
    -2.800 -1.000 1.000 2.800/radii
           2 1 2 / material nos
    1 / index of coarse mesh to use for reaction rates
    0 / number of reaction-rate ratios
    92238 16 92235 16/numerator and denominator
    92238 16 / numerator and denominator

```

A.X. SENSMG Input File for Sec. VI.I (Ref. 12)

(This is the k_{eff} input file. To change to the neutron leakage input, change “keff” to “lkg”.)

```

4 material input file U235 U238 Al Pb
2d keff
mendf70x
4/no. of materials
1 92235 1.000000 /
2 92238 1.000000 /
3 13027 1.000000 /
4 82000 1.000000 /
18.00 18.00 2.70 11.4/
3 7/number of cylinders, number of heights
1. 2. 4./
0. 1. 1.5 2. 4. 4.5 6. 7./heights, bottom to top
    3 3 3/material map, top layer
    1 1 3/material map, next layer

```

0 1 3/material map, next layer
 0 1 4/material map, next layer
 1 1 4/material map, next layer
 4 4 4/material map, next layer
 2 2 2/material map, bottom layer
 1 4/index of coarse mesh to use for reaction
 rates
 0/number of reaction rates
 79197 10 92235 16/

APPENDIX B ISOTOPES AVAILABLE IN THE KYNEA3 LIBRARY

The ZAIDs listed here are the ones to enter in the input file; SENSMG converts them to the names when writing the PARTISN forward input file. The ZAIDS

are intuitive except for four special ones that are needed to distinguish between pairs of the same isotope: 101001 and 1001 for “H1(H2O)” and “H1,” respectively; 104009 and 4009 for “Be9(th)” and “Be9,” respectively; 6012 and 106012 for “C12” and “C12(gph),” respectively; and 95242 and 195242 for “Am242” and “Am242m,” respectively. Note that “H1” does have an $S(\alpha,\beta)$ association: It is “H1(CH2)” from the Vitamin-B6 library.¹⁵

The order listed in Table B.I is the order that they appear in the library. The last seven (from “H3” through “Li7”) do not have atomic weights in the library. Their atomic weights are hard-coded in the SENSMG source code.

The neutron group velocities in Kynea3 are in units of centimeters/microsecond, so divide by 100 to get centimeters/shake.

TABLE B.I
 Isotopes Available in the Kynea3 Library

ZAID	Name	ZAID	Name	ZAID	Name	ZAID	Name	ZAID	Name
1000	H	17000	Cl	40000	Zr	74184	W184	94237	Pu237
101001	H1(H2O)	19000	K	41000	Nb	74186	W186	94238	Pu238
1001	H1	20000	Ca	41093	Nb93	75000	Re	94239	Pu239
1002	H2	22000	Ti	42000	Mo	75185	Re185	94240	Pu240
4000	Be	23000	V	47000	Ag	75187	Re187	94241	Pu241
104009	Be9(th)	24000	Cr	47107	Ag107	79000	Au	94242	Pu242
4009	Be9	24050	Cr50	47109	Ag109	79197	Au197	94243	Pu243
5000	B	24052	Cr52	48000	Cd	82000	Pb	94244	Pu244
5010	B10	24053	Cr53	49000	In	82206	Pb206	95241	Am241
5011	B11	24054	Cr54	50000	Sn	82207	Pb207	95242	Am242
6000	C	25000	Mn	56000	Ba	82208	Pb208	195242	Am242m
6012	C12	25055	Mn55	56138	Ba138	83000	Bi	95243	Am243
106012	C12(gph)	26000	Fe	63000	Eu	83209	Bi209	96241	Cm241
7000	N	26054	Fe54	63151	Eu151	90000	Th	96242	Cm242
7014	N14	26056	Fe56	63152	Eu152	90230	Th230	96243	Cm243
7015	N15	26057	Fe57	63153	Eu153	90232	Th232	96244	Cm244
8000	O	26058	Fe58	63154	Eu154	91000	Pa	96245	Cm245
8016	O16	27000	Co	63155	Eu155	91231	Pa231	96246	Cm246
8017	O17	27059	Co59	72000	Hf	91233	Pa233	96247	Cm247
9000	F	28000	Ni	72174	Hf174	92000	U	96248	Cm248
9019	F19	28058	Ni58	72176	Hf176	92232	U232	1003	H3
11000	Na	28060	Ni60	72177	Hf177	92233	U233	2000	He
11023	Na23	28061	Ni61	72178	Hf178	92234	U234	2003	He3
12000	Mg	28062	Ni62	72179	Hf179	92235	U235	2004	He4
13000	Al	28064	Ni64	72180	Hf180	92236	U236	3000	Li
13027	Al27	29000	Cu	73000	Ta	92237	U237	3006	Li6
14000	Si	29063	Cu63	73181	Ta181	92238	U238	3007	Li7
15000	P	29065	Cu65	73182	Ta182	93237	Np237		
15031	P31	31000	Ga	74000	W	93238	Np238		
16000	S	39000	Y	74182	W182	93239	Np239		
16032	S32	39089	Y89	74183	W183	94236	Pu236		

Acknowledgments

SENSMG was begun in July 2016 with encouragement from M. C. White and P. Talou (LANL) and funding from the U.S. Department of Energy Advanced Simulation and Computing program, Physics and Engineering Models project, at LANL. Since October 2016, SENSMG has been funded by the U.S. National Nuclear Security Administration's Office of Defense Nuclear Nonproliferation Research & Development. C. Perfetti at ORNL ran the SCALE calculations for Sec. VI.A and provided the TSUNAMI-1D numerical results Sec. VI.C.

This manuscript has been authored by Los Alamos National Security under Contract No. De-AC52-06NA25396 with the U.S. Department of Energy. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

References

1. R. E. ALCOUFFE et al., "PARTISN: A Time-Dependent, Parallel Neutral Particle Transport Code System," LA-UR-08-7258 (revised Sep. 2017), Los Alamos National Laboratory (2017).
2. M. L. WILLIAMS, "Perturbation Theory for Nuclear Reactor Analysis," *CRC Handbook of Nuclear Reactor Calculations*, Vol. III, Y. RONEN, Ed., CRC Press, Boca Raton, Florida (1986).
3. W. M. STACEY JR., *Variational Methods in Nuclear Reactor Physics*, Chap. 1, Academic Press, New York (1974).
4. D. G. CACUCI, "A Paradigm-Shifting Methodology for Sensitivity Analysis of Multiplying Nuclear Systems," *Nucl. Sci. Eng.*, **185**, 3, 361 (2017); <https://doi.org/10.1080/00295639.2016.1272993>.
5. C. J. SOLOMON, "MCNP Intrinsic Source Constructor (MISC): A User's Guide," Internal Memorandum LA-UR-12-20252 (updated Mar. 27, 2012), Los Alamos National Laboratory (2012).
6. W. B. WILSON et al., "SOURCES 4C: A Code for Calculating (α,n), Spontaneous Fission, and Delayed Neutron SOURCES and Spectra," *Proc. American Nuclear Society/Radiation Protection and Shielding Division 12th Biennial Topl. Mtg.*, Santa Fe, New Mexico, April 14–18, 2002, American Nuclear Society (2002).
7. C. M. PERFETTI and B. T. REARDEN, "Development of a Generalized Perturbation Theory Method for Sensitivity Analysis Using Continuous-Energy Monte Carlo Methods," *Nucl. Sci. Eng.*, **182**, 354 (2016); <https://doi.org/10.13182/NSE15-13>.
8. J. A. FAVORITE, "Second-Order Reactivity Worth Estimates Using an Off-the-Shelf Multigroup Discrete Ordinates Transport Code," *Proc. Int. Conf. Physics of Reactors (PHYSOR'08)*, Interlaken, Switzerland, Paul Scherrer Institut, September 14–19, 2008.
9. R. T. EVANS and D. G. CACUCI, "A Parallel Krylov-Based Adjoint Sensitivity Analysis Procedure," *Nucl. Sci. Eng.*, **172**, 216 (2012); <https://doi.org/10.13182/NSE11-110>.
10. J. A. FAVORITE, "Flux Moments and Inner Products (U)," X-1–RN(U)09–02 (Nov. 25, 2008), Los Alamos National Laboratory (2008).
11. J. A. FAVORITE and K. C. BLEDSOE, "Eigenvalue Sensitivity to System Dimensions," *Ann. Nucl. Energy*, **37**, 4, 522 (2010); <https://doi.org/10.1016/j.anucene.2010.01.004>.
12. J. A. FAVORITE, "Adjoint-Based Constant-Mass Partial Derivatives," *Ann. Nucl. Energy*, **110**, 1052 (2017); <https://doi.org/10.1016/j.anucene.2017.08.015>.
13. J. A. FAVORITE and S. L. WEIDENBENNER, "Sensitivity of a Response to the Composition of an (α,n) Neutron Source," *20th Topl. Mtg. Radiation Protection and Shielding Division of the American Nuclear Society (RPSD-2018)*, Santa Fe, New Mexico, August 26–31, 2018, American Nuclear Society (accepted).
14. K. C. BLEDSOE, J. A. FAVORITE, and T. ALDEMIR, "Using the Levenberg-Marquardt Method for Solutions of Inverse Transport Problems in One- and Two-Dimensional Geometries," *Nucl. Technol.*, **176**, 1, 106 (2011); <https://doi.org/10.13182/NT176-106>.
15. E. S. VARLEY and J. MATTINGLY, "Rapid Feynman-Y Synthesis: Kynea3 Cross-Section Library Development," *Trans. Am. Nucl. Soc.*, **98**, 575 (2008).
16. S. FRANKLE, "A Prototype Fine-Group Library for Gamma-Ray Spectroscopy, ACTI," Internal Memorandum XCI:SCF–99–19 (U) (updated Feb. 2, 2004), Los Alamos National Laboratory (2004).
17. J. A. FAVORITE et al., "Adjoint-Based Sensitivity and Uncertainty Analysis for Density and Composition: A User's Guide," *Nucl. Sci. Eng.*, **185**, 2, 384 (2017); <https://doi.org/10.1080/00295639.2016.1272990>.
18. "MCNP® User's Manual, Code Version 6.2," C. J. WERNER, Ed., LA-UR-17-29981, Rev. 0 (Oct. 27, 2017), Los Alamos National Laboratory (2017).
19. B. C. KIEDROWSKI, University of Michigan, Private Communication regarding clarification of CONSTRAIN on KSEN card (2016).
20. "SCALE Code System," B. T. REARDEN and M. A. JESSEE, Eds., ORNL/TM-2005/39, Version 6.2, Oak Ridge National Laboratory (2016).
21. W. HAECK et al., "A Comparison of Monte Carlo and Deterministic Solvers for k_{eff} and Sensitivity

- Calculations,” LA-UR-17-31177 (Dec. 1, 2017), Los Alamos National Laboratory (2017); <http://permalink.lanl.gov/object/tr?what=info:lanl-repo/lareport/LA-UR-17-31177> (current as of Mar. 26, 2018).
22. W. HAECK et al., “Comparison of Monte Carlo and Deterministic Solvers for k_{eff} and Sensitivity Calculations,” *Proc. Int. Conf. Physics of Reactors (PHYSOR’18)*, Cancun, Mexico, April 22–26, Mexican Nuclear Society 2018.
23. J. D. BESS and M. D. DeHART, “TREAT Fuel Assembly Characterization for Modern Neutronics Validation Methods,” *Trans. Am. Nucl. Soc.*, **112**, 373 (2015).
24. C. M. PERFETTI, Oak Ridge National Laboratory, Private Communication regarding TSUNAMI results (2016).
25. J. MATTINGLY, “Polyethylene-Reflected Plutonium Metal Sphere: Subcritical Neutron and Gamma Measurements,” SAND2009-5804 Revision 3 (revised July 2012), Sandia National Laboratories (2012).
26. E. C. MILLER et al., “Computational Evaluation of Neutron Multiplicity Measurements of Polyethylene-Reflected Plutonium Metal,” *Nucl. Sci. Eng.*, **176**, 167 (2014); <https://doi.org/10.13182/NSE12-53>.