

## Two $\beta$ -amylase genes, *OsBAM2* and *OsBAM3*, are involved in starch remobilization in rice leaf sheaths

Tatsuya Hirano, Takayuki Higuchi, Minako Hirano, Yu Sugimura & Hiroyasu Michiyama

To cite this article: Tatsuya Hirano, Takayuki Higuchi, Minako Hirano, Yu Sugimura & Hiroyasu Michiyama (2016) Two  $\beta$ -amylase genes, *OsBAM2* and *OsBAM3*, are involved in starch remobilization in rice leaf sheaths, *Plant Production Science*, 19:2, 291-299, DOI: [10.1080/1343943X.2016.1140008](https://doi.org/10.1080/1343943X.2016.1140008)

To link to this article: <https://doi.org/10.1080/1343943X.2016.1140008>



© 2016 The Author(s). Published by Taylor & Francis



Published online: 16 Mar 2016.



Submit your article to this journal [↗](#)



Article views: 894



View related articles [↗](#)



Citing articles: 4 View citing articles [↗](#)

## Two $\beta$ -amylase genes, *OsBAM2* and *OsBAM3*, are involved in starch remobilization in rice leaf sheaths

Tatsuya Hirano, Takayuki Higuchi, Minako Hirano, Yu Sugimura and Hiroyasu Michiyama

Graduate School of Agricultural Science, Meijo University, Nagoya, Japan

### ABSTRACT

To identify mechanisms of starch degradation in rice leaf sheaths at the post-heading stage, we investigated the function of *OsBAM2* and *OsBAM3*, which encode plastid-targeted active  $\beta$ -amylase isoforms, in starch remobilization in leaf sheaths. The starch content in the second leaf sheaths below the flag leaf (the third leaf sheaths) peaked at the flag leaf emergence stage and gradually decreased until 15 days after heading. The mRNA levels of *OsBAM2* and *OsBAM3* in the third leaf sheaths increased from the flag leaf emergence stage to the heading stage when the starch content began to decrease. However, these mRNA levels did not always remain high during post-heading. Overexpression of *OsBAM2* or *OsBAM3* markedly repressed starch accumulation in the third leaf sheaths, showing that *OsBAM2* and *OsBAM3* function in starch degradation in rice leaf sheaths. In contrast, no significant differences in starch content in the third leaf sheaths were detected between knockdown plants of *OsBAM2* or *OsBAM3* and non-transgenic wild-type plants. Our results suggest that reduced expression of the individual genes, *OsBAM2* or *OsBAM3*, does not result in excess accumulation of starch in the leaf sheaths, probably because of the complementary function of another gene or the action of other genes encoding starch-degrading.

**Abbreviation:** AGPase: ADP-glucose pyrophosphorylase; BSA: bovine serum albumin; EDTA: ethylenediamine tetraacetate; GBSS: granule-bound starch synthase; GWD: glucan, water dikinase; KD: knockdown; OE: overexpression; PCR: polymerase chain reaction; PWD: phosphoglucan, water dikinase; RT-PCR: reverse transcriptase polymerase chain reaction; SBE: starch branching enzyme; SStandard error; SSS: soluble starch synthase; WT: non-transgenic wild-type

### ARTICLE HISTORY

Received 5 November 2015  
Revised 9 December 2015  
Accepted 11 December 2015

### KEYWORDS

$\beta$ -Amylase; leaf sheath;  
*Oryza sativa* L; starch  
degradation

Rice (*Oryza sativa* L.) leaf sheaths serve as temporary storage organs for photoassimilates. Starch is accumulated at a high level in leaf sheaths prior to the heading stage and remobilized for grain filling after heading (Perez et al., 1971). Approximately 30% of the final grain yield is derived from carbohydrates accumulated transiently in the leaf sheaths and culms (Cock & Yoshida, 1972). Thus, identifying the mechanisms of starch accumulation at the pre-heading stage and starch remobilization at the post-heading stage in leaf sheaths is important for improving rice grain yield.

Physiological and biochemical studies have been performed to investigate the correlation between starch accumulation and starch-synthesizing enzymes in rice leaf sheaths (He et al., 2005; Hirano et al., 2005; Hirose et al., 1999, 2006; Perez et al., 1971; Watanabe et al., 1997). The activity of granule-bound starch synthase (GBSS) is positively correlated with the starch content of leaf sheaths (Perez et al., 1971). Watanabe et al. (1997) suggested that

starch branching enzyme (SBE) is primarily involved in the regulation of starch accumulation in the leaf sheaths. The starch content of leaf sheaths at the heading stage was lower in rice plants grown under a high-nitrogen fertilizer regime than that under a low-nitrogen fertilizer regime, which was accompanied by a decrease in SBE activity (Hirano et al., 2005). Hirose et al. (2006) have shown that starch accumulation in leaf sheaths before the heading stage was accompanied by a transient rise in ADP-glucose pyrophosphorylase (AGPase), soluble starch synthase, GBSS, and SBE activities. They analyzed in detail the transcript levels of genes encoding these enzymes to develop and propose a profile of starch synthesis in leaf sheaths for comparison with that in endosperms. In this profile, *AGPL1*, *AGPS1*, *SSI*, *SSIIb*, *SSIIIb*, *GBSSII*, and *BEIIa* were expressed predominantly in leaf sheaths during the pre-heading stage (Hirose et al., 2006).

In contrast, the details of the enzymes involved in starch remobilization in leaf sheaths are largely unknown. Only

**Table 1.** Gene-specific primers used in this study.

Gene name	RAP-ID	Forward primer (5' → 3')	Reverse primer (5' → 3')
<RT-PCR>			
<i>OsBAM2</i>	Os10g0465700	CCTTTTGGGGTGGTTCATCTG	GCCATCGGTAGAAAATCTTAACCCG
<i>OsBAM3</i>	Os03g0141200	CGCCTGTCTGTCTGTTTGATTGATTC	TGAGCCAGGTTGAGCGACATC
<i>OsBAM4</i>	Os01g0236800	GAACAGGTTACCAAGTTCGTCAG	CCGACATCGAGGAGAGGTAATG
<i>OsBAM5</i>	Os10g0565200	TAGTATGGAGGCGAGCTTCGTGGC	CTGCAAACCTCCCCGATAATC
<i>OsBAM6</i>	Os07g0543100	CCGATCAGGATTGCTGTCTC	TCTCGCCGTCGAACGGATAC
<i>OsBAM8</i>	Os09g0569200	ACCCTTGGCAAAGAGTTG	AGGATACCACCTGCTCAC
<i>OsBAM9</i>	Os03g0351300	TTCACGTACCAGCGGATG	GCTGCGTCTACACTGTCTG
<i>OsEF1<math>\alpha</math>1</i>	Os03g0177400	AAGAACGGTGATGCTGGTATGG	CCACTTGAACGAGGCTCATTTTC
<Overexpression vector>			
<i>OsBAM2</i>		CGGGATCCATGGCCTTGAACCTTGGCTCA	CGGGATCCTATGAGCGTCTGACCGTGCTAG
<i>OsBAM3</i>		CGGGATCCATGATGTCGCTCAACCTGGC	CGGGATCCTTTTACCCTTCAACGCGCGAGG
<RNAi vector>			
<i>OsBAM2</i>		CACCCAATCCCCAACCTCCAATA	ACGCCGTCTCCTCTGTCTGCT
<i>OsBAM3</i>		CACCATCGTCGCTTCGGCGTAAC	GCGGCATCATCACGTACACC

a few reports have addressed the correlations between changes in starch content and starch-degrading enzymes in leaf sheaths during the post-heading stage. After heading, the activity of  $\alpha$ -amylase was consistent with the degree of starch degradation in leaf sheaths (Ishimaru et al., 2004). Chen and Wang (2008) reported that  *$\alpha$ -amylase 2A (RAmy2A)* and  *$\beta$ -amylase (BAM)* genes (Accession Number; AK068968, *OsBAM3*) were highly expressed in leaf sheaths at the post-heading stage. Recently, Sugimura et al. (2015) proposed that *RAmy2A* genes play an important role in leaf sheath starch degradation at the post-heading stage. These findings seem to support the involvement of  $\alpha$ -amylase in starch degradation in leaf sheaths during the post-heading stage.

A pathway of starch degradation in leaves has been well elucidated in *Arabidopsis thaliana* (L.) Heynh and potato (Zeeman et al., 2010). The starch degradation depends on the reversible phosphorylation of glucans at the surface of the starch granule (Zeeman et al., 2010). Two classes of enzymes, glucan, water dikinase (GWD) (Mikkelsen et al., 2004; Ritte et al., 2002, 2004; Yu et al., 2001) and phosphoglucan, water dikinase (PWD) (Baunsgaard et al., 2005; Kötting et al., 2005), are implicated in the glucan phosphorylation. Maltose is the major product of starch hydrolysis and is exported from chloroplasts at night (Chia et al., 2004; Fulton et al., 2008; Niittylä et al., 2004; Weise et al., 2004). The hydrolysis of  $\alpha$ -1, 4-glucan chains is catalyzed primarily by  $\beta$ -amylases localized in chloroplasts (Fulton et al., 2008; Kaplan & Guy, 2005; Lao et al., 1999; Scheidig et al., 2002; Sparla et al., 2006). There are nine genes encoding  $\beta$ -amylase-like proteins in the *A. thaliana* genome. Total  $\beta$ -amylase activity was significantly reduced in the leaves of *bam1* and *bam3* mutants as compared with the wild-type (WT) plants (Fulton et al., 2008). The *bam3* mutant accumulated excess starch in leaves and the *bam1 bam3* double mutant had a more severe phenotype than the *bam3* mutant (Fulton et al., 2008).

Both *BAM1* and *BAM3* encode catalytically active, plastid-localized proteins (Kaplan & Guy, 2005; Sparla et al., 2006). These findings show that *BAM3* is a  $\beta$ -amylase isoform that functions predominantly in starch degradation in *A. thaliana* leaves and that *BAM1* also plays an important role in this process, at least in the absence of *BAM3* (Fulton et al., 2008). In contrast, in rice, although two  $\beta$ -amylase isoforms, *OsBAM2* and *OsBAM3*, have been reported to be plastid-targeted active enzymes (Hirano et al., 2011), it is still unclear whether *OsBAM2* and *OsBAM3* are actually involved in starch remobilization in rice leaves.

Our objectives in the present study were to elucidate the function of *OsBAM2* and *OsBAM3* in starch remobilization in rice leaf sheaths during the post-heading stage. We analyzed in detail the changes in *OsBAM2* and *OsBAM3* mRNA levels in leaf sheaths at the pre- and post-heading stages. In addition, we generated *OsBAM2*- and *OsBAM3*-overexpressing plants and *OsBAM2*- and *OsBAM3*-knockdown plants to evaluate the effect of the expression levels of *OsBAM2* and *OsBAM3* in starch remobilization in leaf sheaths, and investigated their phenotypes.

## 2. Materials and methods

### 2.1. Plant materials and sampling

For analyses of carbohydrate content,  $\beta$ -amylase activity and mRNA levels of *OsBAM2* and *OsBAM3*, three rice seedlings (*O. sativa* L., cv. Nipponbare) per pot were transplanted to 1/5000 are Wagner pots (three plants per pot) containing 3.0-kg paddy soil on 5 June, 2014 and were grown outdoors at Meijo University. As a basal fertilizer, chemical fertilizer consisting of N (15%), P (15%), and K (10%) was applied at 2.0 g per pot. Nitrogen fertilizer in the form of ammonium sulfate (0.3-g N per pot) was also applied at the 10th leaf stage. The second leaf sheaths below the flag leaf (the third leaf sheaths) were harvested between 9:30 and 10:30 am at the second leaf emergence

stage, the flag leaf emergence stage, the heading stage, and 5, 10, 15, 20, and 40 days after heading.

Overexpression and knockdown plants of *OsBAM2* and *OsBAM3* (T2 generation) were grown under a photoperiod of 13-h light (845  $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ )/11-h dark at 32/25 °C and approximately 60% relative humidity in a growth chamber. The third leaf sheaths were harvested between 9:30 and 10:30 am at the heading stage and 20 days after heading. All harvested samples were longitudinally divided into two equal parts, immediately frozen in liquid nitrogen, and stored at  $-80$  °C.

## 2.2. Vector construction and transformation of rice

All primers used for vector construction are listed in Table 1. Full-length cDNAs of *OsBAM2* and *OsBAM3* were amplified from cDNA prepared from rice (cv. Nipponbare) leaf blades by polymerase chain reaction (PCR) using PrimeSTAR™ HS DNA Polymerase (TaKaRa) and inserted immediately downstream of the maize *Ubi-1* promoter of pAHC17 (Christensen et al., 1992). The resulting plasmids, pAHC17-BAM2, or pAHC17-BAM3, were digested with *HindIII* and ligated together with a *SacI/HindIII* adapter into the binary vector pBECKS2000.2 (McCormac et al., 1999) digested with *SacI* to generate the constructs for *OsBAM2* and *OsBAM3* overexpression.

The inverted repeat (IR) regions triggering RNA-silencing of *OsBAM2* and *OsBAM3* were amplified by the method described above. The PCR products were cloned into the pENTR/D-TOPO cloning vector (Life Technologies) to generate entry vectors. The RNA-silencing vectors were produced by an LR Clonase reaction (Life Technologies) between each entry vector containing the IR region of *OsBAM2* or *OsBAM3* and the pANDA vector (Miki & Shimamoto, 2004; Miki et al., 2005).

Transgenic rice plants were generated by *Agrobacterium tumefaciens*-mediated transformation of rice (cv. Sasanishiki) calli derived from scutellum (Hiei et al., 1994). Plants (T0 generation) were regenerated from transformed calli selected with hygromycin B and transplanted to plastic pots containing nursery soil. The presence of the transgene was verified by PCR using genomic DNA extracted from the leaf blades of the resistant plants as the template.

## 2.3. Semiquantitative RT-PCR

Total RNA was purified from germinating seeds, leaf blades at the active tillering stage, leaf sheaths at the active tillering stage, leaf sheaths at 10–14 days after heading, and developing caryopses at 10–14 days after heading, following the procedures described by Hirano et al. (2011). The purified total RNA (1  $\mu\text{g}$ ) was reverse-transcribed using

SuperScript™ III Reverse Transcriptase (Life Technologies) with 50 pmol of Oligo(dT)<sub>20</sub> Primer (Life Technologies). After the reaction, the cDNA solution was treated with 2 U of *Escherichia coli* RNase H (Life Technologies) at 37 °C for 20 min. The mRNA levels of  $\beta$ -amylase genes were evaluated by semiquantitative RT-PCR using the gene-specific primers listed in Table 1. An aliquot of cDNA solution corresponding to 20 ng total RNA extracted from the third leaf sheaths was used as the template. The reaction was performed using PrimeSTAR™ HS DNA polymerase (TaKaRa) under the following conditions: 24, 26, or 28 cycles of 10 s at 98 °C, 5 s at 55 °C, and 0.5 min at 72 °C. The amplified fragments were separated by electrophoresis on 1.5% (w/v) agarose gel.

## 2.4. Quantitative RT-PCR


Total RNA purification and cDNA synthesis were performed by the methods described above. The mRNA levels of *OsBAM2* and *OsBAM3* were evaluated by quantitative RT-PCR using the gene-specific primers listed in Table 1. An aliquot of cDNA solution corresponding to 20-ng total RNA extracted from the third leaf sheaths was used as the template. Quantitative RT-PCR was performed using a StepOnePlus Real-Time PCR System (Life Technologies) in a 20- $\mu\text{l}$  reaction mixture containing Power SYBER Green PCR Master Mix (Life Technologies). A serial dilution series of plasmid DNA containing the partial cDNAs of the genes was prepared to construct a standard curve. The mRNA level estimated from the standard curve was standardized to the expression of *OsEF1a1* (RAP-ID: Os03g0177400) (Hirano et al., 2011; Sugimura et al., 2015), which encodes the translation elongation factor 1A.

## 2.5. Determination of starch and sugar contents

Starch, sucrose, and maltose contents in the third leaf sheaths were determined by the method of Sugimura et al. (2015).

## 2.6. Determination of $\beta$ -amylase activity

The activity of  $\beta$ -amylase was determined by the method of Scheidig et al. (2002) with modifications. Frozen leaf sheaths were extracted with 2 ml of ice-cold buffer containing 50 mM MOPS–KOH (pH 7.5), 20 mM  $\text{MgCl}_2$ , 2 mM  $\text{CaCl}_2$ , 1 mM ethylenediaminetetraacetic acid (EDTA), 0.1% (w/v) bovine serum albumin (BSA), and 0.1% (v/v)  $\beta$ -mercaptoethanol and 2% (w/v) polyvinylpyrrolidone. The samples were centrifuged at 20,000 $\times$  *g* for 10 min at 4 °C. For the determination of  $\beta$ -amylase activity, 50  $\mu\text{l}$  of the supernatant was mixed with 200  $\mu\text{l}$  of buffer containing 100 mM sodium acetate (pH 5.4), 2 mM



**Figure 1.** Changes in carbohydrate contents in the third leaf sheaths. The third leaf sheaths were harvested at the second leaf emergence stage, the flag leaf emergence stage, the heading stage, and 5, 10, 15, 20, and 40 days after heading. (a) Starch, (b) sucrose, (c) glucose and fructose, and (d) maltose. Data represent means  $\pm$  SE of three replications.

EDTA, 0.1% (w/v) BSA, and 0.1% (v/v)  $\beta$ -mercaptoethanol. Assays were started by the addition of 50  $\mu$ l of Betamyl-3<sup>o</sup> substrate solution (Megazyme) containing *p*-nitrophenyl- $\beta$ -D-maltotrioxide as a substrate and  $\beta$ -glucosidase as a coupling enzyme at 40  $^{\circ}$ C. After 15 min, the reaction was stopped by the addition of 1.5 ml of 1% (w/v) Trizma base. The absorbance of the reaction solutions was analyzed at 400 nm.

### 3. Results

#### 3.1. Changes in carbohydrate content and $\beta$ -amylase activity in the leaf sheaths


Starch content in the third leaf sheaths peaked at the flag leaf emergence stage and then gradually decreased until 15 days after heading (Figure 1(a)). Beginning at 20 days after heading, starch content rapidly increased until 40 days after heading. The sucrose contents in the third leaf sheaths gradually decreased from the second leaf emergence stage until the heading stage (Figure 1(b)). The sucrose contents rapidly increased from 15 days after heading until 40 days after heading. The glucose and fructose contents exhibited maximum levels at the second leaf emergence stage and gradually decreased until 5 days after heading (Figure 1(c)). The maltose content in the third leaf sheaths remained constant from the flag leaf emergence stage until 20 days after heading (Figure 1(d)).

The  $\beta$ -amylase activity in the third leaf sheaths remained substantially constant throughout the experimental period (Figure 2).

#### 3.2. Expression analysis of $\beta$ -amylase genes

Of the nine genes predicted to encode  $\beta$ -amylase isoforms annotated in the rice genome, *OsBAM1* (RAP-ID: Os07g0543300) and *OsBAM7* (RAP-ID: Os07g0667100) were expressed specifically in germinating seeds and developing caryopses, respectively (Hirano et al., 2011). Accordingly, we initially analyzed the transcript levels of  $\beta$ -amylase genes other than *OsBAM1* and *OsBAM7* in several organs. The transcript levels of *OsBAM2* and *OsBAM3* were highest in leaf sheaths after heading (Figure 3(a)). *OsBAM2* and *OsBAM3* were also highly expressed in leaf blades. *OsBAM4* was clearly expressed in leaf blades and leaf sheaths after heading. *OsBAM5* was expressed in leaf blades, leaf sheaths after heading and developing caryopses. The expression of *OsBAM6* was detected specifically in leaf blades. *OsBAM8* and *OsBAM9* were constitutively expressed in all organs analyzed. *OsBAM9* was strongly expressed in leaf sheaths after heading.

The mRNA levels of *OsBAM2* and *OsBAM3* were analyzed in detail in the leaf sheaths during the pre- and post-heading stages. The mRNA level of *OsBAM2* gradually increased from the flag leaf emergence stage until 5 days after heading (Figure 3(b)). Subsequently, the mRNA levels peaked at 15 days after heading and then decreased following 15 days after heading. The mRNA level of *OsBAM3* decreased from heading stage to 5 days after heading. However, the level rapidly increased starting 10 days after heading, reached a maximum at 15 days after heading. Thereafter, the level decreased markedly from 15 to 20 days after heading.



**Figure 2.** Changes in  $\beta$ -amylase activity in the third leaf sheaths. The third leaf sheaths were harvested at the second leaf emergence stage, the flag leaf emergence stage, the heading stage, and 5, 10, 15 and 20 days after heading. Data represent the means  $\pm$  SE of three replications.

### 3.3. Phenotypic analysis of overexpression and knockdown lines of *OsBAM2* and *OsBAM3*

We generated overexpression lines of *OsBAM2* (*BAM2* OE) and *OsBAM3* (*BAM3* OE) to determine whether the overexpression of *OsBAM2* and *OsBAM3* in fact reduces starch content in rice leaves. The mRNA levels of *OsBAM2* and *OsBAM3* markedly increased in the third leaf sheaths of *BAM2* OE and the *BAM3* OE plants, respectively (Figure 4(a)). The length of *BAM2* OE plants was significantly shorter than that of non-transgenic WT plants (Figure 4(b)). *BAM3* OE plants exhibited a slightly dwarf phenotype. Starch content in the third leaf sheaths was markedly lower in *BAM2* OE and *BAM3* OE plants than in WT plants at the heading stage (Figure 4(c)). The starch content of *BAM3* OE plants was significantly higher than that of *BAM2* OE plants.

Knockdown lines of *OsBAM2* (*BAM2* KD) and *OsBAM3* (*BAM3* KD) were generated by the RNA interference method to evaluate the effect of the repressed expression of *OsBAM2* or *OsBAM3* on the starch content in leaf sheaths. The mRNA level of *OsBAM2* was significantly lower in the third leaf sheaths of *BAM2* KD plants than in those of WT and *BAM3* KD plants (Figure 5(a)). Similarly, the third leaf sheaths of *BAM3* KD plants exhibited drastically lower levels of *OsBAM3* mRNA than those of WT plants and *BAM2* KD plants. Few differences in the growth habits were observed between knockdown and WT plants (Figure 5(b)). No significant differences in panicle weight and yield components among the three plant lines were observed (data not shown). Although  $\beta$ -amylase activities in the third leaf sheaths tended to be lower in *BAM2* KD and *BAM3* KD plants than in WT plants, no significant differences were detected among the three plant


lines (Figure 5(c)). *BAM2* KD and *BAM3* KD plants accumulated starch in the third leaf sheaths at levels similar to those of WT plants at the heading stage (Figure 5(d)). Although, at 20 days after heading, the starch contents of *BAM2* KD and *BAM3* KD plants tended to be slightly higher than that of WT plants, there were no significant differences among the three plant lines.

## 4. Discussion

In rice leaf sheaths, starch is accumulated at a high level prior to the heading stage and remobilized for grain filling during the post-heading stage (Perez et al., 1971). In the present study, starch content in the third leaf sheaths peaked at the flag leaf emergence stage and then gradually decreased (Figure 1(a)), suggesting that starch degradation in the third leaf sheaths was activated to provide carbohydrate to the ripening grains after the flag leaf emergence stage. The starch content was reduced to the lowest levels at 15 days after heading and then rapidly increased from 20 days after heading. These results suggest that starch remobilization was almost complete by 20 days after heading. We accordingly evaluated the  $\beta$ -amylase activity and the mRNA levels of  $\beta$ -amylase genes in the third leaf sheaths until 20 days after heading.

Perez et al. (1971) analyzed  $\beta$ -amylase activity in leaf sheaths harvested at 6, 8, 10, 11 (booting stage), 12 (flowering stage), and 14 weeks after transplanting, but there is little information concerning continuous changes in  $\beta$ -amylase activity in leaf sheaths from the heading to the grain filling stage. In the present study, we analyzed in detail the  $\beta$ -amylase activity in the third leaf sheaths at the pre- and post-heading stages. The activity remained substantially constant from the third leaf emergence stage to 20 days after heading (Figure 2). Thus, no significant correlation between  $\beta$ -amylase activity and the amount of starch loss in the third leaf sheaths during the post-heading stage was detected. In *A. thaliana*, *BAM5*, an active  $\beta$ -amylase protein localized in the cytoplasm, may account for up to 80% (Caspar et al., 1989; Lin et al., 1988) or over 70% (Monroe et al., 2014) of total  $\beta$ -amylase activity in leaves. Thus, analysis using a *bam5* mutant allowed the evaluation of the contribution of each plastid-localized  $\beta$ -amylase isoform to total  $\beta$ -amylase activity (Monroe et al., 2014). In rice, the existence of a cytosolic  $\beta$ -amylase isoform that actively functions in leaves is still undetermined. Identification of a cytoplasm-localized  $\beta$ -amylase protein and analysis of its activity may be needed to evaluate the correlations between changes in starch content and  $\beta$ -amylase activity in the leaf sheaths.

In the rice genome, there are nine genes predicted to encode  $\beta$ -amylase-like proteins (Hirano et al., 2011; Saika et al., 2005). Of these genes, at least *OsBAM2* and *OsBAM3*




**Figure 3.** Expression analyses of  $\beta$ -amylase genes. (a) Semiquantitative RT-PCR of *OsBAM2*, *OsBAM3*, *OsBAM4*, *OsBAM5*, *OsBAM6*, *OsBAM8*, and *OsBAM9* in various rice organs. Leaf blade and leaf sheath before heading were harvested at the active tillering stage, and leaf sheath after heading and developing caryopsis were harvested at 10–14 days after heading. *OsEF1a1* encoding translation elongation factor 1A was used as the internal standard. (b) Changes in transcript levels of *OsBAM2* and *OsBAM3* in the third leaf sheaths by quantitative RT-PCR. The third leaf sheaths were harvested at the second leaf emergence stage, the flag leaf emergence stage, the heading stage, and 5, 10, 15 and 20 days after heading. Values were standardized to the transcript level of *OsEF1a1* and represent means  $\pm$  SE of three replications.

encode plastid-targeted active isoforms (Hirano et al., 2011). The finding that the overexpression of *OsBAM2* or *OsBAM3* markedly reduced starch accumulation in the third leaf sheaths at the heading stage (Figure 4(c)) shows that *OsBAM2* and *OsBAM3* can in fact function in the degradation of starch stored transiently in plastid *in vivo*. The degree of starch loss was more pronounced in *BAM2* OE than in *BAM3* OE plants (Figure 4(c)). This phenomenon may result from the differences in expression

levels between the overexpressed *OsBAM2* and *OsBAM3* or the kinetic properties between *OsBAM2* and *OsBAM3* isoforms. A rice mutant lacking *OsAGPL1* encoding a large subunit of ADP-glucose pyrophosphorylase, which had markedly reduced starch content in the stem, displayed the shorter plant length than WT plants (Okamura et al., 2013). In the present study, the plant length of *BAM2* OE, which accumulated much less starch in the leaf sheaths, was significantly shorter than that of WT plants, similarly to the mutant defective in *OsAGPL1* (Figure 4(b)). These results suggest that the repression of starch accumulation in the leaf sheaths affects the plant growth via the abnormal supply of carbohydrate.

The mRNA levels of *OsBAM2* and *OsBAM3* were higher in leaf sheaths after heading than in those before heading and in leaf blades (Figure 3(a)). In addition, the mRNA levels of both *OsBAM2* and *OsBAM3* increased from the flag leaf emergence stage to the heading stage when the starch content began to decrease in the third leaf sheaths (Figure 3(b)). These results raise the possibility that *OsBAM2* and *OsBAM3* play a role in starch degradation in the leaf sheaths. However, the mRNA level of *OsBAM2* decreased from 5 to 10 days after heading and that of *OsBAM3* rapidly decreased from the heading to 5 days after heading. Subsequently, the mRNA levels of both *OsBAM2* and *OsBAM3* reached a maximum at 15 days after heading. Chen and Wang (2008) showed that the  $\beta$ -amylase gene (cDNA accession number: AK068968, designated as *OsBAM3* in this study) was highly expressed in leaf sheaths coincidentally with a decrease in starch content during the post-heading stage. However, our results indicate that the mRNA levels of *OsBAM2* and *OsBAM3* did not always remain high from the heading stage to 15 days after heading, when the starch content gradually decreased in the third leaf sheaths. In *A. thaliana*, expression of *BAM5* is induced by sugar (Mita et al., 1995). Similarly in rice leaves, it is possible that the expression of *OsBAM2* and *OsBAM3* may be tightly regulated in response to changes in carbon status in cells. Such a mechanism may at least in part cause the drastic changes in the mRNA levels of *OsBAM2* and *OsBAM3* in the third leaf sheaths during the post-heading stage.


Although the starch contents in the third leaf sheaths tended to be higher in both *BAM2* KD and *BAM3* KD than in WT plants at 20 days after heading, no significant differences were detected among their starch contents (Figure 5(d)). These results indicate that the knockdown of the individual genes, *OsBAM2* or *OsBAM3*, did not lead to the hyperaccumulation of starch in the leaf sheaths. In fact, the  $\beta$ -amylase activities in the third leaf sheaths were not significantly lower in *BAM2* KD and *BAM3* KD than in WT plants at the heading stage (Figure 5(c)). In *A. thaliana*, a *bam3* mutant deficient for *BAM3*, one of the  $\beta$ -amylase



**Figure 4.** Phenotypic analyses of overexpression plants of *OsBAM2* and *OsBAM3*. (a) Semiquantitative RT-PCR of *OsBAM2* and *OsBAM3* in the third leaf sheaths at the heading stage. *OsEF1a1* was used as internal standard. (b) Plant lengths at the heading stage. Data represent means  $\pm$  SE of five replications. Data followed by different letters represent significant differences at the 5% level by Tukey's test. (c) Starch contents in the third leaf sheaths at the heading stage. Data represent means  $\pm$  SE of three replications. Data followed by different letters represent significant differences at the 5% level by Tukey's test.

isoforms targeted to plastids, accumulated an excess of starch in leaves, whereas the *bam1* mutant deficient for plastid-targeted BAM1 showed normal accumulation of starch (Fulton et al., 2008). Monroe et al. (2014) reported that only the *bam3* mutant among the single  $\beta$ -amylase mutants (*bam1*, *bam2*, *bam3*, and *bam6*) accumulated an excess of starch in leaves of five-week-old plants, but in eight-week-old plants all mutants displayed a starch-excess phenotype. Moreover, the *bam1 bam3* (Fulton et al., 2008), the *bam2 bam3*, and the *bam3 bam6* double mutants (Monroe et al., 2014) displayed a more extreme phenotype for leaf starch accumulation than the *bam3* mutant. Phenotypic analysis of double knockdown of *OsBAM2* and *OsBAM3* may reveal the contribution of these genes to starch degradation in leaf sheaths.

Asatsuma et al. (2005) showed that  $\alpha$ -amylase I-1 encoded by *RAmy1A* plays a role in starch degradation in rice leaves. The expression level of *RAmy2A*, encoding



**Figure 5.** Phenotypic analyses of knockdown plants of *OsBAM2* and *OsBAM3*. (a) Quantitative RT-PCR of *OsBAM2* and *OsBAM3* in the third leaf sheaths at the heading stage. *OsEF1a1* was used as the internal standard. Data are relative to the transcript level of wild-type (WT) plants and represent means  $\pm$  SE of three replications. Data followed by different letters represent significant difference at the 5% level by Tukey's test. (b) Photographs of knockdown plants of *OsBAM2* and *OsBAM3* at booting stage. (c)  $\beta$ -Amylase activity in the third leaf sheaths at the heading stage. Data represent means  $\pm$  SE of three replications. n.s., no significant difference at the 5% level by Tukey's test. (d) Starch contents in the third leaf sheaths at the heading and 20 days after heading. Data represent means  $\pm$  SE of three replications. n.s., no significant difference at the 5% level by Tukey's test.



one of the  $\alpha$ -amylase isoforms, increased significantly in rice leaf sheaths at the post-heading stage (Chen & Wang, 2008; Sugimura et al., 2015). The disruption of *OsGWD1* encoding  $\alpha$ -glucan water dikinase displayed an excess-starch phenotype in rice leaves (Hirose et al., 2013). In the present study, *OsBAM4*, *OsBAM5*, *OsBAM8*, and *OsBAM9* were distinctly expressed in leaf sheaths after heading (Figure 3(a)), raising the possibility that these  $\beta$ -amylase genes may contribute to starch degradation in leaf sheaths during the post-heading stage. These results suggest the involvement of several other genes as well as *OsBAM2* and *OsBAM3* in starch degradation in leaf sheaths, perhaps accounting for the absence of hyperaccumulation of starch in the leaf sheaths of *BAM2* KD and *BAM3* KD.

## Acknowledgments

We thank Dr. Ko Shimamoto (Nara Institute of Science and Technology, Japan) for providing the pANDA vector.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported in part by JSPS KAKENHI [grant number 24580027] and [grant number 26292009].

## References

- Asatsuma, S., Sawada, C., Itoh, K., Okito, M., Kitajima, A., & Mitsui, T. (2005). Involvement of  $\alpha$ -amylase I-1 in starch degradation in rice chloroplasts. *Plant and Cell Physiology*, *46*, 858–869.
- Baunsgaard, L., Lütken, H., Mikkelsen, R., Glaring, M. A., Pham, T. T., & Blennow, A. (2005). A novel isoform of glucan, water dikinase phosphorylates pre-phosphorylated  $\alpha$ -glucans and is involved in starch degradation in *Arabidopsis*. *The Plant Journal*, *41*, 595–605.
- Caspar, T., Lin, T. -P., Monroe, J., Bernhard, W., Spilatro, S., Preiss, J., & Somerville, C. (1989). Altered regulation of  $\beta$ -amylase activity in mutants of *Arabidopsis* with lesions in starch metabolism. *Proceedings of the National Academy of Sciences USA*, *86*, 5830–5833.
- Chen, H.-J., & Wang, S.-J. (2008). Molecular regulation of sink-source transition in rice leaf sheaths during the heading period. *Acta Physiologiae Plantarum*, *30*, 639–649.
- Chia, T., Thorneycroft, D., Chapple, A., Messerli, G., Chen, J., Zeeman, S. C., ... Smith, A. M. (2004). A cytosolic glucosyltransferase is required for conversion of starch to sucrose in *Arabidopsis* leaves at night. *The Plant Journal*, *37*, 853–863.
- Christensen, A. H., Sharrock, R. A., & Quail, P. H. (1992). Maize polyubiquitin genes: Structure, thermal perturbation of expression and transcript splicing, and promoter activity following transfer to protoplasts by electroporation. *Plant Molecular Biology*, *18*, 675–689.
- Cock, J. H., & Yoshida, S. (1972). Accumulation of <sup>14</sup>C-labelled carbohydrate before flowering and its subsequent redistribution and respiration in the rice plant. *Japanese Journal of Crop Science*, *41*, 226–234.
- Fulton, D. C., Stettler, M., Mettler, T., Vaughan, C. K., Li, J., Francisco, P., ... Zeeman, S. C. (2008).  $\beta$ -amylase4, a noncatalytic protein required for starch breakdown, acts upstream of three active  $\beta$ -amylases in *Arabidopsis* chloroplasts. *The Plant Cell*, *20*, 1040–1058.
- He, H. Y., Koike, M., Ishimaru, T., Ohsugi, R., & Yamagishi, T. (2005). Temporal and spatial variations of carbohydrate content in rice leaf sheath and their varietal differences. *Plant Production Science*, *8*, 546–552.
- Hiei, Y., Ohta, S., Komari, T., & Kumashiro, T. (1994). Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *The Plant Journal*, *6*, 271–282.
- Hirano, T., Saito, Y., Ushimaru, H., & Michiyama, H. (2005). The effect of the amount of nitrogen fertilizer on starch metabolism in leaf sheath of japonica and indica rice varieties during the heading period. *Plant Production Science*, *8*, 122–130.
- Hirano, T., Takahashi, Y., Fukayama, H., & Michiyama, H. (2011). Identification of two plastid-targeted  $\beta$ -amylases in rice. *Plant Production Science*, *14*, 318–324.
- Hirose, T., Aoki, N., Harada, Y., Okumura, M., Hashida, Y., Ohsugi, R., ... Terao, T. (2013). Disruption of a rice gene for  $\alpha$ -glucan water dikinase, *OsGWD1*, leads to hyperaccumulation of starch in leaves but exhibits limited effects on growth. *Frontiers in Plant Science*, *4*, 147.
- Hirose, T., Endler, A., & Ohsugi, R. (1999). Gene expression of enzymes for starch and sucrose metabolism and transport in leaf sheaths of rice (*Oryza sativa* L.) during the heading period in relation to the sink to source transition. *Plant Production Science*, *2*, 178–183.
- Hirose, T., Ohdan, T., Nakamura, Y., & Terao, T. (2006). Expression profiling of genes related to starch synthesis in rice leaf sheaths during the heading period. *Physiologia Plantarum*, *128*, 425–435.
- Ishimaru, K., Kosone, M., Sasaki, H., & Kashiwagi, T. (2004). Leaf contents differ depending on the position in a rice leaf sheath during sink-source transition. *Plant Physiology and Biochemistry*, *42*, 855–860.
- Kaplan, F., & Guy, C. L. (2005). RNA interference of *Arabidopsis* beta-amylase8 prevents maltose accumulation upon cold shock and increases sensitivity of PSII photochemical efficiency to freezing stress. *The Plant Journal*, *44*, 730–743.
- Kötting, O., Pusch, K., Tiessen, A., Geigenberger, P., Steup, M., & Ritte, G. (2005). Identification of a novel enzyme required for starch metabolism in *Arabidopsis* leaves. The phosphoglucan, water dikinase. *Plant Physiology*, *137*, 242–252.
- Lao, N. T., Schoneveld, O., Mould, R. M., Hibberd, J. M., Gray, J. C., & Kavanagh, T. A. (1999). An *Arabidopsis* gene encoding a chloroplast-targeted  $\beta$ -amylase. *The Plant Journal*, *20*, 519–527.
- Lin, T. -P., Spilatro, S. R., & Preiss, J. (1988). Subcellular localization and characterization of amylases in *Arabidopsis* leaf. *Plant Physiology*, *86*, 251–259.
- McCormac, A. C., Elliott, M. C., & Chen, D.-F. (1999). pBECK2000: A novel plasmid series for the facile creation of complex binary vectors, which incorporates 'clean-gene' facilities. *Molecular and General Genetics*, *261*, 226–235.

- Miki, D., Itoh, R., & Shimamoto, K. (2005). RNA silencing of single and multiple members in a gene family of rice. *Plant Physiology*, *138*, 1903–1913.
- Miki, D., & Shimamoto, K. (2004). Simple RNAi vectors for stable and transient suppression of gene function in rice. *Plant and Cell Physiology*, *45*, 490–495.
- Mikkelsen, R., Baunsgaard, L., & Blennow, A. (2004). Functional characterization of  $\alpha$ -glucan, water dikinase, the starch phosphorylating enzyme. *Biochemical Journal*, *377*, 525–532.
- Mita, S., Suzuki-Fujii, K., & Nakamura, K. (1995). Sugar-inducible expression of a gene for  $\beta$ -amylase in *Arabidopsis thaliana*. *Plant Physiology*, *107*, 895–904.
- Monroe, J. D., Storm, A. R., Badley, E. M., Lehman, M. D., Platt, S. M., Saunders, L. K., ... Torres, C. E. (2014).  $\beta$ -amylase1 and  $\beta$ -amylase3 are plastidic starch hydrolases in *Arabidopsis* that seem to be adapted for different thermal, pH, and stress conditions. *Plant Physiology*, *166*, 1748–1763.
- Niittylä, T., Messerli, G., Trevisan, M., Chen, J., Smith, A. M., & Zeeman, S. C. (2004). A previously unknown maltose transporter essential for starch degradation in leaves. *Science*, *303*, 87–89.
- Okamura, M., Hirose, T., Hashida, Y., Yamagishi, T., Ohsugi, R., & Aoki, N. (2013). Starch reduction in rice stems due to a lack of *OsAGPL1* or *OsAPL3* decreases grain yield under low irradiance during ripening and modifies plant architecture. *Functional Plant Biology*, *40*, 1137–1146.
- Perez, C. M., Palmiano, E. P., Baun, L. C., & Juliano, B. O. (1971). Starch metabolism in the leaf sheaths and culm of rice. *Plant Physiology*, *47*, 404–408.
- Ritte, G., Lloyd, J. R., Eckermann, N., Rottmann, A., Kossmann, J., & Steup, M. (2002). The starch-related R1 protein is an  $\alpha$ -glucan, water dikinase. *Proceedings of the National Academy of Sciences USA*, *99*, 7166–7171.
- Ritte, G., Scharf, A., Eckermann, N., Haebel, S., & Steup, M. (2004). Phosphorylation of transitory starch is increased during degradation. *Plant Physiology*, *135*, 2068–2077.
- Saika, H., Nakazono, M., Ikeda, A., Yamaguchi, J., Masaki, S., Kanekatsu, M., & Nemoto, K. (2005). A transposon-induced spontaneous mutation results in low  $\beta$ -amylase content in rice. *Plant Science*, *169*, 239–244.
- Scheidig, A., Fröhlich, A., Schulze, S., Lloyd, J. R., & Kossmann, J. (2002). Downregulation of a chloroplast-targeted  $\beta$ -amylase leads to a starch-excess phenotype in leaves. *The Plant Journal*, *30*, 581–591.
- Sparla, F., Costa, A., Schiavo, F. L., Pupillo, P., & Trost, P. (2006). Redox regulation of a novel plastid-targeted  $\beta$ -amylase of *Arabidopsis*. *Plant Physiology*, *141*, 840–850.
- Sugimura, Y., Michiyama, H., & Hirano, T. (2015). Involvement of  $\alpha$ -amylase genes in starch degradation in rice leaf sheaths at the post-heading stage. *Plant Production Science*, *18*, 277–283.
- Watanabe, Y., Nakamura, Y., & Ishii, R. (1997). Relationship between starch accumulation and activities of the related enzymes in the leaf sheath as a temporary sink organ in rice (*Oryza sativa*). *Australian Journal of Plant Physiology*, *24*, 563–569.
- Weise, S. E., Weber, A. P. M., & Sharkey, T. D. (2004). Maltose is the major form of carbon exported from the chloroplast at night. *Planta*, *218*, 474–482.
- Yu, T. S., Kofler, H., Häusler, R. E., Hille, D., Flügge, U. I., Zeeman, S. C., ... Weber, A. (2001). The *Arabidopsis* *sex1* mutant is defective in the R1 protein, a general regulator of starch degradation in plants, and not in the chloroplast hexose transporter. *The Plant Cell*, *13*, 1907–1918.
- Zeeman, S. C., Kossmann, J., & Smith, A. M. (2010). Starch: Its metabolism, evolution, and biotechnological modification in plants. *Annual Review of Plant Biology*, *61*, 209–234.