



SUBSTANCE USE & MISUSE  
An International Interdisciplinary Forum

## Substance Use & Misuse

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/ism20>

# The Role of Alcohol-Related Proactive Dietary Restriction to Feel the Psychoactive Effects of Alcohol Faster on Binge Drinking Frequency among Freshmen College Students

Gail Castañeda, Sarah E. Colby, Melissa D. Olfert, Tracey E. Barnett, Wenjun Zhou, Megan D. Engel & Anne E. Mathews

To cite this article: Gail Castañeda, Sarah E. Colby, Melissa D. Olfert, Tracey E. Barnett, Wenjun Zhou, Megan D. Engel & Anne E. Mathews (2021): The Role of Alcohol-Related Proactive Dietary Restriction to Feel the Psychoactive Effects of Alcohol Faster on Binge Drinking Frequency among Freshmen College Students, Substance Use & Misuse, DOI: [10.1080/10826084.2021.1914104](https://doi.org/10.1080/10826084.2021.1914104)

To link to this article: <https://doi.org/10.1080/10826084.2021.1914104>



© 2021 The Author(s). Published with license by Taylor and Francis Group, LLC



Published online: 19 May 2021.



Submit your article to this journal [↗](#)






View related articles [↗](#)



View Crossmark data [↗](#)

# The Role of Alcohol-Related Proactive Dietary Restriction to Feel the Psychoactive Effects of Alcohol Faster on Binge Drinking Frequency among Freshmen College Students

Gail Castañeda<sup>a</sup>, Sarah E. Colby<sup>b</sup>, Melissa D. Olfert<sup>c</sup>, Tracey E. Barnett<sup>d</sup> , Wenjun Zhou<sup>e</sup> , Megan D. Engelf<sup>f</sup> and Anne E. Mathews<sup>f</sup> 

<sup>a</sup>Department of Occupational Therapy, College of Public Health and Health Professions, University of Florida, Gainesville, Florida, USA;

<sup>b</sup>Department of Nutrition, College of Education, University of Tennessee, Knoxville, Tennessee, USA; <sup>c</sup>Human Nutrition and Foods, West Virginia University, Morgantown, West Virginia, USA; <sup>d</sup>School of Public Health, University of North Texas Health Science Center, Fort Worth, Texas, USA; <sup>e</sup>Business Analytics and Statistics, University of Tennessee, Knoxville, Tennessee, USA; <sup>f</sup>Food Science and Human Nutrition Department, University of Florida, Gainesville, Florida, USA

## ABSTRACT

**Background:** There is a positive cross-sectional relationship between alcohol-related proactive dietary restriction to feel the effects of alcohol faster (APDR) and binge drinking, a health and safety issue impacting college students. **Objective:** To examine: 1) the longitudinal predictive ability of varying levels of APDR on binge drinking frequency; and 1a) the strength of the relationship between varying levels of APDR and binge drinking frequency during freshman year of college ( $n=1,149$ ). **Methods:** Ordinal logistic regression was used to model the relationship between APDR and binge drinking frequency. **Results:** Main findings suggest APDR of students who reported eating less than usual (low APDR) prior to drinking to feel the effects of alcohol faster was a significant predictor of binge drinking frequency (1.27 (95% CI, 0.06 to 0.42), Wald  $\chi^2$  (1) = 8.46,  $p=.009$ ) at baseline, but not at 7-month follow-up (1.02 (95% CI, -0.18 to 0.23), Wald  $\chi^2$  (1) = .51,  $p=.83$ ). APDR for students who reported skipping one or more meals (high APDR) to feel the effects of alcohol faster was not a significant predictor of binge drinking frequency at baseline nor at 7-month follow-up. **Conclusion:** Low APDR is a significant predictor of binge drinking frequency that is established early in the first semester of college with no significant change occurring in binge drinking frequency over the course of students' freshman year at 7-month follow-up. Campus health professionals are urged to emphasize the detrimental health effects of low APDR early in the first semester of college.

## KEYWORDS

College binge drinking; alcohol-related proactive dietary restriction; disordered eating

## Introduction

The weight conscious drinking phenomenon describes the occurrence of disordered eating behaviors meant to compensate for alcohol-related energy intake and/or to augment the psychoactive effects of alcohol (Barry & Piazza-Gardner, 2012; Barry et al., 2013; Peralta, 2002). Weight conscious drinking also accounts for the temporal relation of compensatory behaviors and/or behaviors that enhance the psychoactive effects of alcohol in response to drinking episodes, commonly referred to as 'proactive' and 'reactive' compensatory behaviors (Bryant et al., 2012; Castañeda et al., 2020; Peralta, 2002; Piazza-Gardner & Barry, 2014). Proactive disordered eating behaviors can include self-induced vomiting or dietary restraint *prior to* a planned drinking episode, whereas reactive disordered eating behaviors can include self-induced vomiting or dietary restraint *after* a drinking episode to compensate for the energy intake from alcohol.

While weight conscious drinking embodies a range of hazardous compensatory behaviors in response to the alcohol-related calories consumed, this phenomenon also takes place across varying levels of alcohol consumption. Of special concern for college students is the pairing of weight conscious drinking behaviors with binge drinking episodes (Burke et al., 2010; Kelly-Weeder, 2011). Binge drinking, also known as heavy episodic alcohol consumption, has been empirically identified as a problematic drinking pattern, particularly among the young adult (18-24) college student population (Nelson et al., 2009; Wechsler et al., 2002). In a previous assessment of college students, 42% self-reported engaging in binge drinking behaviors within the past two weeks (Wechsler et al., 2002). Binge drinking is associated with a wide range of health and social problems, including sexually transmitted diseases, unintended pregnancies, accidental injuries, and violent crimes (R. Hingson et al., 2005; Wechsler et al.,

2002; White & Hingson, 2013). Furthermore, while college students have been found to drink less frequently than their non-student counterparts, they drink more heavily at higher frequencies, such that rates of binge drinking are much higher in college student samples, as opposed to non-student samples (R. W. Hingson et al., 2002).

First year college students are at-risk for hazardous drinking (Krieger et al., 2018), and those that endorse weight conscious drinking behaviors have increased odds of binge drinking. (Barry & Piazza-Gardner, 2012; Barry et al., 2013). Previous cross-sectional investigations indicate that college students who intentionally restrict their caloric intake prior to drinking episodes are then more likely to binge drink (Bryant et al., 2012; Burke et al., 2010; Castañeda et al., 2020; Eisenberg & Fitz, 2014). For instance, Burke et al. (2010) found that while 33% of college students who reported engaging in proactive dietary restriction also reported alcohol consumption on 10 to 19 days within the past month, 21% of college students engaging in proactive dietary restriction also reported binge drinking on over 20 days within the past month. That is, students who report engaging in both proactive dietary restriction and binge drinking, drink heavily more frequently. In another study, findings from Castañeda et al. (2020) indicated a strong positive cross-sectional relationship between proactive Alcohol Effects (i.e. behaviors, such as proactive dietary restriction, aimed at enhancing the psychoactive effects of alcohol prior to a drinking episode) and binge drinking frequency among college students.

In a study assessing the range of alcohol-related compensatory behaviors employed by college students, a preliminary set of 34-items were generated based on salient themes identified in Peralta (2002)'s qualitative investigation on weight conscious drinking (Bryant et al., 2012). Findings from Bryant et al. (2012) revealed college binge drinkers scored significantly higher than non-binge drinkers on two weight conscious drinking items assessing alcohol-related proactive dietary restraint to feel the psychoactive effects of alcohol faster. Bryant et al. (2012)'s examination utilized the following two levels of APDR: 1) eating less than usual during one or more meals prior to drinking to feel the effects of alcohol faster; and 2) skipping one or more meals prior to drinking to feel the effects of alcohol faster. While the investigators did not report any comparison between the strength of the relationship between each APDR level and student binge drinking, both items were linked to binge drinking.

The detrimental health effects of alcohol consumption by college students are compounded when binge drinking is paired with proactive dietary restriction intended to feel the psychoactive effects of alcohol faster. Dietary restriction prior to a binge drinking episode is known to increase the the absorption rate of alcohol, blood alcohol concentration, and risk for negative alcohol-related consequences (Cederbaum, 2012; Giles et al., 2009). Furthermore, alcohol-related proactive dietary restraint paired with binge drinking places students at risk for primary (or secondary)

malnutrition due to the body's preferential oxidation of alcohol over other nutrients (Lieber, 2003).

Examining the strength of the relationships between varying levels of alcohol-related proactive dietary restriction and binge drinking frequency can serve to identify APDR as a behavioral marker for college alcohol health prevention screenings and intervention programming. In addition, the investigation of the longitudinal relationship between proactive dietary restraint (aimed at feeling the psychoactive effects of alcohol faster) and binge drinking frequency among nonclinical college freshmen could further inform campus alcohol health prevention and intervention programming by aiding college health professionals in determining the most appropriate programming timeline. To date, no studies have specifically assessed: 1) the strength of the relationships between varying levels of alcohol-related proactive dietary restriction and binge drinking frequency among non-clinical freshman college students; and 2) the longitudinal relationship between varying levels of proactive dietary restraint aimed at feeling the psychoactive effects of alcohol faster and binge drinking frequency among non-clinical freshman college students.

Given a growing body of cross-sectional research has corroborated the relationship between proactive dietary restriction to feel the psychoactive effects of alcohol faster and college student binge drinking, this study specifically assessed the predictive ability of varying levels of alcohol-related proactive dietary restriction to feel the psychoactive effects of alcohol faster (APDR) to contribute to binge drinking frequency at 7-month follow-up among a freshman college cohort. It was hypothesized that: 1) both low and high baseline APDR would remain predictors of binge drinking frequency at 7-month follow-up; and 1a) high baseline APDR would be more strongly associated with binge drinking frequency at 7-month follow-up than low APDR. For the purposes of this paper, low APDR has been defined as "eating less than usual during one or more meals prior to drinking to feel the effects of alcohol faster" and high APDR has been defined as "skipping one or more meals prior to drinking to feel the effects of alcohol faster".

## Method

### *Participants and procedure*

Data for this longitudinal study were obtained from a cohort sample of 1,149 college freshman students at eight U.S. universities during the development phase of 'GetFRUVED', a cohort study to encourage healthful eating habits and increased physical activity among college students. Participating universities are all members of an established multi-state research group (NC1193) and include University of Tennessee, University of Florida, West Virginia University Syracuse University, University of Maine, South Dakota State University, Kansas State University, and Auburn University. APDR was assessed as part of the study's web-based questionnaire covering freshmen student perceptions and

behaviors on a wide range of health-related topics. Participants completed survey questions and participated in anthropometric assessment by trained research assistants at the beginning of the fall 2015 and end of spring 2016 semesters onsite at each campus. The GetFRUVED Study was registered at Clinicaltrials.gov with the identifier code NCT02941497.

### **Inclusion criteria**

Eligibility for the study, previously described in detail by Loso et al. (2018), was defined as individuals age 18 years or older, college enrollment as a freshman student, and low (less than two cups of fruits and/or less than three cups of vegetables) average daily fruit and/or vegetable consumption. Additionally, the participants had to meet at least one of the following criteria: (a) BMI  $\geq 25$  kg/m<sup>2</sup>; (b) self-identified as first-generation college student; (c) student-identified overweight/obese parent; (d) low income background; and (e) self-identified racial minority. All participants provided informed consent and the University of Tennessee Institutional Review Board approved the study for all participating universities, except the University of Florida, Auburn University, and Kansas State University where separate institutional review board approval was obtained.

### **Measures**

Participants completed survey questions onsite via Qualtrics survey software. Survey items included correlates for binge drinking frequency (i.e. geographic region, place of residence, Greek campus organization affiliation, athlete status), APDR to feel the psychoactive effects of alcohol faster, fruit and vegetable intake, eating disorder risk, physical activity, low family affluence, parents overweight or obese, and demographic characteristics, such as sex, race, and ethnic origin. Anthropometric data collected included participant height, body weight, waist, hip, and neck circumferences.

*Alcohol-related proactive dietary restraint.* Two items measuring alcohol-related proactive dietary restraint were extracted from the Compensatory Eating Behaviors in Response to Alcohol Consumption Scale (CEBRACS), a 21-item measure assessing the co-occurrence of disordered eating behaviors, such as dietary restraint, purging, and exercise to compensate for the intake of alcohol-related calories and/or enhance the psychoactive effects of alcohol (Rahal et al., 2012). One item read: "I have eaten less than usual during one or more meals before drinking to feel the effects of alcohol faster". The other item read: "I have skipped one or more meals before drinking to feel the effects of alcohol faster". Both items were rated on a 5-point Likert scale based on behavior frequency: 1 = never; 2 = rarely (approximately 25% of occasions); 3 = sometimes (approximately 50% of occasions); 4 = often (approximately 75% of occasions); 5 = nearly always, with the option to indicate

"choose not to answer". Participant responses were based on alcohol consumed within the past three months.

*Binge drinking frequency.* One item measuring binge drinking frequency was extracted from the Alcohol Use Disorders Identification Test (AUDIT-C), a three-item short form assessing risk for alcohol abuse. The item read: "How often did you have six or more drinks on one occasion in the past year?" Item response options included: 1 = Never; 2 = Less than monthly; 3 = Monthly; 4 = Weekly; 5 = Daily or almost daily, with the option to indicate "choose not to answer". Use of the AUDIT-C's binge frequency item has demonstrated adequate clinical diagnostic accuracy (i.e. area under the receiver operator characteristic curve (AUROC) of 0.8 or greater) for detecting heavy drinking (0.827) and active alcohol abuse and/or alcohol dependence (0.831) (Bush et al., 1998).

*The Eating Attitudes Test (EAT-26).* The EAT-26 is a screening tool commonly used across college student samples to assess eating disorder risk (Garner et al., 1982). Items capture both cognitive and behavioral symptomology related to eating disorders. Total scores of 20 or more are indicative of elevated risk for an eating disorder. The scale also yields a referral index based on the following three criteria: 1) total score; 2) responses to behavioral questions related to eating symptoms and weight loss (e.g. I have gone on eating binges where I feel that I may not be able to stop); and 3) the respondent's BMI. With the current data, Cronbach's alpha coefficients (i.e. internal consistency) for the scale's three subdomains were 0.86 (Dieting), 0.81 (Bulimia and Food Preoccupation), and 0.58 (Oral Control).

*NCI Fruit and Vegetable All-Day screener.* The NCI fruit and vegetable all-day screener (NCIfv) is a 19-item screening tool that estimates median intake of fruit and vegetable servings by obtaining the frequency of participants' fruit and vegetable consumption within the past month, week or day and corresponding serving sizes (Thompson et al., 2002). Frequencies and servings for all the following forms of fruit and vegetable consumption are requested: raw, cooked, snacks, home meals, restaurant meals/take-out, and combinations with other foods. Total number of daily fruit and vegetable servings are computed as the sum of the product of number of times per day and Pyramid/MyPyramid servings for each fruit or vegetable. Our data yielded a Cronbach's alpha coefficient of 0.71 across all 19 items.

*The International Physical Activity Questionnaire Short Form (IPAQ).* The International Physical Activity Questionnaire provides information with respect to health-related physical activity within the past week (Bassett, 2003). The 6-item short form collects data on the frequency and duration of vigorous, moderate, and walking activities performed within the last 7 days. Based on minutes spent on a given physical activity per week (MET) and physical activity duration, average MET scores have been established by the IPAQ Research Committee for each physical activity category (i.e. vigorous, moderate, and low). Internal consistency across all six items rendered an alpha coefficient of 0.59 for the data under study.

**Anthropometric measures.** Physical assessments were conducted at each time point by trained researchers to collect participants' waist, hip, and neck circumference. Measurements were recorded twice and repeated a third time if the difference between the first two measures was greater than 0.5 centimeters, and then the average between the two measures was recorded. Participant height (meters) and weight (kilograms) measurements were taken with calibrated scales and full-length stadiometers at each time point. BMI and waist hip ratio were calculated using measurements recorded during physical assessments. BMI was computed by dividing participants' weight in kilograms by their height in meters squared and waist hip ratio was calculated by dividing participants' waist circumference in centimeters by their hip circumference in centimeters.

**Demographic Characteristics.** Participants reported their age, sex, race, ethnicity (hispanic or non-hispanic), geographic region, athlete status (yes/no), place of residence (on- or off-campus), Greek campus organization affiliation (sorority or fraternity), family affluence, and parent overweight or obese status (yes/no/not sure). Given low race category percentages, only White, Black, and Asian race categories were analyzed. Place of residence was assessed *via* the following question: "Where do you live?" Six response options were provided (Campus residence hall, sorority or fraternity, other college housing, off-campus housing, parent or guardian's home or choose not to answer). Due to low percentages of various off campus response options, place of residence was dichotomized as on campus versus off campus for analyses. Greek campus organization affiliation was derived from respondents who reported living in a sorority or fraternity home. Family affluence was measured with the following four items: "Do you have your own bedroom (for just you, or you and your partner/spouse)?", "How many computers/laptops are in your home?", "How many cars, vans, or trucks does your family own?", and "How many times did you travel away on vacation with your family during the past 12 months?" (Boyce et al., 2006). Bedroom ownership was dichotomized into a yes or no response and the remainder of the items consisted of response options denoting frequencies. Points were assigned per response with a yes being coded as 1 and remaining frequency options paralleled numerical options. A summated score across all four items totaling two points or less was categorized as low affluence.

## Analysis

Descriptive statistics were used to characterize study participants at baseline. An ordinal logistic regression using a complementary log-log link function was conducted to examine the predictive ability of varying levels of APDR to contribute to binge drinking frequency at baseline and 7-month follow up. A dependent sample t-test was conducted to compare mean differences across binge drinking frequency at baseline and 7-month follow up.

## Missing data

Missing data ranged from 0-10% (Bennett, 2001) across baseline demographic predictors for the ordinal regression model. However, missing values were over 30% for APDR among students who reported eating less than usual before drinking to feel the effects of alcohol faster (35.2%), skipping one or more meals before drinking to feel the effects of alcohol faster (35.4%), binge drinking frequency at baseline (35.8%), and binge drinking frequency at 7-month follow up (46.2%). A significant Little's missing completely at random (MCAR) test indicated missing values were not missing completely at random ( $\chi^2 [57] = 142.09, p \leq .001$ ). The missing data mechanism was more closely assessed *via* the use of missing indicators for each of the following items: eating less than usual during one or more meals before drinking to feel the effects of alcohol faster, skipping one or more meals before drinking to feel the effects of alcohol faster, binge drinking frequency at baseline, and binge drinking frequency at follow-up. Missing value indicators were coded as 1 for original missing variables and 0 for data present (Enders, 2010).

A comparison of the ordinal regression model's complete case analysis against two non-parametric multiple imputation methods (missForest with and without missing indicators) for mixed-type variables is provided. Fifty missForest imputations were conducted for improved model validity (Stekhoven & Bühlmann, 2012). Results for the 50 missForest multiple imputation results containing missing data indicators are provided in Tables 2–4. Statistical tests were computed using SPSS statistical software v26 and imputation procedures were conducted using R Version 1.1.453 (Team, 2014).

## Results

### Participant characteristics

The mean age ( $19.2 \pm .9$  years) of all participants and of male ( $19.3 \pm .8$  years) and female participants ( $19.2 \pm .6$  years) were similar. Mean level of APDR for those who reported eating less than usual prior to drinking to feel the effects of alcohol faster ( $1.24 \pm 0.7$ ) was slightly higher than those who reported skipping one or more meals prior to drinking to feel the effects of alcohol faster ( $1.12 \pm 0.5$ ). Mean baseline BMI was normal (healthy weight) across males ( $24.5 \text{ kg/m}^2 \pm 4.2$ ) and females ( $24.0 \text{ kg/m}^2 \pm 4.4$ ). Frequencies for each binge drinking category were the following: never (32.9%), less than monthly (20%), monthly (7.3%), weekly (3.8%), and daily or almost daily (0.2%). Given the daily binge drinking frequency category was not adequately represented in our non-clinical sample, binge drinking frequency categories were collapsed to: 1) never or less than monthly; 2) monthly; and 3) weekly or daily. Age was also not included in the ordered model due to sample age homogeneity (i.e. the sample was composed of freshmen averaging 19 years of age). Descriptive statistics for baseline participant characteristics are provided in Table 1.

**Table 1.** Participant characteristics and behaviors at the start of their freshmen year ( $n = 1,149$ ).

Factor	Frequency <i>n</i> (%)	Missing <i>n</i> (%)
<i>Sex</i>		
Male	377 (32.8)	27 (2.3)
<i>Ethnic origin</i>		
Hispanic or Latino	204 (17.8)	34 (3)
<i>Race</i>		
Black or African American	162 (14.1)	0
Asian/Asian American	137 (11.9)	0
White	817 (71.1)	0
<i>Greek affiliation</i>		
Greek campus organization affiliation	11 (1.0)	28 (2.4)
<i>Athlete status</i>		
Athlete	68 (5.9)	36 (3.1)
<i>Geographic region</i>		
At-risk	528 (46)	98 (8.5)
<i>Residence</i>		
On campus	961 (83.6)	21 (1.8)
<i>Parents overweight/obese</i>	569 (49.5)	164 (14.3)
<i>Family affluence</i>	15 (1.3)	107 (9.3)
<i>Binge drinking frequency</i>		411 (35.8)
Never or less than monthly	608 (52.9)	
Monthly	84 (7.3)	
Weekly or daily	46 (4.0)	
<i>Alcohol-related proactive dietary restriction (APDR)</i>	<b>Mean (SD)</b>	
Eaten less than usual (low)	1.24 (0.67)	405 (35.2)
Skipped one or more meals (high)	1.12 (0.49)	407 (35.4)
<i>BMI (kg/m<sup>2</sup>)</i>	24.17 (4.32)	30 (2.6)
<i>EAT-26 Total</i>	7.99 (8.00)	9 (0.8)
<i>Waist hip ratio</i>	0.79 (0.07)	14 (1.2)
<i>IPAQ Total (met min)</i>	2729.64 (2017.63)	57 (5)
<i>NClfv Total (CE/day)</i>	2.45 (2.20)	5 (0.4)

### Alcohol-related proactive dietary restriction and binge drinking frequency

Model fit for the cross-sectional and longitudinal models examining the predictive ability of varying levels of APDR to contribute to binge drinking frequency was assessed using a chi-square test for statistical significance. A significant chi-square value indicated the model provided a significant improvement over the baseline intercept-only model for the cross-sectional ( $\chi^2(14) = 105.53, p < .001$ ) and longitudinal ( $\chi^2(16) = 182.73, p < .001$ ) models. Goodness of fit was assessed using the deviance chi-square statistics. Non-significant deviance chi-square values ( $\chi^2(479) = 384.35, p = .99$ ) and ( $\chi^2(752) = 534.18, p = .99$ ) indicated the observed data were consistent with the fitted model for the cross-sectional and longitudinal models, respectively. Furthermore, pseudo  $R^2$  indices indicated that the cross-sectional model predicted between 9% and 14% of the variance in baseline binge drinking frequency and the longitudinal model predicted between 15% and 22% of the variance in binge drinking frequency at follow-up. The test of parallel lines was used to assess whether the odds for each binge drinking frequency category were consistent across different thresholds. A non-significant chi-square value indicated the odds for each binge drinking frequency category were consistent across different thresholds for both the cross-sectional ( $\chi^2(14) = 13.99, p = .48$ ) and longitudinal ( $\chi^2(16) = 34.13, p = .20$ ) models.

We hypothesized that both low and high baseline APDR would remain predictors of binge drinking frequency at 7-month follow-up. Our auxiliary hypothesis stated that high

baseline APDR would be more strongly associated with binge drinking frequency at 7-month follow-up than low baseline APDR. Results did not support either of the two hypotheses. The results suggested those who reported eating less than usual prior to drinking to feel the effects of alcohol faster (i.e. low APDR) were associated with greater odds for a higher binge drinking frequency at baseline (1.27 (95% CI, 0.06 to 0.42), Wald  $\chi^2(1) = 8.46, p = .009$ ), but not associated with binge drinking frequency at 7-month follow up (1.02 (95% CI,  $-0.18$  to 0.23), Wald  $\chi^2(1) = .51, p = .83$ ). On the other hand, findings suggested those who reported skipping one or more meals prior to drinking to feel the effects of alcohol faster (i.e. high APDR) were not associated with binge drinking frequency at baseline (1.20 (95% CI,  $-0.07$  to .43), Wald (1) = 2.40,  $p = .16$ ) nor at 7-month follow-up (1.13 (95% CI,  $-0.14$  to 0.39), Wald (1) = 1.42,  $p = .35$ ). As compared to females, males had significantly greater odds for an increased binge drinking frequency at baseline (1.62 (95% CI, 0.31 to 0.65), Wald  $\chi^2(1) = 38.61, p < .001$ ), but not at 7-month follow up (1.19 (95% CI,  $-0.03$  to 0.37), Wald  $\chi^2(1) = 5.17, p = .09$ ). Furthermore, monthly (1.92 (95% CI 0.36 to 0.95), Wald (1) = 28.74,  $p < .001$ ) and weekly or daily (2.47 (95% CI, 0.51 to 1.3), Wald (1) = 28.37,  $p < .001$ ) baseline binge drinking frequencies were significantly associated with greater odds of an increased binge drinking frequency at 7-month follow-up, as compared to never or less than monthly baseline binge drinking frequency.

Non-significant alcohol-related proactive dietary restriction missing indicators suggested individuals who did not self-report APDR at baseline did not significantly differ on

**Table 2.** Ordinal regression analyses to examine the effect of alcohol-related proactive dietary restriction at baseline by college freshmen on binge drinking frequency at 7-month follow up.

Predictor variables	Complete case analysis using negative log-log link function ( <i>n</i> = 465)	50 Random forest multiple imputations using complementary log-log link function ( <i>n</i> = 1,149)	50 Random forest multiple imputations & missing indicators using complementary log-log link function ( <i>n</i> = 1,149)
Baseline binge drinking frequency			
Never or less than monthly	reference category	reference category	reference category
Monthly	4.77 (1.09, 2.04)***	2.00 (.41, .98)***	1.92 (.36, .95)***
Weekly or daily	8.71 (1.62, 2.71)***	2.51 (.54, 1.30)***	2.47 (.51, 1.3)***
APDR <sup>a</sup>			
Low	1.38 (-.14, .78)	1.01 (-.19, .21)	1.02 (-.18, .23)
High	1.12 (-.46, .68)	1.19 (-.09, .43)	1.13 (-.14, .39)
Ethnic origin			
Hispanic or Latino	1.52 (-.06, .89)	1.24 (-.003, .43)	1.18 (-.06, .39)
Race			
Black or African American	0.94 (-.84, .72)	1.09 (-.24, .42)	1.09 (-.28, .44)
Asian/Asian American	3.24 (.36, 1.99)**	1.26 (-.11, .58)	1.28 (-.13, .62)
White	2.56 (.20, 1.68)*	1.25 (-.05, .50)	1.22 (-.11, .50)
Sex			
Male	2.09 (.32, 1.16)**	1.20 (-.004, .36)	1.19 (-.03, .37)
Athlete status			
Athlete	1.62 (-.15, 1.12)	1.13 (-.21, .44)	1.11 (-.24, .44)
Geographic region			
At-risk	1.44 (-.07, .80)	1.17 (-.02, .34)	1.14 (-.07, .32)
Residence			
On campus	0.90 (-.72, .51)	1.01 (-.25, .27)	1.02 (-.26, .30)
Off-campus	reference category	reference category	reference category
Greek affiliation			
Greek	2.55 (-.08, 1.95)	1.99 (-.06, 1.43)	1.87 (-.13, 1.38)
Low APDR <sup>a</sup> – missing indicator			22.20 (-310.38, 316.58)
High APDR <sup>a</sup> – missing indicator			0.04 (-316.72, 310.24)
Binge drinking frequency at follow up – missing indicator			0.78 (-.47, -.03)*

Note: results are presented as odds ratios with 95% confidence intervals in parentheses.

\**p* < .05; \*\**p* < .01; \*\*\**p* < .001.

<sup>a</sup>Alcohol-related proactive dietary restriction (APDR).

APDR from those who provided self-reported APDR: 1) eating less than usual (22 (95% CI, -310 to 316), Wald  $\chi^2$  (1) = 36.07, *p* = .98); and 2) skipping one or more meals (0.04 (95% CI, -316 to 310), Wald  $\chi^2$  (1) = 0.11, *p* = .98). A non-significant baseline binge drinking frequency missing indicator (0.63 (95% CI, -1.27 to -0.34), Wald  $\chi^2$  (1) = 2.86, *p* = .26) suggested individuals who did not self-report binge drinking frequency at baseline did not significantly differ on binge drinking frequency from those who did self-report a baseline binge drinking frequency. A significant binge drinking frequency at follow-up missing indicator (0.78 (95% CI, -0.47 to -0.03), Wald  $\chi^2$  (1) = 0.66, *p* = .03) suggested individuals who did not self-report binge drinking frequency at follow-up had decreased odds for binge drinking frequency.

A non-significant mean difference was observed across baseline and 7-month follow up binge drinking frequencies (*t* (580) = 0.89, *p* = 0.39). All results for the longitudinal ordinal regression model are provided in Table 2. Results for the cross-sectional ordinal regression and paired sample *t*-test are shown in Tables 3 and 4.

## Discussion

This study focused on the strength of the relationship and longitudinal effect of varying levels of APDR on binge drinking frequency among a U.S. college freshman cohort. Findings from this study revealed that students reporting higher frequencies of eating less than usual prior to drinking to feel the psychoactive effects of alcohol faster (low APDR) early in the first semester of college (baseline) had greater odds for increased binge drinking frequency at baseline; however, this predictive relationship was not observed between baseline low APDR and binge drinking frequency at 7-month follow-up. In fact, no significant mean difference was detected between students' binge drinking frequencies at baseline and follow-up. On the other hand, findings suggested those who reported skipping one or more meals prior to drinking to feel the psychoactive effects of alcohol faster (high APDR) were not associated with binge drinking frequency at baseline nor at 7-month follow-up.

**Table 3.** Ordinal regression analyses to examine the effect of alcohol-related proactive dietary restriction at baseline on college students' binge drinking frequency at baseline (cross-sectional).

Predictor variables	Complete case analysis using negative log-log link function (n=672)	50 Random forest multiple imputations using complementary log-log link function (n=1,149)	50 Random forest multiple imputations & missing indicators using complementary log-log link function (n=1,149)
APDR <sup>a</sup>			
Low	1.83 (.28, .93)***	1.33 (.10, .47)**	1.27 (.06,.42) **
High	1.26 (-.15, .61)	1.17 (-.09, .41)	1.20 (-.07, .43)
Ethnic origin			
Hispanic or Latino	0.77 (-.15, .61)	0.94 (-.29, .15)	0.91 (-.33, .13)
Race			
Black or African American	0.94 (-.87, .74)	1.06 (-.27, .37)	1.02 (-.33, .35)
Asian/Asian American	1.44 (-.40,1.14)	1.15 (-.19, .46)	1.09 (-.26, .43)
White	1.66 (-.21, 1.22)	1.20 (-.11, .47)	1.15 (-.17, .44)
Sex			
Male	4.27 (1.04, 1.86)***	1.65 (.32, .66)***	1.62 (.31, .65) ***
Athlete status			
Athlete	1.28 (-.43,.92)	1.19 (-.14, .49)	1.17 (-.17, .48)
Geographic region			
At-risk	1.40 (-.05,.72)	1.14 (.04, .30)	1.11 (-.07, .28)
Residence			
On campus	1.48 (-.32, 1.11)	1.08 (-.20, .35)	1.08 (-.21, .37)
Off-campus	reference category	reference category	reference category
Greek affiliation			
Greek	0.85 (-1.34, 1.02)	1.00 (-.72, .72)	1.01 (-.73, .74)
Low APDR <sup>a</sup> – missing indicator			0.34 (-3.40, 1.22)
High APDR <sup>a</sup> – missing indicator			3.90 (-.81, 3.53)
Baseline binge drinking frequency – missing indicator			0.63 (-1.27, .34)

Note: results are presented as odds ratios with 95% confidence intervals in parentheses.

\* $p < .05$ ; \*\* $p < .01$ ; \*\*\* $p < .001$ .

<sup>a</sup>Alcohol-related proactive dietary restriction (APDR).

**Table 4.** Paired Sample T-test to Compare Binge Drinking Frequency at Baseline and 7-month Follow-up.

	t	df	Sig. (2-tailed)	Mean Difference (SE)	95% C.I.
Complete case analysis (n = 505)					
BDFT2 – BDFT1 <sup>a</sup>	4.6	504	<.001***	0.12 (0.03)	(0.07, 0.17)
Fifty random forest multiple imputations (n = 1,149)					
BDFT2 – BDFT1 <sup>a</sup>	0.87	580	0.39	0.02 (0.02)	(-0.02, 0.06)

<sup>a</sup>The mean difference was calculated as (binge drinking frequency at 7-month follow up – baseline binge drinking frequency).

\*Note: \* $p < .05$ ; \*\* $p < .01$ ; \*\*\* $p < .001$ .

These findings corroborate literature findings suggesting students become acclimated to perceived college drinking norms within their first months of college; thus, it is likely the nonsignificant change in binge drinking frequency observed over the course of students' freshman year was due to the early internalization of perceived college drinking norms (Sher & Rutledge, 2007). However, the strong cross-sectional relationship between APDR and student binge drinking observed by Bryant et al. (2012) is only corroborated for low APDR by this study. Bryant et al. (2012) found that college students who restrict their energy intake prior

to alcohol consumption (to feel the psychoactive effects of alcohol faster) have a greater likelihood to participate in binge drinking. The group compared binge drinkers with non-binge drinkers across dietary restriction prior to alcohol consumption and found increased mean level scores for binge drinkers. It is possible former findings by Bryant et al. (2012) were not examined individually at both APDR thresholds. However, when examined at both APDR thresholds, our study indicates the relationship between high APDR and binge drinking frequency is not present at baseline nor at follow-up for non-clinical college students. The



nonclinical nature of the student cohort under study may explicate why high APDR was not significantly related to binge drinking in our sample at baseline nor follow-up.

Findings from this investigation extend upon current literature findings indicating a cross-sectional relationship between proactive dietary restriction to feel the psychoactive effects of alcohol faster and binge drinking. Specifically, this study addresses a current gap in the college alcohol literature with respect to: 1) the strength of the relationships between varying levels of alcohol-related proactive dietary restriction and binge drinking frequency; and 2) a longitudinal investigation of the predictive ability of APDR on increased binge drinking frequency. This study measured APDR on a severity spectrum, from eating less than usual to skipping one or more meals to feel the effects of alcohol faster. Measuring alcohol-related proactive dietary restriction on a severity spectrum allowed for unique findings with respect to its longitudinal relationship to binge drinking frequency.

Nevertheless, there are limitations to this study. While our study inclusion criteria consisted of freshmen students who were potentially at-risk for poor health behaviors; on average, the final study sample consisted of normal weight freshmen students with a racial composition representative of freshmen attending U.S. universities (Staklis & Chen, 2010). Furthermore, the expected missing data when assessing alcohol intake of college students, most of whom were less than the legal drinking age of 21 years, across binge drinking frequency and alcohol-related proactive dietary restriction substantially lowered sample size and possibly biased model parameter estimates. Thus, study results should be interpreted with caution; however, missing data was accounted for by conducting 50 multiple random forest imputations of the ordinal logistic regression model for comparison against the complete case analysis (CCA) model estimates. Model comparisons rendered similar statistically significant predictors, with more conservative estimates for the imputed models. Significant predictors across both CCA and imputed models also closely corresponded with findings in the current college alcohol literature. For instance, study results corroborated existing evidence in the college alcohol literature regarding an increased risk for binge drinking among males. Furthermore, missing indicators were assessed for baseline alcohol-related proactive dietary restriction and binge drinking frequency at baseline and follow-up. Missing indicator findings across both levels of APDR suggested individuals who did not self-report APDR at baseline did not significantly differ on APDR from those who did provide self-reported APDR. While binge drinking frequency missing indicators suggested individuals who did not self-report binge drinking frequency at baseline were not significantly different in binge drinking frequency from those who did self-report a baseline binge drinking frequency, individuals who did not self-report binge drinking frequency at follow-up had decreased odds for binge drinking frequency. As such, the analysis of the missing data for baseline alcohol-related proactive dietary restriction and binge drinking frequency at follow-up in our sample demonstrated it was not likely that those who skipped these items on the

survey were more likely to engage in APDR or binge drinking than those who provided responses. Moreover, baseline binge drinking frequency was controlled for in the longitudinal regression model. This indicated that significant effects observed were beyond the variance explained by the outcome at baseline, thus improving model validity and fit.

This study establishes that low APDR may, in fact, serve as a significant predictor for binge drinking frequency early in the first semester of college; however, current alcohol-related proactive dietary restriction does not predict binge drinking at 7 month follow-up. With this in mind, campus health professionals are urged to emphasize the hazardous health effects of engaging in alcohol-related proactive dietary restriction early in freshmen students' first weeks of exposure to the college milieu. Future longitudinal studies with more time points are needed to confirm the potential for levels of alcohol-related proactive dietary restriction and binge drinking frequency to remain stable after the first few weeks of exposure to the college milieu. The corroboration of these relationships will further enhance campus health promotion and intervention programming *via* targeted screening efforts. The inclusion of alcohol-related proactive dietary restriction to feel the psychoactive effects of alcohol faster as a correlate for increased binge drinking frequency is suggested for college campus health screenings. Future longitudinal research examining the effect of alcohol-related proactive dietary restriction to feel the psychoactive effects of alcohol faster on binge drinking frequency is warranted with larger sample sizes that are more representative of the U.S. college population.

## Acknowledgements

None.

## Declaration of interest

The authors report no conflicts of interest.

## Authorship

Authors GC, TEB, and AEM designed the study, SEC, MDO, WZ, ME and AEM collected data, GC conducted the statistical analysis, GC wrote the first draft of the manuscript. All authors contributed to and have approved the final manuscript.

## Ethical standards disclosure

This study was conducted according to the guidelines laid down in the Declaration of Helsinki. The Institutional Review Board at the University of Tennessee approved research activities at the University of Tennessee, West Virginia University, South Dakota State University, Syracuse University, and University of Maine. University of Florida, Auburn University, and Kansas State University Institutional Review Boards provided approval for research on their respective campuses. Written informed consent was obtained from all subjects.

## Funding

Funding for this study was provided by USDA Grant 2014-67001-21851. USDA had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

## ORCID

Tracey E. Barnett  <http://orcid.org/0000-0002-2932-1540>

Wenjun Zhou  <http://orcid.org/0000-0003-2346-8151>

Anne E. Mathews  <http://orcid.org/0000-0003-3504-5826>

## References

- Barry, A. E., & Piazza-Gardner, A. K. (2012). Drunkorexia: Understanding the co-occurrence of alcohol consumption and eating/exercise weight management behaviors. *Journal of American College Health: J of ACH*, 60(3), 236–243. <https://doi.org/10.1080/07448481.2011.587487>
- Barry, A. E., Whiteman, S., Piazza-Gardner, A. K., & Jensen, A. C. (2013). Gender differences in the associations among body mass index, weight loss, exercise, and drinking among college students. *Journal of American College Health: J of ACH*, 61(7), 407–413. <https://doi.org/10.1080/07448481.2013.823973>
- Bennett, D. A. (2001). How can I deal with missing data in my study? *Australian and New Zealand Journal of Public Health*, 25(5), 464–469. <https://doi.org/10.1111/j.1467-842X.2001.tb00294.x>
- Boyce, W., Torsheim, T., Currie, C., & Zambon, A. (2006). The family affluence scale as a measure of national wealth: Validation of an adolescent self-report measure. *Social Indicators Research*, 78(3), 473–487. <https://doi.org/10.1007/s11205-005-1607-6>
- Bryant, J. B., Darkes, J., & Rahal, C. (2012). College students' compensatory eating and behaviors in response to alcohol consumption. *Journal of American College Health: J of ACH*, 60(5), 350–356. <https://doi.org/10.1080/07448481.2011.630702>
- Burke, S. C., Cremeens, J., Vail-Smith, K., & Woolsey, C. (2010). Drunkorexia: Calorie restriction prior to alcohol consumption among college freshman. *Journal of Alcohol and Drug Education*, 54(2), 17–34.
- Bush, K., Kivlahan, D. R., McDonell, M. B., Fihn, S. D., & Bradley, K. A. (1998). The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. *Archives of Internal Medicine*, 158(16), 1789–1795. <https://doi.org/10.1001/archinte.158.16.1789>
- Castañeda, G., Colby, S. E., Barnett, T. E., Olfert, M. D., Zhou, W., Leite, W. L., El Zein, A., & Mathews, A. E. (2020). Examining the effect of weight conscious drinking on binge drinking frequency among college freshmen. *Journal of American College Health: J of ACH*, 68(8), 906–908. <https://doi.org/10.1080/07448481.2019.1642204>
- Cederbaum, A. I. (2012). Alcohol metabolism. *Clinics in Liver Disease*, 16(4), 667–685. <https://doi.org/10.1016/j.cld.2012.08.002>
- Eisenberg, M. H., & Fitz, C. C. (2014). "Drunkorexia": exploring the who and why of a disturbing trend in college students' eating and drinking behaviors. *Journal of American College Health: J of ACH*, 62(8), 570–577. <https://doi.org/10.1080/07448481.2014.947991>
- Enders, C. K. (2010). *Applied missing data analysis*. Guilford press.
- Garner, D., Olmsted, M., Bohr, Y., & Garfinkel, P. (1982). The Eating Attitudes Test: Psychometric features and clinical correlates. *Psychological Medicine*, 12(4), 871–878. <https://doi.org/10.1017/s0033291700049163>
- Giles, S. M., Champion, H., Sutfin, E. L., McCoy, T. P., & Wagoner, K. (2009). Calorie restriction on drinking days: An examination of drinking consequences among college students. *Journal of American College Health: J of ACH*, 57(6), 603–610. <https://doi.org/10.3200/JACH.57.6.603-610>
- Hingson, R., Heeren, T., Winter, M., & Wechsler, H. (2005). Magnitude of alcohol-related mortality and morbidity among US college students ages 18–24: Changes from 1998 to 2001. *Annual Review of Public Health*, 26(1), 259–279. <https://doi.org/10.1146/annurev.publhealth.26.021304.144652>
- Hingson, R. W., Heeren, T., Zakocs, R. C., Kopstein, A., & Wechsler, H. (2002). Magnitude of alcohol-related mortality and morbidity among US college students ages 18–24. *Journal of Studies on Alcohol*, 63(2), 136–144. <https://doi.org/10.15288/jsa.2002.63.136>
- Kelly-Weeder, S. (2011). Binge drinking and disordered eating in college students. *Journal of the American Academy of Nurse Practitioners*, 23(1), 33–41.
- Krieger, H., Young, C. M., Anthenien, A. M., & Neighbors, C. (2018). The epidemiology of binge drinking among college-age individuals in the United States. *Alcohol Research: Current Reviews*, 39(1), 23–30.
- Lieber, C. S. (2003). Relationships between nutrition, alcohol use, and liver disease. *Alcohol Research & Health: The Journal of the National Institute on Alcohol Abuse and Alcoholism*, 27(3), 220–231.
- Loso, J., Staub, D., Colby, S. E., Olfert, M. D., Kattelman, K., Vilaro, M., Colee, J., Zhou, W., Franzen-Castle, L., & Mathews, A. E. (2018). Gardening experience is associated with increased fruit and vegetable intake among first-year college students: A cross-sectional examination. *Journal of the Academy of Nutrition and Dietetics*, 118(2), 275–283. <https://doi.org/10.1016/j.jand.2017.09.005>
- Nelson, T. F., Xuan, Z., Lee, H., Weitzman, E. R., & Wechsler, H. (2009). Persistence of heavy drinking and ensuing consequences at heavy drinking colleges. *Journal of Studies on Alcohol and Drugs*, 70(5), 726–734. <https://doi.org/10.15288/jsad.2009.70.726>
- Peralta, R. L. (2002). Alcohol use and the fear of weight gain in college: Reconciling two social norms. *Gender Issues*, 20(4), 23–42. <https://doi.org/10.1007/s12147-002-0021-5>
- Piazza-Gardner, A. K., & Barry, A. E. (2014). A qualitative investigation of the relationship between consumption, physical activity, eating disorders, and weight consciousness. *American Journal of Health Education*, 45(3), 174–182. <https://doi.org/10.1080/19325037.2014.901112>
- Rahal, C. J., Bryant, J. B., Darkes, J., Menzel, J. E., & Thompson, J. K. (2012). Development and validation of the compensatory eating and behaviors in response to alcohol consumption scale (CEBRACS). *Eating Behaviors*, 13(2), 83–87. <https://doi.org/10.1016/j.eat-beh.2011.11.001>
- Sher, K. J., & Rutledge, P. C. (2007). Heavy drinking across the transition to college: Predicting first-semester heavy drinking from pre-college variables. *Addictive Behaviors*, 32(4), 819–835. <https://doi.org/10.1016/j.addbeh.2006.06.024>
- Staklis, S., & Chen, X. (2010). Profile of Undergraduate Students: Trends from Selected Years, 1995–96 to 2007–08. Web Tables. NCES 2010–220. National Center for Education Statistics.
- Stekhoven, D. J., & Bühlmann, P. (2012). MissForest-non-parametric missing value imputation for mixed-type data. *Bioinformatics (Oxford, England)*, 28(1), 112–118. <https://doi.org/10.1093/bioinformatics/btr597>
- Team, R. C. (2014). *A language and environment for statistical computing*. R Foundation for Statistical Computing. 2014. <https://www.R-project.org>.
- Thompson, F. E., Subar, A. F., Smith, A. F., Midthune, D., Radimer, K. L., Kahle, L. L., & Kipnis, V. (2002). Fruit and vegetable assessment: Performance of 2 new short instruments and a food frequency questionnaire. *Journal of the American Dietetic Association*, 102(12), 1764–1772. [https://doi.org/10.1016/S0002-8223\(02\)90379-2](https://doi.org/10.1016/S0002-8223(02)90379-2)
- Wechsler, H., Lee, J. E., Kuo, M., Seibring, M., Nelson, T. F., & Lee, H. (2002). Trends in college binge drinking during a period of increased prevention efforts. Findings from 4 Harvard School of Public Health College Alcohol Study surveys: 1993–2001. *Journal of American College Health: J of ACH*, 50(5), 203–217. <https://doi.org/10.1080/07448480209595713>
- White, A., & Hingson, R. (2013). The burden of alcohol use: Excessive alcohol consumption and related consequences among college students. *Alcohol Research: Current Reviews*, 35(2), 201–218.