



Anti-allergic activity of lotus root (*Nelumbo nucifera*) powder in TDI-sensitized nasal allergy model mice

Mayumi Kaneyasu, Maki Nagata, Hiromi Ikeda, Koichiro Ohnuki & Kuniyoshi Shimizu

To cite this article: Mayumi Kaneyasu, Maki Nagata, Hiromi Ikeda, Koichiro Ohnuki & Kuniyoshi Shimizu (2019) Anti-allergic activity of lotus root (*Nelumbo nucifera*) powder in TDI-sensitized nasal allergy model mice, Food and Agricultural Immunology, 30:1, 968-978, DOI: [10.1080/09540105.2019.1651255](https://doi.org/10.1080/09540105.2019.1651255)

To link to this article: <https://doi.org/10.1080/09540105.2019.1651255>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 07 Aug 2019.



Submit your article to this journal [↗](#)



Article views: 1082



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 3 View citing articles [↗](#)

Anti-allergic activity of lotus root (*Nelumbo nucifera*) powder in TDI-sensitized nasal allergy model mice

Mayumi Kaneyasu^a, Maki Nagata^b, Hiromi Ikeda^b, Koichiro Ohnuki^c and Kuniyoshi Shimizu^b

^aDepartment of Human Nutrition, Faculty of Nursing and Human Nutrition, Yamaguchi Prefectural University, Yamaguchi, Japan; ^bDepartment of Agro-environmental Sciences, Faculty of Agriculture, Kyushu University, Fukuoka, Japan ; ^cDepartment of Biological and Environmental Chemistry, Kinki University, Fukuoka, Japan

ABSTRACT

Lotus root (*Nelumbo nucifera*) is a popular food in Japan and southeast Asia. It is reported that lotus root includes dietary fiber, vitamin C, and abundant polyphenolic compounds. In Chinese medicine, lotus powder is also utilized in anti-tussive or anti-inflammatory drugs. In this study, we focused on the effects of lotus root powder on allergy symptoms in mice. First, we assessed nasal allergy symptoms in 2,4-diisocyanic acid toluene (TDI)-sensitized nasal allergy model mice. We found that lotus root powder alleviated nasal allergy symptoms in the mice. In addition, we estimated serum inflammatory parameters such as serum total IgE, histamine, and leukotriene B4. Lotus root powder intake significantly decreased serum total IgE and leukotriene B4 compared with standard food intake. We suggest that lotus root powder intake may alleviate allergic symptoms.

ARTICLE HISTORY

Received 6 December 2018
Accepted 22 July 2019



KEYWORDS

Lotus root; allergic symptoms; anti-inflammatory

1. Introduction

The rhizome of lotus root (*Nelumbo nucifera*), an aquatic plant, is a very popular food in Japan and southeast Asia (Sridhar & Bhat, 2007; Wang, Yen, Liang, & Wu, 2003). There are some local brands of lotus root on the Japanese market. In Yamaguchi prefecture, an indigenous brand of lotus root is called “Iwakuni-renkon,” which is white, soft, and sticky. Iwakuni-renkon’s morphology is unique in that its root has nine holes (water-conducting tissue), whereas common lotus root has eight holes, although there are individual differences.

The lotus root includes abundant nutrients such as dietary fiber and vitamin C. Some studies have shown that lotus root contains abundant polyphenolic compounds and possesses several beneficial health properties, including hypoglycemic, anti-inflammatory, anti-epileptic, antioxidant activities and therapeutic application in memory disorders (Huang et al., 2011; Khan, Rajput, & Assad, 2019; Mukherjee, Mukherjee,

CONTACT Mayumi Kaneyasu  kaneyasu@yamaguchi-pu.ac.jp  Department of Human Nutrition, Yamaguchi Prefectural University, Yamaguchi 753-0021, Japan

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Maji, Rai, & Heinrich, 2009; Mukherjee, Saha, Pal, & Saha, 1997; Rajput, Khan, & Assad, 2017; Yan, Wang, & Peng, 2009). Furthermore, in Chinese medicine, lotus root is utilized as an anti-tussive and anti-inflammatory drug. Thus, we expected that intake of lotus root could lead to the improvement of allergy symptoms. In this study, we examined the influence of lotus root powder intake on allergy onset using 2,4-diisocyanic acid toluene (TDI)-sensitized nasal allergy model mice (Dearman, Basketter, & Kimber, 1996).

2. Materials and methods

2.1. Animals and housing conditions

Four-week-old female BALB/c mice were reared at $22 \pm 2^\circ\text{C}$ with $50 \pm 10\%$ humidity under a 12-hour light/dark cycle (lights on at 08:00 and off at 20:00). After the mice were acclimated for one week, we divided the mice into a nonsensitization group and a sensitization group. The animals in the nonsensitization group were fed standard food, while those in the sensitization group were further classified into three groups of 5 mice each that were fed as follows for 4 weeks: standard food (based on AIN-93G), standard food containing 5% lotus root powder, and standard food containing 10% lotus root powder. The lotus root powder was processed and provided by Hironaka Food Ltd. (Yamaguchi, Japan).

2.2. Experimental procedure

2.2.1. Schedule of sensitization

After the 4-week feeding period, we applied 2 μL of 5% 2,4-diisocyanic acid toluene (TDI) into a nostril of each animal under anesthesia in the sensitization group for 5 consecutive days. As a control, we applied ethyl acetate, a solvent of TDI solution, into a nostril of each mouse in the nonsensitization group. One week after the first sensitization, we carried out re-sensitization for 5 days in a row using the same method described above. One week after the re-sensitization, we applied the final drop of 5 μL of 2.5% TDI into each animal's nostril under nonanesthetic conditions.

2.2.2. Visual assessment of allergic symptoms

We evaluated the severity of allergic symptoms for 10 min immediately after the final sensitization. As visual characteristics of nasal allergic symptoms, both reddish tinges and hair loss were observed on the nose. The severity of the nasal allergies was scored on a 4-point scale from 0 (no symptoms) to 3 (very severe symptoms) by observing the appearance of each mouse.

2.2.3. Measurement of serum inflammatory parameters

One week after the final sensitization, the mice were sacrificed under anesthesia with pentobarbital (Sumitomo Dainippon Pharma Co., Ltd., Osaka, Japan) and their trunk blood was collected to evaluate the levels of inflammatory parameters such as serum total IgE, histamine, and leukotriene B₄ (LTB₄). The concentrations of serum total IgE, histamine, and LTB₄ were analyzed using three respective ELISA kits: the Mouse IgE ELISA

Quantitation Kit (Bethyl Laboratories, Montgomery, TX, USA), the Histamine ELISA Kit (Neogen, Lansing, MI, USA), and the Leukotriene B₄ EIA Kit (Oxford Biomedical Research, Rochester Hills, MI, USA). The concentrations were measured according to the respective manufacturers' instructions.

2.2.4. Measurement of splenic inflammatory parameters

One week after the final sensitization, the mice were sacrificed under anesthesia with pentobarbital (Sumitomo Dainippon Pharma Co., Ltd., Osaka, Japan) and their spleens were collected. The spleens were weighed, and the ratios of splenic inflammatory parameters, such as CD45R⁺ cells (B-cells) of splenic lymphocytes, a CD3⁺ cell (mature T-cell), a CD4⁺ cell (helper T-cell), and a CD8⁺ cell (suppressor T-cell), were measured. Spleen cells were divided into two test tubes. The cells in one tube were fixed and stained with FITC-conjugated anti-mouse CD4 mAb and PE-conjugated anti-mouse CD8 mAb. The cells in the other tube were fixed and stained with FITC-conjugated anti-mouse CD45R mAb and PE-conjugated anti-mouse CD3 mAb. The samples were analyzed in a flow cytometer (FACS Calibur, Becton-Dickinson, Franklin Lakes, NJ, USA), and the data were processed by CELL Quest software (Becton-Dickinson).

In addition, we measured concanavalin A (Con A)-stimulated interleukin-4 (IL-4) production. Briefly, spleen cells derived from sensitized or nonsensitized mice were cultivated for 48 h at 37°C, 5% CO₂. IL-4 in the supernatant was measured by ANALYZA Immunoassay System mouse IL-4 (Bio-Techne, Minneapolis, MN, USA).

2.3. Statistical analysis

All data were analyzed using one-way ANOVA and were subjected to a post-hoc Fisher's PLSD test. The significance threshold was set at $P < 0.05$. All analyses were performed with StatView (version 5, SAS Institute, Cary, NC, USA).

3. Results and discussion

3.1. Physical conditions

We tested anti-allergic or anti-inflammatory activity using TDI-sensitized nasal allergy model mice. Intranasal application of TDI induced nasal allergic symptoms. Before TDI treatment, mice were given the standard diet or the standard diet supplemented with either 5% or 10% lotus root powder for 4 weeks. TDI treatment was performed intermittently for 4 weeks. During the whole experimental period including before and after intermittent TDI treatment, food intake and body weight gain were quite similar among all groups (data not shown). We inferred from these results that lotus root powder does not affect the food intake or growth of animals.

3.2. Allergic symptoms

We assessed the state of TDI-sensitized nasal allergy symptoms. In TDI-sensitized mice, the hair loss and redness around the nose were prominent, indicating the occurrence of severe allergic reactions. Meanwhile, these allergic symptoms were clearly alleviated by




Figure 1. Typical nasal allergic symptoms in the nonsensitization and sensitization groups. (A) Nonsensitization group; (B) standard food sensitization group; (C) standard food plus 5% lotus root powder sensitization group; (D) standard food plus 10% lotus root powder sensitization group.




Figure 2. Scores of nasal allergic symptoms. *Significant difference compared to nonsensitization group ($P < 0.05$). Each value is the mean \pm SEM ($n = 5$ /group).

lotus root intake in a dose-dependent manner (Figure 1A–D). The severity scores of allergic symptoms were significantly ($P < 0.05$) increased in the TDI-sensitized mice than in the nonsensitized mice (Figure 2). These scores also tended to decrease by lotus root intake in a dose-dependent manner. These results show that lotus root powder alleviates nasal allergic symptoms.

3.3. Serum inflammatory parameters

We investigated serum inflammatory parameters. The concentration of serum histamine, which is released from activated mast cells having an allergic response, differed significantly ($P < 0.05$) between the non-TDI-sensitized mice and the TDI-sensitized mice fed the 10% lotus root powder diet (Figure 3A). Elevated levels of LTB_4 have been reported in people suffering from various allergic diseases, and these levels have been related to disease activity and response to treatment (Ohnishi, Miyahara, & Gelfand, 2008). In this study, the concentration of serum LTB_4 was significantly ($P < 0.05$) decreased in the TDI-sensitized 10% lotus root powder diet group compared with the TDI-sensitized standard diet group (Figure 3B). It is reported that not only histamine but also leukotriene are parameters indicated allergy response. In allergy response, chemical mediators such as histamine and leukotrienes are released from mast cells. The chemical mediators cause sneezing, watery rhinorrhea and nasal blockage as an early phase reaction. In addition, Leukotrienes, produced by inflammatory cells, cause nasal blockage as a late phase reaction, seen at 6–10 h after antigen exposure (Miyake & Karasuyama, 2017; Okubo et al., 2017; Wedi, Novacovic, Koerner, & Kapp, 1999). The Other studies such these reports indicated that histamine release and leukotriene production are regulated differentially. Although lotus powder induce histamine release from mast cells by different regulation, it is suggest that lotus powder may strongly inhibit LTB_4 production in inflammatory cells. The results suggested that the alleviation of allergic symptoms by lotus root powder intake.




Figure 3. Serum inflammatory parameters. (A) Serum histamine concentration, *significant difference compared to nonsensitization group ($P < 0.05$). (B) Serum LTB_4 concentration, *significant difference between standard food sensitization group and 10% lotus root powder sensitization group ($P < 0.05$). Each value is the mean \pm SEM ($n = 5$ /group).




Figure 4. Total serum IgE concentration. Differential letters indicate significant difference ($P < 0.05$). Each value is the mean \pm SEM ($n = 5$ /group).

IgE, an antibody produced by the body's immune system, is associated with allergic responses. The total IgE test is used to diagnose allergic diseases such as atopic disease (Čelakovská & Bukač, 2016). In this study, the total serum IgE concentration was significantly increased by TDI sensitization and significantly decreased in the TDI-sensitized 5% and 10% lotus root powder diet groups (Figure 4).

3.4. Anti-allergic inflammatory parameters

IL-4 was produced by a splenic lymphocyte and leads to the differentiation of B-cells into plasma cells (antibody-producing cells). The serum IL-4 level was no significant difference. However, it was tended to dose-dependent decrease in the TDI-sensitized lotus root powder diet groups compared to the TDI-sensitized standard diet group (Figure 5). These results suggested that the decrease in total IgE level was caused by the decrease in IL-4, which in turn was produced by a splenic lymphocyte.

It is known that $CD4^+$ cells (helper T-cells) and $CD8^+$ cells (suppressor T-cells) can contribute to anti-allergic inflammation. It is well established that $CD4^+$ T-cells are required during sensitization (Corry et al., 1998; Koya et al., 2007, 2009). Our experiments revealed a significant ($P < 0.05$) increase in the $CD4^+$ cell population in the TDI-sensitized 10% lotus root powder diet group compared with the TDI-sensitized standard diet group




Figure 5. IL-4 production by splenic lymphocytes. There is no significant difference. Each value is the mean \pm SEM ($n = 5/\text{group}$).




Figure 6. Ratio of (A) CD4⁺, *significant difference between standard food sensitization group and 10% lotus root powder sensitization group ($P < 0.05$), (B) CD8⁺ to splenic lymphocytes, *significant difference compared to nonsensitization group ($P < 0.05$). Each value is the mean \pm SEM ($n = 5/\text{group}$).

(Figure 6A). The CD8⁺ cell population differed significantly ($P < 0.05$) between the non-TDI-sensitized group and the 10% lotus root powder diet TDI-sensitized group (Figure 6B). Whereas CD4⁺ cells have been demonstrated to have a significant role in the development of allergic diseases, the role of CD8⁺ cells is not clear. Some researchers have failed to find any functional or numerical differences in T-lymphocyte subpopulations in patients with allergic asthma (Schuyler, Gerblich, & Urda, 1985) while others have reported decreased numbers of CD8⁺ T-cells (Jensen, Cramers, & Thestrup-Pedersen, 1981; Kus, Tse, Vedal, & Chan-Yeung, 1985; Strannegard & Strannegard, 1978) or no change in allergic contact dermatitis (Usuda, Fujii, & Nonogaki, 2016). Other groups have found imbalances in the T-cell subpopulations in patients with atopic disease (Canonica, Mingari, Melioli, Colombatti, & Moretta, 1979; Zhang et al., 2017). Moreover, CD4⁺ T-cells/CD8⁺ T-cells was no significant difference between the groups (Figure 7). Based on these results, we hypothesized this mechanism of alleviation nasal allergy by lotus root powder. The lotus root powder could be related to the suppression of IL-4, IgE, and LTB₄ production (Figure 8).

We further measured some factors relating to allergy. The weight of the spleen and the number of splenic lymphocytes did not differ among the groups (data not shown). The CD3⁺ cell (mature T-cell) ratio did not differ significantly among the groups (Figure 9A), whereas the CD45R⁺ cell (B-cell) ratio increased significantly ($P < 0.05$) in the TDI-sensitized 5% and 10% lotus root powder diet group than in the non-TDI-sensitized and TDI-sensitized standard diet groups (Figure 9B).




Figure 7. CD4⁺/CD8⁺, no significant difference between sensitization group. Each value is the mean ± SEM ($n = 4$ or 5 /group).




Figure 8. The mechanism of alleviation nasal allergy by lotus root powder. The lotus root powder could be related to the suppression of IL-4, IgE, and LTB₄ production.




Figure 9. Ratio of (A) CD3⁺, there are no significant differences, (B) CD45R⁺ to splenic lymphocytes, with differential letters indicating significant differences ($P < 0.05$). Each value is the mean \pm SEM ($n = 5$ /group).

4. Conclusion

Our study revealed that TDI-sensitized nasal allergic symptoms in mice, such as hair loss and redness around the nose, were prominently alleviated by the intake of lotus root powder. The results suggested that the alleviation of allergic symptoms by lotus root powder intake could be related to the suppression of IL-4, IgE, and LTB₄ production. These results suggest that lotus root can be utilized to treat allergic symptoms such as hay fever or food allergy.

Acknowledgments

We thank Dr. Satoru Moriguchi for his helpful advice on this experiment. We thank Ms. Junka Oguchi for her technical support of our experimental analysis. We also would like to thank Dr. Wataru Mizunoya and Dr. Shoko Sawano for their valuable and constructive writing assistance.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- Canonica, G. W., Mingari, M. C., Melioli, G., Colombatti, M., & Moretta, L. (1979). Imbalances of T cell subpopulations in patients with atopic diseases and effect of specific immunotherapy. *The Journal of Immunology*, 123(6), 2669–2672.
- Čelakovská, J., & Bukač, J. (2016). Eosinophils in patients suffering from atopic dermatitis and the relation to the occurrence of food allergy and other atopic diseases. *Food and Agricultural Immunology*, 27(5), 700–710.
- Corry, D. B., Grünig, G., Hadeiba, H., Kurup, V. P., Warnock, M. L., Sheppard, D., & Locksley, R. M. (1998). Requirements for allergen-induced airway hyperreactivity in T and B cell-deficient mice. *Molecular Medicine*, 4(5), 344–355.
- Dearman, R., Basketter, D., & Kimber, I. (1996). Characterization of chemical allergens as a function of divergent cytokine secretion profiles induced in mice. *Toxicology and Applied Pharmacology*, 138(2), 308–316.
- Huang, B., He, J., Ban, X., Zeng, H., Yao, X., & Wang, Y. (2011). Antioxidant activity of bovine and porcine meat treated with extracts from edible lotus (*Nelumbo nucifera*) rhizome knot and leaf. *Meat Science*, 87(1), 46–53.
- Jensen, J., Cramers, M., & Thestrup-Pedersen, K. (1981). Subpopulations of T lymphocytes and non-specific suppressor cell activity in patients with atopic dermatitis. *Clinical and Experimental Immunology*, 45(1), 118.
- Khan, R. A., Rajput, M. A., & Assad, T. (2019). Effect of *Nelumbo nucifera* fruit on scopolamine induced memory deficits and motor coordination. *Metabolic Brain Disease*, 34(1), 87–92.
- Koya, T., Matsuda, H., Matsubara, S., Miyahara, N., Dakhama, A., Takeda, K., & Gelfand, E. W. (2009). Differential effects of dendritic cell transfer on airway hyperresponsiveness and inflammation. *American Journal of Respiratory Cell and Molecular Biology*, 41(3), 271–280.
- Koya, T., Miyahara, N., Takeda, K., Matsubara, S., Matsuda, H., Swasey, C., ... Gelfand, E. W. (2007). CD8+ t cell-mediated airway hyperresponsiveness and inflammation is dependent on CD4+ IL-4+ T cells. *The Journal of Immunology*, 179(5), 2787–2796.
- Kus, J., Tse, K. S., Vedal, S., & Chan-Yeung, M. (1985). Lymphocyte sub-populations in patients with allergic and non-allergic asthma. *Clinical & Experimental Allergy*, 15(6), 523–529.
- Miyake, K., & Karasuyama, H. (2017). Emerging roles of basophils in allergic inflammation. *Allergy International*, 66, 382–391.
- Mukherjee, P. K., Mukherjee, D., Maji, A. K., Rai, S., & Heinrich, M. (2009). The sacred lotus (*Nelumbo nucifera*)—phytochemical and therapeutic profile. *Journal of Pharmacy and Pharmacology*, 61(4), 407–422.
- Mukherjee, P. K., Saha, K., Pal, M., & Saha, B. (1997). Effect of *Nelumbo nucifera* rhizome extract on blood sugar level in rats. *Journal of Ethnopharmacology*, 58(3), 207–213.
- Ohnishi, H., Miyahara, N., & Gelfand, E. W. (2008). The role of leukotriene B4 in allergic diseases. *Allergy International*, 57(4), 291–298.
- Okubo, K., Kurono, Y., Ichimura, K., Enomoto, T., Okamoto, Y., Kawauchi, H., ... Masuyama, K. (2017). Japanese guidelines for allergic rhinitis 2017. *Allergy International*, 66, 205–219.
- Rajput, M. A., Khan, R. A., & Assad, T. (2017). Anti-epileptic activity of *Nelumbo nucifera* fruit. *Metabolic Brain Disease*, 32(6), 1883–1887.
- Schuyler, M., Gerblich, A., & Urda, G. (1985). Atopic asthma: T lymphocyte subpopulations. *Clinical & Experimental Allergy*, 15(2), 131–138.
- Sridhar, K., & Bhat, R. (2007). Lotus-A potential nutraceutical source. *Journal of Agricultural Technology*, 3(1), 143–155.
- Strannegard, Ö, & Strannegard, I. L. (1978). T lymphocyte numbers and function in human IgE-mediated allergy. *Immunological Reviews*, 41(1), 149–170.

- Usuda, H., Fujii, H., & Nonogaki, T. (2016). Sasa veitchii extracts suppress 2,4- dinitrofluorobenzene-induced contact hypersensitivity in mice. *Food and Agricultural Immunology*, 27(4), 523–534.
- Wang, L., Yen, J.-H., Liang, H.-L., & Wu, M.-J. (2003). Antioxidant effect of methanol extracts from lotus plumule and blossom (*Nelumbo nucifera* Gertn.). *Journal of Food and Drug Analysis*, 11(1), 60–66.
- Wedi, B., Novacovic, V., Koerner, M., & Kapp, A. (1999). Chronic urticaria serum induces histamine release, leukotriene production, and basophil CD63 surface expression-Inhibitory effects of anti-inflammatory drugs. *Journal of Allergy and Clinical Immunology*, 105(3), 552–560.
- Yan, S.-L., Wang, Q.-Z., & Peng, G.-H. (2009). Determination of catechin in lotus rhizomes by high-performance liquid chromatography. *International Journal of Food Sciences and Nutrition*, 60(5), 432–438.
- Zhang, Z., Zheng, W., Xie, H., Chai, R., Wang, J., Zhang, H., & He, S. (2017). Up-regulated expression of substance P in CD8+ T cells and NK1R on monocytes of atopic dermatitis. *Journal of Translational Medicine*, 15(1), 93.