



Forensic dissection of lip print as an investigative tool in a mixed Egyptian population

S.A. Ahmed, Hanan E. Salem & Marwa M. Fawzy

To cite this article: S.A. Ahmed, Hanan E. Salem & Marwa M. Fawzy (2018) Forensic dissection of lip print as an investigative tool in a mixed Egyptian population, Alexandria Journal of Medicine, 54:3, 235-239, DOI: [10.1016/j.ajme.2017.08.002](https://doi.org/10.1016/j.ajme.2017.08.002)

To link to this article: <https://doi.org/10.1016/j.ajme.2017.08.002>



© 2017 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V.



Published online: 17 May 2019.



Submit your article to this journal [↗](#)



Article views: 269



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Alexandria Journal of Medicine

journal homepage: <http://www.elsevier.com/locate/ajme>

Original Article

Forensic dissection of lip print as an investigative tool in a mixed Egyptian population



S.A. Ahmed*, Hanan E. Salem, Marwa M. Fawzy

Ain Shams University, Faculty of Medicine, Department of Forensic Medicine and Clinical Toxicology, Egypt

ARTICLE INFO

Article history:

Received 21 June 2017

Revised 4 July 2017

Accepted 2 August 2017

Available online 16 August 2017

Keywords:

Lip print

Egypt

Gender

Geographical distribution

Identification

Anthropology

Gender difference

ABSTRACT

Background: Identification is a major problem facing forensic practitioners, DNA and finger prints are highly useful but sometimes aren't easily collected from the crime scene. Lip print could be useful in this field being unique to each individual. Aim: The current study aimed at detecting the frequency and gender relation of lip print pattern in an Egyptian sample. Methodology: Samples were collected on white copy paper, divided into four quadrants then examined with magnifying lens for pattern distribution. Results: The study showed that pattern IV was the most frequently represented pattern in the study sample, pattern I & II were more prevalent in males and females respectively. Prevalent pattern in Cairo and Lower Egypt was I while it was IV in Upper Egypt. Conclusion: The lip print pattern can differ due to gender and geographical origin in Egyptian subjects.

© 2017 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

In various civil, criminal and mass disaster cases, positive identification of unrecognizable deceased individuals could be very difficult. Out of the many existing techniques used for this purpose; comparison of fingerprints, DNA and dental records are probably the most common techniques applied. However, human lip print recognition is one of the most interesting emerging fields in confirming personal identification.¹

Lip prints are defined as normal wrinkles and grooves present in the zone of transition of human lip, between the inner labial mucosa and outer skin, examination of which is known as Cheiloscopia. Those patterns are identifiable as early as the sixth week of intrauterine life. Lip print gained its importance being a unique feature for each individual, as finger prints. Research studies regarding the use of lip prints as evidence in personal identification and criminal investigation are scanty. However, studying it in depth and establishing further information concerning lip prints will certainly help as useful evidence in forensic practice.²

Fischer, 1902 was among the first to take notice of the biological phenomenon of systems of furrows on the red part of human lips in the year 1902. While the use of lip prints in personal identification and criminalization was first recommended in France by Edmond Locard as early as 1932. Le Moyne Snyder was the first to introduce a case in which lip prints helped the crime investigators in an unusual way. Suzuki and Tsuchihashi were among the first to classify the various patterns present on the human lips.³

Lip prints could be left at crime scenes on various objects, e.g. drinking glasses, cigarette butts, and duct tape. Both direct inspection and photography allow more precise and detailed observations needed for lip print investigations. Proper investigation of the individualizing characters of lip prints is essential to identify suspects and further confirm their presence in crime scene.⁴

The oily and moist secretions from sebaceous and salivary glands located at the vermillion border (which is the sharp demarcation between the lip and the adjacent normal skin) and subsequent moisturization from the tongue enables the formation of a latent lip print whenever there is contact with lips leaving behind an important form of transfer evidence. A lip print at the scene of crime can offer a clue for different questions as the type of the crime, the number of the people involved, sexes of suspects, cosmetics used, habits, occupational traits, and the pathological changes of lips themselves.⁵

Peer review under responsibility of Alexandria University Faculty of Medicine.

* Corresponding author at: Abbassia Square, Faculty of Medicine, ASU Cairo, Egypt.

E-mail address: drsamer_ahmed@med.asu.edu.eg (S.A. Ahmed).

<http://dx.doi.org/10.1016/j.ajme.2017.08.002>

2090-5068/© 2017 Alexandria University Faculty of Medicine. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Lip groove patterns rarely change, resisting numerous types of external factors including physical injury, pressure and exposure to hot or cold stimuli. Its permanence enables it to be a reliable tool in forensic field investigations. Especially when combined with other evidence, it greatly adds to the success of crime solving mystery.⁶

Studies related to geographical origin are always an important step for confirming identification. There were no enough data to correlate sex and geographical distribution to lip pattern. Our study aimed to detect lip pattern among some Egyptians of both sexes and of different geographical distribution.

2. Method

2.1. Subjects

The collected specimens in the research belonged to 221 randomly selected adult Egyptian volunteers of both sexes and of various geographical origin.

Exclusion criteria: Any participant suffering from an inflammation, scars, deformity or past history of plastic surgery in the lips was excluded from the study.

Ethical considerations:

- An informed valid consent was taken from each participant.
- An approval was obtained from Ethics committee.
- In order to secure confidentiality, specimens were coded and anonymously stored.

2.2. Recording the lip prints

Dark Red or brown, non-persistent, non-glossy, non-metallic lipstick was used to get clear lip prints. White paper (white A4 Globules classic 80-g copy paper) was used to take the impressions of the lips. A thin film of lipstick was applied onto cleaned and dried lips, left for 5 min, and then the impressions of the lips were taken on the specified papers. The following method was used for taking the impressions from every subject. Direct light pressure was applied by the lips on a standardized properly folded paper then they were stored in a card board box under suitable environmental conditions.

2.3. Examination of the prints

Each lip print was divided into four topographic areas (QI, QII, QIII, QIV) after a study conducted by Santos in 1967,¹ examined by magnifying hand lens with direct light focused on it. The lip print was classified into 4 patterns according to K. Suzuki and Y. Tsuchihashi's classification.³ Figs. 1–4.⁷



Fig. 1. Pattern I: Vertical.



Fig. 2. Pattern II: Branched.



Fig. 3. Pattern III: Intersecting.



Fig. 4. Pattern IV: Reticulate.

3. Results

In our study 221 patients were enrolled. Lip print was studied from 4 lip quadrants yielding 884 samples, 4 types of patterns were analyzed (I–IV) (see Table 1).




Fig. 5. Column chart showing distribution of different lip patterns as regards gender distribution among Egyptian participants involved in the study.




Fig. 6. Column chart showing distribution of different lip pattern as regards origin distribution among Egyptian participants involved in the study.

4. Discussion

Usually personal identification is made through comparing AM and PM records, so if there is a detailed AM record of different parts of upper and lower lips, it could open door to personal identification through thorough examination of lip pattern.²

The research studies and data regarding lip print pattern use in identification are so scanty. The current study was established in order to provide more details regarding the usefulness of lip print pattern, showing its distribution among a sample of Egyptian population. To our knowledge the present study is the first in Egypt to describe lip print pattern among different geographical areas.

It was noticed from the current study that no similar pattern was shown between any two individuals and this was in coincidence with Suzuki et al.³ and Kumar et al.², which supports the role of lip print in personal identification being unique to each person.

The present study showed that the most frequent pattern among persons enrolled in it was pattern IV (Table 2) and this agrees with a study done by Prabhu et al.⁸ which was done on 100 individuals with the aim of extensive examination of lip print patterns with referral to the frequency of each pattern among them.

Sivapathasundharam et al.⁹ reached the same result in their study, confirming that the most prominent pattern was IV. Contrasting a study by Kapoor and Badiye¹⁰ showed that the predominant pattern in Indian sample population of 200 persons was pattern I, this could be attributed to the fact of different location and races of involved populations that might be related to genetic factors.

Prabhu et al.⁸ explained that this variation could be due to geographical variation or strict standards of different pattern identification.

Table 1

Percent (%) of demographic data of the Egyptian participants involved in the study: (N = 221).

Age (Years)		26.8 ± 10.4
Gender	Male	105 (47.5%)
	Female	116 (52.5%)
Origin	Cairo	81 (36.5%)
	Lower Egypt	55 (24.9%)
	Upper Egypt	85 (38.5%)

Table 2

Frequencies of different lip patterns among the Egyptian participants involved in the study: (N = 221).

Pattern	Frequency
I	230 (26%)
II	213 (24.1%)
III	208 (23.5%)
IV	233 (26.4%)

As regards the predominant pattern among both sexes (Table 3 and Fig. 5) that the prevalent pattern in males was pattern I while it was pattern III in females with no statistically significant difference between patterns in both sexes. Non significance could have occurred due to small sample size, so larger samples might be needed to clarify degree of significance (see Table 4).

This coincides with a study in India by Badiye and Kapoor,¹⁰ but differs from the study by Krishnan et al.¹¹ which was focusing on the role of lip prints, finger prints for gender differentiation and showed there was a relationship between lip pattern and sex but the predominant pattern in their study was pattern IV in males and I in females.

This agrees with Prabhu et al.⁸ who showed in their study that there is no correlation between lip pattern and gender as pattern V was predominant in both sexes. Also a study in Dakahlia (Egypt) showed that the prevalent pattern in both sexes was pattern I negating the correlation between lip pattern and sex.⁴

Regarding the geographical distribution of different lip pattern in Egyptian population selected sample (Table 2 & Fig. 6), the predominant pattern in Cairo and lower Egypt was pattern I and this could be attributed to the fact that people tend to leave lower Egypt and settle in Cairo for better living conditions, while the most prevalent pattern in upper Egypt was IV with statistically highly significant difference between different patterns among different geographical origins.

This could be attributed to the closed nature of Upper Egypt where most inhabitants are in closed families that tend to marry from each other and have limited internal immigration to and from other areas in Egypt.




Fig. 7. Column chart showing distribution of different patterns by different quadrants among Egyptian participants involved in the study.

Table 3Chi square (X^2) analysis of different lip patterns as regards gender distribution among Egyptian participants involved in the study.

Pattern		Gender				Total		X^2	P Value
		F		M		Count	%		
		Count	%	Count	%				
I	111	23.9%	119	28.3%	230	26.0%	6.8	0.078	
II	109	23.5%	104	24.8%	213	24.1%			
III	125	26.9%	83	19.8%	208	23.5%			
IV	119	25.6%	114	27.1%	233	26.4%			

P. value = significant < 0.05, non significant > 0.05.

Table 4Chi square (X^2) analysis of different lip patterns as regards origin distribution among Egyptian participants involved in the study.

Pattern		Origin						Total		X^2	P Value
		Cairo		Lower Egypt		Upper Egypt		Count	%		
		Count	%	Count	%	Count	%				
I	89	27.5%	66	30.0%	75	22.1%	230	26.0%	13.34	0.008	
II	88	27.2%	56	25.5%	69	20.3%	213	24.1%			
III	69	21.3%	54	24.5%	85	25.0%	208	23.5%			
IV	78	24.1%	44	20.0%	111	32.6%	233	26.4%			

P. value = significant < 0.05, non significant > 0.05.

Table 5Chi square (X^2) analysis of different patterns distribution by different quadrants among Egyptian participants involved in the study:

Pattern		Quadrants								Total		X^2	P Value
		Quadrant I		Quadrant II		Quadrant III		Quadrant IV		Count	%		
		Count	%	Count	%	Count	%	Count	%				
I	63	28.5%	59	26.7%	55	24.9%	53	24.0%	230	26.0%	4.97	0.837	
II	56	25.3%	51	23.1%	56	25.3%	50	22.6%	213	24.1%			
III	53	24.0%	52	23.5%	46	20.8%	57	25.8%	208	23.5%			
IV	49	22.2%	59	26.7%	64	29.0%	61	27.6%	233	26.4%			

P. value = significant < 0.05, non significant > 0.05.

On the other hand a study by Ragab et al.⁴ in Dakahlia, showed that the prevalent pattern was pattern I but this could be related to the fact that Dakahlia is a part of upper Egypt and their sample didn't show variation as ours as we included persons from different places of upper Egypt.

In the current study the prevalent pattern in each quadrant (Table 5 & Fig. 7) quadrant QI: pattern I, quadrant QII: pattern I and IV, quadrant QIII & QIV: pattern IV, with no statistical significant difference between different patterns in different quadrants.

This partially goes in hand with a study by Chimere et al.¹² on Nepalese population where the most prevalent pattern in quadrants QI & QII was I, but differs in that pattern I was also prominent in quadrant QIII & QIV in their study while in ours the most prominent in quadrant QIII & QIV was pattern IV.

This also disagrees with Amith et al.¹² where quadrant QIII & QIV showed pattern II as the dominant pattern. Also current results differs from a study by Nagrate et al.¹³ where dominant pattern in quadrants QI, QII, QIII & QIV were II, III, III and II respectively. These various results could be attributed to the difference of population

Table 6

ANOVA statistical analysis of relation between age and different lip patterns among Egyptian participants involved in the study.

Pattern		Mean	SD	F	P Value
I		26.60	10.675	0.831	0.477
II		27.11	11.041		
III		26.03	9.292		
IV		27.52	10.549		

P. value = significant < 0.05, non significant > 0.05.

dominance of particular lip print and ethnicity of subjects recruited.¹⁴

There was no significant difference in pattern distribution among different ages (Table 6) which supports the fact that lip print pattern is stable across years. This was confirmed by a study by Nagrale et al.¹³ in India, who concluded that features of lip print relative to surrounding anatomy was proved to be stable, recordable emphasizing on its value as a forensic tool.

A study by Randhawa et al.⁷ threw the light on limitation and changes that occur with advancing age especially after 40 years, due to thinning of lips, perioral area age changes and decrease in lip height. These can influence lip print pattern decreasing its efficacy in identification.

5. Conclusions

1. Lip print can be used among Egyptian subjects for individual identification.
2. There is a variation in lip print based on individual origin and gender.

6. Recommendations

- Further studies on larger sample size of different races for better comparison and could help to create a comprehensive database so that the lip print pattern might serve as an important source of information needed for crime solving and personal identification.

- Digital examination of lip print pattern could be more helpful in other studies.
- Extensive longitudinal study is needed to monitor the changes of lip pattern across years.
- New techniques (e.g. magna brush technique, cobalt oxide) have been evolved for tracing latent lip print in crime scene so further studies are needed to assess feasibility and validity of these new techniques.

Funding

This research didn't receive any special grant from funding agencies in the public commercial or not-for-profit sectors; it was totally and personally funded by authors.

References

1. Reddy LVK. Lip prints: an overview in forensic dentistry. 1st ed. J Adv Dent Res. 2011.
2. Kumar LB, Jayaraman V, Mathew P, Ramasamy S, Austin RD. Reliability of lip prints in personal identification: an inter-racial pilot study. *J Forensic Dent Sci.* 2016;8:178.
3. Suzuki K, Tsuchihashi Y. New attempt of personal identification by means of lip print. *J Indian Dent Assoc.* 1970;42:8–9.
4. Ragab AR, El-Dakroory SA, Rahman RH. Characteristic patterns of lip prints in Egyptian population sample at Dakahlia Governorate. *Int J Legal Med.* 2013;127:521–527.
5. Alzapur A, Nagothu RS, Nalluri HB. Lip prints – a study of its uniqueness among students of MediCiti Medical College. *Indian J Clin Anat Physiol.* 2017;4:68–70.
6. Venkatesh R, David MP. Cheiloscopy: an aid for personal identification. *J Forensic Dent Sci.* 2011;3:67–70.
7. Randhawa K, Narang RS, Arora PC. Study of the effect of age changes on lip print pattern and its reliability in sex determination. *J Forensic Odontostomatol.* 2011;29:45–51.
8. Prabhu RV, Dinkar A, Prabhu V. A study of lip print pattern in Goan dental students – a digital approach. *J Forensic Leg Med.* 2012;19:390–395.
9. Sivapathasundharam B, Prakash PA, Sivakumar G. Lip prints (cheiloscopy). *Indian J Dent Res.* 2001;12:234–237.
10. Kapoor N, Badiye A. A study of distribution, sex differences and stability of lip print patterns in an Indian population. *Saudi J Biol Sci.* 2015;112–131.
11. Krishnan RP, Thangavelu R, Rathnavelu V, Narasimhan M. Gender determination: role of lip prints, finger prints and mandibular canine index. *Exp Ther Med.* 2016;11:2329–2332.
12. Amith HV, Nagesh L, Ankola AV. Lip prints-can it aid in individual identification. *Health Renaissance: Nepal J Online.* 2011;113–118.
13. Nagrale N, Tirpude Bipinchandra, Murkey P, Patond S. Establishing cheiloscopy as a tool for identification: an assessment on 500 subjects in central India. *Al Ameen J Med Sci.* 2014;201–204.
14. Abdel Aziz MH, Badr El Dine FM, Saeed NM. Regression equations for sex and population detection using the lip print pattern among Egyptian and Malaysian adult. *J Forensic Leg Med.* 2016;44:103–110.