

Cytotoxicity of gamma irradiated aflatoxin B₁ and ochratoxin A

Ana-Marija Domijan, Ana Marija Marjanović Čermak, Ana Vulić, Ivana Tartaro Bujak, Ivan Pavičić, Jelka Pleadin, Ksenija Markov & Branka Mihaljević

To cite this article: Ana-Marija Domijan, Ana Marija Marjanović Čermak, Ana Vulić, Ivana Tartaro Bujak, Ivan Pavičić, Jelka Pleadin, Ksenija Markov & Branka Mihaljević (2019) Cytotoxicity of gamma irradiated aflatoxin B₁ and ochratoxin A, Journal of Environmental Science and Health, Part B, 54:3, 155-162, DOI: [10.1080/03601234.2018.1536578](https://doi.org/10.1080/03601234.2018.1536578)

To link to this article: <https://doi.org/10.1080/03601234.2018.1536578>



© 2018 The Author(s). Published with license by Taylor & Francis Group, LLC



Published online: 05 Jan 2019.



Submit your article to this journal [↗](#)



Article views: 1300



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 7 View citing articles [↗](#)

Cytotoxicity of gamma irradiated aflatoxin B₁ and ochratoxin A

Ana-Marija Domijan^a, Ana Marija Marjanović Čermak^b , Ana Vulić^c, Ivana Tartaro Bujak^d, Ivan Pavičić^b, Jelka Pleadin^c, Ksenija Markov^e, and Branka Mihaljević^d 

^aFaculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia; ^bInstitute for Medical Research and Occupational Health, Zagreb, Croatia; ^cCroatian Veterinary Institute, Zagreb, Croatia; ^dRuder Bošković Institute, Zagreb, Croatia; ^eFaculty of Food Technology and Biotechnology, University of Zagreb, Zagreb, Croatia

ABSTRACT

Toxicity of gamma irradiated mycotoxins aflatoxin B₁ (AFB₁) and ochratoxin A (OTA) was investigated *in vitro*. AFB₁ and OTA stock solutions (50 mM, in methanol) were gamma irradiated (5 and 10 kGy) and non-irradiated and irradiated mycotoxins solutions were tested for cytotoxicity on Pk15, HepG2 and SH-SY5Y cell lines (MTT assay, 1–500 μM concentration range; 24 h exposure). Degradation of mycotoxin molecules was examined by liquid chromatography tandem mass spectrometry (HPLC-MS/MS). AFB₁ and OTA radiolytic products were less toxic than the parent mycotoxins to all of the tested cell lines. Gamma irradiation even at 5 kGy had effect on AFB₁ and OTA molecules however, this effect was dependent on chemical structure of mycotoxin. Since gamma irradiation at low dose reduced initial level of both mycotoxins, and gamma irradiated mycotoxins had lower toxicity in comparison to non-irradiated mycotoxins, it can be concluded that gamma irradiation could be used as decontamination method.

ARTICLE HISTORY

Received 19 May 2018
Accepted 11 October 2018

KEYWORDS

Cytotoxicity; food safety; gamma irradiation; mycotoxins; reduction



Introduction

Mycotoxins aflatoxin B₁ (AFB₁) and ochratoxin A (OTA) have been found as contaminants of various commodities all over the world.^[1–3] Recent review article reported AFB₁ concentrations in foodstuffs of up to 42 and 90 μg/kg in Africa and Asia, respectively, and OTA of up to 9 μg/kg in Central and South America and Europe, and 15 μg/kg in Asia.^[3] Toxicity and carcinogenicity of AFB₁ and OTA to experimental animals are well documented.^[4] Due to proven carcinogenicity to humans, International Agency for Research on Cancer (IARC) classified AFB₁ as a Group 1 carcinogen,^[5] while OTA is classified into group 2B (possible human carcinogen).^[6] To protect human and animal health the governments around the world set specific regulations for acceptable concentrations of mycotoxins in food and feed. In the European Union maximum levels and guidance values for mycotoxins in food intended for human consumption have been set by Commission Regulation (EU) 1881/2006 and Commission Recommendation No 2013/165/EU.^[7,8] According to them for AFB₁ maximum level in all cereals and all products derived from cereals is set to 2.0 μg/kg, while for OTA maximum level in all products derived from unprocessed cereals, including processed cereal products and cereals intended for direct human consumption is set to 3.0 μg/kg.^[7]

To reduce mycotoxin contamination and consequently human and animal exposure to them, several studies tested

gamma irradiation as decontamination method.^[9–14] Gamma irradiation is a physical method based on the ability of high-energy photons generated by radiation of ⁶⁰Co to induce chemical changes of target molecule by causing its degradation or remodelling. In mycotoxin molecules the most prone to the interaction with high-energy photons generated by radiation of ⁶⁰Co are aromatic or heterocyclic rings. Studies so far indicate that gamma irradiation can reduce mycotoxin levels in various commodities. Gamma irradiation at low dose (up to 6 kGy) reduced for around 90% AFB₁ level in fruit and red chillies samples,^[9,12] and dose of 10 kGy in maize seed samples reduced AFB₁ level for around 95%.^[14] Similarly, gamma irradiation at dose of 10 kGy almost completely degraded OTA in coffee beans,^[11] and same dose (10 kGy) in dry-cured meat products reduced OTA for around 22.5%.^[13]

Although studies indicate that gamma irradiation could be suitable decontamination method, data on toxicity of radiolytic products of mycotoxins (products formed by gamma irradiation of mycotoxins) are lacking. Therefore, the aim of this study was to test the toxicity of the parent mycotoxin and its radiolytic products on three cell lines, Pk15, HepG2 and SH-SY5Y in order to mimic different target organs (kidney, liver and brain). Additionally, degradation of mycotoxin molecules was monitored by HPLC-MS/MS.

CONTACT Ana-Marija Domijan  adomijan@pharma.hr  Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia.

© 2018 The Author(s). Published with license by Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Materials and methods

Gamma irradiation of mycotoxin stock solutions

Mycotoxins AFB₁ and OTA (Sigma-Aldrich, St. Louis, MO, USA) stock solutions at 50 mM were prepared in methanol and aliquoted in amber glass vials. For each mycotoxin, one sample was not irradiated (non-irradiated sample, positive mycotoxin control) and two samples were subjected to gamma irradiation.

Mycotoxin stock solutions (in closed amber glass vials) were gamma irradiated in a panoramic ⁶⁰Co source at the Radiation Chemistry and Dosimetry Laboratory, Ruđer Bošković Institute (Zagreb, Croatia). The radiation doses were 5 and 10 kGy and the dose rate was 140 Gy/min. The dose rate was established using an ethanol-chlorobenzene dosimetry system^[15] and calculated daily taking into account the radioactive decay of ⁶⁰Co. The irradiation was performed at ambient temperature (about 18 °C) and ambient atmosphere of a gamma chamber.

HPLC-MS/MS analysis

HPLC-MS/MS analysis of non-irradiated and irradiated AFB₁ and OTA stock solutions was performed using an HPLC (Infinity 1260, Agilent, Santa Clara, USA) coupled with a triple quadrupole mass detector (6410, Agilent, Santa Clara, USA). The mobile phase consisted of 0.1% formic acid (in water) and acetonitrile, and the injection volume was 1 µL. The mass spectrometry conditions were as follows: electrospray ionisation (ESI), positive polarity, fragmentor voltage 150 and 110 V for AFB₁ and OTA, respectively, capillary voltage 6 kV (+), source temperature 350 °C, nebuliser 45 psi and gas flow 5 L/min. The mass spectrometer was operated in MS2 scan (full scan mode). Protonated molecular ions of AFB₁ at *m/z* = 313 and OTA at *m/z* = 404 were monitored.

Cell viability

Cell viability was determined by 3-(4,5-dimethylthiazolyl-2)-2,5-diphenyltetrazoliumbromide (MTT assay; Sigma-Aldrich St. Louis, MO, USA), using porcine kidney epithelial (Pk15), human hepatoma (HepG2) and human neuroblastoma (SH-SY5Y) cells. Cells were seeded in 96-well tissue culture plates at a planting density of 8×10^4 cells/mL and maintained at 37 °C in a humidified atmosphere of 5% CO₂ and 95% of air. Cells were treated with non-irradiated AFB₁ or OTA (mycotoxin positive control) or with AFB₁ or OTA irradiated at 5 or 10 kGy at final concentrations of 1–500 µM for 24 h. AFB₁ and OTA stock solutions were prepared in methanol; by diluting stock solutions in PBS buffer, the final methanol concentration in the mycotoxin solutions for cell treatment was less than 1% and did not affect cell viability. In each experiment, a negative control (PBS-buffer treated) and an MTT-positive control were included. After the treatment, the cell medium was removed and cells incubated with MTT solution at a final concentration of 0.5 mg/mL. After incubation (3 h, at 37 °C in an atmosphere of 5% CO₂

and 95% of air), formazan crystals were dissolved in dimethyl sulfoxide (DMSO) and absorbance was read at 570 nm on a multilabel plate reader (VictorTM, PerkinElmer, Waltham, MA, USA).

Statistical analysis

The results of cell viability of two independent sets of experiments (each with minimum six replicates) are expressed as mean percentage (%) ± SD, compared to negative control (PBS-treated cells, set at 100%). The differences among treatments were tested by one-way analysis of variance (ANOVA) followed by a *post hoc* Tukey test using Statistica software version 10.0 (StatSoft Inc. 1984–2011, USA). The statistical significance was set at 95% (*P* = 0.05).

Results and discussion

Cytotoxicity of AFB₁ and OTA

AFB₁ was cytotoxic to all of the tested cell lines, SH-SY5Y, Pk15 and HepG2 (Fig. 1). These results are in agreement with previous studies that showed decreased cell survival after treatment with AFB₁.^[16–19] Cytotoxicity of AFB₁ to SH-SY5Y was recorded at 100 µM (reduction of cell viability for $16 \pm 1.2\%$; *P* < 0.05, compared to negative control; Fig. 1c), while AFB₁ was cytotoxic to Pk15 only at 500 µM (reduction of cell viability for $35 \pm 8.9\%$; *P* < 0.05, compared to negative control; Fig. 1a). As expected, HepG2 cells were the most sensitive to AFB₁ treatment. A decrease in HepG2 viability (by $14 \pm 7.8\%$; *P* < 0.05, compared to negative control; Fig. 1b) was observed even after the lowest applied AFB₁ concentration (1 µM). AFB₁ is a well-documented hepatocarcinogen whose toxicity is mainly connected to its metabolic activation and genotoxic metabolite formation. Since AFB₁ is predominantly metabolised in the liver, the human hepatoma HepG2 cell model is considered to be the most suitable system for testing its *in vitro* toxicity. This system represents liver cells and includes a wide spectrum of phase I and II enzyme activities enabling them to mimic metabolic conditions of xenobiotics in the living organism.^[20,21] The results on cell viability observed in our study after treatment with AFB₁ differ from studies by Costa et al.,^[22] Ghaderi et al.^[23] and Liu et al.^[24] This could be attributed to different cell viability tests, exposure conditions, culture medium as well as supplements added to the medium. However, the results are comparable to the study by Curcuera et al.^[25] in which using the same assay (MTT) similar IC₅₀ on HepG2 cells for AFB₁ was obtained (about 100 µM).

In this study OTA induced a decrease in cell viability to all of the tested cell lines. The cytotoxicity of OTA to Pk-15, HepG2 and SH-SY5Y was observed at a concentration of 50 µM (Fig. 2). However, a more pronounced toxic effect of OTA (at concentration 50 µM) was recorded on SH-SY5Y cells (reduction of cell viability for $31 \pm 3.6\%$, *P* < 0.05, compared to negative control; Fig. 2c) and HepG2 cells (reduction of cell viability for $25 \pm 3.1\%$; *P* < 0.05, compared to negative control; Fig. 2b), while kidney cells (Pk 15) seemed




Figure 1. Cell viability of: (A) Pk15 cells, (B) HepG2 cells, and (C) SH-SY5Y cells after treatment with non-irradiated aflatoxin B₁ (AFB₁, 1–500 μM, 24 h; solid black columns) or gamma irradiated AFB₁ at 5 or 10 kGy (1–500 μM, 24 h; AFB₁-5 kGy, diagonally striped columns, AFB₁-10 kGy, diagonally crossed columns); cell viability is assessed with MTT test. The results are expressed as mean percentage (%) ± standard deviation, compared to negative control (PBS-treated cells, set at 100%; solid white columns) of two independent set of experiments, each with six replicates; column represent mean value and error bar standard deviation. a - different from negative control (PBS-control); b - different from respective non-irradiated control (AFB₁-treated). One-way ANOVA followed by Tukey test ($P < 0.05$).

to be more resistant (reduction of cell viability for $13 \pm 5.6\%$; $P < 0.05$, compared to negative control; Fig. 2a). These results are in agreement with Zhang et al.^[26] who observed on SH-SY5Y cells and primary neurons OTA toxicity at low concentrations ($0.25 \mu\text{M}$ and $0.5 \mu\text{M}$, respectively). Since their results were comparable to those obtained previously by Hundhausen et al.^[27] on hepatocytes, the authors concluded that neurons could be equally susceptible to OTA as

hepatocytes. Moreover, the same authors, by comparing their results to study conducted on kidney tubulus cells^[28,29] concluded that neurons may be even more susceptible to OTA than kidney cells as is observed in our study. Curcuera et al.^[25] and Gayathari et al.^[30] obtained a similar IC₅₀ for OTA on HepG2 as in our study (360 and $210 \mu\text{M}$, respectively). Additionally, as in our study Curcuera et al.^[25] found that AFB₁ was more toxic to HepG2 than OTA.




Figure 2. Cell viability of: (A) Pk15 cells, (B) HepG2 cells, and (C) SH-SY5Y cells after treatment with non-irradiated ochratoxin A (OTA, 1–500 μM, 24 h; solid black columns) or irradiated OTA at 5 or 10 kGy (1–500 μM, 24 h; OTA-5 kGy, diagonally striped columns, OTA-10 kGy, diagonally crossed columns); cell viability is assessed with MTT assay. The results are expressed as mean percentage (%) ± standard deviation, compared to negative control (PBS-treated cells, set at 100%; solid white columns) of two independent set of experiments each with six replicates; column represent mean value and error bar standard deviation. (a) Different from negative control (PBS-control); (b) Different from respective non-irradiated mycotoxin control (OTA-treated). One-way ANOVA followed by Tukey test (significant at level $P < 0.05$).

Cytotoxicity of gamma irradiated AFB₁ and OTA

Results on the cytotoxicity of gamma irradiated mycotoxins demonstrated that radiolytic products of gamma irradiated AFB₁ and OTA in comparison to the parent mycotoxin were less toxic to all of the cell lines tested. In comparison

to the parent molecules at the same concentration observed reduction in toxicity of gamma irradiated mycotoxins to Pk-15, HepG2 and SH-SY5Y cells was by around 20% (from 5 to 40%) (Figs. 1 and 2). For example, while non-irradiated AFB₁ at 100 μM was cytotoxic to SH-SY5Y cells, AFB₁ at the same concentration (100 μM) irradiated at 10 kGy was




Figure 3. Mass spectra of aflatoxin B₁ (AFB₁) stock solutions (50 mM): (A) non-irradiated, and (B) irradiated at 5 kGy dose.

not toxic to SH-SY5Y cells, i.e. a complete reduction of AFB₁ toxicity (reduction by around 16%; $P < 0.05$, compared to non-irradiated AFB₁; Fig. 1c) was observed. These results clearly indicate that gamma irradiation induced formation of less toxic products than parent mycotoxin. In available literature there are only few studies on toxicity of gamma irradiated mycotoxins. In study of Kumar et al.,^[11] human intestinal epithelial (Int-407) cells were treated with gamma irradiated OTA (at 10 kGy) and by applying the same MTT assay an approximately 7-fold reduction of cytotoxicity of irradiated OTA is observed. In recent study cytotoxicity of gamma irradiated OTA (gamma irradiated to up to 10.3 kGy) was tested on HepG2 cells and 2-fold reduction of OTA cytotoxicity (assessed by neutral red assay) was observed.^[31] The difference in reduction of cytotoxicity of gamma irradiated OTA in above mentioned studies (7- and 2- fold) and in our study (only 20%) can be explained by the process of gamma irradiation. Gamma irradiation is more effective in reducing mycotoxin level when mycotoxin is present in lower concentration and when sample has higher moisture content.^[10,32,33] Although gamma irradiation of OTA for *in vitro* testing in the study of Kumar et al.^[11] and Calado et al.^[31] is not clearly described, it could be that they gamma irradiated OTA at lower concentration.

Results of HPLC-MS/MS analysis of non-irradiated and gamma irradiated mycotoxin solutions in our study revealed

that gamma irradiation even at 5 kGy had impact on both, AFB₁ and OTA. The signal intensity of AFB₁ (AFB₁ ion at 313.2 *m/z*) in AFB₁ stock solution gamma irradiated at 5 kGy was reduced around 16 times compared to non-irradiated AFB₁ stock solution (0.2×10^6 vs. 3.2×10^6 ; Fig. 3). On the other hand, the signal intensity of OTA (OTA ion at 404.2 *m/z*) was reduced only twice (3.2×10^5 vs. 7.2×10^5 ; Fig. 4). In the mass spectra of irradiated AFB₁ solutions ions at 285.2, 345.2, 359.2, and 381.2 *m/z* were observed that could have been the product of AFB₁ degradation or molecule remodelling. Ions pointing to molecule degradation or remodelling were not observed in the mass spectra of the irradiated OTA solution probably due to low concentration of these products reaching the instrument detection limit. The lower level of AFB₁ and OTA detected in irradiated mycotoxin solutions confirmed that gamma irradiation has impact on AFB₁ and OTA. Observed difference in the effect of gamma irradiation on AFB₁ and OTA could be attributed to their chemical structure.

HPLC-MS/MS analysis indicates that AFB₁ molecule is more susceptible to degradation and remodelling induced by high-energy photons generated by ⁶⁰Co than OTA. Different mycotoxin susceptibility to gamma irradiation observed in our study was previously reported by Mutler and Erkoc.^[34] They tested different aflatoxins and found that AFB₁ was the most radiosensitive and aflatoxin B₂ the most resistant to gamma irradiation. The high-energy photons generated




Figure 4. Mass spectra of ochratoxin A (OTA) stock solutions (50 mM): (A) non-irradiated, and (B) irradiated at 5 kGy dose.

by ^{60}Co can affect double bonds in aromatic or heterocyclic rings in mycotoxin's molecule. AFB_1 's superior susceptibility to gamma irradiation could be attributed to feasible addition reaction on the 8,9 double bond in the furan ring initiated by the free radicals generated during gamma irradiation. This observation is confirmed by Wang et al.^[33] In study of Wang et al.^[33] AFB_1 was gamma irradiated (up to 10 kGy) and by HPLC-MS/MS analysis seven radiolytic products of gamma irradiated AFB_1 are identified and six of them were considered less toxic than AFB_1 . In that study the toxicity evaluation was based on the structure–activity relationship (SAR) analysis rather than on testing on live, *in vitro* model as in our study. The authors considered radiolytic product in which the addition reaction occurred on the double bond in the terminal furan ring less toxic than AFB_1 . According to quantitative SAR studies, the furofuran moiety of the AFB_1 structure is essential for its toxicity and carcinogenicity. It is demonstrated that the presence of double bond in the terminal furan ring is important determinant of AFB_1 acute and chronic effects in rats.^[33,35,36] Therefore, in our study the lower toxicity of AFB_1 radiolytic products could be explained by lower unsaturation or the number of double-bonds in comparison to non-irradiated AFB_1 .

This study demonstrated that gamma irradiation (at 5 and 10 kGy) has an impact on both, AFB_1 and OTA

molecules. Radiolytic products formed by gamma irradiation of AFB_1 and OTA are found to be less toxic to cells (Pk15, HepG2, SH-SY5Y) than the parent compounds (non-irradiated mycotoxins). Even low gamma irradiation dose (5 kGy) reduced AFB_1 and OTA levels as detected by HPLC-MS/MS. However, AFB_1 was more susceptible to gamma irradiation than OTA. AFB_1 radiolytic products are probably results of addition reaction on double bond in the terminal furan ring. The mechanism of OTA degradation is more complex since its radiolytic products were not detected. Based on the results of studies examining impact of gamma irradiation on nutritional quality of food^[37–39] international food organisations (FAO/IAEA/WHO)^[40] concluded that gamma irradiation dose of up to 10 kGy has no effects on nutritional quality of food and has no adverse effect on human and animal health. Since in this study radiation up to 10 kGy was applied, gamma irradiation can be considered an effective and safe method for mycotoxin detoxification in the production of food and feed.

Acknowledgements

The authors wish to thank to Mr. Makso Herman for editing the manuscript.

ORCID

Ana Marija Marjanović Čermak  <http://orcid.org/0000-0002-9477-1329>

Branka Mihaljević  <http://orcid.org/0000-0003-3412-9219>

References

- [1] Reddy, K. R. N.; Salleh, B.; Saad, B.; Abbas, H. K.; Abel, C. A.; Shier, W. T. An overview of mycotoxin contamination in foods and its implications for human health. *Toxin. Rev.* **2010**, *29*, 3–26.
- [2] Streit, E.; Schatzmayr, G.; Tassis, P.; Tzika, E.; Marin, D.; Taranu, I.; Tabuc, K.; Nicolau, A.; Aprodu, I.; Puel, O.; Oswald, I. P. Current situation of mycotoxin contamination and co-occurrence in animal feed - focus on Europe. *Toxins* **2012**, *4*, 788–809. doi:10.3390/toxins4100788
- [3] Pinotti, L.; Ottoboni, M.; Giromini, C.; Dell'Orto, V.; Cheli, F. Mycotoxin contamination in the EU feed supply chain: A focus on cereal byproducts. *Toxins* **2016**, *8*, 45. doi:10.3390/toxins8020045
- [4] International Programme on Chemical Safety, IPCS. *Safety evaluation of certain mycotoxins in food*. WHO Food Add Ser. **2001**, *47*, 103–415.
- [5] International Agency for Research on Cancer, IARC. *Some traditional herbal medicines, some mycotoxins, naphthalene, and styrene*. IARC Monographs. Vol. 82, IARC, Lyon, **2002**.
- [6] International Agency for Research on Cancer, IARC. *Some naturally occurring substances: Food items and constituents, heterocyclic aromatic amines and mycotoxins*. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Vol. 56, IARC, Lyon, **1993**.
- [7] Commission Regulation (EU) 1881/2006 of 19 December **2006**. setting maximum levels for certain contaminants in foodstuffs. Off. J. L 364:5–24.
- [8] Commission Recommendation No. Off. J. L 2013, Vol. 91, 12–15. /165/EU of 27 March 2013 on the Presence of T-2 and HT-2 Toxin in Cereals and Cereal Products.
- [9] Aziz, N. H.; Moussa, L. A. A. Influence of gamma-radiation on mycotoxin producing moulds and mycotoxins in fruits. *Food Control* **2002**, *13*, 281–288.
- [10] Jalili, M.; Jinap, S.; Noranizan, M. A. Aflatoxins and ochratoxin a reduction in black and white pepper by gamma radiation. *Rad. Phys. Chem.* **2012**, *81*, 1786–1788.
- [11] Kumar, S.; Kunwar, A.; Gautam, S.; Sharma, A. Inactivation of *A. ochraceus* spores and detoxification of ochratoxin a in coffee beans by gamma irradiation. *J. Food Sci.* **2012**, *77*, 44–51.
- [12] Iqbal, S. Z.; Bhatti, I. A.; Asi, M. R.; Zuber, M.; Shahid, M.; Parveen, I. Effect of γ irradiation on fungal load and aflatoxins reduction in red chillies. *Rad. Phys. Chem.* **2013**, *82*, 80–84.
- [13] Domijan, A.-M.; Pleadin, J.; Mihaljević, B.; Vahčić, N.; Frece, J.; Markov, K. Reduction of ochratoxin a in dry-cured meat products using gamma-irradiation. *Food Addit. Contam. Part A Chem. Anal. Control Expo Risk Assess* **2015**, *32*, 1185–1191.
- [14] Markov, K.; Mihaljević, B.; Domijan, A.-M.; Pleadin, J.; Delaš, F.; Frece, J. Inactivation of aflatoxicogenic fungi and the reduction of aflatoxin B₁ *in vitro* and *in situ* using gamma irradiation. *Food Control* **2015**, *54*, 79–85.
- [15] Ražem, D.; Anđelić, L. J.; Dvornik, I. Consistency of ethanol-chlorobenzene dosimetry. *Proceedings of IAEA Symposium on High-Dose Dosimetry*. IAEA: Vienna, **1984**; 143–156.
- [16] Chan, H. T.; Chan, C.; Ho, J. W. Inhibition of glycyrrhizic acid on aflatoxin B₁-induced cytotoxicity in hepatoma cells. *Toxicology* **2003**, *188*, 211–217.
- [17] Lei, M. M.; Zhang, N.; Qi, D. In vitro investigation of individual and combined cytotoxic effects of aflatoxin B₁ and other selected mycotoxins on the cell line porcine kidney 15. *Exp. Toxicol. Pathol.* **2013**, *65*, 1149–1157.
- [18] Nones, J.; Nones, J.; Trentin, A. G. Flavonoid hesperidin protects neural crest cells from death caused by aflatoxin B(1). *Cell Biol. Int.* **2013**, *37*, 181–186.
- [19] Sun, L. H.; Lei, M. Y.; Zhang, N. Y.; Gao, X.; Li, C.; Krumm, C. S.; Qi, D. S. Individual and combined cytotoxic effects of aflatoxin B₁, zearalenone, deoxynivalenol and fumonisin B₁ on BRL 3A rat liver cells. *Toxicol.* **2015**, *95*, 6–12.
- [20] Cao, J.; Liu, Y.; Jia, L.; Jiang, L. P.; Geng, C. Y.; Yao, X. F.; Kong, Y.; Jiang, B. N.; Zhong, L. F. Curcumin attenuates acrylamide-induced cytotoxicity and genotoxicity in HepG2 cells by ROS scavenging. *J. Agric. Food Chem.* **2008**, *56*, 12059–12063.
- [21] Ehrlich, V.; Darroudi, F.; Uhl, M.; Steinkellner, H.; Zsivkovits, M.; Knasmueller, S. Fumonisin B(1) is genotoxic in human derived hepatoma (HepG2) cells. *Mutagenesis* **2002**, *17*, 257–260.
- [22] Costa, S.; Utan, A.; Speroni, E.; Cervellati, R.; Piva, G.; Prandini, A.; Guerra, M. C. Carnosic acid from rosemary extracts: A potential chemoprotective agent against Aflatoxin B₁. An *in vitro* study. *J. Appl. Toxicol.* **2007**, *27*, 152–159.
- [23] Ghaderi, M.; Allameh, A.; Soleimani, M.; Rastegar, H.; Ahmadi-Ashtiani, H. R. A comparison of DNA damage induced by aflatoxin B₁ in hepatocyte-like cells, their progenitor mesenchymal stem cells and CD34+ cells isolated from umbilical cord blood. *Mut. Res. Gen. Toxicol. Environ. Mutag.* **2011**, *719*, 14–20.
- [24] Liu, Y.; Du, M.; Zhang, G. Proapoptotic activity of Aflatoxin B₁ and sterigmatocystin in HepG2 Cells. *Toxicol. Report* **2014**, *1*, 1076–1086.
- [25] Corcuera, L. A.; Arbillaga, L.; Vettorazzi, A.; Azqueta, A.; López de Cerain, A. Ochratoxin a reduces Aflatoxin B₁ induced DNA damage detected by the comet assay in Hep G2 cells. *Food Chem. Toxicol.* **2011**, *49*, 2883–2889.
- [26] Zhang, X.; Boesch-Saadatmandi, C.; Lou, Y.; Wolffram, S.; Huebbe, P.; Rimbach, G. Ochratoxin a induces apoptosis in neuronal cells. *Genes Nutr.* **2009**, *4*, 41–48.
- [27] Hundhausen, C.; Bosch-Saadatmandi, C.; Augustin, K.; Blank, R.; Wolffram, S.; Rimbach, G. Effect of vitamin E and polyphenols on Ochratoxin A-induced cytotoxicity in liver (HepG2) cells. *J. Plant Physiol.* **2005**, *162*, 818–822.
- [28] Boesch-Saadatmandi, C.; Loboda, A.; Jozkowicz, A.; Huebbe, P.; Blank, R.; Wolffram, S.; Dulak, J.; Rimbach, G. Effect of ochratoxin A on redox-regulated transcription factors, antioxidant enzymes and glutathione-S-transferase in cultured kidney tubulus cells. *Food Chem. Toxicol.* **2008**, *46*, 2665–2671.
- [29] Boesch-Saadatmandi, C.; Wagner, A. E.; Graeser, A. C.; Hundhausen, C.; Wolffram, S.; Rimbach, G. Ochratoxin a impairs Nrf2-dependent gene expression in porcine kidney tubulus cells. *J. Anim. Physiol. Anim. Nutr.* **2009**, *93*, 547–554.
- [30] Gayathri, L.; Dhivya, R.; Dhanasekaran, D.; Periasamy, V. S.; Alshatwi, A. A.; Akbarsha, M. A. Hepatotoxic effect of Ochratoxin A and citrinin, alone and in combination, and protective effect of vitamin E: In vitro study in HepG2 cell. *Food Chem. Toxicol.* **2015**, *83*, 151–163.
- [31] Calado, T.; Fernández-Cruz, M. L.; Cabo Verde, S.; Venâncio, A.; Abrunhosa, L. Gamma irradiation effects on Ochratoxin A: Degradation, cytotoxicity and application in food. *Food Chem.* **2018**, *240*, 463–471.
- [32] Calado, T.; Venancio, A.; Abrunhosa, L. Irradiation for mould and mycotoxin control: A review. *Comp. Rev. Food Sci. Food Saf.* **2014**, *13*, 1049–1061.
- [33] Wang, F.; Xie, F.; Xue, X.; Wang, Z.; Fan, B.; Ha, Y. Structure elucidation and toxicity analyses of the radiolytic products of Aflatoxin B₁ in methanol-water solution. *J. Hazard Mat.* **2011**, *192*, 1192–1202.
- [34] Mutluer, B.; Erko, F. U. Effects of gamma irradiation on aflatoxins. *Z. Lebensm. Unters. Forch.* **1987**, *185*, 398–401.
- [35] Mishra, H. N.; Das, C. A review on biological control and metabolism of aflatoxin. *Crit. Rev. Food Sci. Nutr.* **2003**, *43*, 245–264.
- [36] Wogan, G. N.; Edwards, G. S.; Newberne, P. M. Structure-activity relationships in toxicity and carcinogenicity of aflatoxins and analogues. *Cancer Res.* **1971**, *31*, 1936–1942.

- [37] Aziz, N. H.; Souzan, R. M.; Azza, A. Effect of gamma-irradiation on the occurrence of pathogenic microorganisms and nutritive value of four principal cereal grains. *App. Rad. Isotopes* **2006**, *64*, 1555–1562.
- [38] Taipina, M. S.; Garbelotti, M. L.; Lamardo, L. C. A.; Santos, J. S.; Rodas, M. A. B. The effect of gamma irradiation on the nutritional properties of sunflower whole grain cookies. *Proc. Food Sci.* **2011**, *1*, 1992–1996.
- [39] Hamza, R. G.; Afifi, S.; Abdel-Ghaffar, A.-R. B.; Borai, I. H. Effect of gamma-irradiation or/and extrusion on the nutritional value of soy flour. *Biochem. Anal. Biochem.* **2012**, *1*, 118. doi: [10.4172/2161-1009.1000118](https://doi.org/10.4172/2161-1009.1000118)
- [40] World Health Organisation (WHO). *High-dose irradiation: Wholesomeness of food irradiated with doses above 10 KGy*. Joint FAO/IAEA/WHO study group. Technical Report Series, No. 890. **1999**, Geneva.