



## Intercropping influences component and content change of flavonoids in root exudates and nodulation of Faba bean

Ying Chao Liu, Xiao Min Qin, Jing Xiu Xiao, Li Tang, Chi Zhang Wei, Jin Jian Wei & Yi Zheng

To cite this article: Ying Chao Liu, Xiao Min Qin, Jing Xiu Xiao, Li Tang, Chi Zhang Wei, Jin Jian Wei & Yi Zheng (2017) Intercropping influences component and content change of flavonoids in root exudates and nodulation of Faba bean, Journal of Plant Interactions, 12:1, 187-192, DOI: [10.1080/17429145.2017.1308569](https://doi.org/10.1080/17429145.2017.1308569)

To link to this article: <https://doi.org/10.1080/17429145.2017.1308569>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 12 Apr 2017.



[Submit your article to this journal](#)



Article views: 1638



[View related articles](#)



[View Crossmark data](#)



Citing articles: 12 [View citing articles](#)

RESEARCH ARTICLE



## Intercropping influences component and content change of flavonoids in root exudates and nodulation of Faba bean

Ying Chao Liu<sup>a</sup>, Xiao Min Qin<sup>b</sup>, Jing Xiu Xiao<sup>a</sup>, Li Tang<sup>a</sup>, Chi Zhang Wei<sup>b</sup>, Jin Jian Wei<sup>b</sup> and Yi Zheng<sup>a</sup>

<sup>a</sup>College of Resources and Environment, Yunnan Agricultural University, Kunming, People's Republic of China; <sup>b</sup>Tea Breeding Research Center, Guangxi South Subtropical Agricultural Science Research Institute, Chongzuo, People's Republic of China

### ABSTRACT

Flavonoids produced by legume roots are signal molecules acting as nod gene inducers for the symbiotic rhizobium partner. Nevertheless, the changes of flavonoids in root exudates in intercropping system are still unknown. Based on pot experiment of faba bean and wheat intercropping, here we showed that faba bean and wheat intercropping increased the nodules number and dry weight, dry weight per nodule of faba bean compared with those found in monocropping, and the increase of faba bean nodulation was likely caused by the enhancement with flavonol, isoflavone, chalcone and hesperetin in its root exudates. It also promoted exudation of five types of flavonoids by wheat compared with monocropping. Our findings suggest that the flavonoids in root exudates have a positive effect on the nodulation and nitrogen fixation of faba bean in faba bean and wheat intercropping.

### ARTICLE HISTORY

Received 12 January 2017  
Accepted 15 March 2017

### KEYWORDS

Flavonoids; nodulation; intercropping; wheat; faba bean

## 1. Introduction

Intercropping is advanced as one of the foremost conventional production practices used in agro-ecosystem, a set of investigations have clearly demonstrated that compared with monocropping, it could increase natural resources use efficiency (Gao et al. 2009; Rivest et al. 2010; He et al. 2013), boost crop yield and mineral nutrient accumulation (Nataraj et al. 2010; Mao et al. 2012), enhance biological diversity (Nai et al. 2013; Zhang et al. 2015), and lower disease, insect, and weed pressures (Workayehu & Wortmann 2011; Abdel-Monaima & Abo-Elyousr 2012; Boudreau 2013). Recently, it has attracted great attention due to the yield advantage with the serious challenges of resources, environment and food.

Root is the major organ of the plant to uptake nutrients and water, and its root exudates play an important role in rhizosphere processes (Terzano et al. 2015). Flavonoids in root exudates are a class of widely distributed secondary metabolites that play an important function in mediating the interaction of plants with their environment (Pourcel et al. 2007). Studies have demonstrated that it is affected by a range of factors, such as crop species, temperature, UV radiation and nitrogen rates (Coronado et al. 1995; Zhang & Li 2003; Li et al. 2012), and particularly crop species. For instance, the flavonoids in legume root exudates are the inducers of genes responsible for the nodulation process of rhizobium in specific legumes (Broughton et al. 2003; Badri & Vivanco 2009; Paul et al. 2010; Wang et al. 2012). Recently, scientists found that there were different components of flavonoids in legume root exudates in the monocropping system (Miao & Liu 2010; Li 2013), however, when legume is intercropped with cereal does it promote the exudation of the flavonoids by root? How is it promoted? At present, these problems are still not very clear.

Intercropping of wheat (*Triticum aestivum* L.) with faba bean (*Vicia faba*) is one of the most popular intercropping systems in Yunnan province, China (Yang et al. 2014; Xiao et al. 2016), has clearly been demonstrated it could make better use of nutrition, increase crop yield and nodule number, lower disease and change the carbon and nitrogen metabolism (Su et al. 2006; Miao et al. 2009; Dong et al. 2013; Xiao et al. 2015), whereas there is lack of information and knowledge about the effect of it on changes of flavonoids in root exudates and its connection with faba bean nodulation.

The major objective of this study was to elucidate how faba bean and wheat intercropping impacts the flavonoids change in root exudates and the potential role of flavonoids in faba bean nodulation, aimed at providing a new perspective for facilitative root–root interactions and developing sustainable agriculture.

## 2. Materials and methods

### 2.1. Experiment site

Pot trial was carried out in a greenhouse of Yunnan Agricultural University (YNAU) in 2015, Kunming, Yunnan province, southeast China (25°2'N, 102°42'E). This region has a subtropical monsoon climate with an average annual precipitation of 1035 mm, an average annual temperature of 15°C and an altitude of 1962 m. The soil used was taken from a mountain near YNAU, air-dried and ground and sieved to pass through a 5 mm mesh. The soil was red and its initial properties were as follows: pH 6.08; soil organic matter 28.07 g kg<sup>-1</sup>; available nitrogen 68 mg kg<sup>-1</sup>; available phosphorus 16 mg kg<sup>-1</sup> and available potassium 137 mg kg<sup>-1</sup>.

Wheat (*Triticum aestivum* L.cv. Yunmai-42) and Faba bean (*Vicia faba* L. cv. Yuxidalidou) were chosen for the pot experiment. Both seeds were sterilized in 30% (v/v) H<sub>2</sub>O<sub>2</sub> for 20 min,

washed with distilled water, soaked in saturated  $\text{CaSO}_4$  solution for 12 h and then germinated in Petri dishes in the dark for 48 h at 22°C. They were transplanted when the second leaves of wheat and faba bean have expanded to its full size.

## 2.2. Experiment design

Pot trial was conducted with three planting treatments which consisted of wheat monocropping, wheat and faba bean intercropping, faba bean monocropping and three replications, sampling 3 times and giving a total of 36 pots in a randomized block design. Planting ratio was as follows: four plants of wheat seedlings intercropped with two plants of faba bean seedlings in intercropping pot, and eight plants of wheat seedlings, four plants of faba bean seedlings were planted in each monocropping pot, respectively.

Urea, superphosphate and potassium sulfate were applied as N, P and K fertilizers in the experiment. Fifty percent of N rates (150 mg N kg<sup>-1</sup> soil), P and K (100 mg P<sub>2</sub>O<sub>5</sub> and 100 mg K<sub>2</sub>O kg<sup>-1</sup> soil) as basal were incorporated into the soil by mixing with 10 kg soil, and it was then distributed among plastic plant pots (25 cm height × 34 cm diameter), and the remaining 50% of the N dressed in the jointing stage along with irrigation. Soil water content was regularly adjusted during the growth of the plants, and no fungicides or insecticides were applied.

## 2.3. Sampling

The samplings took place at the tillering stage/branching stage, jointing stage/flowering stage and grain-filling stage/seed-filling stage, respectively. According to the methods conducted by Xiao et al. (2014), crop roots were taken off from the soil, shaken off the loose soil, washed the root with water and rinsed three times with distilled water, soaked in 5% thymol solution for 3 min and transferred into containers with 500 ml  $\text{CaCl}_2$  solution for 2 h, and then extracted with 200, 100 and 50 ml ethyl acetate, evaporated and concentrated, titrated to 10 ml with methanol and filtered through 0.45 µm polytetrafluoroethylene filter for high-performance liquid chromatography (HPLC) analysis.

## 2.4. Measurements

### 2.4.1. Number of nodulation

The number of nodules were counted and recorded, while the weight of the oven-dried nodules was obtained with the aid of a sensitive weighing balance.

### 2.4.2. HPLC analysis of flavonoids

An Agilent (Waldbronn, Germany) 1100 HPLC series, which consists of a degasser, binary pump, auto-sampler, thermostat and photodiode array detector, was used to determine the composition and concentration of the flavonoids according to the method reported by Krumbein et al. (2007). A Prodigy column (ODS 3, 150 × 3.0 mm, 5 µm, 100 Å; Phenomenex, Aschaffenburg, Germany) with a security guard C18 column (ODS 3, 4 × 3.0 mm, 5 µm, 100 Å) was used for separating the extracts at 30°C. Solvent A consisted of 99.5% water and 0.5% acetic acid, and solvent B consisted of 100% acetonitrile. The following gradients were used: 30–40% B (5 min), 40–60% B (5 min), 60–90% B (15 min), 90% B (4 min), 90–30% B (5 min) and 30% B (3 min). The chromatogram was

monitored at 270 nm with a flow rate of 0.9 ml min<sup>-1</sup>. Flavonol, isoflavone, flavone, chalcone, quercetin, naringenin and hesperetin were used as external standards for calculating the calibration curve in the range of 0.05–1 mg 100 ml<sup>-1</sup>.

## 2.5. Statistical analysis

Data in the tables and figures were reported as the mean (±) standard deviation (SD) of three replicates, while all other data were reported as the mean alone. We used one-way analysis of variance (ANOVA) to identify differences in the responses of faba bean nodulation characteristics and different flavonoids under intercropping using the SPSS 19.0 software, and applied a significance level of  $P < .05$ .

## 3. Results

### 3.1. Grain yields

In the pot experiment (Table 1), we found that grain yields of faba bean and wheat in intercropping were 8.70%, 14.56% higher than that observed in monocropping, respectively. These results suggest that faba bean intercropped with wheat increased the crop yields compared with monocropping.

### 3.2. Changes in nodules in faba bean and wheat intercropping system

Nodule is an important condition for symbiotic nitrogen fixation of legume crops, and its number can measure the ability of nitrogen fixation. Significant intercropping effect in faba bean nodules was observed (Table 2). The nodules number and dry weight, dry weight per nodule in intercropped faba bean significantly were 80.23%, 133.33% and 29.62% higher than those found in monocropped at the flowering stage (95d), and 5.80%, 139.58% and 126.11% higher than that observed in monocropped at seed-filling stage (131d), which indicated that faba bean and wheat intercropping likely promote faba bean nodulation.

Additionally, the nodules number increased initially and then decreased with the increase in growth stages, and presented the highest number at the flowering stage (95d) which suggested faba bean root formed lots of mature nodules at the flowering stage (95d), and then it slowly rot off with podding. Thus, we need to focus on the changes in flavonoids in faba bean root exudates during the primary period of nodulation to observe whether it was related to faba bean nodulation.

### 3.3. Changes in total flavonoids in faba bean and wheat intercropping system

As shown in Figure 1, the concentrations of the flavonol, isoflavone and hesperetin for four treatments were higher than

**Table 1.** Grain yields of faba bean and wheat (g/pot).

Cropping patterns	Grain yields
MF	22.30
IF	24.24
MW	21.64
IW	24.79

Note: MF: Monocropped faba bean; IF: Intercropped faba bean; MW: Monocropped wheat; IW: Intercropped wheat.

\*Mean significant difference between monocropping and intercropping pattern ( $P < .05$ ).

**Table 2.** The effects of different cropping system on faba bean nodulation ( $\text{g}\cdot\text{plant}^{-1}$ ).

Stages	Cropping patterns	Nodule number	Nodule dry weight	dry weight per nodule
95d	MF	86 $\pm$ 5.57	0.48 $\pm$ 0.06	5.57 $\pm$ 0.36
	IF	155 $\pm$ 4.58*	1.12 $\pm$ 0.09*	7.22 $\pm$ 0.42*
131d	MF	69 $\pm$ 5.29	0.48 $\pm$ 0.05	6.97 $\pm$ 0.67
	IF	73 $\pm$ 4.58	1.15 $\pm$ 0.11*	15.76 $\pm$ 1.21*

Note: MF: Monocropped faba bean; IF: Intercropped faba bean. \*Mean significant difference between monocropping and intercropping pattern ( $P < .05$ ).

others and the concentration levels varied from 0.0% to 40.79%, 16.98% to 22.49%, 13.26% to 27.01%, and 5.04% to 33%, 14.86% to 25%, 12.67% to 18.38%, respectively, while the flavone concentration was lower.

The intercropping effect on total flavonoids is shown in Figure 2, the total flavonoid concentrations in intercropped faba bean root exudates were 23.72%, 29.21% and 45.24% higher than that found in monocropped at the three growth stages, and a considerable difference was found at the flowering stage (95d). Besides, the total flavonoids concentration in intercropped wheat root exudates were 75.44% and 10.04%, respectively, greater than that of monocropped wheat at tillering (60d) and grain-filling stages (131d), while it was significantly lower than monocropped wheat by 7.81% at the jointing stage (95d).

### 3.4. Changes in seven types of flavonoids in faba bean and wheat intercropping system

As shown in Figure 3, the intercropping effect on seven types of flavonoids in the faba bean and wheat root exudates was observed. Faba bean intercropped with wheat increased the


concentration of flavonol, isoflavone, chalcone and hesperetin in its root exudates in intercropping treatment by 114.33% and 853.50%, 13.77% and 44.26%, 101.84% and 72.09%, and 83.79% and 9.24% compared with that found in monocropping during the nodulation period (95d–131d), while it lowered the concentration of quercetin and naringenin. Also, it enhanced the concentration of flavonol, isoflavone, flavone, chalcone, naringenin and hesperetin in intercropping faba bean root exudates at the branching stage (60d) compared with monocropping. Besides, we also found that the concentrations of isoflavone, flavone, chalcone, quercetin and hesperetin in wheat root exudates in intercropped wheat root exudates at the three stages all were greater than that obtained in monocropped wheat.

### 3.5. Cluster analysis


Cluster analysis is used to group the abstract objects with similarity into a category that could be more intuitive to reveal the variations among these objects. As shown in Figure 4, the concentration of seven types of flavonoids for four treatments was used for cluster analysis. Result indicated that intercropping and monocropping treatments were obviously divided into two categories that suggested wheat and faba bean intercropping significantly changed the root exudation of seven types of flavonoids.

### 3.6. Relationship between flavonoids and faba bean nodulation

Recent studies revealed that root exudates played an important role in affecting the legume nodulation and nitrogen



**Figure 1.** Percentage of seven types of flavonoids. MW: monocropped wheat; IW: intercropped wheat; MF: monocropped faba bean and IF: intercropped faba bean.




**Figure 2.** Effects of intercropping on total flavonoids in root exudates. MW: monocropped wheat; IW: intercropped wheat; MF: monocropped faba bean; IF: intercropped faba bean. \*Mean significant difference between monocropping and intercropping patterns ( $P < .05$ ).

fixation in the legume and cereal intercropping systems (Paul et al. 2010). As shown in Table 3, the nodules number and dry weight, dry weight per nodule were positively correlated with the flavonol, isoflavone, flavone, chalcone, quercetin, naringenin and hesperetin. Besides, the flavonol, naringenin and hesperetin were highly significantly positively correlated with above parameters of faba bean nodules which suggested that the flavonol, naringenin and hesperetin of flavonoids in root exudates were more sensitive components that relate to the faba bean nodulation.

## 4. Discussion


### 4.1. Effects of intercropping on faba bean nodulation

Legumes could form nitrogen-fixing nodules through specific recognition of their roots and rhizobium, and it is an important characteristic distinguishing it from other higher plants



**Figure 3.** The effects of intercropping on seven types of flavonoids. MW: monocropped wheat; IW: intercropped wheat; MF: monocropped faba bean and IF: intercropped faba bean. \*Mean significant difference between monocropping and intercropping patterns ( $P < .05$ ).





**Figure 4.** Cluster analysis of seven types of flavonoids. MW: monocropped wheat; IW: intercropped wheat; MF: monocropped faba bean and IF: intercropped faba bean.

(Hirsch 1992), so they can reduce nitrogen input through biological nitrogen fixation. Banik and Sharma (2009) found that baby corn and legume intercropping could promote the ability of legume nodulation. Cun et al. (2014) also reported that faba bean intercropped with garlic clearly improved faba bean nodulation and dry weight of nodules. Similarly, faba bean intercropped with wheat could increase the nodule number, nodules dry weight and dry weight per nodule of faba bean compared with that found in monocropping, and these findings indicated faba bean intercropped with wheat could improve the ability of faba bean nodulation which was in agreement with the results of previous studies (Banik & Sharma 2009; Cun et al. 2014). On the one hand, these results may be due to the interspecific facilitation leading to crops releasing richer exudates which could provide more carbon sources for growth and reproduction of specific rhizobium in faba bean rhizosphere, thus promoting faba bean nodulation in the intercropping system. On the other hand, previous study reported that the nodulation level of legumes was negatively correlated with soil nitrogen content, and known as a 'negative feedback regulation of nitrogen' (Li et al. 2009). Therefore, when it was intercropped with wheat, rhizosphere N availability was decreased caused by stronger competition of wheat for nitrogen, and this decrease presumably derived greater nodulation of faba bean in the intercropping system.

#### 4.2. Changes in root exudation of flavonoids and its role in faba bean nodulation

Flavonoids are a class of widely distributed secondary metabolites that are exuded by plant root, and it can induce specific rhizobium to synthesize and exude a type of Nod factor (Lipo-chit oligosaccharide, LCOs) which plays an important role in the symbiotic nodulation of rhizobium and plant (Wang et al. 2012). For instance, some studies have well shown that several flavonoids, including naringenin,

hesperetin, genistein and 7, 4-dihydroxyflavone, could induce the expression of Nod gens, and then stimulate the nodulation and nitrogen fixation of legume (Broughtone et al. 2003; Giraud et al. 2007; Maj et al. 2010). In our study, the concentrations of flavonol, isoflavone, chalcone and hesperetin in intercropping faba bean root exudates during the nodulation period were increased when compared with monocropping, and we found that flavonol, naringenin and hesperetin of flavonoids in root exudates were more sensitive components relating to faba bean nodulation. Thus, the enhancement of nodulation was likely caused by the enrichment with flavonol, isoflavone, chalcone and hesperetin in root exudates when faba bean was intercropped with wheat, which was similar to the result of Li et al. (2012), who found that maize intercropped with faba bean could change the content of quercetin and luteolin that could promote significantly faba bean nodulation. However, legumes nodulation is a complex process which consists of the interaction between legumes root and soil rhizobium, a series of signal transduction and material synthesis, so further investigation is required to evaluate the intrinsic mechanism.

Furthermore, we observed the changes of different flavonoids in wheat and faba bean root exudates at different growth stages, and found that intercropping also increased the concentration of flavonol, isoflavone, flavone, chalcone, naringenin and hesperetin in faba bean root exudates at the branching stage (60d) compared with monocropping, these enrichments also likely provide the substrate for initiating early nodulation. At the same time, the concentrations of isoflavone, flavone, chalcone, quercetin and hesperetin in intercropping wheat root exudates at the three stages all were higher than that found in monocropping, which suggested faba bean intercropped with wheat also promotes correspondingly the exudation of flavonoids by wheat root through interspecific root interactions. However, the impacts of intercropping on different flavonoids in root exudates were

**Table 3.** Relationship between seven types of flavonoids and faba bean nodulation.

Growth stages	Factors	Flavonol	Isoflavone	Flavone	Chalcone	Quercetin	Naringenin	Hesperetin
95d	Nodule number	0.9464**	0.6502	0.0245	0.9092**	0.4700	0.9928**	0.9928**
	Dry weight of nodule	0.9600**	0.5627	0.0608	0.9160**	0.4159	0.9822**	0.9822**
	Dry weight per nodule	0.9256**	0.4308	0.1530	0.9112**	0.4026	0.9330**	0.9330**
131d	Nodule number	0.9582**	0.7174	0.5693	0.6785	0.4460	0.9928**	0.9928**
	Dry weight of nodule	0.9425**	0.6574	0.5249	0.6170	0.4738	0.9822**	0.9822**
	Dry weight per nodule	0.9726**	0.6396	0.5385	0.5778	0.4241	0.9840**	0.9840**

Note: \* and \*\* mean the differences are significant at level of 5% and 1%, respectively.

observed only at three primary stages in the present study, so further investigation is necessary to understand the flavonoid changes throughout the entire growth period.

## 5. Conclusions

Cereal intercropped with legume is advanced and widespread due to the biological nitrogen fixation of the legume which is beneficial for maintaining and improving soil fertility. The effects of wheat and faba bean intercropping on component and content change of flavonoids and their role in faba bean nodulation remain unclear. In our study, the nodules number, nodules dry weight and dry weight per nodule of faba bean in intercropping were increased compared with those found in monocropped faba bean, and the enhancement of faba bean nodulation was likely caused by the enrichment with flavonol, isoflavone, chalcone and hesperetin in its root exudates. Wheat and faba bean intercropping also promoted exudation of five types of flavonoids by wheat compared with monocropping. However, legumes nodulation is a complex process, so further studies are necessary to gain in-depth knowledge about legume and cereal intercropping underlying the component and content change of flavonoids, particularly the role of flavonoids in legumes nodulation and nitrogen fixation.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

We thank to the National Natural Science Foundation of China (NSFC) [31460551, 31260504, and 41361065] for financial support.

## References

- Abdel-Monaima MF, Abo-Elyousr KAM. 2012. Effect of preceding and intercropping crops on suppression of lentil damping-off and root rot disease in New Valley-Egypt. *Crop Prot.* 32:41–46.
- Badri DV, Vivanco JM. 2009. Regulation and function of root exudates. *Plant, Cell Environ.* 32:666–681.
- Banik P, Sharma RC. 2009. Yield and resource utilization efficiency in baby corn-legume-intercropping system in the eastern plateau of India. *J Sustain Agr.* 33:379–395.
- Boudreau MA. 2013. Disease in intercropping systems. *Annu Rev Phytopathol.* 51:499–519.
- Broughton WJ, Zhang F, Penet X, Staehelin C. 2003. Signals exchanged between legumes and Rhizobium: agricultural uses and perspectives. *Plant Soil.* 252:129–137.
- Coronado C, Zuannazzi JAS, Sallaud C. 1995. Alfalfa root flavonoid production is nitrogen regulated. *Plant Physiol.* 108:533–542.
- Cun ZX, Chen JB, Xiu HX, He XH, Tang DS. 2014. The influence of intercropping mode and planting density on faba bean nodulation and plant growth. *Journal of Yunnan Agricultural University.* 29:514–520.
- Dong Y, Dong K, Tang L, Zheng Y, Yang ZX, Xiao JX, Zhao P, Hu GB. 2013. Relationship between rhizosphere microbial community functional diversity and faba bean Fusarium wilt occurrence in wheat and faba bean intercropping system. *Acta Ecol Sin.* 33:7445–7454.
- Gao Y, Duan A, Sun J, Li F, Liu Z, Liu H, Liu Z. 2009. Crop coefficient and water-use efficiency of winter wheat/spring maize strip intercropping. *Field Crop Res.* 111:65–73.
- Giraud E, Moulin L, Vallenet D, Barbe V, Cytryn E, Avane JC, Jaubert M. 2007. Legumes symbioses, absence of nod genes in photosynthetic Bradyrhizobia. *Science.* 316:1307–1312.
- He Y, Ding N, Shi JC, Wu M, Liao H, Xu JM. 2013. Profiling of microbial PLFAs: Implication for interspecific interactions due to intercropping which increase phosphorus uptake in phosphorus limited acidic soils. *Soil Biol Biochem.* 57:625–634.
- Hirsch AM. 1992. Developmental biology of legume nodulation. *New Phytologist.* 122:211–237.
- Krumbein A, Saeger-Fink H, Schonhof I. 2007. Changes in quercetin and kaempferol concentrations during broccoli head ontogeny in three broccoli cultivars. *J Appl Bot Food Qual-Angew Bot.* 81:136–139.
- Li B, Knunbein A, Neugartb S, Li L, Schreiner M. 2012. Mixed cropping with maize combined with moderate UV-B radiations lead to enhanced flavonoid production and root growth in faba bean. *J Plant Interact.* 7:333–340.
- Li L. 2013. Legume nitrogen fixation ecology principle and application in intercropping system. Beijing: China Agriculture University Press.
- Li YY, Sun JH, Li CJ, Li L, Cheng X, Zhang FS. 2009. Effects of inter-specific interactions and nitrogen fertilization rates on the agronomic and nodulation characteristics of intercropped faba bean. *Scientia Agricultura Sinica.* 42:3467–3474.
- Maj D, Wielbo J, Marek-Kozaczuk M, Skorupska A. 2010. Response to flavonoids as a factor influencing competitiveness and symbiotic activity of *Rhizobium leguminosarum*. *Microbiol Res.* 165:50–60.
- Mao LL, Zhang LZ, Li WQ, Werf WVD, Sun JH, Spiertz H, Li L. 2012. Yield advantage and water saving in maize/pea intercrop. *Field Crops Res.* 138:11–20.
- Miao R, Zhang FS, Li L. 2009. Effects of root barriers on nodulation of faba bean in maize/faba bean, wheat/faba bean and barley/faba bean intercropping systems. *Chin Bull Botany.* 44:197–201.
- Miao SJ, Liu XB. 2010. Molecular physiology of nodulation and nitrogen fixation in soybean. *Soybean Sci.* 29:320–324.
- Nai FJ, Wu LH, Liu HY, Ren J, Liu WX, Luo YM. 2013. Effects of intercropping sedum plumbizincicola and *Apium graveolens* on the soil chemical and microbiological properties under the contamination of zinc and cadmium from sewage sludge application. *Chin J Appl Ecol.* 24:1428–1434.
- Nataraj D, Shashidhar KS, Vinoda KS, Chandrashekara C, Kalyanamurthy. 2010. Profitability and potentiality of baby corn based leguminous vegetable intercropping system. *Environ Ecol.* 28:1433–1436.
- Paul GD, Anthony JM, Penny R. 2010. Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiol Ecol.* 72:313–327.
- Pourcel L, Routaboul JM, Cheyrier V, Lepiniec L, Debeaujon I. 2007. Flavonoid oxidation in plants: from biochemical properties to physiological functions. *Trends in Plant Sci.* 12:29–36.
- Rivest D, Cogliastro A, Bradley R, Olivier A. 2010. Intercropping hybrid poplar with soybean increases soil microbial biomass, mineral N supply and tree growth. *Agroforestry Syst.* 80:33–40.
- Su HP, Tang L, Liu ZH, Li SM, Li YJ, Zheng Y. 2006. Dynamics of nitrogen assimilates of wheat in wheat-faba bean intercropping system. *J Triticeae Crops.* 26:140–144.
- Terzano R, Cesco S, Mimmo T. 2015. Dynamics, thermodynamic and kinetics of exudates: crucial issues in understanding rhizosphere processes. *Plant Soil.* 386:399–406.
- Wang D, Yang SM, Tang F, Zhi HY. 2012. Symbiosis specificity in the legume-thizobial mutualism. *Cell Microbiol.* 14:334–342.
- Workayehu T, Wortmann CS. 2011. Corn-bean intercrop weed suppression and profitability in Southern Ethiopia. *Agron J.* 103:1058–1063.
- Xiao JX, Zheng Y, Tang L. 2014. Effect of wheat and faba bean intercropping on root exudation of low molecular weight organic acids. *Chin J Appl Ecol.* 25:1739–1744.
- Xiao JX, Zheng Y, Tang L, Dong Y. 2015. Effect of wheat and faba bean intercropping on sugar and amino acid exuded by roots. *Ecol Environ Sci.* 24:1825–1830.
- Xiao JX, Zheng Y, Tang L, Wang G, Dong Y. 2016. Changes in organic and phenolic acids in rhizosphere of Interplanted wheat and faba bean with growth stage. *Acta PedologIca Sinica.* 53:686–693.
- Yang ZX, Tang L, Zheng Y, Dong Y, Dong K. 2014. Effects of different wheat cultivars intercropped with faba bean on faba bean Fusarium wilt, root exudates and rhizosphere microbial community functional diversity. *J Plant Nutr Fertilizer.* 20:570–579.
- Zhang FS, Li L. 2003. Using competitive and facilitative interactions in intercropping systems enhance crop productivity and nutrients-use efficiency. *Plant Soil.* 248:305–312.
- Zhang MM, Ao H, Li X, Zhang JY, Wang N, Ju CM, Wang J, Cai DJ, Sun GY. 2015. Effects of intercropping between mulberry and alfalfa on soil enzyme activities and microbial community diversity in rhizosphere. *Acta Agrestia Sinica.* 23:303–309.