



Characterization of soils in terms of pedological variability under different physiography of Damodar command area (part), West Bengal, India

Ranjan Bera, Antara Seal, Thako Hari Das, Dipak Sarkar & Ashis Kumar Chatterjee |

To cite this article: Ranjan Bera, Antara Seal, Thako Hari Das, Dipak Sarkar & Ashis Kumar Chatterjee | (2015) Characterization of soils in terms of pedological variability under different physiography of Damodar command area (part), West Bengal, India, Cogent Food & Agriculture, 1:1, 1014719, DOI: [10.1080/23311932.2015.1014719](https://doi.org/10.1080/23311932.2015.1014719)

To link to this article: <https://doi.org/10.1080/23311932.2015.1014719>



© 2015 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license



Published online: 26 Feb 2015.



Submit your article to this journal [↗](#)



Article views: 1871



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)



SOIL & CROP SCIENCES | RESEARCH ARTICLE

Characterization of soils in terms of pedological variability under different physiography of Damodar command area (part), West Bengal, India

Ranjan Bera, Antara Seal, Thako Hari Das, Dipak Sarkar and Ashis Kumar Chatterjee

Cogent Food & Agriculture (2015), 1: 1014719



Received: 05 October 2014
Accepted: 07 January 2015
Published: 26 February 2015

*Corresponding author: Ranjan Bera,
Department of Soil Science, Visva
Bharati University, Santiniketan 731236,
West Bengal, India
E-mail: bera.ranjan@gmail.com

Reviewing editor:
Manuel Tejada Moral, University of
Seville, Spain

Additional information is available at
the end of the article

SOIL & CROP SCIENCES | RESEARCH ARTICLE

Characterization of soils in terms of pedological variability under different physiography of Damodar command area (part), West Bengal, India

Ranjan Bera^{1*}, Antara Seal¹, Thako Hari Das², Dipak Sarkar² and Ashis Kumar Chatterjee¹

Abstract: Five representative soil profiles occurring on four different physiography under subtropical environment of Damodar command area, India, were studied for soil pedological variability. Two way approaches were taken to evaluate the extent of profile development. Firstly different extractants were used to determine various forms of Fe and Al and their different ratios. Average contents of Fe and Al, extracted by different extracting reagents were found to be in descending order, as follows: $Al_{dith} > Al_{oxa} > Al_{pyr}$ and $Fe_{dith} > Fe_{oxa} > Fe_{pyr}$. Analysis of pyrophosphate (pyr), oxalate (oxa), and dithionate (dith) extractable Fe and Al fractions indicated that with increasing soil age, the content of crystalline Fe and Al oxides increased at the expense of the poorly crystalline forms. The mean content of amorphous Fe and Al, crystalline Fe and Al, and their ratios estimated the degree of soil development. In the second part, elemental analysis was done, silica to sesquioxide ratio as well as ratio of alkali cations was measured and weathering index of each horizon was determined. The ratios and weathering indices indicated that except Madhpur soil series, all other soils were young and pedological development was still in progress in Damodar command area.

Subjects: Environment & Agriculture; Earth Sciences; Environmental Studies & Management

Keywords: Damodar command area; forms of Fe and Al; elemental analysis; weathering index

1. Introduction

Soil properties that change with duration and intensity of weathering provide vital clue toward the pedogenesis of the studied soil. Mineral weathering during pedogenesis results in translocation and/or accumulation of major or trace elements in soils (Kabata-Pendias & Pendias, 1992). The translocation or accumulation of the elements occurred via chemical and biological processes such as leaching, podzolization, and oxidation–reduction (Mausbach & Richardson, 1994; Wilson, Burt,

ABOUT THE AUTHORS

The authors of the research article are working in the arena of soil science. The study was basically PhD work of Dr. Ranjan Bera, corresponding author and other authors attached with two leading Institute of India viz. National Bureau of Soil Survey and Land Use Planning and Department of Soil Science, Viswa-Bharati University joined hands to fulfill the objectivity.

PUBLIC INTEREST STATEMENT

In the research article, an attempt was made to evaluate the soil formation pattern in Damodar Command area of West Bengal, India. The area is known for its high crop productivity and diversified cropping pattern. So an in depth study of its soil formation process might be useful to find the basic reason behind its inherent high soil productivity.

Sobecki, Engel, & Hipple, 1996). Determination of the different soil processes involved elemental evaluation of horizons within and between the solum (Burt, Wilson, Mays, & Lee, 2003). Pedogenic evaluations using elemental analysis include silica to sesquioxide ratio representing soil weathering index (Jackson, 1973) and ratios of alkali cations (e.g. Ca and K) as measures of pedogenic changes and geological uniformity (Muhs, Bettis, Been, & McGeehin, 2001). Comparative movement of silica and sesquioxide down the profile as well as their ratios calculated from the total composition of soils provide basis for understanding profile development in relation to parent material. On the other hand, profile distribution of different forms of Fe and Al oxides particularly dithionite and oxalate extractable Fe and Al serve as useful indicators to identify the horizon of accumulation of secondary oxides (McKeague & Day, 1966). The Fe and Al released during the weathering of Fe and Al bearing parent materials are reprecipitated in the soils as oxides or hydroxides and oxyhydroxides of iron and aluminum. The quantities of these alternation products generally increase with soil age (Dolui & Bera, 2001) and their distribution in soil profiles indicate the stage and degree of soil development (Mahaney & Fahey, 1988). However, very meager information is available on the degree of soil development and the direction of pedogenic processes in the soils of Damodar command area. Hence, in the present study these two approaches were used for pedogenic evaluation and intensity of weathering in the soils occurring under different physiographic positions in the study area.

2. Materials and methods


The study area lies between 23°11'46" to 23°29' 21" N latitude and 87°28'49" to 88°0' 21" E longitude in Vindyan plain of Bardhaman, West Bengal, India. It represents a transition zone of tropical dry subhumid type of climate area and belongs to hot moist subhumid ecological subregion (15.1). The mean summer (March–June) and winter (November–February) temperature are 30.0 and 21.1°C, respectively with a mean annual temperature of 26.5°C. The difference between mean summer and mean winter temperature is more than 5°C, hence the area qualifies for “hyperthermic” temperature regime. The mean annual precipitation is 1,400 mm of which approximately 74.7% is received during June–September.

Based on reconnaissance soil survey, five soil series representing four different physiography were selected to study the soil pedological variability in Damodar command area (part). Among the selected soil series, Khamrp soil series belonged to active floodplain, Karjana soil series belonged to gently sloping floodplain, Banpara soil series belonged to nearly level alluvial plain while Hargram and Madhpur soil series represented gently sloping alluvial plain (Figure 1). Soils of each horizon were processed (<2 mm) and used for analyzing the physical and chemical characteristics of the soil samples viz. particle size distribution, pH, organic carbon, CEC, and base saturation, following standard procedures (Jackson, 1973). The soils were classified as per the Keys to Soil Taxonomy.

Fraction of Fe and Al was determined in the soil by separate extractions with (i) sodium pyrophosphate, overnight extraction at pH 10 (Agriculture Canada, 1984), (ii) acid ammonium oxalate (0.2), four hour extraction at pH 3 (Agriculture Canada, 1984), and (iii) dithionate-citrate-bicarbonate, 20 min extraction twice (CSSC Subcommittee on methods of analysis, 1978). Among the extractants used dithionite-citrate-bicarbonate (DCB) reagent was the most powerful, which dissolved with the extraction capacity of both non-crystalline and crystalline Fe and Al oxides, small amount of water soluble, exchangeable, and organically bound Fe and Al along with limited amount of Fe and Al bearing silicates (Borggard, 1988). This was followed by acid ammonium oxalate, which dissolved both “amorphous” and “organically bound” forms of Fe and Al, but not the crystalline forms (Parfitt & Childs, 1988). In comparison pyrophosphate extractant dissolved only the fraction of iron (Loveland & Digby, 1984) and Al (Driscoll, van Breemen, & Mulder, 1985) that are bound with organic matter. Fe and Al in the extract were determined using atomic absorption spectrophotometer. Total elemental composition of soil samples was carried out following the procedure of Jackson (1973). Weathering index of each horizon was calculated by the formula given by Evans and Cameron (1979).

$$\frac{[(\text{CaO} + \text{MgO} + \text{K}_2\text{O} + \text{Na}_2\text{O})/(\text{CaO} + \text{MgO} + \text{K}_2\text{O} + \text{Na}_2\text{O} + \text{Si}_2\text{O}_3 + \text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3)] \times 100$$

Figure 1. Physiography map of the study area.



3. Result and discussion

Physicochemical properties of the soils from the selected soil series are represented in Table 1. In all the soil series, surface soils exhibited slightly acidic pH (5.4–6.3) which gradually shifted to neutral range in the subsurface and lower horizons except in Hargram soil series. In all the soil series, the lowest clay percentage was observed in the surface horizon and the values increased with increase in soil depth. However, in Hargram and Banpara soil series clay percentage decreased abruptly in the lowest horizon, while no consistent pattern was observed in the soils of Karjana and Khamrp soil series. Such abrupt changes in soil property were in close conformity to the general characteristics of alluvial soils. Cation exchange capacity of the soils also followed a similar trend in the different soil series.

The data on various forms of Fe and Al as extracted by different extractants, their ratios, relationship with soil properties and interrelationship among themselves are presented in Tables 2–4, respectively.

3.1. Pyrophosphate extractable Fe and Al

Pyrophosphate extractant dissolved the fraction of iron (Loveland & Digby, 1984) and Al (Driscoll et al., 1985) bound with organic matter. The fraction of Fe ($0.18\text{--}0.54\text{ g kg}^{-1}$) generally decreased with depth with few exceptions. The fraction of Al followed the same trend varying from 0.49 to 1.78 g kg^{-1} . Comparison of the amount of Fe and Al extracted by pyrophosphate in different soil series indicated the largest extraction from Madhpur series while the lowest extraction from Banpara series. In comparison to Fe more Al was extracted, which might be due to the Al rich materials or because pyrophosphate is not very specific for organic complexed Al, dissolving inorganic Al compounds (like allophane and imogolite) as well (Bera, Seal, Banerjee, & Dolui, 2005).

3.2. Acid ammonium oxalate extractable Fe and Al

Oxalate dissolved both “amorphous” and “organically bound” forms of Fe and Al, but not the crystalline forms (Parfitt & Childs, 1988). Amorphous forms of Fe and Al like hydroxides and oxyhydroxides were dissolved from soil using this extractant (McKeague, Brydon, & Miles, 1971). Al compounds associated with organic matter were also extracted as amorphous alumino-silicates including allophane and imogolite (Kodama & Ross, 1991). In the study area, oxalate extractable of Fe ($0.84\text{--}2.24\text{ g kg}^{-1}$) was more in illuvial horizon of all the soil series indicating translocation of Fe. In these horizons, the fraction of Al

Table 1. Physicochemical properties of five representative soil series

Depth (cm)	pH _w (1:2.5)	pH _{KCl} (1:2.5)	EC(1:5) (dSm ⁻¹)	Org. C (gkg ⁻¹)	Particle size distribution (%)			CEC (cmol(p ⁺)kg ⁻¹)	BS (%)
					Sand	Silt	Clay		
Madhpur series (Very fine mixed hyperthermic Typic Epiaqualfs)									
0–14	6.4	5.5	0.18	5.0	24.0	46.3	29.7	13.7	71
14–38	6.9	5.8	0.12	3.2	27.7	38.6	33.7	15.0	75
38–98	7.2	6.3	0.21	1.9	18.6	33.7	47.7	22.5	79
98–150	7.5	6.4	0.21	1.7	15.2	29.1	55.7	25.6	85
Mean	7.0	6.0	0.2	2.7	21.4	36.9	41.7	19.2	78
Hargram series (Fine mixed hyperthermic Typic Endoaquepts)									
0–13	5.6	4.7	0.14	9.1	31.5	36.3	32.2	13.3	72
13–33	6.2	5.3	0.12	7.1	34.6	32.8	32.6	14.2	75
33–60	6.4	5.5	0.12	2.8	39.3	24.6	36.1	15.9	79
60–78	6.5	5.5	0.11	2.3	45.1	18.0	36.9	16.1	82
78–104	6.7	5.8	0.10	2.1	47.6	16.2	36.2	15.8	85
104–150	6.9	5.7	0.10	1.8	52.8	12.7	34.5	15.1	89
Mean	6.4	5.4	0.1	3.9	41.8	23.4	34.8	15.1	80
Karjana soil series (Fine loamy mixed hyperthermic Typic Endoaquepts)									
0–13	6.3	5.3	0.14	9.4	22.9	50.5	26.6	11.3	69
13–33	7.5	6.4	0.12	7.4	18.2	53.7	28.1	16.1	82
33–60	7.2	6.0	0.14	5.0	17.8	53.8	28.4	17.2	80
60–78	6.8	5.8	0.15	3.8	15.4	56.8	27.8	17.9	75
78–103	6.4	5.5	0.15	2.5	17.1	55.0	27.9	17.2	75
103–150	7.0	6.0	0.16	2.1	15.7	49.7	34.6	18.0	75
Mean	6.9	5.8	0.1	3.7	17.9	53.3	28.9	16.3	76
Banpara soil series (Fine loamy mixed hyperthermic Typic Endoaquepts)									
0–15	6.1	5.1	0.23	8.1	39.5	40.4	20.1	8.2	68
15–32	6.7	5.8	0.11	5.4	21.2	45.8	33.0	16.0	84
32–80	7.0	6.1	0.17	2.1	24.2	43.0	32.8	16.0	93
80–150	7.1	6.2	0.19	1.8	25.0	44.1	30.9	13.5	92
Mean	6.7	5.8	0.2	3.3	27.5	43.3	29.2	13.4	84
Khamrp soil series (Fine loamy mixed hyperthermic Aeric Endoaquepts)									
0–14	5.4	4.7	0.16	8.7	36.4	40.4	23.2	12.0	57
14–42	6.7	5.4	0.14	6.8	29.5	36.3	34.2	15.9	70
42–68	6.8	5.0	0.13	4.8	39.1	36.3	29.6	15.8	72
68–86	6.5	5.5	0.10	2.9	68.3	11.3	20.4	9.4	72
86–113	6.8	5.5	0.10	2.8	75.7	10.3	14.0	6.2	73
113–142	7.5	6.1	0.12	1.9	76.2	8.6	14.8	6.2	77
142–165	6.4	5.2	0.09	1.9	56.2	21.5	22.3	9.8	73
Mean	6.6	5.3	0.1	3.5	54.5	23.5	22.6	10.8	71

Notes: pH_w: pH in water suspension; EC: Electrical conductivity; Org. C: Organic carbon; CEC: cation exchange capacity; BS: Base saturation.

(1.61–7.19 gkg⁻¹) also followed the same trend probably due to the deposition of translocated Al-fulvate complex, protoimogolite, and hydroxy polymers of the element (Farmer, Russell, & Berrow, 1980). Hence, the difference between oxalate and pyrophosphate extractable Fe and Al gave a measure of amorphous inorganic Fe and Al (Dolui & Bera, 2001; Dolui & Chakraborty, 1998) in soils.

Table 2. Fe and Al (g kg^{-1}) extracted by different extractants in selected soil pedons

Depth (cm)	Pyrophosphate extractable (g kg^{-1})		Oxalate extractable (g kg^{-1})		Dithionate extractable (g kg^{-1})	
	Fe _{pyr}	Al _{pyro}	Fe _{oxa}	Al _{oxa}	Fe _{dith}	Al _{dith}
Madhpur series						
0-14	0.54	1.68	1.52	4.72	4.21	12.87
14-38	0.49	1.72	1.68	5.94	4.47	15.41
38-98	0.52	1.78	2.13	6.82	5.37	16.98
98-150	0.42	1.46	2.24	7.19	5.43	17.22
Mean	0.49	1.66	1.89	6.17	4.87	15.62
Hargram series						
0-13	0.46	1.34	0.94	2.65	2.27	6.24
13-33	0.48	1.32	0.98	2.74	2.43	6.56
33-60	0.36	0.98	1.26	3.12	2.86	7.10
60-78	0.34	0.87	1.24	3.04	2.92	7.14
78-104	0.29	0.67	1.20	2.64	2.82	6.98
104-150	0.24	0.56	1.21	2.68	2.98	6.75
Mean	0.36	0.96	1.14	2.81	2.71	6.80
Karjana soil series						
0-13	0.48	1.02	1.02	2.13	2.18	5.06
13-33	0.42	0.98	1.04	2.16	2.32	5.03
33-60	0.40	0.87	1.06	2.34	2.34	5.16
60-78	0.34	0.78	1.12	2.45	2.46	5.21
78-103	0.36	0.79	1.23	2.67	2.57	5.85
103-150	0.28	0.64	1.26	2.76	2.54	5.68
Mean	0.38	0.85	1.12	2.42	2.40	5.33
Banpara soil series						
0-15	0.36	0.69	0.87	1.61	2.14	4.01
15-32	0.32	0.87	0.84	2.18	2.06	5.42
32-80	0.24	0.64	0.96	2.39	2.34	5.68
80-150	0.21	0.54	1.07	2.42	2.41	5.50
Mean	0.28	0.69	0.94	2.15	2.24	5.15
Khamrp soil series						
0-14	0.42	1.08	0.86	2.07	1.94	4.34
14-42	0.38	0.96	0.85	2.01	1.96	4.56
42-68	0.32	0.74	0.94	2.01	1.84	4.02
68-86	0.34	0.82	0.90	2.16	1.98	4.78
86-113	0.26	0.88	1.14	3.64	2.01	6.84
113-142	0.21	0.56	1.24	3.89	2.12	6.74
142-165	0.18	0.49	1.30	3.02	2.23	5.32
Mean	0.32	0.79	1.03	2.69	2.01	5.23

Data (Table 3) revealed that amorphous Fe ($\text{Fe}_{\text{oxa}} - \text{Fe}_{\text{pyr}}$) in the soil ranged between 0.44 and 1.82 g kg^{-1} with a wide variation among the different soil series. Amorphous Al ($\text{Al}_{\text{oxa}} - \text{Al}_{\text{pyr}}$) of the soils varied from 0.92 to 5.73 g kg^{-1} . The data presented provided evidence that an approximate differentiation can be made between organically complexed Fe and Al and amorphous inorganic Fe and Al by selective extraction of soils with pyrophosphate and oxalate. The large apparent increase in the quantities of $\text{Fe}_{\text{oxa}} - \text{Fe}_{\text{pyr}}$ forms and $\text{Al}_{\text{oxa}} - \text{Al}_{\text{pyr}}$ forms suggested a shift toward inorganic pedogenic phases at

Table 3. Comparison of different forms of Fe and Al and their ratio

Depth (cm)	Amorphous (g kg ⁻¹)		Crystalline (g kg ⁻¹)		Degree of activation (g kg ⁻¹)		Co-migration of clay	
	Fe ₂ O ₃ (Fe _{oxa} -Fe _{pyr})	Al ₂ O ₃ (Al _{oxa} -Al _{pyr})	Fe ₂ O ₃ (Fe _{dith} -Fe _{oxa})	Al ₂ O ₃ (Al _{dith} -Al _{oxa})	Fe ₂ O ₃ (Fe _{oxa} /Fe _{dith})	Al ₂ O ₃ (Al _{oxa} /Al _{dith})	Clay/Fe _{dith}	Clay/Al _{dith}
Madhpur series								
0-14	0.98	3.04	2.69	8.15	0.36	0.37	7.05	2.31
14-38	1.19	4.22	2.79	9.47	0.38	0.39	7.54	2.19
38-98	1.61	5.04	3.24	10.16	0.40	0.40	8.88	2.81
98-150	1.82	5.73	3.19	10.03	0.41	0.42	10.26	3.23
Mean	1.40	4.51	2.98	9.45	0.39	0.39	8.43	2.63
Hargram series								
0-13	0.48	1.31	1.33	3.59	0.41	0.42	14.19	5.16
13-33	0.50	1.42	1.45	3.82	0.40	0.42	13.42	4.97
33-60	0.90	2.14	1.60	3.98	0.44	0.44	12.62	5.08
60-78	0.90	2.17	1.68	4.10	0.42	0.43	12.64	5.17
78-104	0.91	1.97	1.62	4.34	0.43	0.38	12.84	5.19
104-150	0.97	2.12	1.77	4.07	0.41	0.40	11.58	5.11
Mean	0.78	1.86	1.58	3.98	0.42	0.41	12.88	5.11
Karjana soil series								
0-13	0.54	1.11	1.16	2.93	0.47	0.42	12.20	5.26
13-33	0.62	1.18	1.28	2.87	0.45	0.43	12.11	5.59
33-60	0.66	1.47	1.28	2.82	0.45	0.45	12.14	5.50
60-78	0.78	1.67	1.34	2.76	0.46	0.47	11.30	5.34
78-103	0.87	1.88	1.34	3.18	0.48	0.46	10.86	4.77
103-150	0.98	2.12	1.28	2.92	0.50	0.49	13.62	6.09
Mean	0.74	1.57	1.28	2.91	0.47	0.45	12.04	5.42
Banpara soil series								
0-15	0.51	0.92	1.27	2.40	0.41	0.40	9.39	5.01
15-32	0.52	1.31	1.22	3.24	0.41	0.40	16.02	6.09
32-80	0.72	1.75	1.38	3.29	0.41	0.42	14.02	5.77
80-150	0.86	1.88	1.34	3.08	0.44	0.44	12.82	5.62
Mean	0.65	1.47	1.30	3.00	0.42	0.42	13.06	5.62
Khamrp soil series								
0-14	0.44	0.99	1.08	2.27	0.44	0.48	11.96	5.35
14-42	0.47	1.05	1.11	2.55	0.43	0.44	17.45	7.50
42-68	0.62	1.27	0.90	2.01	0.51	0.50	16.09	7.36
68-86	0.56	1.34	1.08	2.62	0.45	0.45	10.30	4.27
86-113	0.88	2.76	0.87	3.20	0.57	0.53	6.97	2.05
113-142	1.03	3.33	0.88	2.85	0.58	0.58	6.98	2.20
142-165	1.12	2.53	0.93	2.30	0.58	0.57	10.00	4.19
Mean	0.73	1.90	0.98	2.54	0.51	0.51	11.39	4.70

the expense of organically bound phase (Jersak, McColl, & Hetzel, 1992). Though non-crystalline Fe and Al do not have a definite composition or structure and are only poorly defined, there is probably no precise differentiation between crystalline and non-crystalline material. The microprobe investigation of Norrish and Rossier (1983) indicated that amorphous or microcrystalline Fe and Al oxides were unlikely to exist as separate entities in soils.

Table 4. Correlation matrix for different forms of iron and aluminum and physicochemical properties of selected soil series

Y/X	pH(H ₂ O)	Org. C	Clay	CEC	Fe _{pyr}	Al _{pyr}	Fe _{oxa}	Al _{oxa}	Fe _{dith}
Fe _{pyr}	-0.258	0.557**	0.361	0.447*					
Al _{pyr}	-0.096	0.257	0.468*	0.438*	0.889**				
Fe _{oxa}	0.433*	-0.487**	0.638**	0.591**	0.307	0.579**			
Al _{oxa}	0.385*	-0.406*	0.549**	0.455*	0.372	0.696**	0.954**		
Fe _{dith}	0.335	-0.349	0.741**	0.646**	0.481**	0.719**	0.943**	0.924**	
Al _{dith}	0.330	-0.330	0.636**	0.519**	0.484**	0.773**	0.931**	0.974**	0.972**

*Level of significance at 5% is 0.374.

**Level of significance at 1% is 0.478.

3.3. Dithionite-citrate-bicarbonate (DCB) extractable Fe and Al

Two dissolution reagents in particular are common (i) acid ammonium oxalate (pH 3) for extraction of non-crystalline Fe and Al and (ii) DCB for extraction of non-crystalline plus crystalline Fe and Al. In the DCB procedure (Mehra & Jackson, 1960), dithionite is a powerful reductant, bicarbonate buffers the system (pH 7–9) and sodium citrate prevents the reprecipitation of dissolved Fe and Al (Borggard, 1988). The procedure dissolved both non-crystalline and crystalline Fe and Al oxides and the extracts may also include small amount of water soluble, exchangeable, and organically bound Fe and Al along with limited amount of Fe and Al bearing silicates (Borggard, 1988). The difference between the values obtained by the two methods represented the amount of Fe and Al present in definite crystalline forms (Dolui & Chakraborty, 1998; Dolui, Chandran, & Nayek, 1988). The Fe_{dith} of the soils showed a variation of 1.84–5.43 gkg⁻¹ whereas Al_{dith} of the soils showed a variation of 4.01–17.22 gkg⁻¹ among the different soil series.

3.4. Co-migration of clay

The co-migration of Fe and Al with clay in the different soil profiles is evidenced by the ratios of clay/Fe_{dith} and clay/Al_{dith}. The ratios increased with increase in depth in Madhpur soil series, which can be corroborated with the observations of Dolui and Mazumdar (2003) and Seal, Bera, Bhattacharyya, Mukhopadhyay, and Giri (2006) in alfisols of India. According to Dolui and Bera (2001) these ratios increased with depth within the profile where horizon formation was well expressed and established. However, in Hargram and Karjana soil series, the clay/Fe_{dith} and clay/Al_{dith} ratios showed somewhat decreasing trend with few exceptions. This suggested preferential migration of clay over Fe and Al oxides, which were partially independent of clay movement (Bera et al., 2005). In Banpara and Khamrp soil series, the ratios did not follow any consistent pattern with relevance to the different horizons indicating that horizon formation was still in progress.

3.5. Different forms of Fe and Al and their ratios

In all the soils, DCB extracted Fe and Al values were higher than the acid ammonium oxalate extracted values indicating that a considerable fraction was present in crystalline form. Crystalline Fe (Fe_{dith}–Fe_{oxa}) in the soils ranged between 0.87 and 3.24 gkg⁻¹ while crystalline Al (Al_{dith}–Al_{oxa}) in the soils varied from 2.01 to 10.16 gkg⁻¹.

The crystalline Fe and Al oxides increased at the expense of the poorly crystalline forms with increasing soil age as indicated by the ratios of Fe_{oxa}/Fe_{dith} and Al_{oxa}/Al_{dith} (Mahaney & Fahey, 1988). According to Mahaney, Hancock, and Sanmugasdas (1991), soils with high ratios were younger soils, whereas low ratios indicated older soils. The mean Fe and Al ratios of Madhpur soil series were comparatively lower than other soil series, varying from 0.36 to 0.41 and 0.37 to 0.42, respectively. This suggested that Madhpur soil series could be comparatively older or weathering could have progressed relatively well to an advanced stage in comparison to other soil series. On the other hand, Khamrp soil series (Fe_{oxa}/Fe_{dith} and Al_{oxa}/Al_{dith} ratios varied from 0.43 to 0.58 and 0.44 to 0.58,

respectively) might be comparatively younger in the process of soil development, while in the other soils pedogenic development represented an intermediary stage with respect to Madhpur and Khamrp soil series.

3.6. Interrelationship among different forms of Fe and Al

Fe_{pyr} was positively and significantly correlated with Al_{pyr} (0.889**), Fe_{dith} (0.481**), and Al_{dith} (0.484**) while Al_{pyr} was positively and significantly correlated with Fe_{oxa} (0.579**), Al_{oxa} (0.696**), Fe_{dith} (0.719**), and Al_{dith} (0.773**). Similar relationships were reported by Dolui and Bera (2001), Dolui, Chandran, and Nayek (1987) while working with the tropical soils of India. Highly positive correlation of Fe_{dith} with Fe_{pyr} (0.481**) and Fe_{oxa} (0.943**) as well as Al_{dith} with Al_{pyr} (0.773**) and Al_{oxa} (0.974**) suggested that their relative distribution in soils were functions of the same pedological factor and more importantly all the extractants removed essentially the same forms of Fe and Al (Dolui & Chattopadhyay, 1997).

3.7. Relationship with soil properties

pH of the soils was found to have significant positive correlation with oxalate extractable fractions of Fe (0.433*) and Al (0.385*). This may be correlated with the fact that increase in soil pH increased the adsorptive affinity of clay and iron oxides, therefore reducing the concentration of iron in labile pool (Dolui & Bera, 2001). Organic carbon was positively and significantly correlated with Fe_{pyro} (0.557**) while negatively and significantly correlated with Fe_{oxa} (-0.487**) and Al_{oxa} (-0.406*). Similar observation was also reported by Seal et al. (2006). The highly significant correlation of clay with Fe_{oxa} (0.638**), Fe_{dith} (0.741**), Al_{oxa} (0.549**), and Al_{dith} (0.636**) indicated that these fractions might have been present within the crystal lattice or sorbed on the surface and the interlayer of clay and iron oxides (Dolui & Chattopadhyay, 1997; Howkes & Webb, 1962). Cation exchange capacity showed significant positive correlation with pyrophosphate (0.447* and 0.438*, respectively), oxalate (0.591** and 0.455*, respectively), and dithionate (0.646** and 0.519**, respectively) extractable Fe and Al. Cation exchange capacity indicated the total negative charge of the colloid and in soil may be correlated to the availability of iron and aluminum from charge sites (Bera et al., 2005).

4. Elemental composition

The SiO_2 content of soils ranged between 49.20 and 61.39%, comprising more than half of the total mineral matter (Table 5). These variations in SiO_2 content were mainly associated with the nature of parent material and variation in topography (Sidhu, Ghosh, & Manjaiah, 2000). The SiO_2 content generally tend to decrease down the profile in all the soil series with some exceptions. On the contrary, the chemical composition of the soils indicated higher accumulation of Al_2O_3 and Fe_2O_3 in the subsurface horizons, which generally increased down the profile in all the soil series with some exceptions. The accumulation of Al_2O_3 (23.20–32.16%) and Fe_2O_3 (7.47–14.87%) in the cambic/argillic horizons of the soils indicated the mobilization of these elements within the soil profile due to active soil forming processes (Sidhu et al., 2000). The movement of manganese oxide and its distribution in the soil profiles were significant criteria in pedogenic study as it behaved more or less like free iron oxides (Elahi, Hossin, & Kamal, 1996). Mn_2O_3 content in the soils decreased down the profile with few exceptions while the content of CaO (0.04–0.25%) and MgO (0.16–0.37%) varied within a narrow range. There were also not much variations in ZnO (0.020–0.204%) and CuO (0.001–0.005%) content throughout the profile in all the soil series. The total base percentage (CaO + MgO + Na_2O + K_2O) in Madhpur soil series (mean 2.38%) was lower than that of the other soil series. This indicated advanced stage of weathering in Madhpur soil series when compared with other soil series.

4.1. Molar ratios

Chemical changes occurring in the selected soil series as a result of weathering were determined from changes in the molar ratios of mobile to resistant oxides. These ratios were calculated on the assumption that the resistant oxides of Fe and Al were immobile in the weathering environment (Colman, 1982). Therefore, any changes in the ratios that did occur reflected the extent to which the more mobile oxides of Ca, Mg, Na, and K were removed. In general as weathering progressed, oxide molar ratios decreased as greater quantities of mobile oxides were removed from the weathering

Table 5. Chemical composition (ignition basis) of the soils

Depth (cm)	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	Mn ₂ O ₃	CaO	MgO	K ₂ O	Na ₂ O	ZnO	CuO
	←—————(%)—————→									
Madhpur series										
0-14	53.53	27.89	13.73	0.17	0.04	0.16	1.55	0.46	0.204	0.004
14-38	52.50	30.05	13.65	0.16	0.06	0.17	1.64	0.48	0.172	0.004
38-98	50.15	31.29	14.64	0.09	0.09	0.18	1.75	0.54	0.158	0.005
98-150	49.20	32.16	14.87	0.10	0.09	0.19	1.63	0.57	0.151	0.004
Mean	51.35	30.35	14.22	0.13	0.07	0.18	1.64	0.51	0.171	0.004
Hargram series										
0-13	53.56	27.01	12.86	0.11	0.25	0.31	2.12	1.12	0.020	0.001
13-33	53.92	25.84	12.95	0.13	0.15	0.34	2.50	1.60	0.030	0.001
33-60	52.86	29.35	10.86	0.11	0.15	0.35	2.77	1.65	0.020	0.003
60-78	53.74	28.82	10.75	0.11	0.13	0.39	2.36	1.65	0.020	0.001
78-104	53.49	29.58	12.09	0.12	0.18	0.36	1.95	2.12	0.020	0.002
104-150	54.21	28.92	11.89	0.14	0.20	0.37	1.90	1.93	0.020	0.002
Mean	53.63	28.25	11.90	0.12	0.18	0.35	2.27	1.68	0.022	0.002
Karjana soil series										
0-13	55.01	25.79	11.94	0.09	0.10	0.37	2.24	2.01	0.020	0.002
13-33	54.46	26.35	12.16	0.10	0.10	0.34	2.23	2.11	0.020	0.002
33-60	53.14	26.62	12.14	0.10	0.10	0.36	2.34	2.14	0.030	0.001
60-78	54.08	27.18	12.56	0.10	0.11	0.36	2.36	2.21	0.030	0.001
78-103	52.19	28.40	13.02	0.11	0.11	0.32	2.21	2.94	0.020	0.002
103-150	52.06	28.32	12.87	0.11	0.12	0.32	2.40	2.87	0.020	0.001
Mean	53.49	27.11	12.45	0.10	0.11	0.35	2.30	2.38	0.023	0.002
Banpara soil series										
0-15	56.64	24.50	12.87	0.13	0.09	0.32	2.38	1.72	0.020	0.005
15-32	56.92	26.84	9.35	0.13	0.12	0.34	2.76	2.20	0.030	0.003
32-80	54.74	27.96	10.75	0.11	0.13	0.34	2.92	2.25	0.020	0.002
80-150	53.49	28.18	12.09	0.11	0.15	0.36	3.08	2.30	0.020	0.002
Mean	55.45	26.87	11.27	0.12	0.12	0.34	2.79	2.12	0.023	0.003
Khamrp soil series										
0-14	56.64	23.70	10.60	0.08	0.12	0.30	3.24	3.13	0.020	0.001
14-42	56.06	24.03	10.97	0.09	0.12	0.32	3.26	3.24	0.030	0.001
42-68	55.26	23.95	12.15	0.11	0.11	0.31	3.07	3.24	0.030	0.001
68-86	57.42	23.65	10.86	0.05	0.12	0.32	3.56	3.32	0.030	0.003
86-113	61.39	23.20	7.47	0.05	0.12	0.26	3.39	2.94	0.020	0.001
113-142	61.12	23.62	7.74	0.08	0.11	0.29	3.48	3.01	0.020	0.004
142-165	57.00	23.50	10.90	0.07	0.10	0.32	3.42	3.01	0.020	0.004
Mean	57.84	23.66	10.10	0.08	0.11	0.30	3.35	3.13	0.024	0.002

profile (Birkeland, 1984). The molar ratios SiO₂:R₂O₃ (R₂O₃ = Fe₂O₃ + Al₂O₃), SiO₂:Al₂O₃, CaO:MgO, Al₂O₃:Fe₂O₃, CaO + MgO:R₂O₃, Na₂O + K₂O:R₂O₃, total bases (CaO + MgO + Na₂O + K₂O):R₂O₃, (Table 6) were determined to evaluate most comprehensively the extent of chemical changes occurring in the Vindyan alluvial soils in subtropical environment. According to Sharma, Totawat, and Shyampura (1996) relative accumulation of Fe was considered as one of the most important index for understanding the stage of weathering and the impact of topography on the weatherability of soil.

Table 6. Different ratios and weathering index of selected soil series

Depth (cm)	SiO ₂ /R ₂ O ₃	SiO ₂ /Al ₂ O ₃	Al ₂ O ₃ /Fe ₂ O ₃	CaO/MgO	(CaO + MgO)/R ₂ O ₃	(Na ₂ O + K ₂ O)/R ₂ O ₃	Total bases/R ₂ O ₃	Weathering index
	← Ratio →							
Madhpur series								
0-14	2.48	3.26	3.18	0.18	0.013	0.07	0.08	2.26
14-38	2.30	2.96	3.45	0.25	0.014	0.07	0.08	2.39
38-98	2.09	2.72	3.35	0.36	0.015	0.07	0.08	2.66
98-150	2.00	2.60	3.39	0.34	0.015	0.06	0.08	2.63
Mean	2.22	2.88	3.34	0.29	0.014	0.07	0.08	2.48
Hargram series								
0-13	2.58	3.36	3.29	0.58	0.035	0.12	0.15	4.13
13-33	2.68	3.54	3.13	0.32	0.033	0.16	0.19	4.95
33-60	2.47	3.06	4.23	0.31	0.032	0.16	0.19	5.22
60-78	2.56	3.16	4.20	0.24	0.034	0.15	0.18	4.92
78-104	2.43	3.07	3.83	0.36	0.033	0.15	0.18	5.12
104-150	2.52	3.18	3.81	0.39	0.036	0.14	0.18	4.89
Mean	2.50	3.20	3.72	0.36	0.034	0.15	0.18	4.87
Karjana soil series								
0-13	2.79	3.62	3.38	0.19	0.033	0.17	0.20	5.18
13-33	2.71	3.51	3.39	0.21	0.031	0.17	0.20	5.24
33-60	2.62	3.39	3.43	0.20	0.032	0.18	0.21	5.48
60-78	2.61	3.38	3.39	0.22	0.032	0.18	0.21	5.49
78-103	2.41	3.12	3.42	0.25	0.027	0.20	0.22	6.23
103-150	2.42	3.12	3.45	0.27	0.028	0.20	0.23	6.33
Mean	2.59	3.35	3.41	0.22	0.030	0.18	0.21	5.66
Banpara soil series								
0-15	2.94	3.92	2.98	0.20	0.030	0.17	0.19	4.77
15-32	2.94	3.60	4.50	0.25	0.033	0.20	0.23	5.66
32-80	2.67	3.32	4.07	0.27	0.031	0.20	0.23	5.93
80-150	2.53	3.22	3.65	0.30	0.033	0.20	0.23	6.21
Mean	2.77	3.52	3.74	0.26	0.032	0.19	0.22	5.64
Khamrp soil series								
0-14	3.15	4.06	3.50	0.29	0.032	0.28	0.32	7.15
14-42	3.07	3.96	3.43	0.27	0.033	0.29	0.32	7.34
42-68	2.96	3.92	3.09	0.26	0.031	0.27	0.30	7.21
68-86	3.19	4.12	3.41	0.27	0.034	0.30	0.34	7.55
86-113	3.72	4.49	4.86	0.33	0.031	0.30	0.34	6.70
113-142	3.63	4.39	4.78	0.27	0.033	0.31	0.34	6.87
142-165	3.18	4.12	3.38	0.22	0.033	0.28	0.32	7.12
Mean	3.27	4.15	3.67	0.27	0.032	0.29	0.32	7.13

Decrease in molar ratios of SiO₂/Al₂O₃ and SiO₂/R₂O₃ were noticed down the profile in all the soil series excluding few anomalies in Hargram and Khamrp soil series. The trend exhibited similarities with the observations of Walia and Rao (1996), Kudrat, Manchanda, Prasad, Saxena, and Tayal (1995) and Patil and Dasog (1999). The data also pointed out that SiO₂/Al₂O₃ and SiO₂/R₂O₃ molar ratios in

Madhpur soil series (2.60–3.26 with mean 2.88 and 2.00–2.48, mean 2.22, respectively) were comparatively lower than all other soil series (3.06–4.49 and 2.41–3.72, respectively) indicating that weathering process was at a more advanced stage in Madhpur soil series as compared to the other soil series (Burt et al., 2003; Msanya, Kaaya, Araki, Otsuka, & Nyadzi, 2003).

Alkali earth:resistate ($\text{CaO} + \text{MgO}:\text{R}_2\text{O}_3$) ratio in Madhpur soil series increased down the profile suggesting loss of calcium and magnesium from the upper horizon. However, change of ratios in other soil series did not follow any specific trend. Alkali:resistate ($\text{Na}_2\text{O} + \text{K}_2\text{O}:\text{R}_2\text{O}_3$) ratios ranged between 0.33 and 0.60, while total bases ($\text{CaO} + \text{MgO} + \text{Na}_2\text{O} + \text{K}_2\text{O}$):resistate (R_2O_3) ratios varied from 0.08 to 0.34, respectively, in the different soil series.

4.2. Weathering index

The average weathering index was the highest (7.13) in Khamrp soil series and the lowest (2.48) in Madhpur soil series. This indicated that extent of weathering was comparatively more in Madhpur soil series since lower value of weathering index indicated more exposure to intense weathering (Walia & Rao, 1999). The depth-wise pattern of weathering index showed high values down the profile, which indicated the fall in weathering intensity with depth (Evans & Cameron, 1979). In these alluvial-derived soils, the increase in weathering index was noticed to be erratic in most of the pedons with no specific trend of profile development, which manifested their juvenile stage of pedogenesis (Walia & Rao, 1999). This was expected because their source of materials was mostly transported sediments, which were deposited in different cycles. Similar observations were also reported by Bhowmick, Bera, Seal and Mukhopadhyay (2004) and Raghu Mohan and Rao (1981) in some alluvial-derived soils of India.

5. Conclusion

To ascertain the degree of soil development, two-way approaches were taken up under the present investigation i.e. assessment of different forms of Fe, Al, and their ratios as well as elemental composition of soil. Assessment revealed relative higher content of crystalline Fe and Al oxides vis-a-vis their low ratio values only in Madhpur soil series, which indicated that the series could be comparatively older or weathering could have progressed relatively well to an advanced stage here when compared with other soil series. Elemental composition, sesquioxide ratio as well as weathering indices only corroborated the above findings, revealing the juvenile stage of pedogenesis in all the other pedons and also demonstrated that weathering was still active in these soils.

Funding

The authors received no direct funding for this research.

Author details

Ranjan Bera¹
E-mail: bera.ranjan@gmail.com
Antara Seal¹
E-mail: antaraseal77@gmail.com
Thako Hari Das²
E-mail: acpsb02@yahoo.co.in
Dipak Sarkar²
E-mail: thdasnbss@gmail.com
Ashis Kumar Chatterjee¹
E-mail: dipaksarkar022@gmail.com

¹ Department of Soil Science, Visva Bharati University, Santiniketan 731236, West Bengal, India.

² NBSS & LUP, Block-DK, Sector-II, Salt Lake, Kolkata 700091, West Bengal, India.

Citation information

Cite this article as: Characterization of soils in terms of pedological variability under different physiography of Damodar command area (part), West Bengal, India, Ranjan Bera, Antara Seal, Thako Hari Das, Dipak Sarkar & Ashis Kumar Chatterjee, *Cogent Food & Agriculture* (2015), 1: 1014719.

Cover image

Source: Authors.

References

- Agriculture Canada. (1984). Analytical methods manual. In B. H. Sheldrick (Ed.), *LLRI contribution* (pp. 84–90). Ottawa: Land Resource Research Institute.
- Bera, R., Seal, A., Banerjee, M., & Dolui, A. K. (2005). Nature and profile distribution of iron and aluminum in relation to pedogenic processes in some soils developed under tropical environment in India. *Environmental Geology*, 47, 241–245.
<http://dx.doi.org/10.1007/s00254-004-1149-2>
- Bhowmick, A. K., Bera, R., Seal, A., & Mukhopadhyay, K. (2004). Pedological variability and classification of some acid soils of Bihar. *Indian Journal of Landscape Systems and Ecological Studies*, 27, 186–192.
- Birkeland, P. W. (1984). *Soils and geomorphology*. New York, NY: Oxford.
- Borggard, O. K. (1988). Phase identification by selective dissolution technique. In S. W. Stucki, B. A. Goodman, & U. Schwertmann (Eds.), *Iron in soils and clay minerals* (pp. 93–98). Boston, MA: D. Reidel.
- Burt, R., Wilson, M. A., Mays, M. D., & Lee, C. W. (2003). Major and trace elements of selected pedons in the USA. *Journal of Environment Quality*, 32, 2109–2121.
<http://dx.doi.org/10.2134/jeq2003.2109>

- Canada Soil Survey Committee (CSSC) subcommittee on methods of analysis. (1978). In J. A. McKeague (Ed.), *Manual of soil sampling and methods of analysis* (pp. 98–106). Ottawa: Canadian Society of Soil Science.
- Colman, S. M. (1982). *Chemical weathering of basalts and andesites: Evidence from weathering rinds* (Professional Paper, p. 1246). US Geological Survey (p. 52). USGS Publications.
- Dolui, A. K., Bera, R. (2001). Relation between iron forms and pedogenic processes in some alfisols of Orissa, India. *Agrochimica*, XLV, 161–170.
- Dolui, A. K., & Chakraborty, N. (1998). Differentiation of forms of extractable aluminium in relation to pedogenic processes in two alfisols and an entisols. *International Journal of Tropical Agricultural*, 16, 257–266.
- Dolui, A. K., Chandran, P., & Nayek, A. K. (1987). Comparative studies on different forms of aluminium using various extractants in relation to pedogenic processes in soil profiles. *Journal of the Indian Society of Soil Science*, 35, 103–108.
- Dolui, A. K., Chandran, P., & Nayek, A. K. (1988). Nature and profile distribution of iron in some soil series of Jalpaiguri district, West Bengal. *The Proceedings of the National Academy of Sciences India*, 58(A), 1–6.
- Dolui, A. K., & Chattopadhyay, P. P. (1997). Extraction of forms of iron from some soil series of West Bengal and Bihar. *Agropedology*, 7, 44–47.
- Dolui, A. K., & Mazumdar, A. (2003). Relation between forms of aluminium and pedogenic processes in some alfisols of Bihar and West Bengal. *Crop Research Hisar*, 25, 69–77.
- Driscoll, C. J., van Breemen, N., & Mulder, J. (1985). Aluminium chemistry in a forested spodosol. *Soil Science Society of America Journal*, 49, 437–444. <http://dx.doi.org/10.2136/sssaj1985.03615995004900020033x>
- Elahi, S. F., Hossin, M. F., & Kamal, A. S. M. M. (1996). Characteristics of some soils developed on Madhupur clay in Bangladesh. *Journal of the Indian Society of Soil Science*, 44, 482–488.
- Evans, L. J., & Cameron, B. H. (1979). A chronosequence of soils developed from granitic morainal material, Baffin Island, N.W.T. *Canadian Journal of Soil Science*, 59, 203–210. <http://dx.doi.org/10.4141/cjss79-020>
- Farmer, V. C., Russell, J. D., & Berrow, M. L. (1980). Imogolite and proto-imogolite allophane in spodic horizons: Evidence for a mobile aluminium silicate complex in podzol formation. *Journal of Soil Science*, 31, 673–684. <http://dx.doi.org/10.1111/ejs.1980.31.issue-4>
- Hawkes, H. E., Webb, J. (1962). *Geochemistry of mineral exploration* (p. 9). New York, NY: Harper and Row.
- Jackson, M. L. (1973). *Soil chemical analysis*. New Delhi: Prentice Hall of India.
- Jersak, J. M., McColl, J. G., & Hetzel, J. F. (1992). Changes in extractability of iron, aluminium and silicon and dispersibility by storage of California forest soils. *Communications in Soil Science and Plant Analysis*, 23, 993–1018. <http://dx.doi.org/10.1080/00103629209368645>
- Kabata-Pendias, A., & Pendias, H. (1992). *Trace elements in soils and plants* (2nd ed.). Boca Rotan, FL: CRC Press.
- Kodama, H., & Ross, G. J. (1991). Tiron dissolution method used to remove and characterize inorganic components in soils. *Soil Science Society of America Journal*, 55, 1180–1187. <http://dx.doi.org/10.2136/sssaj1991.03615995005500040047x>
- Kudrat, M., Manchanda, M. L., Prasad, J., Saxena, P. D., & Tayal, H. D. (1995). Morphology, physico-chemical characteristics and classification of lateritic soils of a part of Ajoy catchment in West Bengal. *Agropedology*, 5, 9–14.
- Loveland, P. J., & Digby, P. (1984). The extraction of Fe and Al by 0.1 M pyrophosphate solutions: A comparison of some techniques. *Journal of Soil Science*, 35, 243–250. <http://dx.doi.org/10.1111/ejs.1984.35.issue-2>
- Mahaney, W. C., & Fabey, B. D. (1988). Extractable Fe and Al in late Pleistocene and Holocene paleosols on Niwot Ridge, Colorado Front Range. *Catena*, 15, 17–26. [http://dx.doi.org/10.1016/0341-8162\(88\)90013-6](http://dx.doi.org/10.1016/0341-8162(88)90013-6)
- Mahaney, W. C., Hancock, R. G. V., & Sanmugadas, K. (1991). Extractable Fe-Al and geochemistry of late Pleistocene Paleosol in the Dalijia Shan, Western China. *Journal of Southeast Asian Earth Sciences*, 6, 75–82. [http://dx.doi.org/10.1016/0743-9547\(91\)90098-1](http://dx.doi.org/10.1016/0743-9547(91)90098-1)
- Mausbach, M., Richardson J. L. (1994). Biogeochemical processes in hydric soil formation. In *Current topics in wetland biogeochemistry* (Vol. 1, pp. 68–127). Baton Rouge, LA: Wetland Biogeochem. Inst., Louisiana State Univ.
- McKeague, J. A., Brydon, J. E., & Miles, N. M. (1971). Differentiation of forms of extractable iron and aluminum in soils. *Soil Science Society of America Journal*, 35, 33–38. <http://dx.doi.org/10.2136/sssaj1971.03615995003500010016x>
- McKeague, J. A., & Day, J. H. (1966). Dithionite and oxalate-extractable Fe and Al as aids in differentiating various classes of soils. *Canadian Journal of Soil Science*, 46, 13–22. <http://dx.doi.org/10.4141/cjss66-003>
- Mehra, O. P., & Jackson, M. L. (1960). Iron oxide removal from soils and clays by a dithionite citrate system buffered with sodium bicarbonate. *Clays and Clay Miner*, 7, 317–327.
- Msanya, B. M., Kaaya, A. K., Araki, S., Otsuka, H., & Nyadzzi, G. I. (2003). Pedological characteristics, general fertility and classification of some benchmark soils of Morogoro District, Tanzania. *African Journal of Science and Technology*, 4, 101–112.
- Muhs, D. R., Bettis, E. A., Been, J., & McGeehin, J. P. (2001). Impact of climate and parent material on chemical weathering in loess-derived soils of the Mississippi river valley. *Soil Science Society of America Journal*, 65, 1761–1777. <http://dx.doi.org/10.2136/sssaj2001.1761>
- Norrish, K., & Rossier, H. (1983). Mineral phosphate. In *Soils: An Australian viewpoint* (pp. 335–361). Sponsored by the Division of Soils, Commonwealth Scientific and Industrial Research Organization. Melbourne: Academic Press. CSIRO/London, UK/Australia.
- Parfitt, R. L., & Childs, C. W. (1988). Estimation of forms of Fe and Al—A review, and analysis of contrasting soils by dissolution and Mossbauer methods. *Australian Journal of Soil Research*, 26, 121–144. <http://dx.doi.org/10.1071/SR9880121>
- Patil, P. L., & Dasog, G. S. (1999). Genesis and classification of ferruginous soils in Western Ghat and Coastal region of North Karnataka. *Agropedology*, 9, 1–5.
- Raghu Mohan, N. G., & Rao, A. E. V. (1981). Morphogenesis and classification of ferruginous soils of Goa II. Physico-chemical properties and mineralogy of epipedons and endopedons of Goa. *Mysore journal of Agricultural Science*, 15, 29–39.
- Seal, A., Bera, R., Bhattacharyya, P., Mukhopadhyay, K., & Giri, R. (2006). Degree of soil development in some alfisols of subtropical India with special reference to the nature and distribution of Fe and Al. *International Journal of Agricultural Research*, 1(4), 1–7.
- Sharma, S. S., Totawat, K. L., & Shyampura, R. L. (1996). Characterization and classification of soils in a toposequence over basaltic terrain. *Journal of the Indian Society of Soil Science*, 44, 470–475.
- Sidhu, G. S., Ghosh, S. K., & Manjajiah, K. M. (2000). Pedological variabilities and classification of some dominant soils of Aravallies-Yamuna river transect in semi-arid tract of Haryana. *Agropedology*, 10, 80–87.
- Walia, C. S., & Rao, Y. S. (1996). Genesis, characteristics and taxonomic classification of some soils in Bundelkhand

region of Uttar Pradesh. *Journal of the Indian Society of Soil Science*, 44, 476–481.
Walia, C. S., & Rao, Y. S. (1999). Forms of iron and weathering index of some typical pedons of Bundelkhand region in relation to physiography and parent material. *Agropedology*, 9, 82–87.

Wilson, M. A., Burt, R., Sobecki, T. M., Engel, R. J., & Hipple, K. (1996). Soil properties and genesis of pans in till-derived Andisols, Olympic Peninsula, Washington. *Soil Science Society of America Journal*, 60, 206–218. <http://dx.doi.org/10.2136/sssaj1996.0361599500600010033x>



© 2015 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



Cogent Food & Agriculture (ISSN: 2331-1932) is published by Cogent OA, part of Taylor & Francis Group.

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

