



## Profile and Morphology of Fungal Aerosols Characterized by Field Emission Scanning Electron Microscopy (FESEM)

Komlavi Anani Afanou, Anne Straumfors, Asbjørn Skogstad, Ida Skaar, Linda Hjeljord, Øivind Skare, Brett James Green, Arne Tronsmo & Wijnand Eduard

To cite this article: Komlavi Anani Afanou, Anne Straumfors, Asbjørn Skogstad, Ida Skaar, Linda Hjeljord, Øivind Skare, Brett James Green, Arne Tronsmo & Wijnand Eduard (2015) Profile and Morphology of Fungal Aerosols Characterized by Field Emission Scanning Electron Microscopy (FESEM), *Aerosol Science and Technology*, 49:6, 423-435, DOI: [10.1080/02786826.2015.1040486](https://doi.org/10.1080/02786826.2015.1040486)

To link to this article: <https://doi.org/10.1080/02786826.2015.1040486>



© STAMI National Institute of Occupational Health. Published with license by Taylor & Francis© STAMI National Institute of Occupational Health.



[View supplementary material](#)



Published online: 11 May 2015.



[Submit your article to this journal](#)



Article views: 1160



[View related articles](#)



[View Crossmark data](#)



Citing articles: 11 [View citing articles](#)



# Profile and Morphology of Fungal Aerosols Characterized by Field Emission Scanning Electron Microscopy (FESEM)

Komlavi Anani Afanou,<sup>1</sup> Anne Straumfors,<sup>1</sup> Ashbjørn Skogstad,<sup>1</sup> Ida Skaar,<sup>2</sup> Linda Hjeljord,<sup>3</sup> Øivind Skare,<sup>1</sup> Brett James Green,<sup>4</sup> Arne Tronsmo,<sup>3</sup> and Wijnand Eduard<sup>1</sup>

<sup>1</sup>National Institute of Occupational Health, Department of Chemical and Biological Work Environment, Oslo, Norway

<sup>2</sup>Norwegian Veterinary Institute, Section of Mycology, Oslo, Norway

<sup>3</sup>Norwegian University of Life Science, Institute of Chemistry, Biotechnology and Food Science, Ås, Norway

<sup>4</sup>National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, Allergy and Clinical Immunology Branch, Health Effect Laboratory Division, Morgantown, West Virginia, USA

---

Fungal aerosols consist of spores and fragments with diverse array of morphologies; however, the size, shape, and origin of the constituents require further characterization. In this study, we characterize the profile of aerosols generated from *Aspergillus fumigatus*, *A. versicolor*, and *Penicillium chrysogenum* grown for 8 weeks on gypsum boards. Fungal particles were aerosolized at 12 and 20 L min<sup>-1</sup> using the Fungal Spore Source Strength Tester (FSSST) and the Stami particle generator (SPG). Collected particles were analyzed with field emission scanning electron microscopy (FESEM). We observed spore particle fraction consisting of single spores and spore aggregates in four size categories, and a fragment fraction that contained submicronic fragments and three size categories of larger fragments. Single spores dominated the aerosols from *A. fumigatus* (median: 53%), while the submicronic fragment fraction was the highest in the aerosols collected from *A. versicolor* (median: 34%) and *P. chrysogenum* (median: 31%). Morphological characteristics showed near spherical particles that were only single spores, oblong particles that comprise some spore aggregates and fragments (<3.5 μm), and fiber-like particles that regroup chained spore aggregates and fragments (>3.5 μm). Further, the near spherical particles dominated the aerosols from *A. fumigatus* (median: 53%), while oblong particles were dominant in the aerosols from *A. versicolor* (68%) and *P. chrysogenum* (55%). Fiber-like particles represented 21% and

24% of the aerosols from *A. versicolor* and *P. chrysogenum*, respectively. This study shows that fungal particles of various size, shape, and origin are aerosolized, and supports the need to include a broader range of particle types in fungal exposure assessment.

---

## INTRODUCTION

Indoor environments with a high moisture content often result in saprophytic fungal colonization of structural building materials. Fungal colonization of these indoor environments in combination with indoor dampness has been associated with adverse health outcomes (Institute of Medicine 2004; World Health Organization 2009). However, epidemiological and exposure studies including the assessment of fungal spore exposure in indoor air could not confirm such an association (Eduard 2009). Epidemiological studies based on traditional methods of identification and quantification of fungal spores or fungal colony forming units have not provided insight into health effects from other airborne fungal particles such as fragments of spores and hyphae.

The composition of fungal aerosols is diverse and comprises particles of various sizes, shapes and origin (spores and mycelia), thus with varying aerodynamic behavior (Lacey 1991). Single spores from many molds encountered in indoor environments are globose or ellipsoidal with smooth or ornamented surface (Reponen et al. 2001) while chained spore aggregates are often elongated and fiber-like. Moreover, hyphal fragments have various morphologies: they can be tubular, elongate or fiber-like and irregular.

The aerodynamic diameter of particles is commonly used in the prediction of particle motion and deposition in human

---

© STAMI National Institute of Occupational Health

This is an Open Access article. Non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly attributed, cited, and is not altered, transformed, or built upon in any way, is permitted. The moral rights of the named authors(s) have been asserted.

Received 15 November 2014; accepted 1 April 2015.

Address correspondence to Wijnand Eduard, National Institute of Occupational Health, Department of Chemical and Biological Work Environment, Gydasvei, Oslo 0033, Norway. E-mail: wijnard.eduard@stami.no

respiratory tract. Shape also influences motion and deposition of particles in human respiratory tract (Yeh et al. 1976) and aerosol samplers. For nonspherical or irregular particles, aerodynamic size is dependent on the particle shape and surface structures (Cox 1995) therefore one-dimensional size measurement poorly describes irregular or fibrous shaped particles. To date, little is known about the shape of various particle types reported as fungal aerosols except for the distribution of single spores and aggregates. Moreover, detailed characteristics of the fungal aerosol profile are unknown.

The morphology and size of spores, and hyphal and spore fragments have been assessed by microscopy in attempts to describe the fungal aerosols (Pady and Kramer 1960; Pady and Gregory 1963; Eduard et al. 1988, 1990; Heikkila et al. 1988; Karlsson and Malmberg 1989; Green et al. 2005; Halstensen et al. 2007; Afanou et al. 2014; Vestlund et al. 2014). Quantification and size-characterization of fungal particles have also been performed by measuring the optical or aerodynamic diameter of aerosol particles using automatic particle counters or sizers. For this purpose, a fairly narrow size range has been used to discriminate spores from fragments in aerosols generated from fungal cultures (Kildesø et al. 2000, 2003; Górny et al. 2002; Sivasubramani et al. 2004; Kanaani et al. 2009; Górny and Ławniczek-Wałczyk 2012). This approach led to the discovery of submicronic fragments. Fungal biomarkers such as ergosterol (Rao et al. 2005; Lau et al. 2006; Menetrez et al. 2009), phospholipid fatty acids (Womiloju et al. 2003) and  $\beta$ -glucans (Rao et al. 2005; Reponen et al. 2007; Adhikari et al. 2009; Madsen et al. 2009; Singh et al. 2011; Frankel et al. 2013; Seo et al. 2009, 2014) have also been used to demonstrate the presence of fungal biomass in various sizes of fungal aerosols. Further, enzymes, (e.g., chitinases [Madsen et al. 2005, 2009; Madsen 2012]), antigens (Górny et al. 2002) and allergens (Menetrez et al. 2001) as well as mycotoxins (Sorenson et al. 1987; Brasel et al. 2005) have been used to demonstrate allergenicity and toxicity potential of various size fractions of the fungal aerosols. Actually, the cut-off characteristics of the size-selective samplers are decisive for the separation efficiency.

The toxicological and immunological properties of allergen-bearing fine particles have been found to be strongly correlated with the particle number and overall surface than the particle mass (Nygaard et al. 2004). Thus, it can be hypothesized that the number of fungal particles with aerodynamic diameter  $<2.5 \mu\text{m}$  may show stronger correlations with observed health outcomes, than with mass. Moreover, hyphae have been shown to elicit stronger allergic inflammation as compared to single spores that induced non-allergic inflammatory responses (Bozza et al. 2002; Hohl et al. 2005). Also, Branzk et al. showed that neutrophils with mechanisms of neutrophils extracellular traps (NETosis) and reactive oxidative species (ROS) were involved in clearance of fiber-like particles and large spore aggregates (Branzk et al. 2014). A detailed description of the fungal aerosol by simultaneous

assessment of physical size, shape, and particle numbers is only possible by microscopy (Cox 1995).

Following our previous publication on the enumeration and origin of submicronic fungal fragments by high-resolution microscopy (Afanou et al. 2014), we describe in this study the morphology of fragments and spore particles, and their distribution in *in vitro* generated fungal aerosols. The size, shape and origin characteristics of various fungal particle types are described, as well as the profile of the aerosols generated from pure fungal cultures. The effect of generators and airflows used on the whole-aerosol composition was also investigated.

## MATERIALS AND METHODS

### Fungal Culture

Isolates of *Aspergillus fumigatus* Fresenius 1863 (strain A1258 FGSC) obtained from the Fungal Genetics Stock Center (University of Missouri, Kansas City, KS), *Aspergillus versicolor* Tirobaschi 1908 (strain VI 03554), and *Penicillium chrysogenum* Thom 1910 (strain VI 04528) obtained from the Section of Mycology at the Norwegian Veterinary Institute (Oslo, Norway) were used. The characteristics of these fungal isolates and the procedure for inoculum preparation have been previously described elsewhere (Afanou et al. 2014). Briefly, spores from 2-week-old cultures grown on malt extract agar (MEA) (Samson et al. 2004) were harvested by gently scraping cultures submerged in 20 mL phosphate buffered saline (PBS) containing 0.1% Tween 20 using a sterile inoculation loop. The spore suspension was transferred into 50 mL centrifuge tubes and resuspended by vortexing for  $2 \times 30$  seconds followed by sonication for 3 min in an ultrasonic bath at a frequency of 35 kHz (Sonorex RK 510H, Bandalin Electric, Berlin, Germany). The suspension was filtered through a  $10 \mu\text{m}$  nylon mesh filter (Millipore, Tullagreen, Cork, Ireland), centrifuged at  $1500 \times g$  for 5 min (Sigma 4k15, Osterode, Germany) and the pellet re-suspended in 30 mL sterile milliQ water containing 10% glycerol. The spore concentration was determined by filtration of 0.1 mL of 10 fold diluted suspension through a 25 mm diameter polycarbonate filter with  $0.4 \mu\text{m}$  pore size (Millipore, Tullagreen Cork, Ireland) and subsequent enumeration of deposited spores by field emission scanning electron microscopy (FESEM). One milliliter (containing ca.  $10^8$  spores) of this spore suspension was used to inoculate gypsum board (GB) (Lafarge Gips Dekoform 6 AK, Oberursel, Germany) commonly used as indoor building material in Norwegian residential environments. The GB was cut into 80 mm diameter circular plates that were prepared following the procedure previously described (Afanou et al. 2014). Following 8 weeks of incubation at  $25 \pm 1^\circ\text{C}$  and  $45 \pm 5\%$  relative humidity (in the incubator), the GB plates were utilized for aerosolization experiments.

### Aerosolization and Collection of Fungal Aerosols

The fungal cultures were aerosolized with the Fungal Spore Source Strength Tester (FSSST) (Sivasubramani et al. 2004), and Stami Particle Generator (SPG) (Afanou et al. 2014). The FSSST was built in polyvinyl chloride with a square internal area of 144 cm<sup>2</sup>, and 112 orifices of 0.4 mm diameter that directed air jets perpendicularly toward the culture plates. The SPG was built in aluminum and included a rotating plate of 156 cm<sup>2</sup> as support for the culture plates. A 6 mm diameter tube outfitted with ten orifices of 1.2 mm diameter served to direct air jets perpendicularly toward the culture plate. Schematic illustration for both generators and the experimental setup are shown in Figure 1.

The aerosolization experiments were performed at airflow rates of 12 and 20 L min<sup>-1</sup>. At these airflows, the air velocities at the orifices were 18 and 29 m s<sup>-1</sup> in the SPG, and 14 and 23 m s<sup>-1</sup> in the FSSST, respectively. During aerosolization the GB plates were situated at 14 mm below the orifices. Fungal aerosols were liberated by air currents provided through the orifices. Each aerosolization experiment was run for 120 sec while rotating the plate supporting *A. versicolor* and *P. chrysogenum* culture at 0.5 rpm in the SPG. In

FSSST, the culture dishes were aerosolized without rotation. In the experiment with *A. fumigatus*, a modified approach was used to prevent bioaerosol overload on the sampling filter. Briefly, the *A. fumigatus* culture surface area was covered with a plate that contained a 1 cm diameter central hole. *A. fumigatus* particles were liberated using the same airflow rate as described above. However, the sampling interval was 60 s while rotating at 1 rpm in SPG. Released particles from the culture were transported by the air currents within each system onto a 0.4 μm pore sized polycarbonate filter membrane (Isopore, Millipore, Ireland) and contained in open 37 mm cassettes (SKC Inc., Eighty Four, PA, USA). All experiments were conducted in a Bio Safe Grade II laminar flow hood to prevent external particle contamination and to protect laboratory workers from fungal exposure. The generator chamber was cleaned with 70% ethanol and purged for 10 min with HEPA filtered air prior to each experiment. Each experimental treatment was run in triplicate. Sixteen blank experiments were performed using cleaned empty chamber for background adjustment. Blanks with GB plates were disregarded because the substrate material after fungal bio-deterioration is different from the substrate at start (Sánchez 2009). Four




FIG. 1. Schematic illustration of the aerosolization chambers: Stami particle generator (SPG), fungal spore source strength tester (FSSST) and the experimental set up. One aerosolization chamber is connected to the system set up. Flow at the inlet and outlet of the chamber is measured by flowmeter 1 and 2, respectively. HEPA filter 1 and 2 filter the air to and from the pump, respectively. Constant relative humidity is maintained by the dryer (in line tube with silica beads). P2021 ionizer reduces electrostatic charges. Reproduced with permission from American Society of Microbiology: Afanou et al. 2014; *Applied and Environmental Microbiology*, 80(22): 7122–7130.

blank experiments for each generator  $\times$  flow rate combination were run for 120 s.

### Sample Preparation and Analysis

Briefly, a segment (ca. 2.4 cm<sup>2</sup>) was cut from the collection filter, mounted on a 25 mm aluminum pin stub (Agar Scientific Ltd., Stansted Essex, UK) using double-sided carbon adhesive discs (Ted Pella Inc., Redding, CA, USA), and coated with platinum in a Balzers SDC 050 sputter coater (Balzers, Liechtenstein) as previously described by Afanou and colleges (Afanou et al. 2014). Samples were analyzed using FESEM (SU 6600 Hitachi, Ibaraki-ken, Japan) in the secondary electron imaging (SEI) mode. The microscope was operated at 15 keV acceleration voltage, 1.8 kV extraction voltage and a working distance of 10 mm. Particles were quantified using the counting criteria described by Eduard and Aalen (Eduard and Aalen 1988). The particles were considered to be homogeneously distributed on the filter because an electrically conducting filter holder had been used to reduce electrostatic charging of the filter holder (Eduard et al. 1990).

### Identification and Classification of the Fungal Particles

Spore particles were identified and classified by their morphological features while other indiscernible particles were

defined as fragment particles and classified by length (Figure 2). Spore particles were categorized into five classes according to spore units per aggregate: single spores, and aggregates of 2, 3, 4, and  $\geq 5$  spores. Fragment particles were grouped in four classes based on their length: 0.2–1  $\mu\text{m}$  fragments, 1–2  $\mu\text{m}$  fragments, 2–3.5  $\mu\text{m}$  fragments, and  $\geq 3.5$   $\mu\text{m}$  fragments. Four hundred particles or a maximum of 100 fields were counted at 3000–6000  $\times$  magnification depending on the particle density on the filter and type of particle. The lowest detectable number of particles was  $8 \times 10^3$  and  $2 \times 10^4$  particles/filter at 3000  $\times$  and 6000  $\times$  magnification, respectively. The background numbers of each particle type by cm<sup>2</sup> surface of the supporting culture plate were in the range: 0–200 for single spores, 0 for aggregates of 2 spores, 0–45 for aggregates of 3 spores, 0 for aggregates of 4 spores, 0–40 for aggregates of 5 spores or more, 160–4000 for submicronic fragments, 55–800 for 1–2  $\mu\text{m}$  fragments, 0 for 2–3.5  $\mu\text{m}$  fragments and 0–160 for  $\geq 3.5$   $\mu\text{m}$  fragments.

### Size and Shape Measurements

The largest width and length were additionally measured on 12 randomly selected spore particles (single spores and aggregates of 2–6 spores). Similar measurements were performed on each type of fragment particles (submicronic and larger fragments) counted in 200 randomly selected fields at 3000 $\times$




FIG. 2. Micrographs of various fungal aerosolized particles. Spore particles (a–f); (a) single spores, (b–g) aggregate of 2, 3, 4, 5, and  $>5$  spores from *A. versicolor*. Fragment particles (h–l); (h) submicronic fragments:  $<1$   $\mu\text{m}$ ; (h–l) large fragments; (i) 1–2  $\mu\text{m}$  fragments; (j) 2–3.5  $\mu\text{m}$  fragments, (k and l)  $>3.5$   $\mu\text{m}$  fragments. Scale bar: 10  $\mu\text{m}$  (a, f, k, and l); 30  $\mu\text{m}$  (g); 0.5  $\mu\text{m}$  (h); 5  $\mu\text{m}$  (i); 2  $\mu\text{m}$  (j).

magnification. Measurements were performed at 3000–40,000× magnification depending on particle type and size. Shape of each particle was described by their aspect ratio calculated as length to width ratio.

**Statistical Analysis**

Data from 12 experiments of each fungal species were available for statistical analysis. All counts were adjusted for blank prior to the estimation of the number of particles per cm<sup>2</sup> culture. Negative or zero counts after background adjustment, were arbitrarily assigned 0.1 in order to enable ratio calculations and log ratio transformations for statistical analysis. Primary results were reported as median numbers, 25th and 75th percentiles of particles of each type per exposed culture area. The median numbers of particle types were compared between the fungal species using the nonparametric Kruskal–Wallis (K–W) test for multiple categories followed by *post hoc* Wilcoxon-rank sum test (Mann–Whitney *U* tests) and Bonferroni adjusted *p*-values (significant *p*-values <0.017).

Proportions of each particle type were calculated as the ratio of particle type counted to the total number of particles and were also reported as medians and percentiles (25th and 75th). The effects of the generators and airflows on the particle composition of the fungal aerosols were statistically assessed by applying linear mixed model on centered log ratio (CLR) transformed particle ratios (Aitchison 2003; Pawlowsky-Glahn and Egozcue 2006). CLR transformation was performed with CoDaPack version 2.01.15 (Department of Computer Science and Applied Mathematics, University of Girona, Girona Spain) by dividing each count ratio by the geometric mean of count ratios for each treatment. This transformation has been described to remove the constant sum constraint on compositional data (Aitchison 2003; Pawlowsky-Glahn and Egozcue 2006) prior to analysis.

Schematically the models were:

Generator:  $CLR = constant + Type + Type \times Generator + experiments (random) + residuals$

Airflow rates:  $CLR = constant + Type + Type \times Flow + experiments (random) + residuals$  (CLR: centered log ratio; *Type* represent the nine types of particles; *Generator* represents the two generators used; *Flow* represents the two airflow rates used).

To detect significant changes in the fungal aerosol composition between generators or airflows, we tested the significance of the two interaction terms (*Type* × *Generator*) and (*Type* × *Flow*). This was done by likelihood ratio tests between models with and without the interaction terms. Here, the models without the interaction terms included the two variables of the interaction term as single fixed effects (Jaeger 2008; Fitzmaurice et al. 2011). Furthermore, particle fractions that significantly changed between generators or airflows were pinpointed by the sign and significance of the individual coefficients of these two interaction terms. Significant *p*-values (*p* < 0.05) were adjusted for multiple comparisons using the false discovery rate method (Benjamini and Hochberg 1995; Benjamini 2010). Statistical analysis was performed in Stata SE 13 (Statacorp LP, College Station, TX, USA).

**RESULTS**

**Size and Shape Characteristics of Spores and Fragment Particles**

All counted particles were classified according to their physical length measured by FESEM. The average sizes for spore particles were in the range 2.41–15.04 μm and 2.02–4.26 μm for lengths and widths, respectively (Table 1). The corresponding sizes for fragment particles were 0.52–12.80 μm and 0.33–5.10 μm (Table 2).

Table 1  
Size and aspect ratio of spore particles

	<i>A. fumigatus</i>						<i>A. versicolor</i>						<i>P. chrysogenum</i>					
	S1	S2	S3	S4	S5	S6	S1	S2	S3	S4	S5	S6	S1	S2	S3	S4	S5	S6
N	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12
<b>AM</b>																		
Length	2.68	4.77	6.42	6.83	8.55	12.26	2.97	5.87	8.00	10.61	12.51	15.04	2.41	4.32	6.77	8.88	11.45	12.28
Width	2.25	2.61	3.20	3.64	3.65	4.26	2.53	2.66	3.20	3.65	3.75	4.25	2.02	2.07	2.22	2.78	2.88	3.34
Aspect ratio	1.20	1.83	2.05	1.97	2.45	3.14	1.18	2.22	2.57	3.10	3.70	3.88	1.21	2.16	3.11	3.45	4.19	4.16
<b>SD</b>																		
Length	0.27	0.61	1.61	1.15	1.03	2.48	0.26	0.78	1.15	1.82	2.30	3.17	0.28	0.48	1.10	2.03	1.93	2.11
Width	0.30	0.19	0.46	0.85	0.94	1.08	0.19	0.21	0.50	0.97	1.09	1.10	0.35	0.38	0.31	0.62	0.74	1.13
Aspect ratio	0.15	0.23	0.59	0.56	0.58	1.26	0.20	0.34	0.56	0.94	1.44	1.63	0.20	0.46	0.63	1.33	1.04	1.63

AM: Arithmetic mean; SD: standard deviation; length and width in μm.  
S1: Single spores; aggregates of 2 (S2), 3 (S3), 4 (S4), 5 (S5) and 6 (S6) spores

Table 2  
Size and aspect ratio of fragment particles

	<i>A. fumigatus</i>				<i>A. versicolor</i>				<i>P. chrysogenum</i>			
	<1 $\mu\text{m}$	1–2 $\mu\text{m}$	2–3.5 $\mu\text{m}$	>3.5 $\mu\text{m}$	<1 $\mu\text{m}$	1–2 $\mu\text{m}$	2–3.5 $\mu\text{m}$	>3.5 $\mu\text{m}$	<1 $\mu\text{m}$	1–2 $\mu\text{m}$	2–3.5 $\mu\text{m}$	>3.5 $\mu\text{m}$
N	113	41	27	19	102	44	25	29	67	36	30	68
<b>AM</b>												
Length	0.53	1.38	2.59	11.20	0.52	1.40	2.49	12.80	0.63	1.44	2.59	10.6
Width	0.33	0.77	1.41	4.69	0.34	0.76	1.46	5.10	0.37	0.75	1.42	3.34
Aspect ratio	1.78	2.12	2.26	3.34	1.78	2.19	1.89	3.01	2.00	2.17	2.08	4.68
<b>SD</b>												
Length	0.20	0.27	0.44	10.50	0.21	0.28	0.35	11.77	0.20	0.30	0.43	9.50
Width	0.14	0.34	0.91	5.10	0.22	0.35	0.40	4.53	0.16	0.28	0.64	2.50
Aspect ratio	0.77	1.00	0.87	4.15	0.86	1.04	0.83	3.25	1.47	0.94	0.72	5.82

AM: Arithmetic mean; SD: standard deviation; length and width in  $\mu\text{m}$

<1  $\mu\text{m}$ : submicronic fragments (SF); 1–2  $\mu\text{m}$  fragments (LF1); 2–3.5  $\mu\text{m}$  fragments (LF2) > 3.5  $\mu\text{m}$  fragments (LF3)

Three shape groups were recognized from the aspect ratios (Table 3). Spherical or near-spherical particles with aspect ratio in the range 1:1–1.5:1 group specifically the single spores irrespective to species. The median proportions of this group were 53%, 7%, and 6% of total aerosolized particles from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum*, respectively. A second group of oblong shaped particles has aspect ratios in the range 1.5:1–3:1 and represents 45%, 68%, and 55% (median) of the aerosolized particles from *A. fumigatus*, *A. versicolor* and *P. chrysogenum*, respectively. Further, this group comprises 0.2–1  $\mu\text{m}$  fragments, 1–3.5  $\mu\text{m}$  fragments, and spore aggregates except the largest aggregates. The third group has elongate and fiber-like shape with aspect ratio >3:1. Larger fragments (>3.5  $\mu\text{m}$ ) and larger spore aggregates belong to this group which represents 1%, 21%, and 24% (median) of total particles aerosolized from *A. fumigatus*, *A. versicolor* and *P. chrysogenum*, respectively. The overall average lengths (widths) were in the ranges 2.41–2.97  $\mu\text{m}$  (2.02–2.53  $\mu\text{m}$ ) for near spherical particles, 0.52–8.55  $\mu\text{m}$  (0.33–3.65  $\mu\text{m}$ ) for oblong particles and 3.5–15.04  $\mu\text{m}$  (2.22–5.10  $\mu\text{m}$ ) for the elongate and fiber-like particles. The micrographs of various particle types observed are shown in Figure 2.

### Number of Spores and Fragment Particles

The total median numbers of aerosolized particles were  $4.8 \times 10^5$  per  $\text{cm}^{-2}$  culture of *A. fumigatus*,  $5.6 \times 10^3$  for *A. versicolor* and  $8.9 \times 10^3$  for *P. chrysogenum*. Median sum spore particle numbers were  $2.9 \times 10^5$ ,  $0.8 \times 10^3$ , and  $0.9 \times 10^3$  per  $\text{cm}^{-2}$  culture, while the corresponding numbers for fragment particles were  $1.9 \times 10^4$ ,  $4.2 \times 10^3$ , and  $8.6 \times 10^3$  for per  $\text{cm}^{-2}$  culture for *A. fumigatus*, *A. versicolor* and *P. chrysogenum*, respectively. The median number of spore particles released from *A. fumigatus* cultures differed significantly from *A. versicolor* and *P. chrysogenum* cultures (Table 4).

*A. fumigatus* cultures released over 300 times the median number from *A. versicolor* ( $p < 0.001$ ) and *P. chrysogenum* ( $p < 0.001$ ). The number of fragment particles did not differ significantly between the tested species ( $p = 0.2$ ).

### Fungal Aerosol Composition

The bioaerosols collected from *A. fumigatus* was predominantly composed of single spores (median 53% of total particles) while submicronic fragment fractions were the highest in the aerosols from *A. versicolor* (submicronic fragments, median 34%) and *P. chrysogenum* (submicronic fragments, median 31%). Within the spore particle fraction from *A. fumigatus* and *P. chrysogenum*, single spores were most common: medians 70% and 28%, respectively, whereas the aerosols from *A. versicolor* were mainly composed of larger spore aggregates (aggregates of  $\geq 5$  spores, median 32%) (Table 4). Moreover, the submicronic fragments were dominant in the fragment particle fraction from all three isolates: medians 62%, 70%, and 63% of total fragments from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum*, respectively.

The combined proportions of larger fragments (medians of 1–2  $\mu\text{m}$  fragments, 2–3.5  $\mu\text{m}$ , and  $\geq 3.5 \mu\text{m}$ ) amounted to median 3%, 9%, and 8% of all particles collected from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum*, respectively (Table 4). The median proportions of single spores were 53%, 7%, and 6% while the medians of all spore aggregates combined were 26%, 23%, and 10% of all particles from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum* cultures, respectively (Table 4).

### Effect of Generator Types on the Fungal Aerosol Composition

Significant changes in the aerosol composition related to the type of generator were demonstrated through CLR

Table 3  
Shape characteristics of fungal bioaerosols based on aspect ratio. AM arithmetic means

Range of aspect ratio (AM)	<i>A. fumigatus</i>			<i>A. versicolor</i>			<i>P. chrysogenum</i>		
	Spore particles	Fragment particles	Median (%)	Spore particles	Fragment particles	Median (%)	Spore particles	Fragment particles	Median (%)
1:1-1.5:1 (near spherical)	Single spores	—	53	Single spores	—	7	Single spores	—	6
1.5:1-3:1 (oblong)	Aggregates of 2, 3, 4, and 5 spores	<1 $\mu\text{m}$ , 1-2 $\mu\text{m}$ and 2-3.5 $\mu\text{m}$	45	Aggregates of 2 and 3 spores	<1 $\mu\text{m}$ , 1-2 $\mu\text{m}$ and 2-3.5 $\mu\text{m}$	68	Aggregates of 2 spores	<1 $\mu\text{m}$ , 1-2 $\mu\text{m}$ and 2-3.5 $\mu\text{m}$	55
>3:1 (fiber-like)	Aggregates of 6 spores	>3.5 $\mu\text{m}$	1	Aggregates of 4, 5, and 6 spores	>3.5 $\mu\text{m}$	21	Aggregates of 3, 4, 5, and 6 spores	>3.5 $\mu\text{m}$	24



Table 4  
Particle types aerosolized from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum* (all experiments combined)

Particle types	<i>A. fumigatus</i> (N = 12)		<i>A. versicolor</i> (N = 12)		<i>P. chrysogenum</i> (N = 12)		K–W test <i>p</i> -value
	Number of particles <sup>ab</sup>	% of total <sup>b</sup>	Number of particles <sup>ab</sup>	% of total <sup>b</sup>	Number of particles <sup>ab</sup>	% of total <sup>b</sup>	
<b>Spores</b>							
S1	170 (7.0–1400)	53 (26–62)	0.3 (0.02–1.4)	7 (0.4–14)	0.4 (0.06–2.4)	6 (1–10)	<b>0.001</b>
S2	47 (2.0–240)	12 (8–17)	0.2 (0.02–1.8)	7 (0.3–15)	0.06(0.06–0.40)	1(0.6–5)	<b>&lt;0.001</b>
S3	20 (0.4–107)	4 (2–5)	0.09 (0.02–0.7)	1 (0.3–6)	0.06 (0.06–0.3)	2 (0.7–4)	<b>0.001</b>
S4	11 (0.06–118)	3 (2–4)	0.02 (0.02–0.4)	1 (0.3–4)	0.06 (0.06–0.1)	1(0.6–3)	<b>0.005</b>
≥S5	26 (0.6–140)	5 (2–15)	0.3 (0.09–2.1)	9 (1–19)	0.06(0.06–0.6)	3(0.7–8)	<b>0.01</b>
<b>Sum</b>	<b>290 (9.0–1950)</b>	<b>88 (75–94)</b>	<b>0.9 (0.3–7.0)</b>	<b>39 (5–58)</b>	<b>1.0 (0.3–3.4)</b>	<b>16 (8–42)</b>	<b>&lt;0.001</b>
<b>Fragments</b>							
SF	13.0 (1.0–300)	5 (1–21)	2.0 (0.6–5.0)	34 (7–62)	8.0 (0.6–10)	31 (10–89)	0.2
LF1	2.0 (0.06–4.1)	1 (0.2–1)	0.06(0.06–0.5)	2 (1–8)	0.06 (0.06–0.6)	2 (1–8)	0.2
LF2	2.7 (0.06–4.1)	1(0.2–4)	0.06 (0.06–0.6)	4 (0.5–22)	0.06 (0.06–0.7)	2 (0.7–6)	0.07
LF3	2.0 (0.06–4.1)	1 (0.1–3)	0.06 (0.06–0.2)	3 (1–8)	0.06 (0.06–3.0)	4 (1–38)	0.2
<b>Sum</b>	<b>19.0 (1.8–350)</b>	<b>12 (6–25)</b>	<b>4.0 (1.0–6.0)</b>	<b>61(42–95)</b>	<b>9.0 (2.0–15.0)</b>	<b>84 (58–92)</b>	0.2
<b>TOTAL</b>	<b>480 (17–2659)</b>		<b>5.6 (2.4–14)</b>		<b>8.9 (3–16.5)</b>		

<sup>a</sup>Data represent  $10^3$  particles  $\text{cm}^{-2}$  of culture media. <sup>b</sup>Medians, 25th and 75th percentiles (in parentheses); K–W = Kruskal–Wallis test for the three fungal species. The *post-hoc* Wilcoxon rank sum test of differences between species for median numbers with *A. fumigatus* significantly higher than *A. versicolor* and *P. chrysogenum*. Significance level ( $p \leq 0.017$ ). S1: Single spores; Aggregates of 2 (S2), 3 (S3), 4 (S4) and  $\geq 5$  (S5) spores. SF: Submicronic fragments, LF1: 1–2  $\mu\text{m}$  fragments, LF2: 2–3.5  $\mu\text{m}$  fragments, and LF3:  $\geq 3.5$   $\mu\text{m}$  fragments.

regression models and likelihood ratio tests of the interaction terms '*Generator*  $\times$  *Type*' (Table S1). The test was significant for *A. fumigatus* cultures aerosolized at 12  $\text{L min}^{-1}$  ( $p = 0.005$ ), for the *A. versicolor* and *P. chrysogenum* cultures both aerosolized at 12 and 20  $\text{L min}^{-1}$  ( $p = <0.001$ –0.007). The fraction of submicronic fragments was mainly affected when changes in the profile were observed between SPG and FSSST at 12 and 20  $\text{L min}^{-1}$ . Further, the fraction of submicronic fragments was significantly higher in SPG ( $p = 0.001$ ) as compared to FSSST with *A. fumigatus* cultures aerosolized at 12  $\text{L min}^{-1}$  (Figure 3a) while this fraction was lower in SPG as compared to FSSST at airflow 12  $\text{L min}^{-1}$  for *A. versicolor* ( $p < 0.001$ ) (Figure 3c) and *P. chrysogenum* ( $p = 0.007$ ) (Figure 3e). In addition, the single spore fraction was significantly larger in SPG versus FSSST with *P. chrysogenum* cultures ( $p < 0.001$ ). At 20  $\text{L min}^{-1}$ , no significant difference was observed between SPG and FSSST in the bioaerosol composition from *A. fumigatus* cultures (Figure 3b). But with *A. versicolor* cultures, the 2–3.5  $\mu\text{m}$  fragments fraction was significantly lower in SPG compared to FSSST ( $p = 0.001$ ) (Figure 3d). Furthermore, the submicronic fragment fraction from *P. chrysogenum* cultures was significantly larger in SPG versus FSSST ( $p = 0.002$ ) (Figure 3f).

### Effect of Airflow Rates on the Fungal Aerosol Composition

Fungal cultures subjected to airflow rates of 12  $\text{L min}^{-1}$  and 20  $\text{L min}^{-1}$  showed significantly different aerosol profiles. The

likelihood ratio test of the interaction term '*Flow*  $\times$  *Type*' was significant with both SPG and FSSST for all tested fungal isolates ( $p = <0.001$ –0.003) (Table S2). For *A. versicolor*, the fraction of spore particles was the largest at 20  $\text{L min}^{-1}$  in both generators ( $p = 0.004$ –0.02) whereas the proportion of  $>1$   $\mu\text{m}$  fragment particles was highest at 12  $\text{L min}^{-1}$  ( $p = <0.001$ –0.03) in FSSST (Figures 4c and d). Similar results were observed for *A. fumigatus*, but only in the SPG; the fraction of aggregates of  $\geq 5$  spores increased with increasing airflow ( $p = 0.012$ ), while  $\geq 3.5$   $\mu\text{m}$  fragments fraction ( $p < 0.001$ ) and the submicronic fragments fraction decrease with increasing airflow ( $p = <0.001$ –0.002) (Figure 4a). In the FSSST, the difference between the single spore fractions at 12 and 20  $\text{L min}^{-1}$  was closed to significance ( $p = 0.006$ ) with the largest fraction obtained at 20  $\text{L min}^{-1}$  (Figure 4b). Fractions of single spore and aggregates of  $\geq 5$  spores of *P. chrysogenum* were higher at 20  $\text{L min}^{-1}$  than at 12  $\text{L min}^{-1}$  ( $p = 0.008$ ) in the FSSST but not in SPG (Figures 4e and f). The  $\geq 3.5$   $\mu\text{m}$  fragments fraction increased with increasing airflow in both generators ( $p = <0.001$ –0.013) (Figures 4e and f) as opposed to the findings for the *Aspergillus* isolates. The submicronic fragment fraction decreased with increasing airflow rate in FSSST ( $p < 0.001$ ), but not significantly in SPG ( $p = 0.07$ ).

### DISCUSSION

With FESEM, the aerosols generated from fungal cultures revealed diverse arrays of particles with different size, shape,




FIG. 3. Distribution of mean proportions of particle types aerosolized from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum* cultures; SPG (black columns) and FSSST (white columns) at  $12 \text{ L min}^{-1}$  (a, c, and e) and at  $20 \text{ L min}^{-1}$  (b, d, and f). S1: Single spores; aggregates of 2 (S2), 3 (S3), 4 (S4),  $\geq 5$  (S5) spores. SF: Submicronic fragments, LF1:  $1\text{--}2 \mu\text{m}$  fragments, LF2:  $2\text{--}3.5 \mu\text{m}$  fragments, and LF3:  $\geq 3.5 \mu\text{m}$  fragments. \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.001$  ( $t$ -test of interaction coefficients between generator and particle types in the centered log-ratio mixed effect regression model of the proportions of particles; all  $p$ -values were adjusted for multiple comparisons by the false discovery rate method). Error bars represent standard errors.

and origin. The composition varied substantially between species and aerosolization conditions. The aspect ratio revealed three major particle shapes: near spherical, oblong and fiber-like. All single spores were fitted in the group of near spherical particles which is in close agreement with morphology described for spores from these species (Samson et al. 2004).

The aerodynamic diameter of these particles could therefore be a good predictor of their behavior in the respiratory tract as suggested by Reponen (1995) assuming that surface ornamentation has negligible effects. Further, most submicronic fragments were oblong while most of spores aggregates and larger fragments were elongate and fiber-like in shape. It is therefore likely that motion and deposition models of these particles in the respiratory tract will have to consider their shape characteristics in addition to aerodynamic equivalent diameter as reviewed by Lacey (1991).

Based on the physical sizes of various particles (microscopic length and width), 3–9% (median) of total particle counts were large fragments in the same size range as single spores and spore aggregates. Obviously, the heterogeneity of size and shape of fragment particles will confound the recognition of spore particles by large fragments when automatic counters or sizers are used. For example, Gorny et al. (2002; Gorny and Ławniczek-Wałczyk 2012) used optical diameter

of  $1.6 \mu\text{m}$  (a size delimitation) to discriminate between small fragments and spores from *Aspergillus versicolor*, *Penicillium melinii*, *P. chrysogenum*, and *Cladosporium cladosporioides* culture (Gorny et al. 2002; Gorny and Ławniczek-Wałczyk 2012). Similarly, Madsen described the composition of aerosols from *Botrytis cinerea* by allocating specific aerodynamic diameter ranges to specific particles types such as fragments ( $<1.6 \mu\text{m}$ ), microconidia ( $1.8\text{--}3.3 \mu\text{m}$ ), and macroconidia ( $3.5\text{--}10.4 \mu\text{m}$ ) (Madsen 2012). In these studies, the spore or conidia fractions are likely to comprise both spore aggregates and large fragments.

Using aerosol size fractionation samplers, spores, and fragment fractions have been reported likewise as studies that used automatic particle counters. Further quantification of fungal biomarker in the specific fractions allows the application of this approach in field settings for quantification of fungal submicronic fraction. For example, the mass of  $\beta$ -glucans was estimated in different size fractions of air samples collected by cyclones from indoor environments with molds contamination. The fungal aerosol profile was thus described as submicronic fragments fraction ( $<1 \mu\text{m}$ ), a mixture fraction of spores and fragments ( $1.05\text{--}2.25 \mu\text{m}$ ) and a spore fraction ( $>2.25 \mu\text{m}$ ) (Reponen et al. 2007; Seo et al. 2009, 2014). Moreover, multi-stage impactors (Gorny et al. 2002; Cho et al. 2005) as




FIG. 4. Distribution of mean proportions of particle types aerosolized from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum* cultures. Airflow 12 L min<sup>-1</sup> (black columns) and 20 L min<sup>-1</sup> (white columns) with SPG (a, c, and e) and FSSST (b, d, and f). S1: Single spores; aggregates of 2 (S2), 3 (S3), 4 (S4),  $\geq 5$  (S5) spores. SF: Submicronic fragments, LF1: 1–2  $\mu\text{m}$  fragments, LF2: 2–3.5  $\mu\text{m}$  fragments and LF3:  $\geq 3.5$   $\mu\text{m}$  fragments. \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.001$  (*t*-test of interaction coefficients between airflow and particle types in the centered log-ratio mixed effect regression model of the proportions of particles; all *p*-values were adjusted for multiple comparisons by the false discovery rate method). Error bars represent standard errors.

well as filters with different pore size in series (Brasel et al. 2005) have been used to disclose the fungal aerosol composition. However, the problem of spore bounce associated with impactor and cyclone sampling (Lindsley et al. 2006; Madsen et al. 2009) as well as the cut-off characteristics of impactors and cyclones suggest possible contamination of the submicronic fragment fraction by larger particles.

In any case whether fungal aerosols are studied with automatic counter or size fractionation coupled to biomarkers, characterization of the aerosol composition revealed only a one-dimensional size description of spores and fragment particles. In contrary, two-dimensional size measurements revealing shapes and sizes characteristics of various constituents present in the fungal aerosol was achieved in the present study using field emission scanning electron microscopy. Such visualization of particles favors their classification whether as spore or fragments. The present microscopic approach provides therefore more detailed characteristics of the fungal aerosols as compared to other approaches.

The total numbers of spore particles aerosolized from *A. fumigatus* were significantly higher as compared to *A. versicolor* and *P. chrysogenum* ( $p < 0.001$ ), but no species difference was observed for fragment numbers ( $p = 0.2$ ). We found no significant differences between *A. versicolor* and *P. chrysogenum* either for spore or fragment particles. This is in contradiction to results reported by Górný and Ławniczek-Wałczyk

2012 who also conducted aerosolization experiments on these species and observed significantly higher numbers of spores and fragments from *A. versicolor* as compared to *P. chrysogenum*. We can only speculate about what causes the deviating results: differences in experimental parameters such as generator, airflow, culture age, and the variation between isolates from the same fungal species. Nevertheless, the dominance of the submicronic particle fraction in the aerosols from *A. versicolor* or *P. chrysogenum* is in agreement with previously reported studies based on automatic particle counters or sizers (Górný et al. 2002; Cho et al. 2005; Madsen et al. 2005; Seo et al. 2009; Górný and Ławniczek-Wałczyk 2012). Further, the level of spore propagules (single spores and spore aggregates) aerosolized from *A. versicolor* grown on gypsum board has been previously assessed using microscopic counts (Sivasubramani et al. 2004). They reported higher levels: about  $7 \times 10^3$  per cm<sup>-2</sup> as compared to  $0.9 \times 10^3$  per cm<sup>-2</sup> of total spore particles in our study. This may be due to longer aerosolization period (10 min versus 2 min) and longer incubation time (12 months versus 8 weeks) applied in that study.

The presence of large spore aggregates in the aerosols is of concern because they carry much more antigenic substances into the respiratory tract despite their smaller number in the generated aerosols. The long chained aggregates have fiber-like shape with aerodynamic properties that may increase their chance to reach the trachea-bronchial region. It has been

recently demonstrated that neutrophils sense the microbe size and selectively release neutrophils extracellular traps (NETs) in response to large particles including fungal hyphal fragments and spore aggregates, but not in response to single spores (Branzk et al. 2014). It is therefore likely that different response mechanisms with substantially different cellular pathology will be induced upon inhalation and deposition of fungal particles in the respiratory tract. Large spore aggregates are likely to induce NETosis, whereas single spores are phagocytized. This size-dependent response should be emphasized in future exposure-response studies, and it will be important to include particle size determination in the characterization of fungal aerosols. The present study provides important information of fungal aerosols in this respect. The composition of aerosols generated from *A. fumigatus* including submicronic fragments has, to our knowledge, not been previously reported. Single spore outnumbered other particles suggesting that there is higher probability to detect spores from this species if growing in indoor environments as compared to *A. versicolor* and *P. chrysogenum* with aerosols dominated by fragments.

The present study revealed also significant changes in the bioaerosol composition related to the generators and airflows used for all tested isolates. For example, with *A. fumigatus* cultures the bioaerosol profile changed significantly to relatively fewer single spores and more submicronic fragments in SPG compared to FSSST when the airflow was  $12 \text{ L min}^{-1}$ . Similarly the proportions of submicronic fragments and  $\geq 3.5 \mu\text{m}$  fragments increased at  $12 \text{ L min}^{-1}$  compared to  $20 \text{ L min}^{-1}$  when SPG was used. Gorny and co-authors (Górny and Ławniczek-Wałczyk 2012) reported that generation factors such as air jet velocity, direction and movements have significant effects on the number of small fragments ( $< 1.6 \mu\text{m}$ ) and spores released from fungal cultures of *A. versicolor* and *P. chrysogenum*. Different physical parameters such as the level of turbulence inside the generator, impaction of particles onto the substrate and mycelia, and electrostatic charges on the particles and generator surfaces may collectively influence the release, the transportation, and loss of particles during *in vitro* aerosolization. These factors may even differ between particle types. The particle types (spores, spore aggregates, or fragments) that are mostly affected are likely of great importance for how the final composition of the bioaerosol may look.

The fraction of submicronic fragments was mostly influenced by the generator and airflow used. The most striking change was observed with *P. chrysogenum* aerosolized in the FSSST where the submicronic fragment fraction was 93% at a flow rate of  $12 \text{ L min}^{-1}$  which was reduced to 6% at a flow rate of  $20 \text{ L min}^{-1}$ . Only the results for *A. versicolor* seemed fairly consistent, showing increased spore fractions and lower fragment fractions at higher flow rate in both generators. The other fungal isolates showed more diverse results. Thus, the composition of fungal aerosols in these *in vitro* experiments

was greatly influenced by generation conditions. As changes in the fungal profiles often were in opposite directions, extrapolation of our results to environmental conditions appears difficult. Direct measurements in the environment are therefore required in order to assess exposure to fungal particle types different from spores.

Although detailed characteristics of fungal particles were obtained with FESEM, this microscopic method has been criticized for introducing two biases. First, the imaging that is performed in a vacuum chamber may cause distortion, collapse or disintegration of spore aggregates (Vestlund et al. 2014). Nevertheless, the morphological characteristics and the size measurements found are quite consistent with the literature (Frisvad and Samson 2004). In SEM, naturally collapsed spores have similar size (length and width) as compared to whole spores (Beckett et al. 1984). Furthermore, spores will desiccate when the relative humidity is below 100%, and desiccated spores are likely to occur naturally in the environment (Beckett et al. 1984). Secondly, the analyzed filter area is a small fraction (0.1–1%) of the exposed filter area. Although the distribution of particles in filter samples collected in electrically conducting cassettes is fairly homogenous (Eduard and Aalen 1988) the enumeration at higher resolution (3000–6000 $\times$ ) may lead to very low numbers or zero counted particles leading to highly Poisson distributed particle counts. However, the high resolution microscopic method is the only approach that permits classification and size and shape characterization of a broader range of fungal particles that include submicronic fragments, large fragments and spore particles.

## CONCLUSIONS

The high-resolution microscopic characterization of fungal aerosols from *A. fumigatus*, *A. versicolor*, and *P. chrysogenum* revealed a complex profiles with different shape, origin, and size. Fungal spores were aerosolized as single spores, chained, or clustered spore aggregates with shape and size that favor their deposition in different regions in the respiratory systems. Released fragments were mainly submicronic fragments but also included larger fragments with sizes that partly overlap those of spore particles. The FESEM method used in the present study was satisfactory for assessing shape and size as well as the source of fungal particles aerosolized from pure cultures grown on gypsum board.

The composition of the experimentally generated fungal aerosols was significantly influenced by the generator or the airflow rate used. Changes in the profiles were too diverse to enable extrapolation to real conditions. The complexity and variability of the bioaerosol composition and the dominance of fragments in the fungal aerosol from *A. versicolor* and *P. chrysogenum* supports previous recommendations on detecting broader range of particles types during fungal exposure assessment studies.

## ACKNOWLEDGMENTS

The technical support provided by the Norwegian Veterinary Institute is gratefully acknowledged. The findings and the conclusions in this report are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

## FUNDING

The present study is a part of the project “Fungal particles in indoor air” that is financially supported by the Norwegian Research Council: Grant number NFR196130/H10.

## SUPPLEMENTAL MATERIAL

Supplemental data for this article can be accessed on the publisher’s website.

## REFERENCES

- Adhikari, A., Jung, J., Reponen, T., Lewis, J. S., DeGrasse, E. C., Grimsley, L. F., et al. (2009). Aerosolization of Fungi, (1→3)-β-D-Glucans, and Endotoxin from Flood-Affected Materials Collected in New Orleans Homes. *Environ. Res.*, 109(3):215–24. doi:10.1016/j.envres.2008.12.010
- Afanou, K. A., Straumfors, A., Skogstad, A., Nilsen, T., Synnes, O., Skaar, I., et al. (2014). Submicronic Fungal Bioaerosols: High-Resolution Microscopic Characterization and Quantification. *Appl. Environ. Microbiol.*, 80(22):7122–7130. doi:10.1128/AEM.01740-14
- Aitchison, J. (2003). A concise guide to compositional data analysis. CDA Workshop, Girona. Available at [http://www.leg.ufpr.br/lib/exe/fetch.php/pessoais:abtmartins:a concise guide to compositional data analysis.pdf](http://www.leg.ufpr.br/lib/exe/fetch.php/pessoais:abtmartins:a%20concise%20guide%20to%20compositional%20data%20analysis.pdf). Accessed 17 October 2014
- Beckett, A., Read, N. D., and Porter, R. (1984). Variations in Fungal Spore Dimensions in Relation to Preparatory Techniques for Light Microscopy and Scanning Electron Microscopy. *J. Microsc.*, 136(1):87–95. doi:10.1111/j.1365-2818.1984.tb02548.x
- Benjamini, Y. (2010). Discovering the False Discovery Rate. *J. R. Stat. Soc. Series B: Statistical Methodol.*, 72(4):405–416.
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach of Multiple Testing. *J. R. Stat. Soc.*, 57(1):289–300. doi:10.2307/2346101
- Bozza, S., Gaziano, R., Spreca, A., Bacci, A., Montagnoli, C., Francesco, P., et al. (2002). Dendritic Cells Transport Conidia and Hyphae of *Aspergillus fumigatus* from the Airways to the Draining Lymph Nodes and Initiate Disparate Th Responses to the Fungus. *J. Immunol. (Baltimore, Md. : 1950)*, 168(3):1362–1371.
- Branzk, N., Lubojemska, A., Hardison, S. E., Wang, Q., Guitierrez, M. G., Brown, G. D., and Papayannopoulos, V. (2014). Neutrophils Sense Microbe Size and Selectively Release Neutrophil Extracellular Traps in Response to Large Pathogens. *Nat. Immunol.*, 15(11):1017–1025.
- Brasel, T. L., Douglas, D. R., Wilson, S. C., and Straus, D. C. (2005). Detection of Airborne *Stachybotrys Chartarum* Macrocytic Trichothecene Mycotoxins on Particulates Smaller Than Conidia. *Appl. Environ. Microbiol.*, 71(1):114–122. doi:10.1128/AEM.71.1.114
- Cho, S.-H., Seo, S.-C., Schmechel, D., Grinshpun, S. A., and Reponen, T. (2005). Aerodynamic Characteristics and Respiratory Deposition of Fungal Fragments. *Atmospher. Environ.*, 39(30):5454–5465. doi:10.1016/j.atmosenv.2005.05.042
- Cox, C. S. (1995). Physical Aspect of Bioaerosols Particles. In *Bioaerosols handbook*, C. S. Cox, and C. M. Wathes, eds. CRS Lewis Publishers, Boca Raton, Florida, pp. 15–25.
- Eduard, W. (2009). Fungal Spores: A Critical Review of the Toxicological and Epidemiological Evidence as a Basis for Occupational Exposure Limit Setting. *Crit. Rev. Toxicol.*, 39(10):799–864. doi:10.3109/10408440903307333
- Eduard, W. and Aalen, O. (1988). The Effect of Aggregation on the Counting Precision of Mould Spores on Filters. *Ann. Occupat. Hyg.*, 32(4):471–479.
- Eduard, W., Laguionie, P., Karlsson, K., Palmgren, U., Str , G., and Blomquist, G. (1990). Evaluation of Methods for Enumerating Microorganisms in Filter Samples from Highly Contaminated Occupational Environments. *Amer. Ind. Hyg. Assoc. J.*, 51(8):427–36. doi:10.1080/15298669091369899
- Eduard, W., Sandven, P., Johansen, B. V., and Bruun, R. (1988). Identification and Quantification of Mould Spores by Scanning Electron Microscopy (SEM): Analysis of Filter Samples Collected in Norwegian Saw Mills. *Ann. Occupat. Hyg.*, (32):447–445.
- Fitzmaurice, G., Laird, N., and Ware, J. (2011). *Applied Longitudinal Analysis* (2nd ed.). New Jersey: John Wiley & Sons, Inc., Hoboken.
- Frankel, M., Hansen, E., and Madsen, A. M. (2013). Effect of Relative Humidity on the Aerosolization and Total Inflammatory Potential of Fungal Particles from Dust-Inoculated Gypsum Boards. *Indoor Air*, 1–13. doi:10.1111/ina.12055
- Frisvad, J. C. and Samson, R. A. (2004). Polyphasic Taxonomy of *Penicillium* Subgenus *Penicillium*: A Guide to Identification of Food and Air-Borne Terverticillate *Penicillia* and Their Mycotoxins. *Stud. Mycol.*, 49:1–173.
- Górny, R. L. and Ławniczek-Walczyk, A. (2012). Effect of Two Aerosolization Methods on the Release of Fungal Propagules from a Contaminated Agar Surface. *Ann. Agricult. Environ. Med.*, 19(2):279–84. <http://www.ncbi.nlm.nih.gov/pubmed/22742802>
- Górny, R. L., Reponen, T., Willeke, K., Robine, E., Boissier, M., Sergey, A., et al. (2002). Fungal Fragments as Indoor Air Biocontaminants. *Appl. Environ. Microbiol.*, 68(7):3522–3532. doi:10.1128/AEM.68.7.3522
- Green, B. J., Schmechel, D., Sercombe, J. K., and Tovey, E. R. (2005). Enumeration and Detection of Aerosolized *Aspergillus Fumigatus* and *Penicillium Chrysogenum* Conidia and Hyphae using a Novel Double Immunostaining Technique. *J. Immunol. Methods*, 307(1–2):127–34. doi:10.1016/j.jim.2005.10.001
- Halstensen, A. S., Nordby, K., Wouters, I., and Eduard, W. (2007). Determinants of Microbial Exposure in Grain Farming. *Ann. Occupat. Hyg.*, 51(7):581–92. doi:10.1093/annhyg/mem038
- Heikkilä, P., Salmi, T., and Kotimaa, M. (1988). Identification and counting of fungal spores by scanning electron microscopy. *Scand. J. Work, Environ. Health*, 14(Supplement 1):66–67.
- Hohl, T. M., Van Epps, H. L., Rivera, A., Morgan, L. A., Chen, P. L., Feldmesser, M., and Pamer, E. G. (2005). *Aspergillus Fumigatus* Triggers Inflammatory Responses by Stage-Specific Beta-Glucan Display. *PLoS Pathogens*, 1(3):0232–0240. doi:10.1371/journal.ppat.0010030
- Institute of Medicine. (2004). Human Health Effects. In *Damp Indoor Spaces and Health*. Washington, DC: National Academies Press, pp. 189–243.
- Jaeger, T. F. (2008). Categorical Data Analysis: Away from ANOVAs (transformation or not) and Towards Logit Mixed Models. *J. Memory Language*, 59(4):434–446. doi:10.1016/j.jml.2007.11.007
- Kanaani, H., Hargreaves, M., Ristovski, Z., and Morawska, L. (2009). Fungal Spore Fragmentation as a Function of Airflow Rates and Fungal Generation Methods. *Atmos. Environ.*, 43(24):3725–3735. doi:10.1016/j.atmosenv.2009.04.043
- Karlsson, K. and Malmberg, P. (1989). Characterization of Exposure to Molds and Actinomycetes in Agricultural Dusts by Scanning Electron Microscopy, Fluorescence Microscopy and the Culture Method. *Scand. J. Work, Environ. Health*, 15(5):353–359. doi:10.5271/sjweh.1847

- Kildesø, J., Wurtz, H., Nielsen, K. F., Wilkins, C., Gravesen, S., Thrane, U., et al. (2000). The Release of Fungal Spores from Water Damaged Building Materials. In Proceedings of *Exposure, Human Responses and Building Investigations*, Vol. 1. O. Seppanen and J. Sateri, Eds., Helsinki, Finland, pp. 313–318.
- Kildesø, J., Wütz, H., Nielsen, K. F., Kruse, P., Wilkins, K., Thrane, U., et al. (2003). Determination of Fungal Spore Release from Wet Building Materials. *Indoor Air*, 13:148–155.
- Lacey, J. (1991). Aggregation of Spores and its Effect on Aerodynamic Behaviour. *Grana*, 30(2):437–445. doi:10.1080/00173139109432005
- Lau, A. P. S., Lee, A. K. Y., Chan, C. K., and Fang, M. (2006). Ergosterol as a Biomarker for the Quantification of the Fungal Biomass in Atmospheric Aerosols. *Atmos. Environ.*, 40(2):249–259. doi:10.1016/j.atmosenv.2005.09.048
- Lindsley, W. G., Schmechel, D., and Chen, B. T. (2006). A Two-Stage Cyclone using Microcentrifuge Tubes for Personal Bioaerosol Sampling. *J. Environ. Monit.: JEM*, 8(11):1136–42. doi:10.1039/b609083d
- Madsen, A. M. (2012). Effects of Airflow and Changing Humidity on the Aerosolization of Respirable Fungal Fragments and Conidia of *Botrytis Cinerea*. *Appl. Environ. Microbiol.*, 78(11):3999–4007. doi:10.1128/AEM.07879-11
- Madsen, A. M., Schlünssen, V., Olsen, T., Sigsgaard, T., and Avci, H. (2009). Airborne Fungal and Bacterial Components in PMI Dust from Biofuel Plants. *Ann. Occupat. Hyg.*, 53(7):749–57. doi:10.1093/annhyg/mep045
- Madsen, A. M., Wilkins, K., and Poulsen, O. (2005). Micro-Particles from Fungi. In *Bioaerosols, Fungi, Bacteria, Mycotoxins and Human Health*, E. Johanning, ed., Fungal Research Group Foundation, Inc., Albany, New York, USA, pp. 276–281.
- Menetrez, M. Y., Foarde, K. K., and Ensor, D. S. (2001). An Analytical Method for the Measurement of Nonviable Bioaerosols. *J. Air Waste Manag. Assoc.* (1995), 51:1436–1442. doi:10.1080/10473289.2001.10464365
- Menetrez, M. Y., Foarde, K. K., Esch, R. K., Schwartz, T. D., Dean, T. R., Hays, M. D., et al. (2009). An Evaluation of Indoor and Outdoor Biological Particulate Matter. *Atmos. Environ.*, 43(34):5476–5483. doi:10.1016/j.atmosenv.2009.07.027
- Nygaard, U. C., Samuelsen, M., Aase, A., and Løik, M. (2004). The Capacity of Particles to Increase Allergic Sensitization is Predicted by Particle Number and Surface Area, Not by Particle Mass. *Toxicol. Sci.: Off. J. Soc. Toxicol.*, 82(2):515–24. doi:10.1093/toxsci/kfh287
- Pady, S. and Gregory, P. (1963). Numbers and Viability of Airborne Hyphal Fragments in England. *Trans. Brit. Mycol. Society*, 46(4):609–613.
- Pady, S. and Kramer, C. (1960). Kansas Aeromycology VI: Hyphal Fragments. *Mycologia*, 52:681–687.
- Pawlowsky-Glahn, V. and Egozcue, J. J. (2006). Compositional Data and Their Analysis: An Introduction. *Geolog. Soc., London, Special Publicat.*, 264(1):1–10. doi:10.1144/GSL.SP.2006.264.01.01
- Rao, C. Y., Cox-Ganser, J., Chew, G. L., Doekes, G., and White, S. (2005). Use of Surrogate Markers of Biological Agents in Air and Settled Dust Samples to Evaluate a Water Damaged Hospital. *Indoor Air*, 15(Suppl 9):89–97.
- Reponen, T. (1995). Aerodynamic Diameters and Respiratory Deposition Estimates of Viable Fungal Particles in Mold Problem Dwellings. *Aerosol Sci. Technol.*, 22(1):11–23. doi:10.1080/02786829408959724
- Reponen, T., Grinshpun, S. A., Conwell, K. L., Wiest, J., and Anderson, M. (2001). Aerodynamic Versus Physical Size of Spores: Measurement and Implication for Respiratory Deposition. *Grana*, 40(February 2015):119–125. doi:10.1080/00173130152625851
- Reponen, T., Seo, S.-C., Grimsley, L. F., Lee, T., Crawford, C., and Grinshpun, S. A. (2007). Fungal Fragments in Moldy Houses: A Field Study in Homes in New Orleans and Southern Ohio. *Atmos. Environ. (Oxford, England: 1994)*, 41(37):8140–8149. doi:10.1016/j.atmosenv.2007.06.027
- Samson, R. A., Frisvad, J. C., and Hoekstra, E. (2004). *Introduction to Food and Airborne Fungi*. Utrecht: Centraalbureau voor Schimmelcultures.
- Sánchez, C. (2009). Lignocellulosic Residues: Biodegradation and Bioconversion by Fungi. *Biotechnol. Adv.*, 27(2):185–94. doi:10.1016/j.biotechadv.2008.11.001
- Seo, S.-C., Choung, J. T., Cehn, B. T., Lindsley, W. G., and Kim, K. Y. (2014). The Level of Submicron Fungal Fragments in Homes with Asthmatic Children. *Environ. Res.*, 131C:71–76. doi:10.1016/j.envres.2014.02.015
- Seo, S.-C., Reponen, T., Levin, L., and Grinshpun, S. A. (2009). Size-Fractionated (1→3)- $\beta$ -D-Glucans Concentrations Aerosolized from Different Moldy Building Materials. *Sci. Total Environ.*, 407(2):806–14. doi:10.1016/j.scitotenv.2008.10.018
- Singh, U., Levin, L., Grinshpun, S. A., Schaffer, C., Adhikari, A., and Reponen, T. (2011). Influence of Home Characteristics on Airborne and Dust-borne Endotoxin and  $\beta$ -D-Glucan. *J. Environ. Monit.: JEM*, 13(11):3246–53. doi:10.1039/c1em10446b
- Sivasubramani, S. K., Niemeier, R. T., Reponen, T., and Grinshpun, S. A. (2004). Fungal Spore Source Strength Tester: Laboratory Evaluation of a New Concept. *Sci. Total Environ.*, 329(1–3):75–86. doi:10.1016/j.scitotenv.2004.03.007
- Sorenson, W., Frazer, D. G., Jarvis, B. B., Simpson, J., and Robinson, V. A. (1987). Trichothecene Mycotoxins in Aerosolized Conidia of *Stachybotrys Atra*. *Appl. Environ. Microbiol.*, 53(6):1370–1375.
- Vestlund, T., Al-Ashaab, R., Tyrrel, S. F., Longhurst, P. J., Pollard, S. J. T., and Drew, G. H. (2014). Morphological Classification of Bioaerosols from Composting using Scanning Electron Microscopy. *Waste Manag. (New York, N.Y.)*, 34(22):3833–3842. doi:10.1016/j.wasman.2014.01.021
- Womiloju, T. O., Miller, J. D., Mayer, P. M., and Brook, J. R. (2003). Methods to Determine the Biological Composition of Particulate Matter Collected from Outdoor Air. *Atmos. Environ.*, 37(31):4335–4344. doi:10.1016/S1352-2310(03)00577-6
- World Health Organization. (2009). *Guidelines for Indoor Air Quality: Dampness and Mould*. WHO Regional Office for Europe, Copenhagen O, Denmark. doi:ISBN 978 92 890 4168 3
- Yeh, H. C., Phalen, R. F., and Raabe, O. G. (1976). Factors Influencing the Deposition of Inhaled Particles. *Environ. Health Perspect.*, 15:147–156