



Detection of immunoglobulins containing plasma cells in the thymus, bursa of Fabricius and spleen of vaccinated broiler chickens with Newcastle disease virus vaccine

Md. Abdul Masum, Mohammed Zahirul Islam Khan, Morsheda Nasrin, M. Nazmul Hassan Siddiqi, Mohammed Zubayer Ibna Khan & Md. Nabiul Islam

To cite this article: Md. Abdul Masum, Mohammed Zahirul Islam Khan, Morsheda Nasrin, M. Nazmul Hassan Siddiqi, Mohammed Zubayer Ibna Khan & Md. Nabiul Islam (2014) Detection of immunoglobulins containing plasma cells in the thymus, bursa of Fabricius and spleen of vaccinated broiler chickens with Newcastle disease virus vaccine, International Journal of Veterinary Science and Medicine, 2:2, 103-108, DOI: [10.1016/j.ijvsm.2014.06.001](https://doi.org/10.1016/j.ijvsm.2014.06.001)

To link to this article: <https://doi.org/10.1016/j.ijvsm.2014.06.001>



© Faculty of Veterinary Medicine, Cairo University



Published online: 03 May 2019.



Submit your article to this journal [↗](#)



Article views: 175



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)



Full Length Article

Detection of immunoglobulins containing plasma cells in the thymus, bursa of Fabricius and spleen of vaccinated broiler chickens with Newcastle disease virus vaccine



Md. Abdul Masum ^a, Mohammed Zahirul Islam Khan ^{a,b,*}, Morsheda Nasrin ^a,
M. Nazmul Hassan Siddiqi ^a, Mohammed Zubayer Ibna Khan ^c, Md. Nabiul Islam ^d

^a Department of Anatomy and Histology, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh

^b Faculty of Veterinary Medicine, University Malaysia Kelantan, Locked Bag 36, PengkalanChepa, 16100 Kota Bharu, Kelantan, Malaysia

^c Dhaka Medical College, Dhaka, Bangladesh

^d Division of Neuroanatomy, Yamaguchi University Graduate School of Medicine, Ube, Yamaguchi 755-8505, Japan

Received 10 March 2014; revised 28 May 2014; accepted 2 June 2014

Available online 2 December 2014

KEYWORDS

Igs cells;
Lymphoid tissues;
Newcastle disease vaccine;
Immunohistochemistry;
Broilers

Abstract Mobilization of immunoglobulins (Igs)-containing plasma cells (IgA, IgG and IgM) in the spleen, bursa of Fabricius and thymus was investigated in broiler chickens that were vaccinated with Newcastle disease virus (NDV) vaccine. In the thymus, the Igs-containing plasma cells were distributed in the cortex and medulla. Their frequency and distribution were higher at D₁₄ and at D₂₈. The number of IgG- and IgM-positive cells was greater than IgA-positive cells in thymus. In the bursa of Fabricius, Igs-containing plasma cells were distributed beneath the capsules; within and around the bursal follicles. Their frequency of occurrence significantly peaked at D₁₄ and at D₂₈ in comparison to day-old chickens, and IgG-positive cells were significantly greater than the IgA- and IgM-positive cells in the bursa of vaccinated chickens. In the spleen, Igs-containing plasma cells were distributed in the white pulp, around the trabeculae, and in the periarterial lymphatic sheath. In this secondary lymphatic tissue, IgG- and IgM-positive cell numbers significantly greater

* Corresponding author at: Faculty of Veterinary Medicine, University Malaysia Kelantan, Padang Tembak, 16100 Kota Bharu, Kelantan, Malaysia. Tel.: +60 13 652 7135; fax: +60 9 771 7132.
E-mail address: zahirul@umk.edu.my (M.Z.I. Khan).

Peer review under responsibility of Faculty of Veterinary Medicine, Cairo University.

<http://dx.doi.org/10.1016/j.ijvsm.2014.06.001>

2314-4599 © 2014 Production and hosting by Elsevier B.V. on behalf of Faculty of Veterinary Medicine, Cairo University.

than IgA-positive cells. In conclusion, mobilization of more Igs-positive cells in lymphoid tissues of broiler chickens is due to the effect of NDV vaccine as well as the advancement of age.

© 2014 Production and hosting by Elsevier B.V. on behalf of Faculty of Veterinary Medicine, Cairo University.

1. Introduction

Poultry production, especially chickens and ducks has attained an important place in the agricultural economy of Bangladesh both through contribution to gross domestic product (GDP) and employment, especially in urban and pre-urban areas. Newcastle disease (ND) is the most common and harmful disease for commercial broilers. In addition, ND causes serious outbreaks in commercial layer and parent stocks every year and results in huge losses for the poultry industry. ND leads to necrosis, infiltration of heterophils and heterophilic changes in lymphoid organs [1].

Most living beings manage not only to survive but indeed thrive in a potentially hostile milieu, with seemingly little effort. This freedom from disease is dependent on the existence of a complex and highly sophisticated defense system called lymphoid system [2] which start to develop at embryonic day (ED) 10 in broilers [3,4]. The lymphoid system of fowl is consisting of unique organs and divided into two morphologically and functionally distinct components [5]. The lymphoid tissues have an independent phylogenetic origin, their function being to react to foreign antigens by producing antibodies, thereby providing "adaptive immunity". An obvious characteristic of the lymphatic tissues of mammals and birds is that they are densely populated with lymphocytes. This is because they are involved in the lymphocytes production, immune responses or both of these occurring at the same time [6]. When lymphocytes arrive in the lymphoid tissues or organs they become plasma cells and begin to synthesize immunoglobulins. The plasma cells-containing different classes of immunoglobulins (Igs) are distributed throughout the lymphoid tissues, [7] including Harderian gland of broiler chickens [8].

Chickens have been used as experimental animals for studies of the immune system, because chicken T and B cells mature in the thymus and bursa of Fabricius, respectively, and because these organs are easily manipulated. The development, differentiation, distribution and function of immune cells in these tissues, including the spleen of chicken, have been investigated using a wide variety of methodologies: surgical thymectomy, bursectomy, X-ray irradiation, and the colloidal carbon uptake method, as well as the administration of hormones, drugs, vitamins and minerals [9–14]. In the present study, we used NDV vaccine via both ocular and oral routes to study the frequency and distribution of Igs-containing plasma cells in the thymus, bursa of Fabricius and spleen of broiler chickens.

2. Materials and methods

The experiment was carried out in the Department of Anatomy and Histology, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh, Bangladesh. Part of the research was also conducted in the Laboratory of Anatomy and Histology, Faculty of Veterinary Medicine, University

Malaysia Kelantan, Malaysia. The following materials and methodologies were followed in the present experiment:

2.1. Chickens

A total of 32 day-old "Cob-500" broiler chickens of both sexes were purchased from Kazi Farm Ltd. Mymensingh, Bangladesh. The chickens had no visible developmental disorders or detectable diseases that might influence the distribution of lymphocytes and plasma cells.

2.2. Management

The chickens were reared in a litter system in a poultry shed at the Department of Poultry Science, Faculty of Animal Husbandry, Bangladesh Agricultural University, Mymensingh. All the chickens were provided with broiler feed and mineral water *ad libitum*. Poultry shed bio-security and brooding management were both strictly maintained, as were optimum temperature, lighting and ventilation in the brooder.

2.3. Vaccines

The live attenuated NDV vaccine used in this study was the product of Mohakhali Research Centre, Directorate of Live-stock Services, Dhaka, Bangladesh. The recommended dose is one drop of vaccine per eye in between one to 3-days-old chicken, with a booster dose usually given between 12 and 21 days of age.

2.4. Experimental design

The chickens were grouped into control and treatment groups. Chickens in the control group ($n = 16$) were given a drop of distilled water ocularly at 06:00 on D_1 . From this group 8 chickens were killed at 18:00 on the same day (D_1), and the remaining 8 chickens were sacrificed on D_{10} at 18:00 to ascertain the mobilization of immune cells. Chickens in the treatment group ($n = 16$) were given NDV vaccine ocularly on D_3 at 06:00 with a booster dose of the same vaccine being administered orally on D_{13} at 06:00. The thymus, bursa of Fabricius and spleen were collected from these chickens on D_{14} and D_{28} after sacrificing the chickens (8 broilers per group) through cervical subluxation, and specimens were fixed in Bouin's fluid or ice cold periodate lysine paraformaldehyde (PLP) for histological and immunohistochemical study.

2.5. Antibodies

The following antibodies were used for detecting Igs-positive cells: normal rabbit serum (Biosource international, Inc., Camarillo, CA, USA), goat anti-chicken IgA (Bethyl Laboratories, Montgomery, TX, USA), goat anti-chicken

IgG (Bethyl Laboratories), goat anti-chicken IgM (Bethyl Laboratories) and HRP-conjugated rabbit anti-goat IgG (Bethyl Laboratories).

2.6. Immunohistochemistry

The indirect immunoperoxidase method was performed to study the distribution pattern and frequencies of Igs-positive cells in the bursa of Fabricius, thymus, and spleen, of control and vaccinated broiler chickens. The tissues from bursa of Fabricius, thymus and spleen were fixed in Bouin's fluid or PLP, dehydrated in a series of graded alcohol, cleared in xylene, and embedded in paraffin. Paraffin sections of 6 μm thickness were immunostained by the indirect immunoperoxidase method as described previously [14].

2.7. Histoplanimetry

The Igs-containing plasma cells were counted in 20 fields of the histological sections using a light microscope at a magnification of 400 \times where Igs-plasma cells were diffusely distributed and their relative frequency per focus was calculated according to the point count method [16].

2.8. Statistical analysis

The numbers of Ig-containing plasma cells and their variation in numbers in the thymus, bursa of Fabricius and spleen were compared (randomized block design) among broilers of various stages, and data were evaluated by student's *t*-test [17].

3. Results and discussion

Frequency and distribution of Igs-containing plasma cells (IgA, IgG and IgM) in the lymphoid tissues of broiler chickens:

3.1. Thymus

In the thymus, Igs-containing plasma cells were distributed in the cortex and medulla (Fig. 1a–c). The frequency of Igs-containing plasma cells was statistically greater ($P < 0.05$) in the thymus at D₁₄ and D₂₈ in comparison to the day-old chickens (Figs. 4–6). When comparing frequency of occurrence of different Igs-containing plasma cells, it was found that IgG- and IgM-positive cells were greater ($P < 0.05$) at D₁₄ and D₂₈ ages. In the present study, the population of Igs-containing plasma cells and their infiltration to the lymphoid organs due to inoculations of NDV vaccine were clearly observed. These dynamic changes in the frequency and distribution of Igs-containing plasma cells were also observed in the Harderian gland, cecal tonsil and trachea of our previous studies [14]. The thymus is responsible for the production of T-lymphocytes subsets in the chicken [18] not Igs-containing plasma cells, however existence of statistically higher IgG- and IgM-positive cells in the thymus of the present study indicates that the B lymphocytes infiltrated in the thymus via blood circulation and differentiated into Igs-containing plasma cell for local defensive function as T-lymphocytes subpopulation do in the bursa of Fabricius [19], which is the homing organ for B lymphocytes.




Figure 1 (a–c) Immunostained section of chicken thymus of broiler chickens at D₁(a), D₁₄(b), and D₂₈(c) showing IgM-positive cells (arrows) in cortex (C) and medulla (M). IgM-positive cells are numerous at D₁₄ and at D₂₈. Scale bar = 50 μm .

3.2. Bursa of Fabricius

In the bursa of Fabricius, Igs-containing plasma cells were found principally beneath the capsule; in the cortex, and medulla of the bursal follicles; and around the bursal follicles (Fig. 2a and b). The frequency of Igs-containing plasma cells in this gland was greater at D₁₄ and at D₂₈ in comparison to day-old chickens (Figs. 4–6). When comparing the data, it was found that, the frequency of IgG-positive cells was statistically higher ($P < 0.01$) at D₂₈. In contrast, IgM-positive cells




Figure 2 (a and b) Immunostained sections of bursa of Fabricius of broiler chickens at D₁(a) and D₁₄(b) showing IgA-positive cells (a) and IgG-positive-cells (b) (arrows) in the cortex (C), medulla (M), and beneath the capsule (CP) of the bursal follicle. The immunopositive cells are more numerous at D₁₄ than day-old-chickens. Scale bar = 50 μm.

were insignificantly greater than IgA- and IgG-positive cells at D₁ and at D₁₄. The findings of our present study varied greatly [20]. Honjo et al. [15] reported that most of the bursal lymphoid cells of 4 week old inbred line P chickens were IgM positive. This variation at the same age is possibly due to the strain differences of chickens used in the present study. In the vaccinated chickens of the present study Igs-containing plasma cells were sharply increased from the D₁ chickens and peaked from D₁₄ to D₂₈ and this might be due to immunomodulation of the vaccine.

3.3. Spleen

In the spleen, Igs-containing plasma cells were numerous in the white pulp (Fig. 3a–c), around the trabeculae, around the central artery and in the periarterial lymphatic sheath. These distribution patterns of Igs-containing plasma cells in the vaccinated chickens were similar to our previous study [13], where we used vitamins and minerals to observe the distribution pattern of Igs-containing plasma cells. A similar distribution pattern of Igs-containing plasma cells was also observed in the spleen of the control group in the present study. The frequency and distribution of IgG-positive cells were




Figure 3 (a–c) Immunostained sections of spleens of chickens at D₁(a), D₁₄(b) and D₂₈(c) showing IgA-positive cells (a), IgG-positive cells (b) and IgM-positive cells (c) (arrows) in the white pulp of the spleen. In the spleen, immunopositive cells are more numerous at D₁₄ and D₂₈. Scale bar = 50 μm.

statistically greater ($P < 0.01$) at D₁₄ (Fig. 5); and IgA- and IgM-positive cells were higher at D₂₈ (Figs. 4 and 6). When comparing the data of Igs-containing plasma cells in this gland, it was observed that, both IgG- and IgM-positive cells were statistically higher ($P < 0.01–0.05$) than the IgA-positive cells throughout the period of postnatal development. This finding was in agreement with our previous statement that the NDV vaccine initiates mobilization of similar types of immune cells in the Harderian gland of broiler chickens [14].

Chickens were inoculated with NDV vaccine at D₃, and the booster dose was given at D₁₃ in the present study. We compared the data of Igs-containing plasma cells between vaccinated chickens and control up to D₁₀, and it was found




Figure 4 The frequency of IgA positive cells in the thymus, bursa of Fabricius and spleen of broiler chickens from D₁ to D₂₈. IgA-positive cells are greater at D₁₄ and D₂₈ chickens than in D₁. The frequency of these immune cells is significantly greater ($P < 0.01$) in the spleen, followed by the thymus and bursa of Fabricius at D₂₈ of development. The values are given as the mean \pm standard error ($n = 8$).




Figure 5 The frequency of IgG-positive cells in the thymus, bursa of Fabricius and spleen of chickens during their postnatal growth from D₁ to D₂₈. IgG-positive cells are more frequent in D₁₄ and D₂₈ chickens than in comparison to D₁ chickens. The frequency of these cells are significantly greater ($P < 0.01$) in the spleen, followed by the thymus and bursa of Fabricius at D₁₄ in chickens, whereas at D₂₈, IgG cells are found more numerous in the bursa of Fabricius. The values are given as the mean \pm standard error ($n = 8$).

that the number of Igs-containing plasma cells were significantly greater in vaccinated chickens than in control. In the present study, it was observed that the frequency and distribution of IgA-, IgG- and IgM-positive cells were more in D₁₄ and D₂₈ groups of vaccinated chickens than the chickens of D₁. This finding was similar to the report of Salam et al. [20] and Solcan et al. [21]. Among the Igs-containing plasma cells, the number of IgG- and IgM-positive cells was significantly greater at D₁₄ and D₂₈ groups of chickens. This finding is in partial agreement with the results of Bienenstock et al. [22] and Ohshima et al. [23], who reported that both IgA- and IgG-positive cells formed the bulk of the lymphoid cell




Figure 6 Frequency of IgM-positive cells in the thymus, bursa of Fabricius and spleen of chickens during their postnatal growth from D₁ to D₂₈. The frequency of IgM-positive cells is greater at D₁₄ and D₂₈, and in the spleen these cells are significantly greater ($P < 0.01-0.05$) at D₁ and D₂₈ followed by the thymus and bursa of Fabricius. However, at D₁₄ these immune cells are more numerous in the bursa of Fabricius, but the difference is not significant. The values are given as the mean \pm standard error ($n = 8$).

population in the early stages of life and that IgA-positive cells were predominant in the later stages of life.

IgG and IgM cells protect the body from infection are found in the lymphoid organs, whereas IgA cells are found in the secretory organs [24]. The frequency of these cells and the antibody titre in the serum increased with the NDV vaccine alone [25] or perform synergistic effect with immune adjuvant [26]. In the present study, the number of IgG- and IgM-positive cells was statistically higher than IgA-positive cells in the thymus and bursa of Fabricius. In contrast to the Harderian gland of the duck [27] and chicken [8]; and to the mucosa of the gastrointestinal tract of chickens [7], IgA-positive cells were significantly greater. This observation from avian species was similar to the mammalian context, particularly the rat [24]. This finding suggests that the frequency and distribution of Igs-containing plasma cells are in similar in avian and mammals despite belonging to different phyla.

4. Conclusion

In the present study, immunoglobulins-positive cells were detected in the lymphoid organs using immunohistochemistry. The results demonstrate that IgG- and IgM-containing plasma cells are abundantly found in the thymus, bursa of Fabricius and spleen of broilers chickens. Cells numbers were statistically greater at D₁₄ and at D₂₈ due to an immune response resulting from the NDV vaccine as well as the advancement of age.

Acknowledgements

The University Grants Commission, Bangladesh, supported this project financially. Special Thanks to Gregory Wroblewski (Lecturer, English Communication, Yamaguchi University Graduate School of Medicine, Japan) for a critical reading of this manuscript.

References

- [1] Jungherr EL. Pathology of NDV for chicken. In: Hanson RP, editor. Newcastle disease virus. An evolving pathogens. University of Wisconsin; 1964. p. 252–72.
- [2] Cortan RS, Kumar V, Robbin SL. Robbin's pathogenic basis of disease. 4th edition. Philadelphia: W.B. Saunders; 1998 [p. 163–164].
- [3] Islam MN, Khan MZI, Jahan MR, Fujinaga R, Yanai A, Kokubu K, et al. Prenatal histomorphological development of the lymphoid organs of native chickens of Bangladesh. Pak Vet J 2011;31:1–6.
- [4] Islam MN, Khan MZI, Jahan MR, Fujinaga R, Shinoda K. Ontogenic development of immunoglobulins (Igs)-positive lymphocytes in the lymphoid organs of native chickens of Bangladesh. Int J Vet Sci Med 2013;1:96–101.
- [5] Cooper RD, Peterson A, Good RA. Indication of the thymic and bursal lymphoid system of the chicken. Nature 1965;205:143–6.
- [6] Cormack DH. HAM's histology. 9th ed. Philadelphia: J.B. Lippincott Company; 1987 [p. 234–263].
- [7] Islam MN, Khan MZI, Jahan MR, Karim MR, Kon Y. Comparative studies of mucosa and immunoglobulin (Ig)-containing plasma cells in the gastrointestinal tract of broiler and native chickens of Bangladesh. J Poult Sci 2008;45: 125–31.
- [8] Khan MZI, Jahan MR, Islam MN, Haque Z, Islam MR, Kon Y. Immunoglobulin (Ig)-containing plasma cells in the Harderian gland in broiler and native chickens of Bangladesh. Tissue Cell 2007;39:141–9.
- [9] Papermaster BW, Good RA. Relative contribution of the thymus and the bursa of Fabricius to the maturation of the lympho-reticular system and immunological potential in chicken. Nature 1962;2:838–40.
- [10] Sugimura M, Hashimoto Y. Quantitative histological studies on the spleen of ducks after Neonatal thymectomy and bursectomy. J Anat 1980;131:441–52.
- [11] Khan MZI, Hashimoto Y, Iwami Y, Iwanaga T. Hormonal regulation of T-cell subsets in the oviduct: an immunohistochemical study using sex-hormone-treated chickens. J Vet Med Sci 1996;58:1161–7.
- [12] Karim MR, Khan MZI, Haque Z. Morphometrical analysis of major lymphoid organs of chemotherapy treated chickens. Bangladesh J Vet Med 2005;3:106–9.
- [13] Khan MZI, Akter SH, Islam MN, Karim MR, Islam MR, Kon Y. The effect of Selenium and vitamin E on the lymphocytes and immunoglobulin-containing plasma cells in the lymphoid organ and mucosa-associated lymphatic tissues of broiler chickens. Anat Histol Embryol 2008;37:52–9.
- [14] Honjo K, Hagiwara T, Itoh K, Takahashi E, Hirota Y. Immunohistochemical investigations of lymphocytes in the lymphoid organs of cyclophosphamide treated chickens. J Vet Med Sci 1993;55:895–7.
- [15] Nasrin M, Khan MZI, Siddiqi MNH, Masum MA. Mobilization of immunoglobulin (Ig)-containing plasma cells in Harderian gland, cecal tonsil and trachea of broilers vaccinated with Newcastle disease vaccine. Tissue Cell 2013;45:191–7.
- [16] Weibel ER. Stereological principles for morphology in electron microscopic cytology. Int Rev Cytol 1969;26:235–302.
- [17] Zar JH. Biostatistical analysis 3. Upper Saddle River, New Jersey: Prentice-Hall; 1996 [p 123–129].
- [18] Khan MZI, Hashimoto Y, Asaduzzaman M. Development of T-cells subpopulations in postnatal chicken lymphoid organs. Veterinarski Arh 1998;68:183–9.
- [19] Khan MZI, Hashimoto Y. An immunohistochemical analysis of T-cell subsets in the chicken bursa of Fabricius during postnatal stages of development. J Vet Med Sci 1996;58:1231–4.
- [20] Salam R, Aslam A, Khan SA, Saeed K, Saleem G. Effect of different routes of vaccination against Newcastle disease on lymphoid organs of broilers. Pak Vet J 2003;23:78–83.
- [21] Solcan C, Paul I, Cotea C. The dynamic of the lymphoid tissue associated to the Harderian gland in chickens after the conjunctival vaccination against New Castle disease. Lucra Stiintifice-Medicina-Veterinara,-Universitatea-de-Stiinta-Agricole-si-Medicina-Veterinara-“Ion-Ionescu-de-la-Brad”-Iasi 2000; 43: 433–436.
- [22] Bienenstock J, Gauldie J, Perey DYE. Synthesis of IgG, IgA and IgM by chicken tissues. J Immunol 1973;111:1112–8.
- [23] Ohshima K, Hiramatsu K. Immunohistochemical localization of three different immunoglobulin classes in the Harderian gland of young chicken. Cell Tissue Res 2002;34:129–33.
- [24] McDermott MR, Bienenstock J. Evidence for a common mucosal immunologic system 1. Migration of B Immunoblasts into intestinal, respiratory and genital tissues. J Immunol 1979;122:1892–8.
- [25] Chimenos ZS, Gomez E, Carrillo E, Berinstein A. Locally produced mucosal IgG in chickens immunized with conventional vaccines for Newcastle disease virus. Braz J Med Biol Res 2008;41:318–23.
- [26] Zhang X, Zhang X, Yang Q. Effect of compound mucosal immune adjuvant on mucosal and systemic immune responses in chicken orally vaccinated with attenuated Newcastle-disease vaccine. Vaccine 2007;25:3254–62.
- [27] Oliveira CA, Telles LF, Oliviera AG, Kalapothakis E, Dornelas HG, Mahecha GAB. Expression of different classes of immunoglobulin in intraepithelial plasma cells of the Harderian gland of domestic ducks, *Anas Platyrhynchos*. Vet Immunol Immunopathol 2006;113:257–66.