

# Variations in physiological, biochemical, and structural traits of photosynthesis and resource use efficiency in maize and teosintes (NADP-ME-type C<sub>4</sub>)

Takayuki Yabiku & Osamu Ueno

To cite this article: Takayuki Yabiku & Osamu Ueno (2017) Variations in physiological, biochemical, and structural traits of photosynthesis and resource use efficiency in maize and teosintes (NADP-ME-type C<sub>4</sub>), Plant Production Science, 20:4, 448-458, DOI: [10.1080/1343943X.2017.1398050](https://doi.org/10.1080/1343943X.2017.1398050)

To link to this article: <https://doi.org/10.1080/1343943X.2017.1398050>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



[View supplementary material](#)



Published online: 09 Nov 2017.



[Submit your article to this journal](#)



Article views: 1475



[View related articles](#)



Citing articles: 5 [View citing articles](#)

## Variations in physiological, biochemical, and structural traits of photosynthesis and resource use efficiency in maize and teosintes (NADP-ME-type $C_4$ )

Takayuki Yabiku<sup>a</sup> and Osamu Ueno<sup>a,b</sup>

<sup>a</sup>Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka, Japan; <sup>b</sup>Faculty of Agriculture, Kyushu University, Fukuoka, Japan

### ABSTRACT

$C_4$  plants show higher photosynthetic capacity and resource use efficiency than  $C_3$  plants. However, the genetic variations of these traits and their regulatory factors in  $C_4$  plants still remain to be resolved. We investigated physiological, biochemical, and structural traits involved in photosynthesis and photosynthetic water and nitrogen use efficiencies (PWUE and PNUE) in 22 maize lines and four teosinte lines from various regions of the world. Net photosynthetic rate ( $P_N$ ) ranged from 32.1 to 46.5  $\mu\text{mol m}^{-2} \text{s}^{-1}$ .  $P_N$  was positively correlated with stomatal conductance, transpiration rate, and chlorophyll, nitrogen and soluble protein contents of leaves, but not with specific leaf weight.  $P_N$  was positively correlated with the activities of ribulose-1,5-bisphosphate carboxylase/oxygenase and the  $C_4$ -acid decarboxylases, NADP-malic enzyme and phosphoenolpyruvate carboxykinase, but not with the activity of phosphoenolpyruvate carboxylase. Leaf structural traits (stomatal parameters, leaf thickness, and interveinal distance) were not correlated with  $P_N$ . These data suggest that physiological and biochemical traits are involved in the genetic variation of  $P_N$ , but structural traits are not directly involved. PWUE is in the lower class of values reported for  $C_4$  plants, whereas PNUE is in the highest class of values reported for  $C_4$  plants. PNUE was negatively correlated with leaf nitrogen content but not significantly correlated with  $P_N$ . PWUE was not correlated with  $\delta^{13}\text{C}$  values of leaves, indicating difficulty in using  $\delta^{13}\text{C}$  values as an indicator of PWUE of maize. In general, teosinte lines showed lower  $P_N$  but higher PWUE than maize lines.

### ARTICLE HISTORY

Received 26 July 2017  
Revised 28 September 2017  
Accepted 24 October 2017

### KEYWORDS

Maize; NADP-malic enzyme type; phosphoenolpyruvate carboxykinase; photosynthetic nitrogen use efficiency; photosynthetic rate; photosynthetic water use efficiency; teosinte

### CLASSIFICATION

Crop Physiology


### Introduction

Photosynthetic traits of leaves are one of the most important physiological factors responsible for plant productivity. The improvement of photosynthetic traits promises further increases in plant productivity (Evans, 2013; Zhu et al., 2010). The genetic variation in photosynthetic traits found in both crop and wild species includes a potential to improve crop photosynthesis and ultimately productivity but is largely unexplored (McCouch, 2004; Flood et al., 2011). Photosynthesis is complicatedly regulated by biophysical, biochemical, physiological, and structural traits of leaves. However, our understanding of the regulatory processes is still insufficient (Evans, 2013). The factors that cause the genetic variation in photosynthesis also remain largely unknown (Flood et al., 2011).

It is well known that  $C_4$  plants have higher photosynthetic rate and productivity than  $C_3$  plants (Brown, 1999; Osmond et al., 1982). This is attained by a  $\text{CO}_2$  concentrating mechanism (CCM) operating in  $C_4$  plants. In  $C_4$  leaves,

mesophyll and bundle-sheath (BS) cells are differentiated and surround vascular bundles. In  $C_4$  photosynthesis, atmospheric  $\text{CO}_2$  entering through stomata is fixed by phosphoenolpyruvate carboxylase (PEPC) in mesophyll cells. The  $C_4$  acids produced are transported to BS cells, where they are decarboxylated by  $C_4$  acid decarboxylase. Released  $\text{CO}_2$  is fixed by Rubisco. This biochemical process raises the  $\text{CO}_2$  concentration around Rubisco in BS cells and thus reduces photorespiration (Hatch, 1987). The quantitative balance between mesophyll and BS cells is also required to attain the intimate cooperation between  $C_4$  and  $C_3$  cycles (Dengler et al., 1994). In general,  $C_4$  leaves have a denser vascular system than  $C_3$  leaves. It is thought that this structural trait is associated with rapid translocation of photosynthates in  $C_4$  plants (Leonardos & Grodzinski, 2000; Ueno et al., 2006). Because of the complex mechanism of  $C_4$  photosynthesis, the factors determining photosynthetic rate in  $C_4$  plants still remain to be resolved (von Caemmerer & Furbank, 2016). It is suggested that  $\text{CO}_2$  delivery process in mesophyll cells, activities and

**CONTACT** Osamu Ueno  [uenoos@agr.kyushu-u.ac.jp](mailto:uenoos@agr.kyushu-u.ac.jp)

 Supplementary material for this article can be accessed at <https://doi.org/10.1080/1343943X.2017.1398050>

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

properties of  $C_4$  and  $C_3$  photosynthetic enzymes, substrate regeneration in  $C_4$  and  $C_3$  cycles, transport and diffusion of metabolites, electron transport and light capture in chloroplasts, and  $CO_2$  leakiness from BS cells are involved in the regulation of  $C_4$  photosynthesis (reviewed by von Caemmerer & Furbank, 2016). It also appears that biochemical, physiological, and structural traits of leaves are intricately involved in the genetic variation in photosynthetic rate of  $C_4$  plants (Tsutsumi et al., 2017).

Depending on the  $C_4$  acid decarboxylation system in BS cells, the  $C_4$  photosynthetic pathway is classified into three types: NADP-malic enzyme (NADP-ME), NAD-malic enzyme (NAD-ME), and phosphoenolpyruvate carboxylase (PCK) (Hatch, 1987). The NADP-ME-type  $C_4$  grasses include crops with high biomass productivity, such as maize, sorghum, and Napier grass (Brown, 1999; Carpita & McCann, 2008). In general, each  $C_4$  species is thought to use only one of the 3 decarboxylation systems (Hatch, 1987). However, some  $C_4$  species may use more than one. Maize (*Zea mays* ssp. *mays*), a model NADP-ME-type  $C_4$  grass, uses both NADP-ME and PCK (Walker et al., 1997; Wingler et al., 1999). However, the physiological significance of the dual  $C_4$  acid decarboxylation system in  $C_4$  plants is unclear (Furbank, 2011; Koteyeva et al., 2015).

Because  $C_4$  plants have a CCM, their photosynthetic water and nitrogen use efficiencies (PWUE and PNUE) are higher than those of  $C_3$  plants (Brown, 1977; Osmond et al., 1982). These traits of  $C_4$  plants provide advantages for survival in natural habitats and are also useful for sustainable agriculture. PNUE values of NADP-ME-type  $C_4$  grasses are higher than those of NAD-ME-type  $C_4$  grasses, whereas

PWUE values are not significantly different (Ghannoum et al., 2001, 2005, 2011). Further extensive studies would be required to clarify the genetic variation in resource use efficiency among  $C_4$  plants.

Maize is important as a grain, forage, and bioenergy crop (Carpita & McCann, 2008). Previous studies have addressed the genetic variation in the photosynthetic rate of maize. Some studies reported a large genetic difference in photosynthetic rate (Crosbie et al., 1977; Heichel & Musgrave, 1969), whereas others found a small difference (Baer & Schrader, 1985; Duncan & Hesketh, 1968). Previous studies suggested that Rubisco and pyruvate, Pi dikinase (PPDK) are rate-limiting enzymes in  $C_4$  photosynthesis in maize (Baer & Schrader, 1985; Usuda, 1984; Usuda et al., 1985; von Caemmerer & Furbank, 2016). However, it is still uncertain whether other physiological, biochemical, and structural factors are involved in the variation in photosynthetic rate of maize.

The aim of this study is to investigate the genetic variations in photosynthetic rate and resource use efficiency in maize lines from various regions of the world. Another aim is to determine which factors in physiological, biochemical, and structural traits of leaves regulate these genetic variations in maize lines. We examined here various traits of leaves such as gas exchange traits, chlorophyll (Chl) and nitrogen (N) contents, carbon isotope ratio, activities of  $C_3$  and  $C_4$  enzymes, stomatal parameters, and interveinal distance (IVD) for these maize lines. On the other hand, because wild lines are valuable as genetic resources to improve physiological traits of crops (Flood et al., 2011), four lines of teosintes (*Z. mays* ssp. *mexicana*, *Z. diploperennis*, *Z. perennis*, *Z. nicaraguensis*) were added for comparison.

**Table 1.** Maize and teosinte lines examined in this study.

Species	Line	Country
<i>Zea mays</i> ssp. <i>mays</i>	B73	Iowa, USA
	HP301	Indiana, USA
	IL14H	Illinois, USA
	Ky21	Kentucky, USA
	Mo18W	Missouri, USA
	OH7B	Ohio, USA
	P39	Indiana, USA
	WF9	Indiana, USA
	CM109	Canada
	CML69	Mexico
	Pipoca 4	Brazil
	Pisinga Purpura	Peru
	CB44	Netherlands
	Kuma Mais	Netherlands
	F575	France
	I.C.A.R. 54	Romania
	TZ13	Nigeria
	PI 195114	Ethiopia
	Homedale	South Africa
	Georgian local 1	USSR
Pakistan	Pakistan	
Ki3	Thailand	
<i>Z. mays</i> ssp. <i>mexicana</i>	Ames 8083	Mexico
<i>Z. diploperennis</i>	PI 441930	Mexico
<i>Z. perennis</i>	Ames 21875	Mexico
<i>Z. nicaraguensis</i>	PI 615697	Nicaragua

## Materials and methods

### Plant materials and growth conditions

Twenty-two maize lines (*Z. mays* ssp. *mays*) and four teosinte lines (one line per species: *Z. mays* ssp. *mexicana*, *Z. diploperennis*, *Z. perennis*, and *Z. nicaraguensis*) were used in this study (Table 1). These lines were selected from a wide range of countries. Seeds were provided by the Plant Introduction Station, Agricultural Research Service, USDA, and the NARO Genebank, Tsukuba, Japan. They were germinated in nursery boxes filled with loam granules and were grown for 10 days in a greenhouse in an experimental field of Kyushu University (33°35'N, 130°23'E) during summer 2014. Five seedlings per line were transplanted into 5-L pots (one plant per pot) filled with a sandy loam mixed with chemical fertilizer containing 1.0 g each of nitrogen, phosphorus, and potassium. Plants were then grown outdoors at a mean air temperature of 26 °C and midday

PPFD of full sunlight exceeding 2000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Plants were watered daily to avoid drying of soil. At 2 weeks after transplanting, the chemical fertilizer containing .6 g each of elements was supplied. At 3–4 weeks after transplanting, physiological, biochemical, and structural traits of photosynthesis were examined in fully expanded upper leaves of 3–5 plants per line. At this time, plant height was 80–120 cm in maize lines and 50–80 cm in teosinte lines, and all plants except for CM109 and CB44 were in vegetative stage. However, these two maize lines initiated to develop tassels (Supplemental data 1).

### **Gas exchange and PWUE**

Gas exchange in leaves was measured using a portable photosynthesis system (Li-6400XT; Li-COR, Lincoln, NE, USA). Gas exchange parameters – net photosynthetic rate ( $P_N$ ), stomatal conductance ( $g_s$ ), transpiration rate ( $T_r$ ), and intercellular  $\text{CO}_2$  concentration ( $C_i$ ) – were measured at a PPFD of 2000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , leaf temperature of  $30.0 \pm .5^\circ\text{C}$ , relative humidity of  $60\% \pm .5\%$ , and ambient  $\text{CO}_2$  concentration ( $C_a$ ) of 380  $\mu\text{mol mol}^{-1}$ . PWUE was calculated by dividing  $P_N$  by  $T_r$ .

### **Chl and N contents, specific leaf weight, and PNUE**

Chl content and specific leaf weight (SLW) were measured in the same leaves used for gas exchange measurements. Chl was extracted from the leaves (3.4  $\text{cm}^2$ ) in 80% acetone, and Chl content was measured spectrophotometrically according to Arnon (1949). Leaves (5.7  $\text{cm}^2$ ) were air-dried for 2 days at  $80^\circ\text{C}$  and weighed, and SLW was calculated by dividing dry mass by leaf area. Leaf N content was determined in lower leaves next to the ones used for gas exchange measurements. Leaves were air-dried as described above and ground to powder. The N content in .3 g of leaf powder was determined using a micro-Kjeldahl procedure (Tsutsumi et al., 2017). PNUE was calculated by dividing  $P_N$  by N content.

### **Enzyme assays and leaf soluble protein content**

Parts of the same leaves used for gas exchange measurements were sampled between 10:00 and 12:00 on a clear day, immediately frozen in liquid nitrogen, and stored at  $-80^\circ\text{C}$ . For enzyme assay, leaves (.2 g fresh mass) were ground on ice with a pestle in a mortar containing 1 mL of grinding medium [50 mM HEPES-KOH (pH 7.5), 1 mM EDTA-2Na, 5 mM dithiothreitol, 10 mM  $\text{MgCl}_2$ , and .02% (v/v) Triton X-100] containing .5% (w/v) bovine serum albumin, 5 mg of polyvinylpyrrolidone, and .1 g of quartz sand. The homogenates were filtered through two layers of gauze, the filtrates were centrifuged for 5 min at  $10,000 \times g$

at  $4^\circ\text{C}$ , and the supernatants were used for the enzyme assay. An aliquot of the filtrate was taken for determination of Chl content.

Activities of photosynthetic enzymes were assayed spectrophotometrically in 1-mL reaction mixtures at  $30^\circ\text{C}$ . Activities of PEPC and NADP-ME were assayed as described by Ueno and Sentoku (2006). The activity of PCK was assayed in the carboxylase direction following NADH oxidation according to Sharwood et al. (2014). The total activity of Rubisco was measured as described by Ueno and Sentoku (2006) except that 5 U phosphoglycerate kinase, 5 U glyceraldehyde 3-phosphate dehydrogenase, and 5 U phosphocreatine kinase were used. In the assay of Rubisco activity, the supernatant was preincubated in the presence of 10 mM  $\text{NaHCO}_3$  and 10 mM  $\text{MgCl}_2$  at  $25^\circ\text{C}$  for 10 min.

For measurements of leaf soluble protein (LSP) content, leaves (.1 g fresh mass) were ground on ice, and supernatants were obtained as for enzyme assays except that bovine serum albumin was omitted and 1 mM phenylmethylsulfonyl fluoride and .002% (w/v) leupeptin were added to the grinding medium. The LSP content was measured according to Bradford (1976).

### **Carbon isotope ratio**

A part of each leaf used for gas exchange measurement was air-dried at  $80^\circ\text{C}$  and separately ground in a mortar with a pestle. The same amounts of powder from each leaf were thoroughly mixed, and 2 mg of the mixture was used for measurement of  $^{12}\text{C}$  and  $^{13}\text{C}$  contents as described by Sato and Suzuki (2010). The isotope ratio was expressed in  $\delta$  notation in parts per million (‰) with respect to the Pee Dee belemnite standard.

### **Structural traits**

The middle portions of leaves used for gas exchange measurements were fixed in formalin–acetic acid–alcohol solution for 1 day and cleared according to Ueno et al. (2006). Stomatal density (SD), guard cell length (GL), and IVD were measured under a light microscope. Stomata were counted on each leaf surface in four .38- $\text{mm}^2$  fields per leaf at  $300\times$  magnification. SD was calculated as the sum of the number of stomata on both sides per unit leaf area. GL of 20 cells on each side (40 cells in total) was measured with a micrometer at  $600\times$  magnification. IVD was the mean of 10 measurements of the distance between centers of adjacent small longitudinal veins.

The middle portions of leaves were also fixed in 3% (v/v) glutaraldehyde in 50 mM sodium phosphate buffer (pH 6.8) at room temperature for 1.5 h. After washing with phosphate buffer, they were post-fixed in 2% (w/v)  $\text{OsO}_4$

in sodium phosphate buffer (pH 6.8) for 2 h, dehydrated through an acetone series, and embedded in Quetol resin (Nisshin-EM Co. Ltd., Shinjuku, Tokyo, Japan) at 70 °C. The samples were transversely sectioned at 1  $\mu\text{m}$  thickness with a glass knife on an ultramicrotome and stained with 1% toluidine blue O. Leaf thickness was measured in ImageJ software (Schneider et al., 2012) as the mean of 10 points per section.


### Statistical analysis

Data were analyzed using BellCurve for Excel (Social Survey Research Information Co., Ltd., Shinjuku, Tokyo, Japan). One-way analysis of variance (ANOVA) was used for all parameters. Pearson's correlation coefficients between the parameters were calculated.

## Results

### Gas exchange and PWUE


Gas exchange traits differed significantly among the maize and teosinte lines examined ( $p < .01$ ; Supplemental data 2).  $P_N$  ranged from 32.1  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (teosinte line Ames 21875) to 46.5  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (maize line PI 195114) with a mean of 38.9  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (Figure 1(A); Supplemental data 2). The three teosinte lines of species other than *Z. mays* (PI 441930, Ames 21875, PI 615697) had lower  $P_N$  than most *Z. mays* lines had. However, teosinte line Ames 8083 (*Z. mays* ssp. *mexicana*) had an intermediate  $P_N$  value (Figure 1(A)).  $P_N$  was positively correlated with  $g_s$  (Figure 2(A)) and  $T_r$  (Figure 2(B)) but not with  $C_i/C_a$  (Table 2). PWUE ranged from 5.25  $\text{mmol mol}^{-1}$  (maize line IL14H) to 7.02  $\text{mmol mol}^{-1}$  (Ames 21875), with a mean of 6.07  $\text{mmol mol}^{-1}$  (Figure 1(B);



**Figure 1.** Variations in (A) net photosynthetic rate ( $P_N$ ), (B) photosynthetic water use efficiency (PWUE), and (C) photosynthetic nitrogen use efficiency (PNUE) in leaves of maize and teosinte lines.

Notes: Means  $\pm$  SD ( $n = 3-5$ ). Dashed lines show mean values. Black, maize (*Z. mays* ssp. *mays*); gray, *Z. mays* ssp. *mexicana*; white, *Z. diploperennis*, *Z. perennis*, and *Z. nicaraguensis*.





**Figure 2.** Relationships between  $P_N$  and (A) stomatal conductance ( $g_s$ ), (B) transpiration rate ( $T_r$ ), (C) chlorophyll (Chl) content, (D) leaf soluble protein (LSP) content, (E) leaf N content, and (F) specific leaf weight (SLW) in leaves of maize and teosinte lines.

Notes: Circles are the mean values of each line ( $n = 3-5$ ): black, maize (*Z. mays* ssp. *mays*); gray, *Z. mays* ssp. *mexicana*; white, *Z. diploperennis*, *Z. perennis*, and *Z. nicaraguensis*. Significant at  $P$ : \*\* $< .01$ ; \*\*\* $< .001$ . <sup>NS</sup>not significant.

Supplemental data 2). PWUE was high in all teosinte lines except PI 615697 (Figure 1(B); Supplemental data 2). PWUE was negatively correlated with  $P_N$ ,  $g_s$ ,  $T_r$  and  $C_i/C_a$  (Table 2).

### Other physiological traits

All these traits except  $\delta^{13}\text{C}$  values showed significant differences among lines ( $p < .001$ ; Supplemental data 3). Chl and LSP contents showed large variations among lines, with maximum-to-minimum ratios of 2.1 and 2.6, respectively (Supplemental data 3). The contents of Chl (Figure 2(C)), LSP (Figure 2(D)), and N (Figure 2(E)) were positively correlated with  $P_N$  whereas that of SLW (Figure 2(F)) was not. PNUE ranged from 510  $\mu\text{mol mol N}^{-1} \text{s}^{-1}$  (CM109) to 862  $\mu\text{mol mol N}^{-1} \text{s}^{-1}$  (Pipoca 4), with a mean of 700  $\mu\text{mol mol N}^{-1} \text{s}^{-1}$  (Figure 1(C); Supplemental data 3). Among the four teosinte lines, PNUE was high in PI

441903 and Ames 8083 and low in PI 615697 and Ames 21875. PNUE was not correlated with any gas-exchange or physiological traits except leaf N content (Table 2). The  $\delta^{13}\text{C}$  values were not correlated with any gas exchange or physiological traits (Figure 3; Table 2).

### Activities of photosynthetic enzymes

PEPC activity was not significantly correlated with  $P_N$  (Figure 4(A)), whereas NADP-ME and Rubisco activities were positively correlated with  $P_N$  (Figure 4(B) and (D)). PCK activity varied considerably among the lines (Supplemental data 4) and was high in two lines with high  $P_N$  (PI 195114 and CM 109; Supplemental data 4). The sum of NADP-ME and PCK activities (capacity for  $\text{C}_4$  acid decarboxylation) varied among lines, with a maximum-to-minimum ratio of 3.0 (Supplemental data 4). PCK activity and the capacity for  $\text{C}_4$  acid decarboxylation were positively correlated with  $P_N$  (Figure 4(C); Table 2). However, the contribution ratio of PCK activity to the total  $\text{C}_4$  acid decarboxylation capacity (PCK ratio) was not correlated with  $P_N$  (Table 2). Chl content and LSP were positively correlated with the activities of Rubisco, NADP-ME, and PCK but not with that of PEPC (Table 2).

### Structural traits

Variations in leaf thickness, IVD, and GL were small, with maximum-to-minimum ratios of 1.52 for leaf thickness, 1.40 for IVD, and 1.35 for GL (Supplemental data 5). Variations in SD and SD  $\times$  GL were larger, with maximum-to-minimum ratios of 2.43 for SD and 2.26 for SD  $\times$  GL (Supplemental data 5). The ratio of adaxial to abaxial GL was  $1.04 \pm .02$  and the ratio of adaxial to abaxial SD was  $.71 \pm .06$  (data not shown). There were no correlations between these structural traits and gas-exchange or physiological traits (Supplemental data 6), except that PWUE was weakly negatively correlated with SD ( $r = -.412$ ;  $p < .05$ ) and SD  $\times$  GL ( $r = -.415$ ;  $p < .05$ ). On the other hand, leaf thickness was positively correlated with IVD, whereas SD was negatively correlated with GL (Table 3).

## Discussion

### Physiological traits

Our study showed that  $P_N$  in maize and teosinte lines ranged from 32.1–46.5  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , a factor of 1.45 times. Although some studies have reported a variation of  $>2$  times in  $P_N$  of maize genotypes (Crosbie et al., 1977; Heichel & Musgrave, 1969), we found no such large difference. On the other hand, Duncan and Hesketh (1968) reported a variation in  $P_N$  among maize cultivars similar

**Table 2.** Correlation coefficients (*r*) from linear regression analysis and statistical significance of the relationships between physiological and biochemical traits in maize and teosinte lines. *P<sub>N</sub>* net photosynthetic rate; *g<sub>s</sub>* stomatal conductance; *T<sub>r</sub>* transpiration rate; *C<sub>i</sub>/C<sub>a</sub>* intercellular CO<sub>2</sub> to ambient CO<sub>2</sub> concentration; *PWUE*, photosynthetic water use efficiency; *LSP*, leaf soluble protein; *Chl*, chlorophyll; *SLW*, specific leaf weight; *PNUE*, photosynthetic nitrogen use efficiency;  $\delta^{13}\text{C}$ , carbon isotope ratio; *PEPC*, phosphoenolpyruvate carboxylase; *ME*, malic enzyme; *PCK*, phosphoenolpyruvate carboxykinase; *DC*, capacity for C<sub>4</sub> acid decarboxylation (=NADP-ME activity + PCK activity); *PCK* ratio, [=PCK activity/(NADP-ME activity + PCK activity)].


	<i>P<sub>N</sub></i>	<i>g<sub>s</sub></i>	<i>T<sub>r</sub></i>	<i>C<sub>i</sub>/C<sub>a</sub></i>	<i>PWUE</i>	<i>LSP</i> content	<i>Chl</i> content	<i>Leaf N</i> content	<i>SLW</i>	<i>PNUE</i>	$\delta^{13}\text{C}$	<i>Rubisco</i> activity	<i>PEPC</i> activity	<i>NADP-ME</i> activity	<i>PCK</i> activity	<i>DC</i>	<i>PCK</i> ratio
<i>P<sub>N</sub></i>	1																
<i>g<sub>s</sub></i>	.813***	1															
<i>T<sub>r</sub></i>	.878***	.961***	1														
<i>C<sub>i</sub>/C<sub>a</sub></i>	.169NS	.632***	.557**	1													
<i>PWUE</i>	-.510***	-.805***	-.819***	-.835***	1												
<i>LSP</i> content	.718***	.671***	.724***	.276NS	-.592**	1											
<i>Chl</i> content	.726***	.558**	.671***	.129NS	-.433*	.809***	1										
<i>Leaf N</i> content	.581**	.372NS	.487*	.080NS	-.342NS	.576**	.614***	1									
<i>SLW</i>	-.095NS	-.092NS	-.102NS	-.076NS	.069NS	-.233NS	-.279NS	-.030NS	1								
<i>PNUE</i>	-.098NS	.040NS	-.056NS	-.010NS	.135NS	-.287NS	-.311NS	-.832***	-.161NS	1							
$\delta^{13}\text{C}$	-.034NS	.095NS	.037NS	.045NS	.022NS	-.026NS	-.268NS	-.247NS	.252NS	.214NS	1						
<i>Rubisco</i> activity	.610***	.652***	.692***	.419*	-.615***	.668***	.667***	.301NS	-.003NS	-.070NS	.176NS	1					
<i>PEPC</i> activity	-.055NS	.013NS	-.055NS	.051NS	.018NS	-.099NS	-.234NS	.114NS	.205NS	-.129NS	-.008NS	-.211NS	1				
<i>NADP-ME</i> activity	.501**	.492*	.519**	.223NS	-.440*	.541**	.389*	.344NS	.212NS	-.141NS	.039NS	.519**	.368NS	1			
<i>PCK</i> activity	.521**	.439*	.469*	.117NS	-.225NS	.438*	.423*	.633***	-.104NS	-.382NS	-.149NS	.188NS	.396*	.437*	1		
<i>DC</i>	.457*	.491*	.509**	.267NS	-.438*	.467*	.326NS	.385NS	.150NS	-.182NS	.012NS	.414*	.486*	.958***	.597**	1	
<i>PCK</i> ratio	.387NS	.186NS	.225NS	-.107NS	.001NS	.344NS	.319NS	.560**	-.088NS	-.400*	-.100NS	.119NS	.313NS	.175NS	.860***	.288NS	1

\*Significant at *P* < .05;


\*\*\*< .01;

\*\*\*< .001;

NS not significant.



**Figure 3.** Relationship between photosynthetic water use efficiency (PWUE) and  $\delta^{13}\text{C}$  in leaves of maize and teosinte lines. Notes: Circles are the mean values of each line ( $n = 3\text{--}5$ ). Circle shading is as in Figure 2. <sup>NS</sup>not significant.



**Figure 4.** Relationships between  $P_N$  and activities of (A) PEPC, (B) NADP-malic enzyme (NADP-ME), (C) PEPC, and (D) Rubisco in leaves of maize and teosinte lines.

Notes: Circles are the mean values of each line ( $n = 3\text{--}5$ ). Circle shading is as in Figure 2. Significant at  $P$ : \*\*  $< .01$ ; \*\*\*  $< .001$ . <sup>NS</sup>not significant. Broken lines ( $y = x$ ) show enzyme activities that would be required for equal  $P_N$ .

to that in our study. It seems likely that the differences in growth condition, plant age, and measurement method of  $P_N$  bring about such different results.

We found a positive correlation between  $P_N$  and  $g_s$  (Figure 2(A)). This relationship has been reported in various  $C_3$  and  $C_4$  species (Fernandez et al., 2015; Tsutsumi et al., 2017; Wong et al., 1985). Because maize has a CCM,  $\text{CO}_2$  diffusion in stomata may not be the primary  $P_N$  regulator. The  $P_N$  of  $C_4$  plants is saturated at the  $C_i$  observed at

the atmospheric  $\text{CO}_2$  concentration, whereas the  $P_N$  of  $C_3$  plants is not saturated until a much higher  $\text{CO}_2$  concentration (Percy & Ehleringer, 1984; Wong et al., 1985). In preliminary measurements of some maize lines, we also confirmed that  $P_N$  is saturated at  $C_i \approx 100 \mu\text{mol mol}^{-1}$ , which is observed at  $C_a = 380 \mu\text{mol mol}^{-1}$  (data not shown). This suggests that  $P_N$  would not greatly increase even though  $g_s$  alone increased and thereby higher  $C_i$  was attained within the leaf. Therefore, the effect of non-stomatal factors on the genetic variation in  $P_N$  of maize and teosinte lines cannot be ruled out. On the other hand, there was a close relationship between  $P_N$  and  $g_s$  in maize lines. The reason is unknown. However, the coordinated mechanism between mesophyll photosynthesis and stomata (Lawson et al., 2014) may be involved in this relationship. According to this hypothesis, the concentration of  $\text{CO}_2$  inside the leaf would help maintain the coordination of the mesophyll photosynthesis with stomatal aperture (Lawson et al., 2014).

The Chl, LSP, and leaf N contents were positively correlated with  $P_N$  (Figure 2(C)–(E)), as reported in many plant species, including maize (Wong et al., 1985; Sage & Percy, 1987; Tsutsumi et al., 2017), because all these parameters are closely associated with the contents of photosynthetic pigments and enzymes. No significant correlation was detected between  $P_N$  and SLW (Figure 2(F)), as in *Saccharum* (Nose et al., 1994) and *Oryza* species (Kiran et al., 2013), although positive correlations were found in leguminous (Pearce et al., 1969), cruciferous (Suresh et al., 1997), and *Amaranthus* species (Tsutsumi et al., 2017). Thus, the relationship between  $P_N$  and SLW varies with species.

### Biochemical traits

$P_N$  was positively correlated with activities of NADP-ME, PCK, and Rubisco, but not with that of PEPC (Figure 4). A positive correlation between  $P_N$  and Rubisco activity was found in  $C_4$  and  $C_3$  species (Ghannoum et al., 2011; Tsutsumi et al., 2017; von Caemmerer et al., 1997). In our study, a considerable number of maize and teosinte lines showed lower Rubisco activities than  $P_N$  (Figure 4(D)). Baer and Schrader (1985) reported that, in Rubisco of maize lines, total activities are lower than initial activities. Therefore, this enzymatic trait may be involved in the lower Rubisco activities relative to  $P_N$ , because we measured total activities. However, the possibility of deactivation and/or degradation of Rubisco during extraction cannot be ruled out. In NADP-ME-type  $C_4$  grasses, a positive correlation between  $P_N$  and NADP-ME activity was also reported (Nose et al., 1994; Usuda, 1984). Using antisense RNA, Pengelly et al. (2012) suggested that NADP-ME is unlikely to be a rate-limiting enzyme in  $C_4$



**Table 3.** Correlation coefficients ( $r$ ) from linear regression analysis and statistical significance of the relationships between structural traits of maize and teosinte lines. IVD, interveinal distance; SD, stomatal density; GL, guard cell length.

	Leaf thickness	IVD	SD	GL	SD×GL
Leaf thickness	1				
IVD	.645***	1			
SD	-.371 <sup>NS</sup>	-.487*	1		
GL	.257 <sup>NS</sup>	.225 <sup>NS</sup>	-.686***	1	
SD×GL	-.352 <sup>NS</sup>	-.526	.953***	-.439*	1

\*Significant at  $P < .05$ ;

\*\* $< .01$ ;

\*\*\* $< .001$ ;

<sup>NS</sup>not significant.

photosynthesis of *Flaveria bidentis*, an NADP-ME-type  $C_4$  dicot, because  $P_N$  of transgenic plants was not significantly reduced until 40% reduction of NADP-ME activity in the wild-type plants. At present, it is unknown whether this relationship between NADP-ME activity and  $P_N$  found in the  $C_4$  dicot is applicable to NADP-ME-type  $C_4$  grasses as well. However, these data suggest that the relationship between NADP-ME activity and  $P_N$  may more or less differ among species.

Maize has two  $C_4$  acid decarboxylation enzymes, NADP-ME and PCK (Walker et al., 1997; Wingler et al., 1999). Our study has revealed, for the first time, a considerable genetic variation in PCK activity among maize and teosinte lines. The PCK system in maize has been estimated to contribute 10–14% of the carbon in BS (Arrivault et al., 2017). This ratio was within the range found in our study (3.6–19.8%; Supplemental data 4).

The physiological significance of  $C_4$  acid decarboxylation depending on PCK in maize remains to be understood (Furbank, 2011). Sharwood et al. (2014) reported that the ratio of PCK activity to NADP-ME activity changes in maize exposed to shade and salinity. The parallel operation of the two  $C_4$  acid decarboxylation systems may compensate each other under certain environmental conditions (Bellasio & Griffiths, 2014) and may be energetically profitable. The  $C_4$  acid decarboxylation depending on PCK generates phosphoenolpyruvate (PEP). If PEP is transported to mesophyll cells, ATP needed for the conversion of pyruvate to PEP by PPDK may be saved. We detected no positive correlation between  $P_N$  and PCK ratio in maize and teosinte lines (Table 2). Therefore, the difference in  $P_N$  among maize and teosinte lines would not result from the difference in the contribution ratio of PCK activity to the total  $C_4$  acid decarboxylation capacity. Further investigation would be required to understand the physiological significance of decarboxylation by PCK in maize.

Our study suggests that biochemical processes from  $C_4$  acid decarboxylation to re-fixation of  $CO_2$  by Rubisco

are involved in the genetic variation in  $P_N$ . However, we did not examine the activity of PPDK, which is well known to be the rate-limiting factor in  $C_4$  photosynthesis (Baer & Schrader, 1985; Usuda, 1984; von Caemmerer & Furbank, 2016), because a preliminary study on several lines of maize showed that their activities of PPDK were unstable and unreliable. The relationships between  $P_N$  and PPDK activity and between PPDK and PCK activities remain to be explored.

### Structural traits

$P_N$  was not correlated with leaf thickness or IVD (Supplemental data 6). In  $C_3$  plants, photosynthesis is performed within single mesophyll cells. Therefore,  $C_3$  leaves can change their structure in response to environmental change. The genetic variation in leaf structure of  $C_4$  plants is restricted to a narrower range than that of  $C_3$  plants, because  $C_4$  photosynthesis requires a strict quantitative balance between two types of photosynthetic cells (Dengler et al., 1994; Ghannoum et al., 2011; Tsutsumi et al., 2017). The maximum-to-minimum ratios of leaf thickness and IVD in maize and teosinte lines were generally smaller than those of physiological and biochemical traits. This may conceal the possible relationships between  $P_N$  and these structural traits.

We found a negative correlation between SD and GL (Table 3), as reported in other species (Büßis et al., 2006; Lawson & Blatt, 2014; Tsutsumi et al., 2017). The genetic variation in SD (2.43 times) was greater than that in GL (1.35 times; Supplemental data 5), suggesting physical and genetic limitations on the range of alterations in GL, whereas SD is much more flexible (Tsutsumi et al., 2017).

The stomatal parameters were not correlated with  $P_N$ ,  $g_s$ , or  $T_r$  (Table 3). It seems that an increase in SD would increase  $P_N$  by increasing  $g_s$ . Several studies reported positive correlations between SD and  $P_N$  in various species, but other studies reported no such correlations (reviewed by Lawson & Blatt, 2014). Importantly, stomatal anatomical features such as SD and GL define the maximum theoretical conductance, whereas  $P_N$  is the actual physiological outcome (Dow et al., 2014; Lawson & Blatt, 2014). The relationships between stomatal anatomical features and gas exchange parameters would also be affected by environmental factors such as vapor pressure deficit (Kawamitsu et al., 2002). Therefore, more detailed analyses under different conditions would be needed.

### Resource use efficiency

$C_4$  plants can maintain high  $CO_2$  concentration within BS cells owing to the CCM. This allows  $C_4$  plants to have higher PNUE than that of  $C_3$  plants (Brown, 1977; Ghannoum

et al., 2011). A comparative study of  $C_4$  subtypes in grasses showed higher PNUE in the NADP-ME type than in the NAD-ME type, because Rubisco turnover rate is faster in the former than in the latter (Ghannoum et al., 2005). In our study, the maximum-to-minimum PNUE ratio of maize and teosinte lines was 1.69, and the mean was  $700 \mu\text{mol mol N}^{-1} \text{s}^{-1}$  (Supplemental data 3), which was far higher than PNUE in  $C_3$  plants and was in the highest class of PNUE values previously reported in  $C_4$  plants (Ghannoum et al., 2005; Taylor et al., 2010; Togawa & Ueno, 2015; Tsutsumi et al., 2017; Vogan & Sage, 2011). Our study suggests that the genetic variation in PNUE in maize and teosinte lines depends on leaf N content but not on  $P_N$  (Table 2), because lines with lower leaf N content showed higher PNUE. Wild relatives of cultivated crops often inhabit more severe environment than cultivated conditions and possess useful traits that have been lost in cultivated crops during domestication (Hamaoka et al., 2013; Scafaro et al., 2010). Therefore, we expected that some teosinte lines would have higher PNUE than maize lines, but found no such trend (Figure 1(C)).

The maximum-to-minimum PWUE ratio of maize and teosinte lines was 1.34 times (Supplemental data 2), which was lower than that of PNUE. The mean PWUE was  $6.07 \mu\text{mol mol}^{-1}$  (Supplemental data 2) and was in the lower class of PWUE values previously reported in  $C_4$  plants (Osmond et al., 1982; Togawa & Ueno, 2015; Tsutsumi et al., 2017).

In  $C_3$  plants, there is a positive correlation between  $\delta^{13}\text{C}$  values and PWUE, and  $\delta^{13}\text{C}$  is useful for screening cultivars for high PWUE (Farquhar & Richards, 1984). In this study, the  $\delta^{13}\text{C}$  values did not vary greatly among maize and teosinte lines (Supplemental data 3) and were not significantly correlated with PWUE (Figure 3). In sorghum ( $C_4$ ), O'Leary (1988) also found no significant correlation between  $\delta^{13}\text{C}$  values and PWUE in 120 lines, whereas Henderson et al. (1998) reported a weak but significant correlation in 30 lines. The carbon isotope ratio in plant dry matter reflects carbon isotope discrimination ( $\Delta$ ) during photosynthesis (Cernusak et al., 2013). In  $C_3$  plants,  $\Delta$  is well correlated with  $C_i/C_a$  and hence can be used as an index of PWUE. In  $C_4$  plants, the variation in  $C_i/C_a$  is smaller than in  $C_3$  plants and  $\Delta$  is related to both  $\text{CO}_2$  leakiness from BS cells and to  $C_i/C_a$  (Cernusak et al., 2013; Ghannoum et al., 2011; Henderson et al., 1998). For these reasons, it would not be easy to use  $\delta^{13}\text{C}$  values as an indicator of PWUE of  $C_4$  plants.

We found weak negative correlations between PWUE and SD or  $\text{SD} \times \text{GL}$  (Table 3). A decrease in SD would reduce water loss from leaves. However, it may also decrease  $P_N$  while increasing PWUE, because  $P_N$  is negatively correlated with PWUE (Table 2). It is worth noting that all teosinte

lines except *Z. nicaraguensis* (PI 615697) showed higher PWUE values than those of cultivated maize lines. *Z. nicaraguensis* is known to grow in wet habitats such as coastal, estuarine, and river environments (Bird, 2000; Iltis & Benz, 2000). The lower PWUE in *Z. nicaraguensis* may reflect these ecological traits.

## Conclusion

This study investigated the genetic variations in photosynthetic rate and resource use efficiency in maize and teosinte lines and the regulatory factors involving in these variations.  $P_N$  was positively correlated with physiological traits of leaves such as  $g_s$ ,  $T_r$ , and Chl and N contents. However,  $g_s$  may not be the primary  $P_N$  regulator, because maize has a CCM.  $P_N$  was also positively correlated with activities of NADP-ME, PCK, and Rubisco, but not with that of PEPC. These data suggest that biochemical processes from  $C_4$  acid decarboxylation to re-fixation of  $\text{CO}_2$  by Rubisco are involved in the genetic variation in  $P_N$ . On the other hand, structural traits of leaves such as leaf thickness, IVD, and stomatal parameters were not correlated with  $P_N$ . It is suggested that physiological and biochemical traits are involved in the genetic variation of  $P_N$  in maize and teosinte lines but structural traits are not directly involved. In maize and teosinte lines, PNUE was in the highest class and PWUE was in lower class of values previously reported in other  $C_4$  plants. It is worth noting that PNUE was negatively correlated with leaf N content. Some teosinte lines showed higher PWUE and have a value as genetic resources. The photosynthetic traits of maize may be also regulated by other factors not examined in this study, such as electron transport and  $\text{CO}_2$  leakiness from BS cells. Further studies will be required for our better understanding of the genetic variation in photosynthetic traits in maize.

## Acknowledgements

We thank the Plant Introduction Station (Agricultural Research Service, USDA) and the NARO Genebank, Tsukuba, Japan, for their kind gifts of maize and teosinte seeds.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the Japan Society for the Promotion of Science KAKENHI [grant number JP 16H04868], [grant number JP24380010] to O.U.

## References

- Annon, D. I. (1949). Copper enzymes in isolated chloroplast: Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology*, *24*, 1–15. doi:10.1104/pp.24.1.1
- Arrivault, S., Obata, T., Szcówka, M., Mengin, V., Guenther, M., Hoehne, M., ... Stitt, M. (2017). Metabolite pools and carbon flow during C<sub>4</sub> photosynthesis in maize: <sup>13</sup>CO<sub>2</sub> labeling kinetics and cell type fractionation. *Journal of Experimental Botany*, *68*, 283–298. doi:10.1093/jxb/erw414
- Baer, G. R., & Schrader, L. E. (1985). Relationships between CO<sub>2</sub> exchange rates and activities of pyruvate, Pi dikinase and ribulose biphosphate carboxylase, chlorophyll concentration, and cell volume in maize leaves. *Plant Physiology*, *77*, 612–616. doi:10.1104/pp.77.3.612
- Bellasio, C., & Griffiths, H. (2014). The operation of two decarboxylases, transamination, and partitioning of C<sub>4</sub> metabolic processes between mesophyll and bundle sheath cells allows light capture to be balanced for the maize C<sub>4</sub> pathway. *Plant Physiology*, *164*, 466–480. doi:10.1104/pp.113.228221
- Bird, R. M. C. K. (2000). A remarkable new teosinte from Nicaragua: Growth and treatment of progeny. *Maize Genetic Cooperation Newsletter*, *74*, 58–59. Retrieved from <https://www.cabdirect.org/cabdirect/abstract/20001613170>
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, *72*, 248–254. doi:10.1016/0003-2697(76)90527-3
- Brown, R. H. (1977). A difference in N use efficiency in C<sub>3</sub> and C<sub>4</sub> plants and its implications in adaptation and evolution. *Crop Science*, *18*, 93–98. doi:10.2135/cropsci1978.0011183X001800010025x
- Brown, R. H. (1999). Agronomic implications of C<sub>4</sub> photosynthesis. In R. F. Sage & R. K. Monson (Eds.), *C<sub>4</sub> Plant Biology* (pp. 473–507). San Diego, CA: Academic Press.
- Büßis, D., von Groll, U., Fisahn, J., & Altmann, T. (2006). Stomatal aperture can compensate altered stomatal density in *Arabidopsis thaliana* at growth light conditions. *Functional Plant Biology*, *33*, 1037–1043. doi:10.1071/FP06078
- Carpita, N. C., & McCann, M. C. (2008). Maize and sorghum: Genetic resources for bioenergy grasses. *Trends in Plant Science*, *13*, 415–420. doi:10.1016/j.tplants.2008.06.002
- Cernusak, L. A., Ubierna, N., Winter, K., Holtum, J. A. M., Marshall, J. D., & Farquhar, G. D. (2013). Environmental and physiological determinants of carbon isotope discrimination in terrestrial plants. *New Phytologist*, *200*, 950–965. doi:10.1111/nph.12423
- Crosbie, T. M., Mock, J. J., & Pearce, R. B. (1977). Variability and selection advance for photosynthesis in Iowa stiff stalk synthetic maize population. *Crop Science*, *17*, 511–514. doi:10.2135/cropsci1977.0011183X001700040007x
- Dengler, N. G., Dengler, R. E., Donnelly, P. M., & Hattersley, P. W. (1994). Quantitative leaf anatomy of C<sub>3</sub> and C<sub>4</sub> grasses (Poaceae): Bundle sheath and mesophyll surface area relationships. *Annals of Botany*, *73*, 241–255. doi:10.1006/anbo.1994.1029
- Dow, G. L., Bergmann, D. C., & Berry, J. A. (2014). An integrated model of stomatal development and leaf physiology. *New Phytologist*, *201*, 1218–1226. doi:10.1111/nph.12608
- Duncan, W. G., & Hesketh, J. D. (1968). Net photosynthetic rates, relative leaf growth rates, and leaf numbers of 22 races of maize grown at eight temperatures. *Crop Science*, *8*, 670–674. doi:10.2135/cropsci1968.0011183X000800060009x
- Evans, J. R. (2013). Improving photosynthesis. *Plant Physiology*, *162*, 1780–1793. doi:10.1104/pp.113.219006
- Farquhar, G. D., & Richards, R. A. (1984). Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. *Australian Journal of Plant Physiology*, *11*, 539–552. doi:10.1071/PP9840539
- Fernandez, M. G. S., Strand, K., Hamblin, M. T., Westgate, M., Heaton, E., & Kresovich, S. (2015). Genetic analysis and phenotypic characterization of leaf photosynthetic capacity in a sorghum (*Sorghum* spp.) diversity panel. *Genetic Resources and Crop Evolution*, *62*, 939–950. doi:10.1007/s10722-014-0202-6
- Flood, P. J., Harbinson, J., & Aarts, M. G. M. (2011). Natural genetic variation in plant photosynthesis. *Trends in Plant Science*, *16*, 327–335. doi:10.1016/j.tplants.2011.02.005
- Furbank, R. T. (2011). Evolution of the C<sub>4</sub> photosynthetic mechanism: Are there really three C<sub>4</sub> acid decarboxylation types? *Journal of Experimental Botany*, *62*, 3103–3108. doi:10.1093/jxb/err080
- Ghannoum, O., von Caemmerer, S., & Conroy, J. P. (2001). Carbon and water economy of Australian NAD-ME and NADP-ME C<sub>4</sub> grasses. *Australian Journal of Plant Physiology*, *28*, 213–223. doi:10.1071/PP00078
- Ghannoum, O., Evans, J. R., Chow, W. S., Andrews, T. J., Conroy, J. P., & von Caemmerer, S. (2005). Faster rubisco is the key to superior nitrogen-use efficiency in NADP-malic enzyme relatives to NAD-malic enzyme C<sub>4</sub> grasses. *Plant Physiology*, *137*, 638–650. doi:10.1104/pp.104.054759
- Ghannoum, O., Evans, J. R., & von Caemmerer, S. (2011). Nitrogen and water use efficiency of C<sub>4</sub> plants. In A. S. Raghavendra & R. F. Sage (Eds.), *C<sub>4</sub> photosynthesis and related CO<sub>2</sub> concentrating mechanisms* (pp. 129–146). Dordrecht: Springer.
- Hamaoka, N., Uchida, Y., Tomita, M., Kumagai, E., Araki, T., & Ueno, O. (2013). Genetic variations in dry matter production, nitrogen uptake, and nitrogen use efficiency in the AA genome *oryza* species grown under different nitrogen conditions. *Plant Production Science*, *16*, 107–116. doi:10.1626/ppls.16.107
- Hatch, M. D. (1987). C<sub>4</sub> photosynthesis: A unique blend of modified biochemistry, anatomy and ultrastructure. *Biochimica et Biophysica Acta (BBA) – Reviews on Bioenergetics*, *895*, 81–106. doi:10.1016/S0304-4173(87)80009-5
- Heichel, G. H., & Musgrave, R. B. (1969). Varietal difference in net photosynthesis of *Zea mays* L. *Crop Science*, *9*, 483–486. doi:10.2135/cropsci1969.0011183X000900040029x
- Henderson, S., von Caemmerer, S., Farquhar, G. D., Wade, L., & Hammer, G. (1998). Correlation between carbon isotope discrimination and transpiration efficiency in lines of the C<sub>4</sub> species *Sorghum bicolor* in the glasshouse and the field. *Australian Journal of Plant Physiology*, *25*, 111–123. doi:10.1071/PP95033
- Iltis, H. H., & Benz, B. F. (2000). *Zea nicaraguensis* (Poaceae), a new teosinte from pacific coastal Nicaragua. *Novon*, *10*, 382–390. doi:10.2307/3392992
- Kawamitsu, Y., Hiyane, S., Tamashiro, Y., & Hakoyama, S. (2002). Regulation of photosynthesis and water use efficiency in relation to stomatal frequency and interveinal distance in C<sub>3</sub>- and C<sub>4</sub>-Grass Species. *Environmental Control in Biology*, *40*, 365–374. doi:10.2525/ecb1963.40.365
- Kiran, T. V., Rao, Y. V., Subrahmanyam, D., Rani, N. S., Bhadana, V. P., Rao, P. R., & Voleti, S. R. (2013). Variation in leaf photosynthetic characteristics in wild rice species. *Photosynthetica*, *51*, 350–358. doi:10.1007/s11099-013-0032-3



- Koteyeva, N. K., Voznesenskaya, E. V., & Edwards, G. E. (2015). An assessment of the capacity for phosphoenolpyruvate carboxykinase to contribute to C<sub>4</sub> photosynthesis. *Plant Science*, 235, 70–80. doi:10.1016/j.plantsci.2015.03.004
- Lawson, T., & Blatt, M. R. (2014). Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. *Plant Physiology*, 164, 1556–1570. doi:10.1104/pp.114.237107
- Lawson, T., Simkin, A. J., Kelly, G., & Granot, D. (2014). Mesophyll photosynthesis and guard cell metabolism impacts on stomatal behavior. *New Phytologist*, 203, 1064–1081. doi:10.1111/nph.12945
- Leonardos, E. D., & Grodzinski, B. (2000). Photosynthesis, immediate export and carbon partitioning in source leaves of C<sub>3</sub>, C<sub>3</sub>-C<sub>4</sub> intermediate and C<sub>4</sub> *Panicum* and *Flaveria* species at ambient and elevated CO<sub>2</sub> levels. *Plant, Cell and Environment*, 23, 839–851. doi:10.1046/j.1365-3040.2000.00604.x
- McCouch, S. (2004). Diversifying selection in plant breeding. *PLoS Biology*, 2, e347. doi:10.1371/journal.pbio.0020347
- Nose, A., Uehara, M., & Kawamitsu, Y. (1994). Variations in leaf gas exchange traits of saccharum including feral sugarcane, *Saccharum spontaneum* L. *Japanese Journal of Crop Science*, 63, 489–495. doi:10.1626/jcs.63.489
- O'Leary, M. H. (1988). Carbon isotopes in photosynthesis. *BioScience*, 38, 328–336. doi: 10.2307/1310735
- Osmond, C. B., Winter, K., & Ziegler, H. (1982). Functional significance of different pathways of CO<sub>2</sub> fixation in photosynthesis. In O. L. Lange, P. S. Nobel, & C. B. Osmond (Eds.), *Encyclopedia of plant physiology, New series*, Vol. 12B (pp. 479–547). Berlin: Springer Verlag.
- Pearce, R. B., Carlson, D. K., Barnes, D. K., Hart, R. H., & Hanson, C. H. (1969). Specific leaf weight and photosynthesis in alfalfa. *Crop Science*, 9, 423–426. doi:10.2135/cropsci1969.0011183X000900040010
- Pearcy, R. W., & Ehleringer, J. (1984). Comparative ecophysiology of C<sub>3</sub> and C<sub>4</sub> plants. *Plant, Cell and Environment*, 7, 1–13. doi:10.1111/j.1365-3040.1984.tb01194.x
- Pengelly, J. J. L., Tan, J., Furbank, R. T., & von Caemmerer, S. (2012). Antisense reduction of NADP-malic enzyme in *Flaveria bidentis* reduces flow of CO<sub>2</sub> through the C<sub>4</sub> cycle. *Plant Physiology*, 160, 1070–1080. doi:10.1104/pp.112.203240
- Sage, R. F. & Pearcy, R. W. (1987). The nitrogen use efficiency of C<sub>3</sub> and C<sub>4</sub> plants. II. Leaf nitrogen effects on the gas exchange characteristics of *Chenopodium album* (L.) and *Amaranthus retroflexus* (L.). *Plant Physiology*, 84, 959–963. doi:10.1104/pp.84.3.959
- Sato, R., & Suzuki, Y. (2010). Carbon and nitrogen stable isotope analysis by EA/IRMS. *Researches in Organic Geochemistry*, 26, 21–29. doi:10.20612/rog.26.0\_21
- Scafaro, A. P., Haynes, P. A., & Atwell, B. J. (2010). Physiological and molecular changes in *Oryza meridionalis* Ng., a heat-tolerant species of wild rice. *Journal of Experimental Botany*, 61, 191–202. doi:10.1093/jxb/erp294
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 Years of image analysis. *Nature Methods*, 9, 671–675. doi:10.1038/nmeth.2089
- Sharwood, R. E., Sonawane, B. V., & Ghannoum, O. (2014). Photosynthetic flexibility in maize exposed to salinity and shade. *Journal of Experimental Botany*, 65, 3715–3724. doi:10.1093/jxb/eru130
- Suresh, K., Rao, K. L. N., & Nair, T. V. R. (1997). Genetic variability in photosynthetic rate and leaf characters in *Brassicaceae* coenospecies. *Photosynthetica*, 33, 173–178. doi:10.1023/A:1022147910426
- Taylor, S. H., Hulme, S. P., Rees, M., Ripley, B. S., Woodward, F. I., & Osborne, C. P. (2010). Ecophysiological traits in C<sub>3</sub> and C<sub>4</sub> grasses: A phylogenetically controlled screening experiment. *New Phytologist*, 185, 780–791. doi:10.1111/j.1469-8137.2009.03102.x
- Togawa, Y., & Ueno, O. (2015). Comparison of resource use efficiencies involved in photosynthesis among biochemical subtypes of C<sub>4</sub> grasses. Abstracts of 239th Meeting of the Crop Science Society of Japan, Tokyo. (p. 209).
- Tsutsumi, N., Tohya, M., Nakashima, T., & Ueno, O. (2017). Variations in structural, biochemical, and physiological traits of photosynthesis and resource use efficiency in *Amaranthus* species (NAD-ME-type C<sub>4</sub>). *Plant Production Science*, 20, 300–312. doi:10.1080/1343943X.2017.1320948
- Ueno, O., & Sentoku, N. (2006). Comparison of leaf structure and photosynthetic characteristics of C<sub>3</sub> and C<sub>4</sub> *Alloterospis semialata* subspecies. *Plant, Cell and Environment*, 29, 257–268. doi:10.1111/j.1365-3040.2005.01418.x
- Ueno, O., Kawano, Y., Wakayama, M., & Takeda, T. (2006). Leaf vascular systems in C<sub>3</sub> and C<sub>4</sub> grasses: A two-dimensional analysis. *Annals of Botany*, 97, 611–621. doi:10.1093/aob/mcl010
- Usuda, H. (1984). Variations in the photosynthetic rate and activity of photosynthetic enzymes in maize leaf tissue of different ages. *Plant and Cell Physiology*, 25, 1297–1301. doi:10.1093/oxfordjournals.pcp.a076838
- Usuda, H., Ku, M. S. B., & Edwards, G. E. (1985). Influence of light intensity during growth on photosynthesis and activity of several key photosynthetic enzymes in a C<sub>4</sub> plant (*Zea mays*). *Physiologia Plantarum*, 63, 65–70. doi:10.1111/j.1399-3054.1985.tb02819.x
- Vogan, P. J., & Sage, R. F. (2011). Water-use efficiency and nitrogen-use efficiency of C<sub>3</sub>-C<sub>4</sub> intermediate species of *Flaveria* Juss (Asteraceae). *Plant, Cell and Environment*, 34, 1415–1430. doi:10.1111/j.1365-3040.2011.02340.x
- von Caemmerer, S., & Furbank, R. T. (2016). Strategies for improving C<sub>4</sub> photosynthesis. *Current Opinion in Plant Biology*, 31, 125–134. doi:10.1016/j.pbi.2016.04.003
- von Caemmerer, S., Millgate, A., Farquhar, G. D., & Furbank, R. T. (1997). Reduction of ribulose-1,5-bisphosphate carboxylase/oxygenase by antisense RNA in the C<sub>4</sub> plant *Flaveria bidentis* leads to reduced assimilation rates and increased carbon isotope discrimination. *Plant Physiology*, 113, 469–477. doi:10.1104/pp.113.2.469
- Walker, R. P., Acheson, R. M., Técsi, L. I., & Leegood, R. C. (1997). phosphoenolpyruvate carboxykinase in C<sub>4</sub> plants: Its role and regulation. *Australian Journal of Plant Physiology*, 24, 459–468. doi:10.1071/PP97007
- Wingler, A., Walker, R. P., Chen, Z. H., & Leegood, R. C. (1999). phosphoenolpyruvate carboxykinase is involved in the decarboxylation of aspartate in the bundle sheath of maize. *Plant Physiology*, 120, 539–545. doi:10.1104/pp.120.2.539
- Wong, S. C., Cowan, I. R., & Farquhar, G. D. (1985). Leaf conductance in relation to rate of CO<sub>2</sub> assimilation. 1. Influence of nitrogen nutrition, phosphorus nutrition, photon flux density, and ambient partial pressure of CO<sub>2</sub> during ontogeny. *Plant Physiology*, 78, 821–825. doi:10.1104/pp.78.4.830
- Zhu, X. G., Long, S. P., & Ort, D. R. (2010). Improving photosynthetic efficiency for greater yield. *Annual Review of Plant Biology*, 61, 235–261. doi:10.1146/annurev-arplant-042809-112206