



SCHOOL of
GRADUATE STUDIES
EAST TENNESSEE STATE UNIVERSITY

East Tennessee State University
Digital Commons @ East
Tennessee State University

Electronic Theses and Dissertations

Student Works

5-2018

The Effect of Alcohol Consumption on Adipokine Secretion

Ashley DeGroat

East Tennessee State University

Follow this and additional works at: <https://dc.etsu.edu/etd>



Part of the [Cellular and Molecular Physiology Commons](#), and the [Endocrinology Commons](#)

Recommended Citation

DeGroat, Ashley, "The Effect of Alcohol Consumption on Adipokine Secretion" (2018). *Electronic Theses and Dissertations*. Paper 3425.
<https://dc.etsu.edu/etd/3425>

This Thesis - Open Access is brought to you for free and open access by the Student Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

The Effect of Alcohol Consumption on Adipokine Secretion

A thesis

presented to

the faculty of the Department of Biological Sciences

East Tennessee State University

In partial fulfillment

of the requirements for the degree

Master of Science in Biology

Ashley R. DeGroat

May 2018

Dr. Jonathan Peterson, Chair

Dr. W. Andrew Clark

Dr. Yongke Lu

Keywords: Alcoholic Fatty Liver Disease, Adipokines, Adipose Tissue

ABSTRACT

The Effect of Alcohol Consumption on Adipokine Secretion

by

Ashley R. DeGroat

Alcoholic Fatty Liver Disease (AFLD) is caused by excessive alcohol consumption and is a leading cause of liver related mortalities, with currently no treatments available. The goal of this project was to establish the effect of alcohol consumption on adipose tissue-derived secreted factors, adiponectin and C1q TNF Related Proteins 1-3 (CTRP1-3). We propose that excessive alcohol consumption will reduce circulating levels of adiponectin and CTRPs 1-3. Mice were fed a Lieber-DeCarli control or alcohol diet for 10-days with a gavage (NIAAA model) or 6-weeks with no gavage (chronic model). Serum and adipose tissue were collected and CTRPs 1-3 and adiponectin levels were examined by immunoblot analysis. Our results indicate that long-term alcohol consumption effects adipokine secretion in a sex specific manner. Further research will be needed to explore the physiological relevance of these findings, to determine if these changes are beneficial to combat the negative effects of excessive alcohol consumption.

ACKNOWLEDGEMENTS

This research was supported in part by the National Institute on Alcohol Abuse and Alcoholism of the National Institutes of Health under Award Number R03AA023612 and East Tennessee State University Research Development Committee (E82262). I thank East Tennessee State University for the opportunity to continue my education and complete my Master's degree. I would like to thank my mentor Dr. Peterson for the opportunity to participate in his research, his guidance, and support during the completion of my research project and thesis. I would also like to thank my committee members Dr. Clark and Dr. Lu for their support and guidance while completing my thesis. I owe a huge thanks to everyone who works in our lab for help and assistance. Lastly, I give all of my praise and thanks to God for His guidance in leading me to East Tennessee State University, giving me this opportunity, and for being my hope and strength, for without Him this would not have been possible.

TABLE OF CONTENTS

	Page
ABSTRACT.....	2
ACKNOWLEDGEMENTS.....	3
LIST OF TABLES.....	6
LIST OF FIGURES.....	7
Chapter	
1. LITERATURE REVIEW.....	8
Introduction.....	8
Adipokines.....	10
CTRP1.....	10
CTRP2.....	11
CTRP3.....	11
Adiponectin.....	12
Leptin.....	13
Cytokines.....	13
PAI-1.....	13
IL-6.....	14
TNF- α	14
2. THE EFFECT OF ALCOHOL CONSUMPTION ON THE CIRCULATING LEVELS OF NOVEL ADIPOKINES: CTRP1, CTRP2, AND CTRP3.....	15
Abstract.....	16
Introduction.....	17
Methods.....	18

Results.....	21
Discussion.....	28
Conclusion.....	30
Figure Legends.....	31
3. METHODS.....	40
Adipose Tissue Lipid Analysis.....	40
Fatty Acid Methyl Ester Preparation and Analysis.....	40
Serum Evaluations.....	41
4. RESULTS.....	42
Effects of Ethanol on Circulating Glucose.....	42
Effects of Ethanol on Circulating Triglycerides.....	43
Effects of Ethanol on Serum Transaminases.....	44
Chronic Adipose Tissue Lipid Analysis.....	45
5. DISCUSSION.....	46
Leptin and Adiponectin.....	47
CTRP1.....	48
CTRP2.....	49
CTRP3.....	49
6. CONCLUSION.....	50
REFERENCES.....	51
VITA.....	61

LIST OF TABLES

Table	Page
1.1. Adipokines and Cytokines.....	9
2.1. Number of Animals in Each Experimental Group.....	19
2.2. PCR Primer Sequences.....	20
2.3. Serum Multiplex Analysis.....	23

LIST OF FIGURES

Figure	Page
2.1. Animal Characteristics.....	22
2.2. Circulating Adipokines in the NIAAA Model of Ethanol Feeding.....	24
2.3. Circulating Adipokines in the Chronic Model of Ethanol Feeding.....	25
2.4. Tissue Adipokine Gene Expression.....	26
2.5. Tissue Adipokine Protein Content.....	27
4.1. Chronic Glucose Levels.....	43
4.2. Chronic Triglycerides.....	43
4.3. NIAAA and Chronic Serum Transaminases.....	44
4.4. Chronic Adipose Tissue Lipid Analysis.....	45

CHAPTER 1

LITERATURE REVIEW

Alcoholic Fatty Liver Disease (AFLD) is one of the leading causes of mortality in the United States and around the world (Bataller and Gao 2011; Parry 2011; Kema et al. 2015). Fatty liver is defined as the accumulation of excess lipids in the liver. According to the NIAAA (National Institute on Alcohol Abuse and Alcoholism), a standard drink contains about 14 grams of pure alcohol. The NIAAA defines binge drinking as having a blood alcohol concentration (BAC) level of 0.08 g/dL, which typically occurs after 4 drinks for women and 5 drinks for men at a time. Heavy drinking is defined by the NIAAA as binge drinking for 5 or more days within a month. Chronic alcohol consumption disrupts lipid synthesis and can lead to hepatic steatosis, hepatitis, and cirrhosis of the liver (Nagy et al. 2016). Hepatic steatosis is primarily asymptomatic, but the accumulation of lipids can serve as the beginning of more serious forms of fatty liver disease such as: hepatitis (fatty liver with inflammation), cirrhosis (hepatitis with fibrosis), and liver failure. Not only does this disease cause damage to the liver but it also affects the body as a whole. And a specific area of interest is the adipose tissue.

Adipose tissue has been found to be more than just an organ for storing fat, as it also plays a role in whole-body metabolism and is responsible for synthesis and secretion of many hormones (Ahima and Flier 2000; Coelho et al. 2013). The balance between lipogenesis (fat synthesis) and lipolysis (breakdown of fat) determines fat accumulation. Lipogenesis occurs in the adipose tissue as well as the liver. It is stimulated by an increase in calories and inhibited by fasting. Lipolysis occurs in adipose tissue and breaks down fat into fatty acids for energy production. Once broken down fatty acids are

transported from the adipose tissue through the bloodstream to the liver, muscle, and other tissues for oxidation (Coelho et al. 2013). Alcohol consumption affects adipose tissue mass, adipokine secretion, adipose tissue hydrolysis, and results in the release of excess fatty acids that are transported to the liver and deposited as triglycerides (Kema et al. 2015). Changes to the adipose tissue have an opportunity to affect the levels of adipokines, cell-signaling proteins secreted from the adipose tissue. Adipokines have an effect on insulin sensitivity, glucose and fatty acid metabolism, and the inflammatory process (Peterson et al. 2013). There are many different adipokines secreted from adipose tissue but this project aims to look at a select few because of their roles in lipid metabolism and fatty acid oxidation. C1q/TNF-related proteins (CTRPs) are highly conserved paralogs of adiponectin consisting of a signal peptide, a short variable region, a collagen domain, and a globular C1q (Complement Component 1q) domain (Wong et al. 2008). Because the adiponectin globular domain closely resembles TNF α , proteins with the C1q domain are classified as the C1q/TNF protein family (Wong et al. 2008).

Table 1.1 Adipokines and Cytokines

Molecule	Function/effect	Molecular Weight Observed
CTRP1	Metabolic and cardiovascular functions, lowers blood glucose levels and protects from diet-induced obesity and insulin resistance. Promotes glucose uptake and fatty acid oxidation in skeletal muscle.	35 kDa
CTRP2	Promotes lipid uptake from the blood	38kDa
CTRP3	Stimulates liver lipid metabolism and attenuates diet-induced fatty liver disease.	25kDa
Adiponectin	Increases fatty acid oxidation in the liver and stimulates glucose uptake in skeletal muscle.	34kDa

Table 1.1 (continued)

Leptin	Signals to the brain about body fat stores. Regulation of appetite and energy expenditure. Stimulates fatty acid oxidation in the liver and skeletal muscle. Stimulates glucose uptake in skeletal muscle. Prevents lipotoxicity Pro-fibrogenic
PAI-1	Inhibitor of the fibrinolytic system by inhibition of activation of plasminogen
IL-6	Pro-inflammatory, lipid and glucose metabolism, regulation of body weight Induces insulin and leptin resistance
TNF-α	Pro-inflammatory cytokine that induces PAI-1, increases fatty acid release from adipocytes, and increases lipogenesis in hepatocytes.

Adipokines

CTRP1

C1q/TNF-related protein 1 (CTRP1) is expressed at its highest levels in adipose tissue (Rodriguez et al. 2016). CTRP1 has been found to lower blood glucose and activate AMPK (AMP-activated protein kinase) to control fatty acid metabolism in skeletal muscle (Peterson et al. 2012; Rodriguez et al. 2016). And chronic over expression of CTRP1 has been found to enhance skeletal muscle fat oxidation and reduce insulin resistance caused by a high-fat diet (Rodriguez et al. 2016). CTRP1 has also reduces the formation of plaque and increases aldosterone production (Shabani et al. 2016). In a CTRP1 KO model, the loss of CTRP1 in mice fed a high fat diet resulted in decreased expression of multiple genes associated with lipid metabolism in the adipose tissue (Rodrgiuez et al. 2016).

CTRP2

C1q/TNF-related protein 2 (CTRP2) is a mouse paralog closely related to adiponectin, which promotes glycogen accumulation and activates the AMPK signaling pathway to increase fatty acid oxidation. Because CTRP2 induces glycogen accumulation it may also lower blood glucose (Wong et al. 2008). And because of the role of adiponectin in enhancing insulin resistance, CTRP2 could also play a role in improving insulin resistance.

CTRP3

CTRP3 (C1q/TNF-related protein 3) increases liver lipid metabolism and inhibits inflammation. Previous work has shown that overexpressing CTRP3 as well as daily injections of CTRP3 reduces high-fat diet induced fatty liver (Peterson et al. 2013). Over expressing CTRP3 in a high fat model showed a decrease in the synthesis of triglycerides and a decrease in circulating levels of TNF- α (Peterson et al. 2013). CTRP3 decreases blood glucose by suppressing gluconeogenic expression in the liver (Peterson et al. 2010). It has been found that human patients with non-alcoholic fatty liver disease (NAFLD) exhibited reduced levels of circulating CTRP3 (Zhang et al. 2017). Therefore, restoring CTRP3 levels has been shown as a possible treatment for NAFLD (Peterson et al. 2013). But it is still a question if it could be used to alleviate AFLD, because AFLD is caused by ethanol-induced lipogenesis and decreased lipid oxidation and NAFLD is primarily caused by an excessive accumulation of lipids in the liver (Breitkopf 2009; Coelho et al. 2013; Fujii 2014; Parker 2018). Identifying the affect of alcohol on levels of CTRP3 will provide further insight to this question.

Adiponectin

Adiponectin was one of the first adipokines to be discovered and is well studied. Adiponectin is an adipokine involved in regulating glucose levels as well as the breakdown of fatty acids (Karbowska and Kochan 2006). Adiponectin is exclusively secreted by adipose tissue and abundantly present in the blood stream. Adiponectin stimulates insulin secretion, fatty acid oxidation in the liver, glucose uptake in skeletal muscle, and suppresses TNF- α and IL-6 expression, all factors that are disrupted with ETOH feeding. Chronic ETOH consumption causes a significant decrease in circulating levels of adiponectin, and correlated with the development of liver injury (Xu et al. 2003; Song et al. 2008; Tan et al. 2012). Adiponectin is believed to also play a protective role against alcoholic liver disease in mice as levels increase significantly with consumption of a high fat diet with ethanol (You et al. 2005). Adiponectin is an anti-inflammatory adipokine known to promote appropriate lipid storage, preventing ectopic fat storage in places such as the liver (Lang and Steiner 2017). Circulating levels of adiponectin are shown to be affected by alcohol consumption, although, there is some variability as adiponectin levels have been shown to be unaffected (Tan et al. 2012), suppressed (Chen et al. 2007; Yu et al. 2010), or increased (Sierksma et al. 2004; Pravdova et al. 2009; Mandrekar and Fulham 2016) with the consumption of alcohol. It is suspected that oxidative stress induced by acute alcohol exposure reduces the secretion of adiponectin (Tang et al. 2003), indicating that time since last dose of ethanol can affect results.

Leptin

Leptin is one of the most studied adipokine (Hiney et al. 1999; Roth et al. 2003; Strbák et al. 2003; Lang and Steiner 2017). It plays a role in food intake, energy expenditure, lipolysis, fatty acid oxidation, and lipogenesis (Lang and Steiner 2017).

Leptin also prevents lipotoxicity, which causes cell damage due to the accumulation of lipids in areas other than adipose tissue, such as the liver (Bertolani and Morra 2008). Leptin has been shown to be pro-fibrogenic, as an absence of leptin resulted in reduced liver fibrosis (Leclercq et al. 2002). The effects of alcohol on leptin levels vary among studies; it has been shown to increase (Kiefer et al. 2002; Obradovic 2002; He et al. 2015), decrease (Hiney et al. 1999), and be unchanged (Strbák et al. 2003) using a range of chronic alcoholic models (Lang and Steiner 2017). Circulating levels of leptin have been shown to be affected by alcohol consumption, although, similarly to adiponectin, there is variability. Some models have shown leptin levels to increase (He et al. 2003; Roth et al. 2003; Sierksma et al. 2004; Pravdova et al. 2009) with alcohol consumption while others have shown leptin levels to decrease (Hiney et al. 1999; Tan et al. 2012) with alcohol consumption.

Cytokines

PAI-1

Plasminogen Activator Inhibitor-1 (PAI-1) inhibits plasminogen activation that breaks down fibrin and is regulated by levels of TNF- α (Hou et al. 2004). As alcoholic fatty liver disease progresses, there is an accumulation of extracellular matrices that leads to fibrosis. Plasma activator inhibitor-1 (PAI-1) regulates fibrinolysis. (Arteel 2008).

IL-6

Interleukin 6 (IL-6) acts as a pro inflammatory cytokine. Studies have shown IL-6 to play an important part in protection of the liver through liver repair and preventing apoptosis (Hong et al. 2002).

TNF- α

TNF- α is a pro-inflammatory cytokine and increased levels have been documented in animal models of AFLD (McClain et al. 1998). TNF- α contributes to hepatic steatosis by inducing the expression PAI-1. It increases fatty acid release from adipocytes increasing lipogenesis in hepatocytes, and inhibits the β -oxidation of fatty acids (Arteel 2008).

Because of adipokines' positive affects on lipid oxidation, alcohol consumption will result in decreased circulating levels of adiponectin, CTRP1, CTRP2, and CTRP3.

CHAPTER 2

THE EFFECT OF ALCOHOL CONSUMPTION ON THE CIRCULATING LEVELS OF THE NOVEL ADIPOKINES: CTRP1, CTRP2, AND CTRP3

Ashley R. DeGroat¹, Christina K Fleming², Samantha M Dunlay², Kendra L. Hagood¹, Jonathan M. Peterson^{3,4}.

1: Graduate student, Department of Biomedical Sciences, East Tennessee State University, Johnson City, Tennessee

2. Work completed as an undergraduate student, East Tennessee State University, Johnson City, Tennessee

3: College of Public Health, Department of Health Sciences, East Tennessee State University, Johnson City, Tennessee

4: Quillen College of Medicine, Department of Biomedical Sciences, East Tennessee State University, Johnson City, Tennessee

*Corresponding author

Email: peteronjm1@etsu.edu (JMP)

ABSTRACT:

The goal of this project was to establish the effect of alcohol consumption on a group of novel adipose tissue-derived secreted factors: C1q TNF Related Proteins 1-3 (CTRP1, CTRP2, and CTRP3). Adipose tissue secretes several circulating proteins, called adipokines, which exert a multitude of biological effects important for human health. However, adipose tissue is extremely sensitive to alcohol consumption, leading not only to disrupted fat storage, but also to disruptions in adipokine production. Changes to adipokines could have widespread biological effects and potentially contribute to alcohol-induced ailments. To test the effects of alcohol consumption on adipokines, male and female mice were randomized to a Lieber-DeCarli control diet or Lieber-DeCarli 5% (v/v) ethanol diet for either: 1) 10-days followed by a single gavage of 5 g/kg ethanol on the 11th day (the NIAAA model); or 2) 6-weeks with no binge added (chronic model). In response to the NIAAA model, female mice fed ethanol had an ~200% increase in circulating levels of adiponectin, an ~25% increase in CTRP1, a 25% decrease in CTRP2 and 75% reduction in CTRP3. Whereas, in the male mice ethanol decreased circulating CTRP2 by ~25%, with no changes observed in adiponectin, CTRP1, or CTRP3. The effects of ethanol on CTRP2 levels disappeared after 6-weeks of ethanol feeding (chronic model), as CTRP2 levels were not different between control and ethanol fed mice of either sex. Whereas, in the chronic model, ethanol more than doubled circulating adiponectin levels in both male and female mice. Surprisingly, chronic ethanol feeding resulted in a dimorphic effect on circulating CTRP1. Briefly, CTRP1 levels were increased by ~125% in ethanol fed female mice but were 50% lower in ethanol fed male mice. Lastly, circulating CTRP3 levels were decreased in female mice, with no change in male mice. Combined, this is the first study to document the effects of alcohol on the circulating levels of CTRP1, CTRP2, and CTRP3. Understanding the impact of excessive alcohol consumption on adipokine production and secretion could identify novel alcohol-induced mechanisms of human disease and identify novel potential pharmaceutical targets for treatment development. Lastly, these results confirm earlier findings that alcohol consumption has sex-specific effects.

INTRODUCTION

The detrimental effects of chronic alcohol abuse have been well-documented with established long-term health conditions such as cardiovascular disease [1], respiratory distress [2], gastrointestinal dysfunction, alcoholic liver disease [3-6], cancer [7], and metabolic dysfunction [8]. Excessive alcohol consumption not only causes initial injury via direct toxic effects (i.e. oxidative stress) to the individual tissues, but also results in secondary indirect injury through elevations in inflammatory cytokines and ectopic fat deposition [9, 10]. Although little can be done to prevent the acute toxic effects of alcohol consumption, understanding and reducing secondary alcohol-induced injury is a key component of treating the long-term health conditions associated with chronic alcohol consumption.

Adipose tissue is not only the primary location for the storage of excess lipids, but is also a major contributor to the production of circulating inflammatory cytokines [11]. Furthermore, chronic alcohol consumption results in high levels of adipose-tissue oxidative stress, leading to elevations in inflammation and hyperlipolysis [11]. Therefore, alcohol-induced disruptions to adipose tissue function contribute to the wide-spread development of secondary alcohol-related health conditions. Females have a higher amount of adipose tissue (higher percent body fat), and this may account for the increased susceptibility of females to the chronic effects of alcohol [5, 7, 12-14].

In addition to the direct effects of alcohol on adipose tissue, alcohol consumption can also lead to disruptions in adipokine secretion. Adipokines are bioactive proteins secreted by adipose tissue which can have significant endocrine effects on regulating human health and disease. Circulating levels of the two most widely studied adipokines, leptin and adiponectin, are significantly affected by alcohol consumption [10, 15-29]. Leptin is primarily a satiety signal; therefore, its role in ethanol-related health effects is unclear. However, adiponectin stimulates fatty acid oxidation and inhibits both the activity and production of inflammatory cytokines [10, 30, 31], thus supporting our hypothesis that alcohol-induced disruptions to adipokine production can contribute to the development alcoholic related disease.

In 2004 Wong et al identified a novel family of adipokines, referred to as Complement C1q Tumor Necrosis Factor-Related Proteins (CTRPs) [32, 33]. Reflecting profound biological potency, the initial characterization of these adipose tissue-derived CTRP factors demonstrate wide-ranging effects upon metabolism, inflammation, and survival-signaling [32-52]. As alcohol alters the expression of both leptin and adiponectin, we hypothesized that chronic alcohol abuse would also affect other adipokines. To test this hypothesis, we chose to examine three novel adipokines that have not been examined in relation to alcohol consumption: CTRP1, CTRP2, and CTRP3. All three of these proteins have been demonstrated to improve lipid handling in unique ways in response to high fat diet. Briefly, elevated circulating CTRP1 levels attenuate body weight gain in response to a high fat feeding through increased lipid oxidation in skeletal muscle [36], CTRP2 improves serum lipid clearance in obese mice [37, 46], and CTRP3 attenuates high fat diet-induced hepatic steatosis [38, 53]. In summary, all three of these adipokines have documented, unique methods to

improving lipid handling in ways that could diminish the secondary damage caused by chronic alcohol abuse. Therefore, the purpose of this project was to establish the effects of alcohol on the circulating levels of these proteins. The results of this study could produce new understanding to the mechanism by which chronic alcohol consumption leads to ectopic fat deposition and other health issues and identify novel potential pharmaceutical targets for treatment of alcohol-induced disease.

METHODS

Animal model

Forty female mice (C57BL/6) and thirty-seven male mice (C57BL/6) were used for this study. Mice were housed in polycarbonate cages on a 12-h light-dark photocycle with *ad libitum* access to water and food, except as specified. At the time points indicated, animals were anesthetized with isoflurane and euthanized via cardiac puncture. Serum samples were prepared according to manufacture's instructions (Sarstedt, Cat#41.1500.005). The gonadal fat pads were excised, snap frozen in liquid nitrogen, and stored at -80°C until further analysis. All animal procedures were conducted in accordance with institutional guidelines, and ethical approval was obtained from the University Committee on Animal Care (protocol #P151201; East Tennessee State University, Animal Welfare Assurance number is A3203-01). Animals were checked/weighed daily and euthanized (counted as dead), via CO_2 inhalation, based on the presence of any of the following criteria for humane endpoints: unconsciousness, intractable seizures, labored breathing or respiratory distress, inability to ambulate or maintain upright position, diarrhea or constipation, or the inability to eat or drink.

Ethanol Feeding

Two independent ethanol feeding models were employed: The first model was the 11-day chronic plus binge model, also known as the NIAAA model [54]. This model reportedly mimics hepatic steatosis and liver injury, which occurs in many alcoholic hepatitis patients (26). Briefly, 12-week old mice were acclimatized to a control liquid diet (Bio-serv; cat# F1259SP) for 4 days followed by 10 days on the Lieber-DeCarli ethanol diet (5% v/v ethanol; Bio-serv; cat# F1258SP) *ad libitum*. On the morning of the 11th day (1 hour into light cycle) food was removed and replaced with water and the mice were given a single gavage of ethanol (5 g kg^{-1}). After gavage, cages were placed on heating pads to prevent hypothermia, as described [54]. Nine hours post gavage, mice were anesthetized with isoflurane, until the absence of reflex was observed, and the euthanized by exsanguination, and tissue/serum samples were collected and processed for analysis.

In the second model (chronic model), 8-week old male and 12-week old female mice were acclimatized to a liquid diet *ad libitum*, without the addition of alcohol for 1-

week and then gradually transitioned from 1-5% Lieber-DeCarli ethanol diet (v/v ethanol) over the course of the next 2 weeks, then maintained on 5% ethanol (v/v ethanol) for the remaining 4 weeks. This feeding protocol is believed to reflect chronic ethanol abuse, beginning with low volumes and increasing over time [55]. On the morning of the final day, food was removed, and mice were fasted 9 hours, anesthetized with isoflurane until the absence of reflex was observed, and the euthanized by exsanguination, and tissue/serum samples were collected and processed for analysis. The 9-hour time point was selected to be consistent with the NIAAA model protocol [54].

Control Fed mice were placed on an ethanol free isocaloric control diet (Bio-serv; cat# F1259SP) supplemented with maltose dextrin (to match the calories of ethanol), for use as experimental controls. Food intake in ethanol fed mice was measured daily and mice on the control diet had their food intake limited to match the daily intake for the previous day of the corresponding ETOH-fed mice. Overview of animals in each experimental group are listed in table 1.

Table 2.1 Number of animals in each experimental group

Model	Male		Female	
	Control	ETOH	Control	ETOH
NIAAA model	10	8	6	6
Chronic Model	6	13	8	20

ETOH, ethanol fed; Control, fed matched control diet without ethanol

Immunoblot Analysis

Serum samples were diluted, and adipose tissues (gonadal fat pad) were homogenized in assay buffer (50 mM Tris HCl, pH 8.0, 150 mM NaCl, 0.1% Triton x-100, 0.5% sodium deoxycholate, 0.1% SDS), plus the addition of protease and phosphatase inhibitors (Bimake Cat#B14001 & B15001). Protein concentrations were measured by commercial assay (Pierce™; Cat#PI23236). Afterward, equal proportions of each sample were denatured at 95°C in SDS loading buffer (final concentration: 1% SDS, 5% 2-mercaptoethanol, 10% glycerol, 0.004% bromophenol blue, 0.125 M Tris HCl, pH 6.8). 30 µg per sample (adipose) or 1 µl serum was separated by gel electrophoresis (BioRad; cat#456-1046) and transferred to a nitrocellulose membrane (BioRad; Cat#162-0115), according the manufacturer's instructions. To confirm appropriate protein migration a protein standard was loaded with each blot (BioRad cat#1610374 or Thermo Scientific cat#26616). Membranes were blocked in 2% non-fat milk and probed with primary antibodies: CTRP1 (GW Wong Lab; Johns Hopkins University, Cat# anti-gCTRP1, RRID:AB_2716247), CTRP2 (Abnova Corporation Cat# H00114898-M01, RRID:AB_426121), CTRP3 (R and D Systems Cat# AF2436, RRID:AB_2067713), and Adiponectin (R and D Systems Cat# MAB1119, RRID:AB_2305045). After incubation with primary antibodies membranes were washed and probed with appropriate HRP-labeled secondary antibodies: Goat anti-rabbit (Thermo Fisher Scientific Cat#

31460 RRID:AB_228341), rabbit anti-goat (Thermo Fisher Scientific Cat# 31402, RRID:AB_228395), rabbit anti-rat (Thermo Fisher Scientific Cat# PA1-28786, RRID:AB_10983740), or goat anti-mouse (Cell Signaling Technology Cat# 7076, RRID:AB_330924). Chemiluminescence signals were visualized with Millipore (Cat# 17010A2). Quantification of signal intensity was performed using Alphaview Software (Alpha Innotech).

RNA Isolation

RNA was isolated according to commercial assay following manufacturer's instructions (Direct-zol Cat# R2070). Isolated RNA was eluted in 50 µl RNase-free water; purity (RIN ≥ 7.0) and concentrations were confirmed by microfluidic capillary electrophoresis (Agilent RNA 6000 Nano kit, #5067-1511, Agilent Technologies). 1 µg RNA was reverse transcribed according to manufacturer's instructions (Promega, Cat#A5001).

Quantitative real-time PCR

A 10-fold dilution series of DNA amplicons generated from a prepared sample was employed as a standard curve for each gene of interest, and the qPCR efficiency was determined for each gene (Bio-Rad Cfx thermocycler). All qRT-PCR primers displayed a coefficient of correlation greater than 0.95 and efficiencies between 90% and 110%. Primer sequences are listed in table 2. Briefly, 25 ng of cDNA was incubated in SYBR Green qPCR Master mix (Bimake.com, Cat# B21203) for an initial denaturation at 95 °C for 10 min, followed by 40 PCR cycles each consisting of 95 °C for 15 s, and 60 °C for 1 min. After the last cycle specificity of amplification products were confirmed by analyzing melting curve profiles for primers and products. Data is reported as copy number normalized to the geometric mean of the reference genes Beta-actin (Actb) and Hypoxanthine-guanine phosphoribosyltransferase (Hprt1).

Table 2.2 PCR primer sequences

Gene Name	Forward	Reverse
Actb	CCTCCCTGGAGAAGAGCTATG	TTACGGATGTCAACGTCACAC
Hprt1	CAAACCTTGCTTCCCTGGT	TCTGGCCTGTATCCAACACTTC
Adipoq	CCTGGCCACTTTCTCCTCATT	ATCCTGAGCCCTTTTGGTGT
CTRP1	TCCGAGCTCTGTTGACATGC	AAAGATTGACCAGCCCCTGG
CTRP2	TCCTGGGTACTCTTGGCCTG	AAGCATTGGGTCAGCAGCA
CTRP3	CATCTGGTGGCACCTGCTG	TGACACAGGCAAATGGGAG

Abbreviations: Actb, Beta-actin; Hprt1, Hypoxanthine-guanine phosphoribosyltransferase; Adipoq, Adiponectin; CTRP, C1q and tumor necrosis factor related protein

Multiplex Serum Analysis

Circulating concentrations of Interleukin-6 (IL-6), Tumor necrosis factor alpha (TNF), Plasminogen activator inhibitor-1 (PAI-1) and leptin were determined using commercially available assays (Bio-Plex® Multiplex Immunoassay System, Bio-rad Cat# 171F7001M, 171G5023M, 171I50001, 171G5007M).

Statistical Analysis

Descriptive statistics (mean and standard deviations) were calculated for all measured variables. As the feeding models were not performed concurrently, each model was analyzed independently. Body weight and food intake data were analyzed by two-way repeated measures ANOVA followed by Tukey's multiple comparisons test. Survival curve was determined by log-rank (Mantel-Cox) test. An unpaired t test was used to compare immunoblot and gene expression data between control and ethanol fed groups. All statistical analysis was performed by Graphpad Prism 6.

RESULTS

Animal Characteristics

As expected, with pair feeding no differences in total food intake occurred between control and ethanol fed groups. Although during weeks 2 and 3 the body weights of the ethanol fed female mice were significantly higher than control fed, this difference normalized by week 4 and there was no further difference in body weights between the control and ethanol fed groups of either sex (Fig 1A-B). Further, no differences were observed in ethanol consumption between male and female mice when normalized to body mass (Fig 1C-D). All animals on the NIAAA feeding protocol survived through the end of the experiment. Unexpectedly, with 6-weeks of ethanol feeding the female mice had an approximate 50% mortality rate compared with a non-significant difference in mortality rate in identically treated male mice (Fig 1E-F).

Figure 2.1




Figure 2.1 Animal Characteristics.

Separate cohorts of male and female mice were exposed to two separate ethanol feeding protocols: The NIAAA model (10 days, plus a binge on day 11) or the chronic model (6 weeks). No differences in body weights (A & C) or food intake normalized to body weight (B & D) were observed. There was no difference in the survival curve in ethanol compared with control fed male mice (E), however there was a significant difference in the female survival curve (F). Data reported as mean \pm SD (A-D), data reported as absolute numbers (E & F), $*=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male; NIAAA, National Institute on Alcohol Abuse and Alcoholism.

Multiplex serum analysis

Data from multiplex analysis is reported in Table 2. Briefly, in both models ethanol increased circulating IL-6 levels in female mice, and 6 weeks of ethanol feeding increased TNF levels in female mice. On the other hand, PAI-1 levels were increased with ethanol feeding regardless of sex or feeding model. Lastly, leptin levels were significantly elevated with ethanol feeding in both sexes with the NIAAA model, with no differences noted after 6 weeks of feeding (Chronic model).

Table 2.3 Serum Multiplex analysis

The NIAAA Model				
	Con-M	ETOH-M	Con-F	ETOH-F
IL-6 (pg/mL)	186 ± 61	292 ± 315	82 ± 112	3627 ± 1875*
TNF (pg/mL)	215 ± 57	443 ± 274	529 ± 410	602 ± 318
PAI-1 (ng/mL)	4.0 ± 1.2	10.9 ± 4.5*	3.8 ± 1.3	12.4 ± 7.2*
Leptin (ng/mL)	6.0 ± 3.4	16.3 ± 9.5*	2.4 ± 1.8	41.6 ± 26.7*
Chronic model				
	Con-M	ETOH-M	Con-F	ETOH-F
IL-6 (pg/ml)	57 ± 54	45 ± 23	36 ± 10	81 ± 55*
TNF (pg/ml)	427 ± 117	514 ± 532	338 ± 79	590 ± 263*
PAI-1 (ng/ml)	2.87 ± 0.32	3.9 ± 0.90*	3.66 ± 0.47	6.96 ± 3.2*
Leptin (ng/mL)	12.8 ± 11.6	3.4 ± 2.2	10.4 ± 4.7	7.7 ± 5.0

All data is reported as mean ± standard deviations. *=p< 0.05 Con compared with ETOH. Abbreviations: ETOH, ethanol fed; Con, control fed, M, Male; F, Female; IL-6, Interleukin-6; TNF, Tumor necrosis factor alpha; PAI-1, Plasminogen activator inhibitor-1.

Circulating Adipokines

In response to the NIAAA model, female mice fed ethanol had an increase in circulating levels of both adiponectin and CTRP1 and a decrease in both CTRP2 and CTRP3. Whereas, in the male mice ethanol decreased circulating CTRP2 with no changes observed in adiponectin, CTRP1, or CTRP3 (Fig 2).

In the chronic model, circulating CTRP2 levels were not different between control and ethanol fed mice regardless of sex. Whereas, 6 weeks of ethanol feeding more than doubled circulating adiponectin levels in both male and female mice. Unexpectedly, chronic ethanol feeding had a dimorphic effect on CTRP1 levels: CTRP1 levels were increased in female mice but lowered by ~50% in male mice. Lastly, circulating CTRP3 levels were decreased in female mice, with no change in male mice.

Figure 2.2





Figure 2.2 Circulating Adipokines in the NIAAA model of ethanol feeding.

Circulating adiponectin, CTRP1, CTRP2, and CTRP3 levels were determined by immunoblot in serum collected from mice on the 10-day plus binge model of ethanol feeding (the NIAAA model). Data reported as mean \pm SD. Male and female blots were performed and analyzed independently, and values were normalized to control fed within each sex. $n=6$, and $*=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male; NIAAA, National Institute on Alcohol Abuse and Alcoholism. Raw data files for data presented in Figure 2 are attached as supplemental data (S1).

Figure 2.3

Figure 2.3 Circulating Adipokines in the Chronic model of ethanol feeding.



Serum was collected from the mice after 6 weeks of ethanol feeding and circulating levels of adiponectin, CTRP1, CTRP2, and CTRP3 were determined by immunoblot analysis. Data reported as mean \pm SD. Male and female blots were performed and analyzed independently, and values were normalized to control fed within each sex, $n=6$, and $*=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male. Raw data files for data presented in Figure 3 are attached as supplemental data (S2).

Adipokine Gene Expression and Tissue Content

Gonadal adipose adipokine gene expression and protein content levels were analyzed to determine if differences in serum could be contributed to changes in tissue expression. However, no differences were observed in adipokine gene expression (Fig 4), indicating that changes to gene expression, at least in the gonadal fat pads, were not responsible for the changes in circulating adipokine levels. The adipose tissue protein content (Fig 5) was also analyzed and overall there was no difference in protein levels in the adipose tissue in response to either alcohol-feeding model. These data indicate that the alcohol feeding models used in this study did not alter tissue level expression of these adipokines, at least in the gonadal fat pads.

Figure 2.4




Figure 2.4 Tissue Adipokine Gene Expression.

After 6 weeks gonadal adipose tissue samples were collected from the mice. There were no significant differences found in the gene expression of adiponectin, CTRP1, CTRP2, or CTRP3 (A-D). Data reported as mean \pm SD and normalized to geometric mean. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

Figure 2.5




Figure 2.5
Adipokine Protein Content.

Tissue

Gonadal adipose tissue samples were collected from the mice after ethanol feeding for 6 weeks or 10 days (NIAAA) and adiponectin, CTRP1, CTRP2, and CTRP3 and levels were examined by immunoblot analysis. Data reported as mean \pm SD Male and female blots were performed and analyzed independently, and values were normalized to control fed within each sex, n=6, and *= $p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

DISCUSSION

The major finding of this study is that excessive ethanol consumption effects adipokine levels in a sex specific manner. The expanding family of adipokines, with their multiple biologically relevant functions, provide abundant research targets for the development of novel therapies for the treatment/prevention of human disease. The purpose of this project was to specifically examine the effects of ethanol abuse on three novel adipokines (CTRP1, CTRP2, and CTRP3). These adipokines have documented lipid regulating functions and their disruption could contribute to the development of ectopic fat deposition. Ectopic fat deposition is a major contributor to the secondary injury caused to tissues due to excessive alcohol consumption [9, 10].

The secondary finding of this study is that female mice are more sensitive to ethanol feeding than male mice. This observation has been repeatedly noted in the literature on a variety of experimental outcomes [5, 13, 56]. Specifically, in this study male and female mice consumed similar amounts of ethanol, normalized to body weight. However, female mice had significant inflammation (elevations in TNF and IL-6) and a significant mortality rate in ethanol-fed compared with control-fed mice. Conversely, there was no difference in TNF, IL-6, or mortality between control and ethanol fed male mice. It is important to note that ethanol consumption led to increased levels of PAI-1, indicating in both sexes alcohol consumption leads to damage tissues and disruptions to overall tissue homeostasis [57].

Leptin and Adiponectin

Leptin and adiponectin were the first adipokines discovered and are currently the most well studied. Circulating levels of both leptin and adiponectin have been documented to be affected by alcohol consumption, although to what extent alcohol alters these proteins is unclear. Briefly, leptin levels have been shown to increase with alcohol feeding in some models [10, 15-24], with no change in other feeding models [25], while decreasing in other alcohol feeding models [16, 23, 26]. Specifically, acute alcohol consumption decreases leptin levels in both human and animal models [21-23], however, acute reductions in leptin levels are gone approximately 9-hours post exposure [23]. On the other hand, leptin levels were significantly increased with withdrawal in alcohol patients [58, 59]. Combined, these data indicate that the time from last alcohol exposure could influence serum leptin results. Further variability in serum leptin levels with alcohol consumption could be associated to the feeding model. For example, ethanol increased leptin compared with pair-fed rats, but showed no difference compared with control ad libitum rats [25]. Our findings demonstrate that leptin levels can vary depending on the exposure model as leptin levels were increased in the NIAAA model (which includes a final ethanol bolus), but showed no difference in leptin levels after 6 weeks of chronic ethanol feeding.

Similarly, adiponectin levels have been shown to be suppressed [18, 24], not different [26], or elevated [12, 18, 19, 26-28] in response to alcohol consumption. It is suspected that oxidative stress induced by acute alcohol exposure reduces the secretion of adiponectin [29], indicating that time since last dose of ethanol can affect results. The cause of elevated levels of serum adiponectin seen in many alcohol feeding protocols has not been established. However, chronic alcohol abuse has demonstrated associations between circulating adiponectin levels and the severity of liver damage and elevated adiponectin levels in cases of cirrhosis [10, 60-62]. Our data supports the finding that circulating adiponectin levels increase with both NIAAA and chronic alcohol consumption. As circulating adiponectin levels increase with ethanol exposure, activating adiponectin and adiponectin mediated signaling pathways may not be a successful strategy for the prevention/treatment of ALD, as its levels are already increased with chronic ethanol exposure.

CTRP1

We hypothesized that the alcohol-induced loss of CTRP1 may exacerbate the effects of alcohol abuse due to decreased skeletal muscle lipid oxidation [36], resulting in excessive circulating lipid levels, thus promoting ectopic lipid accumulation. Both the NIAAA and chronic models showed CTRP1 levels were elevated in the female ethanol fed mice, compared with control fed mice, indicating that alcohol consumption does not inhibit CTRP1 levels in females. On the other hand, in the male mice, chronic ethanol feeding resulted in a significant reduction in CTRP1 levels. At this time, the mechanism responsible for the dimorphic effect of alcohol is unknown. This effect may be secondary to the observed increase in pro-inflammatory cytokines (TNF and IL-6). Regardless, alcohol-induced reduction in circulating CTRP1 levels requires further analysis to determine if similar results are shown in human alcoholics and whether restoration of CTRP1 would generate any protective effect in male mice.

CTRP2

As predicted CTRP2 levels decreased with ETOH feeding in both male and female mice during the 10-day plus binge (the NIAAA model) ethanol feeding protocol. However, there was no long-term alcohol-induced difference in CTRP2 in either sex. Therefore, the data does not support that alcohol-induced changes to circulating CTRP2 levels contribute to the adverse effects of chronic alcohol abuse.

CTRP3

Ethanol feeding reduced circulating CTRP3 levels in female, but not male, mice in response to both the NIAAA model and chronic feeding model. Alcoholic cirrhosis occurs at a higher rate in female alcoholic patients, at an earlier age, with a lower proportional amount of alcohol consumption [5, 12, 13]. As CTRP3 levels are

selectively reduced in ethanol fed female mice, this provides a novel mechanism to explore the increase susceptibility of females to alcoholic cirrhosis. In support of this hypothesis, our previous work has shown that CTRP3 acts directly on liver tissue to stimulate lipid oxidation and attenuate diet-induced fatty liver disease [38]. In fact, in human subjects CTRP3 levels are reduced with diet-induced hepatic steatosis, or non-alcoholic fatty liver disease [63]. Further, we have also previously shown that CTRP3 suppresses lipid-induced elevations in pro-inflammatory cytokines, such as TNF. However, the effects of CTRP3 on reducing alcoholic fatty liver disease are yet to be explored. Chronic alcohol consumption disrupts lipid synthesis which leads the buildup of hepatic lipids, resulting alcoholic fatty liver and eventually alcoholic cirrhosis [5, 9, 13], the leading causes of liver failure and a leading cause of death in the United States [5, 13, 14, 56]. Thus, there has been renewed interest in developing effective therapeutic strategies to prevent alcoholic fatty liver disease [4, 6, 9, 13, 64, 65]. Our data, combined with the literature, identifies CTRP3 as an ideal candidate to develop novel treatments for alcoholic fatty liver disease.

Conclusion

Since the discovery of leptin and adiponectin, research into understanding the role of adipokines in human health has become a popular topic. The expanding family of adipokines, with their multiple functions, provides abundant research targets for the development of novel therapies. This study demonstrates that sex and mode of ethanol exposure can significantly influence the results indicating that the role of alcohol on adipokines should be studied via multiple models. Lastly, we have identified the sex specific alcohol-induced reduction in CTRP3 as a potential mechanism for the increased susceptibility of females to alcoholic cirrhosis. These findings warrant further study.

Figure Legends

Figure 1: Animal characteristics.

Separate cohorts of male and female mice were exposed to two separate ethanol feeding protocols: The NIAAA model (10 days, plus a binge on day 11) or the chronic model (6 weeks). No differences in body weights (A & C) or food intake normalized to body weight (B & D) were observed. There was no difference in the survival curve in ethanol compared with control fed male mice (E), however there was a significant difference in the female survival curve (F). Data reported as mean \pm SD (A-D), data reported as absolute numbers (E & F), $*=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male; NIAAA, National Institute on Alcohol Abuse and Alcoholism.

Figure 2: Circulating Adipokines in the NIAAA model of ethanol feeding.

Circulating adiponectin, CTRP1, CTRP2, and CTRP3 levels were determined by immunoblot in serum collected from mice on the 10-day plus binge model of ethanol feeding (the NIAAA model). Data reported as mean \pm SD. Male and female blots were performed and analyzed independently, and values were normalized to control fed within each sex. $n=6$, and $*=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male; NIAAA, National Institute on Alcohol Abuse and Alcoholism. Raw data files for data presented in Figure 2 are attached as supplemental data (S1).

Figure 3: Circulating Adipokines in the Chronic model of ethanol feeding.

Serum was collected from the mice after 6 weeks of ethanol feeding and circulating levels of adiponectin, CTRP1, CTRP2, and CTRP3 were determined by immunoblot analysis. Data reported as mean \pm SD. Male and female blots were performed and analyzed independently, and values were normalized to control fed within each sex, $n=6$, and $*=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male. Raw data files for data presented in Figure 3 are attached as supplemental data (S2).

Figure 4: Tissue adipokine gene expression.

After 6 weeks gonadal adipose tissue samples were collected from the mice. There were no significant differences found in the gene expression of adiponectin, CTRP1, CTRP2, or CTRP3 (A-D). Data reported as mean \pm SD and normalized to geometric mean. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

Figure 5: Gonadal adipose tissue adipokine protein content.

Gonadal adipose tissue samples were collected from the mice after ethanol feeding for 6 weeks or 10 days (NIAAA) and adiponectin, CTRP1, CTRP2, and CTRP3 and levels were examined by immunoblot analysis. Data reported as mean \pm SD. Male and female blots were performed and analyzed independently, and values were normalized to control fed within each sex, n=6, and * $=p < 0.05$ ETOH vs Con. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male. Raw data files for data presented in Figure 5 are attached as supplemental data (S3).

References

1. Matsumoto C, Miedema MD, Ofman P, Gaziano JM, Sesso HD. An expanding knowledge of the mechanisms and effects of alcohol consumption on cardiovascular disease. *J Cardiopulm Rehabil Prev.* 2014;34(3):159-71. doi: 10.1097/HCR.000000000000042. PubMed PMID: 24667667.
2. Baumeister SE, Finger JD, Glaser S, Dorr M, Markus MR, Ewert R, et al. Alcohol consumption and cardiorespiratory fitness in five population-based studies. *Eur J Prev Cardiol.* 2017;2047487317738594. doi: 10.1177/2047487317738594. PubMed PMID: 29056079.
3. Baraona E, Lieber CS. Effects of ethanol on lipid metabolism. *J Lipid Res.* 1979;20(3):289-315. PubMed PMID: 87483.
4. Gao B, Bataller R. Alcoholic liver disease: pathogenesis and new therapeutic targets. *Gastroenterology.* 2011;141(5):1572-85. doi: 10.1053/j.gastro.2011.09.002. PubMed PMID: 21920463; PubMed Central PMCID: PMC3214974.
5. Rehm J, Taylor B, Mohapatra S, Irving H, Baliunas D, Patra J, et al. Alcohol as a risk factor for liver cirrhosis: a systematic review and meta-analysis. *Drug Alcohol Rev.* 2010;29(4):437-45. doi: 10.1111/j.1465-3362.2009.00153.x. PubMed PMID: 20636661.
6. Stickel F, Seitz HK. Alcoholic steatohepatitis. *Best Pract Res Clin Gastroenterol.* 2010;24(5):683-93. doi: 10.1016/j.bpg.2010.07.003. PubMed PMID: 20955970.
7. Choi YJ, Lee DH, Han KD, Kim HS, Yoon H, Shin CM, et al. The relationship between drinking alcohol and esophageal, gastric or colorectal cancer: A nationwide population-based cohort study of South Korea. *PLoS One.* 2017;12(10):e0185778. doi: 10.1371/journal.pone.0185778. PubMed PMID: 28973012; PubMed Central PMCID: PMC5626486.
8. Yoo MG, Kim HJ, Jang HB, Lee HJ, Park SI. The Association between Alcohol Consumption and beta-Cell Function and Insulin Sensitivity in Korean Population. *Int J Environ Res Public Health.* 2016;13(11). doi: 10.3390/ijerph13111133. PubMed PMID: 27854254; PubMed Central PMCID: PMC5129343.
9. Parker R, Kim SJ, Gao B. Alcohol, adipose tissue and liver disease: mechanistic links and clinical considerations. *Nat Rev Gastroenterol Hepatol.* 2017. doi: 10.1038/nrgastro.2017.116. PubMed PMID: 28930290.
10. Steiner JL, Lang CH. Alcohol, Adipose Tissue and Lipid Dysregulation. *Biomolecules.* 2017;7(1). doi: 10.3390/biom7010016. PubMed PMID: 28212318; PubMed Central PMCID: PMC5372728.

11. Kema VH, Mojerla NR, Khan I, Mandal P. Effect of alcohol on adipose tissue: a review on ethanol mediated adipose tissue injury. *Adipocyte*. 2015;4(4):225-31. doi: 10.1080/21623945.2015.1017170. PubMed PMID: 26451277; PubMed Central PMCID: PMC4573182.
12. Fulham MA, Mandrekar P. Sexual Dimorphism in Alcohol Induced Adipose Inflammation Relates to Liver Injury. *PLoS One*. 2016;11(10):e0164225. doi: 10.1371/journal.pone.0164225. PubMed PMID: 27711160; PubMed Central PMCID: PMC45053524.
13. Nagy LE, Ding WX, Cresci G, Saikia P, Shah VH. Linking Pathogenic Mechanisms of Alcoholic Liver Disease With Clinical Phenotypes. *Gastroenterology*. 2016;150(8):1756-68. doi: 10.1053/j.gastro.2016.02.035. PubMed PMID: 26919968; PubMed Central PMCID: PMC4487335.
14. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380(9859):2095-128. doi: 10.1016/S0140-6736(12)61728-0. PubMed PMID: 23245604.
15. He Z, Li M, Zheng D, Chen Q, Liu W, Feng L. Adipose tissue hypoxia and low-grade inflammation: a possible mechanism for ethanol-related glucose intolerance? *Br J Nutr*. 2015;113(9):1355-64. doi: 10.1017/S000711451500077X. PubMed PMID: 25989996.
16. Hiney JK, Dearth RK, Lara F, 3rd, Wood S, Srivastava V, Les Dees W. Effects of ethanol on leptin secretion and the leptin-induced luteinizing hormone (LH) release from late juvenile female rats. *Alcohol Clin Exp Res*. 1999;23(11):1785-92. PubMed PMID: 10591595.
17. Obradovic T, Meadows GG. Chronic ethanol consumption increases plasma leptin levels and alters leptin receptors in the hypothalamus and the perigonadal fat of C57BL/6 mice. *Alcohol Clin Exp Res*. 2002;26(2):255-62. PubMed PMID: 11964566.
18. Yu HC, Li SY, Cao MF, Jiang XY, Feng L, Zhao JJ, et al. Effects of chronic ethanol consumption on levels of adipokines in visceral adipose tissues and sera of rats. *Acta Pharmacol Sin*. 2010;31(4):461-9. doi: 10.1038/aps.2010.12. PubMed PMID: 20208553; PubMed Central PMCID: PMC4007662.
19. Pravdova E, Macho L, Fickova M. Alcohol intake modifies leptin, adiponectin and resistin serum levels and their mRNA expressions in adipose tissue of rats. *Endocr Regul*. 2009;43(3):117-25. PubMed PMID: 19817506.
20. Szkudelski T, Bialik I, Szkudelska K. Adipocyte lipolysis, hormonal and metabolic changes in ethanol-drinking rats. *J Anim Physiol Anim Nutr (Berl)*. 2004;88(7-8):251-8. doi: 10.1111/j.1439-0396.2004.00478.x. PubMed PMID: 15274689.

21. Rojdmarm S, Calissendorff J, Brismar K. Alcohol ingestion decreases both diurnal and nocturnal secretion of leptin in healthy individuals. *Clin Endocrinol (Oxf)*. 2001;55(5):639-47. PubMed PMID: 11894976.
22. Calissendorff J, Brismar K, Rojdmarm S. Is decreased leptin secretion after alcohol ingestion catecholamine-mediated? *Alcohol Alcohol*. 2004;39(4):281-6. doi: 10.1093/alcalc/agh054. PubMed PMID: 15208157.
23. Otaka M, Konishi N, Odashima M, Jin M, Wada I, Matsuhashi T, et al. Effect of alcohol consumption on leptin level in serum, adipose tissue, and gastric mucosa. *Dig Dis Sci*. 2007;52(11):3066-9. doi: 10.1007/s10620-006-9635-x. PubMed PMID: 17406835.
24. Chen X, Sebastian BM, Nagy LE. Chronic ethanol feeding to rats decreases adiponectin secretion by subcutaneous adipocytes. *Am J Physiol Endocrinol Metab*. 2007;292(2):E621-8. doi: 10.1152/ajpendo.00387.2006. PubMed PMID: 17047161; PubMed Central PMCID: PMC1794258.
25. Strbak V, Benicky J, Macho L, Jezova D, Nikodemova M. Four-week ethanol intake decreases food intake and body weight but does not affect plasma leptin, corticosterone, and insulin levels in pubertal rats. *Metabolism*. 1998;47(10):1269-73. PubMed PMID: 9781633.
26. Tan X, Sun X, Li Q, Zhao Y, Zhong W, Sun X, et al. Leptin deficiency contributes to the pathogenesis of alcoholic fatty liver disease in mice. *Am J Pathol*. 2012;181(4):1279-86. doi: 10.1016/j.ajpath.2012.06.013. PubMed PMID: 22841822; PubMed Central PMCID: PMC3463622.
27. Imhof A, Plamper I, Maier S, Trischler G, Koenig W. Effect of drinking on adiponectin in healthy men and women: a randomized intervention study of water, ethanol, red wine, and beer with or without alcohol. *Diabetes Care*. 2009;32(6):1101-3. doi: 10.2337/dc08-1723. PubMed PMID: 19244090; PubMed Central PMCID: PMC2681019.
28. Sierksma A, Patel H, Ouchi N, Kihara S, Funahashi T, Heine RJ, et al. Effect of moderate alcohol consumption on adiponectin, tumor necrosis factor-alpha, and insulin sensitivity. *Diabetes Care*. 2004;27(1):184-9. PubMed PMID: 14693987.
29. Tang H, Sebastian BM, Axhemi A, Chen X, Hillian AD, Jacobsen DW, et al. Ethanol-induced oxidative stress via the CYP2E1 pathway disrupts adiponectin secretion from adipocytes. *Alcohol Clin Exp Res*. 2012;36(2):214-22. doi: 10.1111/j.1530-0277.2011.01607.x. PubMed PMID: 21895711; PubMed Central PMCID: PMC3235233.
30. Ohashi K, Shibata R, Murohara T, Ouchi N. Role of anti-inflammatory adipokines in obesity-related diseases. *Trends Endocrinol Metab*. 2014;25(7):348-55. doi: 10.1016/j.tem.2014.03.009. PubMed PMID: 24746980.

31. Scherer PE, Williams S, Fogliano M, Baldini G, Lodish HF. A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem.* 1995;270(45):26746-9. PubMed PMID: 7592907.
32. Wong GW, Wang J, Hug C, Tsao TS, Lodish HF. A family of Acrp30/adiponectin structural and functional paralogs. *Proc Natl Acad Sci U S A.* 2004;101(28):10302-7. doi: 10.1073/pnas.0403760101. PubMed PMID: 15231994; PubMed Central PMCID: PMCPMC478567.
33. Shapiro L, Scherer PE. The crystal structure of a complement-1q family protein suggests an evolutionary link to tumor necrosis factor. *Curr Biol.* 1998;8(6):335-8. PubMed PMID: 9512423.
34. Byerly MS, Petersen PS, Ramamurthy S, Seldin MM, Lei X, Provost E, et al. C1q/TNF-related protein 4 (CTRP4) is a unique secreted protein with two tandem C1q domains that functions in the hypothalamus to modulate food intake and body weight. *J Biol Chem.* 2014;289(7):4055-69. doi: 10.1074/jbc.M113.506956. PubMed PMID: 24366864; PubMed Central PMCID: PMCPMC3924272.
35. Petersen PS, Wolf RM, Lei X, Peterson JM, Wong GW. Immunomodulatory roles of CTRP3 in endotoxemia and metabolic stress. *Physiol Rep.* 2016;4(5). doi: 10.14814/phy2.12735. PubMed PMID: 26997632; PubMed Central PMCID: PMCPMC4823594.
36. Peterson JM, Aja S, Wei Z, Wong GW. CTRP1 protein enhances fatty acid oxidation via AMP-activated protein kinase (AMPK) activation and acetyl-CoA carboxylase (ACC) inhibition. *J Biol Chem.* 2012;287(2):1576-87. doi: 10.1074/jbc.M111.278333. PubMed PMID: 22086915; PubMed Central PMCID: PMCPMC3256898.
37. Peterson JM, Seldin MM, Tan SY, Wong GW. CTRP2 overexpression improves insulin and lipid tolerance in diet-induced obese mice. *PLoS One.* 2014;9(2):e88535. doi: 10.1371/journal.pone.0088535. PubMed PMID: 24586339; PubMed Central PMCID: PMCPMC3930646.
38. Peterson JM, Seldin MM, Wei Z, Aja S, Wong GW. CTRP3 attenuates diet-induced hepatic steatosis by regulating triglyceride metabolism. *Am J Physiol Gastrointest Liver Physiol.* 2013;305(3):G214-24. doi: 10.1152/ajpgi.00102.2013. PubMed PMID: 23744740; PubMed Central PMCID: PMCPMC3742855.
39. Peterson JM, Wei Z, Seldin MM, Byerly MS, Aja S, Wong GW. CTRP9 transgenic mice are protected from diet-induced obesity and metabolic dysfunction. *Am J Physiol Regul Integr Comp Physiol.* 2013;305(5):R522-33. doi: 10.1152/ajpregu.00110.2013. PubMed PMID: 23842676; PubMed Central PMCID: PMCPMC3763026.
40. Peterson JM, Wei Z, Wong GW. C1q/TNF-related protein-3 (CTRP3), a novel adipokine that regulates hepatic glucose output. *J Biol Chem.* 2010;285(51):39691-

701. doi: 10.1074/jbc.M110.180695. PubMed PMID: 20952387; PubMed Central PMCID: PMC3000950.
41. Seldin MM, Tan SY, Wong GW. Metabolic function of the CTRP family of hormones. *Rev Endocr Metab Disord.* 2014;15(2):111-23. doi: 10.1007/s11154-013-9255-7. PubMed PMID: 23963681; PubMed Central PMCID: PMC3931758.
42. Wei Z, Peterson JM, Wong GW. Metabolic regulation by C1q/TNF-related protein-13 (CTRP13): activation OF AMP-activated protein kinase and suppression of fatty acid-induced JNK signaling. *J Biol Chem.* 2011;286(18):15652-65. doi: 10.1074/jbc.M110.201087. PubMed PMID: 21378161; PubMed Central PMCID: PMC3091174.
43. Wei Z, Seldin MM, Natarajan N, Djemal DC, Peterson JM, Wong GW. C1q/tumor necrosis factor-related protein 11 (CTRP11), a novel adipose stroma-derived regulator of adipogenesis. *J Biol Chem.* 2013;288(15):10214-29. doi: 10.1074/jbc.M113.458711. PubMed PMID: 23449976; PubMed Central PMCID: PMC3624406.
44. Wolf RM, Lei X, Yang ZC, Nyandjo M, Tan SY, Wong GW. CTRP3 deficiency reduces liver size and alters IL-6 and TGFbeta levels in obese mice. *Am J Physiol Endocrinol Metab.* 2016;310(5):E332-45. doi: 10.1152/ajpendo.00248.2015. PubMed PMID: 26670485; PubMed Central PMCID: PMC4773650.
45. Wong GW, Krawczyk SA, Kitidis-Mitrokostas C, Ge G, Spooner E, Hug C, et al. Identification and characterization of CTRP9, a novel secreted glycoprotein, from adipose tissue that reduces serum glucose in mice and forms heterotrimers with adiponectin. *FASEB J.* 2009;23(1):241-58. doi: 10.1096/fj.08-114991. PubMed PMID: 18787108; PubMed Central PMCID: PMC2626616.
46. Wong GW, Krawczyk SA, Kitidis-Mitrokostas C, Revett T, Gimeno R, Lodish HF. Molecular, biochemical and functional characterizations of C1q/TNF family members: adipose-tissue-selective expression patterns, regulation by PPAR-gamma agonist, cysteine-mediated oligomerizations, combinatorial associations and metabolic functions. *Biochem J.* 2008;416(2):161-77. doi: 10.1042/BJ20081240. PubMed PMID: 18783346; PubMed Central PMCID: PMC3936483.
47. Hofmann C, Chen N, Obermeier F, Paul G, Buchler C, Kopp A, et al. C1q/TNF-related protein-3 (CTRP-3) is secreted by visceral adipose tissue and exerts antiinflammatory and antifibrotic effects in primary human colonic fibroblasts. *Inflamm Bowel Dis.* 2011;17(12):2462-71. doi: 10.1002/ibd.21647. PubMed PMID: 21351204.
48. Kopp A, Bala M, Buechler C, Falk W, Gross P, Neumeier M, et al. C1q/TNF-related protein-3 represents a novel and endogenous lipopolysaccharide antagonist of the adipose tissue. *Endocrinology.* 2010;151(11):5267-78. doi: 10.1210/en.2010-0571. PubMed PMID: 20739398.

49. Kopp A, Bala M, Weigert J, Buchler C, Neumeier M, Aslanidis C, et al. Effects of the new adiponectin paralogous protein CTRP-3 and of LPS on cytokine release from monocytes of patients with type 2 diabetes mellitus. *Cytokine*. 2010;49(1):51-7. doi: 10.1016/j.cyto.2009.10.001. PubMed PMID: 19955001.
50. Schmid A, Kopp A, Aslanidis C, Wabitsch M, Muller M, Schaffler A. Regulation and function of C1Q/TNF-related protein-5 (CTRP-5) in the context of adipocyte biology. *Exp Clin Endocrinol Diabetes*. 2013;121(5):310-7. doi: 10.1055/s-0032-1333299. PubMed PMID: 23430573.
51. Schmid A, Kopp A, Hanses F, Bala M, Muller M, Schaffler A. The novel adipokine C1q/TNF-related protein-3 is expressed in human adipocytes and regulated by metabolic and infection-related parameters. *Exp Clin Endocrinol Diabetes*. 2012;120(10):611-7. doi: 10.1055/s-0032-1323803. PubMed PMID: 23174996.
52. Schmid A, Kopp A, Hanses F, Karrasch T, Schaffler A. C1q/TNF-related protein-3 (CTRP-3) attenuates lipopolysaccharide (LPS)-induced systemic inflammation and adipose tissue Erk-1/-2 phosphorylation in mice in vivo. *Biochem Biophys Res Commun*. 2014;452(1):8-13. doi: 10.1016/j.bbrc.2014.06.054. PubMed PMID: 24996172.
53. Li Y, Wright GL, Peterson JM. C1q/TNF-Related Protein 3 (CTRP3) Function and Regulation. *Compr Physiol*. 2017;7(3):863-78. doi: 10.1002/cphy.c160044. PubMed PMID: 28640446.
54. Bertola A, Mathews S, Ki SH, Wang H, Gao B. Mouse model of chronic and binge ethanol feeding (the NIAAA model). *Nat Protoc*. 2013;8(3):627-37. doi: 10.1038/nprot.2013.032. PubMed PMID: 23449255; PubMed Central PMCID: PMC3788579.
55. Cohen JI, Roychowdhury S, McMullen MR, Stavitsky AB, Nagy LE. Complement and alcoholic liver disease: role of C1q in the pathogenesis of ethanol-induced liver injury in mice. *Gastroenterology*. 2010;139(2):664-74, 74 e1. doi: 10.1053/j.gastro.2010.04.041. PubMed PMID: 20416309; PubMed Central PMCID: PMC3273045.
56. Shoreibah M, Raff E, Bloomer J, Kakati D, Rasheed K, Kuo YF, et al. Alcoholic liver disease presents at advanced stage and progresses faster compared to non-alcoholic fatty liver diseases. *Ann Hepatol*. 2016;15(2):183-9. doi: 10.5604/16652681.1193707. PubMed PMID: 26845595.
57. Ghosh AK, Vaughan DE. PAI-1 in tissue fibrosis. *J Cell Physiol*. 2012;227(2):493-507. doi: 10.1002/jcp.22783. PubMed PMID: 21465481; PubMed Central PMCID: PMC3204398.

58. Kraus T, Reulbach U, Bayerlein K, Mugele B, Hillemacher T, Sperling W, et al. Leptin is associated with craving in females with alcoholism. *Addict Biol.* 2004;9(3-4):213-9. doi: 10.1080/13556210412331292541. PubMed PMID: 15511715.
59. Kiefer F, Jahn H, Schick M, Wiedemann K. Alcohol intake, tumour necrosis factor-alpha, leptin and craving: factors of a possibly vicious circle? *Alcohol Alcohol.* 2002;37(4):401-4. PubMed PMID: 12107045.
60. Tietge UJ, Boker KH, Manns MP, Bahr MJ. Elevated circulating adiponectin levels in liver cirrhosis are associated with reduced liver function and altered hepatic hemodynamics. *Am J Physiol Endocrinol Metab.* 2004;287(1):E82-9. doi: 10.1152/ajpendo.00494.2003. PubMed PMID: 15010338.
61. Xu A, Wang Y, Keshaw H, Xu LY, Lam KS, Cooper GJ. The fat-derived hormone adiponectin alleviates alcoholic and nonalcoholic fatty liver diseases in mice. *J Clin Invest.* 2003;112(1):91-100. doi: 10.1172/JCI17797. PubMed PMID: 12840063; PubMed Central PMCID: PMCPMC162288.
62. You M, Considine RV, Leone TC, Kelly DP, Crabb DW. Role of adiponectin in the protective action of dietary saturated fat against alcoholic fatty liver in mice. *Hepatology.* 2005;42(3):568-77. doi: 10.1002/hep.20821. PubMed PMID: 16108051; PubMed Central PMCID: PMCPMC1398076.
63. Zhang J, Zhang B, Cheng Y, Xu J. Low serum CTRP3 levels are associated with nonalcoholic fatty liver disease in patients with type 2 diabetes mellitus. *Cytokine.* 2017. doi: 10.1016/j.cyto.2017.10.023. PubMed PMID: 29113741.
64. Sanyal AJ. Novel therapeutic targets for steatohepatitis. *Clin Res Hepatol Gastroenterol.* 2015;39 Suppl 1:S46-50. doi: 10.1016/j.clinre.2015.05.012. PubMed PMID: 26160474.
65. Eguchi A, Povero D, Alkhoury N, Feldstein AE. Novel therapeutic targets for nonalcoholic fatty liver disease. *Expert Opin Ther Targets.* 2013;17(7):773-9. doi: 10.1517/14728222.2013.789502. PubMed PMID: 23600493.

CHAPTER 3

METHODS

Adipose Tissue Lipid Analysis

Lipids were extracted as described by Bligh and Dyer and as previously performed (16, 18). Briefly, adipose tissue samples were weighed then homogenized in phosphate-buffered saline, 3.75 ml/ml of sample homogenate 1:2 (vol/vol) chloroform-methanol was added, followed by the addition of 1.25 ml/ml chloroform, followed by 1.25 ml distilled water. Samples were vortexed for 30 s between each addition. Samples were then centrifuged at 1,100x g for 10 min at room temperature to give a two-phase solution (aqueous phase on top and organic phase below). The lower phase was collected with a glass pipette with gentle positive pressure. Samples were then divided into 2 aliquots and dried under nitrogen gas at 60°C. To measure total triglyceride levels one aliquot from each sample was dissolved in tert-butyl alcohol-Triton X-100 (3:2 vol/vol) solution. Triglycerides were quantified via colorimetric assay according to manufactures directions (Infinity Triglycerides, Fisher Diagnostics, Cat# TR22421). The remaining aliquot was prepared for Fatty Acid Methyl Ester analysis.

Fatty Acid Methyl Ester Preparation and Analysis

Fatty acids (FA) of snap-frozen adipose tissue samples were extracted using the fatty acid methyl ester (FAME) method and measured with GC-MS. Briefly, boron trifluoride-methanol reagent (B1252; Sigma-Aldrich, St Louis, MO, USA) was added to a prepared aliquot of isolated adipose tissue lipids. The tube was then closed and heated in a block heater at 100°C for 1 hour before returning to room temperature. 1.5 mL distilled water was added and samples were centrifuged for 1 minute at 4000x g. FAMEs were

extracted in the hexane phase, dried under nitrogen gas, suspended in 275 μ L hexane, 5 μ L C17 internal standard (1:9 hexane dilution) was added, and samples were stored at -80° C until further analysis. Gas chromatography (GC) using a flame-ionization detector, (Shimadzu GC- 2010; Shimadzu Corporation, Kyoto Japan) was performed on the samples using a capillary column (Zebron ZB-WAX, 30 m length, 0.25 mm i.d., 0.25 μ m film thickness; Phenomenex, Torrance, CA, USA). The peaks were identified by comparison with Supelco 37 component FAME mix fatty acid standard (Sigma-aldrich Cat# 47885-U).

Serum Evaluations

Serum glucose and triglyceride concentrations were determined using commercially available assays according to manufactures directions (Glucose: Calbiochem. Cat# CBA086; Triglycerides: Infinity Triglycerides, Fisher Diagnostics. Cat# TR22421). Serum ALT levels were measured using an enzymatic assay kit (Cat# TR71121) according to manufacturer's instructions. Serum AST levels were measured using an enzymatic assay kit (Cat# A7561) according to manufacturer's instructions.

CHAPTER 4

RESULTS

Effects of Ethanol on Circulating Glucose

The chronic ethanol feeding model showed no differences in levels of glucose in serum when comparing ETOH fed male and female mice to their respective controls (Figure 1).




Figure 1.

Figure 4.1 Chronic Serum Glucose

Circulating levels of glucose were determined using a commercially available assay. Serum was collected from the mice after 10-day plus binge model of ethanol feeding (NIAAA) and after 6 weeks of ethanol feeding (Chronic). Data is reported as mean \pm SD. * $p < 0.05$ ETOH vs. Control. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

Effects of Ethanol on Circulating Triglycerides

The chronic ethanol feeding model showed no differences in circulating triglycerides when comparing ETOH fed male and female mice to their respective controls (Fig 2).

Figure 2.




Figure 4.2 NIAAA and Chronic Serum Triglycerides

Circulating levels of triglycerides were determined using a commercially available assay. Serum was collected from the mice after the NIAAA model of ethanol feeding (A) and after the chronic model of ethanol feeding (B). Data is reported as mean \pm SD. * $p < 0.05$ ETOH vs. Control. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

Effects of Ethanol on Serum Transaminases

The NIAAA model showed no differences in ALT levels when comparing ETOH fed male and female mice to their respective controls (Fig 3 A). However, the chronic model showed a significant increase ($p < 0.05$) in ALT and AST levels in female ETOH fed mice compared to the female control group (Fig 3 B-C) while there was no difference found in ALT and AST levels for chronic male ETOH fed mice compared to the male control group (Fig 3 B-C).

Figure 3.




Figure 4.3 NIAAA and Chronic Serum Transaminases

ALT (A-B) and AST (C) levels were measured in the serum using a commercially available assay. Serum was collected from the mice after the NIAAA model of ethanol feeding and after the chronic model of ethanol feeding. Data is reported as mean \pm SD. * $p < 0.05$ ETOH vs. Control. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

Chronic Adipose Tissue Lipid Analysis

The chronic model male mice showed no significant differences between the ETOH fed male mice and the control fed male mice. However, the ETOH fed female mice showed a significant increase in levels of oleic acid with ETOH consumption compared to the control fed female mice.

Figure 4.




Figure 4.4 Chronic Adipose Tissue Lipid Analysis

Adipose tissue lipids were isolated, extracted using the FAME method, and measured with gas-chromatography mass spectrometry. Data was analyzed using a 1-way ANOVA and is represented by mean + S.D. * $p < 0.05$ ETOH vs. Control. Abbreviations: Con-F, pair-fed female; ETOH-F, ethanol fed female; Con-M, pair-fed male; ETOH-M, ethanol fed male.

CHAPTER 5

DISCUSSION

The aim of this study was to determine the effect alcohol consumption has in mice on levels of adipokines: CTRP1, CTRP2, and CTRP3. Because adipokines have lipid-regulating functions, their disruption could contribute to ectopic fat deposition, which is a major contributor to the secondary injury caused to tissues due to excessive alcohol consumption (Lang and Steiner 2017; Parker 2018). The major finding of this study is that excessive alcohol consumption affects adipokine levels in a sex specific manner, as female mice appear to be more sensitive to ethanol consumption. Alcoholic cirrhosis occurs at a higher rate in female alcoholic patients, at an earlier age, with a lower proportional amount of alcohol consumption (Rehm et al. 2010; Mandrekar and Fulham 2016; Nagy et al. 2016). In this study, male and female mice consumed similar amounts of ethanol, however, ethanol fed female mice showed significant inflammation (elevations in TNF and IL-6) and a significantly higher mortality rate when compared to control-fed mice. Where as, there was no difference in TNF, IL-6, or mortality between control and ethanol fed male mice. One potential reason for this is because women have less body water than men, so they exhibit a higher concentration of alcohol in the blood when similar amounts of alcohol are consumed (Rehm et al. 2010; Ghosh and Vaughan 2012). It is important to note that ethanol consumption led to increased levels of PAI-1, indicating in both sexes alcohol consumption leads to damage tissues and disruptions to overall tissue homeostasis (Ghosh and Vaughan 2012). The levels of alanine transaminase (ALT) and aspartate transaminase (AST) were analyzed in both the NIAAA and chronic models. Transaminases are liver enzymes that can be measured in the serum

to determine the degree of injury to the liver (Barone et al. 2016; Kim et al. 2016). In the chronic model, female ETOH fed mice showed a significant increase in both ALT and AST levels, twice as much as the male ETOH fed mice. This contributes to our finding that female mice were more sensitive to the ETOH than the male mice.

Leptin and Adiponectin

It has been documented that circulating levels of leptin and adiponectin are affected by alcohol consumption. Our findings demonstrate that leptin levels can vary depending on the feeding model, as leptin levels were increased in the NIAAA model (which includes a final ethanol bolus) but showed no difference after 6 weeks of chronic ethanol feeding. Considering the chronic model showed no difference in leptin levels between control and ETOH groups, the increase in leptin levels found in ETOH fed mice in the NIAAA model could be a result of the final ethanol gavage given to the mice before serum and tissue samples were collected. The chronic model showed that alcohol consumption did not affect the circulating levels of leptin, suggesting that over a longer period of time, leptin levels will be unchanged by alcohol consumption. This variability could be due to the time of the last alcohol exposure, or possibly dependent upon the feeding model used in the study.

Adiponectin levels have also been shown to be unaffected (Tan et al. 2012), suppressed (Chen et al. 2007; Yu et al. 2010), or increased (Sierksma et al. 2004; Pravdova et al. 2009; Mandrekar and Fulham 2016) with the consumption of alcohol. The cause of elevated levels of serum adiponectin seen in many alcohol feeding protocols has not been established. However, chronic alcohol abuse has demonstrated associations

between circulating adiponectin levels and the severity of liver damage and elevated adiponectin levels in cases of cirrhosis (Xu et al. 2003; Tietge et al. 2004; You et al. 2005; Lang and Steiner 2017). Contrary to our hypothesis, our data supports the finding that circulating adiponectin levels increase with both NIAAA and chronic alcohol consumption. As circulating adiponectin levels increase with ethanol exposure, activating adiponectin and adiponectin mediated signaling pathways may not be a successful strategy for the prevention/treatment of ALD, as its levels are already increased with chronic ethanol exposure.

CTRP1

We hypothesized that the alcohol-induced loss of CTRP1 may enhance the effects of alcohol abuse due to decreased skeletal muscle lipid oxidation (Scherer and Shapiro 1998), resulting in excessive circulating lipid levels, thus promoting ectopic lipid accumulation. In our study, the NIAAA model showed increased circulating levels of CTRP1 in female ETOH fed mice and no difference in the circulating levels of CTRP1 in the male ETOH fed mice. This could be due to the increase levels of CTRP1 found in the NIAAA adipose tissue for ETOH fed female mice. After 6 weeks of ethanol feeding CTRP1 levels were elevated in the female ethanol fed mice, compared with control fed mice; indicating that chronic alcohol consumption does not inhibit CTRP1 levels in females. On the other hand, after 6 weeks of ethanol feeding there was a significant decrease in CTRP1 levels in ETOH fed male mice. The mechanism responsible for the dimorphic effect of alcohol is unknown. This effect may be secondary to the observed increase in pro-inflammatory cytokines (TNF and IL-6). Alcohol-induced reduction in circulating CTRP1 levels requires further analysis to determine if similar results are

shown in human alcoholics and whether restoration of CTRP1 would generate any protective effect in male mice.

CTRP2

As predicted, CTRP2 levels decreased with ETOH feeding in both male and female mice during the 10-day plus binge (the NIAAA model) ethanol feeding protocol. However, there was no long-term alcohol-induced difference in CTRP2 in either sex. Therefore, the data does not support that alcohol-induced changes to circulating CTRP2 levels contribute to the adverse effects of chronic alcohol abuse.

CTRP3

Ethanol feeding reduced circulating CTRP3 levels in female, but not male mice, in both the NIAAA model and chronic feeding model. In fact, in human subjects CTRP3 levels are reduced with diet-induced hepatic steatosis, or non-alcoholic fatty liver disease (Zhang et al. 2017). We have previously shown that CTRP3 suppresses lipid-induced elevations in pro-inflammatory cytokines, such as TNF. However, the effects of CTRP3 on reducing alcoholic fatty liver disease are yet to be explored. Our data, combined with the literature, identifies CTRP3 as an ideal candidate to develop novel treatments for alcoholic fatty liver disease.

CHAPTER 6

CONCLUSION

In conclusion, alcohol consumption does have an effect on novel adipokines: CTRP1, CTRP2, and CTRP3. Chronic alcohol consumption disrupts lipid synthesis, which leads to the buildup of hepatic lipids, resulting alcoholic fatty liver and eventually alcoholic cirrhosis (Nagy et al. 2016; Parker 2018; Rehm et al. 2010), the leading causes of liver failure and a leading cause of death in the United States (Nagy et al. 2016; Rehm et al. 2010). Based on our research, there are some differences in the effects of excessive alcohol consumption between males and females. Females are more sensitive to excessive alcohol consumption, but CTRP3 could provide an ideal candidate to develop novel treatment. Because CTRP3 levels are selectively reduced in ETOH fed female mice, this provides a novel mechanism to explore the increase susceptibility of females to alcoholic cirrhosis. Overexpressing CTRP3 in female mice fed an ETOH diet would prove to be a beneficial trial for the effects of CTRP3 on ameliorating AFLD in females. Based on our findings for the male ETOH fed mice, with more research, CTRP1 could prove to be a target for treating AFLD in males.

REFERENCES

- Ahima RS, Flier JS. 2000 Oct. Adipose tissue as an endocrine organ. *Trends in endocrinology and metabolism: TEM*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=10996528>
- Arteel GE. 2008 Mar. New role of plasminogen activator inhibitor-1 in alcohol-induced liver injury. *Journal of gastroenterology and hepatology*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=PMC2413330>
- Baumeister SE, Finger JD, Glaser S, Dorr M, Markus MR, Ewert R, et al. Alcohol consumption and cardiorespiratory fitness in five population-based studies. *Eur J Prev Cardiol*. 2017;2047487317738594. doi: 10.1177/2047487317738594. PubMed PMID: 29056079.
- Baraona E, Lieber CS. Effects of ethanol on lipid metabolism. *J Lipid Res*. 1979;20(3):289-315. PubMed PMID: 87483.
- Barone, R., Rappa, F., Macaluso, F., Caruso, C., Sangiorgi, C., Di, G., . . . Marino, A. (2016). Alcoholic Liver Disease: A Mouse Model Reveals Protection by *Lactobacillus fermentum*. *Clinical and Translational Gastroenterology*. doi:10.1038/ctg.2015.66.
- Bertola A, Mathews S, Ki SH, Wang H, Gao B. Mouse model of chronic and binge ethanol feeding (the NIAAA model). *Nat Protoc*. 2013;8(3):627-37. doi: 10.1038/nprot.2013.032. PubMed PMID: 23449255; PubMed Central PMCID: PMC3788579.
- Bertolani, C., & Marra, F. (2008). The role of adipokines in liver fibrosis. *Pathophysiology : The Official Journal of the International Society for Pathophysiology*, 15(2), 91-101. doi:10.1016/j.pathophys.2008.05.001.
- Breitkopf, K., Nagy, L. E., Beier, J. I., Mueller, S., Weng, H., & Dooley, S. (2009). Current experimental perspectives on the clinical progression of alcoholic liver disease. *Alcoholism Clinical and Experimental Research*, 33(10), 1647-1655. doi:10.1111/j.1530-0277.2009.01015.x.
- Byerly MS, Petersen PS, Ramamurthy S, Seldin MM, Lei X, Provost E, et al. C1q/TNF-related protein 4 (CTRP4) is a unique secreted protein with two tandem C1q domains that functions in the hypothalamus to modulate food intake and body weight. *J Biol Chem*. 2014;289(7):4055-69. doi: 10.1074/jbc.M113.506956. PubMed PMID: 24366864; PubMed Central PMCID: PMC3924272.
- Calissendorff J, Brismar K, Rojdmarm S. Is decreased leptin secretion after alcohol ingestion catecholamine-mediated? *Alcohol Alcohol*. 2004;39(4):281-6. doi:10.1093/alcalc/agh054. PubMed PMID: 15208157.

- Chen, X., Sebastian, B. M., & Nagy, L. E. (2007). Chronic Ethanol Feeding to Rats Decreases Adiponectin Secretion by Subcutaneous Adipocytes. *American Journal of Physiology*, 292(2). doi:10.1152/ajpendo.00387.2006
- Cohen JI, Roychowdhury S, McMullen MR, Stavitsky AB, Nagy LE. Complement and alcoholic liver disease: role of C1q in the pathogenesis of ethanol-induced liver injury in mice. *Gastroenterology*. 2010;139(2):664-74, 74 e1. doi: 10.1053/j.gastro.2010.04.041. PubMed PMID: 20416309; PubMed Central PMCID: PMC3273045.
- Choi YJ, Lee DH, Han KD, Kim HS, Yoon H, Shin CM, et al. The relationship between drinking alcohol and esophageal, gastric or colorectal cancer: A nationwide population-based cohort study of South Korea. *PLoS One*. 2017;12(10):e0185778. doi: 10.1371/journal.pone.0185778. PubMed PMID: 28973012; PubMed Central PMCID: PMC5626486.
- Coelho M, Oliveira T, Fernandes R. 2013 Apr 20. Biochemistry of adipose tissue: an endocrine organ. *Archives of medical science : AMS*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=23671428>
- Eguchi A, Povero D, Alkhouri N, Feldstein AE. Novel therapeutic targets for nonalcoholic fatty liver disease. *Expert Opin Ther Targets*. 2013;17(7):773-9. doi: 10.1517/14728222.2013.789502. PubMed PMID: 23600493.
- Fujii, H. (2014). Fibrogenesis in alcoholic liver disease. *World Journal of Gastroenterology*, 20(25), 8048-8054. doi:10.3748/wjg.v20.i25.8048
- Fulham, M. A., & Mandrekar, P. (2016). Sexual Dimorphism in Alcohol Induced Adipose Inflammation Relates to Liver Injury. *PLOS One*, 10(11). Retrieved from <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0164225>
- Gao B, Bataller R. 2011 Nov. Alcoholic liver disease: pathogenesis and new therapeutic targets. *Gastroenterology*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/21920463>
- Ghosh, A. K., & Vaughan, D. E. (2012). PAI-1 in tissue fibrosis. *Journal of Cellular Physiology*, 227(2), 493-507. doi:10.1002/jcp.22783
- He Z, Li M, Zheng D, Chen Q, Liu W, Feng L. 2015 May 14. Adipose tissue hypoxia and low-grade inflammation: a possible mechanism for ethanol-related glucose intolerance? *The British journal of nutrition*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=25989996>

- Hiney JK, Dearth RK, Lara 3 r, Wood S, Srivastava V, Les W. 1999 Nov. Effects of ethanol on leptin secretion and the leptin-induced luteinizing hormone (LH) release from late juvenile female rats. *Alcoholism, clinical and experimental research*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=10591595>
- Hong F, Kim WH, Tian Z, Jaruga B, Ishac E, Shen X, Gao B. 2002 Jan 3. Elevated interleukin-6 during ethanol consumption acts as a potential endogenous protective cytokine against ethanol-induced apoptosis in the liver: involvement of induction of Bcl-2 and Bcl-x(L) proteins. *Oncogene*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=11791174>
- Hou B, Eren M, Painter CA, Covington JW, Dixon JD, Schoenhard JA, Vaughan DE. 2004 Apr 30. Tumor necrosis factor alpha activates the human plasminogen activator inhibitor-1 gene through a distal nuclear factor kappaB site. *The Journal of biological chemistry*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=14963043>
- Imhof A, Plamper I, Maier S, Trischler G, Koenig W. Effect of drinking on adiponectin in healthy men and women: a randomized intervention study of water, ethanol, red wine, and beer with or without alcohol. *Diabetes Care*. 2009;32(6):1101-3. doi: 10.2337/dc08-1723. PubMed PMID: 19244090; PubMed Central PMCID:PMCPMC2681019.
- Hofmann C, Chen N, Obermeier F, Paul G, Buchler C, Kopp A, et al. C1q/TNFR-related protein-3 (CTRP-3) is secreted by visceral adipose tissue and exerts antiinflammatory and antifibrotic effects in primary human colonic fibroblasts. *Inflamm Bowel Dis*. 2011;17(12):2462-71. doi: 10.1002/ibd.21647. PubMed PMID:21351204.
- Karbowska J, Kochan Z. 2006 Nov. Role of adiponectin in the regulation of carbohydrate and lipid metabolism. *Journal of physiology and pharmacology : an official journal of the Polish Physiological Society*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=17228091>
- Kema VH, Mojerla NR, Khan I, Mandal P. 2015 Apr 2. Effect of alcohol on adipose tissue: a review on ethanol mediated adipose tissue injury. *Adipocyte*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=26451277>
- Kiefer F, Jahn H, Schick M, Wiedemann K. 2002 Jul. Alcohol intake, tumour necrosis factor-alpha, leptin and craving: factors of a possibly vicious circle? *Alcohol and alcoholism (Oxford, Oxfordshire)*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=12107045>

- Kim, W. R., Flamm, S. L., Di, A. M., Bodenheimer, H. C., & Public, D. I. (2008). Serum activity of alanine aminotransferase (ALT) as an indicator of health and disease. *Hepatology*, 47(4), 1363-1370. doi:10.1002/hep.22109.
- Kopp A, Bala M, Buechler C, Falk W, Gross P, Neumeier M, et al. C1q/TNF-related protein-3 represents a novel and endogenous lipopolysaccharide antagonist of the adipose tissue. *Endocrinology*. 2010;151(11):5267-78. doi: 10.1210/en.2010-0571. PubMed PMID: 20739398.
- Kopp A, Bala M, Weigert J, Buchler C, Neumeier M, Aslanidis C, et al. Effects of the new adiponectin paralogous protein CTRP-3 and of LPS on cytokine release from monocytes of patients with type 2 diabetes mellitus. *Cytokine*.
- Kraus T, Reulbach U, Bayerlein K, Mugele B, Hillemacher T, Sperling W, et al. Leptin is associated with craving in females with alcoholism. *Addict Biol*. 2004;9(3-4):213-9. doi: 10.1080/13556210412331292541. PubMed PMID: 15511715.
- Leclercq, I. A., Farrell, G. C., Schriemer, R., & Robertson, G. R. (2002). Leptin is essential for the hepatic fibrogenic response to chronic liver injury. *Journal of Hepatology*, 37(2), 206-213. doi:10.1016/S0168-8278(02)00102-2.
- Li Y, Wright GL, Peterson JM. C1q/TNF-Related Protein 3 (CTRP3) Function and Regulation. *Compr Physiol*. 2017;7(3):863-78. doi: 10.1002/cphy.c160044. PubMed PMID: 28640446.
- Lieber CS. 2000 Dec. Hepatic, metabolic, and nutritional disorders of alcoholism: from pathogenesis to therapy. *Critical reviews in clinical laboratory sciences*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/11192332>
- Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380(9859):2095-128. doi: 10.1016/S0140-6736(12)61728-0. PubMed PMID:23245604.
- McClain CJ, Barve S, Deaciuc I, Hill DB. 1998 Aug. Tumor necrosis factor and alcoholic liver disease. *Alcoholism, clinical and experimental research*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=9727645>
- Matsumoto C, Miedema MD, Ofman P, Gaziano JM, Sesso HD. An expanding knowledge of the mechanisms and effects of alcohol consumption on cardiovascular disease. *J Cardiopulm Rehabil Prev*. 2014;34(3):159-71. doi: 10.1097/HCR.000000000000042. PubMed PMID: 24667667.

- Nagy LE, Ding WX, Cresci G, Saikia P, Shah VH. 2016 Jun. Linking Pathogenic Mechanisms of Alcoholic Liver Disease With Clinical Phenotypes. *Gastroenterology*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=26919968>
- Obradovic T, Meadows GG. 2002 Feb. Chronic ethanol consumption increases plasma leptin levels and alters leptin receptors in the hypothalamus and the perigonadal fat of C57BL/6 mice. *Alcoholism, clinical and experimental research*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=11964566>
- Ohashi K, Shibata R, Murohara T, Ouchi N. Role of anti-inflammatory adipokines in obesity-related diseases. *Trends Endocrinol Metab*. 2014;25(7):348-55. doi: 10.1016/j.tem.2014.03.009. PubMed PMID: 24746980.
- Otaka M, Konishi N, Odashima M, Jin M, Wada I, Matsushashi T, et al. Effect of alcohol consumption on leptin level in serum, adipose tissue, and gastric mucosa. *Dig Dis Sci*. 2007;52(11):3066-9. doi: 10.1007/s10620-006-9635-x. PubMed PMID:17406835.
- Parker, R., Kim, S. J., & Gao, B. (2018). Alcohol, adipose tissue and liver disease: Mechanistic links and clinical considerations. *Nature Reviews Gastroenterology & Hepatology*, 15(1), 50-59. doi:10.1038/nrgastro.2017.116
- Parry CD, Patra J, Rehm J. 2011 Oct. Alcohol consumption and non-communicable diseases: epidemiology and policy implications. *Addiction (Abingdon, England)*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/21819471>
- Peterson JM, Aja S, Wei Z, Wong GW. 2012 Jan 6. CTRP1 protein enhances fatty acid oxidation via AMP-activated protein kinase (AMPK) activation and acetyl-CoA carboxylase (ACC) inhibition. *The Journal of biological chemistry*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=22086915>
- Peterson JM, Seldin MM, Wei Z, Aja S, Wong GW. 2013 Aug 1. CTRP3 attenuates diet-induced hepatic steatosis by regulating triglyceride metabolism. *American journal of physiology. Gastrointestinal and liver physiology*. [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=23744740>
- Peterson JM, Seldin MM, Tan SY, Wong GW. CTRP2 overexpression improves insulin and lipid tolerance in diet-induced obese mice. *PLoS One*. 2014;9(2):e88535. doi: 10.1371/journal.pone.0088535. PubMed PMID: 24586339; PubMed Central PMCID: PMC3930646.

- Peterson JM, Wei Z, Seldin MM, Byerly MS, Aja S, Wong GW. CTRP9 transgenic mice are protected from diet-induced obesity and metabolic dysfunction. *Am J Physiol Regul Integr Comp Physiol*. 2013;305(5):R522-33. doi: 10.1152/ajpregu.00110.2013. PubMed PMID: 23842676; PubMed Central PMCID:PMCPMC3763026.
- Peterson JM, Wei Z, Wong GW. 2010 Dec 17. C1q/TNF-related protein-3 (CTRP3), a novel adipokine that regulates hepatic glucose output. *The Journal of biological chemistry*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=20952387>
- Petersen PS, Wolf RM, Lei X, Peterson JM, Wong GW. Immunomodulatory roles of CTRP3 in endotoxemia and metabolic stress. *Physiol Rep*. 2016;4(5). doi: 10.14814/phy2.12735. PubMed PMID: 26997632; PubMed Central PMCID: PMCPMC4823594.
- Pravdova, E., Macho, L., & Fickova, M. (2009). Alcohol intake modifies leptin, adiponectin and resistin serum levels and their mRNA expressions in adipose tissue of rats. *Endocrine Regulations*, 43(3), 117-125.
 doi:10.4149/endo_2009_03_117
- Rehm, J., Taylor, B., Mohapatra, S., Irving, H., Baliunas, D., Patra, J., & Roerecke, M. (2010). Alcohol as a risk factor for liver cirrhosis: A systematic review and meta-analysis. *Drug and Alcohol Review*, 29(4), 437-445. doi:10.1111/j.1465-3362.2009.00153.x
- Rodriguez S, Lei X, Petersen PS, Tan SY, Little HC, Wong GW. 2016 Oct 1. Loss of CTRP1 disrupts glucose and lipid homeostasis. *American journal of physiology. Endocrinology and metabolism*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=2755298>
- Rojdmark S, Calissendorff J, Brismar K. Alcohol ingestion decreases both diurnal and nocturnal secretion of leptin in healthy individuals. *Clin Endocrinol (Oxf)*. 2001;55(5):639-47. PubMed PMID: 11894976.
- Roth, M. J., Baer, D. J., Albert, P. S., Castonguay, T. W., Dorgan, J. F., Dawsey, S. M., . . . Taylor, P. R. (2003). Relationship between serum leptin levels and alcohol consumption in a controlled feeding and alcohol ingestion study. *Journal of the National Cancer Institute*, 95(22), 1722-1725. Retrieved from
<https://www.ncbi.nlm.nih.gov/pubmed/14625264>

- Sanyal AJ. Novel therapeutic targets for steatohepatitis. *Clin Res Hepatol Gastroenterol.* 2015;39 Suppl 1:S46-50. doi: 10.1016/j.clinre.2015.05.012. PubMed PMID: 26160474.
- Scherer PE, Williams S, Fogliano M, Baldini G, Lodish HF. A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem.* 1995;270(45):26746-9. PubMed PMID: 7592907.
- Schmid A, Kopp A, Aslanidis C, Wabitsch M, Muller M, Schaffler A. Regulation and function of C1Q/TNF-related protein-5 (CTRP-5) in the context of adipocyte biology. *Exp Clin Endocrinol Diabetes.* 2013;121(5):310-7. doi: 10.1055/s-0032-1333299. PubMed PMID: 23430573.
- Schmid A, Kopp A, Hanses F, Bala M, Muller M, Schaffler A. The novel adipokine C1q/TNF-related protein-3 is expressed in human adipocytes and regulated by metabolic and infection-related parameters. *Exp Clin Endocrinol Diabetes.* 2012;120(10):611-7. doi: 10.1055/s-0032-1323803. PubMed PMID: 23174996.
- Schmid A, Kopp A, Hanses F, Karrasch T, Schaffler A. C1q/TNF-related protein-3 (CTRP-3) attenuates lipopolysaccharide (LPS)-induced systemic inflammation and adipose tissue Erk-1/-2 phosphorylation in mice in vivo. *Biochem Biophys Res Commun.* 2014;452(1):8-13. doi: 10.1016/j.bbrc.2014.06.054. PubMed PMID:24996172.
- Seldin MM, Tan SY, Wong GW. Metabolic function of the CTRP family of hormones. *Rev Endocr Metab Disord.* 2014;15(2):111-23. doi: 10.1007/s11154-013-9255-7. PubMed PMID: 23963681; PubMed Central PMCID: PMC3931758.
- Shabani, P., Emamgholipour, S., & Doosti, M. (2017). CTRP1 in Liver Disease. *Advances in Clinical Chemistry, 79*, 1-23. doi:10.1016/bs.acc.2016.10.002.
- Shapiro, L., & Scherer, P. E. (1998). The crystal structure of a complement-1q family protein suggests an evolutionary link to tumor necrosis factor. *Current Biology, 8*(6), 335-340. doi:10.1016/S0960-9822(98)70133-2.
- Shoreibah M, Raff E, Bloomer J, Kakati D, Rasheed K, Kuo YF, et al. Alcoholic liver disease presents at advanced stage and progresses faster compared to non-alcoholic fatty liver diseases. *Ann Hepatol.* 2016;15(2):183-9. doi: 10.5604/16652681.1193707. PubMed PMID: 26845595.
- Sierksma, A., Patel, H., Ouchi, N., Kihara, S., Funahashi, T., Heine, R. J., . . . Hendriks, H. F. (2004). Effect of moderate alcohol consumption on adiponectin, tumor necrosis factor-alpha, and insulin sensitivity. *Diabetes Care, 27*(1), 184-189. doi:10.2337/diacare.27.1.184

- Song Z, Zhou Z, Deaciuc I, Chen T, McClain CJ. 2008 Mar. Inhibition of adiponectin production by homocysteine: a potential mechanism for alcoholic liver disease. *Hepatology* (Baltimore, Md.). [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=18167065>
- Steiner JL, Lang CH. 2017 Feb 16. Alcohol, Adipose Tissue and Lipid Dysregulation. *Biomolecules*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=PMC5372728>
- Stickel F, Seitz HK. Alcoholic steatohepatitis. *Best Pract Res Clin Gastroenterol*. 2010;24(5):683-93. doi: 10.1016/j.bpg.2010.07.003. PubMed PMID: 20955970.
- Strbák V, Benický J, Macho L, Jezová D, Nikodémová M. 1998 Oct. Four-week ethanol intake decreases food intake and body weight but does not affect plasma leptin, corticosterone, and insulin levels in pubertal rats. *Metabolism: clinical and experimental*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=9781633>
- Szkudelski T, Bialik I, Szkudelska K. Adipocyte lipolysis, hormonal and metabolic changes in ethanol-drinking rats. *J Anim Physiol Anim Nutr (Berl)*. 2004;88(7-8):251-8. doi: 10.1111/j.1439-0396.2004.00478.x. PubMed PMID: 15274689.
- Tan X, Sun X, Li Q, Zhao Y, Zhong W, Jia W, McClain CJ, Zhou Z. 2012 Oct. Leptin deficiency contributes to the pathogenesis of alcoholic fatty liver disease in mice. *The American journal of pathology*. [accessed 2017 Dec 4].
<https://www.ncbi.nlm.nih.gov/pubmed/?term=PMC3463622>
- Tang, H., Sebastian, B. M., Axhemi, A., Chen, X., Hillian, A. D., Jacobsen, D. W., & Nagy, L. E. (2012). Ethanol-induced oxidative stress via the CYP2E1 pathway disrupts adiponectin secretion from adipocytes. *Alcoholism Clinical and Experimental Research*, 36(2), 214-222. doi:10.1111/j.1530-0277.2011.01607.x
- Tietge, U. J., Boker, K. H., Manns, M. P., & Bahr, M. J. (2004). Elevated circulating adiponectin levels in liver cirrhosis are associated with reduced liver function and altered hepatic hemodynamics. *American Journal of Physiology*. doi:10.1152/ajpendo.00494.2003
- Wei Z, Peterson JM, Wong GW. Metabolic regulation by C1q/TNF-related protein-13 (CTRP13): activation OF AMP-activated protein kinase and suppression of fatty acid-induced JNK signaling. *J Biol Chem*. 2011;286(18):15652-65. doi: 10.1074/jbc.M110.201087. PubMed PMID: 21378161; PubMed Central PMCID: PMC3091174.
- Wei Z, Seldin MM, Natarajan N, Djemal DC, Peterson JM, Wong GW. C1q/tumor

- necrosis factor-related protein 11 (CTRP11), a novel adipose stroma-derived regulator of adipogenesis. *J Biol Chem.* 2013;288(15):10214-29. doi: 10.1074/jbc.M113.458711.
- Wong GW, Krawczyk SA, Kitidis-Mitrokostas C, Ge G, Spooner E, Hug C, et al. Identification and characterization of CTRP9, a novel secreted glycoprotein, from adipose tissue that reduces serum glucose in mice and forms heterotrimers with adiponectin. *FASEB J.* 2009;23(1):241-58. doi: 10.1096/fj.08-114991. PubMed PMID: 18787108; PubMed Central PMCID: PMCPMC2626616.
- Wolf RM, Lei X, Yang ZC, Nyandjo M, Tan SY, Wong GW. CTRP3 deficiency reduces liver size and alters IL-6 and TGFbeta levels in obese mice. *Am J Physiol Endocrinol Metab.* 2016;310(5):E332-45. doi: 10.1152/ajpendo.00248.2015. PubMed PMID: 26670485; PubMed Central PMCID: PMCPMC4773650.
- Wong GW, Wang J, Hug C, Tsao TS, Lodish HF. 2004 Jul 13. A family of Acrp30/adiponectin structural and functional paralogs. *Proceedings of the National Academy of Sciences of the United States of America.* [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=PMC478567>
- Wong, G. W., Krawczyk, S. A., Kitidis-Mitrokostas, C., Revett, T., Gimeno, R., & Lodish, H. F. (2008). Molecular, biochemical and functional characterizations of C1q/TNF family members: Adipose-tissue-selective expression patterns, regulation by PPAR- γ agonist, cysteine-mediated oligomerizations, combinatorial associations and metabolic functions. *Biochemical Journal*, 416(2), 161-177. doi:10.1042/BJ20081240
- Xu A, Wang Y, Keshaw H, Xu LY, Lam KS, Cooper GJ. 2003 Jul. The fat-derived hormone adiponectin alleviates alcoholic and nonalcoholic fatty liver diseases in mice. *The Journal of clinical investigation.* [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=PMC162288>
- You M, Considine RV, Leone TC, Kelly DP, Crabb DW. 2005 Sep. Role of adiponectin in the protective action of dietary saturated fat against alcoholic fatty liver in mice. *Hepatology (Baltimore, Md.).* [accessed 2017 Dec 4]. <https://www.ncbi.nlm.nih.gov/pubmed/?term=PMC1398076>
- Yoo MG, Kim HJ, Jang HB, Lee HJ, Park SI. The Association between Alcohol Consumption and beta-Cell Function and Insulin Sensitivity in Korean Population. *Int J Environ Res Public Health.* 2016;13(11). doi: 10.3390/ijerph13111133. PubMed PMID: 27854254; PubMed Central PMCID: PMCPMC5129343.

- Yu, H., Li, S., Cao, M., Jiang, X., Feng, L., Zhao, J., & Gao, L. (2010). Effects of chronic ethanol consumption on levels of adipokines in visceral adipose tissues and sera of rats. *Acta Pharmacologica Sinica*, 31(4), 461-469. doi:10.1038/aps.2010.12
- Zhang, J., Zhang, B., Cheng, Y., & Xu, J. (2017). Low serum CTRP3 levels are associated with nonalcoholic fatty liver disease in patients with type 2 diabetes mellitus. *Cytokine*. doi:10.1016/j.cyto.2017.10.023.

VITA

ASHLEY R. DEGROAT

- Personal Data: Date of Birth: October 15, 1993
Place of Birth: Kingsport, Tennessee
Marital Status: Single
- Education: Public Schools, Johnson City, Tennessee

B.S. Biological Sciences, East Tennessee State University,
Johnson City, Tennessee 2016

M.S. Biology, with a concentration in Biomedical Sciences,
East Tennessee State University, Johnson City, Tennessee
2018
- Professional Experience: Undergraduate Research Lab Assistant; Johnson City,
Tennessee, 2012-2016

Graduate Assistant, East Tennessee State University,
College of Public Health, 2016-2018
- Publications: DeGroat, Ashley R. and Peterson, Jonathan M. (2017).
“The Effect of Alcohol Consumption on Adipokine
Secretion”. JOUR. *The FASEB Journal* 31 (1_supplement).
Federation of American Societies for Experimental
Biology: 887.12-887.12.
doi: 10.1096/fasebj.31.1_supplement.887.12.

DeGroat, Ashley R., Clark, William A., Hagood, Kendra L.,
and Peterson, Jonathan M. (2017). “CTRP3 Alters Lipid
Profile in Response to Ethanol Feeding”. JOUR. *The
FASEB Journal* 31 (1_supplement). Federation of
American Societies for Experimental Biology: 887.11-
887.1
doi: 10.1096/fasebj.31.1_supplement.887.11
- Honors and Awards: Second Place Poster Presentation, Biomedical and
Health Sciences, Appalachian Student Research Forum.
“The Effect of Alcohol Consumption on Adipokine
Secretion”. East Tennessee State University, Johnson
City, TN.