

# A major interspecies difference in the ionic selectivity of megakaryocyte $\text{Ca}^{2+}$ -activated channels sensitive to the TMEM16F inhibitor CaCCinh-A01

Kirk A. Taylor & Martyn P. Mahaut-Smith

To cite this article: Kirk A. Taylor & Martyn P. Mahaut-Smith (2019) A major interspecies difference in the ionic selectivity of megakaryocyte  $\text{Ca}^{2+}$ -activated channels sensitive to the TMEM16F inhibitor CaCCinh-A01, Platelets, 30:8, 962-966, DOI: [10.1080/09537104.2019.1595560](https://doi.org/10.1080/09537104.2019.1595560)

To link to this article: <https://doi.org/10.1080/09537104.2019.1595560>



© 2019 The Author(s). Published with license by Taylor & Francis Group, LLC.



Published online: 22 Apr 2019.



Submit your article to this journal [↗](#)



Article views: 1090



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

## PLENARY PAPER AND SHORT COMMUNICATION



# A major interspecies difference in the ionic selectivity of megakaryocyte $\text{Ca}^{2+}$ -activated channels sensitive to the TMEM16F inhibitor CaCCinh-A01

Kirk A. Taylor <sup>1,2</sup> & Martyn P. Mahaut-Smith<sup>1</sup>

<sup>1</sup>Department of Molecular and Cell Biology, University of Leicester, Leicester, UK and <sup>2</sup>National Heart and Lung Institute, Cardio-respiratory Section, Imperial College London, London, UK

## Abstract

TMEM16F is a surface membrane protein critical for platelet procoagulant activity, which exhibits both phospholipid scramblase and ion channel activities following sustained elevation of cytosolic  $\text{Ca}^{2+}$ . The extent to which the ionic permeability of TMEM16F is important for platelet scramblase responses remains controversial. To date, only one study has reported the electrophysiological properties of TMEM16F in cells of platelet/megakaryocyte lineage, which observed cation-selectivity within excised patch recordings from murine marrow-derived megakaryocytes. This contrasts with reports using whole-cell recordings that describe this channel as displaying either selectivity for anions or being relatively non-selective amongst the major physiological monovalent ions.

We have studied TMEM16F expression and channel activity in primary rat and mouse megakaryocytes and the human erythroleukemic (HEL) cell line that exhibits megakaryocytic surface markers. Immunocytochemical analysis was consistent with surface TMEM16F expression in cells from all three species. Whole-cell recordings in the absence of  $\text{K}^+$ -selective currents revealed an outwardly rectifying conductance activated by a high intracellular  $\text{Ca}^{2+}$  concentration in all three species. These currents appeared after 5–6 minutes and were blocked by CaCC<sub>inh</sub>-A01, properties typical of TMEM16F. Ion substitution experiments showed that the underlying conductance was predominantly  $\text{Cl}^-$ -permeable in rat megakaryocytes and HEL cells, yet non-selective between monovalent anions and cations in mouse megakaryocytes. In conclusion, the present study further highlights the difference in ionic selectivity of TMEM16F in platelet lineage cells of the mouse compared to other mammalian species. This provides additional support for the ionic “leak” hypothesis that the scramblase activity of TMEM16F does not rely upon its ability to conduct ions of a specific type.

## Introduction

Procoagulant activity resulting from exposure of anionic membrane phospholipids is critical for thrombosis and haemostasis. Although the underlying mechanisms of this lipid redistribution process are incompletely understood, a  $\text{Ca}^{2+}$ -activated phospholipid scramblase is known to be important [1–3]. TMEM16F is a ubiquitously expressed  $\text{Ca}^{2+}$ -dependent ion channel and phospholipid scramblase [2,4]. Missense mutations of the TMEM16F gene occur in patients with Scott syndrome, a rare bleeding diathesis characterised by defective  $\text{Ca}^{2+}$ -dependent phospholipid scrambling [5–7]. Furthermore, this disease is phenocopied in TMEM16F<sup>-/-</sup> mice [3,8].

The nature of the ionic permeability of TMEM16F channels is controversial. Studies of native cells and expression systems con-

## Keywords

Calcium, megakaryocyte, phospholipid scrambling, platelet, Scott syndrome, TMEM16F

## History

Received 16 November 2018  
Revised 12 February 2019  
Accepted 4 March 2019  
Published online 8 April 2019

clude that human and murine TMEM16F channels are predominantly anion-permeable [4,9–11]. However, multiple reports demonstrate that heterologously expressed TMEM16F displays significant permeability to monovalent cations ( $P_{\text{Na}}/P_{\text{Cl}}$ : 0.3 [12] 0.5 [13] and 1.38 [14]). Such studies have principally relied upon the whole-cell patch clamp configuration. Contrastingly, using excised inside-out membrane patch recordings, one group has reported that murine TMEM16F forms non-selective cation channels with a greater permeability to  $\text{Ca}^{2+}$  than monovalent cations both endogenously in the native megakaryocyte (MK) and in expression systems [8,15]. This raises the question of whether the properties of TMEM16F are influenced by patch excision or by the environment of a specific cell type. MKs are responsible for generating all proteins within their anuclear product and exhibit functional platelet responses [16,17], thus are frequently used as a surrogate for electrophysiological studies of the tiny, fragile platelet.

We have employed whole-cell patch clamp recordings to investigate the biophysical and pharmacological properties of  $\text{Ca}^{2+}$ -activated TMEM16F channels in freshly isolated primary rat and mouse MKs and the human erythroleukemia (HEL) cell line. HEL cells express a number of megakaryocytic glycoproteins [18], have been used for the study of native MK ion channels and express TMEM16F transcripts [19].

Color versions of one or more of the figures in the article can be found online at [www.tandfonline.com/iplt](http://www.tandfonline.com/iplt).

Correspondence: Martyn P. Mahaut-Smith, Department of Molecular and Cell Biology, University of Leicester, Henry Wellcome Building, Lancaster Road, Leicester, LE1 7RH. E-mail: [mpms1@le.ac.uk](mailto:mpms1@le.ac.uk)  
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Methods

**Ethics:** Ethical Approval for this study was granted by the University of Leicester College of Life Sciences Research Ethics Committee for Human Biology (non-NHS).

**Primary MK isolation:** MKs were prepared as previously described [20,21] from adult Wistar rats and C57bl/6 mice following euthanasia in accordance with the UK Animals (Scientific Procedures) Act 1986.

**Cell culture:** HELs (ATCC, Middlesex, UK) were cultured in RPMI 1640 (Invitrogen, Paisley, UK) supplemented with foetal calf serum (10%; Invitrogen) and penicillin/streptomycin (250U/mL; Invitrogen).


**Immunocytochemistry:** Cell suspensions were prepared as described previously [22]. Samples were incubated with anti-*TMEM16F* primary (2 $\mu$ g/ml; Santa Cruz, California, USA) and alexafluor647 (AF647)-conjugated secondary antibodies (1:1000; Invitrogen). Fluorescence was assessed with an Olympus

FV1000 confocal microscope (635nm excitation, 650-750nm emission; Olympus, UK).

**Electrophysiology:** Whole-cell patch clamp recordings were conducted as described previously with  $\geq 70\%$  series resistance compensation and *a priori* liquid junction potential correction [23]. Bath solutions contained 150mM NaCl, 1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 10mM glucose, 10mM HEPES (pH 7.35; NaOH). Where indicated, [Cl<sup>-</sup>]<sub>o</sub> and/or [Na<sup>+</sup>]<sub>o</sub> were reduced by equimolar substitution with gluconate<sup>-</sup> or NMDG<sup>+</sup>, respectively. Internal solutions contained 150mM NaCl, 1mM MgCl<sub>2</sub>, 10mM glucose, 10mM HEPES, 50 $\mu$ M Na<sub>2</sub>-GTP, 1mM EGTA (pH 7.2; NaOH). [Ca<sup>2+</sup>]<sub>i</sub> was set at  $\approx 5$ nM (1mM EGTA, no added Ca<sup>2+</sup>) or 100 $\mu$ M (by addition of CaCl<sub>2</sub>), calculated using Maxchelator (<http://web.stanford.edu/~cpatton/webmaxcS.htm>). The effect of CaCC<sub>inh</sub>-A01 (A01; Merck, Watford, UK) was compared with vehicle (DMSO) control. Statistical analysis was by two-way ANOVA (Prism7, GraphPad Software Inc., CA, USA).

Figure 1. **Detection of Ca<sup>2+</sup>-activated and A01-sensitive *TMEM16F*-like currents in HEL cells and rat and mouse MKs.**

A) *TMEM16F* expression by permeabilised HEL cells (left), primary rat (centre) and mouse (right) MKs assessed by immunocytochemistry with a primary antibody raised against an intracellular epitope of *TMEM16F*. Strong fluorescence was observed at the periphery of cells treated with primary ( $\alpha$ *TMEM16F*) and secondary (AF647) antibodies, whilst no fluorescence was detected in secondary antibody only controls. B-D) whole-cell patch clamp recordings of Ca<sup>2+</sup>-activated currents from HEL cells (B), rat (C) and mouse (D) MKs. Intracellular and bath solutions contained 150mM NaCl and were K<sup>+</sup>-free. [Ca<sup>2+</sup>]<sub>i</sub> was set at either  $\approx 5$ nM (1mM EGTA) or 100 $\mu$ M as indicated. After 10 minutes in the whole-cell mode, currents were recorded in response to voltage steps of 1s duration (-120 to +120mV, 20mV increments) in the presence of vehicle control (0.04% DMSO) or the *TMEM16F* antagonist CaCC<sub>inh</sub>-A01 (A01; 20 $\mu$ M). Representative traces are shown for control or A01-treated MKs in the presence of intracellular EGTA or 100 $\mu$ M [Ca<sup>2+</sup>]<sub>i</sub>. Summary current density-voltage relationship data are shown in the right hand panels for EGTA-subtracted currents under control (solid line) or A01-treated (dashed line) conditions. For immunocytochemistry experiments, scale bars represent 10 $\mu$ m. Data are representative of a minimum of three independent experiments for each condition. \*, \*\* and \*\*\* denote  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively.




## Results

TMEM16F expression in HELs and rat and mouse MKs was assessed by immunocytochemistry with an antibody previously used in mouse dendritic cells [9]. Fluorescence was detected in primary MKs from both species and HELs, with a pattern indicating strong surface expression and no signal from secondary antibody-only controls (Figure 1A).

Previous electrophysiological studies demonstrate that primary MKs and HELs display robust  $K^+$  currents activated by depolarisation and/or intracellular  $Ca^{2+}$  [23–26]. Within whole-cell recordings from other cell types, TMEM16F channels typically activate in response to sustained (5–6 min) elevation of high  $[Ca^{2+}]_i$  ( $EC_{50}$  of  $\approx 100\mu M$ ) [12,14]. Therefore, in our study,  $K^+$ -free bath and pipette salines were used to abolish  $K^+$  currents prior to the predicted activation of TMEM16F. Voltage ramps from  $-120$  to  $+120$  mV were applied at regular intervals and  $[Ca^{2+}]_i$  set at either  $\approx 5$  nM or  $100\mu M$ . At  $100\mu M$   $[Ca^{2+}]_i$ , outwardly rectifying currents developed at similar timepoints in cells from all species:  $439 \pm 25$  s ( $n = 5$ ) in HELs,  $418 \pm 18$  s ( $n = 5$ ) in rat MKs and  $368 \pm 20$  s in mouse MKs ( $n = 4$ ). This current was not

observed at  $\approx 5$  nM  $[Ca^{2+}]_i$  and its activation time course is consistent with reports of TMEM16F [12,14]. After 10 minutes, 1 s voltage steps were applied across the range  $-120$  to  $+120$  mV, as previously used for further characterisation of TMEM16F [8,12]. In the rat MK,  $100\mu M$   $[Ca^{2+}]_i$  induced large outwardly rectifying currents that reversed ( $E_{rev}$ ) at  $-3.3 \pm 0.5$  mV (Figure 1C) and were blocked by the TMEM16F antagonist A01 [8,9,27,28] ( $39.4 \pm 3.8$  vs  $-3.6 \pm 0.7$  pA/pF at  $+120$  mV;  $P < 0.001$ ,  $n = 5$ , Figure 1C). The conductance in HELs and mouse MKs possessed similar properties, although the current amplitudes and extent of outward rectification were smaller in these species compared to the rat. The current amplitude at  $+120$  mV compared to  $-120$  mV was  $18.8 \pm 3.1$  ( $n = 5$ ),  $43.9 \pm 10.3$  ( $n = 5$ ) and  $13.7 \pm 4.7$  ( $n = 4$ ) in HELs, rat and mouse MKs, respectively. In HELs the  $Ca^{2+}$ -activated currents reversed at  $-1.7 \pm 0.3$  mV and were reduced from  $14.1 \pm 2.3$  to  $2.0 \pm 0.4$  pA/pF at  $+120$  mV by A01 (Figure 1B). For mouse MKs,  $Ca^{2+}$ -induced currents reversed close to 0 mV and were reduced from  $9.0 \pm 3.8$  to  $-1.0 \pm 0.8$  pA/pF at  $120$  mV by A01;  $P < 0.01$ ,  $n = 4$  [control] and  $n = 3$ , Figure 1D).



**Figure 2. Comparison of ionic selectivity between human, rat and mouse megakaryocytic  $Ca^{2+}$ -activated, A01 sensitive TMEM16F-like channels.** Whole-cell patch clamp recordings of  $Ca^{2+}$ -activated currents from HEL cells (left), rat (centre) and mouse (right) MKs under conditions designed to isolate TMEM16F channel activity. Intracellular solutions were  $K^+$ -free (150 mM NaCl), and membrane currents were assessed after 10 mins in the whole cell configuration with  $[Ca^{2+}]_i$  set at either  $\approx 5$  nM (1 mM EGTA, not shown) or  $100\mu M$ . Currents were recorded in response to voltage steps of 1 s duration ( $-120$  to  $+120$  mV, 20 mV increments) in the presence of vehicle control (0.04% DMSO) or the TMEM16F antagonist CaCCinh-A01 (A01;  $20\mu M$ ). For experiments in panels A–B extracellular  $Cl^-$  was substituted by equimolar gluconate $^-$  and in panels C–D, extracellular  $Na^+$  and  $Cl^-$  were substituted by NMDG $^+$  and gluconate, respectively. B, D) Summary current density-voltage relationship data for  $Ca^{2+}$ -activated currents under vehicle control (solid line) or A01-treated (dashed line) conditions. Data are representative of a minimum of three independent experiments for each condition. \*, \*\* and \*\*\* denote  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively.



Under the ionic conditions of Figure 1, the equilibrium potential for both  $\text{Na}^+$  and  $\text{Cl}^-$  is 0mV and thus  $\text{Ca}^{2+}$ -activated currents may reflect movement of cations and/or anions. Following reduction of external chloride (gluconate substitution),  $100\mu\text{M}$   $[\text{Ca}^{2+}]_i$  evoked large A01-sensitive currents in each species (Figure 2A). Strikingly,  $E_{\text{rev}}$  shifted in HELs ( $+37.2 \pm 2.1\text{mV}$ ) and rat MKs ( $+33.5 \pm 2.7\text{mV}$ ), whereas mouse MK currents continued to reverse close to 0mV (Figure 2B). Under low  $[\text{Cl}^-]_o$  conditions  $E_{\text{Na}} = 0\text{mV}$  and  $E_{\text{Cl}} = +85.7\text{mV}$ , indicating that the underlying conductance displays greater permeability to  $\text{Cl}^-$  compared to other ions in HELs and rat MK, but not in the mouse MK. A lower, yet significant, permeability to cations or large anions may explain why the shift of  $E_{\text{rev}}$  was less than that expected for a perfectly  $\text{Cl}^-$ -selective conductance, as reported previously for  $\text{Cl}^-$  channels in human platelets [29,30].

In low  $\text{Cl}^-$  external saline, substitution of  $[\text{Na}^+]_o$  with the large cation NMDG<sup>+</sup> failed to alter  $E_{\text{rev}}$  in mouse MKs but shifted this value to a slightly more positive potential in rat MKs ( $+41.9 \pm 3.4\text{mV}$ ) and HELs ( $+40.4 \pm 3.9\text{mV}$ ; Figure 2C,D). These data further suggest a major difference in the ionic selectivity of the TMEM16F-like conductance in megakaryocytic cells from mouse compared to rat or human. They also indicate that the mouse channel is highly non-selective amongst the major monovalent ions used in this study.

## Discussion

The ionic selectivity of  $\text{Ca}^{2+}$ -activated TMEM16F channels has been variably reported as anionic [4,9–12], cationic [8,15] and non-selective between monovalent anions and cations [13,14]. Here, we compared the ionic selectivity of the  $\text{Ca}^{2+}$ -activated conductances of primary megakaryocytes from rat and mouse, and a human megakaryocyte cell line, under conditions previously used to characterise TMEM16F. We observed that endogenous TMEM16F-like conductances are relatively selective for  $\text{Cl}^-$  in rat MKs and a human megakaryocytic cell line but are non-selective in mouse MKs (Figure 2). Although cation-selective TMEM16F channels have been reported in the mouse MK [8] and following heterologous expression of the murine clone in oocytes and HEK cells [8,15], a study of mouse dendritic cells concluded that this conductance is more permeable to  $\text{Cl}^-$  than cations [9]. Plausible explanations for this difference include cell-specific expression of alternative splice variants [13,31] or an effect of the cell environment on channel properties. A further possibility is that the channel may behave differently in excised inside-out patches, which have been exclusively used in studies that conclude TMEM16F is cation-selective [8,15], compared to whole-cell recordings used for most other studies. Interestingly, in excised patch recordings, TMEM16F was also observed to show greater permeability to  $\text{Ca}^{2+}$  than monovalent cations, which may play a role in scramblase activation or inactivation [8,15]. We were unable to test the relative  $\text{Ca}^{2+}$  permeability of the channel due to poor viability of whole-cell recordings in low or high  $[\text{Ca}^{2+}]_o$  solutions. Nevertheless, it seems unlikely that  $\text{Ca}^{2+}$  entry through TMEM16F is crucial for its own activation since this requires sustained high  $[\text{Ca}^{2+}]_i$  levels. It has also been proposed that  $\text{Cl}^-$ -selective TMEM16F channels cause membrane hyperpolarisation, thereby enhancing  $\text{Ca}^{2+}$  influx through an increased driving force [28]. However, activated platelets have a negative membrane potential ( $-45\text{mV}$  to  $-80\text{mV}$ ) set by voltage-gated or  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels [23,30,32] which drives  $\text{Ca}^{2+}$  influx primarily via Orai1, P2X1 and TRPC6 [23,30,33]. Since the platelet  $\text{Cl}^-$  equilibrium potential has been estimated to sit at  $\approx -35\text{mV}$  [30], activation of either a  $\text{Cl}^-$ -selective or non-selective channel ( $E_{\text{rev}} \approx 0\text{mV}$ ) would depolarise rather than hyperpolarise the cell. This argues against a role for TMEM16F in promoting  $\text{Ca}^{2+}$  entry via regulation of the membrane potential.

Although we demonstrate an interspecies difference of ion selectivity for MK TMEM16F-like channels, further work is required to understand the functional significance of the resultant ionic movements for lipid scrambling and/or other functional events in this cell type. Furthermore, the possible relevance of expression of other members of this scramblase/channel family should also be examined with knock-down studies. It is worth noting however that a transcriptomic analysis of purified platelets detected <2% expression of other TMEM16 family members compared to TMEM16F [19]. Furthermore, the same study detected only TMEM16K in addition to TMEM16F in HEL cells, and TMEM16K is inhibited at the high intracellular  $\text{Ca}^{2+}$  concentrations used in the present work [4]. A study of HEK293 cells reported that TMEM16F-dependent currents and phosphatidylserine exposure occur contemporaneously, and that preventing net ionic movement by clamping the membrane voltage to zero does not affect phosphatidylserine scrambling [14]. Furthermore, despite the species difference in ionic selectivity between mouse and human MK TMEM16F, it is worth noting the similarities between the altered scramblase activity of TMEM16F<sup>-/-</sup> mice [3,8] and Scott Syndrome patients [1]. This suggests that the ability to conduct specific ionic species, or an influence on membrane potential, are unlikely to be crucial determinants of TMEM16F-mediated lipid scrambling and thus platelet procoagulant activity. Future studies should therefore directly assess the relationship between platelet and/or megakaryocyte lipid scrambling and TMEM16F ionic selectivity. Overall, the present work provides further support for the “leak” hypothesis of ionic movement through TMEM16F, in which ions move because of phospholipid translocation rather than being critical for lipid redistribution [14].

## Authorship Contributions

KAT and MPMS designed the experiments, analysed data and wrote the manuscript. KAT performed experiments. MPMS secured funding.

## Declaration of Interest

The authors report no declarations of interest.

## Funding

This work was supported by the Medical Research Council [DTG].

## ORCID

Kirk A. Taylor  <http://orcid.org/0000-0002-4599-7727>

## References

- Zwaal RF, Schroit AJ. Pathophysiologic implications of membrane phospholipid asymmetry in blood cells. *Blood* 1997;89(4):1121–1132.
- Suzuki J, Umeda M, Sims PJ, Nagata S. Calcium-dependent phospholipid scrambling by TMEM16F. *Nature* 2010;468(7325):834–838. doi:10.1038/nature09583.
- Fujii T, Sakata A, Nishimura S, Eto K, Nagata S. TMEM16F is required for phosphatidylserine exposure and microparticle release in activated mouse platelets. *Proc Natl Acad Sci U S A* 2015;112(41):12800–12805. doi:10.1073/pnas.1516594112.
- Schreiber R, Uliyakina I, Kongsuphol P, Warth R, Mirza M, Martins JR, Kunzelmann K. Expression and function of epithelial anoctamins. *J Biol Chem* 2010;285(10):7838–7845. doi:10.1074/jbc.M109.065367.
- Castoldi E, Collins PW, Williamson PL, Bevers EM. Compound heterozygosity for 2 novel TMEM16F mutations in a patient with Scott syndrome. *Blood* 2011;117(16):4399–4400. doi:10.1182/blood-2011-01-332502.
- van Kruchten, R, Mattheij NJ, Saunders C, Feijge MA, Swieringa F, Wolfs JL, Collins PW, Heemskerk JW, Bevers EM. Both TMEM16F-dependent and TMEM16F-independent pathways contribute to phosphatidylserine exposure in platelet apoptosis and platelet activation. *Blood* 2013;121(10):1850–1857. doi:10.1182/blood-2012-09-454314.

7. Boisseau P, Bene MC, Besnard T, Pachcek S, Giraud M, Talarmin P, Robillard N, Gourlaouen MA, Bezieau S, Fouassier M. A new mutation of ANO6 in two familial cases of Scott syndrome. *Br J Haematol* 2018;180(5):750–752. doi:10.1111/bjh.14439.
8. Yang H, Kim A, David T, Palmer D, Jin T, Tien J, Huang F, Cheng T, Coughlin SR, Jan YN, et al. TMEM16F forms a Ca<sup>2+</sup>-activated cation channel required for lipid scrambling in platelets during blood coagulation. *Cell* 2012;151(1):111–122. doi:10.1016/j.cell.2012.07.036.
9. Sztayn K, Schmid E, Nurbaeva MK, Yang W, Münzer P, Kunzelmann K, Lang F, Shumilina E. Expression and functional significance of the Ca<sup>2+</sup>-activated Cl<sup>-</sup> channel ANO6 in dendritic cells. *Cell Physiol Biochem* 2012;30(5):1319–1332. doi:10.1159/000343321.
10. Martins JR, Faria D, Kongsuphol P, Reisch B, Schreiber R, Kunzelmann K. Anoctamin 6 is an essential component of the outwardly rectifying chloride channel. *Proc Natl Acad Sci U S A* 2011;108(44):18168–18172. doi:10.1073/pnas.1108094108.
11. Shimizu T, Iehara T, Sato K, Fujii T, Sakai H, Okada Y. TMEM16F is a component of a Ca<sup>2+</sup>-activated Cl<sup>-</sup> channel but not a volume-sensitive outwardly rectifying Cl<sup>-</sup> channel. *Am J Physiol Cell Physiol* 2013;304(8):C748–59. doi:10.1152/ajpcell.00228.2012.
12. Grubb S, Poulsen KA, Juul CA, Kyed T, Klausen TK, Larsen EH, Hoffmann EK. TMEM16F (Anoctamin 6), an anion channel of delayed Ca<sup>2+</sup> activation. *J Gen Physiol* 2013;141(5):585–600. doi:10.1085/jgp.201210861.
13. Scudieri P, Caci E, Venturini A, Sondo E, Pianigiani G, Marchetti C, Ravazzolo R, Pagani F, Galletta LJV. Ion channel and lipid scramblase activity associated with expression of TMEM16F/ANO6 isoforms. *The Journal of Physiology* 2015;593(17):3829–3848. doi:10.1113/JP270691.
14. Yu K, Whitlock JM, Lee K, Ortlund EA, Cui YY, Hartzell HC. Identification of a lipid scrambling domain in ANO6/TMEM16F. *Elife* 2015;4:e06901. doi:10.7554/eLife.06901.
15. Ye W, Han TW, Nassar LM, Zubia M, Jan YN, Jan LY. Phosphatidylinositol-(4, 5)-bisphosphate regulates calcium gating of small-conductance cation channel TMEM16F. *Proc Natl Acad Sci U S A* 2018;115(7):E1667–E1674. doi:10.1073/pnas.1718728115.
16. Tolhurst G, Vial C, Léon C, Gachet C, Evans RJ, Mahaut-Smith MP. Interplay between P2Y<sub>1</sub>, P2Y<sub>12</sub>, and P2X<sub>1</sub> receptors in the activation of megakaryocyte cation influx currents by ADP: evidence that the primary megakaryocyte represents a fully functional model of platelet P2 receptor signaling. *Blood* 2005;106(5):1644–1651. doi:10.1182/blood-2005-02-0725.
17. Shattil SJ, Leavitt AD. All in the family: primary megakaryocytes for studies of platelet alphaIIb beta3 signaling. *Thromb Haemost* 2001;86(1):259–265.
18. Tabilio A, Rosa JP, Testa U, Kieffer N, Nurden AT, Del Canizo MC, Breton-Gorius J, Vainchenker W. Expression of platelet membrane glycoproteins and alpha-granule proteins by a human erythroleukemia cell line (HEL). *Embo J* 1984;3(2):453–459.
19. Wright JR, Amisten S, Goodall AH, Mahaut-Smith MP. Transcriptomic analysis of the ion channelome of human platelets and megakaryocytic cell lines. *Thromb Haemost* 2016;116(2):272–284. doi:10.1160/TH15-11-0891.
20. Hussain JF, Mahaut-Smith MP. ADP and inositol trisphosphate evoke oscillations of a monovalent cation conductance in rat megakaryocytes. *J Physiol* 1998;511(Pt 3):791–801.
21. Osman S, Taylor KA, Allcock N, Rainbow RD, Mahaut-Smith MP. Detachment of surface membrane invagination systems by cationic amphiphilic drugs. *Sci Rep* 2016;6:18536. doi:10.1038/srep18536.
22. Taylor KA, Wright JR, Vial C, Evans RJ, Mahaut-Smith MP. Amplification of human platelet activation by surface pannexin-1 channels. *J Thromb Haemost* 2014;12(6):987–998. doi:10.1111/jth.12566.
23. McCloskey C, Jones S, Amisten S, Snowden RT, Kaczmarek LK, Erlinge D, Goodall AH, Forsythe ID, Mahaut-Smith MP. Kv1.3 is the exclusive voltage-gated K<sup>+</sup> channel of platelets and megakaryocytes: roles in membrane potential, Ca<sup>2+</sup> signalling and platelet count. *J Physiol* 2010;588(Pt 9):1399–1406. doi:10.1113/jphysiol.2010.188136.
24. Lu X, Fein A, Feinstein MB, O'Rourke FA. Antisense knock out of the inositol 1,3,4,5-tetrakisphosphate receptor GAP1(IP4BP) in the human erythroleukemia cell line leads to the appearance of intermediate conductance K(Ca) channels that hyperpolarize the membrane and enhance calcium influx. *J Gen Physiol* 1999;113(1):81–96.
25. Stoneking CJ, Shivakumar O, Thomas DN, Colledge WH, Mason MJ. Voltage dependence of the Ca<sup>2+</sup>-activated K<sup>+</sup> channel K<sub>Ca</sub>3.1 in human erythroleukemia cells. *Am J Physiol Cell Physiol* 2013;304(9):C858–72. doi:10.1152/ajpcell.00368.2012.
26. Uneyama C, Uneyama H, Akaike N. Cytoplasmic Ca<sup>2+</sup> oscillation in rat megakaryocytes evoked by a novel type of purinoceptor. *J Physiol* 1993;470:731–749.
27. De La Fuente R, Namkung W, Mills A, Verkman AS. Small-molecule screen identifies inhibitors of a human intestinal calcium-activated chloride channel. *Mol Pharmacol* 2008;73(3):758–768. doi:10.1124/mol.107.043208.
28. Harper MT, Poole AW. Chloride channels are necessary for full platelet phosphatidylserine exposure and procoagulant activity. *Cell Death Dis* 2013;4:e969. doi:10.1038/cddis.2013.495.
29. MacKenzie AB, Mahaut-Smith MP. Chloride channels in excised membrane patches from human platelets: effect of intracellular calcium. *Biochim Biophys Acta* 1996;1278(1):131–136.
30. Mahaut-Smith MP. Chloride channels in human platelets: evidence for activation by internal calcium. *J Membr Biol* 1990;118(1):69–75.
31. Acheson K. TMEM16F: function from (iso)form. *J Physiol* 2016;594(11):2785–2786. doi:10.1113/JP271980.
32. Mahaut-Smith MP, Hussain JF, Mason MJ. Depolarization-evoked Ca<sup>2+</sup> release in a non-excitable cell, the rat megakaryocyte. *J Physiol* 1999;515(Pt 2):385–390.
33. Sullivan R, Kunze DL, Kroll MH. Thrombin receptors activate potassium and chloride channels. *Blood* 1996;87(2):648–656.