



Multistep soluble polymer-supported, microwave-assisted synthesis of quinoxalines

Farid Fouad

To cite this article: Farid Fouad (2013) Multistep soluble polymer-supported, microwave-assisted synthesis of quinoxalines, Green Chemistry Letters and Reviews, 6:3, 249-253, DOI: [10.1080/17518253.2013.772245](https://doi.org/10.1080/17518253.2013.772245)

To link to this article: <https://doi.org/10.1080/17518253.2013.772245>



Copyright Farid Fouad



Published online: 12 Mar 2013.



Submit your article to this journal [↗](#)



Article views: 903



View related articles [↗](#)



Citing articles: 3 View citing articles [↗](#)

Multistep soluble polymer-supported, microwave-assisted synthesis of quinoxalines

Farid Fouad*

Department of Chemistry & Biochemistry, Kent State University at East Liverpool, East Liverpool, OH, USA

(Received 3 July 2012; final version received 17 January 2013)

A multistep liquid-phase, polymer-supported, microwave-assisted synthesis of quinoxalines on polyethylene glycol, PEG-4500, is demonstrated. The usage of microwave energy to accelerate organic reactions reduces reaction time from hours (under reflux) to only minutes. The solution-phase chemistry on soluble polyethylene glycol, PEG, support facilitates the use of microwave energy (since PEG easily absorbs microwaves) and avoids the silica gel chromatography purifications of intermediate compounds, supporting parallel, combinatorial, and ultimately automated syntheses.

Keywords: quinoxalines; microwave organic synthesis; liquid-phase synthesis; polymer-supported synthesis

Introduction

Liquid-phase synthesis on soluble polymer supports is a growing field in combinatorial chemistry [1–3]. Besides the advantages of solid-phase chemistry (use of excess reagents and simple isolation and purification), it has the advantages of solution chemistry conventional analysis. Polyethylene glycol, PEG, is successful as a polymer support and as a delivery vehicle for many low-molecular-weight drugs, polypeptides, and oligonucleotides [4–5]. Conjugation to PEG increases water solubility and retention time and protects drugs against degrading enzymes. Using microwave energy and water-soluble substrate-PEG conjugates permit the use of water-based organic transformations.

Nitrogen, oxygen, and sulfur containing heterocyclic compounds are key building blocks used to develop medicinally and biologically active compounds. They are predominant among all types of pharmaceuticals and agrochemical products. Heterocyclic compounds also have many industrial uses as components in dyes, antioxidants, polymers, bases, and ligands [6]. Quinoxaline nitrogen heterocycles exhibit a broad spectrum of activities, including antimicrobial, antidiabetic, and antitumor activities [7]. Quinoxalines are also used in materials science as dyes, efficient electroluminescent materials, and organic semiconductors [8].

Enormous numbers of classical synthetic methods of quinoxalines are reported [9]. In a combinatorial chemistry approach, two libraries of quinoxalines were successfully produced via solid-phase syntheses on Rink amide and Wang resins [10,11]. Ecofriendly syntheses of quinoxalines utilizing Nano-BF₃·SiO₂

catalyst [12], zirconium (IV) oxide chloride octahydrate catalyst [13], and catalytic magnetic Fe₃O₄ nanoparticles in water [14] were reported. Green syntheses of quinoxalines, using PEG-400 [15], PEG-600/water [16], and PEG-400/microwave irradiation [17,18] were reported. Recently, an efficient ecofriendly synthesis of biologically active quinoxalines using easily recyclable TiO₂/SO₄²⁻ and microwave irradiation was reported [19–21].

Results and discussion

The present work demonstrates an efficient synthesis of quinoxalines on a PEG-4500 soluble polymer support under microwave irradiation (Figure 1). For simplicity, commercially available 3,4-dinitrobenzoic acid was chosen as the starting material because it possesses a carboxyl group that can be hooked to the hydroxyl termini of PEG-4500, and two adjacent *nitro* groups that can be reduced to the desired 1,2-diamine. Esterification of 2,3-dinitrobenzoic acid to PEG-4500 under open-vessel microwave conditions, afforded the PEG-Substrate conjugate **1** in only 20 minutes. Catalytic hydrogenation reduction of the dinitro, conjugate **1** did not form the desired diamine **2**, instead, partial reduction products were observed by NMR. Reduction of the two nitro groups was successful using excess ammonium formate and catalytic 10% palladium/carbon under open-vessel microwave irradiation for only 7 minutes. The reaction of the 1,2-diamine, **2** and α -diketones was carried out in 9:1 methanol/acetic acid under microwave irradiation in a closed vessel for only 10 minutes. It was easy to monitor this sequence of reactions, using ¹H NMR by tracking the position of the aromatic

*Email: ffouad@kent.edu




Figure 1. Liquid phase synthesis of quinoxalines.

protons that change from the PEG-dinitrosubstrate, conjugate **1** to the PEG-diamine, substrate **2** and then to respective PEG-quinoxalines **3**. Finally, cleavage of the quinoxaline product from the PEG support was successful using sodium methoxide in methanol. The suppression of reaction time from hours to minutes and the elimination of chromatography purification, minimizing the quantities of organic solvents used, make this method more environment friendly than conventional syntheses and opens the door to parallel and combinatorial syntheses.

Experimental

Chemicals were purchased from various suppliers, including Sigma Aldrich and TCI America. ^1H NMR spectra were recorded using a Bruker Avance 400 MHz spectrometer with tetramethylsilane as the internal standard. Mass spectra were recorded using a Bruker Esquire-3000 ESI-MS. HR mass spectra were measured at Mass Spectrometry & Proteomics Facility, Ohio State University at Columbus, Ohio. The microwave irradiation experiments were undertaken in a focused microwave CEM Discover reactor fitted with a rotational system and an IR detector of temperature.

Synthesis of conjugate 1

A 100 mL round-bottomed flask fitted with a reflux condenser was charged with PEG-4500 (5.00 g, 1.11 mmol), 3,4-dinitrobenzoic acid (1.40 g, 6.66 mmol), N,N' -Dicyclohexylcarbodiimide (1.37 g, 6.66 mmol),

and 4-dimethylaminopyridine (0.22 mmol, 27 mg) and dry dichloromethane (40 mL). The flask was subjected to microwave irradiation (open-vessel mode, 80 w) for 15 minutes, cooled at 0°C for 30 minutes and then filtered to remove dicyclohexyl urea. Cold ether (400 mL) was added to precipitate the PEG-substrate, conjugate **1** as a yellowish powder. The precipitate was isolated by filtration and washed with cold ether (yield, 4.86 g).

Synthesis of conjugate 2

To a solution of conjugate **1** (1.30 g, 0.29 mmol) and ammonium formate (366 mg, 5.80 mmol) in methanol (20 mL) was added 10% palladium/carbon (184 mg, 0.087 mmol). The mixture was subjected to microwave irradiation (open-vessel mode, 80 w, 7 minutes). It was filtered over a bed of cellite. Methanol was removed under reduced pressure, and the residue was dissolved in toluene (10 mL) and filtered to remove any insoluble ammonium formate. Cold ether was added to precipitate the PEG-diaminosubstrate, conjugate **2** as a white precipitate (yield 0.98 g)

General procedure for the synthesis of conjugates 3, and cleavage of quinoxalines 4

A CEM designed 10 mL pressure-rated vial was charged with PEG-diaminosubstrate, **2** (300 mg, 0.17 mmol), the respective α -diketones (0.35 mmol), and 9:1 MeOH/AcOH (3 mL). The mixture was irradiated in a CEM Discover Focused Synthesizer (150 w, 130°C , 200 psi, 8 minutes). The mixture was

cooled to room temperature by passing compressed air through the microwave cavity for 2 minutes. It was poured into cold ether (40 mL) and the formed precipitate filtered, washed with ether, and air dried to give **3**. A solution of **3** in NaOMe/MeOH (1 M, 2 mL) was left to stand at room temperature for overnight. Cleaved quinoxalines **4** were precipitated, filtered, and washed with methanol. They were purified by passing through a short column eluted with 10% ethyl acetate/dichloromethane (Table 1).

Methyl-2,3-diphenylquinoxaline-6-carboxylate, 4a [22]

¹H NMR (400 MHz, CDCl₃): δ = 8.92 (dd, *J* = 1.2, 0.4 Hz, 1*H*), 8.39 (dd, *J* = 8.8, 2.0 Hz, 1*H*), 8.24 (dd, *J* = 8.8, 0.4 Hz, 1*H*), 7.52 (m, 4*H*), 7.42 (m, 4*H*), 4.01 (s, 3*H*). ¹³C NMR (100 MHz): δ = 166.4, 155.2, 154.4, 143.2, 140.4, 138.7, 138.6, 132.0, 131.2, 129.9, 129.8, 129.3, 129.2, 129.1, 129.0, 128.4, 52.7. MS-ESI (m/z), 341(M⁺ + 1).

Methyl-2,3-di(4-bromophenyl)quinoxaline-6-carboxylate, 4b

¹H NMR (400 MHz, CDCl₃): δ = 8.87 (d, *J* = 1.2 Hz, 1*H*), 8.39 (dd, *J* = 8.8, 2.0 Hz, 1*H*), 8.24 (dd, *J* = 8.8 Hz, 1*H*), 7.58 (m, 4*H*), 7.39 (m, 6*H*), 4.00 (s, 3*H*). ¹³C NMR (100 MHz): δ = 166.2, 153.5, 152.9, 143.2, 140.4, 137.3, 131.8, 131.4, 129.9, 129.4, 124.3, 124.1, 52.7. MS-EI: 497 (M + H). HRMS (EI): calcd. for C₂₂H₁₄Br₂N₂NaO₂ (M + Na) 518.9320, found 518.9318.

Methyl-2,3-di(4-difluorophenyl)quinoxaline-6-carboxylate, 4c

¹H NMR (400 MHz, CDCl₃): δ = 8.88 (d, *J* = 1.6 Hz, 1*H*), 8.38 (dd, *J* = 8.8, 2.0 Hz, 1*H*), 8.20 (dd, *J* = 8.4, 0.4 Hz, 1*H*), 7.54 (m, 4*H*), 7.01 (m, 4*H*), 4.01 (s, 3*H*). ¹³C NMR (100 MHz): δ = 166.3, 164.7, 164.6, 153.8, 153.1, 143.0, 141.1, 134.5, 134.4, 131.9, 131.9, 131.8, 131.7, 131.6, 129.7, 129.3, 52.7. MS-EI: 377 (M + H). HRMS (EI): calcd. for C₂₂H₁₄F₂N₂NaO₂ (M + Na) 399.0921, found 399.0918.








Methyl-2,3-di(4-methoxyphenyl)quinoxaline-6-carboxylate, 4d

¹H NMR (400 MHz, CDCl₃): δ = 8.84 (d, *J* = 1.6 Hz, 1*H*), 8.31 (dd, *J* = 8.8, 1.6 Hz, 1*H*), 8.15 (d, *J* = 8.8 Hz, 1*H*), 7.52 (m, 4*H*), 6.91 (m, 4*H*), 4.01 (s, 3*H*), 3.85 (s, 6*H*). ¹³C NMR (100 MHz): δ = 166.5, 160.6, 160.4, 154.7, 154.0, 143.0, 140.2, 131.6, 131.3, 131.2, 131.1, 131.0, 130.7, 129.1, 113.8, 55.1, 52.7. MS-EI: 401 (M + H). HRMS (EI): calcd. for C₂₄H₂₀N₂NaO₄ (M + Na) 423.1321, found 423.1333.

Methyl-2,3-di(2-furyl)quinoxaline-6-carboxylate, 4e [23]

¹H NMR (400 MHz, CDCl₃): δ = 8.85 (d, *J* = 1.6 Hz, 1*H*), 8.34 (dd, *J* = 8.8, 1.6 Hz, 1*H*), 8.17 (d, *J* = 8.8 Hz, 1*H*), 7.65 (ddd, *J* = 5.2, 1.6, 0.8 Hz, 2*H*), 6.76 (dd, *J* = 5.2, 1.6 Hz, 2*H*), 6.60 (m, 2*H*), 4.02 (s, 3*H*). ¹³C NMR (100 MHz): δ = 166.1, 150.6, 143.9, 143.3, 142.6, 139.7, 131.8, 131.3, 129.9, 129.3, 114.3, 113.5, 112.1, 112.0, 52.7. MS-ESI: 321(M⁺ + 1).

Table 1. Melting points and yields for quinoxalines **4a-g**.

Run	Product	Yield ^a (%)	MP (°C)
1		30	144–145 ^b
2		34	193–194
3		38	188–189
4		26	163–164
5		28	164–165 ^c
6		36	183–184
7		31	193–195

^aOver the four steps, based on PEG-4500.

^bKnown compound [22], no mp reported.

^cKnown compound [23], no mp reported.

Methyl-2,3-di(2-chlorophenyl)quinoxaline-6-carboxylate, 4f

¹H NMR (400 MHz, CDCl₃): δ = 8.97(d, *J* = 1.6 Hz, 1*H*), 8.46 (dd, *J* = 8.9, 1.4 Hz, 1*H*), 8.29 (d, *J* = 6.4 Hz, 1*H*), 7.48 – 7.25 (m, 8*H*). ¹³C NMR (100 MHz): δ = 166.1, 154.4, 153.7, 143.1, 140.5, 136.9, 136.8, 132.9, 132.1, 131.8, 131.3, 131.1, 130.4, 130.1, 130.0, 129.7, 129.6, 126.4, 52.7. MS-EI: 409 (M + H). HRMS (EI): calcd. for C₂₂H₁₄Cl₂N₂NaO₂ (M + Na) 431.0330, found 431.0326.

Methyl-2,3-bis(4-dimethylaminophenyl)quinoxaline-6-carboxylate, 4g

¹H NMR (400 MHz, CDCl₃): δ = 8.87(d, *J* = 1.5 Hz, 1*H*), 8.34 (dd, *J* = 8.2, 1.3 Hz, 1*H*), 8.15 (d, *J* = 8.1 Hz, 1*H*), 7.55 (m, 4*H*), 6.8 (m, 4*H*), 4.03 (s, 3*H*), 3.91 (s, 6*H*), 3.89 (s, 6*H*). ¹³C NMR (100 MHz): δ = 166.5, 160.6, 160.5, 154.7, 154.0, 143.1, 140.3, 132.4, 132.3, 131.7, 131.4, 131.3, 130.7, 129.2, 129.1, 113.6, 110.9, 55.7, 55.3, 52.6. MS-EI: 427 (M + H). HRMS (EI): calcd. for C₂₆H₂₇N₄O₂ (M + H), 427.2134, found, 427.2156.

Conclusion

Multistep microwave-assisted synthesis of quinoxalines on soluble PEG-4500 polymer support is presented. A sequence of microwave-assisted esterification, reduction, and condensation is demonstrated. In each step, the product is separated simply by precipitation, filtration, and washing. This work demonstrates the validity of liquid-phase, polymer-supported synthesis in combination with microwave energy to carry out multistep synthesis in a convenient and efficient way, paving the way to a more environment friendly, automated, parallel, and combinatorial chemistry approaches.

Acknowledgements

Farid Fouad would like to thank Professor Robert Twieg at the Department of Chemistry and Biochemistry, Kent State University, for providing his lab facility to carry out this work.

References

- [1] Sun CM. Recent advances in liquid-phase combinatorial chemistry. *Combin Chem High Throughput Screen.* 1999;2:299–318.
- [2] Wentworth P Jr. Recent developments and applications of liquid-phase strategies in organic synthesis. *Trends in Biotechnol.* 1999;17:448–452.

- [3] Gravert DJ, Janda KD. Synthesis on soluble polymers: new reactions and the construction of small molecules. *Curr Opin Chem Biol.* 1997;1:107–113.
- [4] Toy PH, Janda KD. Soluble Polymer-Supported Organic Synthesis. *Acc Chem Res.* 2000;33:546–554.
- [5] Bonora GM. Polyethylene glycol: a high-efficiency liquid phase (HELP) for the large-scale synthesis of the oligonucleotides. *Appl Biochem Biotechnol.* 1995; 54:3–17.
- [6] Katritzky AR. *Advances in Heterocyclic Chemistry.* 1992;53:249.
- [7] Lv M, Xu H. The combinatorial synthesis of bioactive quinoxalines, quinoxalinones and quinoxalinols. *Combin Chem High Throughput Screen.* 2010;13:293–301, and references therein.
- [8] More SV, Sastry MNV, Yao C. Cerium (IV) ammonium nitrate (CAN) as a catalyst in tap water: a simple, efficient and green approach to the synthesis of quinoxalines. *Green Chem.* 2006;8:91–95, and references therein.
- [9] Saifina DF, Mamedov VA. New and modified classical methods for the synthesis of quinoxalines. *Russian Chem Rev.* 2010;79:351–370.
- [10] Staszewska A, Stefanowicz P, Szewczuk Z. Direct solid-phase synthesis of quinoxaline-containing peptides. *Tetrahedron Lett.* 2005;46:5525–5528.
- [11] Jeon MK, Kim DS, La HJ, Gong YD. Solid-phase synthesis of quinoxaline derivatives using 6-amino-2,3-dichloroquinoxaline loaded on AMEBA resin. *Tetrahedron Lett.* 2005;46:4979–4983.
- [12] Mirjalili BF, Bamoniri A, Akbari A. Nano-BF₃·SiO₂: a reusable and eco-friendly catalyst for synthesis of quinoxalines. *Chem Het Com.* 2011;47:487–491.
- [13] Jafarpour M, Rezaeifard A, Heidari M. A new catalytic method for eco-friendly synthesis of quinoxalines by zirconium (IV) oxide chloride octahydrate under mild conditions. *Lett Org Chem.* 2011;8:202–209.
- [14] Lue H, Yang S, Deng J, Zhang Z. Magnetic Fe₃O₄ Nanoparticles as New, Efficient, and Reusable Catalysts for the Synthesis of Quinoxalines in Water. *Aust J Chem.* 2010;63:1290–1296.
- [15] Nagarapu L, Mallepalli R, Arava G, Yeramanchi L. Polyethylene glycol (PEG-400)-mediated synthesis of quinoxalines. *Eur J Chem.* 2010;1:228–231.
- [16] Chavan HV, Adsul LK, Bandgar BP. Polyethylene glycol in water: a simple, efficient and green protocol for the synthesis of quinoxalines. *JCS.* 2011;123:477–483.
- [17] Zhang XZ, Wang JX, Bai L. Microwave-assisted synthesis of quinoxalines in PEG-400. *Synth Commun.* 2011;41:2053–2063.
- [18] Zhao Z, Wisnoski DD, Wolkenberg SE, Leister WE, Wang Y, Craig W, Lindsley CL. *Tetrahedron Lett.* 2004;45:4873–4876.
- [19] Krishnakumar B, Velmurugan R, Jothivel S, Swaminathan M. An efficient protocol for the green synthesis of quinoxaline and dipyrindophenazine derivatives at room temperature using sulfated titania. *Catal Commun.* 2010;11(12):997–1002.

- [20] Krishnakumar B, Swaminathan M. Solvent free synthesis of quinoxalines, dipyridophenazines and chalcones under microwave irradiation with sulfated Degussa titania as a novel solid acid catalyst. *J Mol Catal A: Chem.* 2011;350(1–2):16–25.
- [21] Krishnakumar B, Swaminathan M. A recyclable solid acid catalyst sulfated titania for easy synthesis of quinoxaline and dipyridophenazine derivatives under microwave irradiation. *Bull Chem Soc Japan.* 2011;84(11):1261–1266.
- [22] Chun Z, Zejun X, Liangren Z, Ning J. Et₃N-catalyzed oxidative dehydrogenative coupling of α -unsubstituted aldehydes and ketones with aryl diamines leading to quinoxalines using molecular oxygen as oxidant. *Tetrahedron.* 2012; 68:5258–5262.
- [23] You L, Cho EJ, Leavitt J, Ma L-C, Montelione GT, Anslyn EV, Krug RM, Ellington A, Robertus JD. Synthesis and evaluation of quinoxaline derivatives as potential influenza NS1A protein inhibitors. *Bioorg Med Chem Lett.* 2011;21:3007–3011.