## University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Dissertations and Theses in Biological Sciences

Biological Sciences, School of

December 2006

# Molecular systematics and population structure in the North American endemic fish genus *Cycleptus* (Teleostei: Catostomidae)

Michael Lee Bessert University of Nebraska - Lincoln, bessertm@uwstout.edu

Follow this and additional works at: http://digitalcommons.unl.edu/bioscidiss Part of the <u>Life Sciences Commons</u>

Bessert, Michael Lee, "Molecular systematics and population structure in the North American endemic fish genus *Cycleptus* (Teleostei: Catostomidae)" (2006). *Dissertations and Theses in Biological Sciences*. 1. http://digitalcommons.unl.edu/bioscidiss/1

This Article is brought to you for free and open access by the Biological Sciences, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Dissertations and Theses in Biological Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Molecular systematics and population structure in the North American endemic fish genus *Cycleptus* (Teleostei: Catostomidae)

by

Michael L. Bessert

### A DISSERTATION

Presented to the Faculty of

the Graduate College at the University of Nebraska

In partial Fulfillment of Requirements

For the Degree of Doctor of Philospophy

Major: Biological Sciences

Under the Supervision of Professor Guillermo Ortí

Lincoln, Nebraska

December, 2006

## Molecular systematics and population structure in the North American endemic fish genus *Cycleptus* (Teleostei: Catostomidae)

Michael L. Bessert, Ph.D.

University of Nebraska, 2006

Advisor: Guillermo Ortí

The overarching theme of this research was to investigate hierarchical levels of relatedness in natural populations of the cycleptid fishes (blue suckers), a widespread genus in North America that is of conservation concern throughout. Phylogenetic analysis of mitochondrial DNA sequences revealed that the two described *Cycleptus* species, C. elongatus and C. meridionalis are not reciprocally monophyletic, yet do not share any haplotypes. Although lineage sorting is incomplete, Bayesian coalescent analyses indicate that the two groups diverged early in the Pleistocene and have been reproductively isolated since. Whether they should be synonymized as *C.elongatus* and recognized as subspecies is open for debate. Due to differing morphological and allozyme profiles, the author is hesitant to call for this revision. Phylogenetic analyses also revealed that cycleptids in the Rio Grande are monophyletic and clearly divergent from the C. meridionalis/C. elongatus clade. Morphology is being revisited in Rio Grande specimens and a species description is underway. A novel suite of genusspecific, diploid microsatellite markers was also developed for population genetic analyses. Due to the tetraploid nature of these fishes, a single primer pair often

coamplifies multiple paralogous loci, limiting the usefulness of such markers; thus, a new technique was developed to isolate paralogous loci from one another. Genotypic data from isolated paralogs is consistent with functional diploidy and an allotetraploid origin for the family (Catostomidae). Population genetic analyses revealed significant isolation by distance and reduced allelic richness in the upper Missouri River, which has been fragmented by six dams and reservoirs for 50-60 years. These results are in contrast with those from the comparably-sized, but unimpounded, Mississippi River. These differences are noteworthy because only 5-6 generations have passed since the dams were constructed, rendering this one of the earliest detections of genetic effects of habitat fragmentation. Additional monitoring of this system with repeated genetic surveys is strongly recommended. Finally, rangewide population genetic analyses detected nine distinct subpopulations. Four occur in drainages outside of the Mississippi Basin while five occur within the basin. Whether or not the five intrabasin groups are ecologically exchangeable is unknown. Further study of specific subpopulations is encouraged.

### ACKNOWLEDGEMENTS

Where would I be without my beautiful wife, Gianna? Gianna, you are a blessing I do not deserve. I am not sure where I would be without you, but almost assuredly, it would not be here; that is, on the brink of receiving a doctoral degree at UNL. You have kept me going through all the challenges of graduate school in a very selfless and loving manner, and have taken on virtually all the 'home responsilities.' This degree really ought to have your name on it as much as mine because you did at least as much work - it was simply behind the scenes. Now that the crazy hours of graduate school are past, I am truly looking forward to getting to know you again and spending more time being a 'daddy' to our little blessings, Anna and William.

I was born and raised a Christian (and still am) and I would be remiss if I did not thank God for all the blessings in my life, especially an upbringing by parents that were and are the best role models one could have. Thank you, Mom and Dad, for raising me in the Christian faith and modeling honesty, kindness, and perseverance in the face of challenges – especially, in facing such challenges with a calm demeanor. Your examples – and your support and encouragement - were valuable to me throughout graduate school.

Speaking of families, I am also thankful for all the support from our extended families, especially my brother, Tim, and his wife, Betsy, who also live here in Lincoln. I can't count how many times they dropped whatever they were doing to help me - or us out. It was always a great stress relief to come over and watch games, grill out, or just relax. Thank you for your generosity. I am also thankful for my in-laws, Tom and Georgie Hoem. While Gianna was a great support for me, you were a great support for Gianna and helped ease the burden for both of us. The next time you come to visit, I will actually be able to spend time visiting rather than going off to the lab or hiding out writing somewhere. I am looking forward to that. Of course, all of our other brothers and sisters, nephews, nieces, aunts, uncles, and grandparents have all provided meaningful support throughout. Thank you very much.

What can I say about my advisor, Guillermo Ortí? When I came to UNL, I started out as a MS student under Royce Ballinger, who really helped get me in the door. When Royce left, Guillermo 'adopted' me even though I had very little knowledge of molecular techniques. I am forever grateful for that. I remember when I finished my Master's degree and was in the lab late one evening, I told Guillermo I wanted to pursue a PhD and thanked him for giving me a chance. He seemed very surprised and said, "Well, everyone deserves a chance." That may be true, but it certainly is not the way higher academia usually works. Not everyone gets a chance, especially with the limited background that I had. I feel very fortunate for having been allowed to learn molecular techniques – and the ways of graduate school - in your lab, Guillermo. What I appreciate most is how you treat all of your students with respect, not as mindless inferiors. In particular, I really appreciate the level of trust you placed in me; that is, how you allowed me to work fairly independently, but never had any doubts (at least I don't think) that I was making progress. Of course, I also have great respect for you as a scientist. You are one of those rare people who possesses top notch skills and scientist intellect, yet has a personality that I enjoy being around. In short, I think you were the perfect advisor for me.

In addition to Guillermo, I must express my sincere gratitude to all the members of my supervisory committee, Larry Harshman, Mary Liz Jameson, Guoqing Lu, and Etsuko Moriyama. All took time out of their busy schedules to meet with me, provide feedback on my project and proposals, and continually encourage me. They helped me navigate my way through the waters of graduate school and they continue to invest in me with letters, advice, and continued support. Thank you so much!

What are some of the things I will not miss about graduate school? I will not miss spending all night in the lab, often from 10:00pm to 6:00am. Neither will I miss spending night after night writing. I will not miss filling tip boxes, running the infamous 'black box' (BaseStation sequencer), or Sunday evening lab meetings for 101. However, I WILL miss all the great friends I have come to know in the Ortí lab: Paul Bates, Sara Brant, Chad Brock (a great person to talk philosophy with), Jeremy Brozek (we will get that paper out soon), Wei-Jen Chen (who helped me struggle through my MS), Federico Hoffman (who saved me from computer chaos on multiple occasions), Jeff Huebschman (always encouraging), Agustin Jimenez, Chenhong Li (my true confidant in the lab), Jason Macrander (a very hard worker), Federico Ocampo, Annie Paradis (I made it!), Cory Ross (another great friend – thanks for listening to me rant), Craig Sitzman (yeah! we are coauthors.), Julie Sommer (Kurtus STILL puzzles me), Michelle Steinauer, Obdulia Segura Leon (one of the nicest people I know), Adela Roa Varon (another one of the nicest people I know – on par with Obdulia), Stuart Willis, and Matt Bolek (my buddy from the parasit world who is going to make a great professor wherever he ends up).

If I had to single one person out in the lab, though, it would be Chenhong. I was a little unsure what to think when he joined the lab because he was so quiet. I am sure that is what people often think of me, too. What a great friend you have been, Chenhong! It's rare to meet someone who comes from such a vastly different culture, yet, shares so many common interests as well as a common mindset – that is, quiet determination. You know you have a great friend when you can communicate volumes without saying a single word. Chenhong, you set a high standard in the lab and it really pushed me – not in a negative way, but in a positive way. I hope the positive impact has been mutual. You have accomplished some amazing things. I can't believe you tamed the black box before it disappeared! You also got me to eat jellyfish, and now you can outfish me, too! I am going to have to do something about that. Seriously, I have no doubt that you will achieve all the goals you set. You are a tireless worker. It has been a real pleasure working with you on all our side projects and I look forward to collaborating on others in the future, wherever each of us ends up.

And with that, I will close this long journey. It has been pretty arduous at times, but also very satisfying. As I said in the beginning, I am a very blessed man and feel fortunate to have had the experience. I wonder which door will open next.

### TABLE OF CONTENTS

Dissertation The Lage	1
Abstract	ii
Acknowledgements	iii
Table of	
Contents	viii
List of Tables	X
List of Figures	xi
INTRODUCTION.	1
CUADTED ONE Melacular Systematics of the Erschwater Eigh Comus Cueles	
(Talaastai: Catastamidae) as informed from Mitashandrial DNA	pius 15
(Teleoster. Catostoffidae) as inferred from Mitochondrial DNA	13
Abstract	15
Introduction	16
Materials and Methods	10
Results	10 22
Discussion	22
A cknowledgements	24 20
Tables	2)
Figure	
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (Cvcle	eptus
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish ( <i>Cycle elongatus</i> , Catostomidae) with tests of cross-species amplification and i paralogous loci	<i>eptus</i> isolation of 48
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish ( <i>Cycle elongatus</i> , Catostomidae) with tests of cross-species amplification and i paralogous loci.	<i>eptus</i> isolation of 48
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish ( <i>Cycle elongatus</i> , Catostomidae) with tests of cross-species amplification and i paralogous loci	<i>eptus</i> solation of 48
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish ( <i>Cycle elongatus</i> , Catostomidae) with tests of cross-species amplification and i paralogous loci Abstract Introduction	<i>eptus</i> isolation of 48 48 49 49
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish ( <i>Cycle elongatus</i> , Catostomidae) with tests of cross-species amplification and i paralogous loci Abstract Introduction	<i>eptus</i> solation of 48 48 49 49 49
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion</li> </ul>	<i>eptus</i> solation of 48 48 49 51 51
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion.</li> <li>Acknowledgements</li> </ul>	<i>eptus</i> isolation of 48 48 49 51 51 53
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion.</li> <li>Acknowledgements.</li> <li>Tables</li> </ul>	<i>eptus</i> isolation of 48 48 49 51 51 53 54
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion.</li> <li>Acknowledgements.</li> <li>Tables.</li> </ul>	<i>eptus</i> isolation of 48 48 49 51 51 53 54
CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish ( <i>Cycle elongatus</i> , Catostomidae) with tests of cross-species amplification and i paralogous loci Abstract Introduction Materials and Methods Results Discussion Acknowledgements Tables	<i>eptus</i> isolation of 48 48 49 51 51 53 54
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion.</li> <li>Acknowledgements.</li> <li>Tables.</li> </ul>	<i>eptus</i> isolation of 48 48 49 51 51 53 54 <i>us</i> of habitat 56
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion.</li> <li>Acknowledgements.</li> <li>Tables.</li> </ul> CHAPTER THREE - Population genetic structure of the Blue Sucker ( <i>Cycleptu elongatus</i> Lesueur, 1918) in the Upper Missouri River: genetic effects of fragmentation.	<i>eptus</i> isolation of 48 48 49 51 51 53 54 <i>us</i> of habitat 56
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>Discussion.</li> <li>Acknowledgements.</li> <li>Tables.</li> </ul> CHAPTER THREE - Population genetic structure of the Blue Sucker ( <i>Cycleptu elongatus</i> Lesueur, 1918) in the Upper Missouri River: genetic effects of fragmentation.	<i>eptus</i> isolation of 48 49 51 51 53 54 <i>us</i> of habitat 56 56
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results</li> <li>Discussion</li> <li>Acknowledgements</li> <li>Tables.</li> </ul> CHAPTER THREE - Population genetic structure of the Blue Sucker ( <i>Cycleptu elongatus</i> Lesueur, 1918) in the Upper Missouri River: genetic effects of fragmentation. Abstract Introduction. Materials and Methods	<i>eptus</i> isolation of 48 48 49 51 51 53 54 <i>us</i> of habitat 56 56 58
<ul> <li>CHAPTER TWO – Microsatellite loci for allotetraploid blue sucker fish (<i>Cycle elongatus</i>, Catostomidae) with tests of cross-species amplification and i paralogous loci.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> <li>Results.</li> <li>CHAPTER THREE - Population genetic structure of the Blue Sucker (<i>Cycleptu elongatus</i> Lesueur, 1918) in the Upper Missouri River: genetic effects of fragmentation.</li> <li>Abstract.</li> <li>Introduction.</li> <li>Materials and Methods.</li> </ul>	<i>eptus</i> isolation of 48 48 49 51 51 53 54 <i>us</i> of habitat 56 56 56 56 56 

Discussion	
Acknowledgements	
Tables	72
Figures	79
CHAPTER FOUR – Rangewide population structure and	I intermediate polyphyly in the
genus Cycleptus (Teleostei: Catostomidae)	
Abstract	91
Introduction	
Materials and Methods	
Results	
Discussion	
Acknowledgements	
Tables	
Figures	
CONCLUSIONS	
REFERENCES	
APPENDIX ONE	
APPENDIX TWO	

## LIST OF TABLES

TABLE 1.1	Taxa used for cytochrome b primer design
TABLE 1.2	Sequence divergence within and among <i>Cycleptus</i> taxa31
TABLE 2.1	Primer sequences for eleven <i>Cycleptus</i> microsatellite loci
TABLE 2.2	Cross-species amplification results with <i>Cycleptus</i> primers55
TABLE 3.1	Missouri River samples and summary statistics72
TABLE 3.2	Microsatellite characteristics and labeling scheme73
TABLE 3.3	Pairwise R <sub>ST</sub> comparisons for Missouri River samples74
TABLE 3.4	Missouri River AMOVA results75
TABLE 3.5	Missouri River BOTTLENECK results
TABLE 3.6	Pairwise distance matrix for Missouri River sample sites77
TABLE 3.7	Pairwise distance matrix for Mississippi River sample sites78
TABLE 4.1	Rangewide sample localities and summary statistics111
TABLE 4.2	Rangewide AMOVA summary

## LIST OF FIGURES

FIGURE 1.1	Cytochrome <i>b</i> translation in <i>Cycleptus</i>	32
FIGURE 1.2	Cycleptus cytochrome b maximum likelihood phylogram	34
FIGURE 1.3	Cycleptus control region maximum parsimony tree	36
FIGURE 1.4	MCMC convergence assessment with AWTY	38
FIGURE 1.5	Cycleptus control region Bayesian phylogram	40
FIGURE 1.6	Cycleptid Bayesian phylogram based on concatenated data	42
FIGURE 1.7	C. elongatus and C. meridionalis morphology comparison	44
FIGURE 1.8	Cycleptid lip morphology and tuberculation	46
FIGURE 3.1	Missouri River sample localities	79
FIGURE 3.2	Multiplex reaction 'A' chromatogram	81
FIGURE 3.3	Multiplex reaction 'B' chromatogram	83
FIGURE 3.4	Results from Missouri River STRUCTURE analysis	85
FIGURE 3.5	Missouri River IBD results: log(M) vs geographic distance	87
FIGURE 3.6	Mississippi River IBD results: log(M) vs geographic distance.	89
FIGURE 4.1	<i>C. elongatus</i> state conservation status	113
FIGURE 4.2	Rangewide sampling locality map	115
FIGURE 4.3	Rangewide Bayesian STRUCTURE analysis results	117
FIGURE 4.4	<i>C. elongatus</i> + <i>C. meridionalis</i> unrooted haplotype network	119
FIGURE 4.5	Isolation with Migration model	121
FIGURE 4.6	Demographic parameter estimates from IM analysis	123

### **INTRODUCTION**

### Rationale

The overarching theme of this research effort was to investigate hierarchical levels of relatedness in natural populations of the genus *Cycleptus* (blue suckers), an enigmatic group of freshwater fishes native to large rivers in the United States and Mexico. These hydrodynamic fishes are benthic in habit and attain lengths up to 93 cm and weights greater than 10 kg. Breeding colors range from olive blue in *C. elongatus* to 'almost black' in *C. meridionalis* while individuals from the Rio Grande (currently recognized as *C. elongatus*) have a more golden or brassy appearance (Burr and Mayden 1999). Prominent breeding tubercles cover much of the body of spawning males and are present in reduced numbers on females. Both recognized species are known to migrate long distances to spawn. Spawning occurs in fast current over gravel beds. *Cycleptus elongatus* has an approximate 10-year generation time (Becker 1983) while *C. meridionalis* may live considerably longer (Boschung and Mayden 2004). Although the genus is known from 22 states, populations are not considered stable in any one of them.

The research provides important contributions to a range of biological subdisciplines, including systematics, molecular techniques, and conservation biology. For the chosen study taxon, the latter is of greatest urgency given the current status of the group. In carrying out this work, I have assembled one of the most expansive tissue collections for any single freshwater fish taxon in North America – both numerically and in terms of geographic coverage. Novel genetic markers were developed and employed to estimate phylogeny, demography, and phylogeographic patterns in the group. At the time of this writing, there are approximately 20,000-28,000 described fish species and approximately 40% of these occur in freshwater systems (Eschmeyer 1998; Moyle and Cech 2000; Nelson 1994). In the past 100 years, anthropogenic modifications to freshwater lakes, rivers, and streams has have negatively impacted many species while others have faced direct threats from overexploitation (Duncan and Lockwood 2001; Richter *et al.* 1997). Furthermore, there are very few remaining systems that have not been subject to the introduction of nonindigenous species, often with catastrophic results for native taxa (Pimentel *et al.* 2000). Exotic fishes may compete directly for resources or, if closely related, may "swamp out" native stocks via introgressive hybridization (Koppelman 1994; Weigel *et al.* 2003). As a result of these cumulative threats, an estimated 20% of freshwater fish species are at risk for extinction (Leidy and Moyle 1998). In North America, trends of freshwater fish decline are consistent with other regions as approximately ¼ of all species are considered imperiled - and at least 10 species have recently become extinct (Williams *et al.* 1989).

North America harbors approximately 1000 freshwater fish species if the Mexican transition zone is included (Mayden *et al.* 1992; Moyle and Cech 2000). These fishes are distributed among 50 families and 201 genera, but almost 70% of the diversity lies within only four families: Cyprinidae (minnows, 34%), Percidae (perches and darters, 18%), Poeciliidae (livebearers, 8%), and Catostomidae (suckerfishes, 8%). The latter family, which includes the genus *Cycleptus*, is comprised of 80 species that are endemic to North America with the exception two: *Catostomus catostomus* (longnose sucker), which also occurs (naturally) in Asia, and *Myxocyprinus asiaticus* (Chinese sucker), which is restricted to the Yangtze Basin in China.

There are a number of motivating factors for the selection of *Cycleptus* as a study taxon. To begin with, the genus occupies an interesting phylogenetic position within Catostomidae. It has been hypothesized to be sister to the Asian genus *Myxocyprinus* (Ferris and Whitt 1978; Harris and Mayden 2001; Smith 1992). This relationship, if real, reflects an interesting biogeographic pattern that is repeated in paddlefishes (*Polyodon* + *Psephurus*, family Polyodontidae) (Grande and Bemis 1991), giant salamanders (*Cryptobranchus* + *Andrias*, family Cryptobranchidae), and alligators (family Alligatoridae) (Burr and Mayden 1999).

Secondly, there is comparative merit in studying cycleptid fishes. The Mississippi Basin has provided a relatively stable environment for freshwater fish evolution since the late Cretaceous / early Cenozoic (Briggs 1986). As a result, various basal actinopterygian lineages persist in the basin, including Semionotiformes (gars), Polyodontiformes (paddlefishes – one species), and Acipenseriformes (sturgeons). While it is blatantly incorrect from a "tree thinking" standpoint (O' Hara 1988) to group cycleptids with the aforementioned, these fishes are, in fact, commonly included with the others in a group termed the "old river ichthyofauna" of North America (Smith 1981). Perhaps this is due to co-occurrence (similar range) or a common ecology. In any event, several of these lineages are of considerable economic importance (e.g., caviar trade), but most are poorly understood in terms of evolutionary history (ex. Simons et al. 2001). Furthermore, a firm understanding of population genetic structure in these groups is complicated by the fact that many have already been translocated (i.e. "stocked" from one location to another) prior to the collection of any genetic data (see Gardner 2004; Rutledge 1989); thus, naturally occurring allele frequencies, for example, have been

obscured. *Cycleptus* is an important exception. Although they were subject to intense harvest in the early 1900s (Coker 1930) to my knowledge, none have been translocated.

Another, personally motivating challenge in the studying cycleptid population genetics regards the makeup of their nuclear genome. All catostomids, including the genus *Cycleptus*, are tetraploid. The family is thought to have an allotetraploid origin (genome duplication via hybridization), having descended from a cyprinid-like ancestor some 50 mya (Uyeno and Smith 1972). A major complicating factor is that while analytical techniques abound for codominant, diploid loci, polyploid data remain difficult to analyze and interpret, at least in a population genetics context (Pearse and Crandall 2004).

Microsatellite loci (short, tandemly repeated sections of DNA – e.g., CACACA) are the markers of choice for population studies due to their high mutation rate via slipstrand mispairing (Goldstein and Schlötterer 1999). As with any other taxon, these markers are interspersed throughout the *Cycleptus* genome. They may be amplified with primers that match the flanking region on either side of a given locus. Unfortunately, in a polyploid situation, the priming sites may be conserved across multiple sets of chromosomes; thus, in a tetraploid, four fragments may be amplified instead of two, rendering common analytical techniques useless or, at best, difficult. One way to get around this problem is to screen a greater number of loci in the hopes that at least some priming sites are not conserved across duplicate pairs of chromosomes. I have successfully used this approach. In addition, I have developed a method for circumventing the problem by isolating paralogous (co-amplifying) loci from one another (Chapter 3 (already published) - see Bessert *et al.* 2006). A final rationale for the study of cycleptid fishes regards the general conservation status of the genus. They are widely distributed throughout the interior of central North America and extend southward into the Rio Grande basin, a range that includes 22 conterminous states, yet, as previously mentioned, populations are not considered stable in any one of them. At best, they have a natural heritage ranking of S3 (vulnerable) and at worst, SH (possibly extirpated). Most authorities subscribe to trends of population decline resulting from impoundments (blocked spawning routes) and pollution because *C. elongatus* is thought to be highly sensitive (Becker 1983; Pflieger 1997; Robison and Buchanan 1998). However, others suggest that these apparent declines are not an immediate cause for concern and may, in part, be an artifact of inadequate sampling, such as in remote areas of the Rio Grande Basin (Burr and Mayden 1999). Such is the impetus for further investigations such as this one.

#### Literature Review

In the established tradition of thesis preparation, I provide a literature review in the following pages. Rather than provide an exhaustive list of any publication that has ever mentioned cycleptid fishes, I have decided to winnow it down to only include literature that is relevant to the topics at hand – or that may be of future use to researchers building upon this work. For instance, simple redundant appearances on state lists with no additional information were excluded. The review begins with the original species description and concludes with contemporary studies, including the first publication from this dissertation.

*Cycleptus elongatus* was originally described by Charles Alexander Lesueur as *Catostomus elongatus* (Lesueur 1817) and placed within the family Catostomidae. His

effort was based on a dried specimen from the Ohio River that has since been reported lost or discarded (Burr and Mayden 1999). The species was subsequently transferred to a new subgenus *Cycleptus* (Rafinesque 1820). Approximately 25 years later, *Cycleptus* was elevated to generic status (Agassiz 1855).

An abundance of locality and additional descriptive records appeared throughout the mid-1800s to early 1900s, including: presence in Indiana (Jordan 1875); listing for Illinois (Nelson 1876); clarification of name (Jordan 1877); listing in North American catalog (Jordan 1878); presence in Cumberland River, Tennessee (Jordan and Brayton 1878); appearance in the Mississippi Valley (Jordan and Gilbert 1883); presence in the Baraboo River, Wisconsin (Hoy 1883); presence in the Kansas River, Kansas (Cragin 1885; Graham 1885); presence in Whitewater River, Indiana (Evermann 1886); presence in Ohio and Wabash Rivers (Jordan and Evermann 1886); uncommon in Ohio River (Henshall 1888); uncommon in Iowa (Meek 1890); presence in Cumberland River, Kentucky (Woolman 1892); presence at Falls of the Ohio River (Call 1896); uncommon in Missouri River (Evermann and Cox 1896); general description and Illinois distribution (Forbes and Richardson 1908); presence in Río Solado, Nuevo Léon, Mexico (Fowler 1913; Meek 1908); presence in Kiskeminetas River, Pennsylvania (Fowler 1913); description and summary of range (Jordan 1929); spring and fall migration patterns in the upper Mississippi River (Coker 1930); Illinois distribution reviewed (Donnell 1935).

From 1935-1945, coincidental with World War II, no studies were published on blue suckers. After that time, locality records continued but were intermixed with comments on population trends, anatomical studies, etc.: distribution in Minnesota (Eddy and Surber 1947); description of the Weberian ossicles in *Cycleptus* (Nelson 1948); presence in Red River drainage, Oklahoma (Moore and Cross 1950); decline in main stem of upper Mississippi River (Barnickol and Starrett 1951); presence in Big Sioux River, Iowa (Harrison and Speaker 1954); presence in Big Muddy River, Illinois (Lewis 1955); comments on biology and range in Iowa (Harlan and Speaker 1956); presence in Texas Gulf Slope drainages (Hubbs 1957); presence in New Mexico (Koster 1957); distribution in Ohio (Trautman 1957); distribution in Mississippi (Cook 1959); presence in Neosho River, Kansas (Metcalf 1959); presence in lower Vermillion River and Missouri River, South Dakota (Underhill 1959); presence in the Big Blue River, Kansas (Minckley 1959); Cycleptus bones found at Bandelier National Monument, New Mexico - several hundred kilometers beyond their extant distribution in the state (Gehlbach and Miller 1961); studies of osteology and tuberculation (Branson 1962a; Branson 1962b); presence in the lower Missouri River (Fisher 1962); brain and lip morphology (Miller and Evans 1965); comments on spawning dates (Breder and Rosen 1966); update of Kansas distribution (Cross 1967); age and growth (Carlander 1969); Montana distribution (Brown 1971); occurrence in Lake Sharpe (Missouri River impoundment), South Dakota (Elrod and Hassler 1971); distribution in Missouri (Pflieger 1971); status in upper Mississippi River – uncommon (Smith *et al.* 1971); large numbers in tailwaters of Gavins Point dam, South Dakota (Walburg et al. 1971); threatened in six states (Miller 1972); occurrence in Río Bravo (Rio Grande) (Contreras-Balderas and Rivera 1972); karyotype - polyploidy (Uyeno and Smith 1972); distribution in Arkansas (Buchanan 1973); description of gill parasites (Leiby et al. 1973); Oklahoma distribution (Miller and Robison 1973); common in lower reaches of Chippewa and Red Cedar Rivers, Wisconsin (Christenson 1974); accounts from the Red and Sabine Rivers in Louisiana (Douglas

1974); updated Minnesota distribution (Eddy and Underhill 1974); presence in Kiamichi River, Oklahoma (Pigg and Hill 1974); considered rare in Kansas and Oklahoma (Platt et al. 1974; Robison et al. 1974); Kentucky distribution (Clay 1975); distribution and status update in Kansas (Cross and Collins 1975); description of biology and updated Missouri distribution (Pflieger 1975); threatened in Oklahoma (Hubbs and Pigg 1976); evolution of pectoral fins (Lundberg and Marsh 1976); loss of duplicate gene expression – return to functional diploidy, based on allozymes (Ferris and Whitt 1977); pharyngeal bone structure (Eastman 1977); qualities of bile salts as phylogenetic markers (Bussjaeger and Briggs 1978); catostomid phylogeny based on loss of duplicate gene expression allozymes (Ferris and Whitt 1978); occurrence in the Tennessee River in Alabama (Etnier and Starnes 1979); threatened status in Indiana (McReynolds et al. 1979); Illinois range update (Smith 1979); structure of caudal skeleton (Eastman 1980); genetic variability compared to diploid teleosts – based on allozymes (Ferris and Whitt 1980); biology in upper Mississippi River (Rupprecht and Jahn 1980); threatened in Tennessee and Kentucky (Branson et al. 1981; Starnes and Etnier 1980); Holocene fossil record (Smith 1981); updated Ohio distribution (Trautman 1981); status in the lower Mississippi River (Guillory 1982); updated Minnesota distribution (Phillips et al. 1982); description of biology and Wisconsin distribution (Becker 1983); polyploidy confirmed with isozymes (Buth 1983); extirpated from Pennsylvania (Cooper 1983); presence in the Poteau River, Oklahoma (Lindsay et al. 1983); natural history (incl. spawning) from the Neosho River, Kansas (Moss *et al.* 1983); common along revetments in the lower Mississippi main stem (Pennington *et al.* 1983); present in the Green River, Kentucky (Retzer *et al.* 1983); distribution and abundance in Wisconsin (Fago 1984); tetraploidy (Ferris 1984);

phylogenetic position based on larval characters (Fuiman 1985); decline in Rio Grande (Williams et al. 1985); range reduced in Ohio drainage, member of 'old river' fauna (Burr and Page 1986); distribution in Gulf Slope drainages (Conner and Suttkus 1986); presence in Missouri River drainage (Cross *et al.* 1986); presence in the Rio Grande, Conchos, and Pecos rivers (Smith and Miller 1986); presence in Tennessee and Cumberland rivers (Starnes and Etnier 1986); found in proximity to wing dikes and revetments in middle Missouri River (Sandheinrich and Atchison 1986); member of 'old river' faunal group (Robison 1986); Cycleptus images on ancient pottery in New Mexico, far from extant range (Jett and Moyle 1986); collection from the Spring River, Arkansas (Baker and Armstrong 1987); diet in the Black River, New Mexico and distribution in New Mexico (Cowley and Sublette 1987a; Cowley and Sublette 1987b); updated Iowa distribution (Harlan and Speaker 1987); protected in the U.S. (Johnson 1987); doubtful from the Rio Grande in New Mexico (Propst et al. 1987); reproduction in upper Mississippi River (McInerny and Held 1988); description of biology and Arkansas distribution (Robison and Buchanan 1998); stranding and mortality below McAlpine dam, Ohio River (Pearson and Froedge 1989); species of special concern throughout range (Williams *et al.* 1989); range, status, and color image (Tomelleri and Eberle 1990); description and distribution in New Mexico (Sublette et al. 1990); not secure in Tennessee (Etnier and Starnes 1991); description, range map and habitat (Page and Burr 1991); distribution in Mississippi (Ross and Brenneman 1991); Wisconsin distribution (Fago 1992); phylogenetic position within Catostomidae, sister taxon to Myxocyprinus (Smith 1992); presence in James River, not secure in Dakotas (Berry et al. 1993); extirpated from Big Muddy River in Illinois (Burr and Page 1993); description, status in

Tennessee (Etnier and Starnes 1993); status in St. Croix River system, Wisconsin (Fago and Hatch 1993); status in the Wabash River, Indiana (Gammon 1993); status in Big Black and Yazoo Rivers, Mississippi (Holman et al. 1993; Jackson et al. 1993); present in the Arkansas River (Limbird 1993); rare in Kaskaskia River, Illinois (Larimore and Fritz 1993); present in western tributaries of the Missouri River, South Dakota (Ruelle et al. 1993); rare in the Kansas River system, Kansas (Sanders et al. 1993); presence in the Vermillion River, South Dakota (Schmulbach and Braaten 1993); healthy populations in the Yellowstone River, Montana (White and Bramblett 1993); present in the Lamine River, Missouri – spawning (Brown and Coon 1994); possible record in Virginia (Jenkins and Burkhead 1994); description of eggs and larvae (Kay et al. 1994); monotypic subfamily Cycleptinae (Nelson 1994); threatened status in Mexico (Contreras-Balderas et al. 1995); status update in Kansas (Cross and Collins 1995); distribution in West Virginia (Stauffer et al. 1995); new records in Illinois (Burr et al. 1996); Alabama distribution and biology (Mettee *et al.* 1996); update of Missouri distribution and comments on biology (Pflieger 1997); status below Miller's Ferry lock and dam, Alabama(Mettee and Shepard 1997); allozyme variation (Buth and Mayden 1998); stock dynamics in Yazoo River tributaries, Mississippi (Hand 1999); description and threatened status in New Mexico (Propst 1999); historic occurrence in Little Miami River, Ohio (Harrington 1999); formal description of Cycleptus meridionalis, summary opinion of conservation status - global populations not in imminent danger (Burr and Mayden 1999); negative impact of dams (Pringle *et al.* 2000); possible benefit from modifications to seasonal flow regime (Galat and Lipkin 2000); reproduction in backwaters of upper Mississippi River (Fisher and Willis 2000); life history characteristics in unchannelized Missouri River below Gavins

Point dam (Carreiro 2000); considered potential invader to Great Lakes (Cudmore-Vokey and Crossman 2000); gill parasites (Marcogliese 2001); recent accounts in Minnesota (Schmidt and Talmage 2001); allozymes in Rio Grande population distinct (Buth and Mayden 2001); use of main channel habitat in upper Mississippi River (Dettmers et al. 2001); Cycleptinae is either para- or polyphyletic (Harris and Mayden 2001); preference for fast current (Pegg and Pierce 2002); imperiled, but still occurs in upper Tennessee River (Butler 2002); swimming performance may give advantage in channelized waterways (Wolter and Arlinghaus 2003); stock characteristics in upper Yazoo basin, Mississippi (Hand and Jackson 2003a); spawning pattern in Missouri (Vokoun et al. 2003); abundance and biology in James and Big Sioux Rivers, North and South Dakota (Morey and Berry 2003); comparative abundance in two segments of upper Missouri River (Welker and Scarnecchia 2003); larval cycleptids in unimpounded upper Missouri River (Barko et al. 2004); larval densities in lower Missouri River (Galat et al. 2004); appearance in lower Missouri trawling efforts (Herzog 2004); status in Kansas (Haslouer et al. 2005); higher than expectected abundance in upper Kansas River, slow growth rates (Eitzmann et al. 2005); historic occurrence in middle and upper Rio Grande, New Mexico (Cowley 2006); included in general biogeographic study of Great Plains rivers (Hoagstrom and Berry 2006); larval blue sucker ecology in Mississippi River (Adams et al. 2006); development of molecular markers for population genetic studies (Bessert et al. 2006).

### Ongoing Studiesin the Genus

It may be apparent to the reader that most of these records are due to coincidental encounters with cycleptid fishes during other endeavors. The papers that appear in the

following chapters are four of only a handful of studies that have focused exclusively on cycleptid biology. In addition to my efforts, there are two other ongoing investigations in the genus that are worthy of note. In Texas, BioWest consulting company, under the direction of Dr. Ed Oborny, is investigating various life history aspects of C. elongatus in the Colorado River. Those studies are intended to assess potential impacts of modifications to the lower Colorado River (flow diversion to provide water to large cities) on C. elongatus populations. Absolute range within the Colorado, migration patterns, and spawning areas are being defined via radio telemetry. I have assisted this group in one respect, using my molecular markers to confirm the identity of embryos from at least one purported spawning site. A second group, under the leadership of Dr. Trent Sutton at Purdue University, is investigating the age and growth characteristics of blue suckers in the Wabash River of Indiana. The Wabash River population is relatively healthy and is said to provide an important food base for larger predators. This is in contrast to most drainages throughout the global range, so the outcome of Trent's work will hold comparative interest for numerous state agencies.

### Overview

The remainder of this dissertation is divided into four chapters, each with a particular focus and written to be published as a separate journal article. Chapter one examines intrageneric phylogeny using mitochondrial markers. It illuminates the divergence of the Rio Grande population from the two described species and provides an approximate time of divergence between the two major clades. The results of this work, along with previous morphological and allozyme studies (Burr and Mayden 1999; Buth and Mayden 2001), have provided the impetus to delineate

cycleptids from the Rio Grande as a new species. I am presently collaborating with Drs. Brooks Burr of Southern Illinois University (Carbondale) and Richard Mayden of St. Louis University on this work. Having personally visited a good portion of this species' range (including northern Mexico) and seen the effects of exotic salt cedar, golden algae, and drought, I may lobby to name it *Cycleptus speratus*. The Latin word *spero* means "to hope" or "to hope for," and the fishes and the people of this region are both in desparate need of some hope.

Chapter two characterizes the isolation and screening of a unique suite of microsatellite markers that I developed for population genetic studies in the genus. It also describes a new technique for isolating paralogous loci from one another. This can be a significant benefit to anyone attempting to use such markers in polyploid organisms. This work has already been published online (Bessert *et al.* 2006) and will soon appear in *Conservation Genetics*.

Chapter three details various measures of population structure in *C. elongatus* populations in the upper Missouri River. This drainage is unique relative to the remainder of the species range in that it has been subdivided by a series of six major dams and impoundments which have presumably isolated inter-reservoir populations for the last 50-60 years. Most of the population structural analyses show no startling aberrations with the exception of two: 1) The Missouri River populations show a pronounced signal of isolation by distance. This is in contrast to other continuous portions of the range (namely, the unimpounded Mississippi River); 2) Inter-reservoir samples from the upper Missouri show a significant reduction in allelic richness compared to samples from the lower Missouri and from throughout the range. These two

results may, indeed, be the first detectable deviations from normal / natural genetic patterns.

Finally, chapter four examines rangewide population structure using some 600 samples and 15 microsatellite markers. A Bayesian analysis of population structure shows nine distinct genetic clusters ('populations'). In addition to recognizing populations from disjunct drainages (Mobile Basin, Rio Grande Basin, Colorado River, and Sabine River) as distinct, there is clear structure within the Mississippi Basin as five distinct clusters are recognized there (i.e. a total of nine distinct genetic clusters from throughout the range). A second aspect of this chapter includes an investigation of the timing of divergence between *C. elongatus* and *C. meridionalis*, a pair of polyphyletic taxa in which lineage sorting is incomplete. A powerful coalescent approach was used to determine that these species diverged in early Pleistocene time. Associated results also reject the hypothesis of any recent gene flow, lending further support to the species designation of *C. meridionalis*.

## Molecular Systematics of the Freshwater Fish Genus *Cycleptus* (Teleostei: Catostomidae) as inferred from Mitochondrial DNA

### Abstract

Fishes in the genus *Cycleptus* occur in large rivers throughout central North America. The genus was considered monotypic as *Cycleptus elongatus* for over 175 years until a sister species, *C. meridionalis*, was described from southeastern Gulf Coastal drainages based on morphological differences. Subsequent allozyme data revealed evidence for remaining polytypy within *C. elongatus*, but intrageneric relationships have never been assessed with direct genetic evidence. Complete mitochondrial control region sequences (920 base pairs) were collected from 151 specimens and cytochrome *b* sequences were collected from a subset of 48 specimens representing the known range of the genus. Results indicate polyphyly between the two described species while the Rio Grande population, currently recognized as *C. elongatus*, is monophyletic and clearly divergent from the rest. These results, in combination with previous data sets, support designation of the Rio Grande group as a distinct species.

### Introduction

The blue sucker (*Cycleptus elongatus*) was originally described by Charles Alexander Lesueur (1817) as *Catostomus elongatus* from a dried museum specimen collected from the Ohio River. Three years subsequent, the species was placed within the subgenus *Cycleptus* (Rafinesque 1820) and twenty-five years later, *Cycleptus* was elevated to the level of genus (Agassiz 1855). For over 140 years the genus was considered monotypic as *Cycleptus elongatus* until Burr and Mayden (1999) described a sister species, *Cycleptus meridionalis*, from the Pearl and Mobile river basins (Mississippi and Alabama) based on an array of bimodally distributed morphological characters. Rio Grande specimens were divergent for some characters, most notably lip morphology and coloration, but not to the mutual exclusion of all other *Cycleptus* populations. A subsequent allozyme study (Buth and Mayden 2001) revealed further evidence for polytypy within *C. elongatus* because Rio Grande populations displayed unique profiles relative to those from the Mississippi Basin. Still, there have been no direct assessments of phylogeny with genetic (DNA sequence) data to this point.

Why do these fishes remain such an enigma? To be sure, cycleptid fishes have been the direct focus of only a handful of studies compared to most of their large riverine counterparts (e.g., sturgeons, paddlefishes). Although they occur over such a wide range, they are benthic in habit and their hydrodynamic morphology is associated with a preference for rapid current in main stem channels; thus, requiring heavy gear to collect them in reasonable numbers. In addition, unlike sturgeons and paddlefishes, which play a major role in the caviar trade, cycleptids hold no contemporary economic value (although they were commercially harvested in the upper Mississippi during the early 1900's).

Why, then, should we seek a better understanding of cycleptid relationships? Aside from the basic value in fleshing out the tree of life, the study of cycleptid evolution may hold comparative merit for conservation of other, more heavily targeted taxa. They have likely been present since the early Cenozoic (Smith 1981); yet, have not radiated to the degree that other lineages have.

Given the extant distribution of these fishes, multiple hypotheses of unrecognized biodiversity (and relationships among) become apparent. The Mississippi basin is the most stable system inhabited by these fishes; thus, it is likely that cycleptids originated here and dispersed to other areas. The Rio Grande basin is the most disjunct point from the Mississippi basin; therefore, it is logical to hypothesize that each drainage between harbors a distinct, monophyletic clade; i.e. the Sabine (closest to the Mississippi) forms a basal branch from which the Colorado / Rio Grande clades diverged, etc. In this scenario, we would expect the Rio Grande population to be the most divergent from C. *elongatus* inhabiting the Mississippi basin. Conversely, the nature of this distribution is poorly known, especially given the difficulties in reconstructing drainage evolution along the Gulf slope (Conner and Suttkus 1986; Conner 1977). Another possibility is that cycleptids dispersed more recently along coastal margins during periods of reduced salinity resulting from Pleistocene glacial melt. In a recent study, Burr and Mayden (Burr and Mayden 1999) distinguished C. meridionalis from C. elongatus based on an array of discrete morphological differences, yet failed to do so with the Rio Grande, Colorado, or Sabine populations; therefore, if mtDNA divergence is concordant with morphological

divergence, we may expect the two described species to form reciprocally monophyletic clades with western Gulf slope populations nested within *C. elongatus*. Here, I approach these questions with the first molecular systematic study of the genus *Cycleptus* using mitochondrial markers.

### Materials and methods

### Samples

Tissue samples were collected from 151 specimens throughout the known range of the genus (see Materials Examined; Appendix 2). All are vouchered in the personal collection of the author and freely available upon written request. Acquisition of samples was challenging because it requires large river gear such as benthic gill nets and electroshocking devices; thus, assistance was solicited from a combination of more than 100 individuals representing 22 state agencies, two federal agencies, and 15 academic institutions. All samples were acquired in an approximate one year time span from November 2004 to November 2005. A small fin clip, approximately 1-2 cm<sup>2</sup>, was excised from the anal fin of each specimen and fixed in 95% ethanol (EtOH) prior to shipment to the University of Nebraska.

### DNA extraction

Genomic DNA was isolated from fin clips employing a phenol-chloroform extraction protocol (Sambrook *et al.* 1989) or the DNeasy® Tissue purification kit (Qiagen). Extractions were quantified with a GeneQuant II spectrophotometer (Pharmacia Biotech) and diluted to a standard concentration of 100ng/µl to facilitate consistency in PCR amplification.

### Data collection

Primers Cyt b-F (5'-ATGGCAAGCCTACGAAAAA-3) and Cyt b-R (5'-GGCTCATTCTAGTGCCTTGTT-3') were designed from an alignment of other catostomid sequences acquired from GenBank (Table 1.1) and used to amplify 1020 base pairs of the cytochrome b gene (residues 16-356 of 380 total). A 25  $\mu$ l cocktail was prepared for each as follows: 5.0µl 1mM dNTPs, 2.5µl 10X PCR buffer, 1.0µl 50mM MgCl<sub>2</sub>, 1.0 µl 10mM Cyt *b*-F primer, 1.0 µl 10mM Cyt *b*-R primer, 0.3µl (0.5 units) Taq DNA polymerase (Invitrogen), 11.7 µl sterile ddH<sub>2</sub>O, and 2.5µl (250 ng) DNA. Reactions were incubated for 31 cycles of 94° (60 sec), 54° (45 sec), 72° (2 min) followed by a final 10 minute extension at 72°. Residual primers were removed from PCR products via incubation with shrimp alkaline phosphatase (SAP) and exonuclease I at 37° C for 20 minutes. Cycle sequencing reactions were carried out according to a modified BigDye Terminator (Applied Biosystems) protocol. Each reaction contained the following: 1.6µl primer (1mM), 1.2µl BigDye RR mix, 1.0-2.0 µl PCR products. Cycle sequencing reaction conditions were as follows: 96°C (5 min), 99 cycles of 96°C (45 sec), 62°C (1 min), 72°C (2min) followed by a final extension at 72°C (2 min). Products were visualized with a MJ Research BaseStation 51 Automated DNA Fragment Analyzer. Internal primers were also designed and used in some cases to ensure quality signal strength throughout the fragment.

Primers FTTF (5' GCCTA AGAGCATCGGTCTTGTAA 3') and F12R (5' GTCAGGA CCATGCCTTTGTG 3') were used to amplify the mitochondrial control region (922 bp). A 25 µl cocktail was prepared similar to that described above for

cytochrome *b*. Reactions were incubated with a MJ Research thermal cycling regime: 10 cycles at 94° (60 sec), 55° (60 sec), and 72 ° (60 sec) followed 21 cycles at 94° (60 sec), 53° (60 sec), 72 ° (60 sec) and a final extension at 72 ° (5 min). Products were prepared and sequenced according to the conditions highlighted above for cytochrome *b* or were shipped to the high throughput sequencing facility at the University of Washington (http://www.genome.washington.edu/UWGC/).

### Analyses

Raw sequences were edited with Sequencher 4.2 (Gene Codes) or Contig Express (Invitrogen). Multiple alignments were performed with ClustalX (Thompson *et al.* 1997) and checked by eye for obvious misalignments. Since it is a protein-coding gene, cytochrome *b* sequences were translated and the polypeptide checked for erroneous stop codons. SplitsTree 4.0 (Huson and Bryant 2006) was used to identify identical haplotypes and collapse both data sets to exclude them. Hierarchical likelihood ratio tests (hLRTs) were performed with Modeltest 3.7 (Posada and Crandall 1998) and used to determine which of 56 models of molecular evolution best explained each data set.

Phylogenetic analysis of the cytochrome *b* data set was performed with PAUP 4.0 using both maximum likelihood (ML) and maximum parsimony (MP) criteria using *Myxocyprinus* and *Carpiodes* as outgroup sequences (accession numbers AB126083 and AB223007 in Table 1.1). For both analyses, the heuristic search option (100 random addition replications using tree bisection-reconstruction) was used to search for optimal trees. Bootstrap support was estimated with 100 pseudo-replicates for ML and 1000 pseudoreplicates for MP.

Phylogenetic analysis of the control region data set was conducted using a MP approach with PAUP 4.0. In addition, the best-fit model of sequence substitution (determined with Modeltest 3.07) was used in a Bayesian analysis conducted with MrBayes 3.1.1 (Huelsenbeck and Ronquist 2001; Ronquist and Huelsenbeck 2003). This software uses metropolis-coupled Markov chain Monte Carlo (MCMC) to explore tree space, seeking topologies with the highest likelihood. In using multiple heated chains with frequent swapping, tree space is explored much more quickly than with other approaches. At this time, it is the most computationally feasible approach for large data sets – at least if one wishes to implement a particular model of molecular evolution to correct for multiple hits. Two replicates were run with three heated chains and one cold chain with sampling every 100<sup>th</sup> generation. The progress of runs was checked periodically with the program AWTY ("Are We There, Yet?") (Wilgenbusch et al. 2004), a web tool to graphically assess convergence of the MCMC algorithm by plotting posterior probabilities of a given set of 20 splits during the course of a run (Figure 1.4). Once convergence occurs, these probabilities will remain fairly constant over the duration of a run (i.e. they 'flatten out').

In an attempt to gain greater resolution at shallow nodes, a final analysis was performed with concatenated sequences for 48 individuals in whom both regions were amplified. This analysis was also performed with MrBayes3.1.1. Data partitioning was applied to account for differences in molecular evolution between the two markers (using the APLYTO command). Again, two runs were carried out with multiple heated chains to efficiently explore tree space and convergence was monitored with AWTY.

### Results

### Molecular Evolution

Collapsing identical sequences reduced the control region set to 79 unique haplotypes from 151 individuals and for cytochrome b, a reduction to 20 unique haplotypes from 48 individuals. Sequence data may be downloaded directly from GenBank (http://www.ncbi.nlm.nih.gov/Genbank/inde.g.,html) at the following accession numbers: cytochrome b: EF062360-EF062379; Control Region: EF062380-EF062458. Using the hierarchical likelihood ratio test criterion implemented in Modeltest, the TrN + I + G model (Kumar et al. 2004) was chosen for the control region data set. This model is a case of the GTR + I + G model where the parameters controlling the rates of the different types of transitions are equal. The resulting rate matrix parameter estimates were as follows: R(a) [A-C]=1.0000, R(b) [A-G]=30.9075, R(c) [A-T]=1.0000, R(d) [C-G]=1.0000, R(e) [C-T]=15.1074, R(f) [G-T]=1.0000; the proportion of invariable sites (I) was 0.8378; base frequencies were A=0.3239, C=0.2091, G=0.1578, and T=0.3092; gamma distribution shape parameter = 0.5416. The TrN+G model was chosen for cytochrome b. Rate matrix parameter estimates were R[A-C]=1.0000, R[A-G]=73.2022, R[A-T]=1.0000, R[C-G]=1.0000, R[C-T]=24.8720, and R[G-T]=1.0000; base frequencies were A=0.2605, C=0.2956, G=0.1716, and T=0.2724; gamma distribution shape parameter = 0.0153.

The cytochrome *b* gene was highly conserved in the 48 individuals examined. Among the 20 unique haplotypes, there was only a single non-synonymous substitution (Figure 1.1) that occurred in three individuals from the Rio Grande basin. Still, there were 38 variable sites among ingroup haplotypes. Phylogenetic analysis with ML (Figure 1.2) and MP (not shown) criteria yielded consistent results. The inferred topology was interesting in two respects: 1) the hypothesis of reciprocal monophyly between the two described species (*C. elongatus* and *C. meridionalis*) was not supported; 2) instead, the two described species were polyphyletic (incomplete lineage sorting) and formed a sister clade to the monophyletic Rio Grande group (Figure 1.2). Mean pairwise sequence divergence within *C. meridionalis* (0.3%) and *C. elongatus* (0.4%) was virtually no different than the mean divergence between them (0.4%); however, each was markedly divergent from the Rio Grande group (2.0% and 2.1%, respectively, see Table 1.2).

The mitochondrial control region exhibited greater variability than cytochrome b, as mean pairwise differences within *C. meridionalis*, *C. elongatus*, and the Rio Grande clade reached 0.8%. Divergence between *C. meridionalis* and *C. elongatus* was slightly higher at 1.1%, while each was 2.5% and 2.7% divergent from the Rio Grande clade, respectively (Table 1.2). MP analysis of this data set resulted in 36 most parsimonious trees with a length of 157 steps (Figure 1.3). Inspection of split posteriors during Bayesian analysis of the control region data set indicated convergence at approximately  $1.5 \times 10^6$  generations (see Figure 1.4), so the analysis was continued for another  $4.5 \times 10^6$  generations. Posterior probabilities of nodes were estimated from all sampled generations after removal of the initial  $(1.5 \times 10^6)$  burn-in and results were summarized with a 50% majority rule consensus tree (roughly equivalent to bootstrap support) (Figure 1.5). As with the cyt *b* analysis, MP topologies differed only slightly from Bayesian results at shallow nodes, while the overall patterns of *C. meridionalis / elongatus* polyphyly and divergence of the Rio Grande clade was retained.

Finally, the concatenated data set consisted of 48 individuals that, when collapsed, yielded 43 unique haplotypes. Bayesian analysis of the partitioned data set reached convergence at approximately  $1.0 \times 10^6$  generations (as indicated with AWTY) and data collection was continued for another  $3.0 \times 10^6$  generations. The analysis did result in slightly improved resolution at shallow nodes as two strongly supported *C*. *meridionalis* clades were recovered within *C. elongatus* (Figure 1.6).

### Discussion

A fundamental goal of molecular systematics is to place biodiversity in a logical evolutionary context and to enable a better understanding of the underlying principles and processes. The same is true of morphogical systematics. Unfortunately, neither approach is without caveats. To be sure, morphological studies are extremely important in tracing the evolution of any number of specific physical traits - and are essential to the field of taxonomy. At the same time, morphology may not reveal the presence of lineages that diverged millions of years ago, especially in systems that are characterized by long-term environmental stability (Colborn et al. 2001). At the other end of the spectrum, molecular studies may also fail to reveal all relevant biodiversity (Simons *et al.* 2001). While it is true that any heritable divergence in morphology must be accompanied by an underlying change in the genetic code, identifying those changes within an entire genome can be a daunting task; thus, neutral markers (e.g., mtDNA control region, introns, and microsatellites), which may accumulate mutations in the absence of morphological divergence, have become a workhorse in studies of closely related taxa. In particular, mtDNA has become the common marker of choice because its haploid nature and lack of recombination facilitate rapid lineage sorting (Avise 2004).
Here, sequences from two mitochondrial regions were used to investigate phylogenetic structure in the genus *Cycleptus*. There are two known species in the genus, C. elongatus and C. meridionalis. They are geographically disjunct and they exhibit a mutally exclusive, bimodal distribution in a wide array of meristic and morphometric characters. Nevertheless, they fail to pass the test of reciprocal monophyly, even at the mtDNA control region. Should the taxonomy be revised to synonymize C. meridionalis under C. elongatus? If a strict phylogenetic species concept is applied (e.g., Mishler and Theriot 2000), the answer is "yes," but in the author's opinion, no. Indeed, Funk and Omland (2003) recently demonstrated in a survey of more than 2319 animal species assayed with mitochondrial markers that species-level paraphyly or polyphyly occurred in 23% of the cases. If not reciprocal monophyly, then what criterion should be used to delineate a new species? While it is not my intent to burden the reader with a long diatribe on species concepts (for a relevant discussion, see Wheeler and Meier 2000), I will say that no single species concept has yet placed an appropriate frame around each and every taxon (and only those taxa) we recognize as a 'species.' The inherent problem is that we attempt to put a box around a process that is a dynamic continuum. For instance, if we were to view the tree of life with a sliding window, that window could never be small enough to exclusively – and discretely - capture only those groups we recognize as species at any given point in time. While the biological species concept (Mayr 1942) is palatable to many who study sexually-reproducing organisms (and to nonbiologists), it too, is difficult to test, as in this case.

The degree of morphological divergence between *C. elongatus* and *C. meridionalis* (Figure 1.7) (Burr and Mayden 1999) certainly suggests that they are on

independent evolutionary trajectories (Wiley 1978). So, too, does recent evidence for long-term genetic isolation between the two taxa (Chapter 4, p. 103-104).

Given the degree of genetic divergence and reciprocal monophyly between the Rio Grande and the C. elongatus/meridionalis clade, it is also clear that the Rio Grande clade has been genetically isolated for some time. In a recent study, Peng et al. (2006) analyzed mitogenomic data using a Bayesian relaxed clock approach (allowing for some rate variation among lineages) to estimate divergence times within Otocephali, a group that includes clupeomorphs (herrings, anchovies, etc.) and ostariophysans (minnows and suckers, characids, and catfishes). Their estimates were calibrated with five fossil records at various points in the tree. Five catostomid taxa were included in the data set. Among them were *Myxocyprinus asiaticus* and *Carpiodes carpio*, both of which have been hypothesized as sister to *Cycleptus* (Harris and Mayden 2001; Smith 1992). Peng et al. estimated a divergence time of 101 mya between the two taxa. This correlates to a divergence rate of approximate 0.17% / mya for cytochrome b. If we assume that a similar rate occurs in cycleptid fishes and apply it to the <u>net</u> sequence divergence (Nei 1987 p. 276) among the Rio Grande and C. elongatus / meridionalis clades (  $\approx 1.7\%$ ; also see Table 1.2), it translates to a mid-Miocene divergence at approximately 10 mya. This is not unreasonable given that one of the two aforementioned genera almost certainly harbors the closest extant relatives to *Cycleptus*. Of course, the most appropriate course would be to combine *Cycleptus* data with that from Peng *et al.* and rerun their analyses, but that is beyond the scope of this study.

Although cycleptid fishes inhabit a vast range and have been formally known for almost two centuries, limited attention has been placed on their systematics. Accurate knowledge of these relationships advances our understanding of the evolution of these fishes and may shed light on unresolved patterns in other 'old river' ichthyofauna (Simons *et al.* 2001; Smith 1981). The phylogenetic patterns revealed here corroborate and advance previous findings of Burr and Mayden (1999) and Buth and Mayden (2001). The former recognized character differences in Rio Grande specimens (e.g., color patterns and lip papillae), but used caution in interpreting these as diagnosable traits since they are subject to environmental plasticity. Based on the previous recognition of *C. elongatus* and *C. meridionalis* as valid species and the phylogenetic pattern revealed here, renewed efforts are underway to study the morphology of these creatures, especially (lip) papillae structure and tuberculation patterns (Figure 1.8). A formal recognition of the Rio Grande clade as a distinct species is in progress. This will accurately reflect evolutionary history and bolster efforts to conserve these fishes.

#### **Materials Examined**

The list of specimens included in each analysis follows. Specimen numbers follow state (and river drainage) collected from; '>' indicates a continuous series of specimens. More detailed locality and collection information appear in Appendix 2. *Micochondrial control region* 

*C. meridionalis*. Alabama, (Alabama) MLB1-1>MLB1-30; Louisiana, (Pearl) MLB51-7; Mississippi, (Leaf) MLB4-1>MLB4-5.

*C. elongatus*. Arkansas, (Red) MLB2-1>MLB2-3, (White) MLB8-21, MLB8-28; Indiana, (Wabash) MLB9-7, MLB9-22, MLB9-27; Iowa, (Mississippi) MLB10-1; Kansas, (Kansas) MLB34-3; Kentucky, (Ohio) MLB24-1>MLB24-3; Louisiana, (Mississippi) MLB40-1, MLB40-2, (Red) MLB51-5, MLB51-6, (Sabine) MLB58-

1>MLB58-7; Minnesota, (Minnesota) MLB12-9, MLB12-10, (Mississippi) MLB13-

2>MLB13-4; Mississippi, (Black) MLB59-1, MLB59-2; Missouri, (Missouri) MLB17-7;

Montana, (Missouri) MLB18B-1>MLB18B-4, (Yellowstone) MLB20-23; Nebraska,

(Missouri) MLB50-1>MLB50-4, MLB50-7>MLB50-9; North Dakota, (Missouri)

MLB23-1, MLB23-2; Oklahoma, (Red) MLB33-31; South Dakota, (Missouri) MLB25-1,

MLB25-2; Tennessee, (Cumberland) MLB28-2, (Duck) MLB51-2, (French Broad)

MLB30-9, (Nolichucky) MLB30-1>MLB30-4, (Tennessee) MLB28-1; Texas,

(Colorado) MLB45-1>MLB45-12, MLB45-14, MLB45-16, MLB45-17, MLB45-19,

MLB45-20, MLB45-23>MLB45-25, MLB45-27>MLB45-30, (Sabine) MLB51-3,

MLB51-4; Wisconsin, (Chippewa) MLB39-1, (Red Cedar) MLB39-29, (Wisconsin) MLB33-30, MLB51-1.

*C. elongatus* (Rio Grande clade). Mexico, (Conchos) MLB57-1; New Mexico, (Pecos) MLB22-1, MLB22-31>MLB22-34; Texas, (Rio Grande) MLB36-1>MLB36-20. *Cytochrome b* 

*C. meridionalis*. Alabama, (Alabama) MLB1-1, MLB1-2, MLB1-7, MLB1-9, MLB1-12, MLB1-19, MLB1-21; Mississippi, (Leaf) MLB4-3.

C. elongatus. Arkansas, (Red) MLB2-3; Indiana, (Wabash) MLB9-7; Kansas, (Kansas)

MLB34-3; Kentucky, (Ohio) MLB24-2; Louisiana, (Mississippi) MLB40-1; Minnesota,

(Minnesota) MLB12-9; Montana, (Missouri) MLB18B-4; North Dakota, (Missouri)

MLB23-2; Tennessee, (Cumberland) MLB28-2, (French Broad) MLB30-9; Texas,

(Colorado) MLB45-1, MLB45-2; Wisconsin, (Red Cedar) MLB39-30, (Wisconsin)

MLB33-30.

*C. elongatus* (Rio Grande clade). Mexico, (Conchos) MLB57-1; New Mexico, (Pecos) MLB22-1, MLB22-31>MLB22-34; Texas, (Rio Grande) MLB36-1>MLB36>20.

#### Combined data set

See cytochrome *b* set above.

#### Acknowledgements

This work was supported by generous funding from the following entities: The American Museum of Natural History (Theodore Roosevelt Fund), Sigma Xi Scientific Society, The Center for Great Plains Studies, and the University of Nebraska School of Biological Sciences. A special note of appreciation is also extended to all of the outstanding people that assisted in the lab and in the field, especially my former supervisor, Gerald Mestl, of the Nebraska Game and Parks Commission, who first introduced me to cycleptid fishes, to my graduate advisor, Dr. Guillermo Orti, who introduced me to molecular systematics, and to Dr. Richard Mayden, who enthusiastically supported my interest in this project.

Taxon	GenBank Accession Number	Source
Moxostoma erythrurum	AY253421	(Berendzen et al. 2003)
Moxostoma anisurum	AF454880	(Harris et al. 2002)
Thoburnia atripinnis	AF454911	(Harris et al. 2002)
Hypentelium roanokense	AY253420	(Berendzen et al. 2003)
Minytrema melanops	AF454879	(Harris et al. 2002)
Carpiodes carpio	AB126083	(Saitoh <i>et al.</i> 2003)
Carpiodes carpio	NC_00525	(Broughton et al. 2006)
Cycleptus elongatus	AF454868	(Harris et al. 2002)
Myxocyprinus asiaticus	AB223007	(Saitoh et al. 2005)

**Table 1.1** Taxa used for cytochrome b primer design. Sequence data listed below werealigned to identify conserved terminal regions in the catostomid cytochrome b gene.

**Table 1.2** Range and mean pairwise sequence differences within and among *Cycleptus* 

 taxa and outgroups.

	Cytochro	ome b	Control region			
Taxa	Range	Mean	Range	Mean		
ELO-ELO	0.002-0.006	0.004	0.001-0.014	0.008		
MER-MER	0.001-0.005	0.003	0.001-0.016	0.008		
CRG-CRG	0.001-0.005	0.003	0.001-0.013	0.008		
ELO-MER	0.002-0.006	0.004	0.007-0.016	0.011		
ELO-CRG	0.018-0.023	0.021	0.019-0.033	0.025		
MER-CRG	0.019-0.022	0.020	0.020-0.035	0.027		
ELO-OGR	0.146-0.159	0.109				
MER-OGR	0.146-0.159	0.108				
CRG-OGR	0.146-0.160	0.103				

ELO = *Cycleptus elongatus*; MER = *Cycleptus meridionalis*;

CRG = proposed *Cycleptus sp.* from Rio Grande Basin

OGR = Outgroup: *Myxocyprinus asiaticus*; *Carpiodes cyprinus* 

**Figure 1.1.** Cytochrome *b* translation in *Cycleptus* (amino acid residues 16-356). All substitutions were synonymous with the exception of a single point mutation at position 295 (shaded), which caused a change from serine to tyrosine in three of forty-eight individuals (all three from the Rio Grande Basin).

DALVDLPTPSNISVWWNFGSLLGLCLITQILTGLFLAM HYTSDISTAFSSVAHICRDVSYGWLIRNIHANGASFFFI CIYMHIARGLYYGSYLYKETWNIGVILLLLVMMTAFV GYVLPWGQMSFWGATVITNLLSAVPYVGNELVQWIW GGFSVDNATLTRFFAFHFLLPFVVAAATIIHLLFLHETG SNNPAGINSDADKISFHPYFSYKDLLGFAAMLLALTSL ALFSPNLLGDPENFTPANPLVTPPHIKPEWYFLFAYAIL RSIPNKLGGVLALLFSILVLMVVPILHT KQRGLTFRPI TQFLFWTLVADMIILTWIGGMPVEHPFIIIGQIASA **Figure 1.2**. Maximum likelihood phylogram for *Cycleptus* based on 20 unique cytochrome *b* haplotypes and two outgroup taxa. Tree with highest ML score is shown. Numbers to the left of nodes indicate bootstrap values (50% majority rule) greater than 50. The analysis performed with PAUP 4.0 (Swofford 1998) under the TrN+G model of molecular evolution.



**Figure 1.3**. Strict concensus of 36 most parsimonious trees (157 steps) resulting from maximum parsimony analysis of the *Cycleptus* control region alignment. Numbers at nodes indicate percent recovery in 1000 bootstrap pseudoreplicates.



**Figure 1.4.** Visual exploration of MCMC convergence with the software AWTY (Wilgenbusch *et al.* 2004). Each line represents the posterior probability of splits 1-20 during the course of a Bayesian phylogenetic analysis of mtDNA control region sequences performed with MrBayes3.1.1 (Ronquist and Huelsenbeck 2003). In this case, convergence occurred slightly beyond  $1.5 \times 10^6$  generations (X axis on the following page).



Plot of splits 1 to 20 from tmp5b443/Cumulative/out0sKZY7 sorted by widest range

**Figure 1.5**. Unrooted fifty percent majority rule phylogram resulting from Bayesian analysis of 79 unique mitochondrial control region haplotypes identified among 151 individuals. See text for specific models and prior distributions. Numbers at nodes indicate frequency of occurrence among sampled (post burn-in) tree space for highly supported clades ( $\geq$ 85). *C. elongatus* haplotypes from the Rio Grande basin are termed 'C rg.'



- 0.05 substitutions per site

**Figure 1.6**. Unrooted fifty percent majority rule phylogram resulting from Bayesian analysis of 43 unique concatenated (cytochrome b + control region) haplotypes identified among 48 individuals. Data was analyzed using a partitioned model approach. See text for specific models and prior distributions. Numbers at nodes indicate frequency of occurrence among sampled (post burn-in) tree space for highly supported clades (>85). Haplotypes from the Rio Grande basin are termed 'C rg.'



<u>0.1</u> substitutions per site

Figure 1.7. General morphological comparison of *C. elongatus* (top) and *C. meridionalis* (bottom). Images by Joseph R. Tomelleri. High resolution images available from the artist at: joe@americanfishes.com, phone 913-383-9771.



**Figure 1.8**. Lip morphology comparison in the three species of *Cycleptus*: a) *Cycleptus elongatus*, Missouri River, Nebraska (photo M. Bessert); b) *Cycleptus meridionalis*, Pearl River, Lousiana; c) *Cycleptus sp*, Rio Grande, Texas; (b) and (c) from Burr and Mayden (1999); also, (d) tuberculation in *C. elongatus*, French Broad River, Tennessee. Note that (a), (c), and (d) are from live specimens.



**(d)** 



## CHAPTER TWO

Microsatellite loci for allotetraploid blue sucker fish (*Cycleptus elongatus*, Catostomidae) with tests of cross-species amplification and isolation of paralogous loci

## Abstract

The blue sucker (*Cycleptus elongatus*) is a widespread North American catostomid fish that appears to be declining throughout much of its range. Here, we describe the isolation and characterization of eleven microsatellite loci developed for population genetic studies in the genus. We show that an additional step of cloning and sequencing can be useful in isolating paralogous loci that often co-amplify in polyploid organisms. Finally, we present results of cross-species amplifications tests in nine other taxa, including four catostomids.

Keywords: Catostomidae, Cycleptus elongatus, microsatellite, polyploidy, paralog

#### Introduction

*Cycleptus elongatus* (Catostomidae: Cypriniformes) are large fishes that inhabit main stem river channels throughout the Mississippi and Rio Grande basins as well as several disjunct gulf coastal drainages in North America. Although distributed over such a vast range, populations are not considered stable in any of 21 states where they occur. The species is endangered or extirpated in four states (New Mexico, West Virginia, Ohio, and Pennsylvania) and is a candidate for federal listing (Elstad and Werdon 1993). Some authorities suggest that these apparent trends are an artifact of inadequate sampling (Burr and Mayden 1999). Here, we describe the isolation and characterization of eleven microsatellite loci developed for population genetic studies in the species.

#### **Materials and methods**

Genomic DNA was isolated from hypaxial muscle tissue using a standard phenolchloroform extraction protocol (Sambrook *et al.* 1989), followed by ethanol precipitation. Purified samples were sent to Savannah River Ecology Laboratory (SREL - University of Georgia) for microsatellite enrichment with GATA, CATA, CA, and CT probes. The enrichment protocol is summarized by Hauswaldt and Glenn (2003). Subsequent revisions may be obtained directly from Travis C. Glenn (<u>glenn@srel.edu</u>).

Enriched DNA was amplified with the following cocktail: 2.0µl of 1mM dNTPs, 2.5µl of 10X PCR buffer, 2.5µl BSA (250ng/µl), 1.3µl SuperSNX-24 linker (10  $\mu$ M) provided by SREL, 1.0µl of 50mM MgCl<sub>2</sub>, 0.3µl (0.5 units) of Taq DNA polymerase (Invitrogen), 13.4µl sterile ddH<sub>2</sub>O, and 2.0µl eluted DNA fragments from SREL. Cycling conditions were: 95°C (2 min), 35 cycles at 95°C (20 sec), 60°C (20 sec), 72°C (1.5 min), followed by a final extension at 72°C (30 min). PCR products were cloned using the pGEM-T-Easy Vector System I (Promega). PCR-based screening was performed to determine insert length in approximately 200 positive clones. Inserts were amplified using flanking primers M13F and M13R according to the following conditions: 96°C (5 min), 30 cycles of 96°C (45 sec), 60°C (1 min), and 72°C (1min) with a final extension of 72°C (5 min). Products were electrophoresed on a 2% agarose gel to visualize insert length. Approximately 100 clones with inserts > 350bp were selected for sequencing. Clones were sequenced in one direction with a BigDye Terminator cycle sequencing reaction (Applied Biosystems) using the M13F primer. Complimentary (reverse) sequences were obtained for fifty-nine clones that contained microsatellite sequences using the M13R primer. Complimentary pairs were aligned and edited with Sequencher<sup>™</sup> 4.2.2 (Gene Codes Corp.).

PrimerSelect© (DNASTAR Inc) was used to design primer pairs for 19 candidate loci that possessed adequate flanking regions (at least 50bp) and at least seven repeat units of the microsatellite motif. PCR optimization was conducted using a gradient thermal cycler (MJ Research PTC-200) with an annealing step of 48-66°C to determine optimum annealing temperature. Reaction mixtures had a total volume of 10µl and contained 2.0µl of 1mM dNTPs, 1.0µl of 10X PCR buffer, 0.4µl of 50mM MgCl<sub>2</sub>, 0.4µl 10mM forward primer, 0.4µl 10mM reverse primer, 0.1µl (0.5 units) of Taq DNA polymerase (Invitrogen), 4.7µl sterile ddH<sub>2</sub>O, and 1.0µl (≈100ng) DNA. Reactions were denatured at 94°C for 2 minutes, then carried out for 30 cycles at 94°C (30 sec), annealing temperature (30 sec), 72°C (40 sec), followed by a final extension of 72°C (2 min). Fifteen loci amplified cleanly. For these, an ABI Prism 310 Genetic Analyser (Applied Biosystems) was used to score allele lengths in 30 individuals collected from a contiguous 150 km stretch of the upper Missouri River, Montana, USA.

## **Results and Discussion**

Six loci exhibited tetrasomic genotypes (more than 2 alleles) while the other nine were disomic (Table 2.1, loci Ce35-215). The number of alleles per individual was always one or two and the number of alleles per locus ranged from 1-27 with a mean of 10.2 alleles per locus. Three loci (Ce49, Ce126, and Ce215) deviated significantly from Hardy-Weinberg (HW) equilibrium as determined by the HW probability test implemented in GENEPOP 3.1 (Raymond and Rousset 1995).

Although sophisticated techniques abound for population genetic analysis of diploid, codominant markers (Pearse and Crandall 2004), there remains a paucity of methods to analyze polyploid data. Therefore, we sought to increase the number of diploid markers by isolating two co-amplifying (paralogous) loci from one another. A candidate locus (primer pair Ce13F: 5'-GTTCTGGGACTTAACAAAGGGATTT-3', Ce13R: 5'-ACAAACATGAGGTATCAAGTAGTCTAA-3') was amplified by PCR and the products from a single individual that possessed four alleles of distinct length (at least 10 bp different from one another) were cloned and sequenced. Pair-wise length differences (in base pairs) for each combination of alleles were as follows:  $\{1,2\}=13$ ;  $\{1,3\}=25$ ;  $\{1,4\}=55$ ;  $\{2,3\}=12$ ;  $\{2-4\}=42$ ;  $\{3,4\}=30$ . Allele sequences were aligned to identify regions that differed between paralogs and new primers (Ce13S and Ce13L, Table 2.1) were designed at these sites. Each new primer pair amplified only two alleles in the sequenced individual and paralog identities were confirmed by comparing allele lengths to those amplified by the original Ce13 primer pair. Ce13S products differed by 55 bp, corresponding to pair  $\{1,4\}$ ; Ce13L products differed by 12 bp, corresponding to pair  $\{2,3\}$ .

Loci Ce13S and Ce13L were also helpful in determining chromosomal inheritance patterns in *C. elongatus*. In an organism of autopolyploid origin, a multivalent formation (of more than two chromosomes) may occur during meiosis (Wua *et al.* 2001); thus, some individuals would fail to inherit at least one priming site for one of the two loci (resulting in no amplification at that locus). Every individual in this screening set possessed either one or two alleles and the results did not differ significantly from HW expectations (Table 2.1). These results confirm functional diploidy in *C. elongatus* and are consistent with an allotetraploid origin as suggested by Uyeno and Smith (1972). Finally, cross-species amplifications were attempted to determine primer efficacy in nine other taxa. Reactions were conducted in single individuals with positive and negative controls. All products were verified by genotyping as described previously. As expected, some loci are well-conserved within Catostomidae (Table 2.2). In conclusion, these novel markers will be useful for population genetic studies within the genus and may have broader utility within Catostomidae.

#### Acknowledgements

This research was made possible by generous support from the following sources: Initiative for Ecological and Evolutionary Analysis (University of Nebraska), Center for Great Plains Studies (Lincoln, NE), the American Museum of Natural History (Roosevelt Fund), Sigma Xi, Special Funds (UNL) School of Biological Sciences, and the Nebraska Game and Parks Commission.

Locus/ GenBank								
Accession no.	Primers (5 <sup>2</sup>	→ 3')	Repeat motif	Size (bp)	$T_a$ (°C)	k	$H_O$ †	$H_E$
Ce13S DQ401677	F: ATAACAAG R: CCGAGGA	CTTGTCATGCATTCCTGA CAGCGGTTTAAAATAT	(GATA) <sub>20</sub>	114-186	60	24	0.900	0.942
Ce13L DQ401678	F: GTAACAA R: GCCGAGG	ITTTTCATGCATTCCTGGA ACAGCGGTTTA	(GACG) <sub>5</sub> (GATG) <sub>3</sub> (GATA) <sub>4</sub>	132-154	60	6	0.800	0.687
Ce35 DQ401679	F: CTTCACAC R: TGGCAGC	CCAGCTCAAGTCACAT CTAAGCTTAATGCTCTA	(GA) <sub>17</sub>	123-151	60	11	0.833	0.856
Ce49 DQ401680	F: TTTAAGAT R: GAATGTG	TTTCTTCCTTCGACTAA CCCGTGCGCATGAACA	(CAA) <sub>7</sub>	107-113	60	3	0.533*	0.538
Ce52 DQ401681	F: ATGACAG R: GTTTCCAT	CATCCATGCACATTTA 'GGATACCAATTTACCC	(CAAT) <sub>8</sub> CACT(CAAT) <sub>13</sub>	229-241	64	4	0.433	0.534
Ce63 DQ401682	F: CCAAAAG R: CAGACGG	CGTCTTGAAATGTTCA CGAGAGGAGATGGA	(GT) <sub>8</sub> (GA) <sub>14</sub>	145-255	64	18	0.733	0.829
Ce104 DQ401683	F: CACACCCA R: GATACAG	TTACGGCAGGATTA CAATGAGCTTTCATAACACA	(CT) <sub>18</sub>	145-153	60	5	0.567	0.491
Ce126 DQ401684	F: TTCGCTCT R: TGGAGAG	CCGTCCCTTTCATTCT CGAAAAAGAGACATTATCA	(CT) <sub>12</sub>	155-179	60	9	0.967**	0.832
Ce146 DQ401685	F: AACCCAAA R: TGCTCGCT	AAATGAAAATTGTGTTA TATTAAGAGACTCTGATT	(GATA) <sub>8</sub>	150-164	60	4	0.733	0.697
Ce195 DQ401686	F: ACATTGCC R: TCCATCCT	GATTAATTGCATTCATT TCTTCTGCCATTACATT	(CATA) <sub>2</sub> CG(CATA) <sub>18</sub>	243	60	1	0.000	0.000
Ce215 DQ401687	F: TTGTCACA R: CACTCTCA	CCTTTATGGGATTCAT ATAGCGAAATGTAGTTCTT	(TATC) <sub>16</sub> C(TATC) <sub>14</sub>	211-287	60	27	0.767***	0.963

**Table 2.1** Primer sequence, repeat motif, and allelic diversity for eleven microsatellite markers in *Cycleptus elongatus*, each screened in 30 individuals.  $T_a$ , annealing temperature; k, number of alleles;  $H_O$ , observed heterozygosity;  $H_E$ , expected heterozygosity

†Three loci deviated significantly from Hardy-Weinberg equilibrium: Hardy-Weinberg equilibrium probability test, GENEPOP 3.1 (Raymond and Rousset 1995). \*P=0.036, \*\*P=0.020, \*\*\*P=0.010.

**Table 2.2** Cross-taxon amplification to determine priming site conservation. Numbers below each locus represent actual genotypes (allele lengths in base pairs) in a single individual of the given taxon. "X" indicates no successful amplication. All reactions were conducted under conditions optimized for *C. elongatus*.

Taxon	Locus Ce13S	Ce13L	Ce35	Ce49	Ce52	Ce63	Ce104	Ce126	Ce146	Ce195	Ce215
Cycleptus meridionalis	99/123	145/145	Х	113/113	202/202	159/173	152/152	175/185	Х	245/245	179/213
Carpiodes cyprinus	155/175	Х	131/146	113/113	171/171	Х	127/127	155/155	148/148	245/245	Х
Catostomus commersoni	200/204	Х	Х	113/113	200/200	Х	140/140	154/156	Х	Х	Х
Moxostoma macrolepidotum	Х	Х	Х	113/113	Х	Х	Х	Х	Х	245/245	Х

Additional taxa for which all amplification attempts failed: *Pimephales promelas, Ameiuras melas, Prochilodus lineatus, Lepomis cyanellus,* and *Culaea inconstans.* 

# Population genetic structure of the Blue Sucker (*Cycleptus elongatus* Lesueur, 1918) in the Upper Missouri River: genetic effects of habitat fragmentation

#### Abstract

In recent times, anthropogenic modifications have caused profound changes to North American waterways. Channelization and impoundment have homogenized seasonal flow regimes and obstructed upstream migration of many large riverine fishes. The blue sucker, Cycleptus elongatus, is a large catostomid fish that occurs in main stem rivers throughout the Mississippi basin of North America. Although not federally listed as threatened or endangered, populations are not considered stable in any of 21 states where they occur. Included in this range is the Missouri River, which flows more than 3200 kilometers from its headwaters in Montana to its confluence with the Mississippi River at St. Louis, Missouri. Historically, C. elongatus was distributed continuously throughout the mainstem Missouri and its major tributaries, but from 1952-1963, six major impoundments were constructed on the upper Missouri by the US Army Corps of Engineers. The resulting reservoirs have inundated and fragmented riverine habitat from Yankton, South Dakota to the headwaters in Montana. Cycleptus elongatus still occurs in the remnant unimpounded stretches between reservoirs; however, little is known of reproduction, recruitment, and whether inter-reservoir populations are genetically isolated. In order to assess genetic diversity and connectedness, 231 individuals from nine sites in the drainage were genotyped at 15 variable microsatellite loci. Genetic data

were used to calculate traditional summary statistics and fixation indices. In addition, four techniques were used to investigate population structure. The results indicate shallow, 'graded' structure from the lower reaches toward the headwaters. Mantel tests revealed a highly significant pattern of isolation by distance. This is noteworthy because such a pattern does not exist in the unobstructed Mississippi River main stem, a river of comparable size. These results are consistent with reduced intradrainage gene flow in the Missouri River and may represent the first traces of impoundment effects on genetic structure. This information will assist governing agencies in making well-informed decisions regarding conservation and management of *C. elongatus* in the Missouri River basin.

### Introduction

The past 100 years have witnessed a great deal of anthropogenic change to natural waterways in North America. Some rivers have been impounded to store water and regulate flow while others have been channelized to facilitate navigation. In addition, locks and dams have been constructed on many others to further control water levels for navigation. Migration and reproduction in many riverine fishes has been negatively impacted by such modifications because routes to historical spawning grounds have been compromised (Ickes *et al.* 2001; Jungwirth *et al.* 1998; Laroche and Durand 2004).

The blue sucker, *Cycleptus elongatus*, is a large catostomid fish native to main stem rivers throughout the Mississippi basin and is one of the most widespread lotic fish taxa in North America. Its elongate, hydrodynamic body form, with paired fins that are anteriorly rounded, enables it to maintain its position in swift current with little energy expenditure (Wolter and Arlinghaus 2003). Historically, the species occurred in 21 conterminous states, but it is now endangered or extirpated in four (New Mexico, West Virginia, Ohio, and Pennsylvania) (NatureServe 2005) and is a candidate for listing at the federal level (Elstad and Werdon 1993). The range of *C. elongatus* has diminished greatly over the past 100 years, and it is thought that the impoundment of main stem rivers has played a major role (Boschung and Mayden 2004; Robison and Buchanan 1998). In the upper Missouri drainage, *C. elongatus* inhabited virtually all reaches of the mainstem Missouri River from its confluence with the Mississippi River at St. Louis to the far northern headwaters of the Missouri and Yellowstone rivers in Montana (Brown 1971; Underhill 1959), a distance of more than 3200 kilometers.

From 1952-1963, seven major impoundments were constructed on the upper Missouri River by the Army Corps of Engineers in an effort to control flooding and provide hydroelectric power (Hesse *et al.* 1982). The resulting reservoirs have inundated and fragmented large riverine habitat from Yankton, South Dakota to the Missouri River headwaters in Montana, reducing mainstem riverine habitat in the intervening stretches by some 70% (estimate based on length of reservoirs at full pool – see USACE 2001). Following construction of the dams, blue suckers and other migratory species have amassed in the tailwaters at times coinciding with seasonal spawning movements (Eitzmann *et al.* 2005; Walburg *et al.* 1971). Blue suckers are, indeed, riffle spawners that swim up tributaries to reach suitable habitat (i.e. flooded gravel bars).

Although *C. elongatus* still occurs in the reaches between reservoirs, it is unknown whether these represent viable (sustainable) populations and whether any gene flow occurs between them. At best, there is enough reproduction and recruitment to maintain genetic diversity and allow for at least some level of downsream gene flow and at worst there is little to no reproduction and recruitment and complete genetic isolation between inter-reservoir stretches. Here, the fundamental working hypothesis was that little, if any, downstream gene flow occurs because adults are not suited to the lentic characteristics of reservoirs and probably avoid them. Also, larval drift from tributaries above reservoirs may settle to the bottom or be consumed by predators before it can pass through floodgates or turbines, thus limiting gene flow and recruitment.

From this point hereafter, the term "population" refers to the evolutionary paradigm set forth by Waples and Gaggiotti (2006), that is, a population is defined as, "*a group of individuals of the same species living in close enough proximity that any* 

*member of the group can potentially mate with any other member.*" In this context, the following hypotheses regarding impoundment effects were tested in Missouri River populations: 1) inter-reservoir populations in the upper Missouri River will show an overall reduction in genetic diversity compared to those in open sections of the lower river; 2) genetic variance between populations will be non-negligible, arbitrarily accounting for at least 5% of the total variance observed (due to reduced gene flow and increased effects of genetic drift); 3) Missouri River populations will show a significant pattern of isolation by distance due to a lack of homogenizing gene flow; and 4) interreservoir populations will show a signal of recent decline in number as evidenced with an excess in observed heterozygosity.

#### Materials and methods

#### Study location and individual collection

A total of 231 blue suckers were collected from six main stem localities and three tributaries of the Missouri River from November 2004 to November 2005 (Figure 3.1, Table 3.1). To ensure adequate representation throughout, I specifically targeted all inter-reservoir populations and employed stratified sampling in the lower reaches (below Lewis and Clark Reservoir, the first impoundment). Fish were captured using hoop nets, gill nets, and electroshocking devices. A small (1 cm<sup>2</sup> or less) fin clip was removed from each fish and preserved in 95% EtOH for subsequent genetic work. All tissues are vouchered in the personal collection of the author (see Appendix 2) and are freely available upon written request.

#### DNA preparation and amplification
DNA was extracted from tissue samples using either a standard phenolchloroform protocol (Sambrook *et al.* 1989) or a DNeasy® Tissue purification kit (Qiagen). DNAs were eluted in either water of buffer (EB) supplied by Qiagen. A small number of samples were randomly selected and (1-2  $\mu$ l) electrophoresed through a 1% agarose gel to check for quality (high molecular weight). All samples were quantified with a GeneQuant II spectrophotometer (Pharmacia Biotech) and a portion of each was diluted to a working stock concentration of 100ng/ $\mu$ l. The remainder of each elution was placed in -70° C for long-term storage. As with tissue samples, DNA samples are maintained in the personal collection of the author housed at the University of Nebraska (lab of G. Ortí) and are freely available upon written request.

Eleven microsatellite loci previously isolated from *Cycleptus elongatus* (Bessert *et al.* 2006) were chosen for this study. In addition, sixteen primer pairs designed for other catostomid taxa were screened and optimized in *C. elongatus*. This resulted in the addition of four more loci: Mox294, Mox306, Mox329 (Lippe *et al.* 2004), and Dlu4235 (Tranah *et al.* 2001).

In order to increase the efficiency of data collection, primer pairs were screened for potential cross-reactivity in multiplex reactions using the software AutoDimer (Vallone and Butler 2004) and two multiplex reactions were optimized according to guidelines provided by Henegariu *et al.* (1997). If two loci (in the same reaction) were labeled with the same fluorophore, special care was taken that the allelic size ranges were non-overlapping. In this way, I was able to successfully amplify all 15 loci in only two reactions, as follows (also see Table 3.2). Note that all primers were at 10µM "working stock" concentrations. Reaction "A" (Figure 3.2) contained the following: 1.60 µl dNTPs (1mM), 1.20 µl PCR reaction buffer (10X), 0.80 µl MgCl (50mM), 0.30 µl 13S-F\*, 13S-R, Ce52F\*, and Ce52R, 0.32 µl Ce35F\*, Ce35R, Ce126F\*, Ce126R, Mox329F\*, and Mox329R, 0.50 µl Ce215F\* and Ce215R, 0.48 µl Mox306F\* and Mox306R, 0.20 µl Ce104F\* and Ce104R, 0.12 µl Taq DNA polymerase (Gibco BRL), and 1.0 µl DNA (=100ng) for a total volume of 10.20 µl. Thermal cycling conditions were as follows: 94°C (1 min), 30 cycles of 94°C denature for 30 sec, 55.8°C anneal for 40 sec, and 65°C extension for 2 minutes, followed by a final extension of 65°C for 5 minutes and a holding temperature of 4°C. Reaction "B" (Figure 3.3) contained the following: 1.60 µl dNTPs (1mM), 1.20 µl PCR reaction buffer (10X), 0.88 µl MgCl (50mM), 0.26 µl 13L-F\* and 13L-R, 0.44 µl Dlu4235F\* and Dlu4235R, 0.30 µl Ce49F\* and Ce49R, 0.48 µl Ce63F\* and Ce63R, 0.28 µl Ce146F\* and Ce146R, 0.32 µl Ce195F\* and Ce195R, 0.54 µl Mox294F\* and Mox294R, 0.14 µl Tag DNA polymerase (Gibco BRL), and 1.0  $\mu$ l DNA (=100ng) for a total volume of 10.06  $\mu$ l. Thermal cycling conditions were the same as those for reaction "A" except the annealing temperature was 54.8° C. Note that 'pigtail' modifications were added to the 5' end of reverse (unlabeled) primers for loci Ce35, Ce63, and Ce104 in order to reduce stutter and ease scoring (see Brownstein *et al.* 1996). For the specific labeling scheme of each reaction, see Table 3.2.

## Sample preparation and data collection

Prior to genotyping, PCR products were purified with a Mini-Elute (Qiagen) or Microarray PCR purification kit (Telechem International, Inc.) to remove residual unincorporated dyes that can obscure allele signatures in an electropherogram (see Butler 2002). Products were prepared for electrophoresis by mixing (0.5 µl PCR product) with 0.5 μl LIZ 500 size standard (Applied Biosystems) and 9.0 μl deionized formamide. Samples were denatured at 95° C for 3 minutes and quenched on ice for two minutes prior to capillary electrophoresis on an ABI 310 fragment analyzer. Allele sizes were scored with GeneMapper 3.7 (Applied Biosystems) and the data transformed to GenePop format (Raymond and Rousset 1995) with GMCONVERT 0.32 (Faircloth 2006).

Genotypic data were checked for typographical errors, evidence of null alleles (Hardy-Weinberg equilibrium), and adherence to the Stepwise Mutation Model (Ohta and Kimura 1973) with MicroChecker 2.2.3 (VanOosterhout *et al.* 2004) and MSAnalyzer 4.0 (Dieringer and Schlötterer 2002). Additional transformations of the data set to other formats was performed with CONVERT 1.3.1 (Glaubitz 2004).

# Data Analysis – Summary statistics and fixation indices

In addition to observed and expected heterozygosities for each locality, deviations from Hardy-Weinberg equilibrium and pairwise  $F_{ST}$  estimates (Weir and Cockerham 1984) were calculated for each locality using GenePop3.3 (Raymond and Rousset 1995). The program FSTAT 2.9.3.2 was used to calculate the average allelic richness (A) at each locality (Goudet 1995; Goudet 2002). This is a useful method for comparing samples of different sizes because it scales all results to the smallest sample. Since 12 loci did not deviate from expectations of the stepwise mutation model (SMM) (Ohta and Kimura 1973; Valdes *et al.* 1993; Wehrhahn 1975), they were used to calculate *rho*, an estimator of R<sub>ST</sub> values (Valdes *et al.* 1993), with RSTCALC (Goodman 1997) and to perform pairwise comparisons. Note that Bonferroni corrections were applied to adjust for multiple pairwise comparisons (Rice 1989). Pairwise t-tests were performed to test for differences in mean allelic richness between free ranging populations (sites A-D) and impounded populations (E-I).

## Population structure

In order to assess the distribution of genetic variation within and among collection sites as well as that between impounded (sites E-I) and unimpounded (sites A-D), an analysis of molecular variance (AMOVA) (Excoffier *et al.* 1992) was performed with Arlequin 3.1 (Excoffier *et al.* 1992; Schneider *et al.* 2000). This test was executed under the standard model ("different alleles are considered mutationally equidistant from each other") as well as the stepwise mutation model. Two loci were excluded from the latter due to inconsistencies with the SMM (gaps in allele frequency distribution).

Another useful method that does not require *a priori* assumptions regarding population structure is the Bayesian clustering technique implemented in STRUCTURE 2.0 (Pritchard *et al.* 2000). Here, genetic data is used to define the number of subpopulations (k) in the absence of any locality data. The method assumes Hardy-Weinberg equilibrium (HWE) within subpopulations and seeks combinations of individuals that maximize HWE in a pre-determined number of populations. To use the method, one performs several runs at values of k from 1 to n where n = the maximum number of subpopulations to be considered. Assuming quality control in preparation and performance of these runs (unambiguous data, quality markers, adequate burn-in, etc.), the k value eliciting the highest mean posterior probability over several runs indicates the true number of genetic subdivisions within the data set. Four simulations were executed at value of k from one to nine under the admixture model with independent allele frequences and  $\lambda$  fixed at 1; that is, the model assumes that allele frequencies in each subpopulation are independent from one another (i.e. independent draws from a distribution defined by  $\lambda$ ). A set of preliminary simulations was performed to determine appropriate burn-in and run lengths. Real time plots of  $\alpha$  (parameter for the degree of admixture) and other parameter estimates were monitored to determine when convergence occurred. A conservative burn-in period of 250,000 generations was selected for subsequent simulations and was followed by 1,000,000 generations of data collection. Posterior probabilities for *k* were computed based on the mean log likelihood of the data from 4-5 simulations at each value of *k* (see Pritchard and Wen 2004).

Finally, I referred back to the summary statistics and performed several analyses to determine whether the data fit a model of isolation by distance (IBD) (Wright 1943). This model predicts that gene flow will be negatively correlated with geographic distance between populations. When the homogenizing effects of gene flow are reduced, the effects of drift become more pronounced and distant populations tend to diverge at neutral genetic markers. In order to test for IBD, central localities (CL) for each sampling area were estimated from locality plots of each individual collected, and a matrix of linear distances between the central locality for all pairs of populations was calculated (Table 3.5). The web-based software package IBDWS (Jensen *et al.* 2005) was used to calculate a matrix of pairwise genetic differentiation between populations between the matrix of linear distance and the matrix of pairwise genetic differentiation between populations between the matrix of linear distance and the matrix of pairwise genetic differentiation between each of the

following matrix combinations: genetic differentiation vs ln (linear distance); ln (genetic differentiation vs linear distance; and ln (genetic differentiation) vs ln (linear distance).

# Demography

BOTTLENECK 1.2.02 (Cornuet and Luikart 1996) was used to investigate the possibility of recently reduced effective population sizes in the sampling areas. The method compares gene diversity (expected heterozygosity,  $H_e$ ) with the expected equilibrium gene diversity ( $H_{eq}$ ) that is computed from *na* (the observed number of alleles) under mutation/drift equilibrium. If a significant number of loci show an excess in gene diversity then the population has likely undergone a recent bottleneck. A Wilcoxon sign-rank test is used to make this determination. The two-phase mutation model (TPM) was used because it more realistically describes microsatellite evolution than either the strict stepwise mutation (SMM) or infinite alleles (IAM) model. In the TPM, a 95% frequency of stepwise mutations was assumed with a 12% variance of multiple-step mutations.

#### Results

### Marker screening

One locus, Ce63, showed a heterozygote deficite across all sample sites. This pattern is consistent with the presence of null alleles; thus, we did not use this locus in any analyses. In addition, two loci, Mox329 and Ce215 had significant gaps in their allele frequency distributions. While they did not deviate from HW expectations, the mutational pattern clearly deviated from a stepwise (SMM). Consequently, these two markers were withheld from any analyses that assumed the stepwise model.

## Summary statistics and fixation indices

Summary statistics are presented in Table 3.1. Note that site B was excluded from these analyses due to a very small sample size (n=6). For the remaining sites, gene diversity (=observed heterozygosity) was only slightly reduced in impounded populations, and not significantly so (mean<sub>A-D</sub> = 0.701, mean<sub>E-I</sub> = 0.697; t = 0.544, df = 6, p = 0.61). In a similar site comparison of mean allelic richness, the upper Missouri did prove to be significantly reduced in comparison to the lower Missouri (mean<sub>A-D</sub> = 7.31 mean<sub>E-I</sub> = 6.75; t = 2.68, df = 6, p = 0.036), supporting the hypothesis of reduced genetic diversity in the impounded populations. When this analysis was extended to include additional unimpounded sites from throughout the Mississippi basin (see Chapter 4, Table 4.1), the difference was even greater (mean<sub>others</sub> = 7.31, mean<sub>E-I</sub> = 6.75; t = 3.68, df = 15, p = 0.002). Pairwise R<sub>ST</sub> comparisons appear in Table 3.3. Following Bonferroni correction, five significant pairwise differences remained. Four of these occurred between pairs that were among the most distant from one another, hinting at the possibility of isolation by distance.

#### *Population structure*

Both analyses of molecular variance revealed that more than 98% of the total variation occurred within populations, that is, within sample sites (Table 3.4). Roughly 1% occurred among populations above and below the lowest impoundment (sites A-D vs E-I), while a very modest amount (est. 0.38% and 0.31%) occurred between populations; thus, refuting hypothesis 2.

For the Bayesian analysis with STRUCTURE 2.0 (Pritchard *et al.* 2000), the setting of k = 2 yielded the highest mean posterior probability over 4-5 separate

simulations for each value of k from 1-5. The results of the single run with the highest posterior probability appear in Figure 3.4. Pritchard and Wen (2004) suggest caution in interpreting results when a value of k = 3 or less is indicated; however, the slightly graded pattern here is valid as the number of individuals assigned to each cluster is unequal. A questionable signal would include an approximately equal number of individuals in each cluster (Pritchard and Wen 2004). Interestingly, this pattern is unique to the Missouri River when analyzed as part of a range-wide data set (see Chapter 4, Figure 4.3), further suggesting that it represents real structure.

Mantel tests for isolation by distance were highly significant (p < 0.001) for all four matrix combinations, i.e. genetic distance and ln(genetic distance) vs geographic distance and ln(geographic distance). Results for ln(genetic distance) vs geographic distance are presented in Figure 3.5. Slopes and negative correlations were similar for the other three tests, indicating a clear pattern of IBD in the Missouri River.

## Demography

Multilocus frequencies of observed heterozygosity were slightly higher than expected at seven of the nine sampling locations; however, no recent bottlenecks were detected at any of the sample sites (see Table 3.5).

## Discussion

Is there any reason to believe that *C. elongatus* populations in the Missouri River are declining – or that dams have had any genetic impact to this point? At first glance, the answer is unclear. Gene diversity is somewhat uniform throughout. In addition, no significant heterozygosity excess was detected in any of the sampled areas; thus, with this method, there is no evidence for any recent bottlenecks (Table 3.5). However, allelic richness differed significantly between populations above and below the lowest impoundment (see Table 3.1). A reduction in allelic diversity is frequently associated with invasive (Genton *et al.* 2005) or founding populations (Ramstad *et al.* 2004), but may also indicate population decline (Faugeron *et al.* 2004).

Should we expect to see any deviations from 'normal' (undifferentiated) genetic patterns given the brief time frame of approximately 60 years? To be sure, the array of molecular tools employed is powerful enough to detect any of these effects, if present (Table 2; mean number of alleles (na) = 122 in each population); however, it is also important to consider the generation time for *C. elongatus*, which is approximately 10 years (Becker 1983). Given the accompanying lag time, detection of a significant bottleneck (Cornuet and Luikart 1996) would indicate an extreme situation for the affected populations, and this is not the case (Table 3.5).

Conversely, a highly significant pattern of isolation by distance was detected among Missouri River populations (Table 3.6, Figure 3.5). While this may seem intuitive or trivial given that sampled populations span a distance of nearly 3,050 river kilometers, cycleptid fishes are known to migrate hundreds of kilometers to spawning grounds (Mettee and Shepard 1997). In an unobstructed waterway of equal length, it is conceivable that these behaviors could mitigate distance effects and elicit a population signature of - or tending toward - panmixia. In this light, it is worthy to note that the majority of main stem tributaries (e.g., Red, Missouri, Ohio, and Tennessee) within the Mississippi Basin have been impounded at some point (Benke and Cushing 2005) while the main stem Mississippi itself has not. A long series of locks and dams was constructed over a distance of several hundred kilometers in the upper Mississippi, but fish passage is still possible through these corridors (Ickes *et al.* 2001).

To test for potential differences between the two major rivers, equivalent genotypic data was collected from an additional 135 individuals from 6 sites on or near the Mississippi's main stem (Table 3.7; also see Figure 4.2, Chapter 4) and combined with data from the lowest Missouri River site (Table 3.1, site A) for a total of 172 individuals distributed over seven sites spanning 2,493 unobstructed river kilometers. The results of a Mantel test for isolation by distance comparing genetic differentiation and geographic distance in this data set were nonsignificant (Figure 3.6). The other three possible matrix combinations - genetic differentiation vs ln (linear distance); ln (genetic differentiation) vs linear distance; and ln (genetic differentiation) vs ln (linear distance) were tested and also proved nonsignificant.

While this case provides only a single - albeit expansive - comparison with the Missouri River, it warrants concern. A recent study in gallinaceous birds with a similar generation time (10 years) revealed declines in genetic diversity and concomitant shifts toward a metapopulation structure with isolation by distance after only 50 years of anthropogenic fragmentation (Segelbacher *et al.* 2003). In the Missouri River, these highly significant correlations may be the first detectable signal of genetic change due to isolation by impoundment. Careful monitoring of populations in the upper Missouri with periodic combinations of physical and genetic surveys is encouraged.

#### Acknowledgements

A major portion of this work was generously supported by a State Wildlife Grant from the Nebraska Game and Parks Commission. Other significant sources included the Center for Great Plains Studies, the UNL School of Biological Sciences, and the Initiative for Ecological and Evolutionary Analysis. Again, a special note of thanks is extended to the many people who assisted with sample collection (see listed individuals and agencies in Appendix 2) as well as all the members of the Orti lab, who patiently allowed exclusive access to our fragment analyzer for weeks on end.

**Table 3.1** Site-specific summary statistics; CL = central locality (latitude/longitude) of sampling area; <math>n = number ofindividuals sampled; na = number of alleles; Gene diversity ( $H_{O}$ ) = observed heterozygosity;  $H_E$  = expectedheterozygosity; A = allelic richness.

	River / State	CL	п	na	Gene diversity (H <sub>0</sub> )	$H_{\rm E}$	A
А	Missouri / Missouri	38°68'48" N 90°66'96" W	30	140	0.700	0.688	7.340
В	Kansas / Kansas	39°11'20" N 96 °31'03" W	6	62	0.576	0.729	-
С	Missouri / Nebraska	40°81'49" N 95°84'43" W	38	138	0.704	0.745	7.290
D	Platte / Nebraska	41°05'13" N 96°10'83" W	23	128	0.698	0.661	7.311
E	Missouri / South Dakota	42°76'47" N 97°98'81" W	29	142	0.717	0.692	7.271
F	Missouri / North Dakota	47°25'42" N 101°24'34" W	14	96	0.690	0.698	6.343
G	Yellowstone / Montana	46°43'36" N 96°31'03" W	30	134	0.691	0.732	6.629
Н	Missouri / Montana	48°03'42" N 106°91'72" W	31	132	0.699	0.727	6.902
Ι	Missouri / Montana	47°73'71" N 109°61'83" W	29	125	0.686	0.675	6.591
totals			n = 231	total k = 194	mean = 0.698	0.702	mean = 7.554

**Table 3.2.** Locus name, repeat type, allelic size range in screening population,fluorophore label, and color of fifteen microsatellite markers used in this study.

	Locus	Repeat type	Allelic size range (bp)	Fluorophore	Color
	Ce13S	tetra	120-192	6-FAM	blue
	Ce35	di	124-140	PET*	red
	Ce52	tetra	233-245	6-FAM	blue
"Y,,	Ce126	di	166-178	NED*	yellow
ction	Ce215	tetra	215-303	NED*	yellow
Read	Mox306	tetra	175-227	VIC*	green
	Mox329	tetra	158-218	PET*	red
	Ce104	di	144-152	VIC*	green
	Ce13L	tetra	142-162	6-FAM	blue
	Dlu4235	tetra	131-187	PET*	red
B"	Ce49	tri	106-109	6-FAM	blue
ion "	Ce63	di	152-268	VIC*	green
eacti	Ce146	tetra	144-156	NED*	vellow
R	Ce195	tetra	240-244	NED*	vellow
	Mox294	tetra	227-271	PET*	red

\*Proprietary dyes from Applied Biosystems, Inc.

					R <sub>ST</sub>				
		А	С	D	Е	F	G	Н	Ι
ction	А	-	0.0019	-0.0028	0.0014	0.0078	*0.0126	0.0113	* 0.0572
corre	С		-	-0.0003	-0.0026	0.0127	0.0072	*0.0118	* 0.0488
rroni	D			-	0.0057	0.0136	0.0057	0.0074	* 0.0686
Bonfe	Е				-	-0.0052	-0.0038	-0.0029	0.0263
after	F					-	-0.0110	0.0043	0.0120
cance	G	*					-	-0.0009	0.0258
ignifi	Н							-	*0.0572
S.	Ι	*	*	*				*	-

**Table 3.3**. Pairwise  $R_{ST}$  comparisons for all collection sites in the Missouri River. Bold text indicates significant differencesafter Bonferroni corrections for multiple comparisons.

Table 3.4. AMOVA summary under the standard model (A) and the microsatellite model B), which assumes the SSM model of microsatellite evolution. The group level refers to (1) impounded and (2) unimpounded populations.

Source of variation	degrees of freedom	Sum of squares	Variance components	Percentage of variation
Among groups	1	17.177	0.05165 Va	1.05
Among populations within groups	7	40.303	0.01871 Vb	0.38
Within populations	437	2128.168	4.86995 Vc	98.58
Total	445	2185.648	4.94031	
Fixation Indices				

FST : 0.01424

FSC : 0.00383

FCT : 0.01045

B)

A)

Source of variation	degrees of freedom	Sum of squares	Variance components	Percentage of variation
Among groups	1	13.270	0.03944 Va	0.98
Among populations within groups	7	31.978	0.01237 Vb	0.31
Within populations	437	1739.923	3.98152 Vc	98.72
Total	445	1785.170	4.03333	

**Fixation Indices** 

FST : 0.01285

FSC : 0.00310

FCT : 0.00978

Table 3.5. Results from tests for genetic bottlenecks (i.e. tests for significant heterozygote excess) performed with BOTTLENECK 1.2.02 (Cornuet and Luikart 1996);
H<sub>o</sub>, observed heterozygosity; H<sub>E</sub>, expected heterozygosity; P, p-value for one-tailed test of heterozygote excess.

Collection site	$H_E$	Ho	Р
A	0.688	0.700	0.596
В	0.729	0.576	
C	0.745	0.704	0.095
D	0.661	0.698	0.271
Е	0.692	0.717	0.213
F	0.698	0.690	0.188
G	0.732	0.691	0.892
Н	0.727	0.699	0.776
I	0.675	0.686	0.122

Table 3.6	. Matrix of linear pairwise distances between populations (sampling areas) based on the central locality (CL) for
each. Un	its are in (river) kilometers.

Site	А	В	С	D	Е	F	G	Н	Ι
A - Missouri River, MO	Х	644	700	795	1183	1955	2681	2599	3021
B - Kansas River, KS		Х	507	602	990	1762	2488	2406	2828
C - Missouri River, NE			Х	95	483	1255	1981	1899	2321
D - Platte River, NE				х	420	1193	1192	1918	2258
E - Niobrara River, SD					х	772	1498	1416	1838
F - Missouri River, ND						Х	726	644	1065
G - Yellowstone River, MT							Х	526	948
H - Missouri River, MT								Х	422
I - Missouri River, MT									х

**Table 3.7.** Matrix of linear pairwise distances from samples on or near the mainstem Mississippi River. Site

 numbers correspond to Table 4.1 and Figure 4.2 in Chapter 4. Units are in river kilometers.

Site	8	11	20	16	17	18	19
8 – Mississippi River, LA	х	827	1456	1844	2029	2227	2493
11 – Hatchie River, TN		х	790	1178	1363	1561	1827
20 – Missouri River, MO			Х	726	911	1109	1374
16 – Mississippi River, IA				х	185	383	649
17 – Wisconsin River, WI					X	278	544
18 – Chippewa River, WI						X	314
19 – Minnesota River, MN							Х

**Figure 3.1.** Sampling area for nine localities (A-I) in the Missouri River basin. Letters correspond to localities listed in Table 3.1.



**Figure 3.2.** GeneMapper screen image of multiplex "A" reaction for individual 17-16. Eight co-amplified loci appear in the top four lines with LIZ500 internal size standard appears on the bottom line. True allele peaks are circled.



**Figure 3.3.** GeneMapper screen image of multiplex "B" reaction for individual 24-. Eight co-amplified loci appear in the top four lines with LIZ500 internal size standard appears on the bottom line. Again, true allele peaks are circled.



**Figure 3.4.** Results of Bayesian inference of population structure in the Missouri River drainage as determined by STRUCTURE 2.0 (Falush *et al.* 2003; Pritchard *et al.* 2000); k = 2 elicited the highest posterior probability for the number of genetic subgroups. These results represent the outcome when k was set at two. Vertical colored bars indicate the fraction of an individual's genome that has ancestry in a given subgroup (in this case, one of two subgroups).



**Figure 3.5.** Results of Isolation by Distance analysis on Missouri River samples. The graph indicates results of a Mantel test for matrix correlation between log(M) and geographic distance: Z = 44589.1996, r = -0.6186, one-sided  $p \le 0.9999$  from 10000 randomizations (for test of negative correlations, **one-sided p**  $\le$  **0.0001**)



**Figure 3.6.** Results of Isolation by Distance analysis on Mississippi River samples. The graph shows the results of a Mantel test for matrix correlation between log(M) and geographic distance: Z = 26403.4071, r = -0.4225, one-sided p <= 0.9140 from 1000 randomizations (for test of negative correlations, one-sided p <= 0.0860)



# Rangewide population structure and intermediate polyphyly in the genus *Cycleptus* (Teleostei: Catostomidae)

## Abstract

What are the evolutionary mechanisms responsible for the diversification and extant distribution of large riverine fishes? The attempt to unite observable patterns with evolutionary processes is not a trivial task. Recent advances in molecular and geological data analysis have provided many answers to such questions. Still, our knowledge is far from complete and, owing to wholesale anthropogenic changes, the window of opportunity for investigation is rapidly closing on some systems. Such is the case for large riverine fishes that occupy the Mississippi Basin and Gulf Coastal drainages in North America. The Mississippi Basin is characterized by long-term stability, and, for that reason, harbors many ancient lineages of freshwater fishes (e.g., Polyodon, Scaphirhynchus, and others). Unfortunately, many have been translocated from one area to another by humans within the past century, thus obscuring the naturally occurring genetic signal. The focal taxon of this research is the freshwater fish genus *Cycleptus*, a highly migratory group of fishes that has not been subject to stocking efforts. These fishes occupy a vast portion of North America and are of prominent conservation concern throughout. Previous studies revealed incomplete lineage sorting in the two described species, Cycleptus elongatus and C. meridionalis, while the Rio Grande population was reciprocally monophyletic and clearly divergent from the others. The aim of this study

was to characterize inter- and intrabasin population structure throughout the range, to test competing hypotheses regarding the divergence of the two described species, *C. elongatus* and *C. meridionalis*, and to determine whether they are, in fact, genetically isolated. A total of 589 specimens were collected from throughout the known range of the genus. Mitochondrial DNA sequence (control region) and nuclear microsatellite genotypic data (15 loci) were acquired from a total of 151 and 589 individuals, respectively. Bayesian analysis of microsatellite genotypic data indicates distinct subpopulations of *C. elongatus* within the Mississippi basin while mitochondrial markers reveal a pattern of intermediate polyphyly with no gene flow between the two described species.

# Introduction

In recent times, increasingly sophisticated methods of inference from molecular genetic data have enabled a better understanding of the evolutionary mechanisms responsible for speciation (Avise 2004; Felsenstein 2004). The attempt to unite observable patterns with evolutionary processes is not a trivial task. New advances in molecular and geological data analysis have provided many answers to such questions. The issue becomes particularly challenging when the taxa in question recently diverged, placing the investigator at the interface between questions of phylogeny and population genetics.

Many recent investigations have illuminated the history of North American freshwater fish taxa; yet, most of these occur east of the Great Plains or west of the Rocky Mountains (Berendzen *et al.* 2003; Near and Keck 2005; Wilson *et al.* 1996). Only a handful of studies involve more wide ranging taxa (Kreiser *et al.* 2001) in the intervening and surrounding landscape, and few have dealt explicitly with troublesome groups that exhibit incomplete lineage sorting. The present study is the first to address these issues in the North American freshwater fish genus *Cycleptus*, a group that is distributed over much of the continent, but whose two described species do not sort at mtDNA loci.

Blue suckers (genus *Cycleptus*) are members of the family Catostomidae (suckers), a group of tetraploid fishes that is broadly distributed throughout North America and includes a single species endemic to Asia, *Myxocyprinus asiaticus* (*Catostomus catostomus* occurs on both continents). Most researchers hypothesize that the family originated in Asia, dispersed across Beringia during the Eocene, and subsequently diversified in North America (Harris and Mayden 2001; Smith 1981; Uyeno and Smith 1972). Although the phylogenetic position of *Cycleptus* within Catostomidae is still debated, most researchers place it as a basal group either sister to *Myxocyprinus* or to the subfamily Catostominae (Ferris and Whitt 1978; Harris and Mayden 2001), which includes all extant suckers except the subfamily Ictiobinae (buffalofishes and carpsuckers). In either case, it is likely that the cycleptid lineage has been in existence since at least the mid-Cenozoic, although supporting fossil evidence has not been found (Cavender 1986; Swift 1968).

Although they occur over such a vast range in North America (Figures 4.1 and 4.2), intrageneric diversity has only been studied on three previous occasions. Following the elevation of *Cycleptus* to genus (Rafinesque 1820), it was considered monotypic as *C. elongatus* for over 175 years until Burr and Mayden (1999) described a sister species, *C. meridionalis*, from a number of disjunct southeastern Gulf Coast drainages based on an array of bimodally-distributed, non-overlapping morphological characters. A subsequent allozyme study based on 23 individuals collected from four disjunct drainages (Buth and Mayden 2001) showed three distinct groups: i) Mississippi Basin, ii) Pascagoula River plus Mobile Basin, and iii) Rio Grande Basin. Most recently, phylogenetic analysis of mtDNA by Bessert and Orti corroborated previous allozyme work and initiated a formal description of the Rio Grande species (see Chapter 1).

Knowledge of *Cycleptus* diversity, both intrageneric and intraspecies, is important for a number of reasons. Awareness of *Cycleptus* population dynamics will enable a better understanding of other large riverine fish taxa in central North America and, in particular, the role of interdrainage exchange in maintaining species boundaries.

Although geological evidence indicates that drainage patterns within the Mississippi basin have evolved significantly in recent (Quaternary) time (Burr and Page 1986; Cross et al. 1986), there has been a relatively stable large riverine presence here since the late Cretaceous / early Cenozoic (Briggs 1986). This is evidenced by the persistence of multiple basal Actinopterygian lineages in the system, including Amiiformes (bowfin), Acipenseriformes (sturgeons and paddlefishes), and Semionotiformes (gars), which, along with *Cycleptus*, are often characterized as the "old river ichthyofauna." None of these lineages exhibit a great deal of extant diversity and most, including Cycleptus, are of imminent conservation concern throughout their ranges (Becker 1983; Warren et al. 2000) (e.g., *Cycleptus*, see Figure 4.1). Unfortunately, multiple translocations have already been performed in a number of these fishes prior to the collection of any genetic data (see Gardner 2004; Rutledge 1989); as such, the natural genetic signals have been obscured. Cycleptus is an important exception. To my knowledge, none of these fishes have been stocked from one location to another in any portion of their range; therefore, examination of extant and historic population structure will not be confounded by recent events and may provide new insights into the history of other problematic groups, such as the Scaphirhynchus sturgeons (Simons et al. 2001).

Assuming a mid-Cenozoic origin and the likelihood that cycleptid predecessors occupied similar large riverine habitats in North America, there are a number of possibilities for vicariance and divergence, especially during Pleistocene time (Briggs 1986; Cross *et al.* 1986). One specific case pertains to the two described species, *C. elongatus* and *C. meridionalis*. The former occurs throughout the Mississippi Basin and extends westward to the disjunct Colorado River of Texas while the latter occurs only in a handful of disjunct drainages east of the Mississipi River, up to and including the Mobile Basin (Boschung and Mayden 2004). These ranges are nonoverlapping and physically disjunct (Figure 4.2). If the genus was already present in the upper Tennessee at the time, one possible vicariant mechanism contributing to the divergence and extant distribution of these fishes is the breakup of the ancient Appalachian River, which hypothetically connected the upper Tennessee River to the Coosa River (part of the Mobile Basin) during the mid-Miocene, approximately 16.4 to 11.2 million years ago (mya) (Mills and Kaye 2001; Near and Keck 2005). More plausible explanations exist for concordance with the latter breakup of connections between the lower Tennessee and other rivers in the Mobile Basin (e.g., Tombigbee), which occurred in the Pliocene and Pleistocene (Mills and Kaye 2001). Finally, there is the very real possibility of more recent dispersal along the coast via freshets and reduced coastal salinity following Pleistocene glacial melting.

Is there any level of recurrent gene flow among these disjunct populations? Quite possibly there is, given the close proximity of the mouths of these drainages. The mouth of the Mississippi River is less than 200 kilometers from Mobile Bay, and the Pearl and Pascagoula drainages lie in-between. The account of a single *C. meridionalis* specimen captured several kilometers from fresh water at Dauphin island, Alabama (Swingle 1971) suggests these fishes have enough salinity tolerance for limited marine dispersal, possibly accompanying significant freshets. This individual was distressed when captured and its appearance at this locality was, in fact, attributed to a freshet (Burr and Mayden 1999)

As previously-mentioned, understanding intraspecies (indeed, intrabasin) population structure is paramount for appropriate conservation efforts of these fishes.
Both species of *Cycleptus* are known to migrate considerable distances to spawn (Elstad and Werdon 1993; Hand and Jackson 2003b; Peterson *et al.* 1999), but little is known of site fidelity and other aspects of their natural history. If site fidelity is low and dispersal (i.e. gene flow) high, one would expect to see a lack of distinct population structure in a given basin, with the possible exception of groups isolated by impoundments or other barriers (see Chapter 3).

The aims of this study were, therefore, threefold: i) to estimate intraspecies population structure, particularly that of *C. elongatus* in the Mississippi Basin; ii) to test competing hypothesis regarding the *C. elongatus / C. meridionalis* split; and iii) to test the hypothesis of recurring gene flow between the two species. In conducting these analyses, we gain a better understanding of the evolution of the genus - and of large riverine fishes in general. Finally, and equally important, the population genetic information gathered here (and in Chapter 3) will facilitate informed decisions regarding conservation of these fishes.

## **Materials and Methods**

#### Study location and sample collection

Previous microsatellite genotypic and mtDNA sequence data from 231 individuals captured in the Missouri River drainage (Chapter 3: Table 3.2, Figure 3.1) was incorporated into this project. A coordinated effort by more than 150 academic, governmental, and private researchers led to the addition of another 360 individuals from throughout the remaining range, (Table 4.1, Figure 4.2). Fishes were captured with hoop nets, gill nets, and electroshocking devices. A small (1 cm<sup>2</sup> or less) fin clip was removed from each fish and preserved in 95% EtOH for shipment to the University of Nebraska. Tissues are vouchered in the personal collection of the auther (see Appendix 2) and are available upon written request.

# DNA preparation and amplification

DNA was extracted from tissue samples using either a standard phenolchloroform protocol (Sambrook *et al.* 1989) or DNeasy® Tissue purification kit (Qiagen). DNAs were eluted in either water or EB buffer (Qiagen). A small number of samples were randomly selected and (1-2  $\mu$ l) electrophoresed through a 1% agarose gel to check quality. All samples were quantified with a GeneQuant II spectrophotometer (Pharmacia Biotech) and a portion diluted to a standard working stock concentration of 100ng/ $\mu$ l. The remainder of each elution was placed in -70° C for long-term storage. The mitochondrial control region and 15 polymorphic microsatellite markers were amplified, purified, and scored according to the procedures outlined in chapter one (p. 17-19) and chapter three (p. 59-61).

## Sample preparation and data collection

For microsatellite loci, genotypes were scored with GeneMapper 3.7 (Applied Biosystems) and the data was transformed to GenePop format (Raymond and Rousset 1995) with GMCONVERT 0.32 (Faircloth 2006). Genotypic data were checked for syntax errors, evidence of null alleles (Hardy-Weinberg equilibrium), and adherence to the Stepwise Mutation Model (Ohta and Kimura 1973) with MicroChecker 2.2.3 (VanOosterhout *et al.* 2004) and MSAnalyzer 4.0 (Dieringer and Schlötterer 2002). Additional transformations of the data set for analysis with STRUCTURE 2.0 (Pritchard *et al.* 2000) and Arlequin 3.1 (Schneider *et al.* 2000) were performed with CONVERT 1.3.1 (Glaubitz 2004).

For mtDNA control region data, raw sequences were edited with Sequencher 4.2 (Gene Codes) or Contig Express (Invitrogen). Alignment was performed with ClustalX (Thompson *et al.* 1997) and checked by eye for obvious misalignments. For analysis with NETWORK 4.2 (Bandelt *et al.* 1999) and IM (Hey and Nielsen 2004), the alignment was formatted by hand.

#### Data Analysis – Population structure

In addition to observed and expected heterozygosities for each locality, deviations from Hardy-Weinberg equilibrium and pairwise  $F_{ST}$  estimates (Weir and Cockerham 1984) were calculated for each locality using GenePop3.3 (Raymond and Rousset 1995). The program FSTAT 2.9.3.2 (Goudet 1995; Goudet 2002) was also used to calculate the average allelic richness (A) at each locality with 10 or more samples. The distibribution of genetic variation within and among (i) the three species and (ii) within and among basins and disjunct coastal drainages was assayed with a hierarchical analysis of molecular variance (AMOVA – see Table 4.2) (Excoffier *et al.* 1992) performed in Arlequin 3.1 (Excoffier *et al.* 1992; Schneider *et al.* 2000). This test was performed under the standard data model ("different alleles are considered mutationally equidistant from each other").

A second approach that does not require assignment of invidividuals to groups *a priori* (as in AMOVA) is the Bayesian clustering technique implemented in the software STRUCTURE 2.0 (Pritchard *et al.* 2000). Here, the data are used to define the number of subpopulations (*k*). The method assumes HW and linkage equilibrium within

subpopulations and seeks combinations of individuals which maximize these values. To use the method, one performs several runs at *k* values from 1 to n. Assuming quality control in preparation and performance of these runs (e.g., unambiguous data, adequate burn-in, etc.), the value of *k* eliciting the highest mean posterior probability over several runs indicates the true number of genetic subpopulations present. Four simulations were conducted at each value of *k* from one to twelve. A burn-in period of  $2.5 \times 10^5$ generations was followed by  $1.0 \times 10^6$  generations of data collection.

Phylogenetic structure was previously estimated in the genus with mitochondrial markers (Chapter 1); yet, I wished to visualize the distribution of haplotypes among the two species with an unrooted haplotype network. This was calculated with the software NETWORK 4.2 (see Bandelt *et al.* 1999; www.fluxusengineering.com) using the MJ algorithm (Figure 4.4). The advantage in doing this is that it allows the investigator to quickly assess the degree of allele sharing across populations (again, across species in this case) and provides an alternate view of phylogenetic structure. Note that specimens from the Rio Grande were excluded from this analysis due to the previously-determined level of divergence (Chapter 1). In essence, they are not pertinent to the question at hand and would have simply appeared as a monophyletic clade separated from the others by a very long branch.

#### Demography

Population divergence times, historical gene flow, and (female) effective population sizes were estimated using a model of Isolation with Migration as implemented in the software IM (Hey and Nielsen 2004). Briefly, the program applies the IM model (Figure 4.5) to genetic data from a pair of closely related populations (or species, in this case). It uses a Markov Chain Monte Carlo (MCMC) simulation process to estimate posterior probability densities for each of the parameters included in the model. In this case, the basic 6-parameter model was chosen. This includes q1, q2, and qA, referring to the effective size of population 1, population 2, and the ancestral population, respectively; m1, the migration rate of genes from population 1 to population 2 in the coalescent (or, as time moves forward, from population 2 to population 1) and m2, migration rate in the opposite direction; and *t*, the time since divergence of the populations. A subroutine also allows the user to obtain an estimate of the time to most recent common ancestor (TMRCA) for all alleles and/or genotypes included in the input file.

The model parameters are scaled to the 'per year' mutation rate ( $\mu$ ) for the entire locus; thus, they can be converted to demographic parameter estimates if the user defines the mutation rate (that is, for each locus considered – in this case, just the mtDNA control region) and provides a generation time for the organism in the input file. The generation time (in the IM software, 'u') for both *C. elongatus* and *C. meridionalis* is approximately ten years (Becker 1983; Boschung and Mayden 2004).

A recent phylogenetic study of Otocephalan fishes based on mitogenomic DNA placed an approximate divergence time of 101 mya for *Myxocyprinus asiaticus* and *Carpiodes carpio* (Peng *et al.* 2006), both of which have been hypothesized as sister (or part of a sister clade) to *Cycleptus* (Harris and Mayden 2001; Smith 1992). A comparison of *M. asiaticus* and *C. carpio* mtDNA sequences from GenBank (Accession numbers; *Carpiodes carpio*: AY366087, NC\_005257, AB126083, AF454867; *Myxocyprinus asiaticus*: AF036176, AY526869, NC\_006401, AB223007, AY986503)

yields a cytochrome *b* divergence of 17%, which equates to a substitution rate of approximately 0.17 % / MY. Considering the total length of cyt *b*, 1140 base pairs, this amounts to  $9.59 \times 10^{-7}$  mutations per year for the locus (' $\mu$ ' in IM). Since the mean control region sequence divergence between *C. elongatus* and *C. meridionalis* is approximately 2.75 times greater than that for cyt *b* (see Chapter 1, Table 1.1), a control region rate of 2.64 x 10<sup>-6</sup> mutations per year (or 2.64 x 10<sup>-5</sup> mutations per generation given an estimated 10 year generation for *Cycleptus* species) was estimated (e.g., Omland *et al.* 2006).

In order to obtain reliable results from this analysis, it is necessary to perform three separate runs (simulations). The first simulation allows the user to determine appropriate burn-in lengths, uninformative prior distributions, and a number of other settings that can be advantageous when collecting real data. An initial run was performed with unrealistic prior settings as follows: t = 300; q1, q2, and qA all = 500, and m1 and m2 = 100. All of these priors were adjusted substantially for the final two runs, which are identical to one another except that each starts from a different 'seed' point. Priors for the final two runs were set to: t = 10; q1 = 100, q2 = 300, qA = 300, and m1 and m2 both = 5. A burn-in of 1.0 x 10<sup>6</sup> generations was used for the final two simulations and was followed by another 3.0 x 10<sup>6</sup> generations of data collection to ensure that the ESS (effective sample size) for each parameter was 50 or higher (Pritchard and Wen 2004). Seed numbers used for the two runs were 416 and 3119, respectively. As anticipated, these two runs converged to similar estimates for all six model parameters. These were converted to more meaningful demographic parameter estimates (see the IM documentation) and plots of posterior distributions were prepared with Microsoft Excel (Figure 4.6, A-F).

### Results

# Summary Statistics and Population structure

The number of alleles (*na*) detected per sample ranged from a low of 40 in nine samples from the Sabine River to a high of 134 among 38 samples from the Wabash River (Table 4.1). Within the Mississippi Basin, gene diversity was relatively high: 0.66 or higher when the sample size was greater than 10 (Table 4.1). In contrast, disjunct drainages, including the Rio Grande, Colorado River, Sabine River, and Alabama River (site numbers 1,4,5, and 6, respectively, in Figure 4.2), showed reduced levels. The measure of allelic richness (A), in which all values are scaled to the smallest sample size, was also reduced in those drainages. Finally, the upper Missouri River showed significantly less allelic richness than the rest of the Mississippi Basin (Table 4.1; also see Chapter 3, p. 65-66).

The first AMOVA, which assessed the distribution of molecular variance within and among the three cycleptid species, indicated that roughly 12% of the variance occurred among species with 4% among populations (sample sites) within species and 84% within populations (Table 4.2A). These numbers changed slightly when the samples were grouped by disjunct drainage (Alabama, Sabine, Colorado, Rio Grande, and Mississippi Basin). When this was done, more than 15% of the variance occurred between drainages while that between populations within drainages dropped to approximately 1% and that within populations remained stable at approximately 84% (Table 4.2B). The second measure of population structure, Bayesian analysis with

STRUCTURE 2.0 (Pritchard *et al.* 2000), was run four times at each value of *k* from one to twelve. The setting of k = 9 best explained the data as it yielded the highest mean posterior probability over the four simulations. The method clearly recovered the disjunct Rio Grande, Colorado, Sabine, and Mobile basins as distinct subpopulations with very little admixture in the genomes of any individuals from those drainages (Figure 4.3). The other five clusters fall within the Mississippi Basin, as follows: 1) one large Tennessee / Ohio / upper Mississippi cluster that includes the lowest sample site from the Missouri River (site number 20 in Figure 4.2); 2) two intergraded clusters in the Missouri River, in a pattern nearly identical to that indicated by exclusive analysis of this drainage (Chapter 3); and 3) two distinct clusters in the lower Mississippi. Note that cluster 6 in Figure 4.3 is actually a sample of 33 individuals from the Red River in the southeast corner of Oklahoma.

The minimum spanning haplotype network (Figure 4.4) reflects the polyphyly of *C. elongatus* and *C. meridionalis*: three distinct, monophyletic *C. meridionalis* clades are nested within the *C. elongatus* network at different points. Note that no haplotypes are shared between the two species (i.e. no multicolored pie diagrams appear in the figure). *Demography* 

The second and third IM simulations converged to similar parameter estimates and the longer of the two runs is presented here. Posterior distribution peaks for each parameter in the IM model yielded were: q1 = 22.2, q2 = 117.5, qA = 4.65, t = 4.65, m1 = 0.005, and m2 = 0.005. The 95% L/H (low and high point; confidence interval) for each were q1 {13.64, 46.52}, q2 {79.95, 184.65}, qA {1.35, 79.05}, t {3.13, 7.21}, m1 {0.005, 0.245}, and m2 {0.005, 0.085}. Conversions to demographic parameter estimates are plotted in Figure 4.5. In summary,  $\theta_{Cm}$ , the effective number of female *C*. *meridionalis*, is approximately 210,000;  $\theta_{Ce}$  (female *C*. *elongatus*) = 1,109,000; and  $\theta_{Anc}$  (female ancestral) = 44,000. The time of population divergence (*t*), is estimated at 1.76 mya, and occurred long after the time to most recent common ancestor (TMRCA = 2.43 mya) for all haplotypes in the analysis. Finally, migration rates (rate at which genes come into a population) in either direction converted to negligible rates of  $1.32 \times 10^{-8}$  per year or  $1.32 \times 10^{-7}$  per generation.

# Discussion

What mechanisms underlie the extant distribution of any given organism and why is it important to know this? As anthropenic change envelopes the global landscape, it is important that we seize opportunities to understand natural population dynamics while windows for study remain open. The results of these kinds of investigations not only allow us to make predictions about sustainability of focal taxa, but may also provide us with interesting glimpses into the past. Outcomes may ultimately provide for betterinformed mitigation efforts (if necessary), or at worst, satisfy a natural curiousity regarding the evolution of the groups in question.

Here, rangewide population structure was investigated in the fish genus *Cycleptus*, a group of fishes that inhabits a vast, yet diminishing range in North America. Given the rangewide conservation status of these fishes, it is important to understand population structure, and, since the only genetic investigations of these fishes appear in earlier chapters here, the opportunity was taken to include all three species in the same analysis. Since the natural genetic signal in cycleptids has not been obscured by

translocations (i.e. 'stocking,') as it has in other 'old river' ichthyofauna, this study may also lend comparative insights regarding the evolution of those groups.

Analyses of molecular variance indicated that there is distinct structure within the genus, and that much of the structure is associated with geography. If the three defined species (two previously defined plus the third in Chapter 1) satisfy the biological species concept of reproductive isolation, we would expect at least a moderate portion of the molecular variance would to occur between them. This proved to be the case as 12% of the variance was attributed to among-species effects. Interestingly, though, geographic isolation increased this effect. When populations were grouped according to disjunct locality rather than current taxonomy, 15% of the variance occurred among them (Table 4.2). This suggests further study of geographically isolated populations and possible classification as distinct management units (MU) following the criteria outlined by Moritz (1994).

Bayesian analysis with STRUCTURE 2.0 (Pritchard *et al.* 2000) corroborated the AMOVA results as all populations inhabiting disjunct drainages were recovered as distinct and separate (see Figure 4.3, clusters 1, 7, 8, and 9) with very little admixture. Within the Mississippi Basin, an additional five distinct clusters were recovered, but with a generally higher degree of admixture (= higher gene flow) as reflected in the individual (vertical) bars in Figure 4.3. The notable exception, as outlined in Chapter 3, is the pattern occurring in the Missouri River, where almost certainly there is an absence of upstream gene flow, but only in very recent time. In terms of ESU or MU designation for subdivisions in the Mississippi basin, these results do not refute a more detailed null hypothesis of exchangeability (Crandall *et al.* 2000; Templeton 1989); however, it is

unclear whether any of the Mississippi Basin subdivisions outlined here possess adaptations to local environmental conditions, as has been seen in other fishes (Cooke and Kassler 2001). If this is the case, the clusters may be rendered ecologically inexchangeable and would further support their designation as distinct units (Crandall *et al.* 2000). Variation in the length of lip papillae among cycleptids in the Mississippi Basin (Burr and Mayden 1999) lends credence to this possibility. The results of this analysis provide a logical starting point for further investigations in this direction.

From a conservation standpoint, a clear understanding of extant population structure is critical; yet, it is equally important to understand the evolutionary history of the taxa in question. For instance, how and when did the *C. elongatus / meridionalis* split come about? More precisely, when did a common ancestor come to reside in the coastal drainages now occupied by *C. meridionalis*, and when did the two lineages begin to diverge? Finally, does the timing of the split coincide with any geological events that could facilitate vicariance?

One approach that has gained recent popularity, particularly in its applicability to taxa that exhibit incomplete lineage sorting, is the Bayesian technique implemented in the software IM (Pritchard *et al.* 2000). This method allows the estimation of parameters in a model of isolation with migration between two closely related populations or species (Figure 4.5). Two long simulations ( $4.0 \times 10^6$  generations) converged to similar estimates for all model parameters and these were converted to more meaningful demographic parameter estimates (Figure 4.6A-F).

The posterior distribution peak for population divergence time occurred at 1.76 mya (95% L/H = 1.19-2.73), post-dating TMRCA (2.43) by an estimated 670,000 years

107

(Figure 4.6D). This suggests that: i) much of the extant genetic diversity among the two species was already present prior to divergence, and ii) divergence occurred in the early to mid-Pleistocene, or, at the earliest, the late Miocene. This contradicts the hypothesis of the mid-Miocene breakup (16.4-11.2 mya) of the ancient Appalachian River (Mills and Kaye 2001; Near and Keck 2005), which hypothetically connected the upper Tennessee River with the Coosa River of the Mobile Basin, as a mechanism for vicariance. However, the estimated time of divergence is consistent with the Pleistocene breakup of other hypothetical connections, such as that between the Hatchie River (current tributary of the lower Mississippi River) and the Pearl and/or Pascagoula drainages in present-day northern Mississippi. Given the reduced effective size of the common ancestral population relative to extant populations of both species (Figure 4.6A-C), it is also reasonable to hypothesize expansion from a reduced Pleistocene refugium in the lower Mississippi. Unfortunately, it is difficult to ascertain the potential impact of marine dispersal in forming the extant distribution of these fishes; however, a more intense effort focused only on those drainages harboring C. meridionalis populations would yield valuable information. Future efforts are planned in this direction.

Another salient features of the IM analysis is the estimated rate of gene flow in each direction (m1 and m2) of 0.005 (Figure 4.6E-F). This equates to  $1.32 \times 10^{-8}$ migrants per generation, which effectively means long-term reproductive isolation since the 'populations' diverged. This, in combination with the pattern of monophyletic subclades in the haplotype network (Figure 4.4), indicates that the two species are in the latter stages of intermediate polyphyly, or, in the recently proposed terminology by Omland *et al.* (2006), they would be classified as allophyletic. This combination of allophyly, reproductive isolation, and morphological differences lends strong support to the species designation of *C. meridionalis* by Burr and Mayden (1999).

Finally, the estimates of effective population size (Figure 4.6A-B) speak to the current conservation status of both species (Figure 4.1). Cycleptid fishes and other round-bodied suckers are considered bioindicators of stream quality (EPA 1989), and their susceptibility to pollutants is a probable cause for (apparent) decline in many portions of the range. IM analysis indicates robust (female) effective population sizes in both species ( $\theta_{Cm} = 210,000$ ;  $\theta_{Ce} = 1,109,000$ ); thus, implying no imminent concern for the persistence of either. While these results are encouraging for the immediate future, caution is suggested in interpreting them as a panacea. Continued monitoring is encouraged throughout, especially given the intrabasin structure implied here – and the possibility of locally adapted populations.

In conclusion, this study sheds important light on evolutionary history, population structure, and demography in the genus *Cycleptus*. Other 'old ichthyofaunal taxa' (Robison 1986) also show ambiguous relationships at mitochondrial markers, yet, are clearly distinguishable morphologically (Simons *et al.* 2001). Future comparative studies using multiple approaches among these taxa may lead to a better understanding of the process of speciation and maintenance of species boundaries in these groups. Less directly, such studies will also serve to unravel the mysteries of drainage evolution in the major basins and coastal drainages of central North America.

# Acknowledgements

This study was generously supported by a variety of funding sources, including the American Museum of Natural History (Theodore Roosevelt Fund), the US Fish and Wildlife Service (through a State Wildlife Grant from the Nebraska Game and Parks Commission), Sigma Xi Scientific Society, the Center for Great Plains Studies, the UNL School of Biological Sciences, and the Initiative for Ecological and Evolutionary Analysis at the University of Nebraska. This work would not have been possible, particularly in the given time frame, without the myriad efforts of cooperating individuals and agencies throughout the U.S. and Mexico (see Appendix 2).

Gene Site **River/drainage** State/Country diversity  $H_E$ А n na number  $(H_0)$ 1 Alabama AL 30 85 0.592 0.661 5.532 2/3Pearl / Pascagoula MS 6 51 0.536 0.635 -4 Sabine TX / LA 9 40 0.389 0.469 5 ΤX Colorado 30 72 0.519 0.520 4.521 6 Pecos / Black NM 5 52 0.715 0.833 -Rio Grande / 7 5.916 TX / Mexico 21 86 0.627 0.706 Conchos 8 Lower Mississippi LA 32 127 0.679 0.700 7.510 9 Red AR 5 48 0.680 0.827 -10 OK 33 Red 113 0.669 0.681 6.718 TN 19 11 Hatchie 101 0.751 0.759 7.088 12 3 TN / Cumberland TN 36 0.950 0.917 -13 TN 9 78 0.758 0.779 Upper Tennessee -14 KY / OH 30 124 0.697 0.720 7.468 Ohio 15 Wabash IN 38 134 0.700 0.706 7.638 16 IA Middle Mississippi 11 92 0.727 0.767 7.667 17 WI 32 123 0.713 0.746 7.226 Wisconsin Chippewa / Red 7.302 18 WI 30 121 0.730 0.765 Cedar 19 MN 101 0.751 0.754 7.149 MN / Upper Miss. 18 20 30 121 Missouri MO 0.678 0.696 7.340 21 KS 51 0.576 0.729 Kansas 6 -22 NE 125 7.290 Missouri 38 0.755 0.764 7.311 23 NE 23 111 0.693 Platte 0.663 24 NE / SD 29 121 7.271 Missouri / Niobrara 0.681 0.705 25 14 ND 83 0.688 0.738 6.343 Missouri 26 Missouri ND/MT 30 112 0.733 0.748 6.629 27 Yellowstone MT 31 117 0.733 0.745 6.902 28 MT 29 104 0.683 0.678 6.591 Missouri Totals 591 202 0.68 0.72 6.870

**Table 4.1.** Sampling localities and numbers for microsatellite genotypic analyses. Site numbers correspond to those in Figure 4.2; n, number of individuals; na, number of alleles;  $H_0$ , observed heterozygosity;  $H_E$ , expected heterozygosity; A, allelic richness.

**Table 4.2.** AMOVA summary under the standard model when (A) populations are grouped by present species designation (3 groups); and (B) by disjunct location (5 groups).

A )					
A)	Source of variation	d.f.	Sum of squares	Variance components	Percentage of variation
	Among species	2	144.613	0.55988 Va	11.95
	Among populations within species	24	276.431	0.17446 Vb	3.72
	Within populations	1153	4557.026	3.95232 Vc	84.33
	Total	1179	4978.070	4.68666	
	Fixation Indie FST : 0 FSC : 0 FCT : 0	ces ).15669 ).04228 ).11946			
B)	Source of variation	d.f.	Sum of squares	Variance components	Percentage of variation
	Among disjunct areas	s 4	277.779	0.72133 Va	15.24
	Among populations within disjunct areas	s 22	143.266	0.05852 Vb	1.24
	Within populations	1153	4557.026	3.95232 Vc	83.52
	Total	1179	4978.070	4.73218	
	Fixation India FST : 0 FSC : 0 FCT : 0	ces ).16480 ).01459 ).15243			

**Figure 4.1.** State conservation status for *Cycleptus elongatus* as determined by the Nature Conservancy. For rank definitions, see Master (1997) and Stein (2002).

# State/Province Conservation Status Rank



**Figure 4.2.** Sampling locality map. Circles indicate sampling localities. Locality numbers correspond to those appearing in Table 4.1. Shaded areas represent the range of the three known *Cycleptus* species: red = *C. meridionalis*, blue = *C. elongatus*, and green = the undescribed species from the Rio Grande basin. Map, courtesy of NatureServe (2005), may be accessed at: <u>http://www.natureserve.org/explorer/servlet/NatureServe</u>.



**Figure 4.3.** Bayesian inference of population structure as determined with STRUCTURE 2.0 (Falush *et al.* 2003; Pritchard *et al.* 2000). The graph represents a range-wide sample of 596 individuals in the genus *Cycleptus*. Four runs were performed with each value of *k* (number of genetic subgroups) from 1 to 12; *k*=9 provided the highest mean posterior probability. The results shown are from the single *k*=9 run with the highest posterior probability. Vertical colored bars indicate the fraction of an individual's genome that has ancestry in a given subgroup. Hydrologic units appear below the x-axis while actual genetic subgroups (*k* = 1-9) appear above. Note the graded structure in the Missouri River drainage as indicated in Chapter 3; the *k1* group (Mobile basin) is *C. meridionalis*; *k7* is the undescribed species from the Rio Grande; all others are *C. elongatus*.



Full range STRUCTURE results: k = 9 (*n*=589)

**Figure 4.4.** Unrooted haplotype network computed with NETWORK 4.2 (Bandelt *et al.* 1999) based on 125 mtDNA control region sequences from throughout the ranges of *C. elongatus* and *C. meridionalis*. Dark shading represents *C. meridionalis*; non-shaded circles represent *C. elongatus*. The area of each circle corresponds to the number of individuals possessing that haplotype. The text in each circle indicates the sampling locality (state or country) and number of individuals (if >1).



**Figure 4.5.** The Isolation with Migration model described by Nielsen and Wakeley (2001) and implemented in the program IM (Hey and Nielsen 2004).  $\theta_A$ ,  $\theta_1$ , and  $\theta_2$  indicate effective population sizes of an ancestral population and two descendent populations, respectively; *m1* and *m2* indicate migration rates in the coalescent; *t* is the estimated time of divergence between the descendent populations.



**Figure 4.6.** Bayesian inference of population parameters for *C. elongatus* and *C. meridionalis* using the coalescent approach implemented in the program IM (Hey and Nielsen 2004). Each graph represents the posterior distribution for the parameter indicated, scaled to the estimated 'per year' mutation rate ( $\mu$ ) of 2.64 x 10<sup>-6</sup> for the control region (see text). (A)  $\theta_{Cm}$ , (B)  $\theta_{Ce}$ , and (C)  $\theta_A$  are the effective number of females for *C. meridionalis*, *C. elongatus*, and the common ancestral population, respectively; (D) *t* is the time since (population) divergence, TMRCA is the time to the most recent common ancestor of all the haplotypes included in the analysis; (E)  $m_I$  is the migration rate from *C. meridionalis* to *C. elongatus*; (F) *m2* is the migration rate from *C. elongatus* to *C. meridionalis*.









# CONCLUSIONS

The overarching theme of this research effort was to investigate hierarchical levels of relatedness in natural populations of the cycleptid fishes (blue suckers), a widespread genus in North America. Formerly, phylogenetic relationships were unknown in the genus. This research has revealed additional biodiversity in the genus (new species, yet to be described, from the Rio Grande Basin) and called into question the existing taxonomy because of incomplete lineage sorting between the two described species at mitochondrial loci.

Other branches of this effort included an examination of rangewide population structure and a more focused assessment of the impacts of dams on population structure in the upper Missouri River. In order to conduct these analyses, it was necessary to develop an array of taxon specific microsatellite markers. In so doing, a useful technique for isolating paralogous loci in a tetraploid genome was discovered. Population screens with isolated paralogs revealed a pattern of chromosomal inheritance that is consistent with an allopolyploid origin.

Highly efficient multiplexed reactions were used to conduct rangewide microsatellite genotypic data. Results of subsequent population structural analyses indicate distinct structure within the Mississippi basin and a distinct pattern of isolation by distance in the Missouri River drainage, a pattern that is not present in the Mississippi River. This is among the first studies to reveal altered genetic patterns resulting from anthropogenic modifications to large river systems. Finally, Bayesian estimates of historical demography and divergence time between the two described species, *C*. *elongatus* and *C. meridionalis*, revealed a pattern of longterm genetic isolation. This lends some credence to the current taxonomy.

In carrying out this work, approximately 600 tissue specimens were collected from throughout the range over a one-year time period. This frozen 'point in time' will be maintained by the author and is freely available for use by others upon request. The intent is that this collection will provide a valuable baseline for future repeated measures in tracking the evolution of, and providing appropriate conservation measures for, fishes in the genus *Cycleptus*.

## REFERENCES

- Adams S, Flinn M, Burr B, Whiles M, Garvey J (2006) Ecology of larval blue sucker (*Cycleptus elongatus*) in the Mississippi River. *Ecology of Freshwater Fish* 15: 291-300.
- Agassiz L (1855) Article XII. Synopsis of the ichtyological fauna of the Pacific Slope of North America, chiefly from the collections made by the U.S. Expl. Exped. under the command of Captain C. Wilkes, with recent additions and comparisons with eastern types. *American Journal of Sciences and Arts*: 71-99, 215-231.
- Avise JC (2004) *Molecular Markers, Natural History, and Evolution* Sinauer Associates, Inc., Sunderland, MA.
- Baker SC, Armstrong ML (1987) Recent collections of fishes from the Spring River
  drainage in northeast Arkansas. *Proceedings of the Arkansas Academy of Science*41: 96.
- Bandelt H, Forster P, Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16: 37-48.
- Barko VA, Herzog DP, Hrabik RA (2004) Relationship among Fish Assemblages and
   Main-Channel-Border Physical Habitats in the Unimpounded Upper Mississippi
   River. *Transactions of the American Fisheries Society* 133: 371-384.
- Barnickol PG, Starrett WC (1951) Commercial and sport fishes of the Mississippi River between Caruthersville, Missouri and Dubuque, IA. *Illinois Natural History Survey Bulletin* 25: 267-350.

Becker GC (1983) Fishes of Wisconsin University of Wisconsin Press, Madison.

- Benke AC, Cushing CE (2005) Rivers of North America. Elsevier Academic Press, San Diego, CA.
- Berendzen PD, Simons AM, Wood RM (2003) Phylogeography of the northern
  hogsucker, *Hypentelium nigricans* (Teleostei: Cypriniformes): genetic evidence
  for the existence of the ancient Teays River. *Journal of Biogeography* 30: 1139-1152.
- Berry CR, Jr., Duffy WG, Walsh R, et al. (1993) The James River of the Dakotas. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR). National Biological Survey, Biological Report 19.
- Bessert ML, Sitzman C, Ortí G (2006) Avoiding paralogy: diploid loci for allotetraploid blue sucker fish (Cycleptus elongatus, Catostomidae). *Conservation Genetics*online first: 1-4.
- Boschung H, Mayden R (2004) *Fishes of Alabama* The Smithsonian Institution, Washington, DC.
- Branson BA (1962a) Comparative cephalic and appendicular osteology of the fish family Catostomidae. Part I, *Cycleptus elongatus* (Lesueur). *Southwestern Naturalist* 7: 81-153.
- Branson BA (1962b) Observations on the distribution of nuptial tubercles in some catostomid fishes. *Transactions of the Kansas Academy of Science* **64**: 360-372.
- Branson BA, Harker DF, Baskin JM, et al. (1981) Endangered, threatened, and rare animals of Kentucky. *Transactions of the Kentucky Academy of Science* 42: 77-89.

- Breder CM, Rosen DE (1966) *Modes of reproduction in fishes* Natural History Press, Garden City, New York.
- Briggs J (1986) Introduction to the Zoogeography of North American Fishes. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt C, Wiley E), pp. 1-16. Wiley-Interscience, New York, NY.

 Broughton R, Renau P, Richter S (2006) Spatial covariation of mutation and nonsynonymous substitution rates in vertebrate mitochondrial genomes.
 *Molecular Biology and Evolution* 23: 1516-1524.

- Brown CJD (1971) *Fishes of Montana* Montana State University, Big Sky Books, Bozeman, MT.
- Brown DJ, Coon TG (1994) Abunance and assemblage structure of fish larvae in the lower Missouri River and its tributaries. *Transactions of the American Fisheries Society* **123**: 718-732.
- Brownstein M, Carpten J, Smith J (1996) Modulation of nontemplated nucleotide addition by Taq DNA Polymerase: Primer modifications that facilitate genotyping. *Biotechniques* **20**: 1004–1010.
- Buchanan T (1973) *Key to the fishes of Arkansas* Arkansas Game and Fish Commission, Little Rock, Arkansas.

Burr B, Page L (1986) Zoogeography of Fishes of the Lower Ohio-Upper Mississippi
Basin. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt
C, Wiley E), pp. 287-324. Wiley-Interscience, New York, NY.

- Burr BM, Cook KM, Eisenhour DJ, et al. (1996) Selected Illinois fishes in jeapardy: new records and status evaluations. *Transactions of the Illinois State Science Academy* 89: 169-186.
- Burr BM, Mayden RL (1999) A new species of *Cycleptus* (Cypriniformes: Catostomidae) from Gulf slope drainages of Alabama, Mississippi, and Louisiana, with a review of the distribution, biology, and conservation status of the genus. *Bulletin of the Alabama Museum of Natural History* **20**: 19-57.
- Burr BM, Page LM (1993) Fishes of the Big Muddy River drainage with emphasis on historical changes. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 186-209. National Biological Survey, Biological Report 19.
- Bussjaeger C, Briggs T (1978) Phylogenetic implications of bile salts in some catostomid fishes. *Copeia* **1978**: 533-535.
- Buth DG (1983) Duplicate isozyme loci in fishes: origins, distribution, phyletic consequences, and locus nomenclature. In: *Isozymes: current topics in biological and medical research. Genetics and Evolution*, pp. 381-400. Alan R. Liss, Inc., New York, New York.
- Buth DG, Mayden RL (1998) Allozymic and isozymic variation in Cycleptus elongatus (catostomidae). *Gene Families and Isozymes Bulletin* **131**: 34.
- Buth DG, Mayden RL (2001) Allozymic and isozymic evidence for polytypy in the North American catostomid genus *Cycleptus*. *Copeia* **2001**: 899-906.
- Butler RS (2002) Imperiled Fishes of the Southern Appalachian Ecosystem, with Emphasis on the Non-Federally Listed Fauna Report to the Southern Appalachian Ecosystem Team U.S. Fish and Wildlife Service.
- Call RE (1896) Fishes and shells of the Falls of the Ohio. *Memorial History of Louisville* 1: 9-20.
- Carlander KD (1969) Handbook of freshwater fishery biology Iowa State University Press, Ames, Iowa.
- Carreiro JM (2000) Life history characteristics of the blue sucker (Cycleptus elongatus) in the unchannelized reach of the Missouri River below Gavins Point Dam, South Dakota State University.
- Cavender T (1986) Review of the Fossil History of North American Freshwater Fishes.
  In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt C, Wiley E), pp. 699-724. Wiley-Interscience, New York, NY.
- Christenson LM (1974) Notes on the blue sucker, Cycleptus elongatus (Lesueur), in the lower Chippewa and Red Cedar Rivers, Wisconsin. In: *Research Report 75*, p. 7.
  Wisconsin Department of Natural Resources, Madison, Wisconsin.
- Clay WM (1975) *The fishes of Kentucky* Kentucky Department of Fish and Wildlife Resources, Frankfort, Kentucky.
- Coker RE (1930) Studies of common fishes of the Mississippi River at Keokuk. *Bulletin* of the U.S. Bureau of Fisheries **45**: 141-225.
- Colborn J, Crabtree RE, Shaklee JB, Pfeiler E, Bowen BW (2001) The evolutionary enigma of bonefishes (Albula spp.): cryptic species and ancient separations in a globally distributed shorefish. *Evolution* **55**: 807-820.

- Conner J, Suttkus R (1986) Zoogeography of Freshwater Fishes of the Western Gulf
  Slope. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt
  C, Wiley E), pp. 413-456. Wiley-Interscience, New York, NY.
- Conner JV (1977) Zoogeography of freshwater fishes in western Gulf Slope drainages between the Mississippi and the Rio Grande dissertation, Tulane University.

Contreras-Balderas S, Lozano-Villano MdL, Garcia-Ramirez ME (1995) Third annotated and revised checklist of the fishes from Nuevo Leon, Mexico. In: *Consejo Consultivo Estatal para la Preservacion y Fomento de la Flora y Fauna Silvestre de Neuvo Leon*. (eds. Contreras-Balderas S, Gonzalez F, Lazcano V, Contreras A), pp. 71-78. Gobierno del Estado de Nuevo Leon, Mexico.

- Contreras-Balderas S, Rivera R (1972) Una localidad neuva para *Cycleptus elongatus* (LeSueur) en el Río Bravo, México y Estados Unidos (Pisces: Catostomidae). *Revista de la Sodiedad Mexicana de Historia Natural* **23**: 47-50.
- Cook FA (1959) Freshwater fishes in Mississippi Mississippi Game and Fish Commission, Jackson, Mississippi.
- Cooke SJ, Kassler TWP, D.P. (2001) Physiological performance of largemouth bass related to local adaptation and interstock hybridization: implications for conservation and management. *Journal of Fish Biology* **59**: 248-268.
- Cooper EL (1983) Fishes of Pennsylvania and the northeastern United States Pennsylvania State University Press, University Park, Pennsylvania.
- Cornuet JM, Luikart G (1996) Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* **144**: 2001-2014.

- Cowley DE (2006) Strategies for Ecological Restoration of the Middle Rio Grande in New Mexico and Recovery of the Endangered Rio Grande Silvery Minnow. *Reviews in Fisheries Science* 14: 169-186.
- Cowley DE, Sublette JE (1987a) Distribution of fishes in the Black River drainage, Eddy County New Mexico. *Southwestern Naturalist* **32**: 213-221.
- Cowley DE, Sublette JE (1987b) Food habits of *Moxostoma congestum* (Baird and Girard) and *Cycleptus elongatus* (Lesueur) (Catostomidae: Cypriniformes) taken from the Black River, Eddy County, New Mexico. *Southwestern Naturalist* 32: 411-413.
- Cragin FW (1885) Preliminary list of Kansas fishes. *Bulletin of the Washburn Collegoe Laboratory of Natural History* **1**: 105-111.
- Crandall KA, Bininda-Emonds OR, Mace GM, Wayne RK (2000) Considering evolutionary processes in conservation biology. *Trends in Ecology and Evolution* 15: 290-295.
- Cross F, Mayden R, Stewart J (1986) Fishes in the Western Mississippi Drainage. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt C, Wiley E), pp. 363-412. Wiley-Interscience, New York, NY.
- Cross FB (1967) Handbook of fishes of Kansas. University of Kansas Publications -Museum of Natural History **45**: 1-357.
- Cross FB, Collins JT (1975) *Fishes in Kansas* University of Kansas Museum of Natural History - Public Education Series 3.
- Cross FB, Collins JT (1995) *Fishes in Kansas*, 2 edn. University of Kansas Museum of Natural History Public Education Series 3.

- Cudmore-Vokey B, Crossman EJ (2000) Checklists of the Fish Fauna of the Laurentian Great Lakes and Their Connecting Channels. Canadian Manuscript Report of Fisheries and Aquatic Sciences 2550, Toronto, Ontario.
- Dettmers JM, Gutreuter S, Wahl DH, Soluk DA (2001) Patterns in abundance of fishes in main channels of the upper Mississippi River system. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 933-942.
- Dieringer D, Schlötterer C (2002) Microsatellite analyser (MSA): a platform independent analysis tool for large microsatellite data sets. . *Molecular Ecology Notes* 3: 167-169.
- Donnell O (1935) Annotated list of the fishes of Illinois. *Illinois Natural History Survey Bulletin* **20**: 473-500.
- Douglas NH (1974) Freshwater fishes of Louisiana Claitor's Publishing Division, Baton Rouge, Louisiana.
- Duncan JR, Lockwood JL (2001) Extinction in a field of bullets: a search for causes in the decline of the world's freshwater fishes. *Biological Conservation* **102**: 97– 105.
- Eastman JT (1977) The pharyngeal bones and teeth of catostomid fishes. *American Midland Naturalist* **97**: 68-88.
- Eastman JT (1980) The caudal skeletons of catostomid fishes. *American Midland Naturalist* **103**: 133-148.
- Eddy S, Surber T (1947) *Northern fishes* University of Minnesota Press, Minneapolis, MN.

- Eddy S, Underhill JC (1974) *Northern fishes*, 2 edn. University of Minnesota Press, Minneapolis, Minnesota.
- Eitzmann J, Makinster A, Paukert C (2005) Population Dynamics of Blue Suckers in the Kansas River. In: *Kansas Department of Wildlife and Parks Wildlife Surveys and Research: Assessment of Streams on Public Lands* (ed. Kansas Department of Wildlife and Parks ESS), p. 35. Kansas Cooperative Fish and Wildlife Research Unit, Division of Biology, Kansas State University
- Kansas Department of Wildlife and Parks, Environmental Services Section, Manhattan, KS.
- Elrod JH, Hassler TJ (1971) Vital statistics of seven fish species in Lake Sharpe, South
  Dakota 1964-1969. In: *Reservoir fisheries and limnology* (ed. Hall GE), pp. 2740. American Fisheries Society Special Publication 8.
- Elstad S, Werdon S (1993) Draft Status Report on Blue Sucker (Cycleptus elongatus), a Candidate Endangered or Threatened Species, p. 55. U.S. Fish and Wildlife Service, Ecological Services, Bismarck, ND, USA.
- EPA (1989) Rapid Bioassessment Protocols for Use in Streams and Rivers: Benthic Macroinvertebrates and Fish. Office of Water, U.S. Environmental Protection
   Agency Washington, D.C. EPA 444/4-89/001. 1989
- Eschmeyer WN (1998) Catalog of Fishes, p. 2905. California Academy of Sciences, San Francisco.

- Etnier DA, Starnes WC (1979) Whatever happened to the silvery minnow (Hybognathus nuchalis) in the Tennessee River? Proceedings of the Southeastern Fishes Council 2: 1-3.
- Etnier DA, Starnes WC (1991) An analysis of Tennessee's jeopardized fish taxa. *Journal* of the Tennessee Academy of Science **66**: 129-133.
- Etnier DA, Starnes WC (1993) *The fishes of Tennessee* University of Tennesse Press, Knoxville, Tennessee.
- Evermann BW (1886) A list of the fishes observed in the vicinity of Brookville, Franklin
  County, Indiana. *Bulletin No. 2 of the Brookville Society of Natural History* 1886:
  1-11.
- Evermann BW, Cox UO (1896) Report upon the the fishes of the Missouri River basin. Report of the United States Commissioner of Fish and Fisheries for 1894: 325-329.
- Excoffier L, Smouse P, Quattro J (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131: 479-491.
- Fago D (1984) Distribution and relative abundance of fishes in Wisconsin III. Red Cedar River basin. Wisconsin Department of Natural Resources Technical Bulletin 143: 1-69.
- Fago D (1992) Distribution and relative abundance of fishes in Wisconsin VIII. Summary report. Wisconsin Department of Natural Resources Technical Bulletin 175: 1-378.

Fago D, Hatch MD (1993) Aquatic resources of the St. Croix River. In: *Restoration* planning for the rivers of the Mississippi River ecosystem (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 23-56. National Biological Survey, Biological Report 19.

Faircloth BC (2006) GMCONVERT 0.32,

http://gallus.forestry.uga.edu/software/index.php, Athens, GA.

- Falush D, Stephens M, Pritchard J (2003) Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* 164: 1567-1587.
- Faugeron S, Martínez EA, Correa JA, et al. (2004) Reduced genetic diversity and increased population differentiation in peripheral and overharvested populations of *Gigartina skottsbergii* (Rhodophyta,Gigartinales) in southern Chile. *Journal of Phycology* **40**: 454-462.

Felsenstein J (2004) Inferring Phylogenies Sinauer Associates, Inc., Sunderland, MA.

- Ferris S, Whitt G (1978) Phylogeny of tetraploid catostomid fishes based on the loss of duplicate gene expression. *Systematic Zoology* 27: 189-206.
- Ferris SD (1984) Tetraploidy and the evolution of the catostomid fishes. In: *Evolutionary genetics of fishes* (ed. Turner BJ), pp. 55-93. Plenum Press, New York, New York.
- Ferris SD, Whitt GS (1977) Loss of duplicate gene expression and polyploidisation. *Nature* **265**: 258-260.
- Ferris SD, Whitt GS (1980) Genetic variability in species with extensive gene duplication: the tetraploid catostomid fishes. *American Naturalist* 115: 650-666.

- Fisher HJ (1962) Some fishes of the lower Missouri River. *American Midland Naturalist*68: 424-429.
- Fisher S, Willis D (2000) Observations of Age-0 Blue Sucker, Cycleptus elongatus Utilizing an Upper Missouri River Backwater *Journal of Freshwater Ecology* 15: 425-427.
- Forbes SA, Richardson RE (1908) *The fishes of Illinois* Illinois State Laboratory of Natural History.
- Fowler HW (1913) Notes on catostomid fishes. *Proceedings of the Academy of Natural Sciences of Philadelphia* **65**: 45-60.
- Fuiman LA (1985) Contributions of developmental characters to a phylogeny of catostomid fishes, with comments on heterochrony. *Copeia* **1985**: 833-846.
- Funk DJ, Omland KE (2003) Species-level paraphyly and polyphyly: frequency, causes, and consequences, with insights from animal mitochondrial DNA. *Annual Reviews in Ecology, Evolution, and Systematics* 34: 397-423.
- Galat DL, Lipkin R (2000) Restoring ecological integrity of great rivers: historical hydrographs aid in defining reference conditions for the Missouri River.
   *Hydrobiologia* 422-423: 29-48.
- Galat DL, Whitledge GW, Patton LD, Hooker J (2004) Larval Fish Use of Lower
   Missouri River Scour Basins in Relation to Connectivity Final Report to Missouri
   Department of Conservation.
- Gammon JR (1993) The Wabash River: progress and promise. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB,

Benson NG, Zuboy JR), pp. 142-161. National Biological Survey, Biological Report 19.

- Gardner B (2004) Pallid sturgeon research and recovery efforts in the upper Missouri River, MT. *Pallid Sturgeon Recovery Update* **14**: 1-2.
- Gehlbach FR, Miller RR (1961) Fishes from archaeological sites in northern New Mexico. Southwestern Naturalist 6: 2-8.
- Genton BJ, Shykoff JA, Giraud T (2005) High genetic diversity in French invasive populations of common ragweed, *Ambrosia artemisiifolia*, as a result of multiple sources of introduction. *Molecular Ecology* 14: 4275-4285.
- Glaubitz J (2004) CONVERT: A user-friendly program to reformat diploid genotypic data for commonly used population genetic software packages. *Molecular Ecology Notes* 4: 309-310.
- Goldstein DB, Schlötterer C (1999) *Microsatellites: evolution and applications* Oxford University Press, Oxford ; New York.
- Goodman S (1997) RST CALC: A collection of computer programs for calculating unbiased estimates of genetic differentiation and determining their significance for microsatellite data. *Molecular Ecology* **6**: 881-885.
- Goudet J (1995) fstat version 1.2: a computer program to calculate F-statistics. *Journal of Heredity* **86**: 485–486.
- Goudet J (2002) FSTAT Version 2.9.3.2, a Program to Estimate and Test Gene Diversities and Fixation Indices, Available at www2.unil.ch/izea/softwares/fstat.html (last accessed 25 June 2006).

- Graham ID (1885) Preliminary list of Kansas fishes. *Transactions of the Kansas* Academy of Science **9**: 69-78.
- Grande L, Bemis WE (1991) Osteology and phylogenetic relationships of fossil and recent paddlefishes (Polyodontidae) with comments on the interrelationships of Acipenseriformes. In: *Society of Vertebrate Paleontology, Memoir 1*, p. 121 pp. Society of Vertebrate Paleontology, Memoir 1.
- Guillory V (1982) Fishes of the Lower Mississippi River near St. Francisville, Louisiana. *Proceedings of the Louisiana Academy of Sciences* **45**: 108-121.
- Hand G, Jackson D (2003a) Blue sucker stock characteristics in the upper Yazoo River basin, Mississippi, USA. *Fisheries Management and Ecology* **10**: 147-153.
- Hand GR (1999) Stock dynamics of the blue sucker (Cycleptus elongatus) in the principal tributaries of the Yazoo River, Mississippi, Mississippi State University.
- Hand GR, Jackson DC (2003b) Blue sucker stock characteristics in the upper Yazoo River basin, Mississippi, USA. *Fisheries Management and Ecology* **10**: 147-154.
- Harlan JR, Speaker EB (1956) *Iowa fish and fishing* Iowa Conservation Commission, Des Moines, IA.
- Harlan JR, Speaker EB (1987) *Iowa fish and fishing* Iowa Department of Natural Resources, Des Moines, Iowa.

Harrington S (1999) Occurrence and Distribution of Fish Species in the Great and Little Miami River Basins, Ohio and Indiana, Pre-1900 to 1998, p. 16. US Geological Survey, Columbus, OH, USA.

- Harris P, Mayden R (2001) Phylogenetic relationships of major clades of Catostomidae (Teleostei: Cypriniformes) as inferred from mitochondrial SSU and LSU rDNA sequences. *Molecular Phylogenetics and Evolution* 20: 225-237.
- Harris P, Mayden R, Espinosa Perez H, Garcia de Leon F (2002) Phylogenetic
  relationships of *Moxostoma* and *Scartomyzon* (Catostomidae) based on
  mitochondrial cytochrome b sequence data. *Journal of Fish Biology* 61: 1433-1452.
- Harrison HM, Speaker EB (1954) An annotated list of the fishes in the streams tributary to the Missouri River in Iowa. *Proceedings of the Iowa Academy of Science* 61: 511-523.
- Haslouer S, Eberle M, Edds D, et al. (2005) Current status of native fish species in Kansas. Transactions of the Kansas Academy of Science 108: 32-46.
- Hauswaldt J, Glenn T (2003) Microsatellite DNA loci from the Diamondback terrapin (*Malaclemys terrapin*). *Molecular Ecology Notes* **3**: 174-176.
- Henegariu O, Heerema N, Dlouhy S, Vance G, Vogt P (1997) Multiplex PCR: Critical Parameters and Step-by-Step Protocol. *BioTechniques* **23**: 504-511.
- Henshall JA (1888) Contributions to the ichthyology of Ohio No. 1. *Journal of the Cincinnati Society of Natural History* **11**: 76-80.
- Herzog DP (2004) Capture efficiency and habitat use of sturgeon chub (Macrhybopsis gelida) and sicklefin chub (Macrhybopsis meeki), Southeast Missouri State University.
- Hesse L, Hergenrader G, Lewis H, Reetz S, Schlesinger A (1982) *The Middle Missouri River* The Missouri River Study Group, Norfolk, NE.

- Hey J, Nielsen R (2004) Multilocus methods for estimating population sizes, migration rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura* and D. persimilis. *Genetics* 167: 747-760.
- Hoagstrom CW, Berry CR, Jr. (2006) Island Biogeography of Native Fish Faunas among
   Great Plains Drainage Basins: Basin Scale Features Influence Composition.
   American Fisheries Society Symposium 48: 221–264.
- Holman T, Skains J, Riecke D (1993) The Big Black River, Mississippi: a case history In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse
  LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 266-281. National Biological
  Survey, Biological Report 19.
- Hoy PR (1883) Catalogue of the cold-blooded vertebrates of Wisconsin, 1873-1879. Part
  III. Fishes. In: *Geology of Wisconsin* (ed. Chamberliain TC), pp. 427-435. D.
  Atwood Publishing, Madison, WI.
- Hubbs C (1957) Distributional patterns of Texas fresh-water fishes. *Southwestern Naturalist* **2**: 89-104.
- Hubbs C, Pigg J (1976) The effects of impoundments on threatened fishes of Oklahoma. Annals of the Oklahoma Academy of Science **5**: 113-117.
- Huelsenbeck J, Ronquist F (2001) MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* **17**: 754-755.
- Huson D, Bryant D (2006) Application of Phylogenetic Networks in Evolutionary Studies. *Molecular Biology and Evolution* 23: 254-267.

- Ickes BS, Wlosinski JH, Knights BC, Zigler SJ (2001) *Fish passage through dams in large temperate floodplain rivers: an annotated bibliography* U.S. Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin.
- Jackson DC, Brown-Peterson NJ, Rhine TD (1993) Perspectives for rivers and their fishery resources in the Yazoo River basin, Mississippi. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 255-263. National Biological Survey, Biological Report 19.
- Jenkins RE, Burkhead NM (1994) *Freshwater fishes of Virginia* American Fisheries Society, Bethesda, Maryland.
- Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. *BMC Genetics* **6**: 1-6.
- Jett SC, Moyle PB (1986) The exotic originsof fishes depicted on prehistoric Mimbres pottery from New Mexico. *American Antiquity* **51**: 688-720.
- Johnson JE (1987) *Protected fishes of the United States and Canada* American Fisheries Society, Bethesda, Maryland.
- Jordan DS (1875) Synopsis of the genera of fishes to be looked for in Indiana. *6th Annual Report for the U.S. Geological Survey of Indiana* [for 1874]: 197-228.

Jordan DS (1877) Contributions to North American ichthyology, based primarily on the collections of the United States National Museum. I. Review of Rafinesque's memoirs on North American fishes. *Bulletin of the U.S. National Museum* 9: 1-53.

Jordan DS (1878) A catalog of fishes of the fresh waters of North America. *Bulletin IV*, *Hayden's Geological Survey Territory* **2**: 407-442.

- Jordan DS (1929) Manual of the vertebrate animals of the northeastern United States. Inclusive of marine species, 13 edn. World Book Company, Yonkers-on-Hudson, New York.
- Jordan DS, Brayton AW (1878) On the distribution of the fishes of the Allegheny Region of South Carolina, Georgia, and Tennessee, with descriptions of new or little known species. *Bulletin of the U.S. National Museum* **12**: 1-96.
- Jordan DS, Evermann BW (1886) The food fishes of Indiana. *36th Report for 1886, Indiana State Board of Agriculture* **28**: 156-173.
- Jordan DS, Gilbert CH (1883) Syunopsis of the fishes of North America. *Bulletin of the* U.S. National Museum **16**: 1-1018.
- Jungwirth M, Schmutz S, Weiss S (1998) *Fish Migration and Fish Bypasses* Blackwell Science Ltd., Oxford, UK.
- Kay LK, Wallus R, Yeager BL (1994) Reproductive biology and early life history of fishes in the Ohio River drainage. Volume 2: Catostomidae Tennessee Valley Authority, Chattanooga, Tennessee.
- Koppelman JB (1994) Hybridization between Smallmouth Bass, Micropterus dolomieu, and Spotted Bass, M. punctulatus, in the Missouri River System, Missouri.*Copeia* 1994: 204-210.
- Koster WJ (1957) *Guide to the fishes of New Mexico* University of New Mexico Press, Albuquerque, New Mexico.
- Kreiser BR, Mitton JB, Woodling JD (2001) Phylogeography of the plains killifish, *Fundulus zebrinus*. Evolution 55: 339-350.

- Kumar S, Tamura K, Nei M (2004) MEGA3: Integrated software for Molecular
   Evolutionary Genetics Analysis and sequence alignment 5:150-163 *Briefings in Bioinformatics* 5: 150-163.
- Larimore RW, Fritz AW (1993) Environmental changes in the Kaskaskia River basin. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 210-240. National Biological Survey, Biological Report 19.
- Laroche J, Durand J (2004) Genetic structure of fragmented populations of a threatened endemic percid of the Rhone river: *Zingel asper. Heredity*: 329-334.
- Leiby PD, Kritsky DC, Bauman DD (1973) Studies on helminths of North Dakota. VII. Ancyrocephalinae (Monogenea) from the fills of the blue sucker, *Cycleptus elongatus* (LeSueur). *Canadian Journal of Zoology* **51**: 777-779.
- Leidy RA, Moyle PB (1998) Conservation status of the world's freshwater fish fauna: an overview. In: *Conservation Biology: For the Coming Decade* eds. Fieldler PL, Karieva PM), pp. 187–227. Chapman and Hall, New York.
- Lesueur CA (1817) A new genus of fishes, of the order Abdominales, proposed, under the name of *Catostomus*; and the characters of this genus, with those of its species, indicated. *Journal of the Academy of Natural Sciences of Philadelphia* 1: 88-96, 102-111.
- Lewis WM (1955) The fish population of the main stream of the Big Muddy River. *Transactions of the Illinois State Science Academy* **47**: 20-24.
- Limbird RL (1993) The Arkansas River a changing river. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB,

Benson NG, Zuboy JR), pp. 282-294. National Biological Survey, Biological Report 19.

- Lindsay H, Randolph C, Carroll J (1983) Updated survey of the fishes of the Poteau River, Oklahoma and Arkansas. *Proceedings of the Oklahoma Academy of Science* **63**: 42-48.
- Lippe C, Dumont P, Bernatchez L (2004) Isolation and identification of 21 microsatellite loci in the Copper redhorse (*Moxostoma hubbsi*; Catostomidae) and there variability in other catostomids. *Molecular Ecology Notes* **4**: 638-641.
- Lundberg JG, Marsh E (1976) Evolution and functional anatomy o fthe pectoral fin rays in cyprinoid fishes, with emphasis on the suckers (family Catostomidae). *American Midland Naturalist* **96**: 332-349.
- Marcogliese DJ (2001) Implications of climate change for parasitism of animals in the aquatic environment. *Canadian Journal of Zoology* **79**: 1331–1352.
- Mayden RL, Burr BM, Page LM, Miller RR (1992) The Native Freshwater Fishes of North America. In: Systematics, Historical Ecology, and North American Freshwater Fishes (ed. Mayden RL), pp. 827-863. Stanford University Press, Stanford, CA.
- Mayr E (1942) *Systematics and the Origin of Species* University Press, New York, New York.
- McInerny MC, Held JW (1988) Collections of young-of-the-year blue suckers (Cycleptus elongatus) in navigation pool 9 of the upper Mississippi River. *Transactions of the Wisconsin Academy of Sciences, Arts, and Letters* **76**: 69-71.

- McReynolds HE, Whitaker JO, Gammon JR (1979) Development of a proposed list of endangered and threatened vertebrate animals in Indiana. *Proceedings of the Indiana Academy of Science* 88: 166-170.
- Meek SE (1890) The native food fishes of Iowa. *Proceedings of the Iowa Academy of Science* **1187-1889**: 68-76.
- Meek SE (1908) Notes on fresh-water fishes from Mexico and Central America. *Field Columbian Museum, Zoological Series* **7**: 133-157.
- Metcalf AL (1959) Fishes of Chautauqua, Cowley, and Elk Counties, Kansas. *University* of Kansas Publications - Museum of Natural History **11**: 345-400.
- Mettee MF, O'Neil PE, Pierson JM (1996) *Fishes of Alabama and the Mobile Basin* Oxnoor House, Birmingham, Alabama.
- Mettee MF, Shepard TE (1997) Status survey of a blue sucker (Cycleptus elongatus)
   population below Millers Ferry lock and dam, Wilcox County, Alabama
   Geological Survey of Alabama. Final Report, Tuscaloosa, Alabama.
- Miller RJ, Evans HE (1965) External morphology of the brain and lips in catostomid fishes. *Copeia* **1965**: 467-487.
- Miller RJ, Robison H (1973) *The fishes of Oklahoma* Oklahoma State University Press, Stillwater, Oklahoma.
- Miller RR (1972) Threatened freshwater fishes of the United States. *Transactions of the American Fisheries Society* **101**: 239-252.
- Mills H, Kaye J (2001) Drainage history of the Tennessee River: Review and new metamorphic quartz locations. *Southeastern Geology* **40**: 75–97.

- Minckley WL (1959) Fishes of the Big Blue River basin, Kansas. University of Kansas Publications - Museum of Natural History **11**: 401-442.
- Mishler BD, Theriot EC (2000) The Phylogenetic Species Concept (*sensu* Mishler and Theriot): Monophyly, Apomorphy, and Phylogenetic Species Concepts. In:
  Species Concepts and Phylogenetic Theory (eds. Wheeler QD, Meier R), pp. 44-54. Columbia University Press, New York.
- Moore GA, Cross FB (1950) Additional Oklahoma fishes with validation of *Poecilichthys parvipinnis* (Gilbert and Swain). *Copeia* **1950**: 139-148.
- Morey N, Berry CJ (2003) Biological Characteristics of the Blue Sucker in the James River and the Big Sioux River, South Dakota *Journal of Freshwater Ecology* **18**: 33-42.
- Mortiz C (1994) Defining 'Evolutionarily Significant Units' for conservation. *Trends in Ecology and Evolution* **9**: 373-375.
- Moss RE, Scanlan JW, Anderson CS (1983) Observations of the natural history of the blue sucker (*Cycleptus elongatus* Le Sueur) in the Neosho River. *The American Midland Naturalist* 109: 15-21.
- Moyle PB, Cech JJ (2000) *Fishes: An introduction to ichthyology*, 4 edn. Prentice Hall, Upper Saddle River, NJ.
- NatureServe (2005) NatureServe Explorer: An online encyclopedia of life [web application]. Version 4.6. NatureServe.
- Near TJ, Keck BP (2005) Dispersal, vicariance, and timing of diversification in *Nothonotus* darters. *Molecular Ecology* **14**: 3485-3496.

- Nei M (1987) *Molecular Evolutionary Genetics* Columbia University Press, New York, New York.
- Nelson EM (1948) The comparative morphology of the Weberrian apparatus of the Catostomidae and its significance in systematics. *Journal of Morphology* **83**: 225-251.
- Nelson EW (1876) A partial catalogue of the fishes of Illinois. *Illinois Museum of Natural History Bulletin* 1: 33-52.

Nelson JS (1994) Fishes of the World, 3 edn. John Wiley and Sons, New York.

- Nielsen R, Wakeley J (2001) Distinguishing migration from isolation: a Markov chain Monte Carlo approach. *Genetics* **158**: 885-896.
- O' Hara RJ (1988) Homage to Clio, or, toward an historical philosophy for evolutionary biology *Systematic Zoology* **37**: 142-155.
- Ohta T, Kimura M (1973) The model of mutation appropriate to estimate the number of electrophoretically detectable alleles in a genetic population. *Genetic Research* 22: 201-204.
- Omland KE, Baker JM, Peters JL (2006) Genetic signatures of intermediate divergence:
   population history of Old and New World Holarctic ravens (*Corvus corax*).
   *Molecular Ecology* 15: 795-808.
- Page LM, Burr BM (1991) A field guide to the freshwater fishes of North America north of Mexico Houghton Mifflin, Boston, Massachusetts.
- Pearse D, Crandall K (2004) Beyond Fst: analysis of population genetic data for conservation. *Conservation Genetics* 5: 585-602.

- Pearson WD, Froedge MA (1989) Stranding of fishes below McAlpine Dam on the Ohio River. *Transactions of the Kentucky Academy of Science* **50**: 183-201.
- Pegg MA, Pierce CL (2002) Fish community structure in the Missouri and lower Yellowstone rivers in relation to flow characteristics. *Hydrobiologia* 479: 155-167.
- Peng Z, He S, Wang J, Wang W, Diogo R (2006) Mitochondrial molecular clocks and the origin of the major Otocephalan clades (Pisces: Teleostei): A new insight. *Gene* 370: 113-124.
- Pennington CH, Baker JA, Potter ME (1983) Fish populations along natural and revetted banks on the lower Mississippi River. North American Journal of Fisheries Management 3: 2204-2211.
- Peterson MS, Nicholson LC, Snyder DJ, Fulling GL (1999) Growth, spawning preparedness, and diet of *Cycleptus meridionalis* (Catostomidae). *Transactions of the American Fisheries Society* **128**: 900-908.
- Pflieger W (1971) A distributional study of Missouri fishes. *University of Kansas Publications - Museum of Natural History* **20**: 225-570.
- Pflieger W (1997) *The Fishes of Missouri* Missouri Department of Conservation, Columbia, MO, USA.
- Pflieger WL (1975) *The fishes of Missouri* Missouri Department of Conservation, Jefferson City, Missouri.
- Phillips GL, Schmid WD, Underhill JC (1982) *Fishes of the Minnesota region* University of Minnesota Press, Minneapolis, Minnesota.

- Pigg J, Hill LG (1974) Fishes of the Kiamichi River, Oklahoma. Proceedings of the Oklahoma Academy of Science 54: 93-94.
- Pimentel D, Lach L, Zuniga R, Morrison D (2000) Environmental and Economic Costs of Nonindigenous Species in the United States *BioScience* 50: 53-65.
- Platt DR, Cross FB, Distler D, et al. (1974) Rare, endangered, and extirpated species in Kansas. Transactions of the Kansas Academy of Science 76: 97-105.
- Posada D, Crandall K (1998) Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817-818.
- Pringle CM, Freeman MC, Freeman BJ (2000) Regional Effects of Hydrologic Alterations on Riverine Macrobiota in the New World: Tropical-Temperate Comparisons. *BioScience* 50: 807-823.
- Pritchard JK, Stephens M, Donnell P (2000) Inference of Population Structure Using Multilocus Genotype Data. *Genetics* 155: 945-959.
- Pritchard JK, Wen W (2004) Documentation for *structure* software: Version 2. University of Chicago, Chicago.
- Propst DL (1999) *Threatened and endangered fishes of New Mexico* New Mexico Department of Game and Fish Technical Report No. 1, Santa Fe, New Mexico.
- Propst DL, Burton GL, Pridgeon BH (1987) Fishes of the Rio Grande between Elephant Butte and Caballo Reservoirs. *Southwestern Naturalist* **32**: 408-411.
- Rafinesque CS (1820) Ichthyologia Ohiensis [Part 7]. *Western Review and Miscellaneous Magazine* **6**: 355-363.

- Ramstad KM, Woody CA, Sage GK, Allendorf FW (2004) Founding events influence genetic population structure of sockeye salmon (*Oncorhynchus nerka*) in Lake Clark, Alaska. *Molecular Ecology* 13: 277-290.
- Raymond M, Rousset F (1995) GENEPOP (Version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86: 248-249.
- Retzer ME, Burr BM, Warren ML (1983) Fishes of the lower Green River drainage, Kentucky. In: Kentucky Nature Preserves Commission Scientific and Technical Series 3, p. 48.
- Rice W (1989) Analyzing tables of statistical test. Evolution 43: 223-225.
- Richter B, Braun D, Mendelson M, Master L (1997) Threats to imperiled freshwater fauna. *Conservation biology* 11: 1081-1093.
- Robison H (1986) Zoogeographic Implications of the Mississippi River Basin. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt C, Wiley E), pp. 267-286. Wiley-Interscience, New York, NY.
- Robison H, Buchanan T (1998) *Fishes of Arkansas* University of Arkansas Press, Fayetteville, AR.
- Robison H, Moore GA, Jahn LA (1974) Threatened fishes of Oklahoma. *Proceedings of the Oklahoma Academy of Science* **54**: 139-146.
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
- Ross ST, Brenneman WM (1991) *Distribution of freshwater fishes in Mississippi* Mississippi Department of Wildlife, Fisheries, and Parks; Bureau of Fisheries and Wildlife, Jackson, Mississippi.

Rousset F (1997) Genetic Differentiation and Estimation of Gene Flow from F-Statistics Under Isolation by Distance. *Genetics* **145**: 1219-1228.

Ruelle R, Koth R, Stone C (1993) Contaminents, fish, and hydrology of the Missouri River and western tributaries, South Dakota. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 449-480. National Biological Survey, Biological Report 19.

- Rupprecht RJ, Jahn LA (1980) Biological notes on blue suckers in the Mississippi River. *Transactions of the American Fisheries Society* **109**: 323-326.
- Rutledge WP (1989) *Fishery management realities versus fishery genetic fantasies* Texas Parks and Wildlife Dept., Fisheries Division, Fish Hatcheries Branch, Austin, TX.
- Saitoh K, Miya M, Nishida M (2003) Complete nucleotide sequence of mitochondrial genome from river carpsucker, *Carpiodes carpio*. NCBI.
- Saitoh K, Miya M, Nishida M (2005) *Myxocyprinus asiaticus* mitochondrial genome, complete sequence. NCBI.
- Sambrook J, Fritsch E, Maniatis T, Nolan C (1989) *Molecular cloning: a laboratory manual*, 2 edn. Cold Spring Harbor Laboratory Press Woodbury, NY.
- Sanders RM, Jr., Huggins DG, Cross FB (1993) The Kansas River system and its biota.
  In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 295-326. National Biological Survey, Biological Report 19.
- Sandheinrich MB, Atchison GJ (1986) Fish associated with dikes, revetments, and abandoned channels in the middle Missouri River. *Proceedings of the Iowa Academy of Science* **93**: 188-191.

- Schmidt K, Talmage P (2001) Fish community surveys of Twin Cities Metropolitan Area streams Minnesota Department of Natural Resources Special Publication 156, St. Paul, Minnesota.
- Schmulbach JC, Braaten PJ (1993) The Vermillion River: neither red nor dead. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 57-69. National Biological Survey, Biological Report 19.
- Schneider S, Roessli D, Excoffier L (2000) Arlequin ver 2.000: a software for population genetic data analysis. Genetics and Biometry Laboratory, University of Geneva, Geneva, Switzerland.
- Segelbacher G, Hoglund J, Storch I (2003) From connectivity to isolation: genetic consequences of population fragmentation in capercaillie across Europe. *Molecular Ecology* **12**: 1773–1780.
- Simons AM, Wood RM, Heath LSK, B.R., Mayden RL (2001) Phylogenetics of Scaphirhynchus based on mitochondrial DNA sequences. Transactions of the American Fisheries Society 130: 359-366.
- Smith GR (1981) Late Cenozoic freshwater fishes of North America. *Annual Reviews in Ecology and Systematics* **12**: 163-193.

Smith GR (1992) Phylogeny and biogeography of the Catostomidae, freshwater fishes of North America and Asia. In: Systematics, Historical Ecology, and North American Freshwater Fishes (ed. Mayden RL), pp. 778-813. Stanford University Press, Stanford, CA. Smith M, Miller R (1986) The Evolution of the Rio Grande Basin as Inferred from Its Fish Fauna. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt C, Wiley E), pp. 457-486. Wiley-Interscience, New York, NY.

Smith PW (1979) The fishes of Illinois University of Illinois Press, Urbana, Illinois.

- Smith PW, Lopinot AC, Pflieger WL (1971) A distributional atlas of upper Mississippi River fishes. *Illinois Natural History Survey Biological Notes* 73: 20.
- Starnes W, Etnier D (1986) Drainage Evolution and Fish Biogeography of the Tennessee and Cumberland Rivers Drainage Realm. In: *The Zoogeography of North American Freshwater Fishes* (eds. Hocutt C, Wiley E), pp. 325-362. Wiley-Interscience, New York, NY.
- Starnes WC, Etnier DA (1980) Fishes. In: *Tennessee's rare wildlife* (eds. Eager DC, Hatcher RM), pp. B1-B4. Tennessee Department of Conservation, Nashville, Tennessee.
- Stauffer JE, Boltz JM, White LR (1995) The fishes of West Virginia. *Proceedings of the Academy of Natural Sciences of Philadelphia* **146**: 1-389.
- Stein BA (2002) States of the Union: Ranking America's Biodiversity, pp. 1-27. NatureServe, Arlington, VA.
- Sublette JE, Hatch MD, Sublette M (1990) *The fishes of New Mexico* University of New Mexico Press, Albuquerque, New Mexico.
- Swift C (1968) Pleistocene freshwater fishes from Ingleside Pit, San Patricio County, Texas. *Copeia* **1968**: 63-69.
- Swingle HA (1971) Biology of Alabama estuarine areas Cooperative Gulf of Mexico estuarine inventory. *Alabama Marine Resources* 5: 1-123.

- Swofford D (1998) "PAUP": Phylogenetic analysis using parsimony (\*and other methods) p. . Sinauer Associates, Sutherland, MA.
- Templeton AR (1989) The meaning of species and speciation: a genetic perspective. In. In: Speciation and its Consequences (eds. Otte D, Endler JA), pp. 3-27. Sinauer Associates, Sunderland, Massachusetts.
- Thompson J, Gibson T, Plewniak F, Jeanmougin F, Higgins D (1997) The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* **24**: 4876-4882.
- Tomelleri JR, Eberle ME (1990) Fishes of the central United States University Press of Kansas, Lawrence, Kansas.
- Tranah GJ, Agresti JJ, May B (2001) New microsatellite loci for suckers (Catostomidae): primer homology in *Catostomus, Chasmistes, and Deltistes. Molecular Ecology Notes* 1: 55-60.
- Trautman MB (1957) The fishes of Ohio Ohio State University Press, Columbus, Ohio.

Trautman MB (1981) The fishes of Ohio, 2 edn. Ohio State University Press, Columbus.

- Underhill JC (1959) Fishes of the Vermillion River, South Dakota. *Proceedings of the South Dakota Academy of Sciences* **38**: 96-102.
- USACE (2001) Missouri River master water control manual: Review and update. US Army Corps of Engineers, Northwest Division, Omaha, NE.
- Uyeno T, Smith GR (1972) Tetraploid origin of the karyotype of catostomid fishes. *Science* **175**: 644-645.
- Valdes AM, Slatkin M, Freimer NB (1993) Allele frequencies at microsatellite loci: the stepwise mutation model revisited. *Genetics* **133**: 737-749.

- Vallone PM, Butler JM (2004) AutoDimer: a screening tool for primer-dimer and hairpin structures. *BioTechniques* 37: 226-231.
- VanOosterhout C, Hutchinson WF, Wills DP, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535-538.
- Vokoun JC, Guerrant TL, Rabeni CF (2003) Demographics and chronology of a spawning aggregation of blue sucker (*Cycleptus elongatus*) in the Grand River, Missouri, USA. *Journal of Freshwater Ecology* 18: 567-576.
- Walburg CH, Kaiser GL, Hudson PL (1971) Lewis and Clark tailwater biota and some relations of the tailwater and reservoir fish populations, pp. 449-467. In: *Reservoir fisheries and limnology* (ed. Hall GE). American Fisheries Society Special Publication 8.
- Waples R, Gaggiotti O (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology* 15: 1419-1439.
- Warren ML, Burr BM, Walsh SJ, *et al.* (2000) Diversity, distribution, and conservation status of the native freshwater fishes of the southern United States. *Fisheries* 25: 7-29.
- Wehrhahn C (1975) The evolution of selectively similar electrophoretically detectable alleles in finite natural populations. *Genetics* **80**: 375-394.
- Weigel DE, Peterson JT, Spruell P (2003) Introgressive hybridization between native cutthroat trout and introduced rainbow trout. *Ecological Applications* **13**: 38-50.

- Weir B, Cockerham C (1984) Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358-1370.
- Welker TL, Scarnecchia DL (2003) Differences in Species Composition and Feeding
  Ecology of Catostomid Fishes in Two Distinct Segments of the Missouri River,
  North Dakota, U.S.A. *Environmental Biology of Fishes* 68: 129-141.
- Wheeler QD, Meier R (2000) Species concepts and phylogenetic theory: a debate, p. 230. Columbia University Press, New York, New York.
- White RG, Bramblett RG (1993) The Yellowstone River: its fish and fisheries. In: *Restoration planning for the rivers of the Mississippi River ecosystem* (eds. Hesse LW, Stalnaker CB, Benson NG, Zuboy JR), pp. 396-414. National Biological Survey, Biological Report 19.
- Wiley EO (1978) The evolutionary species concept reconsidered. *Systematic Zoology* **27**: 17-26.
- Wilgenbusch J, Warren D, Swofford D (2004) AWTY: A system for graphical exploration of MCMC convergence in Bayesian phylogenetic inference.
- Williams JE, Bowman DB, Brooks JE, et al. (1985) Endangered aquatic ecosystems in North American deserts with a list of vanishing fishes of the region. Journal of the Nevada Academy of Science 20: 1-62.
- Williams JE, Johnson JE, Hendrickson DA, *et al.* (1989) Fishes of North America endangered, threatened, or of special concern: 1989. *Fisheries* **14**: 2-20.
- Wilson C, Hebert P, Reist J, Dempson J (1996) Phylogeography and postglacial dispersal of arctic charr *Salvelinus alpinus* in North America. *Molecular Ecology* 5: 187-197.

- Wolter C, Arlinghaus R (2003) Navigation impacts on freshwater fish assemblages: the ecological relevance of swimming performance. *Reviews in Fish Biology and Fisheries* **13**: 63-89.
- Woolman AJ (1892) Report of an examination of the rivers of Kentucky, with lists of the fishes obtained. *Bulletin of the U.S. Fish Commission* **10**: 249-288.

Wright S (1943) Isolation by distance. Genetics 28: 114-138.

Wua R, Gallo-Meagherb M, Littella R, Zeng Z (2001) A general polyploid model for analyzing gene segregation in outcrossing tetraploid species. *Genetics* 159: 869-882. **APPENDIX ONE**: Edited clone sequences for each microsatellite locus

**Bold, underlined** lettering indicates name and primer locations for loci in which primers were designed and optimized for PCR.

# Ce13 original clone

# \*Ce13S allele

# \*Ce13L allele

# Ce35 clone

#### Ce49 clone

ACAAACAACTTCATTAAACACACACAGGTAATCATAATTACAGTTTAAAGAG AGAGAGATGATGCATTT<u>TTAAGATTTTCTTCCTTCGACTAA</u>ACAACAACA ACAACAACAACAACCTCCAAACACATCTGAATTCAATCCTGACATGTATTCT GTTTG<u>TGTTCATGCGCACGGGCACATTC</u>ATCAACTGAACTGACAAAGATAA CAGATTATTTAGACAAAACCCGTGTGAGGCCGTCCTGTTTTGGACGATAGTTT TTGCCCCGAGACTGTCGAATC

### Ce52 clone

### Ce63 clone

#### Ce104 clone

#### Ce126 clone

# Ce146 clone

# Ce195 clone

# Ce215 clone

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
1-1	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-2	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-3	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-4	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-5	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-6	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-7	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-8	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-9	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-10	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-11	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-12	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
1-13	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-14	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-15	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/15/2004	S. Mettee	Alabama GS
1-16	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-17	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-18	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-19	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-20	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-21	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-22	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-23	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-24	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-25	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-26	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
1-27	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/16/2004	S. Mettee	Alabama GS
1-28	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/30/2004	S. Mettee	Alabama GS
1-29	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/30/2004	S. Mettee	Alabama GS
1-30	C. meridionalis	AL	Alabama	132.5	32.08357	87.40307	3/30/2004	S. Mettee	Alabama GS Univ. of
1-31	C. meridionalis	AL	Alabama					R. Mayden	Alabama
1-32	C. meridionalis	AL	Alabama					R. Mayden	Univ. of Alabama Univ. of
1-33	C. meridionalis	AL	Alabama					R. Mayden	Alabama
2-1	C. elongatus	AR	Red		33.61083	93.85639	9/24/2004	W. Layher	Layner Biologics Layher
2-2	C. elongatus	AR	Red		33.61083	93.85639	9/24/2004	W. Layher	Biologics
2-3	C. elongatus	AR	Red		33.61083	93.85639	9/24/2004	W. Layher	Layner Biologics MS Mus.
4-1	C. meridionalis	MS	Leaf		31.2204	89.0285	10/7/2003	T. Slack	Nat. Sci.
4-2	C. meridionalis	MS	Leaf		31.2204	89.0285	10/7/2003	T. Slack	MS Mus. Nat. Sci. Univ.
4-3	C. meridionalis	MS	Leaf		31.19459	89.16099	6/22/2004	B. Krieser	Southern MS
4-4	C. meridionalis	MS	Leaf		31.06121	88.48307	6/22/2004	B. Krieser	Southern MS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
4-5	C meridionalis	MS	Leaf	02			3/16/2004	B Krieser	Univ. Southern MS
8-21	C. elongatus	AR	White	258.7		3945919	3/7/2006	J. Ouinn	AR Game and Fish
0 70	C l en	۸D	White	250 7		2045010	2/7/2006	I Quinn	AR Game
0-20	C. elongatus	AK	w mite	238.7		3943919	5/ //2000	J. Quini	Purdue
9-1	C. elongatus	IN	Wabash	321	40.51317	86.79308	4/21/2004	T. Kennedy	University Purdue
9-2	C. elongatus	IN	Wabash	321	40.51317	86.79308	4/21/2004	T. Kennedy	University Purdue
9-3	C. elongatus	IN	Wabash	315	40.47564	86.87091	4/3/2004	T. Kennedy	University Purdue
9-4	C. elongatus	IN	Wabash	321	40.51317	86.79308	4/21/2004	T. Kennedy	University
9-5	C. elongatus	IN	Wabash	300	40.40429	86.06691	4/10/2004	T. Kennedy	University
9-6	C. elongatus	IN	Wabash	321	40.51317	86.79308	4/21/2004	T. Kennedy	University
9-7	C. elongatus	IN	Wabash	380	40.77552	85.92357	5/19/2004	T. Kennedy	University Purdue
9-8	C. elongatus	IN	Wabash	300	40.40429	86.06691	5/6/2004	T. Kennedy	University
9-9	C. elongatus	IN	Wabash	300	40.40429	86.06691	4/10/2004	T. Kennedy	University
9-10	C. elongatus	IN	Wabash	315	40.47564	86.87091	3/27/2004	T. Kennedy	University
9-11	C. elongatus	IN	Wabash	315	40.47564	87.87091	4/3/2004	T. Kennedy	University
				River	Latitude	Longitude	Date of		Institution or
----------------------	--------------	-------	-----------	----------	-----------	-----------	-------------------------	------------	----------------------
Individual	Species	State	River	mile	(N)	(W)	collection	Collector	agency
<b>0</b> _1 <b>2</b>	C alongatus	IN	Wabash	315	10 17561	86 87001	3/27/2004	T Kennedy	Purdue University
9-12	C. elongalus	111	vv abasii	515	40.47304	00.07071	3/2//2004	1. Kennedy	Purdue
9-13	C. elongatus	IN	Wabash	315	40.47564	87.87091	4/3/2004	T. Kennedy	University
	0							2	Purdue
9-14	C. elongatus	IN	Wabash	300	40.40429	87.06691	6/9/2004	T. Kennedy	University
									Purdue
9-15	C. elongatus	IN	Wabash	300	40.40429	86.06691	5/6/2004	T. Kennedy	University
									Purdue
9-16	C. elongatus	IN	Wabash	315	40.47/564	87.87091	4/3/2004	T. Kennedy	University
0.17	~ .	DI	XX7 1 1	200	40 40 400	06.06601	51612004	TT 12 1	Purdue
9-17	C. elongatus	IIN	wabash	300	40.40429	86.06691	5/6/2004	I. Kennedy	Durdue
9_18	Calanastus	IN	Wabash	321	10 51317	86 79308	4/21/2004	T Kennedy	Liniversity
)-10	C. elongalus	111	vv abasii	$J_{21}$	40.31317	80.77508	4/21/2004	1. Kennedy	Purdue
9-19	C elongatus	IN	Wabash	321	40 51317	86 79308	4/21/2004	T Kennedy	University
, ,,	e. cionganas			0-1		001/2000	.,,		Purdue
9-20	C. elongatus	IN	Wabash	315	40.47564	86.87091	3/27/2004	T. Kennedy	University
	0							5	Purdue
9-21	C. elongatus	IN	Wabash	304	40.40764	87.00857	6/9/2004	T. Kennedy	University
									Purdue
9-22	C. elongatus	IN	Wabash	380	40.77552	85.92357	5/19/2004	T. Kennedy	University
									Purdue
9-23	C. elongatus	IN	Wabash	300	40.40429	86.06691	4/10/2004	T. Kennedy	University
0.04		DI	XX 7 1 1	• • • •		05.05.400	- 11 0 / <b>2</b> 0 0 4		Purdue
9-24	C. elongatus	IN	Wabash	380	40.76648	85.95422	5/19/2004	T. Kennedy	University
0.25		INT	Wahaah	215	10 17561	06 07001	1/2/2004	TVannada	Purdue
9-23	C. elongatus	11N	wabash	313	40.4/304	00.0/091	4/3/2004	I. Kennedy	University

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
9-26	C. elongatus	IN	Wabash	304	40.40764	87.00857	6/9/2004	T. Kennedy	University
9-27	C. elongatus	IN	Wabash	380	40.77552	85.92357	5/19/2004	T. Kennedy	University
9-28	C. elongatus	IN	Wabash	300	40.40429	86.06691	4/10/2004	T. Kennedy	University
9-29	C. elongatus	IN	Wabash	315	40.47564	86.87091	3/27/2004	T. Kennedy	University
9-30	C. elongatus	IN	Wabash	300	40.40429	86.06691	4/10/2004	T. Kennedy	University
10-1	C. elongatus	IA	Mississippi	532.9			8/5/2004	J. Pitlo	IA DNR
10-2	C. elongatus	IA	Mississippi	543.4			8/5/2004	J. Pitlo	IA DNR
10-3	C. elongatus	IA	Mississippi	532.9			8/9/2004	D. Weiss	IA DNR
10-4	C. elongatus	IA	Mississippi	549.9			8/11/2004	D. Weiss	IA DNR
10-5	C. elongatus	IA	Mississippi	549.9			8/11/2004	D. Weiss	IA DNR
10-6	C. elongatus	IA	Mississippi	549.9			8/11/2004	D. Weiss	IA DNR
10-7	C. elongatus	IA	Mississippi	549.7			8/11/2004	D. Weiss	IA DNR
10-8	C. elongatus	IA	Mississippi	540.5			8/11/2004	D. Weiss	IA DNR
10-9	C. elongatus	IA	Mississippi	548			8/26/2004	D. Weiss	IA DNR

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
10-10	C. elongatus	IA	Mississippi	536.9			9/9/2004	D. Weiss	IA DNR
10-11	C. elongatus	IA	Mississippi	556.7			10/13/2004	J. Pitlo	IA DNR
12-9	C. elongatus	MN	Minnesota	217.1	44.57181	95.08796	8/12/2004	?	MN DNR
12-10	C. elongatus	MN	Minnesota	209.5	44.54417	95.00043	8/13/2004	?	MN DNR
12-11	C. elongatus	MN	Minnesota	111.9	44.16522	94.03383	8/17/2004	?	MN DNR
12-12	C. elongatus	MN	Minnesota	111.9	44.16522	94.03383	8/17/2004	?	MN DNR
12-13	C. elongatus	MN	Minnesota	74.5	44.47513	93.90654	8/21/2004	?	MN DNR
12-14	C. elongatus	MN	Minnesota	74.5	44.47513	93.90654	8/21/2004	?	MN DNR
12-15	C. elongatus	MN	Minnesota	74	44.53706	93.9009	9/9/2004	?	MN DNR
12-16	C. elongatus	MN	Minnesota	74	44.53706	93.90027	9/9/2004	?	MN DNR
13-2	C. elongatus	MN	Mississippi	761.4	?	?	7/15/2004	S. DeLain	MN DNR
13-3	C. elongatus	MN	Mississippi	795.6	44.60281	92.59176	7/27/2004	S. DeLain	MN DNR
13-4	C. elongatus	MN	Mississippi	795.6	44.60281	92.59176	7/27/2004	S. DeLain	MN DNR
13-5	C. elongatus	MN	Mississippi	795.6	44.60281	92.59176	7/27/2004	S. DeLain	MN DNR

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
13-6	C. elongatus	MN	Mississippi	795.6	44.60281	92.59176	7/27/2004	S. DeLain	MN DNR
13-7	C. elongatus	MN	Mississippi	795.6	44.60281	92.59176	7/27/2004	S. DeLain	MN DNR
13-8	C. elongatus	MN	Mississippi	795.6	44.60281	92.59176	7/27/2004	S. DeLain	MN DNR
17-1	C. elongatus	MO	Missouri	92.4	38.70524	91.34109	4/19/2004	Mauldin	USFWS
17-2	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
17-3	C. elongatus	MO	Missouri	213.8	38	90	4/12/2004	Mauldin	USFWS
17-4	C. elongatus	MO	Missouri	92.4	38.70524	91.34109	4/19/2004	Mauldin	USFWS
17-5	C. elongatus	MO	Missouri	92.3	38.69938	91.3388	4/19/2004	Mauldin	USFWS
17-6	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
17-7	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
17-8	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
17-9	C. elongatus	MO	Missouri	124	38.61108	91.92516	4/29/2004	W. Doyle	USFWS
17-10	C. elongatus	MO	Missouri	40.7	38.68257	90.61002	3/24/2004	A. Starostka	USFWS
17-11	C. elongatus	MO	Missouri	16	38.87243	90.34079	5/10/2004	W. Doyle	USFWS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
17-12	C. elongatus	МО	Missouri	40	38.68127	90.56851	3/22/2004	A. Starostka	USFWS
17-13	C. elongatus	MO	Missouri	128	38.61208	91.92825	4/29/2004	W. Doyle	USFWS
17-14	C. elongatus	MO	Missouri	0.2	38.80816	90.12071	5/11/2004	W. Doyle	USFWS
17-15	C. elongatus	MO	Missouri	42	38.68481	90.66957	4/21/2004	Mauldin	USFWS
17-16	C. elongatus	MO	Missouri	42	38.63297	90.63062	4/21/2004	Mauldin	USFWS
17-17	C. elongatus	MO	Missouri	92.4	38.70524	91.34109	4/19/2004	Mauldin	USFWS
17-18	C. elongatus	MO	Missouri	188.1	38.96808	92.59406	3/19/2004	W. Doyle	USFWS
17-19	C. elongatus	MO	Missouri	202	38.97929	92.85339	4/2/2004	A. Starostka	USFWS
17-20	C. elongatus	MO	Missouri	92.4	38.70524	91.34109	4/19/2004	Mauldin	USFWS
17-21	C. elongatus	MO	Missouri	0.2	38.80816	90.12071	5/11/2004	W. Doyle	USFWS
17-22	C. elongatus	MO	Missouri	128	38.61462	91.92518	4/29/2004	W. Doyle	USFWS
17-23	C. elongatus	MO	Missouri	92.3	38.69938	91.3388	4/19/2004	Mauldin	USFWS
17-24	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
17-25	C. elongatus	MO	Missouri	202	38.97929	92.85339	4/2/2004	A. Starostka	

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
17-26	C. elongatus	MO	Missouri	128	38.62086	91.91595	4/29/2004	W. Doyle	USF W 5
17-27	C. elongatus	MO	Missouri	202	38.97929	92.85339	4/2/2004	A. Starostka	USFWS
17-28	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
17-29	C. elongatus	MO	Missouri	92.4	38.70524	91.34109	4/19/2004	Mauldin	USFWS
17-30	C. elongatus	MO	Missouri	2	38.82946	90.14062	5/11/2004	W. Doyle	USFWS
18B-1	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-2	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-3	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-4	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-5	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-6	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-7	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-8	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP
18B-9	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	Montana FWP

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	-								Montana
18B-10	C. elongatus	MT	Missouri	2047.3	47.928	110.49	5/17/2004	R. Rodencal	FWP
									Montana
18B-11	C. elongatus	MT	Missouri	1980.8	47.7471	109.5815	5/20/2004	Wente/McCord	FWP
									Montana
18B-12	C. elongatus	MΤ	Missouri	1982.5	47.7374	109.6226	5/26/2004	Rodencal	FWP
100 12				1000 5	10 0001	100 (00)	5 10 ( 10 00 4	D 1 1	Montana
18B-13	C. elongatus	MI	Missouri	1982.5	47.7374	109.6226	5/26/2004	Rodencal	FWP
10D 14		МТ	Missouri	10075	47 7102	100 606	5/26/2004	Dadamaal	Montana
10D-14	C. elongatus	IVI I	IVIISSOULI	1987.3	47.7192	109.090	3/20/2004	Rodelical	r wr Montana
18B-15	C alongatus	МТ	Missouri	1082.8	17 7371	100 6183	6/29/2004	Rodencal	FW/P
10D-15	C. elongulus	1111	Ivii550ui1	1762.0	47.7571	107.0105	0/2//2004	Rouenear	Montana
18B-16	C elongatus	MT	Missouri	20304	48.03	110 221	6/24/2004	Wente	FWP
100 10	C. elongulus	1,111	111550411	2030.1	10.05	110.221	0/21/2001	vv ente	Montana
18B-17	C. elongatus	MT	Missouri	2030.4	48.03	110.221	6/24/2004	Wente	FWP
	e								Montana
18B-18	C. elongatus	MT	Missouri	1983.6	47.7395	109.6306	6/29/2004	Rodencal	FWP
	0								Montana
18B-19	C. elongatus	MT	Missouri	2068.9	47.50489	110.3543	7/15/2004	McCord	FWP
									Montana
18B-20	C. elongatus	MT	Missouri	1909	47.589	108.472	7/9/2004	Gerrity	FWP
									Montana
18B-21	C. elongatus	MT	Missouri	1884.8	47.5872	108.1302	8/3/2004	Rodencal	FWP
									Montana
18B-22	C. elongatus	MT	Missouri	1883.3	47.7331	109.6034	9/8/2004	Rodencal	FWP
100.00				1002.2	15 5000	100 (00)	0.10.10.00.		Montana
18 <b>B-</b> 23	C. elongatus	MI	Missouri	1883.3	47.7333	109.6034	9/8/2004	Kodencal	гwр

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	1								Montana
18B-24	C. elongatus	MT	Missouri	2032.8	48.0004	110.2539	9/9/2004	Rodencal	FWP
	0								Montana
18B-25	C. elongatus	MT	Missouri	2034.1	48.0176	110.2837	9/9/2004	Rodencal	FWP
	0								Montana
18B-26	C. elongatus	MT	Missouri	2029.5	48.03328	110.2062	9/9/2004	Rodencal	FWP
									Montana
18B-27	C. elongatus	MT	Missouri	2034.8	48.00864	110.266	9/9/2004	Rodencal	FWP
									Montana
18B-28	C. elongatus	MT	Missouri	2034.8	48.00864	110.266	9/9/2004	Rodencal	FWP
									Montana
18B-29	C. elongatus	MT	Missouri	2034.8	48.00864	110.266	9/9/2004	Rodencal	FWP
									Montana
18B-30	C. elongatus	MT	Missouri	2031.4	48.0007	110.2479	9/10/2004	Rodencal	FWP
									Montana
19-1	C. elongatus	MT	Missouri	1763.5	48.0269	106.2149	7/29/2004	Dix	FWP
									Montana
19-2	C. elongatus	MT	Missouri	?	?	?		Dix	FWP
									Montana
19-3	C. elongatus	MT	Missouri	?	?	?		Dix	FWP
									Montana
19B-3	C. elongatus	MT	Missouri		48.07592	105.653	10/4/2004	Baxter	FWP
									Montana
19-4	C. elongatus	MT	Missouri	?	?	?		Dix	FWP
									Montana
19-5	C. elongatus	MT	Missouri	1672	48.04629	105.0741	10/27/2004	Dix	FWP
									Montana
19-6	C. elongatus	MT	Missouri	1711.5	48.029	105.3865	11/5/2004	Dix	FWP

Individual	Species	State	River	River	Latitude	Longitude	Date of collection	Collector	Institution or
marviauai	species	State	River	mme	(11)	$(\mathbf{w})$	concetion	Concetor	Montana
19B-6	C. elongatus	MT	Missouri		48.10859	104.5912	10/5/2004	Baxter	FWP
									Montana
19-7	C. elongatus	MT	Missouri	1702	48.04198	105.3303	10/19/2004	Dix	FWP
									Montana
19B-12	C. elongatus	MT	Missouri		48.07592	105.653	10/4/2004	Baxter	FWP
									Montana
19B-14	C. elongatus	MT	Missouri		48.06271	106.3857	9/28/2004	Baxter	FWP
100 15					40.07500	105 (52	10/4/0004		Montana
19B-15	C. elongatus	MI	Missouri		48.07592	105.653	10/4/2004	Baxter	FWP
10 16	C 1	МТ	Missouri	1761	19 02 121	106 0172	4/27/2004	Doutor	Montana EWD
19-10	C. elongatus	IVI I	Iviissouri	1/01	48.03421	100.9172	4/2//2004	Daxtel	Г W P Montana
19-17	C elongatus	МТ	Missouri	1761	48 03421	106 9172	4/27/2004	Baxter	FWP
17 17	C. elongulus	1011	111550 di l	1701	10.05 121	100.9172	1/2//2001	Duriel	Montana
19-18	C. elongatus	MT	Missouri	1761	48.02703	106.172	4/27/2004	Baxter	FWP
									Montana
19-19	C. elongatus	MT	Missouri	1684	48.05296	105.1366	6/8/2004	Baxter	FWP
	0								Montana
19-20	C. elongatus	MT	Missouri	1684	48.04778	105.1317	6/8/2004	Baxter	FWP
									Montana
19-21	C. elongatus	MT	Missouri	1763.5	48.02877	106.2199	6/9/2004	Baxter	FWP
								_	Montana
19 <b>B-</b> 21	C. elongatus	MT	Missouri		48.07592	105.653	10/4/2004	Baxter	FWP
10.00	~ .			1770	40.04170	106 00 41	C 110 1000 A		Montana
19-22	C. elongatus	MI	Missouri	1//0	48.041/2	106.2341	6/10/2004	Baxter	FWP
10.22		МТ	Missouri	1770	49 00021	106 2592	6/10/2004	Doutor	Nontana
17-23	C. elongatus	111	IVIISSOUIT	1//0	40.00921	100.2363	0/10/2004	Daxiel	L' VV L

	c ·	<u>G</u> ( )	D.	River	Latitude	Longitude	Date of		Institution or
Individual	Species	State	River	mile	(N)	(W)	collection	Collector	agency Montana
19B-23	C. elongatus	MT	Missouri		48.07592	105.653	10/4/2004	Baxter	FWP
									Montana
19-24	C. elongatus	MT	Missouri	1770	48.00921	106.2583	6/10/2004	Baxter	FWP
									Montana
19-25	C. elongatus	MT	Missouri	1760	48.02499	106.1813	6/15/2004	Dix	FWP
								_	Montana
19B-25	C. elongatus	MT	Missouri		48.10859	104.5912	10/5/2004	Baxter	FWP
10.20	~ ·	МТ	NC ·	17(0	40 0 2 2 4 1	106 1700	(115/2004	D.	Montana
19-26	C. elongatus	MI	Missouri	1/60	48.02341	106.1/88	6/15/2004	Dix	FWP
10.27	Calanastus	МТ	Missouri	1760	48 02341	106 1788	6/15/2004	Div	FW/D
19-21	C. elongalus	191 1	Ivi1550u11	1700	40.02341	100.1788	0/13/2004	DIX	Montana
19-28	C elongatus	MT	Missouri	1760	48 02484	106 178	6/15/2004	Dix	FWP
17 20	C. cionguius		111050 411	1,00	10.02101	100.170	0,10,2001	Dim	Montana
19-29	C. elongatus	MT	Missouri	1648.5	48.08899	104.5404	6/23/2004	Dix	FWP
	8								Montana
19-30	C. elongatus	MT	Missouri	1648.5	48.08949	104.5287	6/23/2004	Dix	FWP
									Montana
19B-30	C. elongatus	MT	Missouri		48.02766	106.2356	9/28/2004	Baxter	FWP
									Montana
20-1	C. elongatus	MT	Yellowstone	87.6	47.15451	104.6687	6/23/2004	M. Backes	FWP
<b>2</b> 0D 1			X 7 11	(0)	47 20701	104 4622	0/0/0004		Montana
20B-1	C. elongatus	MT	Yellowstone	68	47.30701	104.4632	9/2/2004	M. Backes	FWP
20.2		МТ	Vallaviatora	1070	16 20105	105 014	7/27/2004	M. Dealrea	Montana
20-2	C. elongatus	1111	renowstone	10/.0	40.38493	103.914	//2//2004	IVI. Dackes	Г W P Montono
20-3	C alongatur	MT	Vellowstone	187 8	46 38495	105 914	7/27/2004	M Backer	FWP
20-5	C. elonguius	141 1	1 CHOWStolle	107.0	TU.JUTJJ	105.714	112112004	IVI. DUCKES	T AA T

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	1								Montana
20-4	C. elongatus	MT	Yellowstone	189	46.3778	105.9269	7/27/2004	M. Backes	FWP
									Montana
20-5	C. elongatus	MT	Yellowstone	189	46.3778	105.9269	7/27/2004	M. Backes	FWP
									Montana
20-6	C. elongatus	MT	Yellowstone	186.4	46.39555	105.8973	7/27/2004	M. Backes	FWP
									Montana
20-7	C. elongatus	MT	Yellowstone	184.5	46.41703	105.8639	7/27/2004	M. Backes	FWP
• • •			× 11	0 - 6		101 ( ( ) <b>-</b>	0 / <b>/ / 0</b> 0 0 <b>/</b>		Montana
20-8	C. elongatus	MT	Yellowstone	87.6	47.15451	104.6687	8/4/2004	M. Backes	FWP
20.0	~ .		X7 11 /	07 (	47 15451	104 ((07	0/4/2004		Montana
20-9	C. elongatus	MI	Yellowstone	87.6	47.15451	104.6687	8/4/2004	M. Backes	FWP
20.10		МТ	Vallaviatora	07 (	17 15151	104 6697	0/4/2004	M. Dealwar	Montana
20-10	C. elongatus	IVI I	renowstone	87.0	47.13431	104.008/	8/4/2004	M. Backes	FWP
20.11	C 1	МТ	Vallowstona	976	17 15151	104 6697	8/1/2001	M Doolson	EWD
20-11	C. elongatus	1111	renowstone	87.0	47.13431	104.008/	0/4/2004	WI. Dackes	r wr Montono
20-12	C alongatus	MT	Vellowstone	87.6	17 15451	104 6687	8/4/2004	M Backes	FW/P
20-12	C. elongulus	191 1	Tenowstone	07.0	47.15451	104.0007	0/4/2004	WI. Dackes	Montana
20-13	C elongatus	MT	Yellowstone	87.6	47 15451	104 6687	8/4/2004	M Backes	FWP
20 15	e. ciongunas			01.0	1,110 101	101.0007	0, 1, 2001		Montana
20-14	C elongatus	MT	Yellowstone	71			8/5/2004	M. Backes	FWP
-									Montana
20-15	C. elongatus	MT	Yellowstone	71			8/5/2004	M. Backes	FWP
	8								Montana
20-16	C. elongatus	MT	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP
	0								Montana
20-17	C. elongatus	MT	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	1								Montana
20-18	C. elongatus	MT	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP
									Montana
20-19	C. elongatus	MT	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP
									Montana
20-20	C. elongatus	MT	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP
									Montana
20-21	C. elongatus	MT	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP
									Montana
20-22	C. elongatus	MΤ	Yellowstone	182	46.43355	105.8204	8/10/2004	M. Backes	FWP
20.22			X7 11	007	46.07500	106 (704	0/10/2004		Montana
20-23	C. elongatus	MI	Yellowstone	237	46.27588	106.6794	8/12/2004	M. Backes	FWP
20.24		МТ	Vallavyatana	227	16 27500	106 6704	9/12/2004	M. Dealwar	Montana
20-24	C. elongatus	IVI I	renowstone	237	40.2/388	100.0794	8/12/2004	M. Backes	FWP
20.25	Calanantus	МТ	Vallowstona	197	16 10655	105 8701	8/16/2004	M Baalzas	IVIOIItalia EW/D
20-23	C. elongatus	101 1	I enowstone	10/	40.40035	103.8701	8/10/2004	IVI. Dackes	r wr Montana
20-26	C alongatus	MT	Vellowstone	184	46 42953	105 8423	8/16/2004	M Backes	FWP
20-20	C. elongulus	141 1	I entowstone	104	40.42755	105.0425	0/10/2004	WI. Duckes	Montana
20-27	C elongatus	MT	Yellowstone	184	46 42953	105 8423	8/16/2004	M Backes	FWP
_0 _/	e. cionganas			101		10010.20	0,10,200		Montana
20-28	C. elongatus	MT	Yellowstone	184	46.42953	105.8423	8/16/2004	M. Backes	FWP
									Montana
20-29	C. elongatus	MT	Yellowstone	184	46.42953	105.8423	8/16/2004	M. Backes	FWP
	0								Montana
20-30	C. elongatus	MT	Yellowstone	289.9	46.221	107.4206	8/27/2004	M. Backes	FWP
22.1							7/14/2004	DI	NUCE
22-1	Cycleptus sp.	NM	Black				//14/2004	B. Larson	NMGF

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	I I I I I I I I I I I I I I I I I I I			-					Albuquerque
22-31	Cycleptus sp.	NM	Pecos				7/14/2004	B. Dorn	Aqm.
									Albuquerque
22-32	Cycleptus sp.	NM	Pecos				7/14/2004	B. Dorn	Aqm.
									Albuquerque
22-33	Cycleptus sp.	NM	Pecos				7/14/2004	B. Dorn	Aqm.
									Albuquerque
22-34	Cycleptus sp.	NM	Black				7/14/2004	B. Dorn	Aqm.
							_ / /	,,,,	North Dakota
23-1	C. elongatus	ND	Missouri	1299			7/12/2004	J. Hendrickson	G&F
<u></u>				1200			7/12/2004	T TT 1 · 1	North Dakota
23-2	C. elongatus	ND	Missouri	1299			7/12/2004	J. Hendrickson	G&F
<b>22</b> 2	~ .		Misser	1200			7/12/2004	T TT 4-: -1	North Dakota
23-3	C. elongatus	ND	Missouri	1299			//12/2004	J. Hendrickson	G&F North Dalasta
22.4		ND	Missouri	1220			0/7/2004	I. Handriakaan	North Dakota
23-4	C. elongatus	ND	IVIISSOUIT	1320			9/7/2004	J. HENdrickson	Var North Daleata
23.5	Calanastus	ND	Missouri	1320			9/7/2004	I Handrickson	G&F
25-5	C. elongalus	ND	Iviissouii	1320			9/1/2004	J. HEHMHEKSON	North Dakota
23-6	C elongatus	ND	Missouri	1320			9/7/2004	I Hendrickson	G&F
25 0	C. cionguius	T(D)	111000 di l	1520			<i>JTT</i> <b>2</b> 001	5. Hendrickbon	North Dakota
23-7	C elongatus	ND	Missouri	1320			9/7/2004	J. Hendrickson	G&F
/	er erenguns								North Dakota
23-8	C. elongatus	ND	Missouri	1320			9/7/2004	J. Hendrickson	G&F
	8								North Dakota
23-9	C. elongatus	ND	Missouri	1320			9/7/2004	J. Hendrickson	G&F
	2								North Dakota
23-10	C. elongatus	ND	Missouri	1320			9/7/2004	J. Hendrickson	G&F

Ter dissi dasa 1	Spacing	State	Divor	River	Latitude	Longitude	Date of	Collector	Institution or
Individual	species	State	Kivei	mine	$(1\mathbf{N})$	( •• )	concetion	Conector	North Dakota
23-11	C. elongatus	ND	Missouri	1343			9/8/2004	J. Hendrickson	G&F
									North Dakota
23-12	C. elongatus	ND	Missouri	1371			9/9/2004	J. Hendrickson	G&F
									North Dakota
23-13	C. elongatus	ND	Missouri	1371			9/9/2004	J. Hendrickson	G&F
									North Dakota
23-14	C. elongatus	ND	Missouri	1371			9/9/2004	J. Hendrickson	G&F
			<u>.</u>	<b>a</b>					Kentucky
24-1	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
24.2	<i>.</i> .		01.	(07	20 20565	05 70102	0/07/2004		Kentucky
24-2	C. elongatus	ΚY	Onio	607	38.28565	85./9193	8/2//2004	Crosby, Duvall	DFWR Kantaalaa
24.2		VV	Ohio	607	20 20565	85 70103	8/27/2004	Crashy Duvall	DEWD
24-3	C. elongatus	K I	Ollio	007	36.26303	03./9195	8/27/2004	Closby, Duvali	Dr wr Kontuolau
24_4	C alongatus	KV	Ohio	607	38 28565	85 70103	8/27/2004	Crosby Duvall	DEWR
24-4	C. elongalus	K I	Onio	007	38.28303	05.79195	0/2//2004	Closby, Duvali	Kentucky
24-5	C alongatus	KV	Ohio	607	38 28565	85 79193	8/27/2004	Crosby Duvall	DFWR
27 3	C. elongulus	K I	Onio	007	50.20505	05.77175	0/2//2004	Closey, Duvan	Kentucky
24-6	C elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
	er erenguns								Kentucky
24-7	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
	and a second second							57	Kentucky
24-8	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
	C							•	Kentucky
24-9	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
									Kentucky
24-10	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR

	а ·	<u>G</u> , ,	р.	River	Latitude	Longitude	Date of		Institution or
Individual	Species	State	River	mile	(N)	(W)	collection	Collector	agency
24-11	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
	and a second second							57	Kentucky
24-12	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
									Kentucky
24-13	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
04.14	~ .	1/1/	01.	(0 <b>7</b>	20.00565	05 70102	0/07/0004		Kentucky
24-14	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
24.15	C clanastus	KV	Ohio	607	38 28565	85 70103	8/27/2004	Crosby Duvall	DEWR
24-13	C. elongalus	K I	Ollio	007	38.28303	05.79195	8/2//2004	Closby, Duvali	Kentucky
24-16	C elongatus	KY	Ohio	607	38 28565	85 79193	8/27/2004	Crosby Duvall	DFWR
2.10	e. cionguius		Olife	007	20.20000	00.19190	0/2//2001	crossy, Duvun	Kentucky
24-17	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
	C							•	Kentucky
24-18	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
									Kentucky
24-19	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
24.20	~ .	1/1/	01.	(0 <b>7</b>	20.00565	05 70102	0/07/0004		Kentucky
24-20	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
24 21	C alamantur	VV	Ohio	607	28 28565	85 70102	8/27/2004	Crosby Duvall	DEWP
24-21	C. elongatus	ΚI	Ollio	007	56.26505	03./9195	8/2//2004	Closby, Duvali	DF WK Kentucky
24-22	C elongatus	KY	Ohio	607	38 28565	85 79193	8/27/2004	Crosby Duvall	DFWR
2122	C. cionguius	111	Olife	007	50.20505	00.77175	0/2//2001	erosoy, Duvun	Kentucky
24-23	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
	and a second second							5,	Kentucky
24-24	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
24-25	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	DFWR
24-26	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	Kentucky DFWR
24-27	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	Kentucky DFWR
24-28	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	Kentucky DFWR
24-29	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	Kentucky DFWR
24-30	C. elongatus	KY	Ohio	607	38.28565	85.79193	8/27/2004	Crosby, Duvall	Kentucky DFWR
25-1	C. elongatus	SD	Missouri		42.77984	98.03719	11/1/2004	D. Schuman	USFWS
25-2	C. elongatus	SD	Missouri		42.77386	97.96674	7/20/2004	D. Schuman	USFWS
25-3	C. elongatus	SD	Missouri		42.76314	98.02292	3/16/2005	D. Schuman	USFWS
25-4	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-5	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-6	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-7	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-8	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
25-9	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-10	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-11	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-12	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-13	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-14	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-15	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-16	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-18	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-19	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-20	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-21	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-22	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-23	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
25-24	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-25	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-26	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-27	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-28	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-29	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
25-30	C. elongatus	SD	Missouri		42.76471	97.98806	5/4/2005	Wanner	USFWS
28-1	C. elongatus	TN	Cumberland				3/30/2004	T. St. John	TWRA
28-2	C. elongatus	TN	Tennessee				4/21/2004	P. Bettoli	Univ.
29-1	C. elongatus	TN	Hatchie		35.51197	89.33594	5/24/2005	M. Clark	TWRA
29-2	C. elongatus	TN	Hatchie		35.51317	89.32812	5/24/2005	M. Clark	TWRA
29-3	C. elongatus	TN	Hatchie		35.51504	89.32738	5/24/2005	M. Clark	TWRA
29-4	C. elongatus	TN	Hatchie		35.51378	89.32431	5/24/2005	M. Clark	TWRA
29-5	C. elongatus	TN	Hatchie		35.51173	89.32018	5/24/2005	M. Clark	TWRA

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
29-16	C. elongatus	TN	Hatchie		35.38293	89.15309	5/24/2005	D. Barber	TWRA
29-17	C. elongatus	TN	Hatchie		35.3119	89.1539	5/24/2005	D. Barber	TWRA
29-18	C. elongatus	TN	Hatchie		35.31507	89.15386	5/24/2005	D. Barber	TWRA
29-19	C. elongatus	TN	Hatchie		35.3177	89.15934	5/24/2005	D. Barber	TWRA
29-20	C. elongatus	TN	Hatchie		35.3177	89.15962	5/24/2005	D. Barber	TWRA
29-21	C. elongatus	TN	Hatchie		35.31609	89.15971	5/24/2005	D. Barber	IWKA
29-22	C. elongatus	TN	Hatchie		35.31526	89.15976	5/24/2005	D. Barber	TWRA
29-23	C. elongatus	TN	Hatchie		35.3166	89.16089	5/24/2005	D. Barber	TWRA
29-24	C. elongatus	TN	Hatchie		35.31658	89.161	5/24/2005	D. Barber	TWRA
29-25	C. elongatus	TN	Hatchie		35.31882	89.16582	5/24/2005	D. Barber	TWRA
29-26	C. elongatus	TN	Hatchie		35.36226	89.50298	5/24/2005	D. Barber	TWRA
29-27	C. elongatus	TN	Hatchie		35.36263	89.50322	5/24/2005	D. Barber	TWRA
30-1	C. elongatus	TN	Nolichucky	44.7	36.03558	82.53038	5/7/2004	R. Bivins	TWRA
30-2	C. elongatus	TN	Nolichucky	44.7	36.03558	82.53038	5/7/2004	R. Bivins	TWRA

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
30-3	C. elongatus	TN	Nolichucky	44.15	36.03363	82.5324	5/7/2004	R. Bivins	TWRA
30-4	C. elongatus	TN	Nolichucky	43.7	36.03212	82.53274	5/7/2004	R. Bivins	TWRA
30-5	C. elongatus	TN	Nolichucky	43.7	36.03212	82.53274	5/7/2004	R. Bivins	TWRA
30-6	C. elongatus	TN	Nolichucky	41.9	36.03488	82.543	5/7/2004	R. Bivins	TWRA
30-7	C. elongatus	TN	Nolichucky	35.4	36.0402	82.58315	5/7/2004	R. Bivins	IWKA
30-8	C. elongatus	TN	Nolichucky	30.9	36.05336	83.00345	5/11/2004	R. Bivins	TWRA
30-9	C. elongatus	TN	French Broad	13.2	35.58021	83.43259	5/18/2004	B. Carter	TWRA
33-1	C. elongatus	WI	Wisconsin	30.5	43.1748	90.6877	7/1/2004	J. Lyons	DNR
33-2	C. elongatus	WI	Wisconsin	30.5	43.1748	90.6877	7/1/2004	J. Lyons	DNR
33-3	C. elongatus	WI	Wisconsin	30.5	43.1748	90.6877	7/1/2004	J. Lyons	DNR
33-4	C. elongatus	WI	Wisconsin	30.5	43.1748	90.6877	7/1/2004	J. Lyons	DNR
33-5	C. elongatus	WI	Wisconsin	30	43.1702	90.6928	7/1/2004	J. Lyons	Wisconsin DNR
33-6	C. elongatus	WI	Wisconsin	30	43.1702	90.6928	7/1/2004	J. Lyons	Wisconsin DNR
33-7	C. elongatus	WI	Wisconsin	30	43.1702	90.6928	7/1/2004	J. Lyons	W1scons1n DNR

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
33-8	C elongatus	WI	Wisconsin	30	43 1702	90 6928	7/1/2004	I Lyons	Wisconsin DNR
55 0	e. ciongunas			20	13.1702	<i>y</i> 0.0 <i>y</i> <b>2</b> 0	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	0. 290115	Wisconsin
33-9	C. elongatus	WI	Wisconsin	26	43.1297	90.7478	7/29/2004	J. Lyons	DNR
	0							2	Wisconsin
33-10	C. elongatus	WI	Wisconsin	26	43.1297	90.7478	7/29/2004	J. Lyons	DNR
	-							-	Wisconsin
33-11	C. elongatus	WI	Wisconsin	26	43.1297	90.7478	7/29/2004	J. Lyons	DNR
									Wisconsin
33-12	C. elongatus	WI	Wisconsin	26	43.1297	90.7478	7/29/2004	J. Lyons	DNR
									Wisconsin
33-13	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-14	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-15	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-16	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-17	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-18	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-19	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
									Wisconsin
33-20	C. elongatus	WI	Wisconsin	25.6	43.1231	90.7511	7/29/2004	J. Lyons	DNR
							_ / /		Wisconsin
33-21	C. elongatus	WI	Wisconsin	20.5	43.0909	90.8305	7/29/2004	J. Lyons	DNR

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
				• • •	10 0000				Wisconsin
33-22	C. elongatus	WI	Wisconsin	20.5	43.0909	90.8305	7/29/2004	J. Lyons	DNR
33-23	C elongatus	WI	Wisconsin	20.3	43 085	90 8311	7/29/2004	J Lyons	W1sconsin DNR
00 20	e. ciongunas			2010		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,,_,,_,,_,,	0.29010	Wisconsin
33-24	C. elongatus	WI	Wisconsin	20.3	43.085	90.8311	7/29/2004	J. Lyons	DNR
									Wisconsin
33-25	C. elongatus	WI	Wisconsin	20.3	43.085	90.8311	7/29/2004	J. Lyons	DNR
									Wisconsin
33-26	C. elongatus	WI	Wisconsin	20.3	43.085	90.8311	7/29/2004	J. Lyons	DNR
									Wisconsin
33-27	C. elongatus	WI	Wisconsin	20.3	43.085	90.8311	7/29/2004	J. Lyons	DNR
									Wisconsin
33-28	C. elongatus	WI	Wisconsin	20.3	43.085	90.8311	7/29/2004	J. Lyons	DNR
				• • •	1 <b>2</b> 00 <b>5</b>	00.0011			Wisconsin
33-29	C. elongatus	WI	Wisconsin	20.3	43.085	90.8311	7/29/2004	J. Lyons	DNR
22.20			***	10.1	42 2015	00 4451	0/1/0001	<b>.</b> .	Wisconsin
33-30	C. elongatus	WI	Wisconsin	43.1	43.2017	90.4471	9/1/2004	J. Lyons	DNR
24.2	~ ·	VO	17	140			10/12/2004		KS State
34-3	C. elongatus	KS	Kansas	140			10/12/2004	C. Paukert	Univ.
24.6		VO	V	140			11/4/2004	C Development	KS State
34-6	C. elongatus	KS	Kansas	140			11/4/2004	C. Paukert	Univ.
24 12		VS	Vancas	140			11/4/2004	C Doultort	KS State
34-12	C. elongatus	КS	Nalisas	140			11/4/2004	C. Pauken	UIIIV. VS Stata
24.20	C alamantur	VS	Kongog	140			10/12/2004	C Paukart	KS State
54-20	C. elongalus	KS	Kalisas	140			10/12/2004	C. I auxort	UIIIV. KS State
34-24	C elongatus	KS	Kansas	140			11/4/2004	C. Paukert	Univ.
	c. crongans								

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
34-26	C. elongatus	KS	Kansas	140			10/12/2004	C. Paukert	Univ.
35-1	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-2	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-3	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-4	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-5	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-6	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-7	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-8	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-9	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	031 w 3
35-10	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-11	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-12	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-13	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
35-14	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-15	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-16	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-17	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-18	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-19	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-20	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-21	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-22	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-23	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-24	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-25	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-26	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-27	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
35-28	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-29	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-30	C. elongatus	OK	Red	725	33.49	96.33	3/16/2005	B. Bristow	USFWS
35-31	C. elongatus	OK	Red	123	33.5742.1	95.13.51.1	11/17/2005	B. Bristow	USFWS
35-32	C. elongatus	OK	Red	123	33.5742.1	95.13.51.1	11/17/2005	B. Bristow	USFWS
35-33	C. elongatus	OK	Red	123	33.5742.1	95.13.51.1	11/17/2005	B. Bristow	USFWS
36-01	Cycleptus sp.	ТХ	Rio Grande	Big Bend NP				T. Bonner	Marcos
36-02	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-03	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-04	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-05	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-06	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-07	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-08	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	150 - San Marcos

									TSU – San
36-09	Cycleptus sp.	ТХ	Rio Grande					T. Bonner	Marcos
36-10	Cycleptus sp.	TX	Rio Grande					T. Bonner	Marcos
									TSU – San
36-11	Cycleptus sp.	ΤX	Rio Grande					T. Bonner	Marcos
2(12		$\mathbf{T}\mathbf{V}$						т р	TSU – San
30-12	Cycleptus sp.	IX	Klo Grande					I. Bonner	Marcos
36-13	Cuelentus en	тх	Rio Grande					T Bonner	150 – Sall Marcos
50-15	Cyclepius sp.	17	Kio Grande					T. Donner	TSU – San
36-14	Cvcleptus sp.	ΤX	Rio Grande					T. Bonner	Marcos
									TSU – San
36-15	Cycleptus sp.	ΤX	Rio Grande					T. Bonner	Marcos
									TSU – San
36-16	Cycleptus sp.	ΤX	Rio Grande					T. Bonner	Marcos
26.17	~ .	τv						TD	TSU – San
36-17	Cycleptus sp.	IX	Rio Grande					I. Bonner	Marcos
36.18	Cuelentus en	$\mathbf{T}\mathbf{Y}$	Rio Granda					T Bonner	150 – San Marcos
50-18	Cyclepius sp.	IЛ	Kio Granue					I. Donner	TSU – San
36-19	Cvclentus sn	ТХ	Rio Grande					T. Bonner	Marcos
••••	eyenepins spi								TSU – San
36-20	Cycleptus sp.	ΤX	Rio Grande					T. Bonner	Marcos
									Wisconsin
37-1	C. elongatus	WI	Mississippi	672			6/2/2004	P. Short	DNR
									Wisconsin
37-2	C. elongatus	WI	Mississippi	752			8/5/2004	K. VanRuden	DNR
20.1		W/I	Chinnawa		11 52010	02 04212	8/4/2004	D Uniile	W1SCONSIN
37-1	C. elongatus	VV I	Chippewa		44.33948	92.04212	0/4/2004	. пијік	DINK

Individual	Species	State	River	River mile	Latitude	Longitude	Date of collection	Collector	Institution or
marviauai	species	State	River	mite	(11)	(")	concetion	Concetor	Wisconsin
39-2	C. elongatus	WI	Chippewa		44.52868	92.0481	8/4/2004	B. Hujik	DNR
			11					5	Wisconsin
39-3	C. elongatus	WI	Chippewa		44.32867	92.04809	8/4/2004	D. Johnson	DNR
	-								Wisconsin
39-4	C. elongatus	WI	Chippewa		44.50321	92.05177	8/4/2004	D. Johnson	DNR
									Wisconsin
39-5	C. elongatus	WI	Chippewa		44.50321	92.05177	8/4/2004	D. Johnson	DNR
									Wisconsin
39-6	C. elongatus	WI	Chippewa		44.49423	92.05341	8/4/2004	D. Johnson	DNR
							0 / <b>/ / 0</b> 0 0 <b>/</b>		Wisconsin
39-7	C. elongatus	WI	Chippewa		44.49423	92.05341	8/4/2004	A. Lamm	DNR
20.0	~ .	11/1	C1 ·		44 40 422	02 052 41	0/4/2004	A T	W1sconsin
39-8	C. elongatus	WI	Chippewa		44.49423	92.05341	8/4/2004	A. Lamm	DNK
20.0		WI	Chinneyye		44 40422	02 05241	Q/4/2004	N. Schoff	W ISCONSIN
39-9	C. elongatus	W I	Chippewa		44.49423	92.03341	8/4/2004	IN. SCHall	DINK
30.10	Colongatur	WI	Chinnewa		11 18508	02 05705	8/4/2004	N Schaff	DNP
39-10	C. elongalus	VV 1	Chippewa		44.40390	92.03703	8/4/2004	IN. Schaff	Wisconsin
39-11	C elongatus	WI	Chinnewa		44 48598	92 05705	8/4/2004	N Schaff	DNR
57 11	C. cionguius		emppewa		11.10270	2.00700	0/ 1/2001		Wisconsin
39-12	C elongatus	WI	Chippewa		44,48598	92.05705	8/4/2004	N. Schaff	DNR
• • • • •	er erenguns					/			Wisconsin
39-13	C. elongatus	WI	Chippewa		44.48598	92.05705	8/4/2004	N. Schaff	DNR
	8		11						Wisconsin
39-14	C. elongatus	WI	Chippewa		44.48598	92.05705	8/4/2004	N. Schaff	DNR
	0								Wisconsin
39-15	C. elongatus	WI	Chippewa		44.48598	92.05705	8/4/2004	N. Schaff	DNR

				River	Latitude	Longitude	Date of		Institution or
Individual	Species	State	River	mile	(N)	(W)	collection	Collector	agency
20.16	~ .	<b>XX</b> /T	C1 ·		44 40500	02 05705	0/4/2004		Wisconsin
39-16	C. elongatus	WI	Chippewa		44.48598	92.05/05	8/4/2004	N. Schaff	
20.17		W/I	Chinneyye		11 16062	02 06122	Q/4/2004	N. Schaff	W ISCONSIN
39-17	C. elongatus	VV I	Chippewa		44.40902	92.00132	8/4/2004	IN. Schaff	DINK
30.18	Calanastus	WI	Chinnews		11 16062	02 06132	8/4/2004	N Schaff	DNP
39-10	C. elongalus	VV I	Cilippewa		44.40902	92.00132	0/4/2004	IN. Schaff	Wisconsin
39-19	C alongatus	WI	Chinnewa		44 46962	92 06132	8/4/2004	N Schaff	DNR
57 17	C. elongulus	** 1	emppewa		11.10902	12.00152	0/ 1/2001	iv. Senair	Wisconsin
39-20	C elongatus	WI	Chippewa		44,46962	92.06132	8/4/2004	N. Schaff	DNR
• / - •	er erenguns					/			Wisconsin
39-21	C. elongatus	WI	Chippewa		44.46962	92.06132	8/4/2004	N. Schaff	DNR
	0		11						Wisconsin
39-22	C. elongatus	WI	Chippewa		44.45267	92.06818	8/4/2004	N. Schaff	DNR
									Wisconsin
39-23	C. elongatus	WI	Chippewa		44.45267	92.06818	8/4/2004	N. Schaff	DNR
									Wisconsin
39-24	C. elongatus	WI	Chippewa		44.45267	92.06818	8/4/2004	N. Schaff	DNR
									Wisconsin
39-25	C. elongatus	WI	Chippewa		44.45267	92.06818	8/4/2004	N. Schaff	DNR
		** **					0 / <b>/ / 0</b> 0 0 <b>/</b>		Wisconsin
39-26	C. elongatus	WI	Chippewa		44.45267	92.06818	8/4/2004	N. Schaff	DNR
20.27	~ .	<b>XX</b> /T	C1 ·		44 45067	02 0/010	0/4/2004		Wisconsin
39-27	C. elongatus	WI	Chippewa		44.45267	92.06818	8/4/2004	N. Schaff	DNK
20.20		W/I	Chinnawa		11 12722	02 07104	8/1/2001	N. Schaff	W ISCONSIN
39-28	C. elongatus	VV I	Chippewa		44.43732	92.07194	8/4/2004	IN. Schaff	DINK
30_20	Calamanter	WI	Red Cedar		11 78031	01 0/038	8/17/2004	D. Johnson	DNR
57-47	C. eiongaius	VV 1	Reu Ceudi			71.74030	0/1//2004	D. Johnson	

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	1								Wisconsin
39-30	C. elongatus	WI	Red Cedar		44.76468	91.93135	8/17/2004	D. Johnson	DNR
									Lousiana
40-1	C. elongatus	LA	Mississippi	315	31.0351	91.3537	3/19/2004	B. Reed	DWF
									Lousiana
40-2	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/29/2004	B. Reed	DWF
									Lousiana
40-3	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-4	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-5	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-6	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-7	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-8	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-9	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-10	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-11	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-12	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
	-								Lousiana
40-13	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
		<b>T</b> 1	. <i>.</i>						Lousiana
40-14	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
40-15	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
40-16	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-17	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-18	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-19	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-20	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-21	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-22	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-23	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-24	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-25	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-26	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF
40-27	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	Lousiana DWF

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	1								Lousiana
40-28	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-29	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Lousiana
40-30	C. elongatus	LA	Mississippi	315	31.0351	91.3537	4/19/2005	B. Reed	DWF
									Univ. of
44-1	C. elongatus	NE	Platte		41.06734	96.05474	5/14/2004	E. Peters	Nebraska
			51						Univ. of
44-3	C. elongatus	NE	Platte		41.05132	96.10832	5/14/2004	E. Peters	Nebraska
445	~ .	NIE	D1 44		41.0(72.4	06 05 474	5/14/2004		Univ. of
44-5	C. elongatus	NE	Platte		41.06/34	96.054/4	5/14/2004	E. Peters	Nebraska
116	C 1	NE	Dlatta		41.05122	06 10922	5/14/2004	E Dotora	Univ. of Nobraska
44-0	C. elongatus	INE	Flatte		41.03132	90.10852	3/14/2004	E. Feleis	Ineuraska Univ. of
11_7	C alongatus	NE	Platte		41 02162	06 13811	5/14/2004	F Peters	Nebraska
/	C. elongulus	INL	Tatte		41.02102	70.13011	3/14/2004	L. I CICIS	Univ of
44-8	C elongatus	NE	Platte		41 05132	96 10832	5/14/2004	E Peters	Nebraska
110	C. cionguius	T (L	1 lutte		11.00102	<i>y</i> 0.100 <i>3</i> 2	5/11/2001		Univ of
44-10	C. elongatus	NE	Platte		41.06734	96.05474	5/14/2004	E. Peters	Nebraska
									Univ. of
44-11	C. elongatus	NE	Platte		41.05132	96.10832	5/14/2004	E. Peters	Nebraska
	0								Univ. of
44-13	C. elongatus	NE	Platte		41.05132	96.10832	5/14/2004	E. Peters	Nebraska
									Univ. of
44-14	C. elongatus	NE	Platte		41.05132	96.10832	5/14/2004	E. Peters	Nebraska
									Univ. of
44-16	C. elongatus	NE	Platte		41.05132	96.10832	5/14/2004	E. Peters	Nebraska

<i>a</i> .		D.'	River	Latitude	Longitude	Date of		Institution or
Species	State	River	mile	(N)	(W)	collection	Collector	agency
C elongatus	NE	Platte		41 02162	96 13811	5/14/2004	E Peters	Nebraska
e. cionguius		1 14000		11.02102	<i>y</i> 0.12011	0/11/2001	L. I etells	Univ. of
C. elongatus	NE	Platte		41.06734	96.05474	5/14/2004	E. Peters	Nebraska
C								Univ. of
C. elongatus	NE	Platte		41.02299	96.138	5/18/2004	E. Peters	Nebraska
								Univ. of
C. elongatus	NE	Platte		41.02162	96.13811	5/14/2004	E. Peters	Nebraska
				41.06724	06.05474	5/14/2004		Univ. of
C. elongatus	NE	Platte		41.06/34	96.05474	5/14/2004	E. Peters	Nebraska
	NE	Dlatta		41.05122	06 10922	5/14/2004	E Dotora	Univ. of
C. elongatus	INE	Flatte		41.03132	90.10852	3/14/2004	E. Peters	Univ of
C elongatus	NE	Platte		41 06734	96 05474	5/14/2004	E Peters	Nebraska
C. cionguius		1 futto		11.00751	20.02171	5/11/2001	E. I etelb	Univ of
C. elongatus	NE	Platte		41.02162	96.13811	6/15/2004	E. Peters	Nebraska
0								Univ. of
C. elongatus	NE	Platte		41.02162	96.13811	5/14/2004	E. Peters	Nebraska
C alamaatus	тх	Colorado		20 12561	96 32302	10/19/2004	I Webster	Bio-West
C. elongulus	IA	Colorado		27.42501	70.52502	10/17/2004	J. WEDSTEI	Dio-west
C elongatus	ТΧ	Colorado		29 42561	96 32302	10/19/2004	J Webster	Bio-West
e. cionganas		00101000			,	10/19/2001		210 11 000
C. elongatus	TX	Colorado		29.42561	96.32302	10/19/2004	J. Webster	<b>Bio-West</b>
C								
C. elongatus	TX	Colorado		29.42561	96.32302	10/19/2004	J. Webster	<b>Bio-West</b>
C. elongatus	ТΧ	Colorado		29.42561	96.32302	10/19/2004	J. Webster	Bio-West
C. elongatus	TX	Colorado		29.33229	96.24072	10/20/2004	J. Webster	<b>Bio-West</b>
	Species C. elongatus C. elongatus	SpeciesStateC. elongatusNEC. elongatusTXC. elongatusTX	SpeciesStateRiverC. elongatusNEPlatteC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColoradoC. elongatusTXColorado	SpeciesStateRivermileSpeciesStateRivermileC. elongatusNEPlatteC. elongatusTXColoradoC. elongatusTXColorado	SpeciesStateRiverRiverLatitude (N)C. elongatusNEPlatte41.02162C. elongatusNEPlatte41.06734C. elongatusNEPlatte41.02299C. elongatusNEPlatte41.02162C. elongatusNEPlatte41.02162C. elongatusNEPlatte41.06734C. elongatusNEPlatte41.06734C. elongatusNEPlatte41.05132C. elongatusNEPlatte41.06734C. elongatusNEPlatte41.02162C. elongatusNEPlatte41.02162C. elongatusNEPlatte41.02162C. elongatusNEPlatte41.02162C. elongatusTXColorado29.42561C. elongatusTXColorado29.33229	SpeciesStateRiverRiver mileLatitude (N)Longitude (W)C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0673496.05474C. elongatusNEPlatte41.0229996.138C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0673496.05474C. elongatusNEPlatte41.0513296.10832C. elongatusNEPlatte41.0513296.10832C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0216296.13811C. elongatusNEPlatte41.0216296.13811C. elongatusTXColorado29.4256196.32302C. elongatusTXColorado29.4256196.32302C. elongatusTXColorado29.4256196.32302C. elongatusTXColorado29.4256196.32302C. elongatusTXColorado29.4256196.32302C. elongatusTXColorado29.4256196.32302C. elongatusTXColorado29.4256196.32302C	River SpeciesRiver mileLatitude $(N)$ Longitude $(W)$ Date of collectionC. elongatusNEPlatte $41.02162$ $96.13811$ $5/14/2004$ C. elongatusNEPlatte $41.06734$ $96.05474$ $5/14/2004$ C. elongatusNEPlatte $41.02299$ $96.138$ $5/18/2004$ C. elongatusNEPlatte $41.02162$ $96.13811$ $5/14/2004$ C. elongatusNEPlatte $41.02162$ $96.13811$ $5/14/2004$ C. elongatusNEPlatte $41.06734$ $96.05474$ $5/14/2004$ C. elongatusNEPlatte $41.06734$ $96.05474$ $5/14/2004$ C. elongatusNEPlatte $41.02162$ $96.13811$ $6/15/2004$ C. elongatusNEPlatte $41.02162$ $96.13811$ $5/14/2004$ C. elongatusTXColorado $29.42561$ $96.32302$ $10/19/2004$ C. elongatusTXColorado $2$	SpeciesStateRiverRiverLatitude (N)Longitude (W)Date of collectionCollectorC. elongatusNEPlatte41.0216296.138115/14/2004E. PetersC. elongatusNEPlatte41.0216296.054745/14/2004E. PetersC. elongatusNEPlatte41.0229996.1385/18/2004E. PetersC. elongatusNEPlatte41.0216296.138115/14/2004E. PetersC. elongatusNEPlatte41.0216296.138115/14/2004E. PetersC. elongatusNEPlatte41.0673496.054745/14/2004E. PetersC. elongatusNEPlatte41.0673496.054745/14/2004E. PetersC. elongatusNEPlatte41.0216296.138116/15/2004E. PetersC. elongatusNEPlatte41.0216296.138116/15/2004E. PetersC. elongatusNEPlatte41.0216296.138115/14/2004E. PetersC. elongatusNEPlatte41.0216296.138115/14/2004E. PetersC. elongatusNEPlatte41.0216296.138115/14/2004E. PetersC. elongatusNEPlatte41.0216296.3230210/19/2004J. WebsterC. elongatusTXColorado29.4256196.3230210/19/2004J. WebsterC. elongatusTXColorado29.4256196.3230210/19/2004 </td

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
45-7	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-8	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-9	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-10	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-11	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-12	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-13	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-14	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-15	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-16	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-17	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-18	C. elongatus	ТХ	Colorado		29.33229	96.24072	10/20/2004	J. Webster	Bio-West
45-19	C. elongatus	ТХ	Colorado		29.564	90.5415	10/21/2004	J. Webster	Bio-West
45-20	C. elongatus	TX	Colorado		29.564	90.5415	10/21/2004	J. Webster	Bio-West
45-21	C. elongatus	TX	Colorado		29.564	90.5415	10/21/2004	J. Webster	Bio-West

45-22	C. elongatus	ТХ	Colorado		29.564	90.5415	10/21/2004	J. Webster	Bio-West
45-23	C. elongatus	ТХ	Colorado		29.564	90.5415	10/21/2004	J. Webster	Bio-West
45-24	C. elongatus	ТХ	Colorado		29.564	90.5415	10/21/2004	J. Webster	Bio-West
45-25	C. elongatus	ТХ	Colorado		30.0054	97.0937	10/22/2004	J. Webster	Bio-West
45-26	C. elongatus	ТХ	Colorado		30.0054	97.0937	10/22/2004	J. Webster	Bio-West
45-27	C. elongatus	ТХ	Colorado		30.08529	97.22236	10/27/2004	M. Robertson	Bio-West
45-28	C. elongatus	ТХ	Colorado		30.08529	97.22236	10/27/2004	M. Robertson	Bio-West
45-29	C. elongatus	ТХ	Colorado		30.08529	97.22236	10/27/2004	M. Robertson	Bio-West
45-30	C. elongatus	ТХ	Colorado		30.08529	97.22236	10/27/2004	M. Robertson	Bio-West
50-1	C. elongatus	NE	Missouri	811				M. Bessert	Univ. of Nebraska
50-2	C. elongatus	NE	Missouri				11/11/2003	M. Bessert	Nebraska
50-3	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Univ. of Nebraska
50-4	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Univ. of Nebraska
50-5	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Univ. of Nebraska
50-6	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Univ. of Nebraska

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
	-								Univ. of
50-7	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Nebraska
									Univ. of
50-8	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Nebraska
									Univ. of
50-9	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Nebraska
									Univ. of
50-10	C. elongatus	NE	Missouri	507	40.148	95.433	11/12/2003	M. Bessert	Nebraska
									NE Game
50-11	C. elongatus	NE	Missouri	507.7	40.15415	95.43599	11/12/2003	K. Steffensen	and Parks
									NE Game
50-12	C. elongatus	NE	Missouri	507.7	40.15415	95.43599	11/12/2003	K. Steffensen	and Parks
									NE Game
50-13	C. elongatus	NE	Missouri	507.7	40.15414	95.43598	11/12/2003	K. Steffensen	and Parks
									NE Game
50-14	C. elongatus	NE	Missouri	507.7	40.15414	95.43598	11/12/2003	K. Steffensen	and Parks
									NE Game
50-15	C. elongatus	NE	Missouri	507.7	40.15414	95.43598	11/12/2003	K. Steffensen	and Parks
									NE Game
50-16	C. elongatus	NE	Missouri	507.7	40.15414	95.43598	11/12/2003	K. Steffensen	and Parks
									NE Game
50-17	C. elongatus	NE	Missouri	507.7	40.15414	95.43598	11/12/2003	K. Steffensen	and Parks
									NE Game
50-18	C. elongatus	NE	Missouri	507.7	40.15414	95.43598	11/12/2003	K. Steffensen	and Parks
							/ /		NE Game
50-19	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
					10.01.12			<b>X G D</b>	NE Game
50-20	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks

T., J., J., 1	Species	Stata	Divor	River	Latitude	Longitude	Date of	Collector	Institution or
Individual	species	State	RIVEI	mme	$(\mathbf{N})$	( •• )	conection	Collector	NE Game
50-21	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
	5								NE Game
50-22	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
									NE Game
50-23	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
									NE Game
50-24	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
50.25	~ .		<u>.</u>		40.01.40	05 04420	11/20/2002		NE Game
50-25	C. elongatus	NE	Missouri	5/4.6	40.8149	95.84428	11/20/2003	K. Sterrensen	and Parks
50.26	C alamantar	NE	Missouri	5746	10 8140	05 84428	11/20/2003	V Staffanson	ne Gaille
30-20	C. elongatus	INE	Iviissouri	5/4.0	40.0149	93.04420	11/20/2003	K. Stelleliseli	NE Game
50-27	C alongatus	NE	Missouri	574.6	40 8149	95 84428	11/20/2003	K Steffensen	and Parks
50 21	C. elongulus	T (L	111550 di l	571.0	10.0117	<i>95</i> .01120	11/20/2005	R. Stellensen	NE Game
50-28	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
	el clonguns			• • • • •					NE Game
50-29	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
	0								NE Game
50-30	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
									NE Game
50-31	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
									NE Game
50-32	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
50.00					40.01.40	0504400	11/20/2002		NE Game
50-33	C. elongatus	NE	Missouri	5/4.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
50.24	<b>a</b> 1	NE	Misser	574 (	40.0140	05 94429	11/20/2002	V Ct - ff	NE Game
50-34	C. elongatus	INE	IVI1SSOUT1	5/4.6	40.8149	95.84428	11/20/2003	K. Sterrensen	and Parks
				River	Latitude	Longitude	Date of		Institution or
------------	-----------------	------------------------	------------	-------	----------	-------------	-------------	---------------	----------------
Individual	Species	State	River	mile	(N)	(W)	collection	Collector	agency
50-35	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
									NE Game
50-36	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
									NE Game
50-37	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
50.20		NIE	Misser	571 (	40.0140	05 04 4 2 0	11/20/2002	V. Ctaffanaan	NE Game
50-38	C. elongatus	NE	MISSOURI	5/4.0	40.8149	95.84428	11/20/2003	K. Sterrensen	and Parks
50-39	C alongatus	NF	Missouri	574.6	40 8149	95 84428	11/20/2003	K Steffensen	and Parks
50-57	C. elongulus	1 NL	1vii550uii	574.0	TU.01TJ	JJ.04420	11/20/2005	R. Steffensen	NE Game
50-40	C. elongatus	NE	Missouri	574.6	40.8149	95.84428	11/20/2003	K. Steffensen	and Parks
	8								
50-41	C. elongatus	NE	Missouri		42.58881	96.69419	4/19/2005	D. Shuman	USFWS
									Tulane
51-1	C. elongatus	WI	Wisconsin		43.187	89.902	9/21/1999	H. Bart	University
-1 0						07.000.00		U.D.	Tulane
51-2	C. elongatus	TN	Duck		35.97742	87.82242	6/3/2000	H. Bart	University
51.2	C 1	$\mathbf{T}\mathbf{V}$	Sahina		22 22057	04 25412	12/0/2002	U Dort	Iulane
51-5	C. elongatus	IЛ	Sabille		52.52657	94.55415	12/9/2005	П. Dalt	Tulane
51-4	C elongatus	ТΧ	Sabine		32 32857	94 35413	12/9/2003	H Bart	University
51 1	C. etongutus	171	Subine		52.52057	71.55115	12/ )/ 2005	II. Duit	Tulane
51-5	C. elongatus	LA	Red		32.89277	93.82091	12/9/2003	H. Bart	University
	8								Tulane
51-6	C. elongatus	LA	Red		32.89277	93.82091	12/9/2003	H. Bart	University
									Tulane
51-7	C. meridionalis	LA	Pearl				3/9/2001	V. Todaro	University

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
51-8	C. meridionalis	LA	Pearl				8/28/1990	D. Jackson	Univ.
51-9	C. meridionalis	LA	Pearl				9/19/1990	D. Jackson	MS State Univ.
52-1	C. elongatus	MN	Minnesota	35.9	45.1310	92.45	8/16/2004	D. Ellison	MNDNR
52-2	C. elongatus	MN	Minnesota	29.7	44.4656	93.35	8/16/2004	D. Ellison	MNDNR
52-3	C. elongatus	MN	Minnesota	35.9	45.1310	92.45	8/16/2004	D. Ellison	MNDNR
55-1	C. elongatus	IN	Wabash	?	?	?	4/19/2005	T. Stefenavage	IN DNR
55-2	C. elongatus	IN	Wabash	?	?	?	5/3/2005	T. Stefenavage	IN DNR
55-3	C. elongatus	IN	Wabash	?	?	?	6/1/2005	T. Stefenavage	IN DNR
55-4	C. elongatus	IN	Wabash	?	?	?	6/1/2005	T. Stefenavage	IN DNR
55-5	C. elongatus	IN	Wabash	?	?	?	6/7/2005	T. Stefenavage	IN DNR
55-6	C. elongatus	IN	Wabash	?	?	?	7/19/2005	T. Stefenavage	IN DNR
55-7	C. elongatus	IN	Wabash	?	?	?	7/20/2005	T. Stefenavage	IN DNR
55-8	C. elongatus	IN	Wabash	?	?	?	8/16/2005	T. Stefenavage	IN DNR
57-1	C. elongatus		Conchos, Mexico		29.2620	104.5254	7/23/2005	L. Lozano Vilano	Univ. Autonoma de Neuvo Leon

Individual	Species	State	River	River mile	Latitude (N)	Longitude (W)	Date of collection	Collector	Institution or agency
58-1	C. elongatus	LA	Sabine		30.52179	93.33378	5/17/2005	B. Reed	LDWF
58-2	C. elongatus	LA	Sabine		30.12561	93.40318	8/11/2005	B. Reed	LDWF
58-3	C. elongatus	LA	Sabine		30.52034	93.33472	8/18/2005	B. Reed	LDWF
58-4	C. elongatus	LA	Sabine		30.52034	93.33472	8/18/2005	B. Reed	LDWF Tulane
58-5	C. elongatus	LA	Sabine		30.84501	93.56738	3/29/2006	W. DeRidder	University
58-6	C. elongatus	LA	Sabine		30.84501	93.56738	3/29/2006	W. DeRidder	University
58-7	C. elongatus	LA	Sabine		30.84501	93.56738	3/29/2006	W. DeRidder	University MS Mus
59-01	C. elongatus	MS	Black		32.7073	90.0934	12/12/2005	J.A. Skains	Nat. Sci. MS Mus
59-02	C. elongatus	MS	Black		32.7073	90.0934	12/12/2005	J.A. Skains	Nat. Sci.