

Spring 5-13-2016

# An Analysis of the Microbial Quality of Packaged Water in Four Sites in Latin America

Karla Feeser

Follow this and additional works at: [http://scholarworks.gsu.edu/iph\\_theses](http://scholarworks.gsu.edu/iph_theses)

---

## Recommended Citation

Feeser, Karla, "An Analysis of the Microbial Quality of Packaged Water in Four Sites in Latin America." Thesis, Georgia State University, 2016.

[http://scholarworks.gsu.edu/iph\\_theses/456](http://scholarworks.gsu.edu/iph_theses/456)

This Thesis is brought to you for free and open access by the School of Public Health at ScholarWorks @ Georgia State University. It has been accepted for inclusion in Public Health Theses by an authorized administrator of ScholarWorks @ Georgia State University. For more information, please contact [scholarworks@gsu.edu](mailto:scholarworks@gsu.edu).

**An Analysis of the Microbial Quality of Packaged Water in Four Sites in Latin America**

By

Karla Feeser

B.S. Towson University

A Thesis Submitted to the Graduate Faculty  
of Georgia State University in Partial Fulfillment  
of the Requirements for the Degree  
Master of Public Health

Atlanta, Georgia

2016

**APPROVAL PAGE**

An Analysis of the Microbial Quality of Packaged Water in Four Sites in Latin America

By

Karla Feeser

Approved:

---

Committee Chair

---

Committee Member

---

Date

## **ACKNOWLEDGEMENTS**

Thank you to my committee chair, Dr. Christine Stauber for her dedication and guidance. My appreciation goes also to Dr. Sheryl Strasser, for her valuable feedback, unique perspective and eternally sunny disposition.

Thank you to Water Ecuador staff for their trust, guidance and financial support: James Golden, Alexander Harding, Heriberto Napa Tobar, Kelly Mills. To Erika Tenorio, Rafael Antonio Quijada Landaerde, and Maria Antezana for their hard work to collect data in La Paz and in Tegucigalpa. To all of the staff at Fundacion Runa, and especially to Lindsay McGeehon de Veloz, for their hospitality in Tena.

Thank you to my husband, Nate, for believing in me always.

## Abstract

**Karla Feeser**

**An Analysis of the Microbial Quality of Packaged Water in Four sites in Latin America**

**(Thesis Chair: Dr. Christine Stauber, GSU School of Public Health)**

Much of the death and disease caused by diarrhea in low and middle income countries could be alleviated with better access to safe drinking water; yet, globally 780 million people lack access. Private, small-scale packaged water providers can and do play an important role in meeting the water needs of populations in impoverished or developing countries where public centralized water utilities are not feasible or not trusted; however, recent studies have indicated concerns about the quality and safety of packaged water. This pilot study seeks to identify factors that may be associated with packaged water quality in four sites in Central and South America. It concludes that a large portion of the contamination found in water that is packaged in reusable containers may stem from inadequate disinfection of the containers between uses, and recommends further research on simple, effective disinfection protocols that are practical for use in low-resource settings. Finally, packaged water enterprises should be considered by policymakers who regulate drinking water quality.

**KEY WORDS:** packaged water, drinking water quality, SWE

## Author's Statement

In presenting this thesis as partial fulfillment of the requirements for an advanced degree from Georgia State University, I agree that the Library of the University shall make it available for inspection and circulation in accordance with its regulations governing materials of this type. I agree that permission to quote from, to copy from, or to publish this thesis may be granted by the author or, in his/her absence, by the professor under whose direction it was written, or in his/her absence, by the Associate Dean, College of Health and Human Sciences. Such quoting, copying or publishing must be solely for scholarly purposes and will not involve potential financial gain. It is understood that any copying from or publication of this dissertation which involves potential financial gain will not be allowed without written permission of the author.

---

Signature of Author

**Notice to Borrowers Page**

All theses deposited in the Georgia State University Library must be used in accordance with the stipulations prescribed by the author in the preceding statement.

The author of this thesis is:

Karla Feeser  
834 Argonne Ave NE, #2  
Atlanta, GA 30308

The Chair of the committee for this thesis is:

Christine E. Stauber, PhD  
Institute of Public Health

Georgia State University  
P.O. Box 3995  
Atlanta, GA 30302-3995

Users of this thesis who are not regularly enrolled as students at Georgia State University are required to attest the acceptance of the preceding stipulation by signing below. Libraries borrowing this thesis for the use of their patrons are required to see that each user records here the information requested.

Name of User	Address	Date	Type of User (Examination Only or Copying)

**Karla Feeser**  
**Curriculum Vitae**

---

Home Address: 834 Argonne Ave NE, #2  
Atlanta, GA 30308  
Phone: 443 – 624 -0903

**Education:**

2010-2012 B.S. Towson University, Baltimore, MD  
2008-2010 A.A. Community College of Baltimore County

**Professional Experience:**

2015- Present Graduate Research Assistant  
Georgia State University

2015-Present Surveillance Officer  
Georgia Emerging Infections Program

2015-2015 Water Research Fellow  
Water Ecuador 501c3

2012-2015 Microbiologist  
Division of Parasitic Diseases and Malaria  
Centers for Disease Control and Prevention



## Table of Contents

Acknowledgements.....	iii
Abstract.....	iv
List of Tables .....	ix
List of Figures.....	x
CHAPTER	
1. INTRODUCTION .....	1
1.1. Purpose of the Research.....	3
2. LITERATURE REVIEW.....	5
2.1. Background.....	5
2.2. Access to Healthy Water.....	6
2.3. The Role of Packaged Water and Small Water Enterprises.....	7
2.3.1. Packaged Water Quality.....	8
2.4. Study Sites.....	12
2.4.1. El Alto, La Paz, Bolivia.....	12
2.4.2. Tegucigalpa, Honduras.....	13
2.4.3. Muisne and Tena, Ecuador.....	13
2.4.4. References.....	15
3. MANUSCRIPT.....	20
3.1. Abstract.....	21
3.2. Introduction.....	22
3.3. Materials and Methods.....	24
3.3.1. Study Setting and Design.....	24
3.3.2. Company Surveys.....	25
3.3.3. Sample Collection.....	25
3.3.4. Testing Procedures.....	27
3.3.5. Data Analysis.....	28
3.4. Results.....	29
3.4.1. Survey Results.....	29
3.4.2. Detection of Total Coliforms and <i>E. coli</i> in water samples.....	31
3.4.3. Results of Analyses.....	34
3.5. Discussion.....	38
3.6. Limitations of the Research.....	41
3.7. Manuscript References.....	43
4. CONCLUSIONS.....	47
5. APPENDICES.....	48
5.1. Appendix A: Company Survey.....	48

## LIST OF TABLES

Table 1. Overview of the 20 companies surveyed in three of four study sites (Tegucigalpa, Muisne and Tena).....	30
Table 2. Water Sample Type by Study Site.....	31
Table 3. Descriptive statistics of total coliform and <i>E.coli</i> concentration in each sample type.....	32
Table 4. Associations of company-treated water and coliform contamination.....	33
Table 5. Quality of water by post-visit sampling time point and sample type.....	35
Table 6. Association between sample type and prevalence of total coliform contamination.....	36
Table 7. Association between sample type and prevalence of <i>E. coli</i> contamination.....	37
Table 8. Results of Univariate and Multivariate Analyses for Presence of Total Coliform Bacteria Among Reusable Bottles.....	38
Table 9. Results of Univariate Analyses for Presence of <i>E. coli</i> Among Reusable Bottles.....	39

## LIST OF FIGURES

- Figure 1. Schematic of a typical water treatment plant and testing protocol .....10
- Figure 2. Percent of samples contaminated with Total Coliforms and E. coli by site and type ... 34

## **Chapter I: Introduction**

Much of the death and disease caused by diarrhea in low and middle income countries could be alleviated with better access to safe drinking water; yet, globally 780 million people lack access (World Health Organization (WHO) & UNICEF, 2012; WHO, 2014; Prüss-Ustün et al., 2014). In developing regions, piped drinking water supplies reach only 46% of the population, due, in part, to disparities in access to safe water infrastructure. Piped water systems often do not reach lower income or rural areas in low-middle income countries where the established infrastructure required of a centralized water utility is often missing, and the high cost to develop missing infrastructure may be prohibitive (Dada, 2011). Even where a centralized utility is available, it may be perceived as unsafe to drink (Hatt, 2006).

In Latin America, access to improved drinking water sources and piped water supplies in particular has risen substantially since 1990 to reach approximately 94% coverage, although substantial disparities between urban and rural areas still exist (WHO & UNICEF, 2012). However, throughout the region, there is mistrust in the quality of the water provided through centralized utilities (de Queiroz, Doria, 2013; Espinosa-García et al., 2015; Jain, 2014). ). Indeed, the microbial quality of piped drinking water sources is not globally monitored; they are thought to be clean due to the nature of their construction. If these sources are poorly maintained or constructed, they may be contaminated with fecal pathogens. Thus, it is likely that the number of people with

access to a safe and improved source of drinking water is even less than estimates allow (WHO & UNICEF, 2012).

Private, small-scale packaged water providers can and do play an important role in meeting the water needs of populations in impoverished or developing countries where public centralized water utilities are not feasible (Solo, 1999). Packaged water, intended to be potable, is sold in a variety of vessels—sachets, pouches, boxes, cans, and reusable and disposable bottles—and typically vended through both water distributors and water kiosks (Dada, 2011). The number of people relying on packaged water, particularly in large twenty liter reusable bottles, is on the rise, even among those households with a connection to a public water utility (JMP, 2011). Oftentimes, even the lowest-income families tolerate the expense because they perceive the water as safe to drink, more so than other sources (Kjellen, 2006).

Packaged water is significantly less likely to be contaminated than other water sources, including improved water sources such as piped water (Williams et al., 2015). However, substantial heterogeneity is found seen across study sites—with more than 40% of studies reporting packaged water to be of equal or lesser quality than piped water. Additionally, significant disparities exist between low income countries (LICs) and upper-middle and high income countries (UM/HICs) in contamination levels of packaged water. Poorer countries may face more obstacles in the monitoring and regulation of packaged water enterprises. In fact, even as small water enterprises gain recognition as viable drinking water alternative, in many developing nations SWEs operate either completely unregulated or unregistered (Dada, 2011). Consequently, studies investigating the microbial contamination of packaged water often find that the quality of packaged water

sold by SWE meets neither international guidelines nor national standards, when they are in place (Fisher et al., 2015; Halage et al., 2015).

For packaged water sold in reusable bottles, inadequate cleaning methods may serve as a potential source of contamination before the point of sale or use (Falcone-Dias, 2012; Marzano, 2011). However, despite the existence of international and national drinking water guidelines, there is no available evidence to suggest that these kinds of stringent processes for the disinfection of reusable containers are enforced in many low income countries throughout the world.

This study seeks to characterize the quality of packaged water available for purchase in four sites in Central and South America: La Paz, Bolivia; Tegucigalpa, Honduras; Muisne, Ecuador; and Tena, Ecuador. Furthermore, it identifies factors that may be associated with packaged water quality, particularly in reusable containers, and discusses recommendations for the regulation of distributors that provided packaged water to underserved communities around the world.

## **1.1 Purpose of Research**

As the activity and importance of SWEs and packaged water vendors increase, more research is needed to assess the quality of water sold, and to inform policies that regulate the private water sector throughout the world. This study seeks to identify factors that may contribute to the deterioration of packaged water quality, and to establish

recommendations for business owners and managers of SWEs that provided packaged water to underserved communities around the world.

## Chapter 2: Literature Review

### 2.1 Background

Diarrheal disease contributes to an estimated 1.5 million deaths each year, including 760,000 deaths among children under the age of five (“WHO | Diarrhoeal disease,” n.d.). Globally, these diseases represent the second leading cause of childhood mortality (Liu et al., 2012). In Latin American countries, twenty to thirty percent of children under five experience diarrhea each month, with the burden falling heaviest on those of lower socioeconomic status (Hatt & Waters, 2006).

Bacteria, viruses or parasites transmitted via the fecal-oral route are the most common causes of diarrhea, especially in the developing world. Fecal-oral transmission occurs through contact with food, water or surfaces that have been contaminated with fecal matter from an infected individual (Kelly, 2011). Important pathogens include norovirus and rotavirus, which are highly infectious and easily transmitted. *Shigella* and toxigenic *E. coli* are common bacterial agents of serious diarrhea, and *Cryptosporidium* exemplifies a potentially culpable parasite (Kotloff et al., 2013).

Furthermore, diarrheal diseases impact the absorption of nutrients in the gut, making them a leading cause of malnutrition worldwide. Malnutrition is a complex health outcome of diarrhea because it leads to an increase in both the frequency and duration of diarrheal episodes. This positive feedback loop of malnutrition and diarrhea not only increases childhood mortality, but also leads to stunted growth, lowered physical fitness, and impaired cognition and school performance (Gracey, 1996). The resulting impact on education, especially among young girls,



often manifests into decreased productivity and limited economic opportunity and growth (Jasper, 2012), which bears lifelong economic implications.

It is estimated that 58% of deaths due to diarrhea can be explained by poor water, sanitation and hygiene (WHO, 2014). Of those, approximately 500,000 are attributable to inadequate drinking water (Prüss-Ustün et al., 2014). Thus, much of the death and disease caused by diarrhea in developing nations could be alleviated with better access to safe drinking water.

## **2.2 Access to Healthy Water**

The Millennium Development Goals (MDG) set forth by the United Nations in 2000 embody a commitment by world leaders to end extreme poverty (UN, 2015). The seventh goal—to ensure environmental sustainability—includes Target 7.C, which aims to “halve, by 2015, the proportion of people without sustainable access to safe drinking water and basic sanitation” (UN, 2015). Progress toward achieving this target is monitored by a collaborative group formed by UNICEF and the World Health Organization named the Joint Monitoring Programme (JMP) for Water Supply and Sanitation (WHO/UNICEF, 2010). JMP plays a key role in the development of measurable indicators used to evaluate progress toward the fulfillment of water and sanitation targets. JMP breaks drinking-water sources down into two main categories: improved sources and unimproved sources. To be classified as improved, a drinking- water source must “by nature of its construction and when properly used, adequately protect the source from outside contamination, particularly faecal matter” (WHO/UNICEF, n.d.). For example, improved water sources can include water that is piped directly to the home or to a public tap, tubewells that draw from groundwater supplies, rainwater, or wells that are dug, constructed and covered to

prevent contamination from entering the source water. However, in practice, these definitions do not stipulate that improved water sources be free of microbial contamination in order to be counted toward progress in achieving Target 7.C, thus recent estimates of the number of people with access to safe drinking water may be inflated (Bain et al., 2014).

### **2.3 The Role of Packaged Water and Small Water Enterprises**

Though one may expect packaged and bottled water to be generally safe for consumption, they are not considered improved drinking-water sources by the JMP (WHO/UNICEF, n.d.). Bottled water is not sufficient, affordable or feasible for use as a sole household water source. For basic uses, including drinking, personal hygiene and cooking, the average person needs approximately 50 liters of water per day (Gleick, 1996). For this reason, bottled water is only considered to be an improved drinking-water source if the household uses some other improved water source for other tasks (WHO/UNICEF, 2011). Nevertheless, packaged and bottled drinking water can fill gaps in areas where improved water sources are unavailable or mistrusted (Hatt & Waters, 2006). Private, small-scale providers can and do play an important role in meeting the water needs of populations in impoverished or developing countries where public centralized water utilities are not feasible (Solo, 1999). Historically, these private water providers have been most abundant in areas where piped water was unsafe, unreliable or economically impossible (Zaroff, 1984). Many developing countries lack the established infrastructure required of a centralized water utility, and the high cost to develop missing infrastructure is often prohibitive (Dada, 2011). Small water enterprises (SWEs) have proven to be efficient and sustainable as alternative drinking water providers in low-resources settings (Bhatt, 2014).

Water vendors, or SWEs can take on many forms. Privately owned connection networks pull water from the public utility and pipe it directly to homes or community standpipes. Water distributors deliver water to households in a variety of ways, from large tankers, which truck water in and dispense into household containers, to packaged water vendor who deliver water bottled in reusable containers. Water kiosks provide a stationary point of sale where individuals can purchase water to carry home by re-filling household containers, purchasing or exchanging empty containers for full containers, or purchasing water in disposable containers such as sachets or small bottles (Kjellen, 2006). Packaged water, intended to be potable, is sold in a variety of vessels—sachets, pouches, boxes, cans, and reusable and disposable bottles—and typically vended through both water distributors and water kiosks (Dada, 2011).

The number of people relying on packaged water, particularly in large twenty liter reusable bottles, is on the rise, even among those households with a connection to a public water utility (JMP, 2011). Oftentimes, even the lowest-income families tolerate the expense because they perceive the water as safe to drink, more so than other sources (Kjellen, 2006).

### **2.3.1 Packaged Water Quality**

A 2015 systematic review and meta-analysis found that, overall, packaged water is significantly less likely to be contaminated than other water sources, including improved water sources such as piped water (Williams et al., 2015). However, substantial heterogeneity was seen across study sites—with more than 40% studies reporting that packaged water was of equal or lesser quality than piped water. Additionally, the analysis revealed significant disparities between low income countries (LICs) and upper-middle and high income countries (UM/HICs) in contamination levels of packaged water, even when accounting for more prevalent use of sachet water—which


is more likely to be contaminated than bottled water—among LICs. Poorer countries may face more obstacles in the monitoring and regulation of packaged water enterprises. In fact, even as SWEs gain recognition as viable drinking water alternative, in many developing nations SWEs operate either completely unregulated or unregistered (Dada, 2011). Consequently, studies investigating the microbial contamination of packaged water often find that the quality of packaged water sold by SWE meets neither international guidelines nor national standards, when they are in place (Fisher et al., 2015; Halage et al., 2015).

Packaged drinking water can be contaminated at any point along the supply line—from source to household. First, water is not necessarily treated before being packaged and sold and therefore is sometimes only as safe as the source water. Typically, packaged water sold as mineral water is untreated, but assumed safe for consumption because it is pulled from underground springs and bottled on-site (Falcone-Dias, 2012). Non-mineral water, or water from other sources, is most often treated prior to bottling or packaging to improve its quality. WHO recommends that multiple disinfection strategies be used in the treatment of drinking water to ensure an adequate reduction in pathogens in the event that the source water is highly contaminated or contaminated with resistance microorganisms, such as *Cryptosporidium*, which is not susceptible to chlorination alone (WHO, 2011). Thus, a combination of chemical disinfection and filtration is often used.

A typical treatment system for packaged water is shown in Figure 1. Filtration of water can occur through many types of media, and its efficacy depends largely on pore size. For example, cloth filters are useful for removing large microbes ( $> 20 \mu\text{m}$ ), such as parasitic worms or microbes

that are associated with copepods. Filters made from ceramic or carbon have smaller pores and are able to remove more organisms, such as small protozoa, and viruses. Granular media filtration, such as sand filtration, can filter out organisms based on the coarseness or grain of the sand particles and the rate of filtration. The most popular chemical disinfectants are chlorine and ozone gas. Chlorine is often added so that a residual concentration of the chemical remains in water (Impellitteri, 2007). Ozone is used to disinfect water, but no residual concentration remains. The use of chemical disinfectants can lead to the creation of disinfection by products in drinking water. UV radiation of water can be achieved through solar disinfection technology or through the use of UV lamps. Processes to improve the taste, odor or appearance of water may also be included. For example, demineralization of water—water softening—can be achieved through the use of a cation exchange resin (World Health Organization, 2011).

**Figure 1.** Schematic of a typical water treatment plant and testing protocol.



Purified water can become contaminated upon introduction into an unclean container, such as a reusable bottle that has not been appropriately disinfected. Additionally, purified water, even stored in a clean container can become contaminated by pathogens introduced through use in the household (Levy, 2008).

Only a handful of studies mention reusable bottles as a potential source of contamination before the point of sale or use (Falcone-Dias, 2012; Marzano, 2011). A study on the microbiological quality of bottled mineral water reported a higher concentration of total coliform bacteria, *E. coli* and *P. aeruginosa* in 20L bottles than in 1.5 L bottles, but did not collect sufficient information to test the hypothesis that the higher contamination was the result of reusable bottles that had not been cleaned appropriately (Falcone-Dias, 2012). Similarly, a 2010 study examining bottled water dispensers as a potential source for contamination in large reusable containers noted the presence of bacteria in the bottles, and suggested that it might be attributable to either the water production plant or the plastic bottles themselves (Marzano, 2011).

When testing the quality of drinking water, the drinking water guidelines set forth by WHO dictate that there should be no detectable *E. coli* per 100mL of water intended for drinking, including packaged or bottled water (WHO, 2011).

Companies that produce and sell bottled water in the United States are regulated by the U.S. Food and Drug Administration (FDA) and have standard protocols for inspecting and cleaning reusable containers (Nestle Waters North American Inc., 2010). However, despite the existence of international and national drinking water guidelines, there is no available evidence to suggest

that these kinds of stringent process for the disinfection of reusable containers are enforced in many low income countries throughout the world.

## **2.4 Study Sites**

In Latin American countries, the use of packaged and bottled drinking water is substantial. In 2008, it is estimated that a projected volume of 5,343 million liters of bottled water were consumed in Latin America (Datamonitor, 2004). In parts of Central and South America, the industry has grown as much of 55% over the last five years (“Informe anual bebidas 2015,” 2015). Although exact data are limited, reports suggest that people throughout Central and South America are increasingly dependent on private water enterprises for purified drinking water (Kjellen, 2006; Pacheco-Vega, 2015).

### **2.4.1 El Alto, La Paz, Bolivia**

In the late 1990s, issues of water scarcity and poor water infrastructure led to a massive privatization of Bolivia’s water resources (Baer, 2015). Particularly, in the city of Cochabamba and in the peri-urban areas surrounding La Paz, concessions that allowed large multi-national corporations to control water utilities were met with backlash and social uprisings. The concessions were reversed, and since then the Bolivian government has amended their constitution to include water as a human right (Baer, 2015)). The Bolivian government ministries have gone through several phases of restructuring as it pertains to water resource provisioning, regulation, and policy making. The structure and governance of the public water utility in the area of La Paz and the surrounding community of El Alto, where the current study was conducted, have been in transition since 2007 (Baer, 2015). Piped water coverage has risen in Bolivia—90 percent of households use piped water or another improved water source compared

to just 69 percent in 1990—but there is large disparity between urban and rural areas, where coverage hovers just under 60 percent (WHO/UNICEF, 2015c). Furthermore, reports have indicated that the majority of water provided through the public water utility is untreated and unsafe to drink (Fundacion Abril, 2013). Thus, SWEs still play an important role in the study area. In particular, informal water distributors and water kiosks operate on a largely unregulated basis (Wutich, 2016).

#### **2.4.2 Tegucigalpa, Honduras**

According to most recent estimates, 90 percent of households in Honduras used piped water (WHO/UNICEF, 2015b). Nevertheless, up to thirty percent of households in some communities continue to use bottled water for drinking and cooking (Halder, 2013). Generally, the bottled water industry throughout Honduras is substantial, owing chiefly to mistrust in the quality of water and reliability of service provided by municipal systems (Public-Private Infrastructure Advisory Facility, 2003). In Honduras, the Technical Standard for Drinking Water Quality (TSDWQ) regulates the production of bottled water. Like the WHO guidelines, the allowable limit for *E. coli* in drinking water under the TSDWQ is zero CFU per 100mL (Ministerio de Salud, 1995).

#### **2.4.3 Muisne and Tena, Ecuador**

The percentage of households using improved water sources in Ecuador has risen significantly since 1990—when only 38 percent of rural households and 59 percent of urban households had access to piped water. However, even today nearly 30 percent of rural households are without access to piped water, and coverage throughout the country has reached only 85 percent (WHO/UNICEF, 2015a). Two of the four present study sites were located in rural Ecuador. The first, Muisne, is a coastal town in the Esmeraldas province in the northern part of the country. In



this part of the country, people relying chiefly on untreated water for drinking (Levy, 2012). The second site was located in the interior of the county just east of the Amazon Rainforest's westernmost edge. Tena and the smaller neighboring town of Archidona lie along the River Napo, which also serves as the main drinking water reservoir for the city's municipal water systems.

## References

- Baer, M. (2015). From Water Wars to Water Rights: Implementing the Human Right to Water in Bolivia. *Journal of Human Rights*, 14(3), 353–376.  
<http://doi.org/10.1080/14754835.2014.988782>
- Bain, R., Cronk, R., Wright, J., Yang, H., Slaymaker, T., & Bartram, J. (2014). Fecal Contamination of Drinking-Water in Low- and Middle-Income Countries: A Systematic Review and Meta-Analysis. *PLoS Medicine*, 11(5).  
<http://doi.org/10.1371/journal.pmed.1001644>
- Bhatt, J. (2014). Comparison of small-scale providers' and utility performance in urban water supply: the case of Maputo, Mozambique. *Water Policy*, 16(1), 102–123.  
<http://doi.org/10.2166/wp.2013.083>
- Dada, A. (2011). Packaged water: optimizing local processes for sustainable water delivery in developing nations. *Globalization & Health*, 7(1), 24–31.  
<http://doi.org/10.1186/1744-8603-7-24>
- Falcone-Dias, M. F., Emerick, G. L., & Farache-Filho, A. (2012). Mineral water: a microbiological approach. *Water Science & Technology: Water Supply*, 12(5), 556–562.  
<http://doi.org/10.2166/ws.2012028>
- Fisher, M. B., Williams, A. R., Jalloh, M. F., Saquee, G., Bain, R. E. S., & Bartram, J. K. (2015). Microbiological and Chemical Quality of Packaged Sachet Water and Household Stored Drinking Water in Freetown, Sierra Leone. *PLOS ONE*, 10(7), e0131772.  
<http://doi.org/10.1371/journal.pone.0131772>
- Fundacion Abril. (2013, June 6). Agua y Saneamiento. Retrieved from  
<http://www.fundacionabril.org/about-2/agua-y-saneamiento/>

- Gleick, P. (1996). Basic Water Requirements for Human Activities: Meeting Basic Needs. *Water International*, 21, 83–92.
- Gracey, M. (1996). Diarrhea and Malnutrition: A Challenge for Pediatricians. *Journal of Pediatric Gastroenterology & Nutrition*, 22(1), 6–16.
- Halage, A. A., Ssemugabo, C., Ssemwanga, D. K., Musoke, D., Mugambe, R. K., Guwatudde, D., & Ssempebwa, J. C. (2015). Bacteriological and physical quality of locally packaged drinking water in Kampala, Uganda. *Journal of Environmental and Public Health*, 2015, 942928. <http://doi.org/10.1155/2015/942928>
- Halder, G. E., Bearman, G., Sanogo, K., & Stevens, M. P. (2013). Water sanitation, access, use and self-reported diarrheal disease in rural Honduras. *Rural and Remote Health*, 13(2), 2413.
- Hatt, L. E., & Waters, H. R. (2006). Determinants of child morbidity in Latin America: A pooled analysis of interactions between parental education and economic status. *Social Science & Medicine*, 62(2), 375–386. <http://doi.org/10.1016/j.socscimed.2005.06.007>
- Jasper, C., Le, T.-T., & Bartram, J. (2012). Water and sanitation in schools: a systematic review of the health and educational outcomes. *International Journal Of Environmental Research And Public Health*, 9(8), 2772–2787. <http://doi.org/10.3390/ijerph9082772>
- Kelly, P. (2011). Infectious diarrhoea. *Medicine*, 39(4), 201–206. <http://doi.org/10.1016/j.mpmed.2011.01.008>
- Kjellen, M., & McGranahan, G. (2006). Informal Water Vendors and the Urban Poor. International Institute for Environment and Development (IIED). Retrieved from <http://pubs.iied.org/pdfs/10529IIED.pdf>
- Kotloff, K. L., Nataro, J. P., Blackwelder, W. C., Nasrin, D., Farag, T. H., Panchalingam,

- S., ... Levine, M. M. (2013). Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *The Lancet*, 382(9888), 209–22.  
[http://doi.org/http://dx.doi.org.ezproxy.gsu.edu/10.1016/S0140-6736\(13\)60844-2](http://doi.org/http://dx.doi.org.ezproxy.gsu.edu/10.1016/S0140-6736(13)60844-2)
- Levy, K., Nelson, K. L., Hubbard, A., & Eisenberg, J. N. S. (2008). Following the Water: A Controlled Study of Drinking Water Storage in Northern Coastal Ecuador. *Environmental Health Perspectives*, 116(11), 1533–1540.  
<http://doi.org/10.1289/ehp.11296>
- Levy, K., Nelson, K. L., Hubbard, A., & Eisenberg, J. N. S. (2012). Rethinking Indicators of Microbial Drinking Water Quality for Health Studies in Tropical Developing Countries: Case Study in Northern Coastal Ecuador. *The American Journal of Tropical Medicine and Hygiene*, 86(3), 499–507. <http://doi.org/10.4269/ajtmh.2012.11-0263>
- Liu, L., Johnson, H. L., Cousens, S., Perin, J., Scott, S., Lawn, J. E., ... Child Health Epidemiology Reference Group of WHO and UNICEF. (2012). Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet (London, England)*, 379(9832), 2151–2161.  
[http://doi.org/10.1016/S0140-6736\(12\)60560-1](http://doi.org/10.1016/S0140-6736(12)60560-1)
- Marzano, M. a., Ripamonti, B., & Balzaretto, C. m. (2011). Short communication: Monitoring the bacteriological quality of Italian bottled spring water from dispensers. *Food Control*, 22, 333–336. <http://doi.org/10.1016/j.foodcont.2010.06.014>
- Ministerio de Salud. (1995). Norma tecnica para la calidad del Agua Potable. Gobierno de Honduras.
- Nestle Waters North American Inc. (2010). Poland Spring Water Quality Report.

Stamford: Nestle Waters.

Prüss-Ustün, A., Bartram, J., Clasen, T., Colford, J. M., Cumming, O., Curtis, V., ...

Cairncross, S. (2014). Burden of disease from inadequate water, sanitation and hygiene in low- and middle-income settings: a retrospective analysis of data from 145 countries.

*Tropical Medicine & International Health*, 19(8), 894–905.

<http://doi.org/10.1111/tmi.12329>

Public-Private Infrastructure Advisory Facility. (2003). Private Solutions for Infrastructure in Honduras. World Bank Publications.

Solo, T. M. (1999). Small-scale entrepreneurs in the urban water and sanitation market.

*Environment and Urbanization*, 11(1), 117–132.

<http://doi.org/10.1177/095624789901100120>

United Nations. (2015). The Millennium Development Goals Report 2015. Retrieved from

[http://www.un.org/millenniumgoals/2015\\_MDG\\_Report/pdf/MDG%202015%20rev%20\(July%201\).pdf](http://www.un.org/millenniumgoals/2015_MDG_Report/pdf/MDG%202015%20rev%20(July%201).pdf)

United Nations. (n.d.). Goal 7 Fact Sheet: Millennium Development Goals and Beyond 2015. Retrieved from [http://www.un.org/millenniumgoals/pdf/Goal\\_7\\_fs.pdf](http://www.un.org/millenniumgoals/pdf/Goal_7_fs.pdf)

WHO | Diarrhoeal disease. (n.d.). Retrieved February 24, 2016, from

<http://www.who.int/mediacentre/factsheets/fs330/en/>

WHO/UNICEF. (2010). JMP Technical Task Force Meeting on Monitoring Drinking-water Quality. Villie-Morgon, France. Retrieved from

[http://www.wssinfo.org/fileadmin/user\\_upload/resources/JMP-Task-Force-Meeting-on-Monitoring-Drinking-water-Quality.pdf](http://www.wssinfo.org/fileadmin/user_upload/resources/JMP-Task-Force-Meeting-on-Monitoring-Drinking-water-Quality.pdf)

- WHO/UNICEF. (2015a). Ecuador: Estimates on the use of water sources and sanitation facilities.
- WHO/UNICEF. (2015b). Honduras: Estimates on the use of water sources and sanitation facilities. Joint Monitoring Programme for Water Supply and Sanitation.
- WHO/UNICEF. (2015c, June). Bolivia (Plurinational State of): Estimates on the use of water sources and sanitation facilities. Joint Monitoring Programme for Water Supply and Sanitation.
- WHO/UNICEF. (n.d.). Definitions and Methods: wat/san categories. Retrieved February 3, 2016, from <http://www.wssinfo.org/definitions-methods/watsan-categories/>
- WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation (JMP). (2011). Drinking Water Equity, Safety and Sustainability; Thematic report on drinking water.
- Williams, A. R., Bain, R. E. S., Fisher, M. B., Cronk, R., Kelly, E. R., & Bartram, J. (2015). A Systematic Review and Meta-Analysis of Fecal Contamination and Inadequate Treatment of Packaged Water. *PLoS ONE*, 10(10), 1–27. <http://doi.org/10.1371/journal.pone.0140899>
- World Health Organization (WHO). (n.d.). Guidelines for Drinking-water Quality, Fourth Edition. Retrieved February 24, 2016, from [http://apps.who.int/iris/bitstream/10665/44584/1/9789241548151\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44584/1/9789241548151_eng.pdf)
- Wutich, A., Beresford, M., & Carvajal, C. (2016). Can Informal Water Vendors Deliver on the Promise of A Human Right to Water? Results From Cochabamba, Bolivia. *World Development*, 79, 14–24. <http://doi.org/10.1016/j.worlddev.2015.10.043>

## **Chapter 3: Manuscript**

**An Analysis of the Microbial Quality of Packaged Water in Four sites in Latin America**

Karla R. Feeser

### 3.1 Abstract

**INTRODUCTION:** Diarrheal disease contributes to an estimated 1.5 million deaths each year, including 760,000 deaths among children under the age of five. Of those, approximately 500,000 are attributable to inadequate drinking water. In areas where piped water is unsafe, unreliable or economically impossible, packaged water sold by private vendors can play an important role in meeting the water needs of these populations. As the activity and importance of packaged water vendors grow, more data is needed to assess the quality of water sold, and to inform policies that regulate the private water sector throughout the world.

**AIM:** This pilot study seeks to identify factors that may contribute to the deterioration of packaged water quality.

**METHODS:** Small packaged water enterprises (SWEs) operating in the following cities were visited twice between May 2014 and September 2015: La Paz, Bolivia; Tegucigalpa, Honduras; and Muisne and Tena, Ecuador. A brief survey was conducted with each distributor, and a facility tour was completed. Water samples were collected directly from the purification system and water packaged in both reusable and disposable containers were purchased. Samples were tested for total coliform and *E. coli* bacterial contamination on the day of collection and over the course of 28 days. Data was analyzed using descriptive statistics, including median as the measure of central tendency, and frequency where the main outcome was presence or absence of either total coliform bacteria or *E. coli*. To determine the factors that were most associated with water quality deterioration, logistic regression was performed.

**RESULTS:** A total of 616 samples were collected. This study found that 52% of the packaged water examined was contaminated with total coliform bacteria. Raw, untreated water and treated water packaged in reusable containers were most likely to be contaminated with total coliform bacteria and *E.coli* compared to treated water taken directly from the system. There was no significant association between water treatment or bottle disinfection protocols and total coliform or *E. coli* contamination.

**DISCUSSION:** The study succeeded in identifying at what stage and in what type of container water is most likely to be contaminated with bacterial water-quality indicators. Furthermore, it highlights the heterogeneity that exists in terms of types of water sold, water treatment systems, and sanitizing protocols among SWEs in Central and South America. Reusable containers are vulnerable to contamination with total coliform bacteria and *E. coli*, even when filled with clean water, thus the contamination may be due to inadequate disinfection between uses. These results may have implications for national or international policies that regulate private water enterprises, and can inform guidelines for packaged water distributors in particular. Further research is needed to identify optimal cleaning methods for reusable containers that are practical for use in lower resource settings.



### **3.2 Introduction**

Globally, 780 million people lack access to improved drinking water (WHO & UNICEF, 2012). In developing regions, piped drinking water supplies reach only 46% of the population, due, in part, to disparities in access to safe water infrastructure. Piped water systems often do not reach lower income or rural areas in low-middle income countries where the established infrastructure required of a centralized water utility is often missing, and the high cost to develop missing infrastructure may be prohibitive (Dada, 2011). Where a centralized utility is available, it may be perceived as unsafe to drink (Hatt, 2006). Indeed, the microbial quality of piped drinking water sources is not globally monitored; they are thought to be clean due to the nature of their construction. If these sources are poorly maintained or constructed, they may be contaminated with fecal pathogens. Thus, it is likely that the number of people with access to a safe and improved source of drinking water is even less than estimates allow (WHO & UNICEF, 2012).

In Latin America, access to improved drinking water sources and piped water supplies in particular has risen substantially since 1990 to reach approximately 94% coverage, although substantial disparities between urban and rural areas still exist (WHO & UNICEF, 2012).

However, throughout the region, there is mistrust in the quality of the water provided through centralized utilities (de Queiroz, Doria, 2013; Espinosa-García et al., 2015; Jain, 2014).

A growing number of people throughout the world are using bottled or packaged drinking water to fill gaps in water access and water quality. From 1990 to 2010, the number of people using bottled water rose from 37 million to 228 million and the number of people relying on private water vendors more than doubled (WHO & UNICEF, 2012). Although exact data for Central and South America are limited, reports suggest that people throughout these regions are also

increasingly dependent on private water enterprises for packaged, purified drinking water (Kjellen, 2006; Opryszko, 2009; Pacheco-Vega, 2015). Oftentimes, even where piped water is available, families may tolerate the added expense because they perceive the water as safe to drink, more so than other sources (Kjellen, 2006).

Water can be packaged into a variety of containers—disposable or reusable, bottles or sachets—and the delivery of water to consumers may vary greatly in different settings; however, reusable 20 liter (L) bottles are found in LMICs throughout Central and South America (Liu, 2013; Malkin, 2012). Typically, these plastic, cylindrical, narrow mouthed bottles are filled with water when purchased from a water distributor. When consumers empty the bottle, they bring it back to the distributor and the distributor either refills the bottle or exchanges the empty bottle for a full one. Despite their widespread use, little is known about the quality and safety of these large reusable bottles. In fact, despite many studies regarding the safe treatment of drinking water, there is a substantial gap in the scientific and public health literature concerning the quality of reusable 20L bottles and how to best implement and measure the impact of any associated interventions.

Companies that produce and sell bottled water in the United States are regulated by the U.S. Food and Drug Administration (FDA) and have standard protocols for inspecting and cleaning reusable containers (Nestle Waters North American Inc., 2010). However, despite the existence of international and national drinking water guidelines (World Health Organization, 2011), there is no available evidence to suggest that these kinds of stringent process for the disinfection of reusable containers are enforced in many low income countries throughout the world.

Poorer countries may face more obstacles in the monitoring and regulation of packaged water enterprises. In fact, even as private water distributors gain recognition as viable drinking water alternative, in many developing nations they operate either completely unregulated or unregistered (Dada, 2011). Consequently, studies investigating the microbial contamination of packaged water often find that the quality of packaged water sold by private distributors meets neither international guidelines nor national standards, where they are in place (Fisher et al., 2015; Halage et al., 2015).

This study seeks to characterize the quality of packaged water available for purchase in four sites in Central and South America: La Paz, Bolivia; Tegucigalpa, Honduras; Muisne, Ecuador; and Tena, Ecuador. Furthermore, it identifies factors that may contribute to the deterioration of packaged water quality, particularly in reusable containers, and discusses recommendations for the regulation of distributors that provided packaged water to underserved communities around the world.

### **3.3 Materials and Methods**

#### **3.3.1 Study Setting and Design**

This study was carried out as a pilot project in four study sites across three Central and South American countries between May 2014 and September 2015. The study was undertaken by the non-profit organization Water Ecuador in the summer of 2015 in La Paz, Bolivia and in Tegucigalpa, Honduras with the collaboration of the Zamorano Escuela Agrícola Panamericana. In Ecuador, the study was carried out in Muisne in between May and August 2014 and in Tena in between May and August 2015.

In each study site, packaged water distributors were identified using an informal reconnaissance survey to identify the most popular brands sold in local stores and directly from trucks or kiosks. Once identified, water distributors were visited during their normal operating hours so that researchers could collect samples, purchase water and interview the business manager.

This protocol was approved by the Georgia State University Institutional Review Board (IRB) and designated as not human subjects research (GSU IRB protocol number: H16339).

### **3.3.2 Company Surveys**

A brief survey was conducted with each distributor at the Tegucigalpa, Tena and Muisne study sites to determine how many bottles were sold, how sales varied throughout the year, and where and how bottles were most often sold. The researcher also requested a tour of the facilities to view the treatment and cleaning procedure (Appendix 1)

### **3.3.3 Sample Collection**

Five types of water samples were collected from small packaged water distributors during the study, depending on the consent of the business manager and on the availability of packaged water products for purchase. If available, raw water was collected from source water feeding the treatment system at a water distribution company or from a point along the system prior to any disinfection point. Treated water was sampled directly from the distributor's treatment system. Packaged water products were also sampled and these consisted of: sachets, disposable bottles or reusable containers. Sachets are small, sealed plastic pouches filled with water. These are

intended for single-use and cannot be filled with water again. Disposable bottles were typically available in slightly larger volumes ranging from 500mL to 4L. They were made of a weak plastic, intended for one-time use. The most abundant container types were 20L reusable bottles. They are sold by the water distribution companies, and can be returned and refilled or exchanged for full bottles.

**Tegucigalpa.** In Honduras, eight unique distributors were visited on three occasions with a one-week time interval between visits. At each visit, one sample of crude water and one sample of treated water were collected, and three 20L reusable bottles of water were purchased. Crude and treated water samples were tested on the day of the visit, along with one of the three 20L bottles. The remaining two bottles were tested on day 7 and day 31 following the visit.

**La Paz.** In La Bolivia, eight unique distributors were visited twice each, with a minimum of one week in between visits. Samples were not obtained directly from the purification systems but 20L reusable bottles were purchased from each distributor and sachet water was purchased from six of eight distributors.

**Muisne.** In Muisne, treated water samples were collected, and both small, disposable (4L) bottles and large, reusable (20L) bottles were purchased. Six unique distributors were visited. Four were visited twice each, and two were visited once. At the first visit, researchers conducted an interview with a company representative, collected two samples of treated water directly from the distribution system, and purchased four small (4L) disposable bottles and twelve large (20L) reusable bottles that had been filled with water that same day. Researchers returned to each

distributor a minimum of one week later to collect two additional samples of treated water, and purchase 16 more disposable and reusable bottles—four and twelve, respectively. Disposable and reusable bottles were inspected upon purchase and stored, unopened, away from direct sunlight, in the laboratory where temperature varied from 23.5 – 27<sup>0</sup> C, until testing. Treated water samples were labeled on site, and tested within 3 hours on Day 0. 20L reusable bottles were tested in duplicate 0, 3, 7, 14, 21 and 28 days from the date of collection. Non-reusable bottles were tested 0, 7, 21, and 28 days from the date of collection (Figure 1).

**Tena.** Six unique distributors were visited in Tena. Five were visited twice each, and one was visited one-time only. Sample collection and distributor visits were conducted in Tena as in Muisne with the following change: instead of twelve, ten large (20L) reusable bottles were purchased at each distributor and tested in duplicate on Days 0, 3, 7, 14 and 21.

### **3.3.4 Testing Procedures**

In accordance with the World Health Organization Guidelines for Drinking-water Quality, *E. coli* and total coliforms were used as indicators of fecal contamination and inadequate water treatment, respectively (WHO, 2011). Bottles and sachets were visually inspected for damage, and the water assessed for turbidity and evident contamination, such as mold or insects. A 100mL sample of water from each bottle or sachet was collected and tested per manufacturer's instructions (IDEXX Laboratories, 2015). Briefly, one packet of IDEXX reagent was added to the sample. The sample-reagent mixture was poured into a sterile Quantitray 2000. Trays were sealed using a small iron. Care was taken to ensure that the hot iron did not touch any part of the tray for longer than 10 seconds, to avoid overheating. Samples were incubated at 35±0.5°C for

24 hours. Total coliform and *E. coli* concentrations were determined using the corresponding IDEXX Most Probable Number (MPN) table (IDEXX Laboratories, 2013).

Two negative controls were undertaken for each experiment: one with a 500mL Dasani-brand bottled water, which was available for purchase at all study sites, and one sample of distilled medical-grade water, purchased from a pharmacy. A positive control containing a mixture of sewer and borehole water was performed at the end of each experiment to avoid contaminating testing equipment.

### **3.3.5 Data Analysis**

The data for each sample were entered in Microsoft Excel and imported into SAS 9.4 for further analysis. Data was analyzed using descriptive statistics, including median as the measure of central tendency, and frequency where the main outcome was presence or absence of either total coliform bacteria or *E. coli*. Data was observed for trends in frequency and severity of contamination in relation to storage time. Logistic regression was performed to compare the proportion of treated water contaminated with total coliform or *E. coli* to the proportion of other sample types that were contaminated.

Restricting the analysis to reusable containers, logistic regression was used to compare the proportion of bottles contaminated in Tena (where the highest prevalence of contamination existed) to the proportion of contaminated bottles in other study sites. To determine the effect of the number of interactions between researchers and distributors (i.e. whether repeat visits to the same manufacturer influenced the cleanliness of the samples), logistic regression was used to compare the proportion of contaminated bottles collected on the third visit to the proportion of

contaminated bottles collected on the first visit. Where company survey data was available, each company was classified into exclusive categories for water treatment methods and for bottle cleaning methods. These variables were analyzed as above to determine their associations with total coliform and *E. coli* contamination. Multivariate logistic regression analysis was performed on all categorical variables to produce adjusted odds ratios for total coliform contamination in reusable containers.

### **3.4 Results**

#### **3.4.1 Survey Results**

Surveys were completed at a total of 20 distributors with eight from Tegucigalpa, Honduras, and six each from Muisne and Tena in Ecuador. No surveys or tours were completed during visits to distributors in La Paz. All used carbon activated filtration in their water treatment systems. All but one (95%) used ultraviolet (UV) sterilization, and most companies who used UV in their water treatment systems also used ozone sterilization (75%). Sand filtration was used by all companies in Tegucigalpa, by 66% in Muisne, and by one company in Tena (17%).

Microfiltration was used less frequently in water distribution systems of the companies surveyed (35%). Only one company (5%), based in Tena, used chlorine in its water treatment system.

Ninety-five percent of companies (n=19) indicated some procedure for cleaning reusable bottles. Approximately seventy percent used hot water—either mechanized or manually—to clean bottles and 65% used some kind of detergent. Fifteen percent used only hot water to sanitize bottles. One used only a cold water rinse. Less than a quarter of companies washed bottles with a chlorine solution or used ozone to sterilize reusable bottles (Table 1).



**Table 1. Overview of the 20 companies surveyed in three of four study sites (Tegucigalpa, Muisne and Tena).**

Site	Distributor	Water Treatment Method <sup>1</sup>	Method for Cleaning Containers <sup>2</sup>
Tegucigalpa	A	SF + UV	CW + Detergent
Tegucigalpa	B	SF + MF + UV	CW + Detergent
Tegucigalpa	C	SF + UV	HW only
Tegucigalpa	D	SF only	HW + Detergent
Tegucigalpa	E	SF + MF + UV + Chem	None
Tegucigalpa	F	SF + MF + UV + Chem	HW + Detergent
Tegucigalpa	G	SF + MF + UV + Chem	HW + Detergent
Tegucigalpa	H	SF + MF + UV + Chem	HW + Detergent
Muisne	A	SF + MF + UV + Chem	None
Muisne	B	MF + UV + Chem	HW + Det. + Chlorine
Muisne	C	UV + Chem	HW + Detergent
Muisne	D	SF + MF + UV + Chem	HW + Det. + Chlorine
Muisne	E	SF + MF + UV + Chem	HW + Detergent
Muisne	F	SF + MF + UV + Chem	HW + Detergent
Tena	A	MF + UV + Chem	None
Tena	B	MF + UV	HW + Det. + Chlorine
Tena	C	MF + UV + Chem	CW + Detergent
Tena	D	UV + Chem	HW only
Tena	E	UV + Chem	HW only
Tena	F	SF + UV + Chem	CW only

<sup>1</sup>All distributors surveyed used Carbon Activated Filters to treat water; SF = Sand Filtration, UV = Ultraviolet Irradiation, Chem = Chemical disinfection with Chlorine or Ozone.

<sup>2</sup>CW= Cold Water, HW = Hot Water

### 3.4.2 Detection of Total Coliforms and *E. coli* in water samples

A total of 616 samples were collected or purchased from a total of 28 distributors across the four study sites (Table 2). Overall, the median concentration of total coliform bacteria was 51.9 cfu/100mL (range: <1 – 200.5) in raw water, 19.2 cfu/100mL (range: <1 - >2419.6) in water packaged in reusable bottles, 19.05 cfu/100mL (range: <1 -211.0) in water packaged in disposable bottles, <1 cfu/100mL (max: >2419.6) in sachet water, and <1 cfu/100mL (max: >2419.6) in treated water samples. The median for *E. coli* was <1 cfu/100ml of all types of samples. The median *E. coli* concentration was 0.0 cfu/100mL in raw water (max: 200.5), water packaged in reusable bottles (max: 472.1), water packaged in disposable bottles (max: 0.0), and in treated water samples (max: 154.2) (Table 3).

**Table 2. Water Sample Type by Study Site**

Site (No. of Distributors)	Raw (%)	Treated (%)	Disposable Bottle (%)	Sachet (%)	Reusable Bottle (%)	Total
<b>Tegucigalpa (8)</b>	24 (20.0)	24 (20.0)	0	0	72 (60.0)	120
<b>La Paz (8)</b>	0	0	0	110 (67.9)	52 (32.1)	162
<b>Muisne (6)</b>	0	22 (11.8)	36 (19.4)	0	128 (68.8)	186
<b>Tena (6)</b>	0	22 (14.9)	16 (10.8)	0	110 (17.9)	148
<b>Total</b>	24 (3.9)	68 (11.0)	52 (8.4)	110 (17.9)	362 (58.8)	616

In Tegucigalpa, the median level of total coliform contamination was 0.0. cfu/100mL (max: 25.4) among treated water samples, and 8.7 cfu/mL (range: 0 – 200.5) among samples taken from reusable containers. No *E. coli* was detected in treated water samples collected in Tegucigalpa. The median *E. coli* concentration among samples taken from reusable bottles was 0.0 cfu/100mL (max: 9.9).

**Table 3. Descriptive statistics of total coliform and *E. coli* concentration in each sample type.**

<b>Parameter</b>	<b>Median<sup>1</sup></b>	<b>IQR</b>	<b>Max.</b>	<b>Min.</b>
<b><i>Raw Water (n=24)</i></b>				
Total coliform	51.9	[159.5 – 12.7]	200.5	<1
<i>E. coli</i>	<1	[0.5 – <1]	65.9	<1
<b><i>Treated Water (n= 68)</i></b>				
Total coliform	<1	[2.6 – <1]	2419.6	<1
<i>E. coli</i>	<1	0.0	154.2	<1
<b><i>Disposable Bottle (n=52)</i></b>				
Total coliform	<1	[5.8 – 0.0]	211	<1
<i>E. coli</i>	<1	NA	<1	<1
<b><i>Sachet Water (n=110)</i></b>				
Total coliform	<1	0.0	2419.6	<1
<i>E. coli</i>	<1	0.0	<1	<1
<b><i>Reusable Bottle (n=362)</i></b>				
Total coliform	19.2	[203.7 – <1]	2419.6	<1
<i>E. coli</i>	<1	0.0	472.1	<1

<sup>1</sup>Total coliform bacteria and *E. coli* concentrations are present in units of CFU per 100ml

In Muisne, the median total coliform concentration was <1 cfu/100mL in treated water samples (max: 7.5) and in samples taken from disposable bottles (max: 178.2); it was 23.1 cfu/100mL among samples taken from reusable bottles (range: 0 – >2419.6). No *E. coli* was detected in treated water samples or in samples from disposables bottles collected in Muisne. The median *E. coli* contamination in samples taken from reusable bottles was 0.0 cfu/100mL (max: 24.9).


In Tena, the median total coliform concentration was 13.2 cfu/100mL in treated water samples (max: >2419.6) and 0.0 cfu/100mL among reusable bottle samples (max: 108.9); it was 121.3 cfu/100mL among samples taken from non-reusable bottles (max: >2419.6). The median *E. coli* contamination was 0.0 cfu/100mL in treated water samples (max: 154.2), and in samples taken from reusable containers (max: 472.1). No *E. coli* were detected in samples taken from disposable bottles. In La Paz, the median level of coliform contamination was 0.0 cfu/100mL among samples taken from reusable containers (max: >2419.6).

Total coliform bacteria were detected in 311 (50.5%) water samples. Total coliform bacteria was most frequently detected in raw water (22/24; 91.7%) and in samples taken from reusable bottles (243/362; 67.1%). Sachet and treated water were contaminated with total coliform bacteria least frequently (10.0% and 27.9%, respectively) (Figure 2). *E. coli* contamination was found in 45 (7.3%) water samples. *E. coli* was most frequently found in raw water samples (6/24; 25.0%) and in samples taken from reusable bottles (35/362; 9.7%). No *E. coli* were detected in samples taken from non-reusable containers, such as disposable bottles (0/52) or sachet water (0/110) (Figure 2). Ten of twenty companies sampled at the point of treatment were found to have contamination with total coliforms even after treatment. All samples from reusable bottles that were purchased from these companies were contaminated with total coliforms as well. Of the ten companies whose treated water was free from total coliform contamination, nine samples of water from reusable bottles were contaminated with total coliforms (Table 4). No noticeable degradation in water quality was observed between samples tested on the day of the visit and those tested after being stored for up to 28 days (Table 5).

**Table 4.** Associations of company-treated water and coliform contamination

<b>Quality of Reusable Bottles Sold</b>	<b>Total Coliform Contamination in Treated Water (%)</b>	<b>No Total Coliform Contamination Detected in Treated Water (%)</b>	<b>Total</b>
<b>Total Coliform Contamination Detected</b>	10 (35.7)	9 (53.6)	19
<b>No Contamination Detected</b>	0 (0.0)	1 (10.7)	1
<b>Total</b>	10	10	20

**Figure 2.** Percent of samples contaminated with Total Coliforms and *E. coli* by site and type



<sup>a</sup> Non-reusable water sources include sachet water and disposable bottles.

<sup>b</sup> La Paz-No data for treated water available

<sup>c</sup> Tegucigalpa-No data for reusable bottles available

\*Error bars represent 95% Confidence Intervals

### 3.4.3 Results of Analyses

#### *Association between sample type and Total Coliform or E. coli contamination*

Raw, untreated water and water packaged in reusable bottles were significantly more likely to be contaminated with total coliforms than treated water taken directly from purification systems (ORs= 28.4 and 5.3, respectively). Sachet water was least likely to be contaminated with total coliform bacteria (OR = 0.3 compared to treated water) (Table 6).

**Table 5. Quality of water by post-visit time sampling time point and sample type**

Sample Type	Sampling	n	Total Coliform	
	Time Point		Detected (%)	<i>E. coli</i> Detected (%)
Treated	0	68	19 (27.9)	4 (5.8)
Sachet	0	52	7 (13.4)	0 (0)
	3	28	3 (10.7)	0 (0)
	7	30	1 (3.3)	0 (0)
Disposable Bottles	0	13	4 (30.7)	0 (0)
	7	13	3 (23.0)	0 (0)
	14	4	1 (25.0)	0 (0)
	21	13	4 (30.7)	0 (0)
	28	9	3 (33.3)	0 (0)
Reusable	0	120	58 (48.3)	10 (8.3)
	3	44	36 (81.8)	6 (13.6)
	7	68	58 (85.2)	7 (10.2)
	14	44	33 (75.0)	7 (15.9)
	21	44	36 (81.8)	4 (9.0)
	28	42	22 (52.3)	1 (5.0)

<sup>1</sup>Post-visit sampling time point is in indication of the number of days since the sample was purchased or collected from the distributor. Time-point 0 indicates that the sample was tested on the same day as purchased.

No *E. coli* were detected in any samples taken from disposable bottles or sachet water. Raw, untreated water was most likely to be contaminated with *E. coli* (OR = 5.3 compared to treated water). *E. coli* contamination was found more frequently in reusable bottles than in treated water, but the resulting OR (OR= 1.7) was not statistically significant (95% CI: 0.6 – 5.0) (Table 7).

#### *Factors Associated with Coliform contamination of Reusable Bottles in Univariate Analyses*

When restricting univariate analyses to samples taken from reusable bottles (n=362) as these were the only sample type collected in all study sites and from all distributors, study site was significantly associated with total coliform contamination (p<0.01). Water packaged in reusable

containers was most likely to be contaminated with total coliforms in Tena and least likely to be contaminated in La Paz (OR= 0.04 [0.0- 0.1]).

**Table 6. Association between sample type and prevalence of total coliform contamination**

Sample Type	Total Coliform			OR (95% CI)
	Present	Absent	Total	
Raw Water	22 (91.7)	2 (8.3)	24 (100)	28.4 (6.1 -132.5)
Treated Water	19 (27.9)	49 (72.1)	68 (100)	<i>ref.</i>
Disposable Bottle	16 (30.8)	36 (69.2)	52 (100)	1.1 (0.5 - 2.5)
Sachet Water	11 (10.0)	99 (90.0)	110 (100)	0.3 (0.1 - 0.7)
Reusable Bottle	243 (67.1)	119 (32.9)	362 (100)	5.3 (2.9- 9.3)
Total	311 (50.5)	305 (49.5)	616 (100)	

The treatment methods used in the water distribution system were not significantly associated with total coliform contamination of reusable bottles, but bottle cleaning methods were. Water sampled from reusable bottles that were purchased from distributors who used only hot water to clean bottles were more likely to be contaminated with total coliforms than those that were cleaned with a combination hot water, detergent and chlorine bleach (OR=6.3 [1.9 – 19.9]). Total coliform contamination was found in all reusable bottles that were cleaned with only cold water (OR= >999.9), but due to this complete separation of data the finding is not significant (Table 8). When samples taken from reusable bottles purchased from distributors whose treated water samples tested positive for contamination were removed from analyses, no bottle cleaning methods remained significant (p=0.4).

*Factors Associated with E.coli contamination of Reusable Bottles in Univariate Analyses*

In univariate analyses, water treated using carbon filtration, UV irradiation and chemical disinfection was significantly more likely to be contaminated with *E. coli* than water treated with these methods plus sand and membrane filtration (OR= 16.3[4.6 – 57.9]). Similar to total

coliform contamination, bottles cleaned with HW only were significantly more likely to be contaminated with *E. coli* than those cleaned with HW, detergent and chlorine (OR= 10.3 [2.8 – 37.7]) (Table 9). When samples taken from reusable bottles purchased from distributors whose treated water samples tested positive for contamination were removed from analyses, no bottle cleaning methods remained significant (p=0.7).

**Table 7. Association between sample type and prevalence of *E. coli* contamination.**

Sample Type	<i>E. coli</i>			OR (95% CI)
	Present	Absent	Total	
Raw Water	6 (25.0)	18 (75.0)	24 (100)	5.3 (1.4 - 20.9)
Treated Water	4 (5.9)	64 (94.1)	68 (100)	<i>ref.</i>
Reusable Bottle	35 (9.7)	327 (90.3)	362 (100)	1.7 (0.6 - 5.0)

No *E. coli* was found in any samples collected from disposable bottles or sachet water.

### *Results of Multivariate Analyses*

There was no survey data available for companies visited in La Paz, so the multivariate model was restricted to reusable bottles purchased in Tegucigalpa, Tena, and Muisne. Tena remained the most likely site for contamination of total coliform contamination, compared to Tegucigalpa which was the least likely to be contaminated (aOR = 0.03 [0.0 – 0.5]). No other factors remained significant in multivariate analysis (Table 8). Multivariate analysis was not conducted for *E. coli* contamination due to the low prevalence of *E. coli* across study sites which contributed to a questionable validity of the multivariate model.



**Table 8. Results of Univariate and Multivariate Analyses for Presence of Total Coliform Bacteria Among Reusable Bottles**

Exposure	No Total Coliforms Detected (%)	Total Coliforms Detected (%)	OR [95% CI]	<i>p</i>	aOR <sup>1</sup> [95%CI]	<i>p</i>
<b>Site</b>						
Tegucigalpa	24 (33.3)	48 (19.75)	0.4 [0.2 - 0.8]	0.3	0.0[0.0 - 0.5]	<b>&lt;0.01</b>
La Paz	43 (82.7)	9 (17.3)	0.0 [0.0 - 0.1]	<0.01	--	--
Muisne	34 (26.6)	94 (73.4)	0.5 [0.3 -0.9]	<0.01	0.3[0.0 - 1.5]	0.1
Tena	18 (16.4)	92 (83.6)	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
<b>Site Visit</b>						
1st Visit	52 (28.6)	97( 71.4)	1.3 [0.5 - 3.1]	0.3	1.5[0.5 - 4.6]	0.1
2nd Visit	59 (37.8)	97 (62.2)	0.8 [0.3 - 2.0]	0.3	0.7[0.2 - 2.2]	0.1
3rd Visit	8 (33.3)	16 (66.7)	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
<b>Bottle Damage (n=17)</b>	5 (29.4)	12 (70.6)	1.2 [0.4 - 3.4]	1	0.6 0.1 - 2.8]	0.5
<b>Bottle Cleaning<sup>1</sup></b>						
None	20 (37.7)	33 (62.3)	0.9 [0.4 - 2.0]	0.8	0.6[0.2 - 1.9]	0.4
CW only <sup>2</sup>	0 (0.0)	10 (100.0)	--	--	--	--
HW only	4 (8.2)	45 (91.8)	6.3[1.9 - 19.9]	<0.01	4.1[0.3-54.4]	0.3
CW + Detergent	5 (13.2)	33 (86.8)	3.7[1.2 - 10.9]	0.02	2.3[0.3-21.2]	0.5
HW + Detergent	27 (26.0)	77 (74.0)	1.6 [0.8 - 3.2]	0.2	2.4 [0.8 -7.3]	0.1
HW + Detergent + Chlorine	20 (35.7)	36 (64.3)	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
<b>Water Treatment<sup>1</sup></b>						
SF	3 (33.3)	6 (66.7)	0.8[0.2- 3.5]	0.8	1.2[0.3 - 5.8]	0.8
SF + UV	4 (22.2)	14 (77.8)	1.5 [0.5 - 4.8]	0.5	1.4[0.2-12.9]	0.8
MF + UV	9 (45.0)	11(55.0)	0.5 [0.2 - 1.3]	0.2	0.1[0.0 - 1.3]	0.08
UV + Chem	10 (15.6)	54 (84.4)	2.3 [1.1 - 4.9]	0.03	0.3[0.1 - 1.0]	0.06
SF + MF + UV	1 (11.1)	8 (88.9)	3.4[0.4 - 27.9]	0.3	3.6[0.2 - 67.9]	0.4
MF + UV + Chem	11(21.2)	41 (78.9)	1.6 [0.7 - 3.4]	0.2	0.4 [0.1 - 1.8]	0.3
SF + UV + Chem <sup>2</sup>	0 (0.0)	10 (100.0)	--	--	--	--
SF + MF + UV + Chem	38 (29.7)	90 (70.3)	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>

<sup>1</sup>No surveys were conducted at water companies located in La Paz; reusable bottles collected and sampled in La Paz are excluded from these analyses.

<sup>2</sup>All bottles cleaned with CWO were filled with SF + UV + Chem treated water. These ten bottles were collected from a single distributor and all were contaminated with total coliform bacteria.

### 3.5 Discussion

In this pilot study, we were able to describe and identify substantial differences across 20 packaged water distributors in 4 sites in Central and South America. The quality of packaged water varied significantly between reusable and disposable containers, and between reusable containers purchased in different study sites. More than half of the packaged water examined in this study was contaminated with total coliform bacteria. Twenty liter reusable bottles were

available for purchase at every distributor in all study sites, illustrating their ubiquity. The majority of reusable containers were contaminated with total coliform contamination, and *E. coli* was detected in the bottles sold in three of the four study sites.

**Table 9. Results of Univariate Analyses for Presence of *E. coli* Among Reusable Bottles**

Exposure	No <i>E. coli</i> Detected (%)	<i>E. coli</i> Detected (%)	OR [95% CI]	<i>p</i>
<b>Site<sup>1</sup></b>				
Honduras	68 (94.4)	4 (5.6)	0.2 [0.1 - 0.6]	0.9
La Paz	52 (100)	0 (0)	---	---
Muisne	125 (97.7)	3 (2.3)	0.1 [0.0 - 0.3]	0.9
Tena	84 (76.4)	26 (23.6)	<i>ref.</i>	<i>ref.</i>
<b>Site Visit</b>				
1st Visit	164 (90.1)	18 (9.9)	1.2 [0.3 - 5.6]	0.7
2nd Visit	143 (91.7)	13 (8.3)	1 [0.2 - 4.7]	0.9
3rd Visit	22 (91.7)	2 (8.3)	<i>ref.</i>	<i>ref.</i>
<b>Bottle Damage (n=17)</b>	16 (94.1)	1 (5.9)	0.6 [0.1 - 4.8]	1
<b>Bottle Cleaning<sup>1</sup></b>				
None	52 (98.1)	1 (1.9)	0.3 [0.0 - 3.4]	0.4
CW only	8 (80)	2 (20)	4.4 [0.6 - 30.6]	0.1
HW only	31 (63.2)	18 (36.7)	10.3 [2.8 - 37.7]	<0.01
CW + Detergent	32 (84.2)	6 (15.8)	3.3 [0.8 - 14.2]	0.1
HW + Detergent	101 (97.1)	3 (2.8)	0.5 [0.1 - 2.7]	0.4
HW + Detergent + Chlorine	53 (94.7)	3 (5.4)	<i>ref.</i>	<i>ref.</i>
<b>Water Treatment<sup>1</sup></b>				
SF	8 (88.9)	1 (11.1)	5.2 [0.5 - 55.9]	0.2
SF + UV	17 (94.4)	1 (5.6)	2.5 [0.2 - 24.9]	0.4
MF + UV	19 (95.0)	1(5.0)	2.2 [0.2 - 22.2]	0.5
UV + Chem	46 (14.9)	18 (5.9)	16.3 [4.6 - 57.9]	<0.01
SF + MF + UV	8 (88.9)	1 (11.1)	5.2 [0.5 - 55.9]	0.2
MF + UV + Chem	46 (88.5)	6 (11.5)	5.4 [1.3 - 22.6]	0.02
SF + UV + Chem	8 (80.0)	2 (20.0)	10.4 [1.5 - 71.5]	0.02
SF + MF + UV + Chem	125 (97.7)	3 (2.3)	<i>ref.</i>	<i>ref.</i>

Multivariate analyses were not conducted for *E. coli* due to the low prevalence of *E. coli* contamination across study sites and sample types which contributed the questionable validity of the multivariate model.

<sup>1</sup>No *E. coli* was detected in any sample collected in La Paz and no surveys were conducted at water companies located in this study site; reusable bottles collected and sampled in La Paz are excluded from these analyses.

The treated water samples tested from half of distributors were free from total coliform contamination (n= 10, 50%). Yet of the ten companies whose treated water samples were free from total coliform contamination, nine (90%) of them sold bottles that were positive for total coliform contamination. Based on these observations, it is likely that the primary source of contamination found in water packaged in reusable bottles stems from the bottles themselves, and may be due to inadequate disinfection between uses. The present study found no link between particular bottle cleaning methods and water contamination; although there is some evidence to suggest that the addition of chlorine and detergent to hot water cleaning protocols can prevent contamination. However, data collection assessed bottle cleaning method by a single self-report survey item that did not evaluate adherence to reported cleaning protocols, so it remains unclear whether adherence to these protocols is consistent. It is recommended in future research to collect data on bottle cleaning methods using several survey items that will evaluate adherence to protocols as well.

Total coliforms are indicator organisms, and their presence in treated water can be used to determine if water treatment processes were adequate to destroy bacterial pathogens. It is possible for water containing total coliform bacteria to be free of human pathogens; however, it is unlikely that a water sample would contain pathogenic bacteria and test negative for total coliform bacteria. *E. coli* are also indicator organisms, and can be used to determine whether drinking water has been exposed to a source of fecal contamination such that there could be pathogenic organisms present. The drinking water guidelines set forth by WHO dictate that there should be no detectable *E. coli* per 100mL of water intended for drinking, including packaged or bottled water (WHO, 2011). Despite the added expense and level of trust that consumers have,

the water bottled in these study sites demonstrated coliform and E. coli contamination levels that may pose a substantial health risk to consumer (Bain, 2014). These results may have implications for national or international policies that regulate private water enterprises, and can inform guidelines for packaged water distributors in particular.

The elaborate bottle cleaning techniques used by large bottled water providers that operate in higher income countries may not be feasible for the small distributors that operated in the studied settings. Even when the appropriate technologies are in places, failure in protocol adherence can occur. For example, companies that operate in the U.S. clean their large reusable containers vigorously, and reuse them a certain number of times-- up to 35-- before recycling them (Nestle Waters North American Inc., 2010). In resource poor settings, bottles may be reused more times than manufacturer recommendations in order to save money and resources.

### **3.6 Limitations of the Research**

This study has several limitations. While the study was conducted in four distinct geographical regions in attempt to gain a more representative sample of LMICs in Central and South America, the heterogeneity of packaged water available for purchase in each study site limited the bulk of the analyses to reusable 20L containers. Furthermore, the quality of water differed significantly between companies within each site. Therefore, little can be said regarding how or if the quality of packaged water varies between regions. Additionally, the study was conducted during a single time of year, and samples were collected from only a fraction of the private bottled water enterprises that operate throughout these regions. Finally, the study does not account for the

different kinds of storage and transportation conditions that can occur between bottling and point of use.

## Manuscript References

- Bain, R., Cronk, R., Hossain, R., Bonjour, S., Onda, K., Wright, J., ... Bartram, J. (2014). Global assessment of exposure to faecal contamination through drinking water based on a systematic review. *Tropical Medicine & International Health*, 19(8), 917–927.  
<http://doi.org/10.1111/tmi.12334>
- Dada, A. (2011). Packaged water: optimizing local processes for sustainable water delivery in developing nations. *Globalization & Health*, 7(1), 24–31. <http://doi.org/10.1186/1744-8603-7-24>
- Datamonitor. (2004). Bottled Water Industry Profile: Latin America (p. 1). MarketLine, a Progressive Digital Media business. Retrieved from  
<http://ezproxy.gsu.edu/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=27275369&site=eds-live>
- de Queiroz, J. T. M., Doria, M. de F., Rosenberg, M. W., Heller, L., & Zhouri, A. (2013). Perceptions of bottled water consumers in three Brazilian municipalities. *Journal of Water and Health*, 11(3), 520–531. <http://doi.org/10.2166/wh.2013.222>
- Espinosa-García, A. C., Díaz-Ávalos, C., González-Villarreal, F. J., Val-Segura, R., Malvaez-Orozco, V., & Mazari-Hiriart, M. (2015). Drinking water quality in a Mexico city university community: perception and preferences. *EcoHealth*, 12(1), 88–97.  
<http://doi.org/10.1007/s10393-014-0978-z>
- Falcone-Dias, M. F., Emerick, G. L., & Farache-Filho, A. (2012). Mineral water: a microbiological approach. *Water Science & Technology: Water Supply*, 12(5), 556–562.  
<http://doi.org/10.2166/ws.2012028>
- Fisher, M. B., Williams, A. R., Jalloh, M. F., Saqee, G., Bain, R. E. S., & Bartram, J. K. (2015).

- Microbiological and Chemical Quality of Packaged Sachet Water and Household Stored Drinking Water in Freetown, Sierra Leone. *PLOS ONE*, 10(7), e0131772.  
<http://doi.org/10.1371/journal.pone.0131772>
- Halage, A. A., Ssemugabo, C., Ssemwanga, D. K., Musoke, D., Mugambe, R. K., Guwatudde, D., & Ssempebwa, J. C. (2015). Bacteriological and physical quality of locally packaged drinking water in Kampala, Uganda. *Journal of Environmental and Public Health*, 2015, 942928. <http://doi.org/10.1155/2015/942928>
- Hatt, L. E., & Waters, H. R. (2006). Determinants of child morbidity in Latin America: A pooled analysis of interactions between parental education and economic status. *Social Science & Medicine*, 62(2), 375–386. <http://doi.org/10.1016/j.socscimed.2005.06.007>
- IDEXX Laboratories. (2013). Quanti-Tray/2000 Insert and Most Probable Number (MPN) Table. Retrieved from <https://www.idexx.com/resource-library/water/quant-tray-2000-procedure-en.pdf>
- IDEXX Laboratories. (2015). Colilert. Retrieved from <https://www.idexx.com/resource-library/water/colilert-procedure-en.pdf>
- Impellitteri, C. (2007). US EPA Small Drinking Water Systems: State of the Industry and Treatment Technologies to Meet the Safe Drinking Water Act Requirements September 2007 - lps100419.pdf. Retrieved April 2, 2016, from <http://permanent.access.gpo.gov/lps100419/lps100419.pdf>
- Informe anual bebidas 2015. (2015). *Industria Alimenticia*, 26(8), 12–19.
- Jain, M., Lim, Y., Arce-Nazario, J. A., & Uriarte, M. (2014). Perceptual and socio-demographic factors associated with household drinking water management strategies in rural Puerto Rico. *PloS One*, 9(2), e88059. <http://doi.org/10.1371/journal.pone.0088059>

- J. Liu, N. d&#39;Ozouville. (2013). Water contamination in Puerto Ayora: Applied interdisciplinary research using Escherichia coli as an indicator bacteria. Galapagos Report 2011-2012, 76–83.
- Kjellen, M., & McGranahan, G. (2006). Informal Water Vendors and the Urban Poor. International Institute for Environment and Development (IIED). Retrieved from <http://pubs.iied.org/pdfs/10529IIED.pdf>
- Madeline Baer. (2015). From Water Wars to Water Rights: Implementing the Human Right to Water in Bolivia. *Journal of Human Rights*, 14(3), 353–376.  
<http://doi.org/10.1080/14754835.2014.988782>
- Malkin, E. (2012, July 16). Mexicans Struggle to Kick Bottled-Water Habit. *The New York Times*. Retrieved from <http://www.nytimes.com/2012/07/17/world/americas/mexicans-struggle-to-kick-bottled-water-habit.html>
- Marzano, M. a., Ripamonti, B., & Balzaretto, C. m. (2011). Short communication: Monitoring the bacteriological quality of Italian bottled spring water from dispensers. *Food Control*, 22, 333–336. <http://doi.org/10.1016/j.foodcont.2010.06.014>
- Nestle Waters North American Inc. (2010). Poland Spring Water Quality Report. Stamford: Nestle Waters.
- Opryszko, M. C., Huang, H., Soderlund, K., & Schwab, K. J. (2009). Data gaps in evidence-based research on small water enterprises in developing countries. *Journal of Water and Health*, 7(4), 609–622. <http://doi.org/10.2166/wh.2009.213>
- Pacheco-Vega, R. (2015). Agua embotellada en México: de la privatización del suministro a la mercantilización de los recursos hídricos. *Bottled Water in Mexico: From Privatization of Supply to Marketization of Water Resources.*, 22(63), 221–263.



WHO & UNICEF. (2012). Progress on Drinking Water and Sanitation: 2012 Update. Joint Monitoring Programme for Water Supply and Sanitation. Retrieved from <http://www.unicef.org/media/files/JMPReport2012.pdf>

Williams, A. R., Bain, R. E. S., Fisher, M. B., Cronk, R., Kelly, E. R., & Bartram, J. (2015). A Systematic Review and Meta-Analysis of Fecal Contamination and Inadequate Treatment of Packaged Water. PLoS ONE, 10(10), 1–27. <http://doi.org/10.1371/journal.pone.0140899>

World Health Organization. (2011). Guidelines for Drinking Water Quality, Fourth Edition. WHO. Retrieved from [http://apps.who.int/iris/bitstream/10665/44584/1/9789241548151\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44584/1/9789241548151_eng.pdf)

## Chapter 4: Conclusion

A key finding from this study is that water packaged in reusable containers is significantly more likely to be contaminated than water packaged in other containers, including disposable bottles and sachets. Furthermore, the contamination likely stems from inadequate cleaning of the bottles between uses. This finding is supported by a small number of studies comparing large 20 liter reusable containers to other packaged water types in the body of literature on packaged water quality (Falcone-Dias, 2012; Marzano, 2013; Levesque, 1994). Previous studies have reported on the quality of sachet water, and have found it to be poor—perhaps owing to the fact that sachet water is more of an informal industry and perhaps less often scrutinized by regulatory frameworks (Dada, 2011; Stoler, 2012). This study somewhat contradicts those findings. No fecal contamination was found in any of the sachet water samples, but sachet water was only available for purchase in a single study site, so this discrepancy cannot be addressed with the current data available.

This study reports on packaged water quality in two rural areas in Ecuador, which contributes to the body of literature on packaged water quality which contains few studies conducted in rural areas (Williams et al., 2015). Furthermore, this study highlights the heterogeneity that exists in terms of types of water sold, water treatment systems, and sanitizing protocols among SWEs in Central and South America.

Reusable containers are vulnerable to contamination with total coliform bacteria and *E. coli*, even when filled with clean water. These results may have implications for national or international policies that regulate private water enterprises, and can inform guidelines for

packaged water distributors in particular. For regulating bodies to ignore packaged water's role in filling gaps in access to improved drinking water systems is to allow unregulated and—and perhaps unaware—private enterprises to distributed contaminated water. Further research is needed to identify optimal cleaning methods for reusable containers that are practical for use in lower resource settings.

## 5. Appendices

**APPENDIX A:** Survey questions administered to the business managers at each bottled water distributor.



### Survey Questions (Spanish)

#### *Información básica*

- ¿Cuándo su empresa comenzará a funcionar?
- ¿Cuántos bidones se venden por semana?
- ¿Cómo varían la ventas según la temporada?
- ¿Dónde venden botellas?
- ¿Cuánto venden sus jarras de agua 20L para que los consumidores / empresas?

#### *Información de la empresa*

Hábleme de cómo opera su empresa (por ejemplo, obtiene y purifica el agua). (Nota para el topógrafo - esta pregunta es para minimizar el sesgo de composición abierta.)

Si necesita, use estas sondas: Fuente de agua, proceso de tratamiento (RO, ozono, UV, carbón activado, filtros de sedimentos, cloro, sulfato de aluminio, de reducción de la dureza, etc.) proceso de distribución, proceso de saneamiento jarra, proceso de contratación, el número de empleados

- ¿Cómo ha cambiado la demanda de agua embotellada en los últimos 5-10 años?
- ¿Cómo cree que la demanda de agua embotellada va a cambiar en el futuro?

#### *Información interactiva*

- ¿Puedo tener un tour?
- Nota prácticas de higiene.
- ¿Puedo tener una muestra de su agua purificada directamente de su sistema?