

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Theses, Dissertations, and Student Research:
Department of Psychology

Psychology, Department of

7-2011

Exploratory Analyses Of A Developmental Conceptualization of Insight And Treatment Outcomes Of Individuals With Serious Mental Illness in Psychiatric Rehabilitation

Ashley R. Wynne

University of Nebraska-Lincoln, awynne6@yahoo.com

Follow this and additional works at: <http://digitalcommons.unl.edu/psychdiss>



Part of the [Clinical Psychology Commons](#), and the [Psychiatry and Psychology Commons](#)

Wynne, Ashley R., "Exploratory Analyses Of A Developmental Conceptualization of Insight And Treatment Outcomes Of Individuals With Serious Mental Illness in Psychiatric Rehabilitation" (2011). *Theses, Dissertations, and Student Research: Department of Psychology*. 34.

<http://digitalcommons.unl.edu/psychdiss/34>

This Article is brought to you for free and open access by the Psychology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Theses, Dissertations, and Student Research: Department of Psychology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

EXPLORATORY ANALYSES OF A DEVELOPMENTAL
CONCEPTUALIZATION OF INSIGHT AND TREATMENT
OUTCOMES OF INDIVIDUALS WITH SERIOUS MENTAL ILLNESS
IN PSYCHIATRIC REHABILITATION

by

Ashley R. Wynne

A DISSERTATION

Presented to the Faculty of
The Graduate College at the University of Nebraska
In Partial Fulfillment of Requirements
For the Degree of Doctor of Philosophy

Major: Psychology

Under the Supervision of Professor William D. Spaulding

Lincoln, Nebraska

July, 2011

EXPLORATORY ANALYSES OF A DEVELOPMENTAL CONCEPTUALIZATION
OF INSIGHT AND TREATMENT OUTCOMES OF INDIVIDUALS WITH SERIOUS
MENTAL ILLNESS IN PSYCHIATRIC REHABILITATION

by

Ashley R. Wynne, Ph.D.

University of Nebraska, 2011

Adviser: William D. Spaulding

The purpose of the present study was to further examine the relationship between adolescent psychiatric pathology and SMI by assessing the relationship between prior mental health services before the age of 18 and time of assessment on people's insight into their illnesses. A secondary relationship between adolescent psychiatric pathology and functioning in a variety of domains before, during, and after treatment was assessed. Overall, there was an inconsistent pattern of results and partial support of hypotheses. The current study was a retrospective longitudinal study in which assessments were given to 308 participants in an inpatient psychiatric rehabilitation unit every 6 months. Results indicated that those with service use during adolescence were younger at admission to CTP, have slightly lower levels of education, had an earlier age of onset, and significantly more previous hospitalizations. A relationship between Axis I and II diagnoses and service use was also identified. Improvements in neurocognitive, sociocognitive, insight, and behavioral functioning measures are evident over the course of treatment, however are not consistent for all groups and all measures used within this

study. Contrary to the hypothesis, those adults with service usage in adolescence did not endorse differing overall insight or ability to relabel symptoms scores over the course of treatment. However, as hypothesized, there were no differences between those with and without service usage in adolescence on any measure of insight after one year of psychiatric rehabilitation. In general, the CTP participants endorsed lower insight into need for treatment scores across treatment. Also, analyses revealed no significant relationship between whether or not someone used services in adolescence or APP severity level and rate of rehospitalization or discharge location restrictiveness. In summary, this study was exploratory in nature and inconsistent results and mixed support of hypotheses was found. This field of research has numerous implications for increasing insight and bettering outcomes for persons with SMI.

TABLE OF CONTENTS

LIST OF TABLES	v
LIST OF FIGURES	XI
DEDICATION	XIV
ACKNOWLEDGEMENTS	XV
CHAPTER 1 6 INTRODUCTION AND LITERATURE REVIEW	1
<i>Definitions of Insight</i>	1
<i>Etiology of Impaired Insight</i>	4
<i>Insight and Suicide Risk</i>	7
<i>The Role of Insight in Treatment and Recovery</i>	8
<i>Insight and Human Development</i>	9
<i>Present Study</i>	16
CHAPTER 2 - METHOD	22
<i>Design Overview</i>	22
<i>Setting</i>	22
<i>Participants</i>	24
<i>Measures</i>	25
<i>Demographic and Clinical Characteristics</i>	26
<i>Neurocognition Measures</i>	26
<i>Social Cognition Measures</i>	30
<i>Insight Measures</i>	31
<i>Behavioral Functioning Measures</i>	33
<i>Symptomatology Measures</i>	34
<i>Outcome Measures</i>	36
<i>Procedure</i>	37
<i>Data Collection</i>	37
<i>Data Cleaning</i>	42
<i>Data Analysis</i>	43
CHAPTER 3 6 RESULTS.....	45
<i>Demographic and Clinical Characteristics of Participants of CTP</i>	45
<i>Hypothesis 1</i>	47
<i>Functioning at admission</i>	48
<i>Neurocognitive functioning</i>	48
<i>Social cognitive functioning</i>	49
<i>Insight functioning</i>	50
<i>Behavioral functioning</i>	52
<i>Symptomatology</i>	54
<i>Functioning over the course of treatment</i>	59
<i>Neurocognitive functioning</i>	60
<i>Social cognitive functioning</i>	77
<i>Insight functioning</i>	96
<i>Behavioral functioning</i>	107
<i>Symptomatology</i>	122
<i>Hypothesis 2</i>	139
<i>Outcome</i>	139
<i>Discharge location</i>	140
<i>Rehospitalization rate</i>	140
CHAPTER 4 6 DISCUSSION	144
<i>General Discussion</i>	144
<i>Limitations of the Present Study</i>	156
<i>Future Directions</i>	158
REFERENCES	161
APPENDIX	171

LIST OF TABLES

- TABLE 1. *DEMOGRAPHIC AND CLINICAL CHARACTERISTICS OF PARTICIPANTS (N=308)*
- TABLE 2. *DEMOGRAPHIC AND CLINICAL CHARACTERISTICS OF PARTICIPANTS ACCORDING TO SERVICE USE DURING ADOLESCENCE (N=308)*
- TABLE 3. *DEMOGRAPHIC AND CLINICAL CHARACTERISTICS OF PARTICIPANTS ACCORDING TO APP SEVERITY LEVEL (N=308)*
- TABLE 4. *RELATIONSHIP BETWEEN AXIS I DIAGNOSIS AND SERVICE USE (N=282)*
- TABLE 5. *RELATIONSHIP BETWEEN AXIS II DIAGNOSIS AND SERVICE USE (N = 284)*
- TABLE 6. *RELATIONSHIP BETWEEN RACE AND SERVICE USE (N = 288)*
- TABLE 7. *RELATIONSHIP BETWEEN MARITAL STATUS AND SERVICE USE (N = 272)*
- TABLE 8. *MEAN SCORES AND STANDARD DEVIATIONS FOR DEMOGRAPHIC AND CLINICAL VARIABLES AS A FUNCTION OF SERVICE USE*
- TABLE 9. *BIVARIATE CORRELATIONS FOR NEUROCOGNITIVE VARIABLES*
- TABLE 10. *SERVICE USE BY NEUROCOGNITIVE VARIABLES AT ADMISSION ONE-WAY ANALYSES OF VARIANCE (ANOVAS)*
- TABLE 11. *APP SEVERITY LEVEL BY NEUROCOGNITIVE VARIABLES AT ADMISSION ANOVAS*
- TABLE 12. *BIVARIATE CORRELATIONS FOR SOCIALCOGNITIVE VARIABLES*
- TABLE 13. *SERVICE USE BY THEORY OF MIND, SOCIAL COGNITION AT ADMISSION ANOVAS*
- TABLE 14. *APP LEVEL BY THEORY OF MIND AND SOCIALCOGNITIVE FUNCTIONING AT ADMISSION ANOVAS*
- TABLE 15. *BIVARIATE CORRELATIONS FOR INSIGHT VARIABLES*
- TABLE 16. *SERVICE USE BY INSIGHT MEASURES AT ADMISSION ANOVAS*
- TABLE 17. *APP LEVEL BY INSIGHT MEASURES AT ADMISSION ANOVAS*
- TABLE 18. *BIVARIATE CORRELATIONS OF NOSIE SUBSCALES*
- TABLE 19. *SERVICE USE BY NOSIE SCALE SCORES AT ADMISSION ANOVAS*
- TABLE 20. *APP LEVEL BY NOSIE SCALE SCORES AT ADMISSION ANOVAS*
- TABLE 21. *BIVARIATE CORRELATIONS OF BPRS TOTAL SCORES, BPRS FACTOR SCORES, AND SYMPTOMATOLOGY MEASURES*
- TABLE 22. *SERVICE USE BY BPRS FACTOR SCORES, BPRS TOTAL, AND SYMPTOMATOLOGY MEASURES AT ADMISSION ANOVAS*
- TABLE 23. *APP LEVEL BY BPRS FACTOR SCORES, BPRS TOTAL, AND SYMPTOMATOLOGY MEASURES AT ADMISSION ANOVAS*
- TABLE 24. *SERVICE USE BY BPRS ITEMS AT ADMISSION ANOVAS*
- TABLE 25. *APP SEVERITY LEVEL BY BPRS ITEMS AT ADMISSION ANOVAS*
- TABLE 26. *SERVICE USE BY WCST REPEATED MEASURES ANOVAS*
- TABLE 27. *SERVICE USE BY NEUROCOGNITION MEASURES REPEATED MEASURES ANOVAS*
- TABLE 28. *SERVICE USE BY RCFT REPEATED MEASURES ANOVAS*

TABLE 29. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST CORRECT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 30. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST PERSEVERATIVE ERRORS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 31. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST RNADOM SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 32. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST CONSOLIDATION INDEX SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 33. MEAN SCORES AND STANDARD DEVIATIONS FOR RAVLT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 34. MEAN SCORES AND STANDARD DEVIATIONS FOR NAB TOTAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 35. MEAN SCORES AND STANDARD DEVIATIONS FOR TRAILS A SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 36. MEAN SCORES AND STANDARD DEVIATIONS FOR TRAILS B SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 37. MEAN SCORES AND STANDARD DEVIATIONS FOR COWAT/FAS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 38. MEAN SCORES AND STANDARD DEVIATIONS FOR RBANS TOTAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 39. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT COPY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 40. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT IMMEDIATE MEMORY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 41. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT DELAYED MEMORY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 42. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT RECOGNITION SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 43. APP SEVERITY LEVEL BY WCST REPEATED MEASURES ANOVAS

TABLE 44. APP SEVERITY LEVEL BY NEUROCOGNITION MEASURES REPEATED MEASURES ANOVAS

TABLE 45. SERVICE USE BY RCFT REPEATED MEASURES ANOVAS

TABLE 46. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST CORRECT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 47. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST PERSEVERATIVE ERRORS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 48. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST RANDOM ERRORS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 49. MEAN SCORES AND STANDARD DEVIATIONS FOR WCST CONSOLIDATION INDEX SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 50. MEAN SCORES AND STANDARD DEVIATIONS FOR RAVLT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 51. MEAN SCORES AND STANDARD DEVIATIONS FOR NAB TOTAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

- TABLE 52. MEAN SCORES AND STANDARD DEVIATIONS FOR TRAILS A SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 53. MEAN SCORES AND STANDARD DEVIATIONS FOR TRAILS B SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 54. MEAN SCORES AND STANDARD DEVIATIONS FOR COWAT/FAS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 55. MEAN SCORES AND STANDARD DEVIATIONS FOR RBANS TOTAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 56. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT COPY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 57. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT IMMEDIATE MEMORY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 58. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT DELAYED MEMORY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 59. MEAN SCORES AND STANDARD DEVIATIONS FOR RCFT RECOGNITION SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL
- TABLE 60. SERVICE USE BY FKK REPEATED MEASURES ANOVAS
- TABLE 61. SERVICE USE BY SOCIALCOGNITION MEASURES REPEATED MEASURES ANOVAS
- TABLE 62. SERVICE USE BY SOCIALCOGNITION MEASURES REPEATED MEASURES ANOVAS
- TABLE 63. MEAN SCORES AND STANDARD DEVIATIONS FOR HINTING SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 64. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK INTERNAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 65. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK SELF CONCEPT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 66. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK SELF EFFICACY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 67. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK POWERFUL OTHERS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 68. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK CHANCE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 69. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK EXTERNALITY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 70. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ INTERNAL POSITIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 71. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ PERSONAL POSITIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 72. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ SITUATIONAL POSITIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 73. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ INTERNAL NEGATIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18
- TABLE 74. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ PERSONAL NEGATIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 75. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ SITUATIONAL NEGATIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 76. MEAN SCORES AND STANDARD DEVIATIONS FOR CST SOCIAL SUPPORT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 77. MEAN SCORES AND STANDARD DEVIATIONS FOR CST SELF CONTROLLING SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 78. MEAN SCORES AND STANDARD DEVIATIONS FOR CST ESCAPE AVOIDANCE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 79. MEAN SCORES AND STANDARD DEVIATIONS FOR CST PLANFUL PROBLEM SOLVING SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 80. APP SEVERITY LEVEL BY FKK REPEATED MEASURES ANOVAS

TABLE 81. APP SEVERITY LEVEL BY SOCIALCOGNITION MEASURES REPEATED MEASURES ANOVAS

TABLE 82. APP SEVERITY LEVEL BY SOCIALCOGNITION MEASURES REPEATED MEASURES ANOVAS

TABLE 83. MEAN SCORES AND STANDARD DEVIATIONS FOR HINTING TASK SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 84. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK INTERNAL LOCUS OF CONTROL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 85. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK SELF CONCEPT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 86. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK SELF EFFICACY SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 87. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK POWERFUL OTHERS SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 88. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK CHANCE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 89. MEAN SCORES AND STANDARD DEVIATIONS FOR FKK EXTERNAL LOCUS OF CONTROL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 90. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ INTERNAL POSITIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 91. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ PERSONAL POSITIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 92. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ SITUATIONAL POSITIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 93. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ INTERNAL NEGATIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 94. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ PERSONAL NEGATIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 95. MEAN SCORES AND STANDARD DEVIATIONS FOR IPSAQ SITUATIONAL NEGATIVE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 96. MEAN SCORES AND STANDARD DEVIATIONS FOR CST SOCIAL SUPPORT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 97. MEAN SCORES AND STANDARD DEVIATIONS FOR CST SELF CONTROLLING SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 98. MEAN SCORES AND STANDARD DEVIATIONS FOR CST ESCAPE AVOIDANCE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 99. MEAN SCORES AND STANDARD DEVIATIONS FOR CST PLANFUL PROBLEM SOLVING SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 100. SERVICE USE BY IS REPEATED MEASURES ANOVAS

TABLE 101. MEAN SCORES AND STANDARD DEVIATIONS FOR IS RELABEL SCALE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 102. MEAN SCORES AND STANDARD DEVIATIONS FOR IS AWARENESS OF ILLNESS SCALE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 103. MEAN SCORES AND STANDARD DEVIATIONS FOR IS NEED FOR TREATMENT SCALE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 104. MEAN SCORES AND STANDARD DEVIATIONS FOR IS TOTAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 105. APP SEVERITY BY IS REPEATED MEASURES ANOVAS

TABLE 106. MEAN SCORES AND STANDARD DEVIATIONS FOR IS RELABEL SCALE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 107. MEAN SCORES AND STANDARD DEVIATIONS FOR IS AWARENESS OF ILLNESS SCALE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 108. MEAN SCORES AND STANDARD DEVIATIONS FOR IS NEED FOR TREATMENT SCALE SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 109. MEAN SCORES AND STANDARD DEVIATIONS FOR IS TOTAL INSIGHT SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 110. SERVICE USE BY NOSIE SUBSCALES AND TOTAL ASSETS REPEATED MEASURES ANOVAS

TABLE 111. MEAN SCORES AND STANDARD DEVIATIONS FOR NOSIE SUBSCALES AND TOTAL ASSETS AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 112. APP SEVERITY LEVEL BY NOSIE SUBSCALES AND TOTAL ASSETS REPEATED MEASURES ANOVAS

TABLE 113. MEAN SCORES AND STANDARD DEVIATIONS FOR NOSIE SUBSCALES AND TOTAL ASSETS AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 114. SERVICE USE BY BPRS FACTORS AND BPRS TOTAL REPEATED MEASURES ANOVAS

TABLE 115. APP SEVERITY LEVEL BY BPRS FACTORS AND BPRS TOTAL REPEATED MEASURES ANOVAS

TABLE 116. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS TOTAL SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 117. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS PSYCHOTIC DISORGANIZATION FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 118. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS HALLUCINATIONS/DELUSIONS FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 119. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS PARANOIA FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 120. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS EMOTIONAL BLUNTING FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 121. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS ANXIETY/DEPRESSION FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 122. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS AGITATION/ELATION FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 123. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS TOTAL AND FACTOR SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 124. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS ITEM SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF SERVICE USE BEFORE AGE 18

TABLE 125. MEAN SCORES AND STANDARD DEVIATIONS FOR BPRS ITEM SCORES AT ADMISSION, 6 MONTHS, AND 12 MONTHS AS A FUNCTION OF APP SEVERITY LEVEL

TABLE 126. RELATIONSHIP BETWEEN LEVEL OF DISCHARGE LOCATION RESTRICTIVENESS AND SERVICE USE ($N = 242$)

TABLE 127. RELATIONSHIP BETWEEN LEVEL OF DISCHARGE LOCATION RESTRICTIVENESS AND APP SEVERITY LEVEL ($N = 242$)

LIST OF FIGURES

- FIGURE 3.1. SERVICE MAIN EFFECT FOR SERVICE USAGE GROUPS ON WCST PERSEVERATIVE ERROR SCORE ACROSS TREATMENT
- FIGURE 3.2. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON WCST RANDOM ERROR SCORE ACROSS TREATMENT
- FIGURE 3.3. INTERACTION BETWEEN SERVICE USAGE GROUPS ON RAVLT SCORES ACROSS TREATMENT
- FIGURE 3.4. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON RCFT IMMEDIATE MEMORY SCORES ACROSS TREATMENT
- FIGURE 3.5. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON RCFT DELAYED MEMORY SCORES ACROSS TREATMENT
- FIGURE 3.6. INTERACTION BETWEEN SERVICE USAGE GROUPS ON TRAILS A PERFORMANCE ACROSS TREATMENT
- FIGURE 3.7. INTERACTION BETWEEN SERVICE USAGE GROUPS ON COWAT SCORES ACROSS TREATMENT
- FIGURE 3.8. TIME MAIN EFFECT AND INTERACTION BETWEEN SERVICE USAGE GROUPS ON RBANS TOTAL SCORES ACROSS TREATMENT
- FIGURE 3.9. INTERACTION BETWEEN APP SEVERITY LEVELS ON RAVLT SCORES ACROSS TREATMENT
- FIGURE 3.10. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON RCFT IMMEDIATE MEMORY SCORES ACROSS TREATMENT
- FIGURE 3.11. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON RCFT DELAYED MEMORY SCORES ACROSS TREATMENT
- FIGURE 3.12. TIME MAIN EFFECT AND INTERACTION BETWEEN APP SEVERITY LEVELS ON RBANS TOTAL SCORES ACROSS TREATMENT
- FIGURE 3.13. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON HINTING TASK SCORES ACROSS TREATMENT
- FIGURE 3.14. TIME MAIN EFFECT AND INTERACTION BETWEEN SERVICE USAGE GROUPS ON FKK CHANCE SUBSCALE SCORES ACROSS TREATMENT
- FIGURE 3.15. TIME MAIN EFFECT AND INTERACTION BETWEEN SERVICE USAGE GROUPS ON FKK EXTERNALITY SCALE SCORES ACROSS TREATMENT
- FIGURE 3.16. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON IPSAQ PERSONAL POSITIVE SCALE SCORES ACROSS TREATMENT
- FIGURE 3.17. INTERACTION BETWEEN SERVICE USAGE GROUPS ON IPSAQ INTERNAL NEGATIVE SCALE SCORES ACROSS TREATMENT
- FIGURE 3.18. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON CST SOCIAL SUPPORT SCORES ACROSS TREATMENT
- FIGURE 3.19. SERVICE USAGE GROUP MAIN EFFECT FOR SERVICE USAGE GROUPS ON CST SELF-CONTROLLING SCORES ACROSS TREATMENT
- FIGURE 3.20. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON HINTING TASK SCORES ACROSS TREATMENT
- FIGURE 3.21. TIME MAIN EFFECT AND INTERACTION BETWEEN APP SEVERITY LEVELS ON FKK CHANCE SCORES ACROSS TREATMENT
- FIGURE 3.22. TIME MAIN EFFECT AND INTERACTION BETWEEN APP SEVERITY LEVELS ON FKK EXTERNALITY SCORES ACROSS TREATMENT
- FIGURE 3.23. APP SEVERITY LEVELS MAIN EFFECT ON IPSAQ SITUATIONAL POSITIVE SUBSCALE SCORES ACROSS TREATMENT

FIGURE 3.24. INTERACTION BETWEEN APP SEVERITY LEVELS ON IPSAQ INTERNAL NEGATIVE SUBSCALE SCORES ACROSS TREATMENT

FIGURE 3.25. APP SEVERITY LEVELS MAIN EFFECT ON CST SELF-CONTROLLING SUBSCALE SCORES ACROSS TREATMENT
FIGURE 3.26. SERVICE MAIN EFFECT AND INTERACTION BETWEEN SERVICE USAGE GROUPS ON BIRCHWOOD INSIGHT TOTAL SCORES ACROSS TREATMENT

FIGURE 3.27. INTERACTION BETWEEN SERVICE USAGE GROUPS ON BIRCHWOOD RELABEL SYMPTOMS SCORES ACROSS TREATMENT

FIGURE 3.28. SERVICE USAGE GROUP MAIN EFFECT ON BIRCHWOOD RELABEL SYMPTOMS SCORES ACROSS TREATMENT

FIGURE 3.29. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON BIRCHWOOD NEED FOR TREATMENT SCORES ACROSS TREATMENT

FIGURE 3.30. INTERACTION BETWEEN APP SEVERITY LEVELS ON BIRCHWOOD INSIGHT TOTAL SCORES ACROSS TREATMENT

FIGURE 3.31. INTERACTION BETWEEN APP SEVERITY LEVELS ON BIRCHWOOD RELABEL SYMPTOMS SCALE SCORES ACROSS TREATMENT

FIGURE 3.32. APP SEVERITY LEVEL MAIN EFFECT AND INTERACTION BETWEEN APP SEVERITY LEVELS ON BIRCHWOOD AWARENESS SCALE SCORES ACROSS TREATMENT

FIGURE 3.33. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON BIRCHWOOD NEED FOR TREATMENT SCALE SCORES ACROSS TREATMENT

FIGURE 3.34. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE DAILY SCHEDULE COMPETENCE SCORES ACROSS TREATMENT

FIGURE 3.35. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE SOCIAL INTEREST SCORES ACROSS TREATMENT

FIGURE 3.36. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE NEATNESS SCORES ACROSS TREATMENT

FIGURE 3.37. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE IRRITABILITY SCORES ACROSS TREATMENT

FIGURE 3.38. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE PSYCHOTICISM SCORES ACROSS TREATMENT

FIGURE 3.39. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE MOTOR RETARDATION SCORES ACROSS TREATMENT

FIGURE 3.40. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON NOSIE TOTAL SCORES ACROSS TREATMENT

FIGURE 3.41. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON NOSIE DAILY SCHEDULE COMPETENCE SCORES ACROSS TREATMENT

FIGURE 3.42. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON NOSIE SOCIAL INTEREST SCORES ACROSS TREATMENT

FIGURE 3.43. TIME MAIN EFFECT AND INTERACTION BETWEEN APP SEVERITY LEVELS ON NOSIE NEATNESS SCORES ACROSS TREATMENT

FIGURE 3.44. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON NOSIE IRRITABILITY SCORES ACROSS TREATMENT

FIGURE 3.45. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON NOSIE PSYCHOTICISM SCORES ACROSS TREATMENT

FIGURE 3.46. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON NOSIE MOTOR RETARDATION SCORES ACROSS TREATMENT

FIGURE 3.47. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON NOSIE TOTAL SCORES ACROSS TREATMENT

FIGURE 3.48. INTERACTION BETWEEN SERVICE USAGE GROUPS ON BPRS PSYCHOTIC DISORGANIZATION FACTOR SCORES ACROSS TREATMENT

FIGURE 3.49. SERVICE USAGE GROUP MAIN EFFECT ON BPRS SUICIDALITY ITEM SCORES ACROSS TREATMENT

FIGURE 3.50. INTERACTION BETWEEN SERVICE USAGE GROUPS ON BPRS HALLUCINATIONS ITEM SCORES ACROSS TREATMENT

FIGURE 3.51. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON BPRS UNUSUAL THOUGHT CONTENT ITEM SCORES ACROSS TREATMENT

FIGURE 3.52. INTERACTION BETWEEN SERVICE USAGE GROUPS ON BPRS BIZARRE BEHAVIOR ITEM SCORES ACROSS TREATMENT

FIGURE 3.53. TIME MAIN EFFECT AND INTERACTION BETWEEN SERVICE USAGE GROUPS ON BPRS SELF-NEGLECT ITEM SCORES ACROSS TREATMENT

FIGURE 3.54. TIME MAIN EFFECT FOR SERVICE USAGE GROUPS ON BPRS MOTOR RETARDATION ITEM SCORES ACROSS TREATMENT

FIGURE 3.55. APP SEVERITY LEVELS MAIN EFFECT ON BPRS SUICIDALITY ITEM SCORES ACROSS TREATMENT

FIGURE 3.56. INTERACTION BETWEEN APP SEVERITY LEVELS ON BPRS BIZARRE BEHAVIOR ITEM SCORES ACROSS TREATMENT

FIGURE 3.57. INTERACTION BETWEEN APP SEVERITY LEVELS ON BPRS SELF-NEGLECT ITEM SCORES ACROSS TREATMENT

FIGURE 3.58. APP SEVERITY LEVELS MAIN EFFECT ON BPRS DISORIENTATION ITEM SCORES ACROSS TREATMENT

FIGURE 3.59. TIME MAIN EFFECT FOR APP SEVERITY LEVELS ON BPRS MOTOR RETARDATION ITEM SCORES ACROSS TREATMENT

DEDICATION

For my family, then and now.

All it took was a bushel and a peck.

ACKNOWLEDGEMENTS

I am extremely grateful to my advisor, William Spaulding, for his guidance throughout my undergraduate and graduate careers. Will, you are a source of inspiration and I thank you for your time and support throughout it all. I would like to thank my supervisory committee for being a part of this dissertation and appreciate the time and input of Dave Hansen, Mike Dodd, and Fred Luthans. I am also grateful for the support, mentorship, and wealth of information provided by Cal Garbin, Mary Sullivan, Jocelyn Ritchie, and Joe Swoboda. Thank you to the staff and all the participants at CTP, CMHC, and Midtown. I will be eternally gratefully for the experiences I had in these environments, and more importantly with these people. Finally, Dr. Williams deserves my thanks for making me passionate about the field of psychology and in serving others.

To the UNL Serious Mental Illness Research Group, rock star members past and present, I am grateful to have been in the trenches with you. I especially want to offer many thanks to Jeff Nolting, Jason Vogler, and Robert Johnson who encouraged me to apply to the research lab in the first place, and welcomed me with open arms. I could not have completed half of the analyses and presentations throughout graduate school without aid from Kee-Hong Choi. I am glad that those individuals in FI2007 were able to take the walk through graduate school with me, sharing times of struggle and success. I will forever think of the EBP tripod when in need of a laugh. This project would not have been possible if not for the hard work, including sifting through years of dust and boxes and countless coding hours, by clinical assistants. Thank you to Samantha Volk, Mark Shulte, Antoni Stewart, Ann Tweet, Missy Huenink, and Nikki Witter. My deepest gratitude goes to those who continue to conduct research, work primarily with, and advocate for this population.

I am extremely lucky to be supported by a great group of friends, attained prior to and during graduate school. The countless trips to Zhangs, football games, cupcakes and assorted foods made from a box, and your smiles and hugs made this possible. To my mother, my biggest cheerleader, who believed at times when I did not. Thank you to my brother, whose good heart and kind words were always appreciated and to my grandparents, whose faith is truly awe inspiring. To the Pape family, you have made

Lincoln truly enjoyable over the last seven years and I could not ask for a more supportive family. Finally, I am delighted Darin has been by my side through all this, he has been a constant source of support and I could not be more excited to start life after grad school together.

CHAPTER 1 ó INTRODUCTION AND LITERATURE REVIEW

Exploratory Analyses of a Developmental Conceptualization of Insight and Treatment
outcomes of Individuals with Serious Mental Illness in Psychiatric Rehabilitation

Individuals with Serious Mental Illnesses (SMI) are commonly observed to have an extreme lack of awareness of their condition, the consequences of their mental illness, and their need for treatment. Freud (1940) perceived that those with a diagnosis in the schizophrenia spectrum were acutely unaware of their own affect and cognitions. The WHO International Pilot Study of Schizophrenia validated earlier observations by finding that poor insight is one of 12 symptoms or signs selected to identify Schizophrenia and was found to be a statistically significant discriminator (Carpenter, Strauss, & Bartko, 1973). Since these early findings, there has been resurgence in recent years of research studying the concept of insight and its impact on recovery in the SMI population (Carpenter, Strauss, & Bartko, 1973; Ghaemi & Pope, 1994; Lysaker, Bell, Milstein, Bryson, & Beam-Goulet, 1994; Schwartz, 1998; Smith et al, 1999).

Definitions of Insight

The concept and definition of insight has changed over time. Jaspers (1963) was one of the first researchers to differentiate awareness of illness from insight, with awareness of illness being defined as the feeling of being "ill and changed." Insight was differentiated by Jaspers (1963) as an "objectively correct estimate of the severity of the illness (and) an objectively correct judgment of its particular type" (p. 419). Jaspers

thought that only when all of these features are present could the patient be considered to have insight (Jaspers, 1963; cited in Rusch & Corrigan, 2002).

Recent studies suggest that insight is not dichotomous, but multidimensional (Amador, Strauss, Yale, & Gorman, 1991; David, 1990; David & Kemp, 1997). David (1990) proposed that three dimensions comprise the construct of insight: recognition that one has a mental illness, the ability to re-label unusual mental events as pathological, and the belief that one needs treatment and actual adherence to treatment. These dimensions are not concrete but instead are overlapping, dynamic trends that account for diverse variations along the course of the illness. The model proposed by David (1990) has face validity and thus is commonly used in research studying insight (McGorry & McConville, 1999; Schwartz, 1998).

Amador et al. (1991) proposed another multidimensional model that differentiates between unawareness of illness and incorrect attribution of deficit or consequence of illness, and treats insight as a complex phenomenon with separate but interrelated mechanisms. Amador et al. (1993) concluded that the best way to define poor insight is as a construct consisting of multiple components much like the symptoms associated with SMI.

Insight can also be thought of as the ability to relate to professionals' views and the ability to meaningfully converse about the subject (Rusch & Corrigan, 2002).

Markova and Berrios (1995) conceptualized insight, as assessed in clinical interviews and observations, as a combination of the person's perception of his or her own condition, the clinician's perception of the individual's condition, and the clinician's own conceptual understanding of the construct of insight. Consequently, evaluating a

person's insight can be affected by the person and clinician's attitudes and beliefs, therapeutic alliance, sociocultural differences and backgrounds, and the clinician's working definition of insight.

Many differing techniques have been used to operationally define and measure insight. Objectively measuring and identifying symptoms and the degree of insight is important for reliably using valid results to inform diagnosis and treatment decisions throughout the course of illness. Although most current researchers agree that insight needs to be defined and measured as a multidimensional construct, some researchers continue to treat it as a dichotomous construct. Amador et al. (1991) try to measure the multidimensional nature of insight throughout the course of illness by designing tools like the Scale to Assess Unawareness of Mental Disorders (SUMD), a delineated 5-point scoring method that rates responses to controlled stimuli during a semi-structured interview. The SUMD assesses current and past insight as well as the individual's insight into the future course of their disorder and treatment. The SUMD includes a systematic assessment of historical records and other sources to determine what symptoms have actually been present. Although this may seem a prerequisite to assessing insight, such systematic accounting is not always included, and the SUMD has an advantage over other instruments in this regard. Also, Amador and colleagues (1993) proposed an additional dimension of insight, as assessed by the SUMD, that is capable of measuring the time and memory dimensions of insight, with full insight including the past, present, and possible future course (need for future treatment, risk of relapse, etc.) of the disorder. Thus, for example, an individual may understand that current hallucinations are illusory while fully believing the validity of past hallucinations.

Recently, researchers such as Frith (1992) and Lysaker et al. (2005) have begun to examine metacognition, the ability to think about thinking, and its relationship with the common signs and symptoms of schizophrenia. Some researchers regard metacognition as a component of insight, with the assumption that when a person's ability for metacognition is poor their degree of insight is lower. Examining metacognition often involves theory of mind tasks. Theory of mind is defined as the ability to form a representation of the consciousness of others and be able to draw conclusions about their motives and make inferences as to others' internal feelings (Lysaker et al., 2005). Research examining theory of mind tasks have shown that greater impairment is linked with deficits in visual and verbal memory and poorer flexibility for abstract thought (Greig et al., 2004). In a study by Lysaker et al. (2005), 61 men with schizophrenia were interviewed to examine the relationship between their scores on the metacognition assessment scale and quality of life, neurocognition, and insight. When age and gender were controlled, it was concluded that higher levels of metacognition, as defined as purposeful problem solving, were associated with less emotional withdrawal and paranoia and better social functioning, verbal memory, and insight (Lysaker et al., 2005). Therefore, when an individual with SMI is able to use purposeful problem solving and metacognition he or she is able to better conceptualize his or her illness. However, it is still unclear whether deficits in one's own awareness are a separate property from third person metacognition.

Etiology of Impaired Insight

If insight is a multidimensional construct, a single etiology is unlikely. Several causes are suspected to interact in complex ways and are expressed in different ways in the heterogeneous SMI population. It has been proposed that lack of insight in those that have schizophrenia is analogous to the neurocognitive deficits in the traumatic brain injury (TBI) population. Several researchers have observed and measured the unawareness of deficits in those with TBI, and have linked this lack of awareness to defects in the right hemisphere (Amador et al., 1991; McGlynn & Schacter, 1989; Prigatano & Schacter, 1991 as cited in Rusch and Corrigan, 2002); yet, some researchers studying schizophrenia did not find a relationship between defects in the right hemisphere and lack of insight (David et al., 1995; Kemp & David, 1996; McEvoy et al., 1996). Researchers studying the schizophrenia spectrum have found that the left hemisphere is affected more than the right in the SMI population. There is one finding of a correlation between poor insight and increased ventricle-to-brain ratios (Takai et al., 1992), and poor insight into one's disorder has been associated with lesions in the parietal lobe (McEvoy et al., 1996) and smaller brain size (Flashman, et al. 2000). Using MRI, Flashman et al. (2000) found that patients with poorer insight had smaller brain size and intracranial volumes than patients with higher levels of insight or the comparison subjects.

Researchers are currently divided as to whether or not unawareness of illness in the SMI population is best understood as the consequence of cognitive deficits that complicate the person's ability to understand confusing aspects of their mental illness and everyday life. Lysaker et al. (1994) and McEvoy et al. (1996) found that poor insight is related to frontal lobe deficits as measured by poorer performance on tests of

neurocognition and executive functioning, but others have failed to find a relationship (Cuesta et al., 1995; Freudenreich et al., 2004).

A different view of the etiology of impaired insight, not necessarily incompatible with neuropsychological views, is that it is a coping mechanism. McGlashan et al. (1975) showed that there are two distinct recovery styles for those that are experiencing psychosis. The first group, the "integrators," incorporates their mental illness into their everyday lives and has a high sense of awareness of the course that their illness has taken. The second group, or those that "seal over," refuse to discuss or even think about their mental illness and thus are inclined to have lower levels of awareness of their deficits, course of illness, and consequences of illness. These differences could be understood to reflect different coping style preferences. In a follow-up study of 30 "recovered" patients, McGlashan and Carpenter (1981) found that the absence of a negative attitude towards illness and symptoms was critical to achieving a positive outcome.

Due to the heterogeneous SMI population and the multifaceted nature of insight, insight as a coping mechanism may be more useful for some individuals, as well as more helpful in coping with some symptoms than others. Using the SUMD to measure symptoms, Mohammed et al. (1999) concluded that poor insight into negative symptoms (e.g. alogia, affective flattening, avolition) is associated with poor executive functioning while poor insight into positive symptoms (e.g. delusions, hallucinations, disorganized speech or thinking, grossly disorganized behavior, catatonic behaviors) is not associated with poor executive functioning. Therefore, it can be theorized that unawareness of negative symptoms may be due to neurocognitive deficits related to the frontal lobe and

executive functioning, while unawareness of positive symptoms may mainly act as a coping mechanism.

Poor insight as a coping method may also serve a protective role for individuals grappling with trying to understand their disorder who are experiencing higher levels of depression (Lysaker et al., 2005). Coping by having poor insight can allow the individual to avoid reality and as a result improve self-esteem and circumvent depression. Greater use of a form of coping, labeled accommodation or adaptation, was also a significant predictor of adjustment, providing support for the role of coping as a protective factor for people with schizophrenia (Lecomte & Mercier, 2005).

Insight and Suicide Risk

The role of poor insight as a coping method also suggests that good insight may paradoxically be a risk factor for depression and suicide. Individuals with a disorder in the schizophrenia spectrum have a 20 percent higher chance of attempting suicide than the general population and have a lifetime suicide attempt rate of 20 to 40% (Meltzer, Anad, & Alphas, 2000). In the SMI population, suicidal ideation and behavior have been linked to depression (Amador et al., 1994; Caldwell & Gottesman, 1990). Research has indicated that higher levels of insight lead to lower self-esteem and greater depression, thus increasing risk of suicide attempts (Amador et al., 1994; Caldwell & Gottesman, 1990; Evren & Evren, 2004; Pompili et al., 2004). Baechler (1979; restated in Pompili et al., 2004) stated that "schizophrenics do not kill themselves insofar as they are schizophrenic but insofar as they are persons who know they are schizophrenic or are threatened with becoming so and who wish to avoid this fated outcome." Amador et al.

(1996) reported that in their study of 218 patients with schizophrenia, 169 individuals did not have suicidal behavior whereas 49 had suicidal behaviors or ideologies. Amador and colleagues (1996) found in their study that patients who have more awareness of their delusions, blunted affect, and anhedonia were more likely to commit suicide. In a study by Evren and Evren (2004), sixty individuals with schizophrenia were interviewed and their clinical case summaries were reviewed. About half of the sample, 45%, had experienced suicidal ideation, had made one or more suicide attempts, had lower negative symptom scores, and had been diagnosed as depressed at one point in time. For these individuals, their mean scores on the three components of insight were statistically significantly higher than for those who did not have a history of suicidal ideation or attempts. Evren and Evren (2004) replicated other studies (e.g. Caldwell & Gottesman, 1990) that indicated that those that did attempt suicide and those that did not attempt suicide did not differ on demographic characteristics such as age, gender, marital status, educational status, and employment history.

The Role of Insight in Treatment and Recovery

While increased levels of insight have been shown to be a risk factor for suicide, moderate levels of insight can also act as an important factor in treatment and recovery. Poor insight can create a barrier between practitioner and patient that may affect the alliance during therapy and the patient's chance for long-term success. Poor insight has been associated with poor work skills, more noncompliance, and more readmissions to hospitals (Lysaker et al., 1994). Several studies have shown that high levels of insight are associated with improved outcome, fewer hospitalizations, better post-hospital

adjustment, and better medication compliance both in psychiatric in-patient settings and outpatient settings (Amador et al., 1993; Schwartz, 1998). These findings support the view that increasing insight early during the course of treatment can increase the probability of treatment being successful.

Insight and Human Development

To fully understand insight into adult SMI, attention must be paid to the emotional, physical, and mental health difficulties that people experience as they transition from adolescence to adulthood. Research indicates that one in five children have some form of a diagnosable mental, emotional, or behavioral disorder, yet 70 percent of those children do not receive mental health services (Kenny et al., 2002). In addition to a lack of a common approach to detecting and following these children across systems, the neglect of services seems to be due to factors such as the varying definitions and classifications of childhood psychiatric disorders, research with heterogeneous child samples, using differing age groups for studies examining adolescence, and multi-disciplines disagreeing on what constitutes child and adolescence psychosis and emotional disorders.

The term Serious Emotional Disturbance, applied to people under age 18, is comparable to Serious Mental Illness applied to adults. Both terms are used colloquially and legislatively to describe people who are diagnosed with a psychiatric disorder according to the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) (American Psychiatric Association, 1994), when the diagnosed condition results in severely impaired functioning and disability.

Children and adolescents can be diagnosed with different diagnoses as they progress through the lifespan even though their symptomatology may be similar; they can be deemed as having SED prior to age 18 and after they are 18 they can be deemed again as having SMI. Therefore, it is difficult to estimate the rates of SED and SMI in the population age 16 to 21 (Jonikas, Laris, & Cook, 2003) and hence it is difficult to estimate the amount of the population that is not receiving mental health services.

Research into SMI has demonstrated that disorders in the schizophrenia spectrum are chronic and that poor outcome is often linked to earlier onset (Helgeland & Torgensen, 2005). Helgeland and Torgensen (2005) also found diagnostic stability from adolescence psychiatric pathology to adulthood for schizophrenia spectrum disorders. Other researchers disagree with this finding, concluding that it is impossible to have valid diagnostic consistency between schizophrenia and psychosis in childhood (Chen, Swann, & Burt, 1996). Due to these inconsistent results it is clear that further research into the transition from SED to SMI is needed.

The transition from adolescence to adulthood can be understood in terms of developmental and institutional transitions (Vander Stoep, Davis, & Collins, 2000). Developmental transitions are biological and social in nature while institutional transitions refer to changes in service settings, legal, or bureaucratic status. Research by Vander Stoep et al. (2000) suggests that establishing and maintaining peer and family relationships, employment, high levels of functioning, treatment adherence, and transition back into a community setting can be impeded or delayed throughout the lifespan by SED or psychiatric pathology in childhood or adolescence.

According to Helgeland and Torgersen (2005), SMI in adulthood is associated with developmental hardships such as complications during pregnancy, neurodevelopmental abnormalities, and delayed motor and language development, thus suggesting schizophrenia is a developmental mental illness with relative stability in childhood and adolescence and onset in early adulthood. As a result it can be deduced that individuals with adolescent psychiatric pathology (APP) that have poor levels of insight may continue to have poor levels of insight once in the SMI population. Clinical characteristics such as degree of psychopathology, length of previous hospitalizations, and age of first hospitalization are also associated with patterns of psychosocial treatment response and insight into one's illness (Kupper and Hoffmann, 2000; Peer and Spaulding, 2007). Although research has not conceived a developmental theory of insight, it can be assumed that valuable information that can inform recovery decisions can be gleaned from future research viewing insight in a developmental context.

Simply having APP can hinder an individual's level of functioning and course of illness factors such as treatment adherence, but failure to engage in services can also be attributed to lack of insight (Carpenter, Strauss, & Bartko, 1973; Ghaemi & Pope, 1994; Lysaker et al., 1994; Schwartz, 1998; Smith et al., 1999). Developmental impairments are more ubiquitous in those with symptoms in adolescence or APP than in those with late onset schizophrenia (Helgeland & Torgersen, 2005), thus indicating a possible more biologically severe subtype of the SMI population. Individuals that have the lowest degrees of insight and a history of APP may have more severe symptoms and may, despite receiving mental health services during adolescence, continue to have poor insight as adults receiving in-patient services.

Families are often a source of emotional support for those with a mental illness. Research has shown that individuals with APP have more dysfunctional family relationships as well as increased perceptions that their families are distant and do not offer support (Jonikas et al., 2003). While adolescence individuals may seek support from other sources such as friends or significant others; for adolescents with mental illnesses this may not be possible due to poor social skills, paranoia, stigma associated from having a mental illness, blunted affect, and aggressive behaviors towards peers. Failure to establish or maintain relationships may force the person to manage their illness differently; poor insight may develop as a coping mechanism in adolescence to substitute for the lack of support from family and peers. This method of coping may continue throughout the course of the illness and contribute to the individual's insight remaining low after being admitted to inpatient treatment settings.

In a study Van Meijel et al. (2002) healthcare professionals and families of patients with schizophrenia were interviewed. All interviewees agreed that in order to improve outcomes for those with SMI earlier intervention and symptom recognition is needed. Furthermore, all participants in the study agreed that a certain level of insight is needed to accomplish early recognition and intervention for the SMI population (Van Meijel et al., 2002). Those that have lower levels of insight in adolescence while meeting criteria for SED may, with the help of mental health services and psychopharmacology, increase their levels of insight while making the transition into the SMI population. This increased understanding of the need for services, ability to understand their mental illness, and the ability to re-label unusual mental events as pathological can have important implications for treatment and the chronicity of states of psychosis that the

patient experiences and copes with, as well as the number of times that they must be hospitalized. However, individuals with extremely poor insight during adolescence and into adulthood may, despite intensive intervention, not be able to understand and re-label their illness and may be more likely to have poorer treatment outcomes. Further research needs to accurately longitudinally measure, conceptualize, and define insight in order to inform our decisions for early identification and intervention strategies, as well as help explain the developmental transition from SED to SMI and the impact insight may have on this population with greater symptom severity. The conclusion can be drawn that further analysis into the relationship of insight and the transition from SED or APP to SMI can create new contexts in which SED and SMI and the mental health services for each can be viewed.

To date, there has been only one exploratory study conducted in a clinical setting that compares the characteristics of people with APP to those without. In a study by Wynne (2009) the relationship between APP and SMI was examined by assessing the relationship between prior mental health services before the age of 18 and time of assessment on people's insight into their illnesses as adults participating in a psychiatric rehabilitation. The sample consisted of 112 patients recruited from an inpatient psychiatric rehabilitation unit at a state psychiatric hospital. Results indicated that those with service use during adolescence were found to be younger at admission, have slightly lower levels of education, have an earlier age of onset, and significantly more previous hospitalizations. A relationship between Axis II diagnosis and service use was also identified, indicating that having an Axis II diagnosis in adulthood and APP may signify that individuals had particularly noticeable symptoms that increased their chances of

being served in adolescence.

Wynne (2009) also found that improvements in neurocognitive functioning were not evident over the course of treatment. Although improvements in insight were evident over the course of treatment for the general CTP population, adults with SMI who used mental health services before the age of 18 did not significantly differ from those who did not use services before the age of 18 with respect to level of insight; except for awareness of illness at 6 months as measured by the IS. Results of analyses on insight measures over the course of treatment by APP severity level suggest that increases in measures of insight are not evident for the medium and high APP severity groups and changes in total insight and insight into need for treatment are different for the no services in adolescence and low APP severity groups over time. This field of research has numerous implications for measuring and increasing insight among persons with SMI and may have implications for treatment success within institutions and treatment outcomes once discharged into the community.

A major factor contributing to the paucity of research in this area is the substantial methodological problems associated with this kind of research. The impact of prior mental health services in adolescence on SMI takes years to demonstrate full, measurable effects. Obtaining historical information from treatment providers who served individuals during adolescence also proves problematic, as by the time this information has been requested by the adult's treatment provider, long periods of time may have passed and the clinical information destroyed. Furthermore, it takes years to accumulate the needed data in order to be able to understand this process. Although the study by Wynne (2009) used data from a nine year period, there was low power for several key

analyses due to missing data and/or changes in the clinical assessment battery. The use of multivariate analyses in this study was undermined by a low number of valid cases on many of the variables over time. Therefore, conclusions about differences between groups, or the lack thereof, over the course of treatment are tentative due to insufficient power. Even though there was low power the presence of several interactions between service use and APP severity levels and insight measures across treatment approaching statistical significance indicates that effects were not missed. At any rate, this study explored the relationship between service use and insight such that future studies in this area can make more informed research hypotheses using stricter constraints in research design to circumvent problems related to lack of power and missing data. Thus further study using a greater amount of participants completing repeated measures over time could overcome low power problems and find statistically significant relationships between APP severity groupings and insight over time in treatment.

In summary, previous research has indicated that psychosocial functioning (e.g., social competence, social interest) in SMI is a product of complex relationships between factors such as neurocognitive functioning (Brekke, Kohrt, & Green, 2001) and sociocognitive variables, such as locus of control, and that both of these domains have been implicated in the research on the concept of insight. A greater understanding of the interrelationships between these factors, utilizing the developmental conceptualization of insight proposed by Wynne (2009) may aid not only in improving conceptualization of insight and psychosocial functioning during treatment in adolescence and adulthood, but also treatment outcomes once individuals are in the community functioning as adults (Hoffmann & Kupper, 2002; Kupper & Hoffman, 2000; Smith et al., 1999). Also, better

understanding of these complex relationships may be able to aid in improving treatment outcomes for those with SMI without solely depending on improving a person's insight into their illness.

Present Study

The purpose of the present study was to further examine the relationship between adolescent psychiatric pathology, often captured by the term of art SED, and SMI by assessing the relationship between prior mental health services before the age of 18 and time of assessment on people's insight into their illnesses. Earlier understanding of degrees of insight in the SED population has important ramifications for the lifelong diagnosis and treatment of those with SMI. Whereas poor levels of insight in adolescence in some individuals with more severe symptomatology may continue to be stable into adulthood despite receiving psychological services, some individuals with the help of mental health services may be able to increase their levels of insight over time and improve their likelihood of successful treatment outcome and transition into the community. However, there are also reasons to believe these APP severity groups do not differ. The reasons for which an individual interacts with mental health service systems are complicated and there is no one path to treatment. For example, family involvement for people with SMI varies greatly. If one person has an active family and another does not, the former may be more likely to have a receive services or be screened for APP than the latter, independent of their functional abilities, simply because a family member advocated for the appointment. Alternatively, an individual may repeatedly encounter mental health professionals in adolescence, typifying a more chronic course that may be

better served through guardianship, but no interested party seeks guardianship on behalf of the individual. These confounds suggest that we need to know more about the concept of insight and APP and its correlates to the clinical picture for these same people as they transition and receive adult psychiatric services.

The present study had two primary objectives:

1. To determine if there was a pattern of differential functioning for adults with SMI with or without mental health service use during adolescence across various levels of functioning. In general, it was anticipated that individuals with adolescent psychiatric pathology would demonstrate lower overall functioning than those without adolescent psychiatric pathology. As described above, these are individuals that for various unknown reasons, received mental health services during adolescence. Research indicates that this population may represent individuals experiencing more severe symptomatology, and that adolescent psychiatric pathology can impede treatment success in adulthood. Because of this, hypothesis 1 predicted that those individuals who had various levels of APP will show demonstrably lower functioning in the clinical setting. Thus, it was hypothesized that the APP severity would correlate with clinical functioning. Specifically, the groups would differ in statistically significant ways across all domains of functioning measured (neurocognitive, social cognitive, symptomatology, and behavioral functioning).

A. Upon Admission. Hypothesis 2 predicts that at the time of admission, individuals with adolescent psychiatric pathology would demonstrate lower overall functioning than those without adolescent psychiatric pathology on neurocognitive, social cognitive, and

behavioral functioning measures. However, hypothesis 3 predicted that because of prior contact with services, those individuals with adolescent psychiatric pathology would have higher levels of insight into their disorder at admission than those without adolescent psychiatric pathology. In addition, hypothesis 4 anticipated that those with adolescent psychiatric pathology would have higher levels of symptomatology. When analyses using different levels of adolescent psychiatric pathology were conducted, hypothesis 5 predicted that those with adolescent psychiatric pathology that required hospitalization in adolescence or high mental health service utilization in adolescence would have higher levels of insight into their disorder at admission than those with none or less adolescent psychiatric pathology. However, hypothesis 6 predicted this group would also have the highest levels of symptomatology and would have the lowest scores on neurocognitive and sociocognitive measures.

B. Over the course of treatment. Hypothesis 7 predicted that overall differences in functioning, symptomatology, and treatment adherence would not remain over the course of treatment. It was hypothesized that as a result of psychiatric rehabilitation there would be an increase in neurocognitive functioning, insight into disorder, internal locus of control, and behavioral functioning and a decrease in external locus of control and symptomatology for those with and without adolescent psychopathology. That is, it is hypothesis 8 anticipated that both groups would show improvement in functioning over the course of psychiatric rehabilitation. It was predicted that significant differences at admission or six months would not remain between those with adolescent psychiatric pathology and those without adolescent psychiatric pathology in all areas after 12 months of treatment. However, it was predicted that those individuals with adolescent

psychiatric pathology would demonstrate higher symptomatology after 12 months of rehabilitation. Furthermore, when different levels of adolescent psychiatric pathology were used in analyses, hypothesis 9 predicted that as adolescent psychiatric pathology becomes more severe functioning will decrease, there would be more severe symptomatology. Overall, while it was hypothesized there would be differences in a variety of domains it was also hypothesized that differences in insight upon admission would not remain one-year after beginning psychiatric rehabilitation.

2. *To determine if there is a pattern of differential outcome following discharge for individuals with adolescent psychiatric pathology from those without.* While previous research had not been conducted to inform hypotheses in this area, it was generally hypothesized that there would be differences between groups in terms of treatment outcome. Specifically:

C. *Discharge disposition.* Hypothesis 10 predicts that discharge disposition would differ between the two groups in that individuals with severe adolescent psychiatric pathology would be discharged to a more restrictive setting. This hypothesis was based on the idea that those individuals who used high amounts of mental health services or were hospitalized during adolescence may have represented a group with more severe symptomatology and may have been hospitalized more and for longer periods of time during the course of their illness. These individuals may, through frequent hospitalizations, been reinforced to fulfill the "patient role" and have been institutionalized, thus representing a group or participants who take a longer time to transition into the community and may influence treatment providers' notions such that

there is an assumption that individuals who were hospitalized during adolescence require higher levels of care and supervision.

D. Rehospitalization rate. Hypothesis 11 predicted that there would be a greater rate of rehospitalization for those with APP than those without APP. Furthermore, when analyses were conducted between the different APP severity groups, hypothesis 12 predicted that those individuals in the medium and highest APP severity groups would have a greater rate of rehospitalization than those in the low to no APP severity groups. In addition, hypothesis 12 predicted that those in the highest APP severity group would have the greatest rate of rehospitalization, as these individuals may represent a subgroup experiencing more severe and refractory symptoms that can be potential barriers to living in the community for extended periods.

Based on the results of these analyses, exploratory analyses were undertaken to glean additional information about the role of service use and APP severity level in the psychiatric rehabilitation and insight of individuals with SMI. This data was accumulated over the course of thirteen years, allowing for a preliminary look at the therapeutic consequences of prior mental health service usage and the longitudinal effects of psychiatric rehabilitation treatment for these individuals. The archival database utilized in this proposed study was ideal for this type of research. It afforded the opportunity to study populations with different service histories under a similar set of clinical circumstances over an extended period of time. Although one previous study (Wynne, 2009) sought to better understand the relationship between mental health service utilization during adolescence on insight scores across treatment in adulthood, it should

be reiterated that the proposed study built on the previous study by Wynne (2009), was highly exploratory in nature, and sought to expand the scope of previous research by including a larger sample size and more assessment measures that have been implicated as important to the study of the concept of insight and in the treatment outcome for those with SMI. This was a critical first step in the analysis of the role of mental health service usage during adolescence and adolescent psychiatric pathology in the treatment of adults with SMI and these adults' insight into their mental illness.

CHAPTER 2 - METHOD

Design Overview

The primary purpose of this study was to assess the relationship of mental health service usage and adolescent psychiatric pathology severity with insight before, during, and after treatment. In addition, a secondary relationship between adolescent psychiatric pathology and functioning in a variety of domains before, during, and after treatment was assessed. The participants with adolescent psychiatric pathology were compared to those without to determine if any differences existed. Also, when adolescent psychiatric pathology was separated into three separate groups by the amount and type of services utilized during adolescence, these three different levels of adolescent psychiatric pathology were compared to those without. Univariate and multivariate analyses were conducted within and between the two groups service groups (service use vs. no service use in adolescence) and the four APP severity level groups with respect to overall functioning using multiple measures described below.

Setting

The Psychiatric Rehabilitation Program, formerly known as and will continue to be referred to in this paper as the Community Transition Program (CTP), was an inpatient unit at the Lincoln Regional Center (LRC) a public state psychiatric hospital in Lincoln, Nebraska. This 40-bed inpatient unit hosted a comprehensive psychiatric rehabilitation program for those most disabled by mental illness in the State of Nebraska. Individuals were typically discharged to a less restrictive setting after a 9 to 18 month period of intensive treatment, with the average length of stay being 12 months. Treatment

engagement was encouraged through the use of contingency management based on social learning theory. The regimen included pharmacotherapy, psychoeducational groups and classes to target improved management of symptoms and disorder, and training aimed at improving occupational, leisure, and social skills. The treatment was designed to target multiple levels and domains of functioning for individuals with SMI, rather than only targeting an isolated area of deficit (e.g., symptoms). Thus, treatment did not focus on clinical diagnosis. Rather, treatment plans using functional analysis were individualized and based on making step-wise improvements in deficient areas of functioning and capitalizing on observed and participant perceived areas of relative strength of the individual. Treatment plans were designed by a multidisciplinary treatment team including nurses, social workers, psychiatrists, occupational therapists, psychologists, and psychiatric technicians. In addition, program participants were encouraged to be active members of their own treatment team for the purpose of increased engagement in treatment and to promote the concept of recovery in the therapeutic milieu. Consequently, the CTP referred to individuals as "participants" instead of patients, reinforcing the idea of active engagement of participants in their treatment. For a detailed outline on the theoretical foundation and practical applications of this innovative, state-of-the-art treatment technology read *Treatment and Rehabilitation of Severe Mental Illness* (Spaulding et al., 2003).

The Lancaster County Community Mental Health Center (LCCMHC) also contributed data to the archival database. Since a majority of participants from CTP are served by LCCMHC upon discharge, ongoing program evaluation using records at

LCCMHC yielded data pertaining to outcome such as rehospitalization rate, discharge location, and use of services in the community.

Participants

Participants discharged from the CTP program from 1996 through March 2009 contributed data to the archival database utilized in this study. The CTP program participants were not directly involved in any specific research procedure. Rather, the database included the ongoing clinical data collected as part of the routine assessment process described further below at the CTP and additional data gathered through chart review.

Participants of the CTP must have had as part of the criterion for admission an Axis I major mental disorder administrative designation as SPMI. This sample represents a severe and treatment refractory subpopulation. They were referred to the CTP because of either extended institutionalization in custodial settings or repeated re-hospitalizations with no stable community functioning. The sample consisted of 354 participants. All subjects met DSM-IV (American Psychiatric Association, 1994) criteria for schizophrenia, schizoaffective disorder, or bipolar disorder. Participants were between the ages of 18 and 60.

DSM-IV (American Psychiatric Association, 1994) diagnoses for patients with schizophrenia, schizoaffective disorder, and bipolar disorder were confirmed prior to the study by interviews with the participants, chart reviews, reviews of historical documentation, and consultation with the program director and treatment team. Patient data was used only when ratings from these different sources produced the same

diagnosis. All patients were on various combinations of antipsychotics, mood stabilizers, and anxiolytic medications during the study.

Initially, the archival database had 354 participants. Because available outcome data in the database pertained to the most recent admission to CTP and multiple individuals in the database received rehabilitation at various stages in their life and disorder, all previous admissions before the most recent admission were excluded from analyses to maximize the number of valid cases available for analyses. Also, participants with lengths of stay in the CTP program of less than six months were excluded from analyses since treatment effects were captured as there was not adequate time to participate in the CTP rehabilitation program or have multiple assessment administrations within six months time.

A vast majority of CTP participants were either civilly committed or admitted by a legal guardian. Either way, treatment was typically viewed as involuntary in that most did not decide for themselves to enter the program. This was particularly important in developmental analyses of insight due to individuals being hospitalized in adolescence being made wards of the state at an earlier age. Involuntary treatment may play a role in the concept of "insight," as some researchers have suggested that insight is often viewed in clinical contexts as an individual's degree of agreement with treatment providers (Rusch & Corrigan, 2002). Analyses took this variable into account and determined differences amongst those with and without mental health service use in adolescence and between the different APP severity levels.

Measures

Demographic and Clinical Characteristics

Through chart review at CTP, information regarding the demographic and clinical characteristics of participants was included in the archival database. These variables included gender, age, years of education, race/ethnicity, marital status, legal status, number of previous hospitalizations in the participant's lifetime, age of first hospitalization, Axis I and Axis II psychiatric diagnoses, comorbid Axis I diagnoses, length of stay at CTP, and other relevant variables.

Neurocognition Measures

1) Repeatable Battery for the Assessment of Neuropsychological Status (RBANS; Randolph, 1997). The RBANS is a brief neurocognitive screening assessment (approximately 25 minutes) ideal for individuals who may lack levels of attention needed to complete longer assessments. There are two forms of the RBANS (Form A and Form B), intended to minimize the practice effects of repeated administrations. The twelve subtests of the RBANS are grouped into five neurocognitive domains. The five domain scores of the RBANS include immediate memory, delayed memory, attention, language, and visuospatial/constructional functioning. In addition, a total index score represents overall cognitive functioning. Convergent validity has been established in people diagnosed with schizophrenia for the RBANS with other neuropsychological constructs, like memory and intelligence (Gold, Queern, Iannone, & Buchanan, 1999; Hobart, Goldberg, Bartko, & Gold, 1999). In addition, sensitivity to patterns of cognitive impairment in SMI and general reliability and validity have been evaluated (Gold et al., 1999; Hobart et al., 1999).

2) COGLAB (Spaulding, Garbin, & Dras, 1989). COGLAB was created as a computerized test battery comprised of tests common in the psychopathology literature. The WCST, a measure of executive functioning, was used in an effort to understand the subject's ability for abstract thought, cognitive flexibility, and to replicate past studies that have found a significant relationship between the WCST and insight (Lysaker et al., 1994; McEvoy et al., 1996). CTP standard assessment included two of the tests from the battery; however, for the current study the Card Sorting Task (WCST; an adaptation of the Wisconsin Card Sorting Task, Heaton et al., 1993) was used. The WCST is a computerized neuropsychological test of "set shifting" in which participants are asked to display their ability to be flexible when exposed to changing schedules of reinforcement by matching a target card to one of four stimulus cards without being told the matching principle of color, shape, or number that changes frequently without warning to the subject. The WCST was used to measure executive functioning and its known sensitivity to impairment in concept formation, cognitive flexibility, and abstract thought (Lysaker et al., 1994; Lysaker et al., 1998). The WCST allows clinicians and researchers to assess frontal lobe functions such as strategic planning, organized searching, utilizing environmental feedback to shift cognitive sets, goal oriented or problem solving behavior, and decreasing impulsive responses. The computerized WCST takes approximately 12-20 minutes to complete and provides objective measures of overall success by computing the number of categories achieved, number of trials, number of errors, number of perseverative errors, percentage of perseverative errors, and the consolidation index as a measure of set-shifting.

Good discriminant validity was demonstrated between individuals diagnosed with schizophrenia and controls in an early study of the COGLAB (Spaulding et al., 1989). Results from large multivariate studies conducted with normal and patient populations indicate overall acceptable psychometric properties of the COGLAB (Spaulding, Hargrove, Crinean, & Martin, 1981; Spaulding et al., 1989).

3) Rey Auditory Verbal Learning Test (RAVLT; Schmidt, 1996). The RAVLT, is a seven trial list-learning task with alternative forms consisting of 15 words presented in an auditory format. Participants were instructed to recall as many words as they can from the list immediately following each of five trials. A distractor trial was then presented consisting of a different list of 15 words and participants are required to recall as many words from this distractor list. Finally, participants were required to recall as many words as possible from the original list without it being presented again. In general, the RAVLT provides a measure of verbal memory. The number of words remembered after the fifth trial is the most commonly used RAVLT score. Acceptable discriminant validity between memory impaired vs. memory intact patients and normal vs. neurological patients has been demonstrated, as well as adequate test-retest validity (Schmidt, 1996).

4) Rey-Osterrieth Complex Figure Test (RCFT; Rey, 1941). This is a test of visuoconstructional ability and nonverbal memory. It is comprised of four tasks: a copy trial, immediate recall trial, delayed recall trial, and a recognition task. Figures were scored using the 18-point scoring system, originally developed by Osterrieth (1944), and outlined in Meyers & Meyers (1995).

5) The Trailmaking Test (A&B) (Army Individual Battery, 1944). This two component test assesses attention, visual scanning, and information processing. In Part A, individuals connected circles numbered 1 through 25 by drawing a line sequentially from 1 to 25 as quickly as possible. In Part B, individuals completed a similar task alternating sequentially between numbers and letters (e.g. 1 to A, A to 2, 2 to B, B to 3). Performance was measured by subtracting the number of errors from the total possible score, resulting in two scores, one for each trial.

6) Neuropsychological Assessment Battery Screening Memory Domain (NAB) (Stern & White, 2001). The NAB Screening Memory Domain score is a composite measure of the participant's verbal and visual memory functioning, based on the sum of the Screening Shape Learning Immediate Recognition, Screening Shape Learning Delayed Recognition, Screening Story Learning Immediate Recall, and the Screening Story Learning Delayed Recall scores. The Screening Memory Domain score is used as an indicator of an individual's verbal explicit learning, visual explicit learning, verbal free recall after a brief delay, and visual delayed recognition memory after a brief delay (Stern & White, 2001). The reliability coefficient of the Screening Memory Domain score is .79 (Stern & White, 2001).

7) Controlled Oral Word Association Test (COWAT or FAS) (Benton & Hamsher, 1976). The COWAT measures verbal fluency and word generating ability, or the ability to rapidly generate and organize verbal information. In a series of 60-s trials, participants were asked to name as many words as possible that began with specified letters of the alphabet F, A, and S. Total number of words generated was scored.

Theory of Mind Measures

1) Hinting Task (Corcoran, Mercer & Frith, 1995). The Hinting Task measures theory of mind abilities and requires working memory, verbal comprehension, and verbal production abilities. A summary score was generated based on ability and facility of comprehension and interpretation of social implications. Limited psychometric data is available, however, the Hinting Task is sensitive to deficits in schizophrenia, correlates with other ToM measures and has good face validity in studies with seriously mentally ill participants (Corcoran & Frith, 2003).

Socialcognition Measures

1) Inventory for the Measurement of Self-Efficacy and Externality (I-SEE or FKK; Krampen, 1991). The I-SEE provides a measure of global attributional style, or locus of control. It is comprised of 32 items which were each rated on a six point Likert scale ranging from *strongly disagree* to *strongly agree*. The measure consists of four main scales: "internality" (i.e., "whether I have an accident is based on my own behavior"); "self-concept of one's own competence" or "self-efficacy" (i.e. "I can do many things to protect my self-interest"); "powerful others" in controlling beliefs" (i.e., "other people often prevent my plans from becoming reality"); and "chance in controlling beliefs" (i.e., "whether I fall ill is a matter of chance"). These scales were combined to yield two composite scales a general external scale ("externality") and a general internal scale ("self-concept of one's own efficacy") which represents participant's beliefs about their self-efficacy. Krampen (1991) established reliability for the two composite scales.

2) The Internal, Personal, Situational Attribution Questionnaire (IPSAQ)

(Kinderman & Bentall, 1997) assesses a more interpersonal attributional style based on

participants explanations of positive and negative social scenarios. It yields two measures: an externalizing bias (Eb) score (the degree to which persons attribute negative events to external factors and attribute positive events to themselves) and personalizing bias (Pb) (the degree to which persons attribute negative events to other people as opposed to situational factors).

3) The Coping Strategies Task (CST) (Mindt & Spaulding, 2002). The CST is used for assessing coping-related cognition in individuals with schizophrenia-spectrum disorders. The CST is comprised of four coping subscale scores including: Social Support Seeking, Self-Controlling, Escape Avoidance, and Planful Problem Solving. Reliability analyses reveal that the CST and its subscales demonstrated adequate reliability, although one subscale (Behavior Reaction) demonstrated less robust split-half and test-retest reliabilities. Concurrent validity was evaluated by analyzing the relationship between the CST and measure of stress, observed behavior, and neurocognition. Correlational analyses have revealed that coping attributions, as measured by the CST, were found to be association with perceived stress, observed behavior, and executive functioning.

Insight Measures

1) Insight Scale (IS) (Birchwood, Smith, Drury, Healy, Macmillan, & Slade, 1994). This brief self-report measure (8 items) allows participants to choose one of three responses: agree, disagree or unsure, for each item. It yields a total score and three subscale scores representing Davidø (1990) three domains of insight. The correct answer for each was counted as one point and each dimension is scored on a scale of 0-4, with an

overall insight score ranging from 0 to 12 with higher scores indicating higher levels of insight. The three subscales, therefore, are: "need for treatment" (i.e., "I do not need medication"), "ability to relabel psychotic experiences" ("some of my symptoms were made by my mind"), and "awareness of illness" ("I am mentally well"). This measure of insight focused on insight into functional impairment rather than specific illness categorizations. This self-report questionnaire is well validated and used extensively with the SMI population, with test-retest reliability equaling .90 (Birchwood et al., 1994).

2) The Self-Appraisal of Illness Questionnaire (SAIQ) (Marks, Fastenau, Lysaker, & Bond, 2000). The SAIQ is a self-report instrument designed to assess attitudes toward mental illness among persons receiving psychiatric treatment. The SAIQ is a pencil and paper self-report instrument composed of 17 items. The format for each item is a statement or a question. The items addressed acknowledgment of illness, beliefs about the outcome of illness, acknowledgment of a need for psychiatric treatment, and extent of worry about illness and about illness related issues. Participants were asked to respond to the statements and questions using a four-point Likert scale, which varies according to the statement or question content. The validity of the SAIQ was examined through a factor analysis. Three factors emerged: Need for Treatment, Worry, and Presence/Outcome of Illness. The three SAIQ subscales are correlated with researcher-rated insight scales and neuropsychological tests. Results indicate that the Need for Treatment and Presence/Outcome subscales were significantly correlated with both researcher-rated insight scales and with neuropsychological tests of executive functioning. The Worry subscale has been found to not be related to either researcher-rated insight scales or neuropsychological tests (Marks, et al., 2000).

Behavioral Functioning Measures

1) Nurse Observation Scale for Inpatient Evaluation (NOSIE-30; Honigfeld, Roderic, & Klett, 1966). The NOSIE is a 30-item behavioral checklist format completed by nursing staff at CTP, with responses ranging from zero (never) to four (always). Two psychiatric technicians completed the checklist weekly for each participant. Items cover six areas of unit (öwardö) functioning: social competence (örefuses to do ordinary things expected of him or herö), social interest (ötries to be friendly with othersö), neatness (ökeeps clothes neatö), irritability (ögets angry or easily annoyedö), psychoticism (ötalks, mutters, or mumbles to selfö), and motor retardation (öis slow-moving or sluggishö). When combined, these six areas represent a total assets score. Adaptive functioning scales (i.e., social interest) were positively weighted and maladaptive scales (i.e., irritability) were negatively weighted when determining the total assets score. This measure has been widely used as part of the psychiatric rehabilitation treatment and was a routine assessment in the treatment program. Analyses within the CTP population have yielded Pearson correlations between 0.68 and 0.72 for all scales (Penn, Mueser, Spaulding, Hope, & Reed, 1995; Spaulding et al., 1999b). Also, a more recent reliability update of this measure confirmed it remains reliable in modern treatment settings, with inter-rater reliability on the total assets score at 0.76, on maladaptive scales at 0.68, and adaptive scales at 0.75 (Lyll, Hawley, & Scott, 2004).

Symptomatology Measures Over Course of Treatment

1) The Brief Psychiatric Rating Scale ó Extended Versions (BPRS-E; Lukoff, Nuechterlein, & Ventura, 1986; Van der Does, Linszen, Dingemans, Nugter, & Scholte, 1993). Used to evaluate symptoms, the BPRS-E was used routinely every six months at CTP. In general, the BPRS-E is widely used to assess changes in psychiatric symptoms. Using a Likert scale from 1 (symptom is not present) to 7 (symptom is very severe), clinicians rated individuals based on interview content and general, observed behavior on 24 items. The BPRS-E is a widely used instrument and reliability and validity have been demonstrated (Bailey, Lachar, Rhoades, Diefenbach, Espadas, & Varner, 2004). Factor analyses on the former 18-item version and the newer 24-item version have yielded four, five, and six factor solutions of symptom items (e.g., Spaulding, Reed, Sullivan, Richardson, & Weiler, 1999a; Perlick, et al., 1999; Burger et al., 1997). The six-factor solution validated by Spaulding, Fleming, Reed, Sullivan, Storzbach, & Lam (1999a) was used in this study because the original validation took place with the same population at CTP. A standard principal component analysis of the BPRS (Spaulding et al., 1999a) yielded six factors: Psychotic Disorganization, Hallucinations/Delusions, Paranoia, Emotional Blunting, Agitation/Elation and Anxiety/Depression.

Symptomatology Measures at Admission

The following assessments were administered at time of admission to CTP and as deemed necessary by the treatment team. In order to obtain the largest sample, only admission scores on the following assessments was used.

1) The Beck Hopelessness Scale (BHS) (Beck & Steer, 1988). The BHS is a self-report instrument that consists of 20 true-false statements designed to assess the extent of

positive and negative beliefs about the future during the previous week and takes less than five minutes to complete. Each of the 20 statements was scored 0 or 1. A total score was calculated by summing the pessimistic responses for each of the 20 items. The BHS has been standardized using psychiatric inpatients and outpatients (Beck et al., 1974; Beck & Steer, 1988). Beck and Steer (1988) report high internal reliability across diverse clinical and nonclinical populations with reliabilities ranging from .87 to .93. The BHS has adequate one-week test-retest reliability in a psychiatric outpatient sample ($r = .69$; Beck & Steer, 1988). Correlation coefficients between the BHS and the Beck Depression Inventory pessimism item range from .42 to .64 in clinical samples (Beck & Steer, 1988).

2) Beck Depression Inventory-II (BDI-II) (Beck, Steer, & Brown, 1996). The BDI-II is a 21-item self-report assessment of depressive symptoms. The respondent was asked to rate how much he or she has been bothered by each symptom on a 4-point scale ranging from 0 to 3. Each item consists of four statements that reflect gradations in the intensity of a particular depressive symptom. The respondent chooses the statement that best corresponds to the way that he or she has felt for the past two weeks. The psychometric properties of the inventory have been reviewed by Beck, Steer, and Brown (1996).

3) Suicide Probability Scale (SPS) (Cull & Gill, 1988). The SPS is a 36-item self-report measure of current suicidal ideation, hopelessness, negative self-evaluation, and hostility that takes approximately 10 minutes to administer. Subjects answered each item on a 4-point scale ranging from 1 (‘‘None or a little of the time’’) to 4 (‘‘Most or all of the time’’). There are three summary scores: A Suicide Probability Score, a total weighted score and a normalized T-score. The Suicide Probability Score can be adjusted to reflect

different a priori base rates for particular clinical populations. The internal reliability for the SPS is high (Cronbach alpha = .93) and has high test-retest reliability over a three-week period ($r = .92$; Cull & Gill, 1988). Although the SPS was designed to be a measure of suicide risk, there is a paucity of research studies that have tested the predictive validity of this measure.

Outcome Measures

1) Rehospitalization rate. One of the primary goals, often inappropriately cited as the only goal, of treatment programs is the prevention of future hospitalizations and the decrease of inpatient hospital days and use of emergency services (Cook, Pickett, Razzano, Fitzgibbon, Jonikas, & Cohler, 1996; Anthony, Cohen, & Vitalo, 1978). The inclusion of data from LCCMHC was a critical part of analyzing outcome from the CTP program since a majority of participants were served through LCCMHC upon discharge from CTP. Program evaluation activity at the LCCMHC has in the past, and hopefully in the future, established a rehospitalization data-tracking program. The data available for participants prior to 2005 was cross-checked with chart reviews at CTP and chart reviews and interviews with staff at LCCMHC. The data for participants from 2005-2009 was obtained and/or cross-checked from LCCMHC. Since the archival database in this study included people discharged from CTP in 1996 to people discharged from CTP in 2009, people may range in the possible amount of time since discharge. This study explored ways of addressing this such as the percentage of hospital days out of all hospitals and non-hospital days since discharge and the survival rate (or how long before the first rehospitalization).

2) Discharge disposition. The restrictiveness and nature of the setting to which individuals were discharged from the CTP can also serve as a measure of outcome. There were at least 35 different discharge locations to which people went following CTP. Less restrictiveness (e.g., independent living) was characteristic of better outcome and less symptomatology. For the purposes of this study, there were essentially four categories of discharge location from most restrictive to least restrictive: 1) Regional Center transfer, 2) Psychiatric Residential Rehabilitation, 3) Assisted Living, and 4) Independent living. These categories were based on interviews with the CTP program director and CTP social workers who were primarily responsible for discharge planning and most familiar with community services as part of previous research in the construction of the archival database. These four categories did not encompass all discharges from CTP such as those to nursing homes or developmentally disabled (DD) group homes. Few individuals were discharged to nursing homes, which were considered more restrictive than even psychiatric residential rehabilitation. Discharges to nursing homes from CTP were rare and typically due to the persons' medical rather than psychiatric condition, and therefore were not included in analysis. Similarly, very few people were discharged to DD settings because one of the exclusion criteria from admission to CTP was developmental disability.

Procedure

Data Collection

An archival longitudinal database was used in this study. Approval for construction of this deidentified archival clinical data for research purposes was obtained

from the university IRB and the state hospital research committee. Most of the archival data came from the CTP clinical archives, but additional data pertaining to outcome and community functioning was collected from Lancaster County Community Mental Health Center. Clinical data was routinely collected as part of the CTP program and contributed significantly to the database.

In addition, extensive chart review at both CTP and LCCMHC added additional data regarding hospitalization history and general clinical and demographic characteristics. In addition to the initial interview in which participants were asked about what mental health services they received before the age of 18, a large amount of historical information was gathered from a review of participants' past records and social history reports. Prior mental health services usage before the age of 18, coded as no service use vs. service use, was derived from whether or not a participant received any type of mental health services before age 18 (therapy, medications used for mental illnesses, special behavioral school classes or programs, institutionalization during adolescence, etc.). These prior history variables addressed whether or not the participant had any DSM diagnosis, displayed prolonged characteristics pertinent to the schizophrenia spectrum prior to age 18, or if they could have been termed as having a serious emotional disturbance during pre-adolescence or adolescence.

Information regarding symptoms at onset, whether or not the patient had premorbid behavioral problems, and number of previous hospitalizations was also obtained. Although instances of violence and delinquency in adolescents were noted, individuals were not assessed as having SED or APP in adolescence simply due to delinquency or aggression. These variables, taken into account with the other historical

variables, were used to assess and categorize individuals who used services prior to the age of 18 into different APP severity level groupings. Those participants that did not receive services or have any historical variables indicating APP were coded as not having APP in adolescence. Individuals with one or two instances of mental health service use (i.e. met with a mental health provider once for evaluation,) were coded as the low severity APP group while individuals with more frequent or longer duration of mental health services prior to the age of 18 were coded as the moderate APP group. Those participants that were institutionalized during childhood and or adolescents and had received extensive mental health services at that time were coded as the high severity APP group. , Data from 354 participants discharged from the CTP program between 1996 and March 2009 were used in analyses.

Assessments at CTP. All participants completed a comprehensive clinical assessment upon admission to the CTP program and most of these assessments were repeated at six month intervals throughout a person's hospitalization in order to monitor treatment response and inform future treatment planning decisions. These assessments primarily included measures of neurocognitive and social cognitive functioning. Clinical psychology graduate students or trained clinical assistants administered and scored all measures according to standardized instructions. Scoring was assisted by several computerized scoring programs. One significant change occurred in the routine assessment battery during the thirteen-year period from which the archival data was extracted. The RBANS, the IS, and social cognitive measures were added to the assessment battery between 2000 and 2001 when new admissions arrived during that time. The RAVLT, and SAIQ were phased out at that time in order to maintain that the

assessment battery could be completed in a manageable amount of time or in favor of newer assessment measures with stronger psychometric properties. Therefore, individuals discharged before 2000 did not have RBANS and social cognitive data. Likewise, people who entered the program at the time of the change did not have RAVLT, and SAIQ data. The RBANS was phased out in 2005-2006 and the NAB was used instead. Therefore, individuals discharged after 2000 and before 2006 had RBANS data while those participants entering the program between 2006 and 2009 had NAB data. Results using the RBANS and NAB were analyzed separately. Assessments continued to be collected at six-month intervals and therefore biannual neurocognitive and/or social cognitive data, as well as insight data was available for most participants. Analyses were conducted for the different APP severity groups on the various psychological assessments available at CTP during those participant's involvement with the program. Comparisons between and within groups were conducted across the different domains of assessments for different time points.

In addition, assessments of various areas of functioning and overall functioning were regularly completed in the context of the general milieu. Psychiatric technicians completed NOSIE assessments on a weekly basis. The monthly average of weekly ratings was included in the present database, making monthly NOSIE data pertaining to each person's unit functioning available. Data entry and management was completed by a trained clinical psychology graduate student on a monthly basis to be used in treatment progress meetings. At the time of extraction, the data was subjected to fidelity checks to monitor if the behaviors are being correctly recorded and contingencies implemented as intended by the treatment team.

The current study was a retrospective longitudinal study in which three assessments were taken from all participants in the study during the intake within two weeks of admission, at 6 months, and a year after admission. The 12 month time period was selected because the average length of stay at CTP is around 12 to 14 months which maximizes the amount of data available at any given time point. The greater part of treatment in skills training is also delivered within this time frame. All participants were interviewed and given the BHS, BDI-II, socialcognition battery, neurocognition battery, BPRS, and insight measures within the first two-weeks of their stay in the in-patient unit. Patients were interviewed again at 6-months for their semi-annual review, and at one-year for their annual review and given the BPRS, socialcognition and neurocognition batteries, and insight measures at each time point. The NOSIE was completed by psychiatric technicians weekly for each participant. Average monthly scores were computed for the six areas and total assets score on the NOSIE. NOSIE scores within two weeks of admission, after six months of treatment, and after one year of treatment were used in the present study.

LCCMHC. Comprehensive chart review and interviews with LCCMHC staff were completed by a clinical psychology graduate student involved in program evaluation activity as part of a practicum placement at LCCMHC. After collecting hospitalization data, the data was cross-checked with LRC records through chart review to ensure its accuracy. Not all CTP participants were served by LCCMHC upon discharge, nor has discharge data been obtained for those individuals served from 2006-2009. Whenever possible data for these individuals was obtained through LRC records.

After the archival database was completed with data from both settings, two graduate students completed additional quality assurance checks. Data was subjected to cross-checking with original and computerized archival data to ensure its reliability and accuracy. Once outcome data was obtained or completed for those individuals served from 2006 to 2009 the same methods described above were used to ensure reliability and accuracy.

Data Cleaning

Before analyses, data was examined for skewness and potential outliers. It was necessary to ensure normal distribution of the dependent variables because most of the analyses that were used in this study assume normality. Distributional skewing and asymmetrical outliers can both produce skew and therefore transformation and/or outlier windsorizing was applied only after examining the nature of the skew. Any dependent variables that demonstrated a skewed distribution (skewness $> +/- 1.00$) without outliers was normalized using conservative transformation procedures. Using Tukey's Hinges, the data was systematically screened for outliers. All outliers were included after a windsorizing procedure which replaces extreme values with the highest acceptable value was applied (Hoaglin, Mosteller & Tukey, 1983). All variables were cleaned to within acceptable skewness range with the exception of the rehospitalization and previous hospitalizations data. A few variables required both square root transformation and windsorizing of extreme outliers in order to normalize the distribution. Using the above procedures, all variables were cleaned to within acceptable skewness range with the exception of the rehospitalization data. The nature of this data prevented transformation or windsorizing. Therefore, the rehospitalization data were analyzed disregarding skew.

However, follow-up analyses were conducted by creating categorical variables to further corroborate the results.

Data Analysis

This study was exploratory in nature. In general, the study aimed to determine if there are differences between those who have different levels of adolescent psychiatric pathology and those who do not, and in particular to follow-up preliminary research (Wynne, 2009). The thirteen hypotheses of this study are evaluated with combinations of group contrast and correlational analyses. A hierarchical strategy for data analysis was used, starting with omnibus multivariate analyses of variance (MANOVA) performed within the domains of measurement selected for analysis (clinical, cognitive, behavioral, etc), then continuing to ANOVAs on specific measures and scales, followed by planned comparison cell contrasts, and finally correlational analysis. Analysis of variance (ANOVA) and its variants were the primary statistic used in order to replicate earlier results (Wynne, 2009). One-way ANOVAs with time 1 or time 2 covariates were used whenever possible. Whether or not individuals used services in adolescence, and further analyses utilizing Adolescent psychiatric pathology groupings were the between-subjects factor and each set of measures were the within-subjects factors. One tailed t-tests were used in analyses because of specific directional hypotheses. In addition, evaluating the relationships between categorical variables was accomplished through Chi square analysis. Also, results were compared across the different assessment measures used at different time periods at CTP, possibly providing results for future results on different factors or clusters of neurocognitive, socialcognitive, symptom, and insight measures that can be grouped together to better capture the multidimensional, and perhaps

developmental nature of insight in individuals who had varying levels of APP and are now participating in psychiatric rehabilitation as adults.

Because this is an exploratory study of a very complex database and there was an unusual amount of missing data, appropriate use of MANOVA was prevented.

Significant omnibus MANOVA results were not strictly interpreted as necessary for further group comparisons. Since the specific analyses relate to the respective study hypotheses in complex ways, the results will be organized according to the hypotheses, with the relevant analyses identified and described for the respective hypotheses in turn.

Unequal n correction. Due to the significant differences in the number of participants in each condition, a correction of unequal n s was performed. The most accepted procedure that best fits the collected data was used during analysis.

CHAPTER 3 ó RESULTS

The overall purpose of this study is to determine if there is a pattern of differential functioning for adults with SMI with or without a history of using mental health services in adolescence, and further delineated into APP severity groupings, across various levels of functioning. Before proceeding to evaluate the hypotheses, descriptive analyses of the demographic and clinical characteristics of people with different amounts of service use during adolescence are described.

Demographic and Clinical Characteristics of People at CTP

Of the 320 participants in this study, 171 (53.4%) used or had contact with mental health services during adolescence, see Tables 1-3. Of the people who did not use services during adolescence, 63.1% ($n = 94$) are male and 35.6% ($n = 53$) are female. Of the people who did use services 64.1% ($n = 109$) are male and 35.9% ($n = 61$) are female; the difference in gender proportions between service users and non-users is not statistically significant, $X^2(1) = 0.001, p = .98$.

Analyses using the No Service Use in adolescence vs. Service Use in adolescence distinction reveals a pattern of relationships among several demographic variables. There is a significant relationship between diagnostic subtype and mental health service use, $X^2(8) = 19.14, p = .01$ (see Table 4). Those diagnosed with Schizophrenia, Chronic/Undifferentiated Type, Schizoaffective Disorder, and öotherö are more likely to have used mental health services during adolescence than those diagnosed with Schizophrenia, Paranoid Type. Also, there is a significant relationship between Axis II diagnostic subtype and service use, $X^2(9) = 18.14, p = .03$ (see Table 5). Those diagnosed with Paranoid Personality Disorder are less likely to have received services

during adolescence than those diagnosed with Borderline Personality Disorder. Those diagnosed with Borderline Personality Disorder are more likely to have received services during adolescence than those without an Axis II diagnosis, which are more likely to have not used services during adolescence.

There is not a significant relationship between race and service use, $X^2(5) = 2.37$, $p = .797$ (see Table 6). When those participants identified as African American, Hispanic, Asian American, or Other are combined into one category and compared to those identified as Caucasian, or "White," there is not a significant relationship with service use, $X^2(1) = .007$, $p = 0.93$. This suggests that people who are "Non-White" are just as likely as those who are "White" to receive or have access to mental health services during adolescence. There is not a significant relationship between marital status and service use, $X^2(4) = 6.29$, $p = .18$ (see Table 7).

Finally, analyses between service use and age, length of stay at CTP, age at first hospitalization, number of previous hospitalizations in lifetime, and years of education reveal significant relationships. The means and standard deviations for each group for these variables are shown in Table 8. Analyses using one-way ANOVAs reveal a significant difference between groups in age $F(1, 306) = 42.20$, $p < .001$; number of years of education, $F(1, 281) = 24.31$, $p < .001$; the number of previous hospitalizations, $F(1, 296) = 4.46$, $p = .04$; and in the age of onset, $F(1, 246) = 71.82$, $p < .001$. There is no significant difference between groups on length of stay, $F(1, 135) = 2.33$, $p = .13$.

To summarize the results of the descriptive analyses, those with service use during adolescence tend to be younger at admission to CTP, have slightly lower levels of education, and have an earlier age of onset and more previous hospitalizations, as

expected. A relationship between Axis I and Axis II diagnoses and service use was identified. Those diagnosed with Schizophrenia, Chronic/Undifferentiated Type, Schizoaffective Disorder, and other are more likely to have used mental health services during adolescence than those diagnosed with Schizophrenia, Paranoid Type. Those diagnosed with Paranoid Personality Disorder are less likely to have received services during adolescence than those diagnosed with Borderline Personality Disorder; whereas those diagnosed with Borderline Personality Disorder are more likely to have received services during adolescence with other or no diagnosis. These results are consistent with expectations previously outlined and support the reliability of the distinctions made between who did or did not use mental health services prior to the age of 18 used in this study

Hypothesis 1

The first hypothesis of this study is to determine if there is a pattern of differential functioning between adults with SMI with or without mental health service use during adolescence across various levels of functioning both at admission and across treatment. In general, it is anticipated that individuals with adolescent psychiatric pathology will demonstrate lower overall functioning than those without adolescent psychiatric pathology. Because of this, hypothesis 1 predicts that those individuals who used mental health services in adolescence will show demonstrably lower overall functioning in the clinical setting than individuals who did not use mental health services during adolescence. Thus, it is hypothesized that the service usage will correlate with clinical functioning. Specifically, the groups (No Service Use vs. Service Use) will differ across

all domains of functioning measured (neurocognitive, social cognitive, symptomatology, and behavioral functioning) at admission and over the course of treatment.

Functioning at Admission

Neurocognitive Functioning

The RBANS, RCFT, RAVLT, COGLAB Card Sort, Trails A & B, COWAT/FAS, and NAB were included in analyses as measures of neurocognitive functioning.

Bivariate correlations between whether or not individuals used mental health services prior to age 18 and the neurocognitive variables included in analyses are shown in Table 9.

Between groups ANOVAs were conducted to maximize the power available to detect any differences between individuals who used mental health services in adolescence vs. those who did not use services on the neurocognitive variables (Tables 10 and 11). Contrary to all hypotheses, on all measures of neurocognition individuals with SMI who used services prior to the age of 18 do not demonstrate poorer neurocognitive functioning than those who did not use services, all $F_s < 3.33$, all $p_s > .07$. Of the 14 tests, one significant test is expected. Two reached trend level, $p = .07-.08$. However, when planned comparisons were made as expected better performance on RCFT Recognition was attained by individuals who did not use services prior to the age of 18 ($M = 20.54$) than by individuals who had used services prior to age 18 ($M = 18.17$), $t(71) = 1.82$, $p = .04$. Furthermore, as hypothesized better performance on Trails B was achieved by individuals without service use prior to age 18 ($M = 105.14$) than by individuals who had used services prior to age 18 ($M = 125.24$), $t(114) = -1.63$, $p = .05$.

Also as hypothesized, scores on RBANS Total were greater for individuals without service usage in adolescence ($M = 75.50$) than by individuals who had used services prior to age 18 ($M = 69.74$), $t(92) = 1.77$, $p = .04$. These significant planned comparisons indicate analyses utilizing ANOVAs did not have enough power in some instances to detect significant differences.

Furthermore, when individuals who used services prior to age 18 are grouped based on adolescent psychiatric pathology, APP severity groups do not differ in statistically significant ways on neurocognitive measures at admission, all F s < 2.18 , all p s $> .10$ (Table 11). Of the 14 tests, one significant test is expected. However, no tests reached trend level. Overall, contrary to hypotheses, these results suggest that participants with low, medium, and high APP severities do not significantly differ from those without APP at the time of admission, with respect to neurocognitive functioning.

Social cognitive functioning

The I-SEE (FKK), a measure of attributional style, the Hinting Task, a measure of theory of mind, the IPSAQ, a measure of interpersonal attributional style, and the CST, a measure of coping-related cognition, were used to examine social cognitive differences between individuals with SMI who used services in adolescence and those who did not use services prior to the age of 18. It is hypothesized that people who used services prior to the age of 18 would demonstrate lower overall sociocognitive functioning than those who did not use services. Furthermore, when analyses utilize APP severity groupings it is hypothesized that individuals in the high APP severity group will have the lowest

scores on social cognition measures at admission. Bivariate correlations can be found in Table 12.

Despite not having any significant MANOVA results, univariate ANOVAs were conducted to maximize the power available to detect any differences in the neurocognitive variables because an extremely low number of valid cases in the MANOVA analyses may have prevented any significant differences from being detected. Exploratory analyses using one-way between groups ANOVAs reveal no significant differences between those who did and did not use services prior to the age of 18 on measures of social cognition at admission, all F s < 2.58, all p s > .12 (Table 13). Of the 17 tests, one significant test is expected. However, no tests reached trend level. When planned comparisons were completed, no significant differences exist between the groups.

Furthermore, when APP severity grouping were considered in analyses groups do not differ in statistically significant ways on social cognition measures at admission, all F s < 2.49, all p s > .07 (Table 14). Of the 14 tests, one significant test is expected. Two reached trend level, $p = .07-.08$. Overall, contrary to hypotheses, these results suggest that participants who used mental health services in adolescence do not significantly differ from those without service use in adolescence at the time of admission, with respect to social cognitive functioning. In addition, there do not appear to be substantial differences in social cognitive functioning at admission for the various APP severity levels.

Insight

The Birchwood Insight Scale (IS), a measure of insight representing David's (1990) three domains of insight, and the Self-Appraisal of Illness Questionnaire, a measure designed to assess attitudes towards mental illness among people receiving psychiatric treatment, were used to examine insight differences between individuals with SMI who used services in adolescence and those who did not use services prior to the age of 18. It is hypothesized that because of prior contact with services, those individuals with contact with mental health services in adolescence have higher levels of insight into their disorder at admission than those who did not use mental health services in adolescence. Furthermore, when analyses utilize APP severity groupings, it is hypothesized that individuals in the high APP severity group would have greater insight scores at admission than individuals with no, low, or medium APP. The bivariate correlation matrix for the insight measures and their subscales can be found in Table 15.

Exploratory analyses using one-way ANOVAs reveal no significant differences between those who did and did not use services prior to the age of 18 on measures of insight at admission, all $F_s < 2.56$, all $p_s > .13$ (Table 16). Of the seven tests, one significant test is expected yet none reached trend level. When planned comparisons were completed, as expected scores on SAIQ Presence/Outcome of Illness subscale scores were greater for individuals with service usage in adolescence ($M = 14.29$) than for individuals who did not use services prior to age 18 ($M = 12.20$), $t(15) = -1.60$, $p = .05$.

Furthermore, analyses utilizing APP severity groupings indicate APP severity groups do not differ in statistically significant ways on insight measures at admission, all $F_s < 1.59$, all $p_s > .20$ (Table 17). Overall, contrary to hypotheses, these results suggest that participants who used mental health services prior to the age of 18 do not

significantly differ from those who did not use services in adolescence at the time of admission, with respect to insight. In addition, there are not significant differences in insight at admission for the various APP severity levels. Note in Table 17 one significant F test is expected and only one reached trend level at $p = .06 - .08$.

Behavioral functioning

The NOSIE total assets score was used as a measure of general behavioral functioning. Bivariate correlation of NOSIE subscales can be found in Table 18. It is expected that those who used mental health services prior to the age of 18 would demonstrate poorer behavioral functioning at admission than those who did not use services in adolescence. Contrary to the hypothesis, a one-way ANOVA does not reveal a significant difference, $F(1, 119) = 2.46, p = .12$, on total assets between individuals without mental health service usage in adolescence ($M = 157.68, SD = 24.13$) and those who used mental health services in adolescence ($M = 150.50, SD = 26.26$) (Table 18). Of the seven tests, one significant test is expected yet none reached trend level. When planned comparisons were completed, no significant differences between those with mental health service usage in adolescence and those without mental health service usage in adolescence.

In addition, it is hypothesized that individuals in the high APP severity group demonstrate poorer functioning on behavioral functioning measures. When NOSIE total assets are examined across APP severity groups using a one-way ANOVA, contrary to hypotheses individuals with more severe adolescent psychiatric pathology do not

demonstrate poorer behavioral functioning at admission than those individuals with none, low, and medium APP, $F(3, 117) = 1.60, p = .19$ (Table 19).

Because the NOSIE is comprised of three adaptive functioning scales and three maladaptive functioning scales, further exploratory analyses were conducted to determine if differences were apparent across all areas of functioning assessed by the NOSIE or whether the differences were specific to particular subscales. Pearson correlations reveal significant intercorrelations among all subscales and the total assets score. The bivariate correlation matrix can be found in Table 17.

Follow-up analyses did not reveal significant differences between those with mental health service use prior to age 18 and those without mental health service use prior to age 18 on the NOSIE subscales at admission, all $F_s < 2.46$, all $p_s > .12$ (see Table 18). When NOSIE subscales were examined across APP severity groups using a one-way ANOVA, contrary to hypotheses individuals with more severe adolescent psychiatric pathology do not demonstrate poorer behavioral functioning on five of six subscales at admission than those individuals with none, low, and medium APP, all $F_s < 2.15$, all $p_s > .09$. Differences between APP severity groups are found for the NOSIE Motor Retardation subscale, $F(3, 117) = 2.70, p = .05$ (see Table 19 for means and standard deviations). However, contrary to hypotheses, individuals in the high APP group have lower NOSIE Motor Retardation scores, thus indicating better functioning, than those in the medium and low APP groups and had NOSIE Motor Retardation scores equivalent to those without APP (LSD minimum mean difference = 2.1).

Overall, results of NOSIE analyses do not support hypotheses that those with mental health service use in adolescence demonstrate poorer behavioral functioning upon

admission than those without mental health service use in adolescence. Furthermore, hypotheses are not supported that differences in behavioral functioning at admission exist between the different APP severity groupings. Although significant difference was found on the NOSIE Motor Retardation subscale, results were opposite to hypothesized and the high APP severity group received lower scores than individuals in the low and medium APP severity groups.

Symptomatology

It was hypothesized that individuals who used mental health services in adolescence would have a higher level of symptomatology at admission than those who did not use services prior to the age of 18. Contrary to hypothesis, a one-way ANOVA reveals no significant mean difference, $F(1, 154) = 0.10, p = .75$, in overall symptomatology at admission as measured by the BPRS total score between those who used mental health services prior to age 18 ($M = 48.29, SD = 14.62$) and those who did not use services prior to age 18 ($M = 47.60, SD = 12.15$). Also it was hypothesized that when APP severity groupings are considered those individuals with the most severe APP demonstrate a higher level of symptomatology than those with less severe APP. Contrary to hypothesis, there is no significant mean difference, $F(1, 154) = 0.10, p = .75$, in overall symptomatology between those without APP and the various levels of APP severity (Tables 22 and 23).

Six factor scores for the BPRS were computed in order to evaluate symptom groupings as opposed to the gross overall measure of symptomatology provided by the total score. The six factors used were Psychotic Disorganization,

Hallucinations/Delusions, Paranoia, Emotional Blunting, Agitation/Elation and Anxiety/Depression. One-way ANOVAs for each of the BPRS factor scores were conducted to determine whether differences existed between groups in symptom areas. There are no significant differences between individuals who used mental health services prior to the age of 18 and those who did not use services prior to age 18 for any of the symptom factor scores, all $F_s < 2.65$, all $p_s > .11$, see Table 22. Of the six tests, one significant test is expected yet none reached trend level. When planned comparisons were completed, contrary to hypothesis scores on BPRS Paranoia Factor scores were greater for individuals without service usage in adolescence ($M = 9.00$) than for individuals who did use services in adolescence ($M=8.14$), $t(155) = 1.63$, $p = .05$. Furthermore, when analyses consider differences between APP severity groupings, there are no significant mean differences for any symptom factor scores, all $F_s < 1.48$, all $p_s > .22$, see Table 23. Of the six tests, one significant test is expected yet none reached trend level.

Finally, one-way ANOVAs for each of the 24 BPRS items were conducted to determine if there were differences in any specific symptoms between individuals who used mental health services prior to age 18 and those individuals who did not use services. Two significant differences emerged. People who used mental health services in adolescence ($M = 1.79$, $SD = 1.37$) have a higher rating of Suicidality, $F(1, 156) = 10.36$, $p = .002$, than those who did not use mental health services in adolescence ($M = 1.23$, $SD = .62$). Although a significant difference occurs between the groups on Uncooperativeness, $F(1, 155) = 3.91$, $p = .05$, contrary to hypothesis those who did not use services in adolescence ($M = 1.83$, $SD = 1.23$) have a higher rating of Uncooperativeness than those individuals who used services in adolescence ($M = 1.49$,

$SD = .91$). Results of all the one-way ANOVAs can be found in Table 24. Of the 24 analyses, one is expected to be significant. Two analyses were significant and three analyses reached trend level, $p = .06$. When planned comparisons were completed, as hypothesized scores on BPRS Suicidality item scores were greater for individuals with service usage in adolescence ($M = 1.79$) than for individuals who did not use services in adolescence ($M = 1.23$), $t(156) = -3.22$, $p < .001$. Also, as hypothesized BPRS Guilt item scores were greater for individuals with service usage in adolescence ($M = 2.19$) than for individuals who did not use services in adolescence ($M = 1.84$), $t(156) = -1.69$, $p = .05$. As hypothesized, BPRS Hallucination item scores were greater for individuals with service usage in adolescence ($M = 2.41$) than for individuals who did not use services in adolescence ($M = 1.87$), $t(156) = -1.91$, $p = .03$. Also, as hypothesized BPRS Self-Neglect item scores were greater for individuals with service usage in adolescence ($M = 2.39$) than for individuals who did not use services in adolescence ($M = 2.05$), $t(156) = -1.89$, $p = .03$. Contrary to hypotheses, BPRS Grandiosity item scores were lower for individuals with service usage in adolescence ($M = 1.63$) than for individuals who did not use services in adolescence ($M = 2.13$), $t(156) = 1.88$, $p = .03$. Also contrary to hypotheses, BPRS Suspiciousness item scores were lower for individuals with service usage in adolescence ($M = 2.33$) than for individuals who did not use services in adolescence ($M = 2.77$), $t(156) = 1.62$, $p = .05$. Finally, contrary to hypotheses BPRS Uncooperativeness item scores were lower for individuals with service usage in adolescence ($M = 1.49$) than for individuals who did not use services in adolescence ($M = 1.83$), $t(155) = 1.95$, $p = .03$. Results of these planned comparisons indicated analyses using ANOVAs did not have enough power to detect significant differences.

One-way ANOVAs for each of the 24 BPRS items were conducted to determine if there are differences in any specific symptoms between those with no APP, low APP, medium APP, and high APP and two significant differences emerged. Partial support for hypotheses is found as people with medium or high APP have a higher rating on Suicidality, $F(3, 154) = 5.34, p = .002$, than those without APP or low APP. A significant difference is found between the groups on Disorientation, $F(3, 153) = 3.73, p = .01$. Contrary to hypothesis, those with no, medium, or high APP receive lower ratings of Disorientation than those with low APP. Results of all the one-way ANOVAs can be found in Table 24. Of the 24 analyses, one is expected to be significant. Two analyses were significant and no analyses reached trend level.

Further analyses of symptomatology measures at admission included the SPS, BDI-II, and the BHS. Again, it was anticipated that individuals who used services in adolescence would have a higher level of symptomatology at admission than individuals who did not use services in adolescence. One-way ANOVAs for the BDI-II, BHS, SPS Suicidality scale, SPS Negative Self-Evaluation scale, and the SPS Hostility scale at admission reveal no significant mean differences, all $F_s < 3.30$, all $p_s > .07$ (Table 22). Two significant differences emerge for the SPS Total score and the SPS Hopelessness scale score. As hypothesized, people who used mental health services in adolescence ($M = 61.02, SD = 10.22$) receive higher symptomatology scores as measured by the SPS Total score, $F(1, 122) = 7.41, p = .01$, than those who did not use mental health services in adolescence ($M = 56.02, SD = 10.19$), and further directional planned comparisons also found this result, $t(122) = -2.72, p < .01$. Also, as hypothesized, people who used mental health services in adolescence ($M = 57.48, SD = 13.09$) received higher

symptomatology scores as measured by the SPS Hopelessness score, $F(1, 122) = 3.76, p = .05$, than those who did not use mental health services in adolescence ($M = 53.33, SD = 10.42$), planned comparisons for this variable were also significant, $t(122) = -1.94, p = .03$. Further planned comparisons found as hypothesized SPS Hostility scores were higher for individuals with service usage in adolescence ($M = 56.88$) than for individuals who did not use services in adolescence ($M = 52.81$), $t(121) = -1.82, p = .04$.

One-way ANOVAs were also conducted to determine if there were differences in any specific symptoms between those with no APP, low APP, medium APP, and high APP and two significant differences emerged. Partial support for hypotheses is found as people with medium or high APP (which were equivalent to each other) are rated with a higher level of symptomatology as measured by the SPS Total score, $F(3, 120) = 3.34, p = .02$, than those without APP or low APP. Results of all the one-way ANOVAs and means and standard deviations can be found in Table 23.

Overall, results of analyses using the BPRS total, BPRS factor scores, and BPRS items suggest that only a significant relationship exists on two BPRS items for service use and level of symptomatology. The amount of significant differences between those who used services in adolescence and those who did not use services in adolescence is close to what is expected to occur by chance, thus lowering confidence that these differences occur due to specific differences between the groups on measures of symptomatology. Although those who used services prior to adolescence receive higher suicidality scores than those who did not use services, those who did not use services in adolescence are deemed more uncooperative than individuals who did use mental health services in adolescence. Planned comparisons found that individuals with service usage

in adolescence are rated higher on suicidality, guilt, hallucinations, and self-neglect. However, contrary to hypotheses results from planned comparisons indicate individuals with mental health service usage are rated lower on grandiosity, suspiciousness, and uncooperativeness. When analyses are conducted between APP severity level and BPRS total, BPRS factor scores, and BPRS items only two significant differences emerge. Again as hypothesized, those individuals with medium or high APP receive higher suicidality scores than those without APP or in the low APP severity group.

Results of analyses of symptomatology at admission using the BDI-II, BHS, and SPS Total and SPS scales suggest several differences exist. As hypothesized, individuals who had mental health services in adolescence demonstrate more symptomatology at admission than those who did not have services in adolescence as measured by the SPS Total and SPS Hopelessness scale scores. Planned comparisons also found that individuals with mental health service usage in adolescence have more hostility as measured by the SPS Hostility scale. When analyses are conducted between symptomatology measures and APP severity levels, partial support for hypotheses is found, as individuals with medium or high APP receive higher SPS Total scores than those with no or low APP.

Functioning Over the Course of Treatment

When analyzing difference over the course of treatment, it is predicted that overall differences in functioning and symptomatology, and will not remain over the course of treatment. It is hypothesized that as a result of psychiatric rehabilitation there will be an increase in neurocognitive functioning, insight into disorder, internal locus of

control, and behavioral functioning and a decrease in external locus of control and symptomatology for those who did and did not use services in adolescence. That is, it is hypothesized that both groups will show improvement in functioning over the course of psychiatric rehabilitation. However, it is predicted that those individuals who used services prior to age 18 will demonstrate higher symptomatology after 12 months of rehabilitation. Furthermore, when different levels of adolescent psychiatric pathology are used in analyses, it is predicted that as adolescent psychiatric pathology becomes more severe functioning will decrease and there will be higher symptomatology. Overall, while it is hypothesized there will be differences in a variety of domains it is also hypothesized that differences in insight upon admission will not remain one-year after beginning psychiatric rehabilitation.

Neurocognitive Functioning Across Treatment.

As described above, it was anticipated that while there will be difference in neurocognitive functioning between those who used mental health services in adolescence and those who did not use services in adolescence, neurocognitive functioning will improve over the course of treatment for both groups, and there will be no differences between groups on neurocognitive measures after one year of treatment. A 2 (service use) x 3 (assessment time) repeated measure ANOVA was completed for each measure.

There is not an interaction of assessment time and service use as they relate to WCST Correct score ($F(2, 72) = .64, p = .53$). Contrary to hypotheses, there is no main effect for WCST correct over time ($F(2, 72) = .66, p = .52$) or for service use ($F(1, 36) =$

1.35, $p = .25$). However, when planned comparisons were analyzed contrary to hypothesis WCST correct scores were higher for individuals with service usage in adolescence ($M = 24.66$) than for individuals who did not use services in adolescence ($M = 21.74$), $t(58) = -2.01$, $p = .02$. There is not an interaction between assessment time and service use as they relate to WCST Perseverative Errors ($F(2, 112) = 1.36$, $p = .26$). Contrary to hypotheses, there is no main effect of time for WCST Perseverative Errors ($F(2, 112) = 2.53$, $p = .08$). However, there is a main effect for service use ($F(1, 56) = 5.13$, $p = .03$), with more overall WCST Perseverative errors made by those who did not use services prior to age 18 ($M = 26.72$, $SD = 16.00$) than from those individuals who did use services prior to the age of 18 ($M = 20.52$, $SD = 11.58$) (LSD minimum mean difference = 3.90). However, this pattern of the main effect is only descriptive for performance after six months and one year of participating in a psychiatric rehabilitation program




Figure 3.1 Service Main Effect for Service Usage Groups on WCST Perseverative Error Score Across Treatment

There is no interaction between assessment time and service use as they relate to WCST Random Errors over time ($F(2, 112) = 0.17, p = .84$). As hypothesized, there is a main effect for WCST Random Errors over time ($F(2, 112) = 3.66, p = .03$), with more random errors made at admission than after six to twelve months of rehabilitation, representing an increase in neurocognitive functioning (LSD minimum mean difference = 2.98). This pattern is descriptive for both those that did and did not use mental health services in adolescence. As hypothesized, a main effect for service use is not found ($F(1, 56) = 0.71, p = .41$) and differences do not exist between those who did and did not use mental health services in adolescence after one year of treatment.




Figure 3.2 Time Main Effect for Service Usage Groups on WCST Random Error Score Across Treatment

There is not an interaction between assessment time and service use as they relate to WCST Consolidation Index scores ($F(2, 72) = .10, p = .91$). Contrary to hypothesis

there is no main effect for WCST Consolidation Index scores over time ($F(2, 72) = 1.92$, $p = .15$) or for service use ($F(1, 36) = .71$, $p = .40$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 26, 29 - 32.

There is an interaction of assessment time and service use as they relate to RAVLT scores ($F(2, 18) = 5.25$, $p = .02$, $Mse = 2.34$). The pattern of this interaction is that RAVLT performance increased over time as hypothesized for those that did not use mental health services in adolescence, whereas contrary to hypothesis individuals who did use services in adolescence demonstrate no significant difference on RAVLT scores from admission to 6 and scores decrease at one year of treatment (LSD minimum mean difference = 1.37). However, individuals who used mental health services in adolescence receive higher RAVLT scores at admission and six months than those who did not services in adolescence and as hypothesized this difference between groups does not exist after one year of treatment (Table 27, 33). Planned comparisons found no difference between groups at time one, however contrary to hypothesis RAVLT scores were higher for individuals with service usage in adolescence ($M = 9.50$) than for individuals who did not use services in adolescence ($M=6.43$), $t(30) = -2.61$, $p = .01$. Again, as hypothesized planned comparisons found no differences in RAVLT scores between groups after one year of treatment There is no main effect for RAVLT over time ($F(2, 18) = .24$, $p = .79$) or for service use ($F(1, 9) = 2.19$, $p = .17$).




Figure 3.3 Interaction Between Service Usage Groups on RAVLT Scores Across Treatment

Contrary to hypotheses, there is not an interaction of assessment time and service use as they relate to NAB total score ($F(2, 26) = .47, p = .63, Mse = 52.61$). Contrary to hypotheses, there was no main effect for NAB total scores over time ($F(2, 26) = 1.74, p = .20$) or for service use ($F(1, 13) = .28, p = .61$) (Tables 27 and 34).

There is no interaction of assessment time and service use as they relate to RCFT Copy score ($F(2, 60) = .19, p = .82, Mse = 18.89$) or RCFT Recognition score ($F(2, 60) = .63, p = .54, Mse = 11.58$). There is no main effect for RCFT Copy scores over time ($F(2, 60) = 1.75, p = .18$) or for RCFT Recognition scores over time ($F(2, 60) = .55, p = .58$). Also there is no main effects for service use for RCFT Copy scores ($F(1, 30) = .32, p = .58$) or RCFT Recognition scores ($F(1, 30) = .04, p = .85$).

There is no interaction between assessment time and service use as they relate to RCFT Immediate Memory scores over time ($F(2, 60) = 1.37, p = .26, Mse = 186.63$). As hypothesized, there is a main effect for RCFT Immediate Memory scores over time ($F(2, 60) = 4.62, p = .01$), with no significant difference on scores from admission to six months however improvement in scores occurred at one year (LSD minimum mean difference = 6.83). However, this pattern is only descriptive for those who used mental health services in adolescence. A main effect for service use is not found ($F(1, 30) = 3.60, p = .07$).




Figure 3.4 Time Main Effect for Service Usage Groups on RCFT Immediate Memory Scores Across Treatment

There is no interaction between assessment time and service use as they relate to RCFT Delayed Memory scores over time ($F(2, 60) = .13, p = .88, Mse = 30.90$). As hypothesized, there is a main effect for RCFT Delayed Memory scores over time ($F(2, 60) = 4.89, p = .01$), with no significant difference on scores from admission to six

months and improvement in scores at one year (LSD minimum mean difference = 2.78). This pattern is descriptive for those who did and did not use services in adolescence. A main effect for service use is not found ($F(1, 30) = 1.69, p = .20$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 28, and 39 - 42.




Figure 3.5 Time Main Effect for Service Usage Groups on RCFT Delayed Memory Scores Across Treatment

There is an interaction of assessment time and service use as they relate to Trails A performance ($F(2, 110) = 3.38, p = .04, Mse = 165.13$). As hypothesized, the pattern of this interaction is that Trails A performance was slower, and thus worse, at admission than at 6 months and one year (which were equivalent to each another) (LSD minimum mean difference = 2.78) for those who used mental health services in adolescence. However, partial support is found for hypotheses for those who did not use mental health services prior to age 18, as their performance worsens from admission to six months, however improves from six to twelve months. At admission, those who did not use

mental health services prior to age 18 demonstrate poorer performance on Trails A than those who did not use mental health services in adolescence.. At six months, those who did not use services prior to age 18 demonstrate poorer performance than those who did use services prior to age 18, however, as hypothesized after one year of treatment differences in Trails A performance do not exist between the two groups. There is no overall main effect for Trails A over time ($F(2, 110) = .51, p = .60$) or for service use ($F(1, 55) = .004, p = .95$).




Figure 3.6 Interaction Between Service Usage Groups on Trails A Performance Across Treatment

There is no interaction between assessment time and service use as they relate to Trails B performance ($F(2, 110) = .53, p = .59$). Contrary to predictions, there is no overall main effect for Trails B over time and both groups do not demonstrate better performance over the course of treatment ($F(2, 110) = .12, p = .89$). There is also no

main effect for service use ($F(1, 55) = .20, p = .66$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 27, and 35-36.

There is an interaction between assessment time and service use as they relate to COWAT/FAS score over time ($F(2, 114) = 3.02, p = .05, Mse = 44.09$). Contrary to hypothesis, the pattern of this interaction is that although COWAT/FAS performance improves from six months to one year for those who did not use mental health services in adolescence, performance at one year is lower than at admission (LSD minimum mean difference = 1.41). However, for those who did use mental health services in adolescence performance is equivalent from admission to six months, however improves from six to twelve months. While those who did not use services prior to age 18 perform better on the COWAT/FAS at admission, this difference does not exist at six months. Contrary to hypotheses, after one year of treatment those who did use services prior to age 18 demonstrate better performance than those who did not use services prior to age 18. Also, contrary to hypotheses there is no overall main effect for COWAT/FAS over time ($F(2, 114) = 1.91, p = .15$) or for service use ($F(1, 57) = .06, p = .82$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 27 and 37.




Figure 3.7 Interaction Between Service Usage Groups on COWAT Scores Across Treatment

There is an interaction of assessment time and service use as they relate to RBANS Total score ($F(2, 108) = 3.65, p = .03, Mse = 51.18$). Partial support for hypotheses is found, as the pattern of this interaction is that RBANS Total score improves from admission to six months but does not improve from 6 months to one year for those who used mental health services prior to age 18, although scores at one year are better than at admission (LSD minimum mean difference = 3.82). For those who did not use services prior to age 18, performance does not improve from admission to six months, however improves from six to twelve months. As hypothesized, there is a main effect for overall RBANS Total score over time ($F(2, 108) = 9.82, p < .001$), with performance remaining the same from admission to six months but improving from six to twelve months of rehabilitation (LSD minimum mean difference = 2.70). This pattern is only descriptive for those who did not use services in adolescence. There is no overall main effect for service use ($F(1, 54) = 1.67, p = .20$). Results of repeated measure

ANOVAs and means and standard deviations can be found in Tables 27 and 38. When planned comparisons were completed, as hypothesized and previously stated individuals without mental health service usage had higher RBANS scores at admission than those who used mental health services in adolescence. However, contrary to hypotheses, differences existed between groups after one year of treatment. Individuals without mental health service usage in adolescence had higher RBANS scores ($M = 81.60$) than for individuals who did not use services in adolescence ($M=75.71$), $t(73) = 1.64$, $p = .05$.




Figure 3.8 Time Main Effect and Interaction Between Service Usage Groups on RBANS Total Scores Across Treatment

When 4 (APP Severity Level) x 3 (assessment time) repeated measures ANOVAs are conducted using APP severity levels, no interactions, assessment time main effects, or APP severity level main effects are found for the following neurocognition measures: WCST Correct, WCST Perseverative Errors, WCST Random Errors, WCST Consolidation Index, NAB Screener Total score, RCFT Copy, RCFT Recognition, Trails A and B, and COWAT/FAS (all $F_s < 2.66$, all $p_s > .06$). Contrary to hypotheses,

increases in neurocognitive functioning over treatment as measured by the previously mentioned measures are not found when analyses included APP severity levels. Also, contrary to hypotheses those with more severe levels of APP do not demonstrate poorer neurocognitive functioning on these measures at any assessment time. However, as hypothesized differences do not remain amongst the groups after one year of treatment. Results of repeated measure ANOVAs utilizing APP severity levels and means and standard deviations can be found in Tables 44 - 59.

There is an interaction between assessment time and APP severity level as they relate to RAVLT scores ($F(4, 16) = 3.88, p = .02, Mse = 2.11$). No individuals in the low APP severity group completed RAVLT assessments and therefore are not included in analyses. RAVLT performance for individuals with no APP does not change from admission to six months, however, performance increases from six to twelve months. For individuals in the medium APP severity group, RAVLT performance does not change during treatment, whereas for those in the high APP severity group RAVLT performance increases from admission to six months but decreases from six to twelve months (which was equivalent to performance at admission) (LSD minimum mean difference = 2.27). RAVLT performance at admission is equal for the no APP group and medium APP group and for the medium and high APP groups, however contrary to hypotheses individuals in the high APP severity group demonstrate better performance than those in the no APP group on the RAVLT at admission. At six months, contrary to hypotheses those in the high APP severity group demonstrate the highest performance on the RAVLT, followed by those in the medium APP severity group. As hypothesized, these differences in RAVLT performance do not remain after one year of treatment. There is no main effect

for RAVLT over time ($F(2, 16) = .15, p = .86$) or for APP severity level ($F(2, 8) = 1.06, p = .39$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 45, 50.




Figure 3.9 Interaction Between APP Severity Levels on RAVLT Scores Across Treatment

There is no interaction between assessment time and APP severity levels as they relate to RCFT Immediate Memory scores ($F(6, 56) = 1.76, p = .12, Mse = 175.91$). There is an overall main effect for RCFT Immediate Memory scores over time ($F(2, 56) = 5.66, p = .01$), with no significant difference on scores from admission to six months and partial support for hypotheses as there are improvements in performance from six to twelve months. While performance for all APP severity groups remains the same from admission to six months, improvements in RCFT Immediate Memory scores from six to twelve months only occur for those in the low and high APP severity levels (LSD minimum mean difference = 6.70). A main effect for APP severity level is not found

($F(3, 28) = 1.27, p = .31$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 46, 57.




Figure 3.10 Time Main Effect for APP Severity Levels on RCFT Immediate Memory Scores Across Treatment

There is no interaction between assessment time and APP severity levels as they relate to RCFT Delayed Memory scores ($F(6, 56) = 2.14, p = .06, Mse = 27.06$). There is an overall main effect for RCFT Delayed Memory scores over time and partial support for hypotheses ($F(2, 56) = 6.23, p < .01$), with no significant difference on scores from admission to six months and improvements in performance from six to twelve months. This pattern of performance is only descriptive for no APP and high APP severity groups (LSD minimum mean difference = 2.63). A main effect for APP severity level is not found ($F(3, 28) = .55, p = .65$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 46, 58.




Figure 3.11 Time Main Effect for APP Severity Levels on RCFT Delayed Memory Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to RBANS total scores ($F(6, 104) = 2.40, p = .03, Mse = 49.84$). RBANS performance for individuals with no or low APP does not change from admission to six months, however performance increases from six to twelve months. For individuals in the medium APP severity group, RBANS performance does not change during treatment, whereas for those in the high APP severity group RBANS performance increases from admission to six and twelve months (which were equivalent to each other) (LSD minimum mean difference = 5.34). RBANS performance at admission is equal for the no, low, and medium APP groups; however as hypothesized, individuals in the high APP severity group have lower RBANS total scores than those in the no APP group at admission. At six months, RBANS performance is equivalent for all APP severity

groups. After one year of treatment, contrary to hypotheses, those in the no and low APP severity groups have higher RBANS total scores than those in the medium and high APP severity groups and those in the high APP severity group have higher RBANS total scores than those in the medium APP severity group. There is an overall main effect for RBANS total score over time ($F(2, 104) = 6.37, p < .01$) with RBANS performance remaining the same from admission to six months but improving by one year of treatment. However, this main effect is only descriptive for individuals in the no and low APP severity groups. There is not a main effect for APP severity level ($F(3, 52) = .80, p = .50$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 45, 55.




Figure 3.12 Time Main Effect and Interaction Between APP Severity Levels on RBANS Total Scores Across Treatment

In summary, results of analyses between those who used mental health services prior to age 18 and those who did not use services and neurocognitive functioning at

admission and over the course of treatment suggest that improvements in neurocognitive functioning are evident over the course of treatment on WCST Random Errors, for individuals who did not use services prior to age 18 on the RAVLT, for individuals who did use services in adolescence on RCFT Immediate Memory, for both groups on RCFT Delayed Memory, for individuals that used services in adolescence on Trails A, for those who used services on adolescence on COWAT/FAS, and for both groups on RBANS total. Results of analyses between assessment time and APP severity level indicate improvements in neurocognitive functioning occur over the course of treatment on the RAVLT for those without APP, for those with low or high APP on RCFT Immediate Memory, for those with none or high APP on RCFT Delayed Memory, and for those in the none, low, or high groups on RBANS total performance.

When analyses are conducted between those who used services in adolescence and those who did not use services in adolescence, as hypothesized no differences exist between groups after one year of psychiatric rehabilitation on the following measures: WCST Correct, WCST Perseverative Errors, WCST Random Errors, WCST Consolidation Index, RCFT Copy, RCFT Recognition, RCT Immediate and Delayed Memory, NAB total, RAVLT, and Trails A and B performance. Contrary to hypotheses, individuals who used mental health services in adolescence demonstrate better performance on the COWAT/FAS after one year of treatment than individuals who did not use services in adolescence. However, individuals who did not use services in adolescence demonstrate poorer performance on the RBANS total after one year of treatment than those who did use mental health services in adolescence.

Finally, when analyses are conducted utilizing APP severity level, as hypothesized no differences exist between groups after one year of psychiatric rehabilitation on the following measures: WCST Correct, WCST Perseverative Errors, WCST Random Errors, WCST Consolidation Index, RCFT Copy, RCFT Recognition, RCT Immediate and Delayed Memory, NAB total, RAVLT, and Trails A and B performance. Contrary to hypotheses, individuals with high APP demonstrate better performance on the RAVLT at admission and six months, however as hypothesized this difference does not remain after one year of treatment. As hypothesized, those individuals with high APP demonstrate poorer performance on the RBANS total at admission. However, contrary to hypotheses difference remain between the groups at twelve months and those without APP or low APP perform better on the RBANS than those in the medium or high APP severity groups.

Social Cognitive Functioning Across Treatment

It is anticipated that individuals with and without mental health service usage in adolescence demonstrate an increase in functioning, a decrease in external locus of control, and an increase in internal locus of control. Furthermore, when different levels of adolescent psychiatric pathology are considered, it is hypothesized as APP becomes more severe functioning decreases, however any differences will not remain between groups after one year of treatment. A 2 (service use) x 3 (assessment time) repeated measure ANOVA was completed for each measure.

There is not interaction between assessment time and service use as they relate to Hinting scores over time ($F(2, 102) = .21, p = .81, Mse = 6.32$) (Table 61, 63). As

hypothesized, there is a main effect for Hinting scores over time ($F(2, 102) = 8.32, p < .001$), with improvement on scores from admission to six months and one year (which were equivalent to each other) (LSD minimum mean difference = .98). This pattern was descriptive for both those with and without mental health service usage in adolescence. A main effect for service use is not found ($F(1, 51) = .01, p = .93$).




Figure 3.13 Time Main Effect for Service Usage Groups on Hinting Task Scores Across Treatment

There is not an interaction between service use and assessment time as they relate to FKK Internal Locus of Control ($F(2, 50) = .80, p = .46, Mse = 21.14$). Contrary to hypotheses, there is no main effect for FKK Internal Locus of Control scores over time and internal locus of control does not increase over treatment for either group ($F(2, 50) = .27, p = .77$). There is also not a main effect for service use ($F(1, 25) = .06, p = .82$). There is not an interaction between service use and assessment time as they relate to FKK Self Concept scores ($F(2, 50) = .71, p = .50, Mse = 12.23$). Contrary to hypotheses, there is no main effect for FKK Self Concept, a subscale of FKK Internal Locus of Control, scores over time ($F(2, 50) = 1.14, p = .33$). There is not a main effect for service use

($F(1, 25) = .33, p = .57$). There is not an interaction of assessment time and service use as they relate to FKK Self Efficacy scores ($F(2, 50) = .48, p = .62, Mse = 44.46$). Contrary to hypotheses, there is no main effect for FKK Self Efficacy scores over time ($F(2, 50) = .11, p = .90$). There is no main effect for service use ($F(1, 25) = .25, p = .62$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 60, and 64 - 66.

There is not an interaction between assessment time and service use as they relate to FKK Powerful Others ($F(2, 50) = .90, p = .41, Mse = 18.42$). Contrary to hypotheses that performance on this subscale of FKK External Locus of Control would decrease over time, there is no main effect for FKK Powerful Others scores over time ($F(2, 50) = 1.04, p = .36$). There is also no main effect for service use ($F(1, 25) = .33, p = .57$).

There is an interaction of assessment time and service use as they relate to FKK Chance scores ($F(2, 50) = 5.29, p = .01, Mse = 19.93$). Partially supporting hypotheses, the pattern of this interaction is that FKK Chance scores remained the same across treatment for those without mental health service usage in adolescence, whereas as hypothesized FKK Chance scores were highest (more external locus of control) at admission than at 6 months to one year (which were equivalent to each other) for those who used mental health services in adolescence (LSD minimum mean difference = 3.47). While FKK Chance scores are equivalent to one another at admission for those with and without mental health service usage in adolescence, contrary to the hypothesis those without mental health service usage in adolescence have higher FKK Chance scores at six and twelve months than those who used mental health services in adolescence. Planned comparisons also found this result at six months, $t(46) = 1.87, p = .03$, and at

one year, $t(34) = 1.72, p = .05$. There is a main effect for overall FKK Chance score over time ($F(2, 50) = 4.66, p = .01$), as hypothesized FKK Chance scores are higher at admission than at six to twelve months of rehabilitation (which were equivalent to each other) (LSD minimum mean difference = 2.46). This pattern is only descriptive for those who used mental health services in adolescence. There is no overall main effect for service use ($F(1, 25) = 1.13, p = .30$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 60, 67 - 68.




Figure 3.14 Time Main Effect and Interaction Between Service Usage Groups on FKK Chance Subscale Scores Across Treatment

There is an interaction of assessment time and service use as they relate to FKK External Locus of Control score over time ($F(2, 50) = 3.87, p = .03, Mse = 51.98$). Contrary to the hypothesis, the pattern of this interaction is that FKK External Locus of Control scores remain the same across treatment for those who did not use mental health services in adolescence, whereas as hypothesized FKK Chance scores are highest at admission than at 6 months to one year (which are equivalent to each other) for those

who did use mental health services in adolescence (LSD minimum mean difference = 5.61). While FKK External Locus of Control scores are equivalent to one another at admission for those with and without mental health service usage prior to age 18, contrary to the hypothesis those without service usage in adolescence had higher FKK External Locus of Control scores at six and twelve months than those with APP. Planned comparisons only found this difference at six months, $t(46) = 1.90, p = .03$. There is a main effect for overall FKK External Locus of Control score over time ($F(2, 50) = 3.77, p = .03$), as hypothesized FKK External Locus of Control scores are higher at admission than at six to twelve months of rehabilitation (which are equivalent to each other) (LSD minimum mean difference = 3.97). This pattern is only descriptive for those who used mental health services in adolescence. There is no overall main effect for service use ($F(1, 25) = .74, p = .40$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 60 and 69.




Figure 3.15 Time Main Effect and Interaction Between Service Usage Groups on FKK Externality Scale Scores Across Treatment

There is not an interaction of assessment time and service use as they relate to IPSAQ Internal Positive score ($F(2, 30) = 2.85, p = .07, Mse = 5.67$). There is no main effect for IPSAQ Internal Positive scores over time ($F(2, 30) = .39, p = .68$) or for service use ($F(1, 15) = 2.15, p = .16$). There is no interaction between assessment time and service use as they relate to IPSAQ Personal Positive scores over time ($F(2, 30) = .93, p = .41, Mse = 2.25$). There is a main effect for IPSAQ Personal Positive scores over time ($F(2, 30) = 4.00, p = .03$), with no significant difference in scores from admission to six months and one year (which are equivalent to each other), however scores at 12 months are significantly less than at admission (LSD minimum mean difference = 1.0). This result indicates at one year individuals are less likely to attribute positive events to themselves than at admission. However, this pattern is descriptive for neither group and is therefore misleading as a general description. A main effect for service use is not found ($F(1, 15) = .51, p = .49$).




Figure 3.16 Time Main Effect for Service Usage Groups on IPSAQ Personal Positive Scale Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to IPSAQ Situational Positive ($F(2, 30) = 2.18, p = .13, Mse = 5.05$). There is no main effect for IPSAQ Situational Positive scores over time ($F(2, 30) = 2.32, p = .12$) or for service use ($F(1, 15) = 4.02, p = .06$). However, planned comparisons found contrary to hypothesis IPSAQ Situational Positive scores are higher for individuals without service usage in adolescence after one year of treatment ($M = 5.07$) than for individuals without service usage in adolescence ($M=3.55$), $t(23) = 1.68, p = .05$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 62, and 70 - 72.

There is an interaction between assessment time and service use as they relate to IPSAQ Internal Negative scores ($F(2, 30) = 3.41, p = .05, Mse = 5.97$). As hypothesized, the pattern of this interaction is that IPSAQ Internal Negative scores are highest at admission and six months (which are equivalent to each other (LSD minimum mean difference = 2.42) and decreases at twelve months for those without mental health service usage in adolescence. This result indicates that individuals are more likely to attribute negative events to people or themselves after one year of treatment than at admission or six months. However, contrary to hypotheses, for those with mental health service usage in adolescence scores remain the same across treatment. While IPSAQ Internal Negative scores are equivalent to each other for those with and without mental health service usage in adolescence at admission and six months, contrary to hypotheses at twelve months those with mental health service usage in adolescence have higher IPSAQ Internal Negative scores than those individuals who did not use services in adolescence, $t(23) = -$

2.18, $p = .02$. There is no overall main effect for IPSAQ Internal Negative scores over time ($F(2, 30) = .39, p = .68$) or for service use ($F(1, 15) = 1.41, p = .25$).




Figure 3.17 Interaction Between Service Usage Groups on IPSAQ Internal Negative Scale Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to IPSAQ Personal Negative scores ($F(2, 30) = .47, p = .63, Mse = 5.01$). There is no main effect for IPSAQ Personal Negative scores over time ($F(2, 30) = .33, p = .72$) or for service use ($F(1, 15) = .63, p = .44$). There is not an interaction between assessment time and service use as they relate to IPSAQ Situational Negative scores ($F(2, 30) = 2.14, p = .14, Mse = 5.18$). There is no main effect for IPSAQ Situational Negative scores over time ($F(2, 30) = .45, p = .64$) or for service use ($F(1, 15) = .69, p = .42$). When planned comparisons were completed, contrary to hypothesis IPSAQ Situational Negative scores were lower for individuals with service usage in adolescence ($M = 3.27$) than for individuals who did not use services in adolescence ($M = 6.21$), $t(23) = 3.16, p < .01$.

Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 62, and 73 - 75.

There is not an interaction of assessment time and service use as they relate to CST Social Support scores ($F(2, 40) = .74, p = .48, Mse = 44.41$). There is a main effect for overall CST Social Support scores over time ($F(2, 40) = 4.83, p = .01$). As hypothesized CST Social Support scores are higher after one year of treatment than at admission or six months (which are equivalent to each other) (LSD minimum mean difference = 4.06). However, this pattern is only descriptive for individuals who used services in adolescence and is misleading. There is no main effect for service use ($F(1, 20) = .21, p = .65$).




Figure 3.18 Time Main Effect for Service Usage Groups on CST Social Support Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to CST Self-Controlling scores ($F(2, 40) = .24, p = .79, Mse = 31.86$). There is no main effect for CST Self-Controlling scores over time ($F(2, 40) = .14, p = .87$). There is a

main effect for service use ($F(1, 20) = 7.05, p = .02$). Contrary to hypotheses, those individuals without mental health service usage in adolescence have overall higher CST Self-Controlling scores than individuals with mental health service usage in adolescence and this pattern is consistent across assessment times. However, when planned comparisons were completed individuals without mental health service usage in adolescence only demonstrate higher CST Self-Controlling scores after one year of treatment, $t(35) = 1.77, p = .04$.




Figure 3.19 Service Usage Group Main Effect for Service Usage Groups on CST Self-Controlling Scores Across Treatment

There is not an interaction of assessment time and service use as they relate to CST Escape Avoidance scores ($F(2, 40) = 1.92, p = .16, Mse = 35.23$). There is no main effect for CST Escape Avoidance scores over time ($F(2, 40) = 1.74, p = .19$) or for service use ($F(1, 20) = .13, p = .72$). Also, there is not an interaction between assessment time and service use as they relate to CST Planful Problem Solving scores ($F(2, 40) = .57, p = .57, Mse = 40.89$). There is no main effect for CST Planful Problem Solving

scores over time ($F(2, 40) = .47, p = .63$) or for service use ($F(1, 20) = 1.16, p = .29$).

Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 61, 76 - 79.

When 4 (APP Severity Level) x 3 (assessment time) repeated measures ANOVAs are conducted no interactions, assessment time main effects, or APP severity level main effects are found for the following social cognition measures: FKK Internal Locus of Control, FKK Self Concept, FKK Self Efficacy, FKK Powerful Others, IPSAQ Internal Positive, IPSAQ Personal Positive, IPSAQ Personal Negative, IPSAQ Situational Negative, CST Social Support, CST Escape Avoidance, and CST Planful Problem Solving (all $F_s < 2.71$, all $p_s > .06$) (Tables 80 ó 99).

There is no interaction between assessment time and APP severity level as they relate to Hinting Task scores ($F(6, 98) = .90, p = .50, Mse = 6.26$) (Table 81, 83). There is a main effect for Hinting scores over time as hypothesized ($F(2, 98) = 5.87, p < .01$), with improvement on scores from admission to six months and one year (which are equivalent to each other) (LSD minimum mean difference = .97). This pattern is descriptive for those with no APP. A main effect for APP severity level is not found ($F(3, 49) = .80, p = .50$).




Figure 3.20 Time Main Effect for APP Severity Levels on Hinting Task Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to FKK Chance score ($F(6, 46) = 2.88, p = .02, Mse = 19.09$). Although contrary to hypotheses FKK Chance scores remain the same across treatment for those without APP, as hypothesized FKK Chance scores are highest (more external locus of control) at admission than at 6 months to one year (which are equivalent to each other) for those with low and medium APP (LSD minimum mean difference = 3.47). For individuals in the high APP group, FKK Chance scores remain the same from admission to six months and from six months to one year, but scores at one year are lower than at admission as hypothesized. While FKK Chance scores are equivalent to one another at admission amongst the groups, those without APP have higher FKK Chance scores at six months than those in the low and medium APP groups. Individuals in the high APP group have higher FKK Chance scores at six months than those in the low and medium APP groups.

However, there is partial support for hypotheses as those individuals with more severe symptomatology have higher FKK Chance scores (more external locus of control) at six months than those with medium APP. At one year, FKK Chance scores for those in the low, medium, and high APP severity groups are equivalent while those in the no APP group have higher FKK Chance scores than those in the low and high APP groups. There is a main effect for overall FKK Chance score over time ($F(2, 46) = 7.81, p = .001$). As hypothesized FKK Chance scores are higher at admission than at six to twelve months of rehabilitation (which are equivalent to each other) (LSD minimum mean difference = 2.40). This pattern is only descriptive for those with low APP. There is no overall main effect for APP severity level ($F(3, 23) = .35, p = .79$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 80, 84 - 89.




Figure 3.21 Time Main Effect and Interaction between APP Severity Levels on FKK Chance Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to FKK External Locus of Control score ($F(6, 46) = 2.30, p = .05, Mse = 50.17$). Although, the pattern of this interaction is that FKK External Locus of Control scores remain the same across treatment for those without APP, partial support for hypotheses is demonstrated as FKK Chance scores are highest at admission than at 6 months to one year (which are equivalent to each other) for those with low or medium APP (LSD minimum mean difference = 7.79). For individuals with high APP, FKK External Locus of Control scores at one year are lower than at admission or six months (which are equivalent to each other). This result still supports hypothesizes that those in the high APP group demonstrate a decrease in external locus of control after one year of treatment. While FKK External Locus of Control scores are equivalent to one another at admission and one year for those with and without APP, contrary to the hypothesis those without APP have higher FKK External Locus of Control scores at six months than those with low or medium APP. However, partial support for hypotheses is found as individuals in the high APP group have higher FKK External Locus of Control scores at six months than individuals with low or medium APP. There is a main effect for overall FKK External Locus of Control score over time ($F(2, 46) = 5.47, p = .01$). As hypothesized FKK External Locus of Control scores are higher at admission than at six to twelve months of rehabilitation (which are equivalent to each other) (LSD minimum mean difference = 3.90). This pattern is only descriptive for those with low or medium APP. There is no overall main effect for APP severity level ($F(3, 23) = .29, p = .83$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 80 and 89.




Figure 3.22 Time Main Effect and Interaction between APP Severity Levels on FKK Externality Scores Across Treatment

There is not an interaction between assessment time and APP severity level as they relate to IPSAQ Situational Positive scores ($F(6, 26) = 1.77, p = .15, Mse = 4.74$). There is no main effect for IPSAQ Situational Positive scores over time ($F(2, 26) = 1.05, p = .37$). There is a main effect for APP severity level ($F(1, 13) = 4.23, p = .03$). Overall IPSAQ Situational Positive scores, attributing positive events to situational factors, are highest for those without APP and with high APP (which are equivalent to each other) than for individuals in medium APP groups. Individuals in the low APP groups have the lowest overall IPSAQ Situational Positive scores (LSD minimum mean difference = 1.40). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 82 and 92.

IPSAQ Situational Positive Subscale Scores Across Treatment




Figure 3.23 APP Severity Levels Main Effect on IPSAQ Situational Positive Subscale Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to IPSAQ Internal Negative scores ($F(6, 26) = 3.21, p = .02, Mse = 4.85$). The pattern of this interaction is that IPSAQ Internal Negative scores remain the same across treatment for those with no APP and medium APP (LSD minimum mean difference = 3.15). IPSAQ Internal Negative scores for those in the low APP group remain the same from admission and six months but increase by one year. For individuals with high APP, scores increase from admission to six and twelve months (which are equivalent to each other), indicating as treatment progresses those in the high APP severity group are more likely to attribute negative events to themselves or others than at admission. While IPSAQ Internal Negative scores are equivalent to each other for those with and without APP at admission, at six months those with no APP have higher scores than those in the low APP group. Those in the medium and high APP groups have higher scores at sixth

months than those in the low APP group. Contrary to hypotheses, by one year of treatment, those in the low, medium, and high APP groups have higher scores than those in the no APP group. There is no overall main effect for IPSAQ Internal Negative scores over time ($F(2, 26) = 1.20, p = .32$) or for APP severity level ($F(3, 13) = .66, p = .59$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 82 and 93.




Figure 3.24 Interaction between APP Severity Levels on IPSAQ Internal Negative Subscale Scores Across Treatment

There is not an interaction between assessment time and APP severity as they relate to CST Self-Controlling scores ($F(6, 36) = 1.81, p = .13, Mse = 27.52$). There is no main effect for CST Self-Controlling scores over time ($F(2, 36) = .48, p = .62$). There is a main effect for APP severity level ($F(3, 18) = 3.39, p = .04$). Those individuals without APP have overall higher CST Self-Controlling scores than individuals with low, medium, and high APP. Contrary to hypotheses, those with high APP have lower overall CST

Self-Controlling scores, indicating these individuals may utilize other coping strategies. However, this pattern is not consistent across treatment times except for after one year of treatment (Tables 81 and 97).




Figure 3.25 APP Severity Levels Main Effect on CST Self-Controlling Subscale Scores Across Treatment

In summary, results of analyses between those who used mental health services prior to age 18 and those who did not use services and socialcognitive functioning over the course of treatment suggest that improvements in sociocognitive functioning are evident for both groups over the course of treatment on the Hinting Task and for those with mental health service usage on CST Social Support. Contrary to hypotheses, there is not a decrease in FKK Powerful Others (a subscale of FKK External Locus of Control) over treatment. Furthermore, contrary to hypotheses there is not an increase in FKK Internal Locus of Control or FKK Self Concept or FKK Self Efficacy (both subscales of FKK Internal Locus of Control) over the course of treatment. As hypothesized, FKK

External Locus of Control and FKK Chance, a subscale of FKK External Locus of Control, do decrease over treatment; however, this only occurred for those who used mental health services in adolescence. Additionally, IPSAQ Internal Negative scores, part of a personalizing bias, decrease across treatment for those without mental health service usage in adolescence; however, this remains constant across treatment for those who used mental health services in adolescence.

Results of analyses between assessment time and APP severity level indicate improvements in sociocognitive functioning are over the course of treatment on the Hinting Task. Contrary to hypotheses, there is not a decrease in FKK Powerful Others (a subscale of FKK External Locus of Control) over treatment. Furthermore, contrary to hypotheses there is not an increase in FKK Internal Locus of Control or FKK Self Concept or FKK Self Efficacy (both subscales of FKK Internal Locus of Control) over the course of treatment. As hypothesized, FKK External Locus of Control decrease over treatment, however this only occurs for those in the low, medium, and high groups. For FKK Chance, a subscale of FKK External Locus of Control, decreases over treatment only occur for those with low or medium APP. Additionally, IPSAQ Internal Negative scores, part of a personalizing bias, increase across treatment for those with low or high APP and remain constant across treatment for those with medium or no APP.

As hypothesized, when analyses are conducted between those without and without mental health service usage in adolescence, no differences exist between groups after one year of psychiatric rehabilitation on the following measures: Hinting Task, FKK Internal Locus of Control, FKK Self Concept, FKK Self Efficacy, FKK Powerful Others, IPSAQ Internal Positive, IPSAQ Situational Positive, IPSAQ Personal Negative, IPSAQ

Situational Negative, CST Social Support, CST Escape Avoidance, and CST Planful Problem Solving. Contrary to hypotheses, individuals without mental health service usage prior to age 18 demonstrate higher scores on the FKK Chance scale and FKK External Locus of Control measures after one year of treatment than those without mental health service usage in adolescence. Also, contrary to hypotheses those without mental health service usage demonstrate lower scores after one year of treatment on the IPSAQ Internal Negative and higher scores on the CST Social Support scale than those with APP.

Finally, as hypothesized, when analyses are conducted utilizing APP severity level, as hypothesized no differences exist between groups after one year of psychiatric rehabilitation on the following measures: Hinting Task, FKK Internal Locus of Control, FKK Self Concept, FKK Self Efficacy, FKK Powerful Others, FKK External Locus of Control, IPSAQ Internal Positive, IPSAQ Personal Positive, IPSAQ Personal Negative, IPSAQ Situational Negative, CST Social Support, CST Escape Avoidance, CST Self Controlling, and CST Planful Problem Solving. Contrary to hypotheses, individuals without APP demonstrate higher scores on the FKK Chance scale after one year of treatment than those with low or high APP. Also, contrary to hypotheses those without APP demonstrate lower IPSAQ Internal Negative scores after twelve months than individuals with low, medium, or high APP (which are all equivalent).

Insight Over the Course of Treatment

It is anticipated that adults with SMI who had mental health services before the age of 18 have more insight into their mental illness over the course of treatment than

those that did not have mental health services prior to the age of 18. Furthermore, it is predicted that difference in insight across treatment will not remain after one year of treatment. A 2 (service use) x 3 (time) repeated measure ANOVA with follow-up analyses using the LSD procedure ($p = .05$) was completed for each measure. Analyses utilizing 2 (service use) x 3 (assessment time) repeated measure ANOVAs and 4 (APP Severity Level) x 3 (assessment time) repeated measure ANOVAs were attempted for each SAIQ subscale. However, based on when this assessment was administered at CTP, no group based on these distinctions had more than 10 people and one group had as few as zero. Therefore, no repeated measures analyses were undertaken based on SAIQ subscales and only analyses of the Insight Scale and its subscales will be discussed below.

There is an interaction between assessment time and service use on Insight Scale total score ($F(2, 114) = 3.26, p = .04, Mse = 6.82$). Insight total score remain the same over treatment for those individuals who used mental health services in adolescence, whereas for those individuals who did not use services prior to age 18 Insight Total scores are highest at admission and six months (which are equivalent to each other) and decrease after one year of treatment (LSD minimum mean difference = 1.36). Partial support for hypotheses is found, as those who did not use mental health services prior to age 18 have higher Insight Total scores at admission and six months and scores become equivalent for both groups after one year of treatment. However, contrary to hypotheses, those that had contact with mental health services before age 18 do not receive higher Insight Total scores at admission than those who did not use services prior to age 18. There is no main effect for Insight Total scores over time ($F(2, 114) = 2.37, p = .10$).

There is a significant main effect for service use ($F(1, 57) = 5.47, p = .02$). Individuals who did not use mental health services in adolescence have higher overall Insight Total scores across treatment than individuals who did use mental health services in adolescence; however, this pattern is only descriptive for both groups at admission and six months (LSD minimum mean difference = .96). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 100 and 104.




Figure 3.26 Service Main Effect and Interaction between Service Usage Groups on Birchwood Insight Total Scores Across Treatment

There is an interaction between assessment time and service use on Insight Relabel scale score ($F(2, 114) = 3.98, p = .02, Mse = .97$). Insight Relabel scale scores are highest at admission and six months (which are equivalent to each other) and decrease at one year for individuals without mental health service usage in adolescence, whereas for those individuals with mental health service usage in adolescence Insight Relabel scale scores remain the same over treatment (LSD minimum mean difference = .51). Contrary to hypotheses, no differences are found between individuals with and without

mental health service usage in adolescence on Insight Relabel scale scores at admission, and six months. As hypothesized, no differences are found between groups after twelve months. There is no main effect for Insight Relabel scale scores over time ($F(2, 114) = 2.40, p = .10$) or for service use ($F(1, 57) = .52, p = .48$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 100 and 101.




Figure 3.27 Interaction between Service Usage Groups on Birchwood Relabel Symptoms Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to Insight Awareness scale scores ($F(2, 114) = 2.52, p = .09, Mse = 1.25$). There is no main effect for Insight Awareness scale scores over time ($F(2, 114) = 1.37, p = .26$). Contrary to hypotheses, individuals with mental health service usage in adolescence do not have higher IS Awareness scores at admission than those without mental health service usage in adolescence. There is a main effect for service use ($F(1, 57) = 8.42, p = .01$). Individuals without mental health service usage in adolescence have higher overall

Insight Awareness scale scores across treatment than individuals with mental health service usage in adolescence, however this pattern is only descriptive for both groups at admission and six months (LSD minimum mean difference = .42). When planned comparisons were completed, Insight Awareness scores were lower for individuals with service usage in adolescence ($M = 1.93$) than for individuals who did not use services in adolescence ($M = 2.41$), $t(125) = 2.16$, $p = .02$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 100 and 102.




Figure 3.28 Service Usage Group Main Effect on Birchwood Relabel Symptoms Scores Across Treatment

There is not an interaction of assessment time and service use as they relate to Insight Need for Treatment scale scores ($F(2, 114) = 1.96$, $p = .15$, $Mse = 1.12$). There is a main effect for overall Insight Need for Treatment scale scores over time ($F(2, 114) = 3.22$, $p = .04$). Overall Insight Need for Treatment scale scores remain the same from admission to six months and from six months to one year of treatment, however overall scores after one year of treatment are lower than at admission (LSD minimum mean

difference = .39). However, this pattern is descriptive for neither group and is therefore misleading. There is no main effect for service use ($F(1, 57) = 2.77, p = .10$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 100 and 103.




Figure 3.29 Time Main Effect for Service Usage Groups on Birchwood Need For Treatment Scores Across Treatment

In addition, 4 (APP Severity Level) x 3 (assessment time) repeated measure ANOVAs were conducted for each insight measure. There is an interaction between assessment time and APP severity level on Insight Scale total score ($F(6, 110) = 3.38, p = .004, Mse = 6.31$). Insight total score remain the same over treatment for those individuals with high APP, whereas for those individuals without APP Insight Total scores remain the same from admission to six months and decrease after one year of treatment to scores consistent with those at admission (LSD minimum mean difference = 1.85). Insight total scores for individuals with low APP decrease from admission to six

and twelve months (LSD minimum mean difference = 1.85). Insight scores for individuals with medium APP increase from admission to six and twelve months (which are equivalent to each other). At admission, those with medium APP have lower Insight total scores than those without APP or low APP. After six months, individuals without APP have the highest Insight totals (all other groups have scores equivalent to each other). As hypothesized, after one year of psychiatric rehabilitation all groups have equivalent Insight total scores. There is no main effect for Insight Total scores over time ($F(2, 110) = 1.98, p = .14$). There is not a significant main effect for APP severity level ($F(3, 55) = .43, p = .73$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 105 and 109.




Figure 3.30 Interaction between APP Severity Levels on Birchwood Insight Total Scores Across Treatment

There is an interaction between assessment time and APP severity level on Insight Relabel scale score ($F(6, 110) = 2.90, p = .01, Mse = .93$). Insight Relabel scale scores

are highest at admission and six months (which are equivalent to each other) and decrease at one year for individuals without APP, whereas for those individuals with high APP Insight Relabel scale scores remain the same over treatment (LSD minimum mean difference = .71). Insight Relabel scale scores for individuals with low APP are highest at admission, contrary to hypotheses, and decrease from six to twelve months (which are equivalent to each other). At admission, individuals in the low APP group have higher scores than those in the medium APP group. As hypothesized, no differences are found between individuals without APP and various severities of APP on Insight Relabel scale scores at six and twelve months. There is no main effect for Insight Relabel scale scores over time ($F(2, 110) = .91, p = .41$) or for APP severity level ($F(3, 55) = .43, p = .73$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 105 and 106.




Figure 3.31 Interaction between APP Severity Levels on Birchwood Relabel Symptoms Scale Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to Insight Awareness scale scores ($F(6, 110) = 2.48, p = .03, Mse = 1.20$). Insight Awareness scores remain the same across treatment for those without APP and individuals in the medium and high APP group. Insight Awareness scores for individuals with low APP are higher at admission than at six to twelve months (which are equivalent to each other) (LSD minimum mean difference = .81). At admission, contrary to hypotheses, those with low APP have higher Insight Awareness scores than those with medium or high APP. At six months, those without APP have higher Insight Awareness scores than individuals with low or high APP. As hypothesized, by one year of treatment all groups have equivalent Insight Awareness scores. There is no main effect for Insight Awareness scale scores over time ($F(2, 110) = 2.16, p = .12$). There is a main effect for APP severity level ($F(3, 55) = 2.89, p = .04$). Individuals without APP have higher overall Insight Awareness scale scores across treatment than individuals with each severity of APP, whereas individuals with low or high APP (which are equivalent to each other) have higher overall Insight Awareness scores than individuals with medium APP. However this pattern is not descriptive for any assessment time and is therefore misleading (LSD minimum mean difference = .61). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 105 and 107.




Figure 3.32 APP Severity Level Main Effect and Interaction between APP Severity Levels on Birchwood Awareness Scale Scores Across Treatment

There is not an interaction of assessment time and APP severity level as they relate to Insight Need for Treatment scale scores ($F(6, 110) = 1.79, p = .11, Mse = 1.10$). There is a main effect for overall Insight Need for Treatment scale scores over time ($F(2, 110) = 3.14, p = .04$). Overall Insight Need for Treatment scale scores are highest at admission and decrease at six months and from six months to one year of treatment (LSD minimum mean difference = .39). However, this pattern is not completely descriptive for any group APP at all assessment times and is therefore misleading. There is no main effect for APP severity level ($F(3, 55) = .91, p = .44$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 105 and 109.




Figure 3.33 Time Main Effect for APP Severity Levels on Birchwood Need For Treatment Scale Scores Across Treatment

In summary, results of analyses on insight measures at admission and over the course of treatment suggest that hypothesized improvements in insight did not occur over the course of treatment for the all of the CTP population. Contrary to the hypothesis those adults who used mental health services in adolescence do not endorse differing Insight Totals or ability to relabel symptoms scores over the course of treatment. Furthermore, individuals without mental health service usage in adolescence endorse lower Insight Total scores and their ability to relabel symptoms as part of their illness decreases over the course of treatment. Furthermore, in general the CTP participants endorse lower Insight Need For Treatment scores across treatment. While differences do not exist after one year of treatment between those with and without service usage in adolescence on insight measures, this suggests neither groups' insight changed more than the others after participating in psychiatric rehabilitation.

When results of analyses on insight measures at admission and over the course of treatment utilize APP severity levels, results suggest that contrary to the hypothesis only individuals with low APP endorse increases in Insight Totals over the course of treatment. Insight Totals for individuals with no or low APP decrease over treatment while Insight Total scores remain the same across treatment for those with high APP. As hypothesized, ability to relabel symptoms increases over treatment for those with medium APP but remains the same for those with high APP. In contrast, individuals with no or low APP experience decreases in Insight Relabel scores across treatment. Insight Awareness scores remain the same across treatment for those with no, medium, or high APP while it decreases for those with low APP. Furthermore, insight into need for treatment decreases across treatment for individuals with no, low, or high APP but remains stable across treatment for those with medium APP. However, as hypothesized, there are no differences between those with no, low, medium, and high APP on Insight measures after one year of psychiatric rehabilitation.

Behavioral Functioning Across Treatment

It is hypothesized that as a result of psychiatric rehabilitation, there will be an increase in behavioral functioning for all CTP participants over the course of psychiatric rehabilitation. Although differences across treatment are hypothesized, it is predicted that differences will not remain after one year. A 2 (service use) x 3 (assessment time) repeated measure ANOVA was completed for each measure. There is not an interaction between assessment time and service use as they relate to NOSIE Daily Schedule Competence scores, $F(2, 188) = .57, p = .57, Mse = 14.53$. As hypothesized, the 2

(service) x 3 (time) repeated measures ANOVA reveals a significant main effect for time on the NOSIE Daily Schedule Competence scale scores, $F(2, 188) = 26.86, p < .001$ indicating that, overall, improvements in NOSIE Daily Schedule Competence occur with treatment for both individuals with and without mental health service usage in adolescence (LSD minimum mean difference = 1.10). There is no main effect for service use, $F(1, 94) = 2.54, p = .11$ (Tables 110 and 111). Planned comparisons indicate after six months of treatment NOSIE Daily Schedule Competence scores were lower for individuals with service usage in adolescence ($M = 32.65$) than for individuals who did not use services in adolescence ($M=34.83$), $t(115) = 1.84, p = .03$.




Figure 3.34 Time Main Effect for Service Usage Groups on NOSIE Daily Schedule Competence Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Social Interest scores, $F(2, 188) = 1.30, p = .28, Mse = 14.73$. As hypothesized, repeated measures ANOVA reveal a significant main effect for time on the NOSIE Social Interest scale scores, $F(2, 188) = 39.51, p < .001$ indicating that, overall,

improvements in NOSIE Social Interest scores occur with treatment for both individuals with and without mental health service usage in adolescence (LSD minimum mean difference = 1.10). There is no main effect for service use, $F(1, 94) = .06, p = .80$ (Tables 110 and 111).




Figure 3.35 Time Main Effect for Service Usage Groups on NOSIE Social Interest Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Neatness scores, $F(2, 188) = .33, p = .72, Mse = 8.19$. As hypothesized, analyses reveal a significant main effect for time on the NOSIE Neatness scale scores, $F(2, 188) = 20.89, p < .001$ indicating that, overall, improvements in NOSIE Neatness scores occur with treatment for both individuals with and without mental health service usage in adolescence (LSD minimum mean difference = .82). There is no main effect for service use, $F(1, 94) = 1.20, p = .28$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 110 and 111.




Figure 3.36 Time Main Effect for Service Usage Groups on NOSIE Neatness Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Irritability scores, $F(2, 188) = .15$, $p = .86$, $Mse = 16.27$. Analyses reveal a significant main effect for time on the NOSIE Irritability scale scores, $F(2, 188) = 3.78$, $p = .03$ indicating partial support for hypotheses. NOSIE Irritability scores increase from admission to six months and, as hypothesized, scores decrease from six months to one year of treatment for individuals with and without mental health service usage during adolescence (LSD minimum mean difference = 1.15). There is no main effect for service use, $F(1, 94) = 1.38$, $p = .24$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 110 and 111.




Figure 3.37 Time Main Effect for Service Usage Groups on NOSIE Irritability Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Psychoticism scores, $F(2, 188) = .96, p = .38, Mse = 3.56$. Contrary to hypotheses, results reveal a significant main effect for time on the NOSIE Psychoticism scale scores, $F(2, 188) = 7.47, p = .001$ indicating that, overall, NOSIE Psychoticism scores increase from admission to six months and one year of treatment (LSD minimum mean difference = .54). There is no main effect for service use, $F(1, 94) = .87, p = .35$ (Tables 110, 111).




Figure 3.38 Time Main Effect for Service Usage Groups on NOSIE Psychoticism Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Motor Retardation scores, $F(2, 188) = .34, p = .72, Mse = 7.03$. As hypothesized, results reveal a significant main effect for time on the NOSIE Motor Retardation scale scores, $F(2, 188) = 15.93, p < .001$ indicating that, overall, NOSIE Motor Retardation scores are not significantly different from admission to six months, however scores decrease by one year of treatment (LSD minimum mean difference = .76). There is no main effect for service use, $F(1, 94) = 1.50, p = .22$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 110 and 111.




Figure 3.39 Time Main Effect for Service Usage Groups on NOSIE Motor Retardation Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Total Assets scores, $F(2, 188) = .21, p = .81, Mse = 206.89$. As hypothesized, results reveal a significant main effect for time on the NOSIE Total Assets scores, $F(2, 188) = 21.68, p < .001$ indicating that, overall, improvements in NOSIE Total Assets scores occur with treatment for both individuals with and without mental health service use in adolescence (LSD minimum mean difference = 4.11). There is no main effect for service use, $F(1, 94) = 1.97, p = .16$ (Tables 110 and 111).




Figure 3.40 Time Main Effect for Service Usage Groups on NOSIE Total Scores Across Treatment

Results were also analyzed using a 4 (APP Severity Level) x 3 (assessment time) repeated measure ANOVA for each NOSIE measure. Results indicate there is not an interaction between assessment time and APP severity level as they relate to NOSIE Daily Schedule Competence scores, $F(6, 184) = .85, p = .54, Mse = 14.54$. As hypothesized, the 4 (APP severity level) x 3 (time) repeated measures ANOVA reveals a significant main effect for time on the NOSIE Daily Schedule Competence scale scores, $F(2, 184) = 23.33, p < .001$ indicating that, overall, improvements in NOSIE Daily Schedule Competence occur with treatment (LSD minimum mean difference = 1.09) (Tables 112, 113). This pattern is descriptive for all groups except individuals with high APP, for which scores decrease from admission to six months but are highest at one year. There is no main effect for APP severity level, $F(3, 92) = 1.10, p = .35$.

NOSIE Daily Schedule Competence Scores Across Treatment




Figure 3.41 Time Main Effect for APP Severity Levels on NOSIE Daily Schedule Competence Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to NOSIE Social Interest scores, $F(6, 184) = 1.83, p = .10, Mse = 14.40$. As hypothesized, results reveal a significant main effect for time on the NOSIE Social Interest scale scores, $F(2, 184) = 36.58, p < .001$ indicating that, overall, improvements in NOSIE Social Interest scores occur with treatment (LSD minimum mean difference = 1.08). There is no main effect for APP severity level, $F(3, 92) = .73, p = .54$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 112 and 113.




Figure 3.42 Time Main Effect for APP Severity Levels on NOSIE Social Interest Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to NOSIE Neatness scores, $F(6, 184) = 3.14, p = .01, Mse = 7.62$. For individuals without APP, NOSIE neatness scores remain the same from admission to six months but improve from six months to one year, whereas for individuals with low or medium APP scores continue to improve throughout treatment (LSD minimum mean difference = 1.58). For individuals with high APP, scores decrease from admission to six months but improve from six months to one year of treatment. At admission, individuals with high APP have the highest NOSIE neatness scores followed by those without APP. At six months, individuals without APP have higher scores than those with low or high APP. As hypothesized, after one year of treatment all groups have equivalent NOSIE neatness scores. As hypothesized, results reveal a significant main effect for time on the NOSIE Neatness scale scores, $F(2, 184) = 19.85, p < .001$ indicating that, overall, scores remain the same from admission to six months but improvements in NOSIE Neatness scores

occur from six months to one year (LSD minimum mean difference = .79). However, this pattern is only descriptive for those without APP and is therefore misleading. There is no main effect for APP severity level, $F(3, 92) = .73, p = .54$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 112 and 113.




Figure 3.43 Time Main Effect and Interaction between APP Severity Levels on NOSIE Neatness Scores Across Treatment

There is not an interaction between assessment time and APP severity level as they relate to NOSIE Irritability scores, $F(6, 184) = .40, p = .88, Mse = 16.44$. As hypothesized, results reveal a significant main effect for time on the NOSIE Irritability scale scores, $F(2, 184) = 3.40, p = .04$ indicating that, overall, NOSIE Irritability scores increase from admission to six months and decrease from six months to one year of treatment (LSD minimum mean difference = 1.16) (Tables 112, 113). This pattern is descriptive for all groups except the high APP group, for which NOSIE Irritability scores

remain the same across treatment. There is no main effect for APP severity level, $F(3, 92) = 1.30, p = .28$.




Figure 3.44 Time Main Effect for APP Severity Levels on NOSIE Irritability Scores Across Treatment

There is not an interaction between assessment time and APP severity level as they relate to NOSIE Psychoticism scores, $F(6, 184) = 1.58, p = .16, Mse = 3.49$.

Analyses reveal a significant main effect for time on the NOSIE Psychoticism scale scores, $F(2, 184) = 8.07, p < .001$ indicating that, overall, NOSIE Psychoticism scores increase from admission to six months and one year of treatment (which are equivalent to each other) (LSD minimum mean difference = .53). However, this pattern is only descriptive for those with low APP (Tables 112, 113). There is no main effect for APP severity level, $F(3, 92) = .36, p = .78$.




Figure 3.45 Time Main Effect for APP Severity Levels on NOSIE Psychoticism Scores Across Treatment

There is not an interaction between assessment time and APP severity level as they relate to NOSIE Motor Retardation scores, $F(6, 184) = 2.03, p = .06, Mse = 6.76$. Results reveal a significant main effect for time on the NOSIE Motor Retardation scale scores, $F(2, 184) = 13.55, p < .001$ indicating that, overall, NOSIE Motor Retardation scores are not significantly different from admission to six months, however as hypothesized scores decrease across treatment (LSD minimum mean difference = .74). However, this pattern is only descriptive for individuals with high APP and is misleading. There is no main effect for APP severity level, $F(3, 92) = 1.63, p = .19$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 112 and 113. One tailed planned comparisons indicate contrary to hypotheses NOSIE Motor Retardation scores were higher for individuals with service usage in adolescence ($M = 5.35$) than for individuals who did not use services in adolescence ($M = 4.18$), $t(94) = -1.66, p = .05$.




Figure 3.46 Time Main Effect for APP Severity Levels on NOSIE Motor Retardation Scores Across Treatment

There is an interaction between assessment time and APP severity level as they relate to NOSIE Total Assets scores, $F(6, 184) = 2.20, p = .05, Mse = 197.68$. NOSIE Total Assets scores increase across treatment for individuals with low or medium APP (LSD minimum mean difference = 8.04). For individuals without APP, scores do not change from admission to six months and from six months to one year, at which time scores are greater than at admission (Table 112, 113). NOSIE Total Assets score for individuals with high APP do not change from admission to six months but increase from six months to one year (which is equivalent to scores at admission). At admission, NOSIE Total Assets scores are equivalent for those without APP and those with high APP. However, those without APP or with high APP have higher NOSIE Total Assets than individuals with low or medium APP. At six months, individuals without APP or medium APP have higher scores than those with low or high APP. As hypothesized,

after one year of treatment no differences between groups exist on NOSIE Total Assets. As hypothesized, the 4 (APP severity level) x 3 (time) repeated measures ANOVA reveal a significant main effect for time on the NOSIE Total Assets scores, $F(2, 184) = 22.23, p < .001$ indicating that, overall, improvements in NOSIE Total Assets scores occur with treatment (LSD minimum mean difference = 4.02). There is no main effect for APP severity level, $F(3, 92) = 1.48, p = .23$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 112 and 113.




Figure 3.47 Time Main Effect for APP Severity Levels on NOSIE Total Scores Across Treatment

Overall, these results suggest that improvements in behavioral functioning do occur across treatment on almost all NOSIE subscales and NOSIE Total Assets for those with and without mental health service usage. However, partial support for hypotheses is found for NOSIE Irritability and NOSIE Psychoticism. When service use is utilized in analyses, all participants' NOSIE Irritability scores contrary to hypotheses increase from

admission to six months but, as hypothesized, do decrease by 12 months of treatment. Furthermore, contrary to hypotheses NOSIE Psychoticism scores for all participants continued to increase over the course of treatment. When results are analyzed with regards to APP severity level, the above results are again found. However, NOSIE Motor Retardation scores improve for all APP severity levels. Finally as predicted, no differences exist on NOSIE measures amongst groups after one year of psychiatric rehabilitation.

Symptomatology Across Treatment

It is hypothesized that as a result of psychiatric rehabilitation, improvements will occur in symptomatology across treatment. While differences may exist amongst groups at admission or six months, it is predicted that individuals who used mental health services in adolescence demonstrate higher symptomatology after one year of treatment. However, when different levels of APP are used in analyses, it is predicted that as APP becomes more severe there will be more severe symptomatology. A 2 (service use) x 3 (assessment time) repeated measure ANOVA and a 4 (APP Severity Level) x 3 (assessment time) repeated measure ANOVA was completed for each measure.

There is not an interaction between assessment time and service use as they relate to BPRS Total scores, $F(2, 128) = .86, p = .43, Mse = 97.89$. There is no main effect for BPRS Total scores over time ($F(2, 128) = 1.23, p = .30$) or for service use ($F(1, 64) = .23, p = .64$) (Table 114, 116).

There is an interaction between assessment time and service use on BPRS Psychotic Disorganization Factor score, $F(2, 128) = 3.17, p = .05, Mse = 3.87$. BPRS

Psychotic Disorganization Factor scores remain stable over treatment except for those who used mental health services in adolescence whose score at one year of treatment are lower than at admission as hypothesized (LSD minimum mean difference = .96).

Individuals with mental health service usage in adolescence have higher BPRS Psychotic Disorganization Factor scores at admission than individuals without mental health service usage in adolescence. However, contrary to hypotheses this difference does not persist at 6 months to one year of treatment. There is no main effect for BPRS Psychotic Disorganization Factor scores over time ($F(2, 128) = 2.16, p = .32$) or for service use ($F(1, 64) = 1.20, p = .28$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 114 and 117.




Figure 3.48 Interaction Between Service Usage Groups on BPRS Psychotic Disorganization Factor Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to BPRS Hallucination/Delusions Factor scores, $F(2, 134) = .85, p = .43, Mse = 7.97$.

There is no main effect for BPRS Hallucinations/Delusions Factor scores over time ($F(2, 134) = .82, p = .44$) or for service use ($F(1, 67) = .47, p = .50$) (Tables 114, 118). Also, there is not an interaction between assessment time and service use as they relate to BPRS Paranoia Factor scores, $F(2, 134) = .05, p = .95, Mse = 6.88$. There is no main effect for BPRS Paranoia Factor scores over time ($F(2, 134) = .04, p = .96$) or for service use ($F(1, 67) = 1.75, p = .19$) (Tables 114, 119).

There is not an interaction between assessment time and service use as they relate to BPRS Emotional Blunting Factor scores, $F(2, 134) = .19, p = .83, Mse = 4.41$. There is no main effect for BPRS Emotional Blunting Factor scores over time ($F(2, 134) = 1.03, p = .36$) or for service use ($F(1, 67) = 1.20, p = .28$). However, planned comparisons indicate BPRS Emotional Blunting Factor scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 6.17$) than for individuals who did not use services in adolescence ($M = 5.22$), $t(119) = -1.85, p = .03$.

In addition, there is not an interaction between assessment time and service use as they relate to BPRS Anxiety/Depression Factor scores, $F(2, 134) = .27, p = .77, Mse = 6.82$. There is no main effect for BPRS Anxiety/Depression Factor scores over time ($F(2, 134) = .50, p = .61$) or for service use ($F(1, 67) = .64, p = .43$). However, planned comparisons indicate BPRS Anxiety/Depression Factor scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 9.20$) than for individuals who did not use services in adolescence ($M = 7.82$), $t(118) = -2.09, p = .02$.

Finally there is not an interaction between assessment time and service use as they relate to BPRS Agitation/Elation Factor scores, $F(2, 134) = 1.08, p = .34, Mse = 1.51$. There is no main effect for BPRS Agitation/Elation Factor scores over time ($F(2, 134) =$

.06, $p = .94$) or for service use ($F(1, 67) = .02, p = .90$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 114, 120 - 122.

Because isolated differences between groups on individual BPRS items are found at admission, those analyses are repeated here to determine whether there is a consistent pattern over time on any particular BPRS items. No interactions, assessment time main effects, or service use main effects are found for the following BPRS items: Somatic Concern, Anxiety, Depression, Guilt, Hostility, Elevated Mood, Grandiosity, Suspiciousness, Disorientation, Conceptual Disorganization, Blunted Affect, Emotional Withdrawal, Tension, Uncooperativeness, Excitement, Distractibility, Motor Hyperactivity, and Mannerisms and Posturing (all F s < 3.71 , all p s $> .06$).

However, planned comparisons indicate several differences between individuals who used services in adolescence and those without a history of adolescent service usage. Results indicate BPRS Depression item scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 2.28$) than for individuals who did not use services in adolescence ($M = 1.85$), $t(119) = -1.89, p = .03$. Results also indicate BPRS Guilt item scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 2.10$) than for individuals who did not use services in adolescence ($M = 1.55$), $t(118) = -2.49, p < .01$. Additionally, planned comparisons indicate BPRS Hostility item scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 2.81$) than for individuals who did not use services in adolescence ($M = 2.32$), $t(119) = -1.89, p = .04$. As previously stated, t-tests at admission found a difference between groups on BPRS Grandiosity item scores. This result is also found after one year of treatment, when again results indicate scores are

lower for individuals with service usage in adolescence ($M = 1.57$) than for individuals who did not use services in adolescence ($M = 2.26$), $t(77) = 1.71$, $p = .05$. Results indicate BPRS Blunted Affect item scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 2.77$) than for individuals who did not use services in adolescence ($M = 2.32$), $t(119) = -1.91$, $p = .03$. Analyses also indicate BPRS Tension item scores are lower after one year of treatment for individuals with service usage in adolescence ($M = 1.33$) than for individuals who did not use services in adolescence ($M = 1.76$), $t(77) = 2.07$, $p = .02$. Finally, results indicate BPRS Motor Hyperactivity item scores are lower after one year of treatment for individuals with service usage in adolescence ($M = 1.18$) than for individuals who did not use services in adolescence ($M = 1.51$), $t(77) = 1.72$, $p = .05$. The previously stated results obtained from planned comparisons indicate analyses utilizing ANOVAs over the course of treatment sometimes did not have enough power to detect significant differences amongst the groups.

There is not an interaction between assessment time and service use as they relate to BPRS Suicidality scores, $F(2, 136) = .14$, $p = .87$, $Mse = .64$. There is no main effect for BPRS Suicidality scores over time ($F(2, 136) = .55$, $p = .58$). As hypothesized, there is a main effect for service use ($F(1, 68) = 6.62$, $p = .01$), with individuals with mental health service usage in adolescence receiving higher overall BPRS Suicidality scores than those without mental health service usage in adolescence at all assessment times (Table 124). When planned comparisons were completed, results indicate BPRS Suicidality item scores are higher at six months of treatment for individuals with service usage in adolescence ($M = 1.73$) than for individuals who did not use services in adolescence ($M =$

1.16), $t(119) = -3.35$, $p < .001$. Additionally, after one year of treatment results indicate contrary to hypothesis BPRS Suicidality item scores remain higher for individuals with service usage in adolescence ($M = 2.28$) than for individuals who did not use services in adolescence ($M = 1.85$), $t(77) = -.87$, $p = .01$.




Figure 3.49 Service Usage Group Main Effect on BPRS Suicidality Item Scores Across Treatment

There is an interaction between assessment time and service use on BPRS Hallucinations item score, $F(2, 136) = 3.97$, $p = .02$, $Mse = 1.87$. BPRS Hallucination items scores remain stable over treatment for those without mental health service usage in adolescence, whereas BPRS Hallucination items scores, contrary to hypotheses, decrease over treatment for those with mental health service usage in adolescence (LSD minimum mean difference = .59). As hypothesized, individuals with mental health service usage in adolescence have higher BPRS Hallucination item scores at admission than individuals without mental health service usage in adolescence; however, contrary to hypotheses, this difference does not persist at 6 months to one year of treatment. There is no main effect

for BPRS Hallucination item scores over time ($F(2, 136) = .08, p = .93$) or for service use ($F(1, 68) = .61, p = .44$). Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 124.




Figure 3.50 Interaction Between Service Usage Groups on BPRS Hallucinations Item Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to BPRS Unusual Thought Content item scores, $F(2, 134) = .51, p = .60, Mse = 1.57$. There is a main effect for assessment time, $F(2, 134) = 4.14, p = .02$. Scores remain the same from admission to six months, and from six months to one year, however as hypothesized scores after one year of treatment generally decrease from scores at admission. However, this pattern is only descriptive for individuals who used mental health service use in adolescence. There is no main effect for service use, $F(1, 67) = .57, p = .45$ (Table 124).




Figure 3.51 Time Main Effect for Service Usage Groups on BPRS Unusual Thought Content Item Scores Across Treatment

There is an interaction between assessment time and service use on BPRS Bizarre Behavior item score, $F(2, 136) = 3.24, p = .04, Mse = 1.22$. BPRS Bizarre Behavior items scores remain stable over treatment for those with and without mental health service usage in adolescence; however BPRS Bizarre Behavior items scores at one year are less than scores at admission for those who used mental health services in adolescence (LSD minimum mean difference = .52). Individuals who used mental health services in adolescence have higher BPRS Bizarre Behavior item scores at admission than individuals without mental health service usage in adolescence; however, contrary to hypotheses, this difference does not persist at 6 months to one year of treatment. There is no main effect for BPRS Bizarre Behavior item scores over time ($F(2, 136) = 1.00, p = .37$) or for service use ($F(1, 68) = 1.99, p = .16$) (Table 124).




Figure 3.52 Interaction Between Service Usage Groups on BPRS Bizarre Behavior Item Scores Across Treatment

There is an interaction between assessment time and service use on BPRS Self Neglect item score, $F(2, 136) = 4.46, p = .01, Mse = .61$. As hypothesized, BPRS Self Neglect items scores remain stable from admission to six months and then decrease for individuals who used mental health services in adolescence. However, contrary to hypotheses, scores increase from admission to six and twelve months (which are equivalent to each other) for those without mental health service usage in adolescence (LSD minimum mean difference = .37). As hypothesized, individuals who used mental health services in adolescence have higher BPRS Self-Neglect item scores at admission than individuals without mental health service usage in adolescence; however, contrary to hypotheses, this difference does not persist at 6 months to one year of treatment. There is an overall main effect for BPRS Self Neglect item scores over time, $F(2, 136) = 5.68, p < .01$. Self Neglect items scores increase from admission to six months and then decrease

at one year of treatment (LSD minimum mean difference = .26). However, this pattern is not descriptive for those with or without mental health service usage in adolescence and is therefore misleading. There is no main effect for service use, $F(1, 68) = .07, p = .80$. Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 124.




Figure 3.53 Time Main Effect and Interaction Between Service Usage Groups on BPRS Self-Neglect Item Scores Across Treatment

There is not an interaction between assessment time and service use as they relate to BPRS Motor Retardation item scores, $F(2, 134) = .11, p = .90, Mse = .61$. There is a main effect for assessment time, $F(2, 134) = 4.65, p = .01$. Scores decrease from admission to six months and one year (which are equivalent to each other) for individuals with and without mental health service usage in adolescence (LSD minimum mean difference = .26). There is no main effect for service use, $F(1, 67) = .77, p = .38$ (Table 124).




Figure 3.54 Time Main Effect for Service Usage Groups on BPRS Motor Retardation Item Scores Across Treatment

When 4 (APP Severity Level) x 3 (assessment time) repeated measures ANOVAs were conducted using APP severity levels, no interactions, assessment time main effects, or APP severity level main effects are found for the following assessments: BPRS Total, BPRS Psychotic Disorganization Factor, BPRS Hallucinations/Delusions Factor, BPRS Paranoia Factor, BPRS Emotional Blunting Factor, BPRS Anxiety/Depression Factor, BPRS Agitation/Elation Factor, BPRS Somatic Concern item, BPRS Anxiety item, BPRS Depression item, BPRS Guilt item, BPRS Hostility item, BPRS Elevated Mood item, BPRS Grandiosity item, BPRS Suspiciousness item, BPRS Hallucinations item, BPRS Unusual Thought Content item, BPRS Conceptual Disorganization item, BPRS Blunted Affect item, BPRS Emotional Withdrawal item, BPRS Tension item, BPRS Uncooperativeness item, BPRS Excitement item, BPRS Distractibility item, BPRS Motor

Hyperactivity item, and BPRS Mannerisms and Posturing item (all $F_s < 2.51$, all $p_s > .06$).

Results of repeated measure ANOVAs and means and standard deviations can be found in Tables 115, 123, and 125. Again, there are several significant differences on BPRS items when analyses assess differences amongst APP severity levels. There is not an interaction between assessment time and APP severity level as they relate to BPRS Suicidality scores, $F(6, 132) = .51, p = .80, Mse = .64$. There is no main effect for BPRS Suicidality scores over time, $F(2, 132) = .89, p = .41$. There is a main effect for APP severity level, $F(3, 66) = 4.64, p = .01$, with individuals with each level of APP having equivalent overall BPRS Suicidality scores, however as hypothesized those with high APP have higher overall BPRS Suicidality scores than those with no or low APP. This pattern is descriptive for all assessment times (Table 125).




Figure 3.55 APP Severity Levels Main Effect on BPRS Suicidality Item Scores Across Treatment

There is an interaction between assessment time and APP severity level on BPRS Bizarre Behavior item score, $F(6, 132) = 2.19, p = .05, Mse = 1.20$. BPRS Bizarre Behavior items scores remain stable over treatment for those with no, low, and medium APP whereas for individuals with high APP Bizarre Behavior item scores at one year are, as hypothesized, less than scores at admission (LSD minimum mean difference = .73). Individuals without APP and with medium APP have lower BPRS Bizarre Behavior item scores at admission than those in the low and high APP groups (which are equivalent to each other). At six months, BPRS Bizarre Behavior item scores are equivalent for each group. After one year of treatment, contrary to hypotheses, Bizarre Behavior item scores are highest for individuals with low APP, with those with no, medium, and high APP receiving equivalent BPRS Bizarre Behavior scores. There is no main effect for BPRS Bizarre Behavior item scores over time ($F(2, 132) = .17, p = .84$) or for APP severity level ($F(3, 66) = 1.22, p = .31$). Means and standard deviations are in Tables 125.




Figure 3.56 Interaction between APP Severity Levels on BPRS Bizarre Behavior Item Scores Across Treatment

There is an interaction between assessment time and APP severity level on BPRS Self Neglect item score, $F(6, 132) = 2.56, p = .02, Mse = .60$. BPRS Self Neglect items scores remain stable from admission to six months and then, as hypothesized, these scores decrease for individuals with high APP. Contrary to hypotheses, scores increase from admission to six months and remain the same from six to twelve months for those without APP (LSD minimum mean difference = .52). BPRS Self Neglect item scores remain the same across treatment for individuals with low or medium APP. No differences are found between groups on BPRS Self-Neglect item scores at admission or at six months, however as hypothesized individuals with high APP have lower BPRS Self Neglect scores than all other groups after one year or treatment. There is not an overall main effect for BPRS Self Neglect item scores over time ($F(2, 132) = 1.43, p = .24$) or for APP severity level ($F(3, 66) = .50, p = .68$) (Table 125).




Figure 3.57 Interaction between APP Severity Levels on BPRS Self-Neglect Item Scores Across Treatment

There is not an interaction between assessment time and APP severity level as they relate to BPRS Disorientation item scores, $F(6, 130) = .58, p = .75, Mse = .41$. There is not a main effect for assessment time, $F(2, 130) = .31, p = .74$. There is an overall main effect for APP severity level, $F(3, 65) = 4.97, p < .01$. Overall, contrary to hypotheses individuals with low APP have higher BPRS Disorientation item scores than those with no, medium, or high APP at all assessment times (LSD minimum mean difference = .52) (Table 125).




Figure 3.58 APP Severity Levels Main Effect on BPRS Disorientation Item Scores Across Treatment

There is not an interaction between assessment time and APP severity level as they relate to BPRS Motor Retardation item scores, $F(6, 130) = .34, p = .92, Mse = .62$. There is a main effect for assessment time, $F(2, 130) = 3.12, p = .05$. Scores remain the same from admission to six months and from six months to one year. However, as hypothesized scores at one year are lower than scores at admission (LSD minimum mean

difference = .27). However, this pattern is only descriptive for individuals with high APP and is therefore misleading. There is no main effect for APP severity level, $F(3, 65) = 1.04, p = .38$.




Figure 3.59 Time Main Effect for APP Severity Levels on BPRS Motor Retardation Item Scores Across Treatment

Overall, results of analyses using the BPRS total, BPRS factor scores, and BPRS items suggest that differences amongst groups exist for several symptomatology measures as they relate to assessment time and service use. Contrary to hypotheses, symptomatology for individuals with and without mental health service usage in adolescence does not improve over the course of treatment as measured by BPRS Total Assets, BPRS Hallucination/Delusions factor, BPRS Paranoia factor, BPRS Emotional Blunting factor, BPRS Anxiety/Depression factor, and BPRS Agitation/Elation factor. Partial support for hypotheses is found with the BPRS Psychotic Disorganization factor, as improvements across treatment are only descriptive for those who used mental health

services in adolescence. As hypothesized, individuals with mental health service usage in adolescence demonstrate more symptomatology at admission on the BPRS Psychotic Disorganization factor, however no other differences between groups at admission exist for other BPRS factor scores. In addition, contrary to hypotheses, individuals with mental health service usage in adolescence do not demonstrate more symptomatology after one year of treatment on the BPRS Total Assets score and BPRS factor scores. When BPRS items are analyzed across time, partial support for hypotheses is found. For individuals who used mental health services in adolescence, scores on BPRS Hallucinations/Delusion item, BPRS Unusual Thought Content item, BPRS Bizarre Behavior item, and BPRS Self-Neglect item improve by one year of treatment. However, scores on these same measures remain the same across treatment for those without mental health service usage in adolescence. However, as hypothesized, BPRS Motor Retardation scores improve over the course of treatment for those with and without mental health service usage in adolescence. As hypothesized, individuals who did use mental health services in adolescence demonstrate more symptomatology at admission on the BPRS Suicidality item, BPRS Hallucination/Delusions item, BPRS Bizarre Behavior item, and BPRS Self-Neglect item. However, by one year of treatment the only difference that remains between groups is that individuals who used mental health services in adolescence receive higher suicidality scores than those without mental health service usage in adolescence.

In summary, when analyses were conducted between APP severity level and BPRS total, BPRS factor scores, and BPRS items only significant differences emerge on individual BPRS items. Again, partial support for hypotheses is found. Improvement in

symptomatology across treatment is found only for those with high APP on BPRS Self-Neglect item, BPRS Bizarre Behavior item, and BPRS Motor Retardation item. As hypothesized, differences exist between the groups at various assessment times. As hypothesized, individuals with high APP demonstrate more symptomatology at admission on the BPRS Bizarre behavior item, however individuals with high APP demonstrate equivalent scores to those with low APP at admission. As hypothesized, individuals with high APP exhibit more symptomatology after one year of treatment on BPRS Suicidality item and BPRS Self-Neglect item. However, contrary to hypotheses individuals with low APP receive the highest BPRS Bizarre Behavior item and BPRS Disorientation item scores after one year of treatment and therefore demonstrate more symptomatology on these measures compared to those with high APP.

Hypothesis 2

Outcome

A pattern of differential outcome was hypothesized for those who used services during adolescence versus individuals who did not use services and for each APP severity level. Specifically, it is predicted that that discharge disposition differs between the two groups in that individuals with severe adolescent psychiatric pathology are discharged to a more restrictive setting. In addition, it is predicted that there is a greater rate of rehospitalization for those who used mental health services in adolescence than those without mental health services in adolescence. Furthermore, when analyses are conducted between the different APP severity groups, it is predicted that those

individuals in the medium and highest APP severity groups have a greater rate of rehospitalization than those in the low to no APP severity groups.

Also, it is predicted that those in the highest APP severity group have the greatest rate of rehospitalization, as these individuals may represent a subgroup experiencing more severe and refractory symptoms that can be potential barriers to living in the community for extended periods.

Discharge Location

It was predicted that people who used services during adolescence are discharged to more restrictive levels of care than those who did not use services. Chi square analyses reveal there is no significant relationship between the level of restrictiveness of discharge setting and service use in adolescence, $X^2(3) = 1.85, p = 0.60$ (See Table 126).

It was also predicted that when APP severity level is examined, discharge disposition differs between the groups in that individuals with severe adolescent psychiatric pathology are discharged to a more restrictive setting than individuals in the no, low, or medium APP groups. Chi square analyses reveal there is no significant relationship between the level of restrictiveness of discharge setting and APP severity level, $X^2(9) = 5.86, p = 0.75$ (See Table 127). Contrary to the hypothesis, those individuals with severe adolescent psychiatric pathology are not discharged to more restrictive settings than individuals with less severe adolescent pathology.

Rehospitalization Rate

Following discharge, CTP participants spent an average of 94.86% of days during the first six months after discharge in the community rather than in the hospital, 95.20% of days within the first year, 93.22% of days within eighteen months, and 93.20% of days

within the first two years after discharge. Ninety-three of the 123 people, or 75.6%, with rehospitalization data available are not rehospitalized from the time of discharge through the time the outcome data was collected. There are no significant differences between those who did or did not use services during adolescence on the percentage of days in the first six months after CTP discharge spent in the community ($F(1,122) = .03, p = 0.86$), the percentage of days in the first year after CTP discharge spent in the community ($F(1,109) = .22, p = 0.64$), the percentage of days within eighteen months after CTP discharge spent in the community ($F(1, 88) = .30, p = 0.58$), or the percentage of days within two years after CTP discharge spent in the community ($F(1, 73) = .02, p = 0.90$).

Furthermore, when APP severity level is examined, there are no significant differences between APP severity level on the percentage of days in the first six months after CTP discharge spent in the community ($F(3,122) = .72, p = 0.54$), the percentage of days in the first year after CTP discharge spent in the community ($F(3,109) = 1.02, p = 0.39$), the percentage of days within eighteen months after CTP discharge spent in the community ($F(3, 88) = .67, p = 0.58$), or the percentage of days within two years after CTP discharge spent in the community ($F(3, 73) = .55, p = 0.65$).

As described in the methods section, the rehospitalization data is skewed and could not be transformed or winsorized into an acceptable skewness range. This is largely a function of the large proportion of people who were never rehospitalized during the follow-up period. In order to substantiate the above results since the data used in the above analyses is skewed, categorical variables of "rehospitalized" and "not rehospitalized" were created based on all or none cutoff levels. Chi square analysis reveals no significant relationship between whether or not someone used services in

adolescence and rehospitalization categorization, $\chi^2(1) = .62, p = .43$. Likewise, rehospitalization categorizations were made based on rehospitalization in six-month intervals following discharge. At six months post-discharge, there is not a significant relationship between service use and rehospitalization categorization, $\chi^2(1) = .62, p = .43$. At 12 months post-discharge, again no significant relationships between service use and rehospitalization categorization are found, $\chi^2(1) = .30, p = .59$. Again, at 18 months post-discharge, there remains no relationship between the two variables, $\chi^2(1) = .88, p = .35$. Finally, at 24 months post-discharge, there remains no relationship between the two variables, $\chi^2(1) = .05, p = .82$.

When APP severity level is examined, Chi square analysis again reveals no significant relationship between APP severity level and whether or not they are rehospitalized after discharge from CTP, $\chi^2(3) = 2.87, p = .41$. Likewise, rehospitalization categorizations for each APP severity level are made based on rehospitalization in six-month intervals following discharge. At six months post-discharge, there is not a significant relationship between the various APP severity levels and rehospitalization categorization, $\chi^2(3) = 3.32, p = .35$. At 12 months post-discharge, again no significant relationships between APP severity level and rehospitalization categorization are found, $\chi^2(3) = 4.03, p = .26$. Again, at 18 months post-discharge, there remains no relationship between the two variables, $\chi^2(3) = 4.21, p = .24$. Finally, at 24 months post-discharge, there remains no relationship between the two variables, $\chi^2(3) = 5.21, p = .16$.

In summary, results of analyses of outcome data with regard to service use indicate that at the time of discharge and during the follow-up period after discharge, no

differences between individuals who used services in adolescence and those who did not use services are evident. Moreover, no differences between APP severity groupings are noted. Contrary to all outcome hypotheses, there is no evidence that APP severity levels differ in terms of rehospitalization outcomes.

CHAPTER 5 6 DISCUSSION

The primary purpose of this study was to assess the relationship of adolescent psychiatric pathology with insight before, during, and after treatment. In addition, a secondary relationship between adolescent psychiatric pathology and functioning in a variety of domains before, during, and after treatment were assessed. The participants with adolescent psychiatric pathology were compared to those without to determine if any differences existed. Also, when adolescent psychiatric pathology was separated into three separate groups by the amount and type of services utilized during adolescence, these three different levels of adolescent psychiatric pathology were also be compared to those without. Univariate and multivariate analyses were conducted within and between the two groups and four groups with respect to overall functioning using multiple measures described below. To date, only one study on this topic is known to exist (Wynne, 2009). Because of the paucity of research in this area, this study was highly exploratory in nature, giving a first look at the clinical correlates across treatment between individuals with and without adolescent psychiatric pathology. Furthermore, these clinical correlates across treatment in adulthood were also studied between individuals with various severities of adolescent psychiatric pathology.

Overall, there was mixed support for the hypotheses of the study. In summary results suggest individuals with service use during adolescence were found to be younger at admission to CTP, have slightly lower levels of education, and have an earlier age of onset and more previous hospitalizations, as expected. The results are consistent with earlier work (Herron, 1962) on the process reactive distinction in schizophrenia. The process reactive distinction analyzed the developmental sequence of schizophrenia

spectrum disorders. Specifically, research in this area is concerned with individuals' premorbid patterns, the timeline preceding the onset of the disorder, the precipitant(s) that may engender the onset of the disorder, and symptom presentation during the course of the disorder. Results from this study indicate those who used services in adolescence, as specifically the medium and high APP groups closely resemble the traditional psychopathology "process" group from research in the 1950s-1980. Process type has been associated with an insidious onset, gradual emotional blunting, and withdrawing from daily activities. Research found individuals identified in the process group had poor performance in school, deterioration in functioning in a variety of domains including neurocognitive, hospitalization during adolescence, poor response to treatment, awareness of change in self, and a family history of mental illness. Individuals without mental health service usage in adolescence may represent the reactive type again from research in the 1950s-1980s. Reactive type is usually characterized by a sudden onset of the disorder after the person experiences a particular stressor, have an onset later in life, functioning within normal limits in most domains in life prior to onset, good academic performance in school, good response to treatment, and "no sensation of change" (Haas & Sweeney, 1992). While it is interesting to note analyses utilizing the service use prior to age 18 vs. no service use prior to age 18 found similar results to research using process-reactive distinctions, utilizing either of these promotes a dichotomous typology rather than analyzing adults with SMI on a continuum. Thus, further research utilizing APP severity levels may be able to further delineate differences in premorbid functioning, onset of illness, clinical correlates, and influence favorable treatment outcomes for each individual at all levels of the continuum.

A relationship between Axis I and Axis II diagnoses and service use was identified. Those diagnosed with Schizophrenia, Chronic/Undifferentiated Type, Schizoaffective Disorder, and *other* appear more likely to have used mental health services during adolescence than those diagnosed with Schizophrenia, Paranoid Type. Those diagnosed with Paranoid Personality Disorder appear less likely to have received services during adolescence than those diagnosed with Borderline Personality Disorder; whereas those diagnosed with Borderline Personality Disorder appear more likely to have received services during adolescence. Those who used services and did not use services were more likely to be diagnosed with *other* than to have a diagnosis of Borderline Personality Disorder. Those diagnosed with Borderline Personality Disorder appear more likely to have received services during adolescence than not whereas those without an Axis II diagnosis were more likely to have not used services during adolescence. The differences between groups corroborate previous findings and indicate that exploration of these differences in participant characteristics needs to continue to be explored in future studies. The finding that Axis I and Axis II diagnoses are associated with service use may be a sign those that receive mental health services during adolescence may be more likely to develop comorbid disorders that greatly influence their functioning in adulthood. This may also indicate that those individuals who did not receive services during adolescence were experiencing less severe symptomatology than those who received services and thus these individuals may not have been perceived as high risk or needing services during adolescence by mental health professionals. This finding may have important implications for screening tools used by mental health professionals serving adolescents transitioning into adulthood.

Contrary to all hypotheses, participants with service usage in adolescence do not significantly differ from those without at the time of admission, with respect to neurocognitive, social cognitive functioning, or insight. In addition, there do not appear to be substantial differences in neurocognitive functioning, social cognitive functioning, or insight at admission for the various APP severity levels. Overall, results of NOSIE analyses do not support hypotheses that those with service use in adolescence demonstrate poorer behavioral functioning upon admission than those without, nor were hypotheses supported that differences in behavioral functioning at admission existed between the different APP severity groups. Although a significant difference occurred on the NOSIE Motor Retardation subscale, results were opposite to hypothesized and the high APP severity group received lower scores than individuals in the low and medium APP severity groups.

It was anticipated that individuals with service usage in adolescence would have a higher level of symptomatology at admission than those without, and specifically those individuals with the most severe APP would demonstrate a higher level of symptomatology than those with less severe APP. Individuals with service usage in adolescence received higher suicidality and hopelessness scores at admission than those without. Indeed, as research has indicated, symptoms such as suicidality and depression may all influence or be manifestations or outcomes of various levels of insight (Amador et al., 1991; Amador et al., 1993; Amador et al., 1996; Caldwell & Gottesman, 1990; Lysaker et al., 2005; McGlashan et al., 1975). These results indicate future studies may utilize cluster analysis in order to study the unique differences amongst groups in

variables, as a consistent pattern of differences in performance was not found by the current study.

When analyses were conducted between APP severity levels two significant differences emerged. Again as hypothesized, those individuals with medium or high APP received higher suicidality scores on multiple measures than those without APP or in the low APP severity group. Individuals without service use in adolescence were rated as more uncooperative than individuals with various levels of APP severity, indicating that prior contact with services may actually aid in working with treatment providers upon rehospitalization.

Results of analyses between those who used mental health services prior to age 18 and those who did not use services and neurocognitive functioning over the course of treatment suggest that improvements in neurocognitive functioning are evident over the course of treatment on strategic planning and organized searching, for individuals without service use in adolescence on a measure of verbal memory, for individuals with service use in adolescence on a measure of visuoconstructional ability and nonverbal memory, for both groups in nonverbal memory, for individuals with service use in adolescence on a task assessing attention and information processing, for those with service use in adolescence on verbal fluency and word generating ability, and for both groups on a neurocognitive screening assessment measuring overall cognitive functioning.

Results of analyses between assessment time and APP severity level indicate improvements in neurocognitive functioning are over the course of treatment on a task of verbal memory for those without APP, for those with low or high APP on a task of immediate nonverbal memory and visuoconstructional ability, for those with none or high

APP on delayed nonverbal memory, and for those in the none, low, or high groups on a neurocognitive screener measuring overall cognitive functioning.

As hypothesized, when analyses were conducted between those without service usage before age 18 and those with service usage in adolescence, no differences existed between groups on most measures of neurocognitive functioning. Contrary to hypotheses, individuals with service usage in adolescence were better able to rapidly generate and organize verbal information after one year of treatment than those without service usage in adolescence. However, individuals without APP demonstrated poorer overall cognitive functioning after one year of treatment than those with APP.

Finally, as hypothesized, when analyses were conducted utilizing APP severity level, as hypothesized no differences existed between groups after one year of psychiatric rehabilitation on most neurocognitive measures. Contrary to hypotheses, individuals with high APP demonstrated better verbal memory performance at admission and six months, however as hypothesized this difference did not remain after one year of treatment. As hypothesized, those individuals with high APP demonstrated poorer overall cognitive functioning at admission. However, contrary to hypotheses differences remained between the groups at twelve months and those without APP or low APP performed better on a measure of overall cognitive functioning than those in the medium or high APP severity groups.

Improvements in sociocognitive functioning were also evident over the course of treatment on participant's ability and facility of comprehension and interpretation of social limitations and for those with service usage prior to age 18 on social support coping cognitions. Contrary to hypotheses, there was not a decrease in participant's

beliefs being effected by their beliefs that others are powerful, a component of external locus of control. Furthermore, contrary to hypotheses there was not an increase in internal locus of control, in individual's self-concept of their own competence or participant's self-efficacy (both components of internal locus of control) over the course of treatment. As hypothesized, overall external locus of control and the role of chance in determining one's fate, a subscale of external locus of control, did decrease over treatment. However, this only occurred for those with service usage in adolescence.

When social cognitive was assessed amongst the various APP severity groups results indicate improvements in sociocognitive functioning are inconsistent over the course of treatment on theory of mind abilities. As previously indicated, there was not a decrease in participant's beliefs in powerful others over treatment. Furthermore, contrary to hypotheses there was not an increase in internal locus of control or participant's self-concept or self-efficacy over the course of treatment amongst the various APP severity groups. As hypothesized, external locus of control decreased over treatment, however this only occurred for those in the low, medium, and high groups. For beliefs that chance controls one's fate, decreases over treatment only occurred for those with low or medium APP.

Contrary to the hypothesis those adults with service usage in adolescence did not endorse differing overall insight or ability to relabel symptoms scores over the course of treatment. However, as hypothesized, there were no differences between those with and without service usage in adolescence on any measure of insight after one year of psychiatric rehabilitation. Furthermore, individuals without service usage prior to age 18 endorsed lower overall insight total scores and their ability to relabel symptoms as part of

their illness decreased over the course of treatment. In general, the CTP participants endorsed lower insight into need for treatment scores across treatment, possibly indicating an increased desire to be discharged as length of stay increased.

When results of analyses on insight measures at admission and over the course of treatment utilize APP severity levels, results suggest that differences in insight across treatment existed between groups. However, contrary to hypotheses only individuals with low APP endorsed increases in total insight over the course of treatment. Total insight for individuals with no or low APP decreased over treatment while total insight scores remained the same across treatment for those with high APP. As hypothesized, ability to relabel symptoms increased over treatment for those with medium APP but remained the same for those with high APP. Individuals with no or low APP experienced decreased ability to relabel symptoms across treatment. Insight awareness scores remained the same across treatment for those with no, medium, or high APP while it decreased for those with low APP. Furthermore, insight into need for treatment again decreased across treatment for individuals with no, low, or high APP groups but remained stable across treatment for those with medium APP. However, as hypothesized, there were no differences between those with and without APP on insight measures after one year of psychiatric rehabilitation.

Overall, results suggest that improvements in behavioral functioning do occur across treatment in almost all domains measured within the current study for those with and without service usage in adolescence. However, partial support for hypotheses was found for irritability and psychoticism scores. When service use was utilized in analyses, all participants' irritability scores increased from admission to six months but decreased

by 12 months of treatment. Furthermore, contrary to hypotheses psychoticism scores for all participants continued to increase over the course of treatment. When results were analyzed with regards to APP severity level, the above results were again found. However, motor retardation improved for all APP severity levels. However, as predicted, no differences existed amongst groups after one year of psychiatric rehabilitation.

Results also suggest that differences amongst groups exist for several symptomatology measures as they relate to assessment time and service use. Contrary to hypotheses, symptomatology for individuals with and without service usage in adolescence did not improve over the course of treatment. Partial support for hypotheses was found with the BPRS Psychotic Disorganization factor, as improvements across treatment were only descriptive for those with service usage in adolescence. As hypothesized, individuals with service usage prior to age 18 demonstrated more symptomatology at admission on the BPRS Psychotic Disorganization factor, however no other differences between groups at admission existed for other BPRS factor scores. Also contrary to hypotheses, individuals with service usage in adolescence did not demonstrate more symptomatology after one year of treatment. When specific symptomatology items were analyzed across time, partial support was found. For individuals with mental health service usage in adolescence, items assessing hallucinations/delusions, unusual thought content, bizarre behavior, and self-neglect improved by one year of treatment. However, on scores on these same measures remained the same across treatment for those without service usage in adolescence. However, as hypothesized, BPRS Motor Retardation scores improved over the course of treatment for those with and without service usage in adolescence. As hypothesized,

individuals with service usage in adolescence demonstrated more suicidality, hallucinations/delusions, bizarre behavior, and self-neglect. However, by one year of treatment the only difference that remained between groups was that individuals with service usage in adolescence received higher suicidality scores than those without service usage in adolescence.

Results also suggest that differences amongst groups exist for several symptomatology measures as they relate to assessment time and APP severity level. Again, partial support for hypotheses were found. Improvement in symptomatology across treatment was found only for those with high APP on items measuring self-neglect, bizarre behavior, and motor retardation. As hypothesized, differences existed between the groups at various assessment times. As hypothesized, individuals with high APP demonstrated more symptomatology at admission on items assessing bizarre behavior, however individuals with high APP demonstrated equivalent scores to those with low APP at admission. As hypothesized, individuals with high APP exhibited more suicidality and self-neglect. However, contrary to hypotheses individuals with low APP received the highest bizarre behavior and disorientation item ratings after one year of treatment.

Differences in level of discharge restrictiveness and rehospitalization rates were predicted for those with and without service usage in adolescence and for the various APP severity levels. Contrary to the hypothesis, those individuals with severe adolescent psychiatric pathology were not discharged to more restrictive settings than individuals with less severe adolescent pathology. There were no significant differences between those who did or did not use services during adolescence on the percentage of days spent

in the community in the first six, twelve, eighteen, and twenty-four months after CTP discharge. Furthermore, when APP severity level is examined, there are no significant differences between APP severity level on the percentage of days in the first six, twelve, eighteen, and twenty-four months after CTP discharge spent in the community.

Again, contrary to hypotheses, analyses revealed no significant relationship between whether or not someone used services in adolescence and rehospitalization categorization. Likewise, rehospitalization categorizations were made based on rehospitalization in six-month intervals following discharge. At six, twelve, eighteen, and twenty-four months post-discharge, there was not a significant relationship between service use and rehospitalization categorization.

When APP severity level was examined, analyses again revealed no significant relationship between whether or not someone used services in adolescence and whether or not they were rehospitalized after discharge from CTP at all assessment time periods. Thus, contrary to all outcome hypotheses, there is no evidence that with or without APP differ in terms of rehospitalization outcomes or discharge location restrictiveness.

Overall, similar discharge patterns between those with and without APP were found with discharges from CTP. Treatment teams may be aware of the social history of each person and it is likely that failure at previous discharge locations impact future discharge planning. Of note, several measures of symptomatology and behavioral functioning (i.e. those with service usage in adolescence were rated higher on items of self-neglect and suicidality) were unable to maintain adequate self-care or an appropriate level of care needed for some discharge locations. Based on past rates of rehospitalization, treatment teams may be more likely to discharge to locations seen as

transitional, or intermediate levels of care. Perhaps even more likely, discharges may be dictated by the community providers themselves based on prior working with participants from the program. Therefore, participants from CTP may only be accepted by a limited range of providers leading to the non-significant discharge findings.

The second part of the outcome hypothesis pertained to rate of rehospitalization. Results of this study cannot be used to fully substantiate nor disprove the notion that people with APP have higher rates of rehospitalization and that those with high APP may represent a more population with more treatment refractory symptoms. A possible confound was the community agencies for which rehospitalization data was available. Records were not available if individuals resided or were hospitalized in another state, if they did not use the local CMHC, or if they were rehospitalized anywhere else besides the state hospital. This may have significantly skewed the rehospitalization data that was available, therefore more qualitative analyses or case studies regarding rehospitalization rate for those with and without APP may be warranted to further evaluate this hypothesis.

The inconsistent pattern of results and partial support of hypotheses in this study may be explained by the nature of the population from which the sample was drawn. Because of the severity and chronicity of psychiatric disorder within the CTP population, some of the non-significant differences between groups can be attributable to the overall high level of impairment present in the population at CTP, thus creating a "leveling effect" that may obscure some findings that would occur in other treatment environments. The CTP participants represent a particularly treatment-refractory population and differences in functioning between groups with various amount and types of service use during adolescence may be less apparent than they would be in a less severe or less

chronic psychiatric population with more variability. Discriminating between any groups within the CTP population may require not only the existence of differences, but substantial differences. That is not to say that there is not considerable heterogeneity within this group, but it may mean that fine, subtle differences between groups may be hard to detect, and these differences may or may not be meaningful. This conclusion is congruent with studies which have had difficulty detecting treatment effects between groups in the CTP setting (Spaulding, Reed, Sullivan, Richardson, & Weiler, 1999; Wynne, 2009). A similar study across treatment settings and/or populations may be warranted.

Limitations of the Present Study

The goal of identifying between a developmental conceptualization of insight based on contact with adolescent psychiatric pathology and various domains of functioning while hospitalized and outcomes in the community resulted in a myriad of findings. The lack of consistent patterns of differences between groups may be because of varied conceptualizations of the construct of insight, constitutes service use during adolescence, and the unknown quality of services received. One weakness of this study is that included measures of insight that do not reflect recent advances in multidimensional insight assessments. Furthermore, insight assessments may be influenced by practice effects, as participants are given the same self-report form over the course of treatment. Since this study utilized archival clinical data, only measures that are part of routine assessment at CTP were available. While poor cognitive functioning has been linked to insight (e.g., Amador et al., 1991; Frith, 1992; Lysaker et al., 1994;

Lysaker et al., 1995; McGlynn & Schacter, 1989; Prigatano & Schacter, 1991 as cited in Rusch and Corrigan, 2002) and this study found inconsistent results between groups on neurocognitive functioning. It may be that a more global measurement of insight reflecting current conceptualizations of insight, or a variable derived of various domains from assessments in other domains, would better capture the relationship between service use during adolescence and changes in insight and neurocognitive functioning across treatment.

The current archival database does not lend itself to pre-post analyses. Future studies may consider reformatting the archival database so as to have an admission data point and a discharge data point (or the assessment closest to discharge) as an approximation of pre- and post- rehabilitation functioning. Furthermore, the use of a flexible assessment battery with this clinical population meant there was an unusually large amount of missing data that prevented some use of appropriate statistical analyses. The amount of significant results was also close to that expected to occur by chance, thus lowering confidence in complete interpretation of results. For most analyses examining the between-group differences and within group difference there was enough power to detect differences. Effect size estimates were examined in addition to the statistical significance tests to determine there was enough power to detect all differences.

At any rate, this study explored the relationship between various domains of functioning during the course of psychiatric rehabilitation and service use, APP severity and insight such that future studies in this area can make more informed research hypotheses using stricter constraints in research design to circumvent problems related to possible confounds.

Future Directions

Areas of needed research have already been alluded to in the above discussion. Specifically, a replication or study similar to the one undertaken here, with more defined hypotheses and changes in assessment measures is needed to further clarify the characteristics of those with various amounts of service use during adolescence and the differences in several domains of functioning across treatment after the age of 18 and once discharged into the community. Likewise, a similar study in a broader population would allow for more generalizability of results. Finally, a study or a series of case studies examining different events and service use during adolescence, possibly a longitudinal study following subjects from adolescence to adulthood, is necessary to better understand why some people improve their insight across treatment and others do not given the different trajectories or trends in insight scores and other domains of functioning across treatment for the various severity levels of APP.

While the Birchwood Insight Scale is theoretically based on a multidimensional view of insight (David, 1990), this measure may not be indicative of current multidimensional conceptualizations of insight, or other conceptualizations such as the view that insight is the degree to which an individual agrees with their treatment provider, or insight as a coping mechanism. The current study sought to determine relationships between individual's insight and domains of functioning, however a consistent pattern could not be established. Therefore, using measures of insight to identify correlates to clinical functioning, or insight across treatment, is unresolved until a relationship between the psychological construct of insight and other biopsychosocial

measures is better understood. Studies identifying correlates of insight to various clinical variables have not been conducted within a population whose historical use of services during adolescence has been investigated. Or, if they have, the effect of contact with mental health providers earlier in life on insight into treatment after the onset of illness has not been considered in analyses. Such a study might simply include different biopsychosocial measures within a population such as the one in this study to determine if the measures make distinctions or can predict different APP severity level groups across treatment.

In general, more empirical, as opposed to theoretical investigations, of the concepts found within the prodromal research and service use during adulthood are needed in order to prevent or decrease the revolving door phenomenon commonly found in mental health settings. The current study continues to take essential steps towards identifying the relationship between the access to and use of mental health services by adolescents and the subsequent clinical functioning of those individuals with an onset of mental illness later after transitioning into adulthood.

References

- American Psychiatric Association (1994). *Diagnostic and statistical manual of mental disorders (4th ed.)*. Washington, DC: Author.
- Amador, X.F., Harkavy-Friedman, J., Kasapis, C., Yale, S.A., Flaum, M., & Gorman, J.M. (1996). Suicidal behavior in schizophrenia and its relationship to awareness of illness. *American Journal of Psychiatry, 153*, 1185-1188.
- Amador, X.F., & Strauss, D.H. (1993). Poor insight in schizophrenia. *Psychiatric Quarterly, 64*, 305-318.
- Amador, X.F., Strauss, D.H., Yale, S.A., & Gorman, J.M. (1991). Awareness of illness in schizophrenia. *Schizophrenia Bulletin, 17*, 113-132.
- Army Individual Test Battery. (1944). *Manual of directions and scoring*. Washington DC: War Department, Adjutant General's Office.
- Baechler, J. (1979). *Suicide*. New York: Basic Books.
- Bailey, S., Lachar, D., Rhoades, H.M., Diefenbach, G.J., Espadas, A., & Varner, R.V. (2004). Quantifying symptomatic change during acute psychiatric hospitalizations using new subscales for the anchored Brief Psychiatric Rating Scale. *Psychological Services, 1*, 68-82.
- Beck, A.T., & Steer, R.A. (1988). *Manual for the Beck Hopelessness Scale*. San Antonio, TX: Psychological Corporation.
- Beck, A.T., Steer, R.A., & Brown, G.K. (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Benton, A.L., & Hamsher, K. (1976). *Multilingual aphasia examination*. Iowa City: University of Iowa.

- Bleuler, E. (1950). *Dementia praecox or the group of schizophrenias*. Zinkin J. trans. New York: International Universities Press.
- Brekke, J., Kohrt, B., & Green, M. (2001). Neuropsychological functioning as a moderator of the relationship between psychosocial functioning and the subjective experience of self and life in schizophrenia. *Schizophrenia Bulletin*, 27, 697-708.
- Caldwell, C.B., & Gottsman (1990). Schizophrenics kill themselves too: A review of risk factors for suicide. *Schizophrenia Bulletin*, 16, 571-589.
- Carpenter, W.T., Jr. Strauss, J.S., & Bartko, J.J. (1973). Flexible system for the diagnosis of schizophrenia: Report from the WHO international pilot study of schizophrenia. *Science*, 182:1275-1278.
- Chen, Y.R., Swann, A.C., & Burt, D.B. (1996). Stability of diagnosis in schizophrenia. *American Journal of Psychiatry*, 153, 682-686.
- Cohen, J. (1988). Set correlation and contingency tables. *Applied Psychological Measurement*, 12(4), 425-434.
- Corcoran, R., & Frith, C. D. (2003). Autobiographical memory and theory of mind: Evidence of a relationship in schizophrenia. *Psychological Medicine*, 33, 897-905.
- Corcoran, R., Mercer, G. & Frith, C. D. (1995). Schizophrenia, symptomatology and unique and combined contributions of multiple child abuse types and abuse severity to adult trauma symptomatology social inference: Investigating theory of mind in people with schizophrenia. *Schizophrenia Research*, 17, 5-13.

- Cuestra, M.J., Peralta, V., Caro, F., & de Leon, J. (1995). Is poor insight in psychotic disorders associated with poor performance on the Wisconsin Card Sorting Test? *American Journal of Psychiatry*, *152*, 1380-1382.
- Cull, J. G., & Gill, W. S. (1988). *Suicide Probability Scale Manual*. Los Angeles: Western Psychological Services.
- David, A.S. (1990). Insight and psychosis. *British Journal of Psychiatry*, *156*, 798-808.
- David, A., & Kemp, R. (1997). Five perspectives on the phenomenon of insight in psychosis. *Psychiatry Annual*, *27*, 791-797.
- David, A., van Os, J., Jones, P., Harvey, I., Foerster, A., & Fahy, T. (1995). Insight and psychotic illness. Cross-sectional and longitudinal associations. *British Journal of Psychiatry*, *167*, 621-628.
- De Bruyn, E.E.J., Haassen, P.P., Van Poortinge, Y.H., & Lutje Spelberg, H.C. (1986). *Wechsler Intelligence Scale for Children - Revised, Dutch Translation*. Lisse, The Netherlands: Swets and Zeitlinger.
- Evren, C., & Evren, B. (2004). Characteristics of schizophrenic patients with a history of suicide attempt. *International Journal of Psychiatry in Clinical Practice*, *8*, 227-234.
- Flashman, L.A., McAllister, T.W., Andreasen, N.C., & Saykin, A.J. (2000). Smaller brain size associated with unawareness of illness in patients with schizophrenia. *American Journal of Psychiatry*, *157*, 1167-1169.
- Freud, S. (1940). Outline for the future of psychoanalysis. Strachey A, Strachey J, Trans. *Collected papers*, Vol II. London: Hogarth Press.

- Freudenreich, O., Deckersbach, T., & Goff, D.C. (2004). Insight into current symptoms of schizophrenia. Association with frontal cortical function and affect. *Acta Psychiatrica Scandinavica*, 110, 14-20.
- Friedman, H. (1982). Simplified determinations of statistical power, magnitude of effect and research sample sizes. *Educational & Psychological Measurement*, 42(2), 521-526.
- Frith, C.D. (1992). *The cognitive neuropsychology of schizophrenia*. Sussex: Lawrence Erlbaum Associates.
- Ghaemi, S. N., & Pope, H. G. (1994) Lack of insight in psychotic and affective disorders: a review of empirical studies. *Harvard Review of Psychiatry*, 2, 22-33.
- Gold, J.M., Queern, C., Iannone, V.N., & Buchanan, R.W. (1999). Repeatable Battery for the Assessment of Neuropsychological Status as a screening test in schizophrenia, I: Sensitivity, reliability, and validity. *American Journal of Psychiatry*, 156, 1944-1950.
- Goldman, R.S.; Axelrod, B.N.; and Tompkins, L.M. (1992). Effect of instructional cues on schizophrenic patients' performance on the Wisconsin Card Sorting Test. *American Journal of Psychiatry*, 149(12): 1718-1722.
- Greig, T.C., Bryson, G.J., & Bell, M.D. (2004). Theory of mind performance in schizophrenia: Diagnostic, symptoms and neuropsychological correlates. *Journal of Nervous Mental Disorders*, 192, 12-18.
- Haas, G.L., & Sweeney, J.A. (1992). Premorbid and onset of first-episode schizophrenia. *Schizophrenia Bulletin*, 18(3), 373-386.

- Hoaglin, D.C., Mosteller, F., & Tukey, J.W., (1983). *Understanding robust and exploratory data analysis*. New York: Wiley.
- Heaton, R.K., Chelune, G.J., Talley, J.L., Kay, G.G., & Curtis, G. (1993). *Wisconsin Card Sorting Test Manual: Revised and expanded*. Odessa, FL. Psychological Assessment Resources.
- Helgeland, M.I., & Torgersen, S. (2005). Stability and prediction of schizophrenia from adolescence to adulthood. *European Child and Adolescent Psychiatry, 14 (2)*, 83-94.
- Herron, W.G. (1962). The process-reactive distinction of schizophrenia. *Psychological Bulletin, 59(4)*, 329-343.
- Hobart, M.P., Goldberg, R., Bartko, J.J., & Gold, J.M. (1999). Repeatable Battery for the Assessment of Neuropsychological Status as a screening test in schizophrenia, II: Convergent/discriminant validity and diagnostic group comparisons. *American Journal of Psychiatry, 156*, 1951-1957.
- Honigfeld, G., Roderic, D., & Klett, J. (1966). NOSIE-30: A treatment-sensitive ward behavior scale. *Psychological Review, 19*, 180-182.
- Jaspers, K. (1963). *General psychopathology* (7th edition, originally published 1959, translation by J. Hoenig & M.W. Hamilton). Manchester: Manchester University Press.
- Jonikas, J.A., Laris, A., & Cook, J.A. (2003). The passage to adulthood: Psychiatric rehabilitation service and transition related needs of young adult women with Emotional and psychiatric disorders. *Psychiatric Rehabilitation Journal, 27(2)*, 114-121.

- Kemp, R., & David, A. (1996). Psychological predictors of insight and compliance in psychotic patients. *British Journal of psychiatry*, *169*, 444-450.
- Krampen, G. (1991). Fragebogen zu kompetenz-und kontrolluebersengungen (FKK), *Handanweisung (1 ed.)*. Goettingen: Hogrefe Verlag fuer Psychologie.
- Kupper, Z., & Hoffmann, H. (2000). Course patterns of psychosocial functioning in schizophrenia patients attending a vocational rehabilitation program. *Schizophrenia Bulletin*, *26(3)*, 681-698.
- Kupper, Z., & Hoffmann, H. (2002). Facilitators of psychosocial recovery from schizophrenia. *International Review of Psychiatry*, *14(4)*, 1369-1627.
- Lecomte, Y., & Mercier, C. (2005). The stress process perspective and adaptation of people with schizophrenia: An exploratory study. *Social Psychiatry and Psychiatric Epidemiology*, *40(2)*, 139-148.
- Lewis, L. (2004). Mourning, insight, and reduction of suicide risk in schizophrenia. *Bulletin of the Menninger Clinic*, *68*, 231-244.
- Link, B.G., & Phelan, J.C. (1999). Labeling and stigma. In C.S. Aneshensel & J.C. Phelan (Eds.), *Handbook of sociology of mental health* (pp. 481-494). New York: Kluwer Academic/Plenum Publishers.
- Lukoff, D., Nuechterlein, K. H., & Ventura, J. (1986). Manual for the expanded Brief Psychiatric Rating Scale (BPRS). *Schizophrenia Bulletin*, *12*, 594-602.
- Lyall, D., Hawley, C., & Scott, K. (2004). Nursesø Observation Scale for Inpatient Evaluation: Reliability update. *Journal of Advanced Nursing*, *46 (4)*, 390-394.
- Lysaker, P., Bell, M., Milstein, R., Bryson, G., & Beam-Goulet, J. (1994). Insight and psychosocial treatment compliance in schizophrenia. *Psychiatry*, *57*, 307-315.

- Lysaker, P.H., Bell, M.D., Bryson, G., & Kaplan, E. (1998). Neurocognitive function and insight in schizophrenia: Support for an association with impairments in executive function but not with impairments in global function. *Acta Psychiatrica Scandinavica*, *97*, 297-301.
- Lysaker, P.H., Carcione, A., Dimaggio, G., Johannesen, J.K., Nicolo, G., Procacci, M., & Semerari, A. (2005). Metacognition amidst narratives of self and illness in schizophrenia: Associations with neurocognition, symptoms, insight, and quality of life. *Acta Psychiatrica Scandinavica*, *112*, 64-71.
- Markova, J.S., & Berrios, G.E. (1995). Insight in clinical psychiatry revisited. *Comprehensive Psychiatry*, *36*, 367-376.
- Marks, K.A., Fastenau, P.S., Lysaker, P.H. & Bond, G.R., (2000). Self-appraisal of illness questionnaire: Relationship to researcher-rated insight and neuropsychological function in schizophrenia. *Schizophrenia Research*, *3(27)*, 203-211.
- McEvoy, J.P., Aland Jr., J., Wilson, W.H., Guy, W. & Hawkins, L. (1981). Measuring chronic schizophrenic attitudes toward their illness and treatment, *Hosp. Community Psychiatry*, *12*, 856-858.
- McEvoy, J.M., Freter, S., Everett, F., Geller, J.L., Appelbaum, P., Apperson, L.J. & Roth, L. (1989). Insight and clinical outcome of schizophrenic patients. *Journal of Nervous Mental Disorders*, *177(1)*, 48-51.
- McEvoy, J.P., Hartman, M., Gottlieb, D., Godwin, S., Apperson, L.J., & Wilson, W. (1996). Common sense, insight, and neuropsychological test performance in schizophrenia patients. *Schizophrenia Bulletin*, *22*, 635-641.

- McEvoy, J., Scheifler, P., & Frances, A. (Eds.). (1999). Treatment of schizophrenia. *Journal of Clinical Psychiatry*, 60 (Suppl.11), 4-80.
- McGlashan, T.H., Levy, S.T., & Carpenter, W.T., Jr. (1975). Integration and sealing over. Clinically distinct recovery styles from schizophrenia. *Archives of General Psychiatry*, 32, 1269-1272.
- McGlashan, T. H., Wadeson, H. S., Carpenter, W. T., *et al* (1977). Art and recovery style from psychosis. *Journal of Nervous and Mental Disease*, 164, 182-190.
- McGlashan, T.H., & Carpenter Jr., W.T. (1981). Does attitude toward psychosis relate to outcome? *American Journal of Psychiatry*, 138, 797-801.
- McGlynn, S.M., & Schacter, D.L. (1989). Unawareness of deficits in neuropsychological syndromes. *Journal of Clinical and Experimental Neuropsychology*, 11, 143-205.
- McGorry, P.D., & McConville, S.B. (1999). Insight in psychosis: An elusive target. *Comprehensive Psychiatry*, 40, 131-142.
- Meltzer, H.Y., Anad, R., & Alphs, L. (2000). Reducing suicide risk in schizophrenia: Focus on the role of clozapine. *CNS Drugs*, 14, 355-365.
- Meyers, J. E., & Meyers, K. R. (1995). *The Meyers scoring system for the Rey Complex Figure Test and recognition trial: Professional manual*. Odessa: Psychological Assessment Resources.
- Mindt, M., & Spaulding, W. (2002). The coping strategies task: Assessment of coping in schizophrenia. *Psychiatric Rehabilitation Skills*, 6(3), 428-453.
- Mohammed, S., Fleming, S., Penn, D.L., & Spaulding, W. (1999). Insight in schizophrenia: Its relationship to measures of executive functions. *Journal of Nervous & mental Disease*, 187, 525-531.

- Peer, J., & Spaulding, W. (2007). Heterogeneity in recovery of psychosocial functioning during psychiatric rehabilitation: An exploratory study using latent growth mixture modeling. *Schizophrenia Research, 93*(1), 186-193.
- Penn, D., Mueser, K., Spaulding, W., Hope, D., & Reed, D. (1995). Information processing and social competence in chronic schizophrenia. *Schizophrenia Bulletin, 21*(2), 269-281.
- Pompili, M., Ruberto, A., Kotzalidis, G.D., Girardi, P., & Tatarelli, R. (2004). Suicide and awareness of illness in schizophrenia: An overview. *Bulletin of the Menninger Clinic, 68*, 297-318.
- Prigatano, G.P., & Schacter, D.L. (1991). *Awareness of deficit after brain injury: Clinical and theoretical issues*. New York: Oxford University Press.
- Randolph, C. (1998). *Repeatable battery for the assessment of neuropsychological status*. San Antonio: Harcourt Brace & Company.
- Rey, A. (1941). L'examen psychologique dans les cas de depression et d'encephalopathie traumatique. *Archives de Psychologie, 28*, 286-340.
- Rusch, N., & Corrigan, P.W. (2002). Motivational interviewing to improve insight and treatment adherence in schizophrenia. *Psychiatric Rehabilitation Journal, 26*, 23-32.
- Schmidt M. (1996). *Rey Auditory Verbal Learning Test*. Los Angeles: Western Psychological Services.
- Spaulding, W.D., Fleming, S.K., Reed, D., Sullivan, M.E., Storzbach, D., & Lam, M. (1999a). Cognitive functioning in schizophrenia: Implications for psychiatric rehabilitation. *Schizophrenia Bulletin, 25*, 275-289.

- Spaulding, W., Garbin, C.P., & Dras, S.R. (1989). Cognitive abnormalities in schizophrenic patients and schizotypal college students. *Journal of Nervous and Mental Disease, 177*(12), 717-728.
- Spaulding, W., Johnson, D. & Coursey, R. (2001). Treatment and rehabilitation of schizophrenia. In M. Sammons & N. Schmidt (Eds.) *Combined treatments for mental disorders: A guide to psychological and pharmacological interventions*. Washington, DC: American Psychological Association.
- Spaulding, W.D., Reed, D., Sullivan, M., Richardson, C., & Weiler, M. (1999b). Effects of cognitive treatment in psychiatric rehabilitation. *Schizophrenia Bulletin, 25*, 657-676.
- Spaulding, W.D., Sullivan, M., & Poland, J. (2003). *Treatment and rehabilitation of severe mental illness: A comprehensive approach*. New York: Guildford Publications.
- Smith, T.E., Hull, J.W., Goodman, M., Hedayat-Harris, A., Wilson, D.F., Israel, L.M., & Munich, R.L. (1999). The relative influences of symptoms, insight and neurocognition on social adjustment in schizophrenia and schizoaffective disorder. *Journal of Nervous and Mental Disorders, 187*, 102-108.
- Stern, R.A., & White, T. (2001). *Neuropsychological Assessment Battery: Administration, Scoring and Interpretation Manual*. Psychological Assessment Resources, Inc.
- Takai, A., Uematsu, M., Ueki, H., Sone, K., & Kaiya, H. (1992). Insight and its related factors in chronic schizophrenic patients: A preliminary study. *European Journal of Psychiatry, 6*, 159-170.

- Van Meijel, B., Van der Gaag, M., Kahn, R.S., & Grypdonck, M. (2002). The practice of early recognition and intervention to prevent psychotic relapse in patients with schizophrenia: An exploratory study. Part 2. *Journal of Psychiatric and Mental Health Nursing, 9*, 357-363.
- Van der Does, A.J.W., Linszen, D.H., Dingemans, P.M., Nugter, M.A., & Scholte, W.F. (1993). A dimensional and categorical approach to the symptomatology of recent-onset schizophrenia. *The Journal of Nervous and Mental Disease, 181*, 744-749.
- Vander Stoep, A., Davis, M., & Collins, D. (2000). Transition: A time of developmental and institutional clashes. In H.B. Clark & M. Davis (Eds.), *Transition to adulthood: A resource for assisting young people with emotional or behavioral difficulties* (pp. 3-28). Baltimore, MD: Paul H. Brookes Publishing Co.
- Wynne, A. (2009). Developmental analysis of insight of individuals with serious mental illness in an inpatient psychiatric setting (Unpublished masters equivalency research project, University of Nebraska-Lincoln, 2009).

APPENDIX:

CTP Information Gathering Sheet pg. 1

Name: _____

ID number (please write down if you find they have more than 1 ID #): _____

Date(s) the file was looked at: _____

Clinical Assistant who checked file: _____

Please write down any information about mental health services the patient received before the age of 18 (therapy, medicine, services from other mental health practitioners such as social works, etc; being in-patient hospitalized, whether they were in adolescent services here at the regional center or in another state, etc):

Prior Therapy before the age of 18:

Medicine before the age of 18:

Services form other mental health providers (social workers, OT, psychiatrists, etc.):

CTP Information Gathering Sheet pg. 2

Were they in-patient hospitalized before the age of 18?

Any other information that is useful for determining how many and what type of mental health services they received prior to the age of 18?

Is more information needed to determine the amount or type? If so, what do you feel is needed?

Was there any vague information that did not allow you to accurately rate the amount of previous mental health services? If so describe the information.

Table 1
Demographic and Clinical Characteristics of Participants

<u>Demographic and Clinical Variables</u>	<u>N</u>	<u>Mean (SD) or Percentage</u>
Age (range: 18 to 71)	308	37.91 (12.62)
Education, # of years in school (range: 6 to 20 years)	283	12.23 (2.03)
Length of stay at CTP in days (range:57-2545)	137	637.28 (452.97)
Age of onset (range: 4 to 63)	248	19.92 (8.60)
Number of hospitalizations in lifetime (range: 0 to 105)	298	9.69 (9.77)
Gender (n=317)		
Male	203	64.0%
Female	114	36.0%
Race/Ethnicity (n=288)		
Caucasian	238	82.6%
African American	30	10.4%
Hispanic	6	2.1%
Native American	4	1.4%
Asian American	2	0.7%
Other	8	2.8%
Marital Status (n=272)		
Single	171	62.9%
Married	21	7.7%
Divorced	69	25.4%
Widowed	4	1.5%
Separated	7	2.6%
Legal Status (n=280)		
Mental Health Board Commitment (MHB)	220	78.6%
Voluntary per guardian (VpG)	34	12.1%
Not responsible by reason of insanity (NRRI)	22	7.9%
Voluntary (V)	4	1.4%
Primary Axis I Diagnosis (n=282)		
Schizophrenia, Paranoid Type	77	27.3%
Schizophrenia, Chronic/Undifferentiated Type	66	23.4%
Schizoaffective	78	27.7%
Bipolar	35	12.4%
Psychotic Disorder NOS	4	1.4%
Dementia / Organic Brain Disease	4	1.4%
Impulse Control Disorder	6	2.1%
Other	12	4.3%
Axis II Diagnoses (n=284)		
None	105	37.0%
Borderline	20	7.0%
Paranoid	46	16.2%
Antisocial	13	4.6%
Schizoid	4	1.4%
Borderline Intellectual Functioning / Mild Mental Retardation	19	6.7%
NOS	30	10.6%
Other	47	16.5%

Table 2
Demographic and Clinical Characteristics of Participants according to Service Use During Adolescence

<u>Demographic and Clinical Variables</u>	No Service Use		Service Use	
	<u>N</u>	<u>Mean (SD) or Percentage</u>	<u>N</u>	<u>Mean (SD) or Percentage</u>
Age (range: 19 to 71)	145	42.55 (12.35)	163	33.77 (11.40)
Education, # of years in school (range: 6 to 20 years)	134	12.84 (1.98)	149	11.69 (1.92)
Length of stay at CTP in days (range:11-2545)	72	581.39 (411.34)	65	699.20 (490.81)
Age of onset (range: 4 to 63)	118	24.20 (9.07)	130	16.03 (5.92)
Number of hospitalizations in lifetime (range: 0 to 105)	139	8.42 (7.47)	159	10.81 (11.31)
Gender (n=317)				
Male	94	63.1%	109	64.1%
Female	53	35.6%	61	35.9%
Race/Ethnicity (n=288)				
Caucasian	111	74.5%	127	82.5%
African American	15	10.1%	15	9.7%
Hispanic	1	0.7%	5	3.2%
Native American	2	1.3	2	1.3%
Asian American	1	0.7%	1	0.6%
Other	4	2.7%	4	2.6%
Marital Status (n=272)				
Single	75	58.6%	96	66.7%
Married	10	7.8%	11	7.6%
Divorced	34	26.6%	35	24.3%
Widowed	3	2.3%	1	0.7%
Separated	6	4.7%	1	0.7%
Legal Status (n=280)				
Mental Health Board Commitment (MHB)	103	69.1%	117	77.5%
Voluntary per guardian (VpG)	12	8.1%	22	14.6%
Not responsible by reason of insanity (NRRI)	11	7.4%	11	7.3%
Voluntary (V)	3	2.0%	1	0.7%
Primary Axis I Diagnosis (n=282)				
Schizophrenia, Paranoid Type	51	38.3%	26	17.4%
Schizophrenia, Chronic/Undifferentiated Type	23	17.3%	43	28.9%
Schizoaffective	35	26.3%	43	28.9%
Bipolar	15	11.3%	20	13.4%
Psychotic Disorder NOS	1	0.8%	3	2.0%
Dementia / Organic Brain Disease	2	1.5%	2	1.3%
Impulse Control Disorder	3	2.3%	3	2.0%
Other	3	2.3%	9	6.1%
Axis II Diagnoses (n=284)				
None	58	43.3%	47	31.3%
Borderline	4	3.0%	16	10.7%
Paranoid	20	14.9%	26	17.3%
Antisocial	4	3.0%	9	6.0%
Schizoid	1	0.7%	3	2.0%
Borderline Intellectual Functioning to Mild Mental Retardation	6	4.4%	13	8.1%
NOS	14	10.4%	16	10.7%
Other	27	20.1%	20	13.3%

Table 3
Demographic and Clinical Characteristics of Participants according to APP
Severity Level

Demographic and Clinical Variables	No APP		Low APP		Med APP		High APP	
	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD) or
		or Percentage		or Percentage		or Percentage		Percentage
Age (range: 19 to 71)	145	42.55 (12.35)	23	40.04 (10.68)	34	33.04 (10.23)	106	32.64 (11.53)
Education, # of years in school (range: 6 to 20 years)	134	12.84 (1.98)	22	12.23 (1.77)	31	11.87 (1.82)	96	11.51 (1.97)
Length of stay at CTP in days (range:11-2545)	72	581.39 (411.34)	19	781.68 (633.89)	17	698.18 (382.86)	29	645.76 (448.31)
Age of onset (range: 4 to 63)	118	24.20 (9.07)	20	20.20 (6.41)	31	17.68 (6.91)	79	14.33 (4.60)
Number of hospitalizations in lifetime (range: 0 to 105)	139	8.42 (7.47)	22	11.00 (7.57)	33	7.45 (6.32)	104	11.83 (12.95)
Gender (n=317)								
Male	94	63.1%	13	56.5%	27	77.1%	69	61.6%
Female	53	35.6%	10	43.5%	8	22.9%	43	38.4%
Race/Ethnicity (n=288)								
Caucasian	111	74.5%	18	81.8%	31	93.9%	78	78.8%
African American	15	10.1%	1	4.5%	1	3.0%	13	13.1%
Hispanic	1	0.7%	1	4.5%	1	3.0%	3	3.0%
Native American	2	1.3	-	-	-	-	2	2.0%
Asian American	1	0.7%	1	4.5%	-	-	-	-
Other	4	2.7%	1	4.5%	-	-	3	3.0%
Marital Status (n=272)								
Single	75	58.6%	11	55.0%	19	61.3%	66	71.0%
Married	10	7.8%	1	5.0%	5	16.1%	5	5.4%
Divorced	34	26.6%	8	40.03%	6	19.4%	21	22.6%
Widowed	3	2.3%	-	-	1	3.2%	-	-
Separated	6	4.7%	-	-	-	-	1	1.1%
Legal Status (n=280)								
Mental Health Board Commitment (MHB)	103	69.1%	15	68.2%	22	66.7%	80	83.3%
Voluntary per guardian (VpG)	12	8.1%	6	27.3%	9	27.3%	7	7.3%
Not responsible by reason of insanity (NRR1)	11	7.4%	1	4.5%	2	6.1%	8	8.3%
Voluntary (V)	3	2.0%	-	-	-	-	1	1.0%
Primary Axis I Diagnosis (n=282)								
Schizophrenia, Paranoid Type	51	38.3%	3	13.6%	8	25.0%	15	15.8%
Schizophrenia,Chronic/ Undifferentiated Type	23	17.3%	9	40.9%	12	37.5%	22	23.2%
Schizoaffective	35	26.3%	10	45.5%	2	6.3%	31	32.6%
Bipolar	15	11.3%	-	-	3	9.4%	17	17.9%
Psychotic Disorder NOS	1	0.8%	-	-	1	3.1%	2	2.1%
Dementia/Organic Brain Disease	2	1.5%	-	-	-	-	2	2.1%
Impulse Control Disorder	3	2.3%	-	-	2	5.9%	1	1.1%
Other	3	2.3%	-	-	4	12.5%	5	5.3%
Axis II Diagnoses (n=284)								
None	58	43.3%	7	31.8%	8	25.8%	32	33.0%
Borderline	4	3.0%	-	-	2	6.5%	14	14.4%
Paranoid	20	14.9%	6	27.3	9	29.0%	11	11.3%
Antisocial	4	3.0%	2	9.1%	1	3.2%	6	6.2%
Schizoid	1	0.7%	-	-	-	-	3	3.1%
Borderline Intellectual Functioning ó Mild Mental Retardation	6	4.4%	1	4.5%	3	9.7%	9	9.3%
NOS	14	10.4%	2	9.1%	2	6.5%	12	12.4%
Other	27	20.1%	4	18.2%	6	19.3%	10	10.3%

Table 4
Relationship Between Axis I Diagnosis and Service Use (N=282)

<u>Axis I Diagnosis</u>	<u>No Service</u>	<u>Service Use</u>	<u>Total</u>
Schizophrenia, Paranoid Type	51	26	77
Schizophrenia, Chronic/Undifferentiated Type	23	43	66
Schizoaffective	35	43	78
Other ²	24	37	61
Total	133	149	282

² Other includes diagnoses such as, but not limited to, the following: Bipolar Disorder, Dementia, Psychotic Disorder NOS, Pervasive Developmental Disorder, Impulse Control Disorder, and Aspergerø.

Table 5
Relationship Between Axis II Diagnosis and Service Use (N=284)

<u>Axis II Diagnosis</u>	<u>No Service</u>	<u>Service Use</u>	<u>Total</u>
Borderline	4	16	20
Paranoid	20	26	46
Antisocial	4	9	13
Other ⁴	34	36	70
NOS	14	16	30
No Axis II diagnosis (None)	58	47	105
Total	134	150	284

⁴ Other includes diagnoses such as, but not limited to, the following: Histrionic Personality Disorder, Schizoid, and Borderline Intellectual Functioning.

Table 6
Relationship Between Race and Service Use (N=288)

<u>Race</u>	<u>No Service</u>	<u>Service Use</u>	<u>Total</u>
Caucasian	111	127	238
Non-white	23	27	50
Total	134	154	288

Table 7
Relationship Between Marital Status and Service Use (N=272)

<u>Marital Status</u>	<u>No Service</u>	<u>Service Use</u>	<u>Total</u>
Single	75	96	171
Married	10	11	21
Divorced	34	35	69
Widowed	3	1	4
Separated	6	1	7
Total	128	144	272

Table 8
*Mean Scores and Standard Deviations for Demographic and Clinical Variables as a
 Function of Service Use*

	<u>No Service</u>		<u>Service Use</u>	
	<u>M</u>	<u>SD</u>	<u>M</u>	<u>SD</u>
Age*	42.55	12.35	33.77	11.38
Length of Stay	581.39	411.34	699.20	490.81
Years of Education*	12.84	1.98	11.69	1.92
Age of Onset*	24.20	9.07	16.03	5.92
Number of Previous Hospitalizations*	8.42	7.47	10.81	11.31

* P<.05

Table 9
Bivariate Correlations for Neurocognitive Variables

	1a	2	3	4	5	6	7	8	9	10	11	12	13	14
1 - RAVLT Trial 5	-													
2 - NAB Total	.74	-												
3 - COGLAB Total Correct	.10	-a	-											
4 - Card Sort Random Errors	-.41**	-.15	-.41**	-										
5 - Card Sort Perseverative Errors	-.36**	-.05	-.43**	.58**	-									
6 - Card Sort Consolidation Index	.23	-1.0**	-.10	-.13	.12	-								
7 - RCFT Copy	1.0**	.24	a	-.30*	-.03	.53	-							
8 - RCFT Immediate Memory	1.0**	.55**	a	-.53**	-.42**	-.26	.38**	-						
9 - RCFT Delayed Memory	1.0**	.66**	a	-.48**	-.40**	-.16	.41**	.95**	-					
10 - RCFT Recognition	-1.0**	.36	a	-.29*	-.17	-.45	-.05	.48**	.50**	-				
11 - Trails A	.10	-.54**	a	.22	.29*	-.25	.01	-.23*	-.30*	-.06	-			
12 - Trails B	-.45	-.61	a	.33**	.32*	.20	-.14	-.32**	-.38**	-.03	.54**	-		
13 - COWAT/FAS	-.56	.56**	a	-.38**	-.23	-.17	.35**	.37**	.34**	.08	-.32**	-.27**	-	
14 - RBANS Total	.30	.53*	a	-.31*	-.22	.06	.48**	.52**	.55**	.25*	-.15	-.29**	.41**	-

^a Note: The RAVLT was phased out of regular use at CTP when the Rey Complex Figure Test and RBANS battery were added. As can be seen here, there were no participants with assessments from both time periods from which to compute correlations.

** Correlation is significant at the 0.01 level (1-tailed).

* Correlation is significant at the 0.05 level (1-tailed).

Table 10
Service Use by Neurocognitive Variables at Admission
One-Way Analyses of Variance (ANOVAs)

Neurocognitive Variables	Univariate			
	M(SD)	F	df	p
RAVLT				
No Service	8.07 (3.73)	.18	1, 55	.68
Service Use	8.47 (3.30)			
NAB Total				
No Service	82.10 (17.06)	.53	1, 76	.47
Service Use	79.08 (18.00)			
WCST Correct				
No Service	24.63 (2.01)	.09	1, 60	.77
Service Use	24.75 (1.02)			
WCST Perseverative Errors				
No Service	22.13 (13.01)	.44	1, 106	.51
Service Use	20.51 (12.26)			
WCST Random Errors				
No Service	23.58 (16.65)	.05	1, 106	.83
Service Use	22.85 (18.27)			
WCST Consolidation Index				
No Service	3.46 (4.64)	.61	1, 60	.44
Service Use	2.57 (4.31)			
RCFT Copy				
No Service	29.03 (7.21)	.04	1, 71	.84
Service Use	29.33 (6.00)			
RCFT Immediate Memory				
No Service	12.76 (7.09)	.63	1, 71	.43
Service Use	14.32 (9.60)			
RCFT Delayed Memory				
No Service	12.68 (6.68)	.34	1, 71	.56
Service Use	13.80 (9.55)			
RCFT Recognition				
No Service	20.54 (6.16)	3.33	1, 71	.07
Service Use	18.17 (4.87)			
Trails A				
No Service	41.71 (14.59)	.11	1, 114	.74
Service Use	42.99 (25.71)			
Trails B				
No Service	105.14 (47.61)	2.65	1, 114	.11
Service Use	125.24 (81.07)			
COWAT/FAS				
No Service	30.97 (11.19)	.66	1, 113	.42
Service Use	29.32 (10.57)			
RBANS Total				
No Service	75.50 (16.89)	3.13	1, 92	.08
Service Use	69.74 (14.06)			

Table 11
APP Severity Level by Neurocognitive Variables at Admission
One-Way Analyses of Variance (ANOVAs)

<u>Neurocognitive Variables</u>	<u>Univariate</u>			
	<u>M(SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
RAVLT				
No APP	8.07 (3.73)			
Low APP	8.71 (4.07)	.45	3, 53	.72
Med APP	9.57 (2.76)			
High APP	7.88 (3.22)			
NAB Total				
No APP	82.10 (17.06)			
Low APP	70.00 (13.11)	.73	3, 74	.54
Med APP	87.25 (18.66)			
High APP	78.95 (18.25)			
WCST Correct				
No APP	24.63 (2.01)			
Low APP	25.00 (0.00)	.29	3, 58	.83
Med APP	25.00 (0.00)			
High APP	24.50 (1.41)			
WCST Perseverative Errors				
No APP	22.13 (13.01)			
Low APP	21.50 (11.02)	.49	3, 104	.69
Med APP	17.25 (14.87)			
High APP	21.45 (11.68)			
WCST Random Errors				
No APP	23.58 (16.65)			
Low APP	23.88 (7.79)	1.66	3, 104	.18
Med APP	13.25 (11.23)			
High APP	26.09 (20.98)			
WCST Consolidation Index				
No APP	3.46 (4.64)			
Low APP	1.31 (1.23)	1.04	3, 58	.38
Med APP	4.50 (6.58)			
High APP	2.03 (3.38)			
RCFT Copy				
No APP	29.03 (7.21)			
Low APP	30.07 (3.40)	.15	3, 69	.93
Med APP	28.06 (8.38)			
High APP	29.57 (5.84)			
RCFT Immediate Memory				
No APP	12.76 (7.09)			
Low APP	14.21 (4.72)	1.15	3, 69	.34
Med APP	18.56 (16.09)			
High APP	12.74 (7.36)			
RCFT Delayed Memory				
No APP	12.68 (6.68)			
Low APP	12.14 (4.72)			
Med APP	17.56 (16.10)	.85	3, 69	.47
High APP	12.91 (7.39)			
RCFT Recognition				
No APP	20.54 (6.16)	2.18	3, 69	.10
No APP	18.14 (3.08)			

Low APP	21.13 (8.11)			
Med APP	17.05 (3.28)			
High APP				
Trails A				
No APP	41.71 (14.59)			
Low APP	52.25 (25.14)	1.72	3, 112	.17
Med APP	35.08 (15.67)			
High APP	45.71 (30.05)			
Trails B				
No APP	105.14 (47.61)			
Low APP	122.75 (80.44)	1.36	3, 112	.26
Med APP	111.21 (70.12)			
High APP	134.48 (88.39)			
COWAT/FAS				
No APP	30.97 (11.19)			
Low APP	28.11 (9.61)	.36	3, 111	.78
Med APP	30.67 (12.93)			
High APP	28.87 (9.50)			
RBANS Total				
No APP	75.50 (16.89)			
Low APP	75.00 (15.04)	1.53	3, 90	.21
Med APP	70.91 (14.80)			
High APP	67.35 (13.43)			

Table 12
Bivariate Correlations for Socialcognitive Variables

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
1 - Hinting Task	-																	
2 - FKK Internal	-.03	-																
3 - FKK Self Concept	.06	.28*	-															
4 - FKK Self Efficacy	.01	.86**	.73**	-														
5 - FKK Powerful Others	.09	.07	-.11	-.01	-													
6 - FKK Chance	-.01	.09	-.45**	-.21	.66**	-												
7 - FKK Externality	.05	.08	-.33**	-.12	.92**	.91**	-											
8 - IPSAQ Internal Positive	-.00	.25*	.25*	.31*	.09	.02	.06	-										
9 - IPSAQ Personal Positive	.26*	-.15	-.12	-.17	.15	.22	.20	-.13	-									
10 - IPSAQ Situational Positive	.19	.22	.19	.26*	.08	-.05*	.02	-.39**	-.30*	-								
11 - IPSAQ Internal Negative	-.03	.10	-.18	-.03	.05	.24*	.16	.48**	.09	-.22	-							
12 - IPSAQ Personal Negative	.30*	.004	.30*	.17	-.09	-.17	-.14	.16	.41**	-.09	-.37**	-						
13 - IPSAQ Situational Negative	.02	.31*	.23	.34**	.32*	.07	.21	-.03	-.22	.56**	-.35**	-.24	-					
14 - CST Social Support Seeking	.03	-.03	-.11	-.08	-.18	-.12	-.17	.06	.19	-.10	.18	-.12	-.01	-				
15 - CST Self-Controlling	.04	.05	.41**	.26*	.01	-.22	-.12	.26*	-.01	-.12	-.07	.13	.15	-.11	-			
16 - CST Escape Avoidance	.16	.08	.28*	.20	-.39**	-.47**	-.47**	-.21	.05	.14	-.16	.10	.03	.10	.37**	-		
17 - CST Planful Problem Solving	-.01	.09	.40**	.28*	-.13	-.45**	-.31*	.02	.12	.15	-.23	.16	.31*	.34*	.43**	.30*	-	

** Correlation is significant at the 0.01 level (1-tailed).

* Correlation is significant at the 0.05 level (1-tailed).

Table 13
Service Use by Theory of Mind, Social Cognition at Admission One-Way Analyses of Variance (ANOVAs)

	M(SD)	F	df	p
Theory of Mind Measure				
Hinting Task				
No Service	15.12 (3.32)	1.06	1, 92	.31
Service Use	14.36 (3.81)			
Socialcognition Measures				
FKK Internal				
No Service	34.41 (7.18)	.04	1, 43	.84
Service Use	34.00 (5.33)			
FKK Self Concept				
No Service	33.56 (4.71)	.17	1, 43	.68
Service Use	32.94 (5.16)			
FKK Self Efficacy				
No Service	67.96 (9.95)	.13	1, 43	.72
Service Use	66.94 (7.80)			
FKK Powerful Others				
No Service	25.96 (8.80)	.24	1, 43	.63
Service Use	24.78 (6.51)			
FKK Chance				
No Service	23.48 (7.20)	.55	1, 43	.46
Service Use	25.17 (7.80)			
FKK Externality				
No Service	49.44 (15.53)	.01	1, 43	.91
Service Use	49.94 (11.60)			
IPSAQ Internal Positive				
No Service	8.19 (3.50)	.09	1, 43	.77
Service Use	7.89 (2.78)			
IPSAQ Personal Positive				
No Service	3.52 (2.39)	.00	1, 43	.96
Service Use	3.56 (1.92)			
IPSAQ Situational Positive				
No Service	3.26 (2.35)	.44	1, 43	.51
Service Use	3.78 (2.90)			
IPSAQ Internal Negative				
No Service	5.89 (3.77)	.00	1, 43	1.0
Service Use	5.89 (2.93)			
IPSAQ Personal Negative				
No Service	4.89 (3.48)	.00	1, 43	1.0
Service Use	4.89 (2.83)			
IPSAQ Situational Negative				
No Service	4.07 (2.80)	.23	1, 43	.64
Service Use	4.50 (3.13)			
CST Social Support Seeking				
No Service	35.58 (9.99)	.11	1, 42	.74
Service Use	36.61 (10.03)			
CST Self-Controlling				
No Service	52.65 (6.97)	2.58	1, 42	.12
Service Use	48.44 (10.44)			
CST Escape Avoidance				
No Service	37.12 (6.73)	.84	1, 42	.36
Service Use	35.44 (4.53)			
CST Planful Problem Solving				
No Service	42.08 (8.85)	.00	1, 42	.95
Service Use	41.89 (10.02)			

Table 14
APP Level by Theory of Mind and Social Cognitive Functioning at Admission
One-Way Analyses of Variance (ANOVAs)

	M(SD)	F	df	p
Theory of Mind Measure				
Hinting Task				
No APP	15.12 (3.32)			
Low APP	13.71 (1.98)	0.52	3, 90	.67
Med APP	15.00 (3.46)			
High APP	14.31 (3.55)			
Socialcognition Measures				
FKK Internal				
No APP	34.41 (7.18)			
Low APP	34.43 (5.56)	.12	3, 41	.95
Med APP	34.67 (5.75)			
High APP	32.60 (5.41)			
FKK Self Concept				
No APP	33.56 (4.71)			
Low APP	32.71 (5.19)	.06	3, 41	.98
Med APP	33.17 (5.95)			
High APP	33.00 (5.34)			
FKK Self Efficacy				
No APP	67.96 (9.95)			
Low APP	67.14 (7.71)	.10	3, 41	.96
Med APP	67.83 (9.66)			
High APP	65.60 (6.99)			
FKK Powerful Others				
No APP	25.96 (8.80)			
Low APP	25.71 (4.54)	.19	3, 41	.90
Med APP	25.17 (8.31)			
High APP	23.00 (7.58)			
FKK Chance				
No APP	23.48 (7.20)			
Low APP	24.29 (8.32)	.23	3, 41	.87
Med APP	26.00 (9.59)			
High APP	25.40 (6.11)			
FKK Externality				
No APP	49.44 (15.53)			
Low APP	50.00 (10.61)	.04	3, 41	.99
Med APP	51.17 (16.51)			
High APP	48.40 (7.57)			
IPSAQ Internal Positive				
No APP	8.19 (3.50)			
Low APP	9.00 (3.32)	.53	3, 41	.67
Med APP	6.83 (2.64)			
High APP	7.60 (1.95)			
IPSAQ Personal Positive				
No APP	3.52 (2.39)			
Low APP	2.71 (2.22)			
Med APP	4.50 (2.07)	.70	3, 41	.56
High APP	3.60 (0.55)			
IPSAQ Situational Positive				
No APP	3.26 (2.35)			
Low APP	3.29 (2.43)	.39	3, 41	.76
Med APP	3.67 (4.27)			
High APP	4.60 (1.67)			
IPSAQ Internal Negative				
No APP	5.89 (3.77)			
Low APP	6.00 (3.37)	.01	3, 41	1.0
Med APP	5.67 (3.33)			
High APP	6.00 (2.35)			

IPSAQ Personal Negative				
No APP	4.89 (3.48)			
Low APP	4.29 (3.04)	.38	3, 41	.77
Med APP	4.50 (3.27)			
High APP	6.20 (1.92)			
IPSAQ Situational Negative				
No APP	4.07 (2.80)			
Low APP	4.43 (4.35)	.27	3, 41	.85
Med APP	5.17 (2.71)			
High APP	3.80 (1.64)			
CST Social Support Seeking				
No APP	35.58 (9.99)			
Low APP	37.29 (8.40)	.09	3, 40	.96
Med APP	37.17 (13.17)			
High APP	35.00 (10.03)			
CST Self-Controlling				
No APP	52.65 (6.97)			
Low APP	51.00 (5.60)	2.49	3, 40	.07
Med APP	51.17 (6.08)			
High APP	41.60 (16.99)			
CST Escape Avoidance				
No APP	37.12 (6.73)			
Low APP	34.86 (2.19)	.36	3, 40	.78
Med APP	36.50 (5.58)			
High APP	35.00 (6.21)			
CST Planful Problem Solving				
No APP	42.08 (8.85)			
Low APP	40.57 (3.99)	.51	3, 40	.68
Med APP	45.67 (14.50)			
High APP	39.20 (10.31)			

Table 15
Bivariate Correlations for Insight Variables

	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>
1- Total Insight	-						
2 - Ability to relabel psychotic experiences	.73 **	-					
3 - Awareness of Illness	.89 **	.45 **	-				
4 - Need for Treatment	.83 **	.39 **	.72 **	-			
5 - SAIQ Need for Treatment	.80 **	.76 **	.76 **	.57 **	-		
6 - SAIQ Worry	-.80 **	-.77 **	-.77 **	-.57 **	-.95 **	-	
7 - SAIQ Presence/Outcome of Illness	-.80 **	-.77 **	-.77 **	-.58 **	-.98 **	.95 **	-

** Correlation is significant at the 0.01 level (1-tailed).

* Correlation is significant at the 0.05 level (1-tailed).

Table 16
Service Use by Insight Measures at Admission One-Way Analyses of Variance (ANOVAs)

<u>Insight Measures</u>	<u>M(SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
IS Relabel Symptoms Scale				
No Service	2.46 (1.11)	.05	1, 107	.82
Service Use	2.42 (1.15)			
IS Awareness of Illness Scale				
No Service	2.34 (1.37)	.14	1, 107	.71
Service Use	2.25 (1.28)			
IS Need for Treatment Scale				
No Service	2.37 (1.17)	.17	1, 107	.68
Service Use	2.27 (1.19)			
IS Total Score				
No Service	7.17 (3.11)	.29	1, 107	.59
Service Use	6.86 (2.91)			
SAIQ Need for Treatment				
No Service	6.10 (6.67)	2.27	1, 15	.15
Service Use	1.71 (4.54)			
SAIQ Worry				
No Service	26.90 (8.63)	1.27	1, 15	.28
Service Use	31.71 (8.69)			
SAIQ Presence/Outcome of Illness				
No Service	12.20 (3.05)	2.56	1, 15	.13
Service Use	14.29 (1.89)			

Table 17
APP Level by Insight Measures at Admission One-Way Analyses of Variance (ANOVAs)

<u>Insight Measures</u>	<u>M(SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
IS Relabel Symptoms Scale				
No APP	2.46 (1.11)			
Low APP	2.44 (1.60)	1.29	3, 105	.28
Med APP	1.80 (.92)			
High APP	2.59 (1.05)			
IS Awareness of Illness Scale				
No APP	2.34 (1.37)			
Low APP	2.67 (1.32)	1.19	3, 105	.32
Med APP	1.60 (1.26)			
High APP	2.32 (1.25)			
IS Need for Treatment Scale				
No APP	2.37 (1.17)			
Low APP	2.72 (.97)	1.17	3, 105	.32
Med APP	1.75 (.79)			
High APP	2.31 (1.30)			
IS Total Score				
No APP	7.17 (3.11)			
Low APP	7.83 (3.30)	1.59	3, 105	.20
Med APP	5.15 (1.76)			
High APP	7.10 (2.94)			
SAIQ Need for Treatment				
No APP	6.10 (6.67)			
Low APP	6.00 (8.49)	1.20	3, 13	.35
Med APP	0.00 (0)			
High APP	0.00 (0)			
SAIQ Worry				
No APP	26.90 (8.63)			
Low APP	23.50 (16.26)	1.32	3, 13	.31
Med APP	35.00 (0.0)			
High APP	35.00 (0.0)			
SAIQ Presence/Outcome of Illness				
No APP	12.20 (3.05)			
Low APP	12.50 (3.54)	1.21	3, 13	.35
Med APP	15.00 (0.0)			
High APP	15.00 (0.0)			

Table 18
Bivariate Correlations of NOSIE subscales

	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>
1 ó NOSIE Daily Schedule Competence	-						
2 ó NOSIE Social Interest	.46**	-					
3 ó NOSIE Neatness	.79 **	.50**	-				
4 ó NOSIE Irritability	-.65**	-.29**	-.49**	-			
5 ó NOSIE Psychoticism	-.51**	-.24**	-.39**	.50**	-		
6 ó NOSIE Motor Retardation	-.73**	-.52**	-.68**	.25**	.21*	-	
7 ó NOSIE Total	.88**	.67**	.84**	-.69**	-.54**	-.74**	-

** Correlation is significant at the 0.01 level (2-tailed).

* Correlation is significant at the 0.05 level (2-tailed).

Table 19
Service Use by NOSIE scales scores at Admission One-Way Analyses of Variance (ANOVAs)

<u>NOSIE Subscales</u>	<u>M (SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
Daily Schedule Competence				
No Service	33.57 (6.81)			
Service Use	31.84 (6.72)	1.98	1, 119	.16
Social Interest				
No Service	15.66 (6.45)			
Service Use	15.53 (6.92)	.01	1, 119	.92
Neatness				
No Service	23.77 (5.50)			
Service Use	22.50 (5.77)	1.54	1, 119	.22
Irritability				
No Service	4.00 (5.10)			
Service Use	5.48 (6.52)	1.95	1, 119	.17
Psychoticism				
No Service	1.41 (2.66)			
Service Use	1.66 (2.61)	.28	1, 119	.60
Motor Retardation				
No Service	5.93 (4.13)			
Service Use	6.82 (4.53)	1.29	1, 119	.26
NOSIE Total Score				
No Service	157.68 (24.13)			
Service Use	150.50 (26.26)	2.46	1, 119	.12

Table 20
APP Level by NOSIE scale scores at Admission One-Way Analyses of Variance (ANOVAs)

<u>NOSIE Subscales</u>	<u>M (SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
Daily Schedule Competence				
No APP	33.57 (6.81)			
Low APP	31.79 (8.59)			
Med APP	30.96 (5.96)	.82	3, 117	.49
High APP	32.50 (5.89)			
Social Interest				
No APP	15.66 (6.45)			
Low APP	14.72 (6.92)			
Med APP	14.97 (6.79)	.29	3, 117	.83
High APP	16.49 (7.18)			
Neatness				
No APP	23.77 (5.50)			
Low APP	21.60 (6.67)			
Med APP	20.73 (5.23)	2.18	3, 117	.09
High APP	24.40 (5.08)			
Irritability				
No APP	4.00 (5.10)			
Low APP	5.67 (7.15)			
Med APP	4.00 (5.09)	1.21	3, 117	.31
High APP	6.38 (7.01)			
Psychoticism				
No APP	1.41 (2.66)			
Low APP	1.57 (2.40)			
Med APP	1.84 (3.48)	.13	3, 117	.94
High APP	1.60 (2.12)			
Motor Retardation				
No APP	5.93 (4.13)			
Low APP	7.61 (5.84)	2.70	3, 117	.05
Med APP	8.42 (4.40)			
High APP	5.14 (2.90)			
NOSIE Total Score				
No APP	157.68 (24.13)			
Low APP	144.50 (32.32)	1.60	3, 117	.19
Med APP	148.40 (21.92)			
High APP	156.24 (24.17)			

Table 21
Bivariate Correlations of BPRS Total Scores, BPRS Factor Scores, and Symptomatology Measures

	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>	<u>8</u>	<u>9</u>	<u>10</u>	<u>11</u>	<u>12</u>	<u>13</u>	<u>14</u>
1. BPRS Total	-													
2. Psychotic Disorganization	.75 **	-												
3. Hallucinations/ Delusions	.75 **	.53* *	-											
4. Paranoia	.75 **	.48* *	.52 **	-										
5. Emotional Blunting	.19 *	.07	-.04	-.05	-									
6. Anxiety/ Depression	.56 **	.12	.24 **	.40 **	-.05	-								
7. Agitation/ Elation	.44 **	.32* *	.29 **	.27 **	-.33 **	.35 **	-							
8. BHS Total	.32 **	.07	.14	.18	.25 *	.35 **	-.13	-						
9. BDI-II Total	.26 *	.15	.13	.09	.20	.28 *	-.16	.70 **	-					
10. SPS Total	.46 **	.26* *	.30 *	.28 *	.21	.29 *	.15	.59 **	.62 **	-				
11. SPS Hopelessness Scale	.37 **	.15	.26 *	.21	.17	.30 *	.02	.43 **	.63 **	.80 **	-			
12. SPS Suicidality Scale	.40 **	.27* *	.23	.19	.19	.25 *	.08	.32 **	.55 **	.66 **	.74 **	-		
13. SPS Negative Self-Evaluation Scale	.11	.14	.10	.08	.08	-.02	-.09	.37 **	.27 **	.50 **	.38 **	.49 **	-	
14. SPS Hostility Scale	.21	.08	.12	.03	.19	.16	.13	.36 **	.48 **	.64 **	.71 **	.60 **	.42 **	-

** Correlation is significant at the 0.01 level (2-tailed).

* Correlation is significant at the 0.05 level (2-tailed).

Table 22
Service Use by BPRS Factor Scores, BPRS Total, and Symptomatology Measures at Admission
One-Way Analyses of Variance (ANOVAs)

	<u>M (SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
BPRS Total				
No Service	47.60 (12.15)	.10	1, 154	.75
Service Use	48.29 (14.62)			
<u>BPRS Factor Scores</u>				
Psychotic Disorganization				
No Service	6.35 (2.76)	.99	1, 154	.32
Service Use	6.84 (3.37)			
Emotional Blunting				
No Service	5.96 (3.65)	.04	1, 155	.85
Service Use	6.07 (3.29)			
Paranoia				
No Service	9.00 (3.21)	2.65	1, 155	.11
Service Use	8.14 (3.36)			
Anxiety/Depression				
No Service	9.03 (3.46)	2.03	1, 155	.16
Service Use	9.90 (4.12)			
Hallucinations/Delusions				
No Service	6.71 (4.08)	.23	1, 155	.63
Service Use	6.39 (4.17)			
Agitation/Elation				
No Service	3.49 (1.55)	1.51	1, 155	.22
Service Use	3.82 (1.79)			
BHS Total				
No Service	4.24 (4.55)	2.22	1, 118	.14
Service Use	5.58 (5.19)			
BDI-II Total				
No Service	9.33 (10.10)	2.20	1, 99	.14
Service Use	12.96 (13.82)			
SPS Total				
No Service	56.02 (10.19)	7.41	1, 122	.01
Service Use	61.02 (10.22)			
SPS Hopelessness Scale				
No Service	53.33 (10.42)	3.76	1, 122	.05
Service Use	57.48 (13.09)			
SPS Suicidality Scale				
No Service	51.41 (8.74)	1.97	1, 122	.16
Service Use	54.35 (13.65)			
SPS Negative Self-Evaluation Scale				
No Service	59.48 (9.79)			
Service Use	58.60 (12.56)	.19	1, 121	.67
SPS Hostility Scale				
No Service	52.81 (10.57)	3.30	1, 121	.07
Service Use	56.88 (13.81)			

Table 23
APP Level by BPRS Factor Scores, BPRS Total, and Symptomatology Measures at Admission One-Way Analyses of Variance (ANOVAs)

	<u>M (SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
BPRS Total				
No APP	47.60 (12.15)			
Low APP	45.96 (11.94)	.32	3, 152	.81
Med APP	46.91 (14.91)			
High APP	49.34 (15.28)			
<u>BPRS Factor Scores</u>				
Psychotic Disorganization				
No APP	6.35 (2.76)			
Low APP	6.70 (3.59)	.43	3, 152	.73
Med APP	6.52 (2.94)			
High APP	6.98 (3.51)			
Emotional Blunting				
No APP	5.97 (3.65)			
Low APP	6.92 (2.71)	.44	3, 153	.72
Med APP	5.47 (2.76)			
High APP	6.06 (3.58)			
Paranoia				
No APP	9.00 (3.21)			
Low APP	7.58 (3.83)	1.03	3, 153	.38
Med APP	8.14 (3.52)			
High APP	8.28 (3.24)			
Anxiety/Depression				
No APP	9.03 (3.46)			
Low APP	8.42 (2.95)	1.48	3, 153	.22
Med APP	9.94 (4.64)			
High APP	10.25 (4.18)			
Hallucinations/Delusions				
No APP	6.71 (4.08)			
Low APP	6.15 (3.48)	.19	3, 153	.91
Med APP	5.97 (3.53)			
High APP	6.59 (4.56)			
Agitation/Elation				
No APP	3.49 (1.55)			
Low APP	3.23 (1.20)	1.26	3, 153	.29
Med APP	4.15 (1.89)			
High APP	3.86 (1.88)			
BHS Total				
No APP	4.24 (4.55)			
Low APP	5.69 (6.20)	1.54	3, 116	.21
Med APP	7.38 (3.93)			
High APP	4.98 (5.18)			
BDI-II Total				
No APP	9.33 (10.10)			
Low APP	11.71 (14.84)	1.84	3, 97	.14
Med APP	18.91 (15.75)			

High APP	11.37 (12.43)			
SPS Total				
No APP	56.02 (10.19)			
Low APP	58.43 (11.32)	3.34	3, 120	.02
Med APP	64.50 (8.40)			
High APP	60.68 (10.31)			
SPS Hopelessness Scale				
No APP	53.33 (10.42)			
Low APP	56.71 (13.63)	1.57	3, 120	.20
Med APP	60.29 (12.95)			
High APP	56.74 (13.16)			
SPS Suicidality Scale				
No APP	51.41 (8.74)			
Low APP	54.64 (10.95)	.65	3, 120	.58
Med APP	54.00 (15.51)			
High APP	54.37 (14.17)			
SPS Negative Self-Evaluation Scale				
No APP	59.48 (9.79)			
Low APP	58.14 (9.67)	.20	3, 119	.89
Med APP	57.00 (15.58)			
High APP	59.32 (12.64)			
SPS Hostility Scale				
No APP	52.81 (10.57)			
Low APP	57.71 (10.77)	1.53	3, 119	.21
Med APP	59.92 (16.73)			
High APP	55.53 (13.89)			

Table 24
Service Use by BPRS Items at admission One-Way Analyses of Variance (ANOVAs)

<u>BPRS Items</u>		<u>M (SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
Somatic Concern	No Service	2.50 (1.63)	.30	1, 156	.59
	Service Use	2.36 (1.57)			
Anxiety	No Service	2.63 (1.40)	.14	1, 156	.71
	Service Use	2.72 (1.62)			
Depression	No Service	2.19 (1.19)	2.19	1, 156	.14
	Service Use	2.52 (1.53)			
Suicidality	No Service	1.23 (0.62)	10.36	1, 156	.002*
	Service Use	1.79 (1.37)			
Guilt	No Service	1.84 (1.24)	2.85	1, 156	.09
	Service Use	2.19 (1.37)			
Hostility	No Service	2.53 (1.59)	.42	1, 156	.52
	Service Use	2.70 (1.74)			
Elevated Mood	No Service	1.51 (0.96)	.32	1, 156	.57
	Service Use	1.61 (1.25)			
Grandiosity	No Service	2.13 (1.82)	3.54	1, 156	.06
	Service Use	1.63 (1.52)			
Suspiciousness	No Service	2.77 (1.80)	2.63	1, 156	.11
	Service Use	2.33 (1.58)			
Hallucinations	No Service	1.87 (1.55)	3.66	1, 156	.06
	Service Use	2.41 (1.97)			
Unusual Thought Content	No Service	2.71 (1.99)	1.29	1, 156	.23
	Service Use	2.35 (1.96)			
Bizarre Behavior	No Service	1.93 (1.49)	1.60	1, 156	.21
	Service Use	2.25 (1.61)			
Self-Neglect	No Service	2.05 (0.94)	3.55	1, 156	.06
	Service Use	2.39 (1.24)			
Disorientation	No Service	1.33 (0.70)	.92	1, 155	.34
	Service Use	1.45 (0.90)			
Conceptual Disorganization	No Service	2.06 (1.40)	.65	1, 155	.42
	Service Use	1.88 (1.43)			
Blunted Affect	No Service	2.54 (1.48)	.04	1, 155	.84
	Service Use	2.59 (1.30)			
Emotional Withdrawal	No Service	2.25 (1.41)	.11	1, 155	.74
	Service Use	2.18 (1.21)			
Motor Retardation	No Service	2.00 (1.17)	.01	1, 155	.94
	Service Use	2.01 (1.19)			
Tension	No Service	1.75 (1.16)	1.53	1, 155	.22
	Service Use	1.55 (0.85)			
Uncooperativeness	No Service	1.83 (1.23)	3.91	1, 155	.05*
	Service Use	1.49 (0.91)			
Excitement	No Service	1.63 (1.12)	1.99	1, 155	.16
	Service Use	1.40 (0.93)			
Distractibility	No Service	1.78 (1.15)	.54	1, 155	.46
	Service Use	1.93 (1.33)			
Motor hyperactivity	No Service	1.37 (0.88)	.23	1, 155	.63
	Service Use	1.30 (0.73)			
Mannerisms and Posturing	No Service	1.17 (0.55)	.11	1, 155	.74
	Service Use	1.20 (0.58)			

*p<.05

Table 25
APP Severity Level by BPRS Items at admission One-Way Analyses of Variance (ANOVAs)

<u>BPRS Items</u>		<u>M (SD)</u>	<u>F</u>	<u>df</u>	<u>p</u>
Somatic Concern	No APP	2.50 (1.63)	1.18	3, 154	.32
	Low APP	1.77 (0.93)			
	Med APP	2.12 (1.55)			
	High APP	2.58 (1.67)			
Anxiety	No APP	2.63 (1.40)	.68	3, 154	.57
	Low APP	2.19 (1.15)			
	Med APP	2.74 (2.06)			
	High APP	2.84 (1.56)			
Depression	No APP	2.19 (1.19)	.91	3, 154	.44
	Low APP	2.35 (1.55)			
	Med APP	2.38 (1.57)			
	High APP	2.60 (1.54)			
Suicidality	No APP	1.23 (0.62)	5.34	3, 154	.002*
	Low APP	1.15 (0.56)			
	Med APP	1.94 (1.35)			
	High APP	1.90 (1.49)			
Guilt	No APP	1.84 (1.24)	1.49	3, 154	.22
	Low APP	1.77 (1.09)			
	Med APP	2.29 (1.56)			
	High APP	2.26 (1.37)			
Hostility	No APP	2.53 (1.59)	1.11	3, 154	.35
	Low APP	2.00 (1.29)			
	Med APP	2.68 (1.88)			
	High APP	2.88 (1.77)			
Elevated Mood	No APP	1.51 (0.96)	.45	3, 154	.72
	Low APP	1.35 (0.75)			
	Med APP	1.76 (1.47)			
	High APP	1.62 (1.29)			
Grandiosity	No APP	2.13 (1.82)	1.18	3, 154	.32
	Low APP	1.54 (1.39)			
	Med APP	1.65 (1.69)			
	High APP	1.65 (1.52)			
Suspiciousness	No APP	2.77 (1.80)	.90	3, 154	.45
	Low APP	2.42 (1.66)			
	Med APP	2.24 (1.52)			
	High APP	2.34 (1.61)			
Hallucinations	No APP	1.87 (1.55)	1.70	3, 154	.17
	Low APP	1.92 (1.80)			
	Med APP	2.29 (1.90)			
	High APP	2.57 (2.05)			
Unusual Thought Content	No APP	2.71 (1.99)	.71	3,153	.55
	Low APP	2.69 (1.97)			
	Med APP	2.03 (2.04)			
	High APP	2.37 (1.96)			
Bizarre Behavior	No APP	1.93 (1.49)	.59	3, 154	.63
	Low APP	2.35 (1.75)			
	Med APP	2.12 (1.69)			
	High APP	2.26 (1.58)			
Self-Neglect	No APP	2.05 (0.94)	2.01	3, 154	.11
	Low APP	2.08 (0.95)			
	Med APP	2.71 (1.23)			
	High APP	2.36 (1.30)			
Disorientation	No APP	1.33 (0.70)	3.73	3, 153	.01*
	Low APP	2.08 (1.50)			
	Med APP	1.21 (0.47)			
	High APP	1.38 (0.76)			
Conceptual Disorganization	No APP	2.06 (1.40)	.55	3, 153	.65
	Low APP	2.23 (1.59)			
	Med APP	1.88 (1.46)			
	High APP	1.79 (1.39)			

Blunted Affect	No APP	2.54 (1.48)			
	Low APP	3.00 (1.29)			
	Med APP	2.38 (1.17)	.53	3, 153	.66
	High APP	2.55 (1.34)			
Emotional Withdrawal	No APP	2.25 (1.41)			
	Low APP	2.38 (1.12)			
	Med APP	2.03 (1.07)	.22	3, 153	.88
	High APP	2.17 (1.28)			
Motor Retardation	No APP	2.00 (1.17)			
	Low APP	2.15 (1.07)			
	Med APP	1.82 (1.03)	.22	3, 153	.89
	High APP	2.04 (1.28)			
Tension	No APP	1.75 (1.16)			
	Low APP	1.54 (0.88)			
	Med APP	1.56 (.97)	.50	3, 153	.68
	High APP	1.56 (.81)			
Uncooperativeness	No APP	1.83 (1.23)			
	Low APP	1.62 (1.12)			
	Med APP	1.53 (1.13)	1.38	3, 153	.25
	High APP	1.45 (0.78)			
Excitement	No APP	1.63 (1.12)			
	Low APP	1.23 (0.45)			
	Med APP	1.53 (1.18)	.86	3, 153	.46
	High APP	1.40 (0.94)			
Distractibility	No APP	1.78 (1.15)			
	Low APP	1.54 (0.78)			
	Med APP	1.56 (0.86)	1.64	3, 153	.18
	High APP	2.14 (1.52)			
Motor hyperactivity	No APP	1.37 (0.88)			
	Low APP	1.31 (0.63)			
	Med APP	1.41 (1.00)	.21	3, 153	.89
	High APP	1.27 (0.65)			
Mannerisms and Posturing	No APP	1.17 (0.55)			
	Low APP	1.31 (0.63)			
	Med APP	1.06 (0.24)	.58	3, 152	.63
	High APP	1.23 (0.64)			

*p<.05

Table 26
Service Use by WCST Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>			
	WCST Correct F (2,72)	WCST Perseverative Errors F (2,112)	WCST Random Errors F (2, 112)	WCST Consolidation Index F (2, 72)
Main Effect Assessment Time	.66	2.53	3.66*	1.92
Main Effect Service Use	1.35	5.13*	.71	.71
Interaction Assessment Time * Service Use	.64	1.36	.84	.10

*p<.05, **p<.01, ***p<.001

Table 27
*Service Use by Neurocognition Measures
 Repeated Measures ANOVAs*

<u>Variable</u>	<u>Measures</u>					
	RAVLT F (2,18)	NAB Total F (2,26)	Trails A F (2, 110)	Trails B F (2, 112)	COWAT/ FAS F (2, 114)	RBANS Total F (2, 108)
Main Effect Assessment Time	.24	1.74	.51	.12	1.91	9.82***
Main Effect Service Use	2.19	.28	.004	.20	.06	1.67
Interaction Assessment Time * Service Use	5.25*	.47	3.38*	.53	3.02*	3.65*

*p<.05, **p<.01, ***p<.001

Table 28
Service Use by RCFT Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>			
	RCFT Copy F (2,60)	RCFT Immediate Copy F (2,60)	RCFT Delayed Memory F (2, 60)	RCFT Recognition F (2, 60)
Main Effect Assessment Time	1.75	4.62**	4.89**	.55
Main Effect Service Use	.32	3.60	1.69	.04
Interaction Assessment Time * Service Use	.19	1.37	.13	.63

*p<.05, **p<.01, ***p<.001

Table 29
*Mean Scores and Standard Deviations for WCST Correct Scores at Admission, 6 Months,
 and 12 Months as a Function of Service Use Before Age 18*

<u>Services Before Age 18</u>	<u>WCST Correct Scores</u>		
	<u>At Admission</u> M (SD)	<u>Six Months</u> M (SD)	<u>Twelve Months</u> M (SD)
No Service Use	24.35 (2.67)	23.41 (4.23)	23.94 (3.31)
Service Use	24.86 (.66)	24.81 (.87)	24.33 (2.83)

Table 30

Mean Scores and Standard Deviations for WCST Perseverative Errors Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>WCST Perseverative Errors Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	24.52 (13.52)	30.70 (18.50)	24.93 (15.98)
Service Use	22.10 (11.41)	21.52 (12.09)	17.94 (11.23)

Table 31

Mean Scores and Standard Deviations for WCST Random Errors Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>WCST Random Error Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	27.89 (15.98)	29.81 (18.19)	21.74 (18.47)
Service Use	24.90 (18.71)	25.26 (17.51)	20.23 (17.51)

Table 32

Mean Scores and Standard Deviations for WCST Consolidation Index Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>WCST Consolidation Index Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	4.59 (5.77)	4.58 (8.42)	11.87 (32.49)
Service Use	2.84 (4.90)	2.74 (3.21)	7.43 (19.47)

Table 33
Mean Scores and Standard Deviations for RAVLT Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>RAVLT Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	6.80 (3.19)	5.80 (3.11)	8.20 (3.27)
Service Use	9.00 (2.19)	10.33 (3.14)	8.50 (2.51)

Table 34
Mean Scores and Standard Deviations for NAB Total Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>NAB Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	70.00 (10.23)	75.00 (3.74)	77.75 (9.32)
Service Use	75.91 (15.98)	80.09 (14.96)	78.27 (14.16)

Table 35
Mean Scores and Standard Deviations for Trails A Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>Trails A Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	42.88 (14.57)	50.30 (19.59)	43.03 (14.97)
Service Use	47.04 (28.23)	42.76 (30.58)	45.24 (38.51)

Table 36
Mean Scores and Standard Deviations for Trails B Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>Trails B Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	107.84 (48.26)	112.83 (58.77)	112.03 (93.06)
Service Use	126.91 (84.19)	112.66 (94.25)	117.02 (86.00)

Table 37
Mean Scores and Standard Deviations for COWAT/FAS Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>COWAT/FAS Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	30.71 (9.45)	26.75 (12.70)	28.82 (8.87)
Service Use	26.74 (9.27)	27.68 (9.86)	30.32 (9.62)

Table 38
Mean Scores and Standard Deviations for RBANS Total Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>RBANS Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	73.77 (15.39)	73.19 (13.49)	80.84 (16.03)
Service Use	68.32 (13.57)	72.60 (13.45)	73.04 (14.62)

Table 39
Mean Scores and Standard Deviations for RCFT Copy Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>RCFT Copy Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	29.85 (6.85)	28.46 (5.52)	29.85 (5.97)
Service Use	28.42 (5.95)	26.92 (7.38)	29.55 (6.38)

Table 40
Mean Scores and Standard Deviations for RCFT Immediate Memory Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>RCFT Immediate Memory Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	11.58 (5.20)	12.31 (5.53)	16.12 (6.85)
Service Use	15.82 (1.68)	15.76 (9.97)	29.92 (29.46)

Table 41
Mean Scores and Standard Deviations for RCFT Delayed Memory Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>RCFT Delayed Memory Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	12.12 (4.98)	11.88 (5.14)	15.23 (6.69)
Service Use	15.17 (11.66)	15.39 (9.43)	19.71 (11.14)

Table 42
Mean Scores and Standard Deviations for RCFT Recognition Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>RCFT Recognition Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	19.08 (2.53)	18.92 (1.71)	19.08 (2.84)
Service Use	18.21 (6.00)	19.68 (7.10)	19.95 (2.70)

Table 43
APP Severity Level by WCST Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>			
	WCST Correct	WCST Perseverative Errors	WCST Random Errors	WCST Consolidation Index
	F (6,68)	F (6,108)	F (6, 108)	F (6, 68)
Main Effect				
Assessment Time	.20	.73	1.81	.77
Main Effect				
APP Severity Level	.54	2.66	1.87	.40
Interaction				
Assessment Time * APP Severity Level	.26	1.09	1.30	.08

*p<.05, **p<.01, ***p<.001

Table 44
*APP Severity Level by Neurocognition Measures
 Repeated Measures ANOVAs*

<u>Variable</u>	<u>Measures</u>					
	RAVLT	NAB Total	Trails A	Trails B	COWAT/ FAS	RBANS Total
	F (4,16)	F (4,24)	F (6, 106)	F (6, 106)	F (6, 110)	F (6, 104)
Main Effect Assessment Time	.15	.39	.20	.56	1.16	6.37**
Main Effect APP Severity Level	1.06	.30	1.58	.84	.09	.80
Interaction Assessment Time * APP Severity Level	3.88*	1.40	1.28	.55	1.67	2.40*

*p<.05, **p<.01, ***p<.001

Table 45
Service Use by RCFT Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>			
	RCFT Copy	RCFT Immediate Copy	RCFT Delayed Memory	RCFT Recognition
	F (6,56)	F (6,56)	F (6, 56)	F (6, 56)
Main Effect Assessment Time	2.20	5.66*	6.23**	1.33
Main Effect Service Use	.80	1.27	.55	1.19
Interaction Assessment Time * Service Use	.25	1.76	2.14	1.46

*p<.05, **p<.01, ***p<.001

Table 46
Mean Scores and Standard Deviations for WCST Correct Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>WCST Correct Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	24.35 (2.67)	23.41 (4.23)	23.94 (3.31)
Low APP	25.00 (.00)	25.00 (.00)	25.00 (.00)
Medium APP	25.00 (.00)	25.00 (.00)	24.80 (.45)
High APP	24.75 (.87)	24.67 (1.16)	23.92 (3.75)

Table 47
Mean Scores and Standard Deviations for WCST Perseverative Errors Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>WCST Perseverative Errors Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	24.52 (13.52)	30.70 (18.50)	24.93 (15.98)
Low APP	22.50 (11.79)	26.25 (6.19)	19.50 (12.77)
Medium APP	18.29 (11.80)	10.00 (6.56)	16.14 (14.68)
High APP	23.35 (11.52)	24.60 (12.14)	18.25 (10.18)

Table 48
Mean Scores and Standard Deviations for WCST Random Errors Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>WCST Random Error Scores</u>		
	<u>At Admission</u> M (SD)	<u>Six Months</u> M (SD)	<u>Twelve Months</u> M (SD)
No APP	27.89 (15.98)	29.81 (18.19)	21.74 (18.47)
Low APP	23.00 (8.29)	40.50 (10.76)	22.00 (8.41)
Medium APP	14.00 (12.74)	9.71 (10.42)	17.71 (20.23)
High APP	29.10 (20.68)	27.65 (16.95)	20.75 (18.45)

Table 49
Mean Scores and Standard Deviations for WCST Consolidation Index Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>WCST Consolidation Index Scores</u>		
	<u>At Admission</u> M (SD)	<u>Six Months</u> M (SD)	<u>Twelve Months</u> M (SD)
No APP	4.59 (5.77)	4.58 (8.42)	11.87 (32.49)
Low APP	1.23 (.70)	1.14 (1.31)	1.85 (1.35)
Medium APP	5.31 (8.28)	4.17 (4.52)	8.87 (14.81)
High APP	2.36 (3.82)	2.67 (2.99)	8.69 (24.40)

Table 50
Mean Scores and Standard Deviations for RAVLT Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>RAVLT Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	6.80 (3.19)	5.80 (3.11)	8.20 (3.27)
Low APP	-	-	-
Medium APP	8.50 (2.12)	8.50 (3.54)	9.00 (1.41)
High APP	9.25 (2.50)	11.25 (2.99)	8.25 (3.10)

Table 51
Mean Scores and Standard Deviations for NAB Total Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>NAB Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	70.00 (10.23)	75.00 (3.74)	77.75 (9.32)
Low APP	-	-	-
Medium APP	78.00 (4.24)	75.00 (4.24)	67.00 (7.07)
High APP	75.44 (17.76)	81.22 (16.42)	80.78 (14.34)

Table 52
Mean Scores and Standard Deviations for Trails A Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>Trails A Total Scores</u>		
	<u>At Admission</u> M (SD)	<u>Six Months</u> M (SD)	<u>Twelve Months</u> M (SD)
No APP	42.88 (14.57)	50.30 (19.59)	43.03 (14.97)
Low APP	64.75 (30.39)	65.00 (27.83)	61.26 (23.81)
Medium APP	37.78 (18.91)	31.76 (12.55)	33.03 (14.78)
High APP	48.40 (31.61)	44.08 (36.65)	48.88 (49.41)

Table 53
Mean Scores and Standard Deviations for Trails B Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>Trails B Total Scores</u>		
	<u>At Admission</u> M (SD)	<u>Six Months</u> M (SD)	<u>Twelve Months</u> M (SD)
No APP	107.84 (48.26)	112.83 (58.77)	112.03 (93.06)
Low APP	174.00 (89.80)	160.50 (30.45)	127.34 (47.63)
Medium APP	100.03 (44.58)	96.32 (50.47)	92.53 (54.13)
High APP	132.27 (99.94)	110.81 (122.80)	130.60 (108.47)

Table 54
*Mean Scores and Standard Deviations for COWAT/FAS Scores at Admission, 6 Months,
 and 12 Months as a Function of APP Severity Level*

<u>APP Severity Level</u>	<u>COWAT/FAS Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	30.71 (9.45)	26.75 (12.70)	28.82 (8.87)
Low APP	22.50 (8.89)	30.75 (12.76)	28.25 (11.03)
Medium APP	27.90 (10.56)	26.20 (10.68)	28.80 (11.58)
High APP	27.06 (8.84)	27.82 (9.17)	31.71 (8.41)

Table 55
*Mean Scores and Standard Deviations for RBANS Total Scores at Admission, 6 Months,
 and 12 Months as a Function of APP Severity Level*

<u>APP Severity Level</u>	<u>RBANS Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	73.77 (15.39)	73.19 (13.49)	80.84 (16.03)
Low APP	70.60 (13.81)	73.80 (15.01)	81.40 (8.39)
Medium APP	68.56 (15.31)	70.22 (14.81)	67.44 (13.96)
High APP	67.09 (13.19)	74.00 (12.68)	73.82 (16.28)

Table 56
*Mean Scores and Standard Deviations for RCFT Copy Scores at Admission, 6 Months,
 and 12 Months as a Function of APP Severity Level*

<u>APP Severity Level</u>	<u>RCFT Copy Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	29.85 (6.85)	28.46 (5.52)	29.85 (5.97)
Low APP	27.88 (2.32)	27.25 (3.40)	31.25 (5.56)
Medium APP	26.75 (9.49)	23.67 (11.42)	26.83 (7.80)
High APP	29.78 (4.14)	28.94 (4.90)	30.61 (5.81)

Table 57
*Mean Scores and Standard Deviations for RCFT Immediate Memory Scores at
 Admission, 6 Months, and 12 Months as a Function of APP Severity Level*

<u>APP Severity Level</u>	<u>RCFT Immediate Memory Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	11.58 (5.20)	12.31 (5.53)	16.12 (6.85)
Low APP	14.75 (6.36)	14.25 (6.20)	30.00 (17.22)
Medium APP	19.33 (18.85)	18.25 (15.55)	18.42 (10.52)
High APP	13.94 (7.47)	14.78 (7.16)	37.56 (40.10)

Table 58
Mean Scores and Standard Deviations for RCFT Delayed Memory Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>RCFT Delayed Memory Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	12.12 (4.98)	11.88 (5.14)	15.23 (6.69)
Low APP	10.75 (5.87)	14.00 (6.72)	22.63 (10.48)
Medium APP	18.92 (18.69)	16.83 (14.17)	15.42 (11.56)
High APP	14.63 (7.38)	15.06 (7.40)	21.28 (11.56)

Table 59
Mean Scores and Standard Deviations for RCFT Recognition Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>RCFT Recognition Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	19.08 (2.53)	18.92 (1.71)	19.08 (2.84)
Low APP	16.75 (2.87)	19.25 (1.26)	20.25 (1.50)
Medium APP	21.17 (9.58)	23.50 (11.11)	19.83 (2.32)
High APP	16.89 (3.30)	17.33 (4.27)	19.89 (3.48)

Table 60
Service Use by FKK Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>					
	FKK Internal	FKK Self Concept	FKK Self Efficacy	FKK Powerful Others	FKK Chance	FKK Externality
	F (2,50)	F (2,50)	F (2, 50)	F (2, 50)	F (2, 50)	F (2, 50)
Main Effect Assessment Time	.27	1.14	.11	1.04	4.66**	3.77*
Main Effect Service Use	.06	.33	.25	.33	1.13	.74
Interaction Assessment Time * Service Use	.80	.71	.48	.90	5.29**	3.87*

*p<.05, **p<.01, ***p<.001

Table 61
*Service Use by Socialcognition Measures
 Repeated Measures ANOVAs*

<u>Variable</u>	<u>Measures</u>				
	Hinting F (2,102)	CST Social Support F (2,40)	CST Self Controlling F (2, 40)	CST Escape Avoidance F (2, 40)	CST Planful Problem Solving F (2, 40)
Main Effect Assessment Time	8.32***	4.83**	.14	1.74	.47
Main Effect Service Use	.01	.21	7.05*	.13	1.16
Interaction Assessment Time * Service Use	.21	.74	.24	1.92	.57

*p<.05, **p<.01, ***p<.001

Table 62
*Service Use by Socialcognition Measures
 Repeated Measures ANOVAs*

<u>Variable</u>	<u>Measures</u>					
	IPSAQ Internal Positive F (2,30)	IPSAQ Personal Positive F (2,30)	IPSAQ Situational Positive F (2, 30)	IPSAQ Internal Negative F (2, 30)	IPSAQ Personal Negative F (2, 30)	IPSAQ Situational Negative F (2, 30)
Main Effect Assessment Time	.39	4.00*	2.32	.39	.33	.45
Main Effect Service Use	2.15	.51	4.02	1.41	.63	.69
Interaction Assessment Time * Service Use	2.85	.93	2.18	3.41*	.47	2.14

*p<.05, **p<.01, ***p<.001

Table 63
*Mean Scores and Standard Deviations for Hinting Scores at Admission, 6 Months, and
 12 Months as a Function of Service Use Before Age 18*

<u>Services Before Age 18</u>	<u>Hinting Scores</u>		
	<u>At Admission</u> M (SD)	<u>Six Months</u> M (SD)	<u>Twelve Months</u> M (SD)
No Service Use	14.86 (3.31)	16.48 (3.12)	16.14 (2.90)
Service Use	14.50 (3.08)	16.75 (3.35)	16.04 (3.67)

Table 64
Mean Scores and Standard Deviations for FKK Internal Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

	<u>FKK Internal Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	33.88 (7.98)	33.94 (5.64)	34.88 (5.37)
Service Use	33.36 (6.53)	35.09 (5.63)	32.82 (6.35)

Table 65
Mean Scores and Standard Deviations for FKK Self Concept Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

	<u>FKK Self Concept Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	33.06 (5.08)	32.75 (3.84)	32.88 (3.74)
Service Use	33.55 (5.09)	31.09 (4.16)	31.55 (6.36)

Table 66
Mean Scores and Standard Deviations for FKK Self Efficacy Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

	<u>FKK Self Efficacy Index Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	66.94 (11.11)	66.69 (7.11)	67.75 (6.90)
Service Use	66.91 (8.85)	66.18 (7.01)	64.36 (9.68)

Table 67
Mean Scores and Standard Deviations for FKK Powerful Others Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>FKK Powerful Others Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	26.69 (9.19)	26.88 (7.00)	25.94 (7.23)
Service Use	26.73 (7.14)	23.73 (7.88)	24.36 (7.87)

Table 68
Mean Scores and Standard Deviations for FKK Chance Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>FKK Chance Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	24.69 (7.79)	25.63 (7.60)	24.19 (5.94)
Service Use	26.73 (9.09)	20.09 (6.07)	20.09 (5.19)

Table 69
Mean Scores and Standard Deviations for FKK Externality Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>FKK Externality Total Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	51.38 (16.47)	52.50 (13.29)	50.13 (12.12)
Service Use	53.45 (13.84)	43.82 (13.11)	44.45 (10.31)

Table 70
Mean Scores and Standard Deviations for IPSAQ Internal Positive Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IPSAQ Internal Positive Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	8.50 (2.83)	7.37 (3.29)	7.00 (3.89)
Service Use	7.67 (2.45)	10.11 (1.69)	9.33 (2.18)

Table 71
Mean Scores and Standard Deviations for IPSAQ Personal Positive Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IPSAQ Personal Positive Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	4.87 (2.36)	3.75 (1.49)	2.75 (2.82)
Service Use	3.56 (1.81)	3.44 (1.42)	2.78 (1.64)

Table 72
Mean Scores and Standard Deviations for IPSAQ Situational Positive Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IPSAQ Situational Positive Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	2.63 (1.51)	4.88 (2.42)	5.38 (2.62)
Service Use	3.33 (1.94)	2.44 (1.88)	3.89 (2.32)

Table 73
Mean Scores and Standard Deviations for IPSAQ Internal Negative Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IPSAQ Internal Negative Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	5.63 (4.21)	6.50 (4.72)	4.00 (3.38)
Service Use	6.22 (2.22)	6.78 (3.42)	8.22 (3.11)

Table 74
Mean Scores and Standard Deviations for IPSAQ Personal Negative Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IPSAQ Personal Negative Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	6.13 (4.19)	5.13 (3.68)	4.87 (3.60)
Service Use	4.22 (3.03)	4.56 (1.94)	4.22 (2.91)

Table 75
Mean Scores and Standard Deviations for IPSAQ Situational Negative Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IPSAQ Situational Negative Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	4.25 (2.12)	4.25 (2.05)	6.25 (1.83)
Service Use	4.11 (2.62)	4.56 (4.04)	3.56 (3.21)

Table 76
Mean Scores and Standard Deviations for CST Social Support Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>CST Social Support Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	32.75 (10.11)	37.17 (9.16)	40.67 (8.33)
Service Use	37.10 (11.06)	36.80 (9.88)	41.50 (10.36)

Table 77
Mean Scores and Standard Deviations for CST Self Controlling Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>CST Self Controlling Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	54.50 (5.14)	54.08 (7.10)	54.75 (4.83)
Service Use	48.90 (5.41)	48.00 (9.15)	46.90 (11.18)

Table 78
Mean Scores and Standard Deviations for CST Escape Avoidance Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>CST Escape Avoidance Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	33.67 (5.28)	36.25 (6.08)	38.17 (3.74)
Service Use	35.60 (5.48)	39.70 (4.57)	34.90 (12.21)

Table 79

Mean Scores and Standard Deviations for CST Planful Problem Solving Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>CST Planful Problem Solving Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
<u>Services Before Age 18</u>			
No Service Use	42.08 (8.85)	43.33 (5.03)	45.17 (6.24)
Service Use	41.40 (11.05)	39.10 (7.88)	40.90 (10.78)

Table 80

APP Severity Level by FKK Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>					
	FKK Internal	FKK Self Concept	FKK Self Efficacy	FKK Powerful Others	FKK Chance	FKK Externality
	F (6,46)	F (6,46)	F (6,46)	F (6,46)	F (6,46)	F (6,46)
Main Effect Assessment Time	.54	1.23	.09	1.11	7.81***	5.47**
Main Effect APP Severity Level	.06	.26	.12	.25	.35	.29
Interaction Assessment Time * APP	1.15	1.17	1.19	1.09	2.88*	2.30*

*p<.05, **p<.01, ***p<.001

Table 81
*APP Severity Level by Socialcognition Measures
 Repeated Measures ANOVAs*

<u>Variable</u>	<u>Measures</u>				
	Hinting F (6,98)	CST Social Support F (6,36)	CST Self Controlling F (6, 36)	CST Escape Avoidance F (6, 36)	CST Planful Problem Solving F (6, 36)
Main Effect Assessment Time	5.87**	2.71	.48	2.15	.21
Main Effect APP Severity Level	.80	.12	3.39*	1.12	.67
Interaction Assessment Time * APP	.90	.32	1.81	2.26	.52

*p<.05, **p<.01, ***p<.001

Table 82
*APP Severity Level by Socialcognition Measures
 Repeated Measures ANOVAs*

<u>Variable</u>	<u>Measures</u>					
	IPSAQ Internal Positive F (6,26)	IPSAQ Personal Positive F (6,26)	IPSAQ Situational Positive F (6, 26)	IPSAQ Internal Negative F (6, 26)	IPSAQ Personal Negative F (6, 26)	IPSAQ Situational Negative F (6, 26)
Main Effect Assessment Time	1.57	1.74	1.05	1.20	.13	.02
Main Effect APP Severity Level	1.55	.36	4.23*	.66	.24	.64
Interaction Assessment Time * APP	1.30	1.18	1.77	3.21*	.46	1.85

*p<.05, **p<.01, ***p<.001

Table 83
Mean Scores and Standard Deviations for Hinting Task Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>Hinting Task Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	14.86 (3.31)	16.48 (3.12)	16.14 (2.90)
Low APP	12.33 (.58)	14.67 (3.06)	17.33 (.58)
Medium APP	14.29 (3.64)	15.86 (3.13)	14.43 (4.93)
High APP	15.07 (3.03)	17.64 (3.41)	16.57 (3.23)

Table 84
Mean Scores and Standard Deviations for FKK Internal Locus of Control Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>FKK Internal Locus of Control Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	33.88 (7.98)	33.94 (5.64)	34.88 (5.37)
Low APP	35.33 (9.50)	36.00 (6.56)	32.67 (2.52)
Medium APP	34.60 (6.43)	35.00 (6.60)	31.00 (8.25)
High APP	29.33 (2.52)	34.33 (5.13)	36.00 (6.00)

Table 85
Mean Scores and Standard Deviations for FKK Self Concept Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>FKK Self Concept Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	33.06 (5.08)	32.75 (3.84)	32.88 (3.74)
Low APP	32.00 (2.65)	33.00 (1.00)	30.33 (4.73)
Medium APP	33.80 (6.42)	30.60 (2.88)	30.00 (5.00)
High APP	34.67 (5.86)	30.00 (7.81)	35.33 (10.02)

Table 86
Mean Scores and Standard Deviations for FKK Self Efficacy Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>FKK Self Efficacy Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	66.94 (11.11)	66.69 (7.11)	67.75 (6.90)
Low APP	67.33 (10.26)	69.00 (7.55)	63.00 (5.57)
Medium APP	68.40 (10.69)	65.60 (3.78)	61.00 (6.21)
High APP	64.00 (6.25)	64.33 (11.93)	71.33 (16.01)

Table 87
Mean Scores and Standard Deviations for FKK Powerful Others Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>FKK Powerful Others Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	26.69 (9.19)	26.88 (7.00)	25.94 (7.23)
Low APP	28.33 (3.79)	21.67 (3.79)	25.67 (9.87)
Medium APP	26.20 (8.84)	21.80 (8.32)	22.60 (9.21)
High APP	26.00 (8.89)	29.00 (10.00)	26.00 (5.29)

Table 88
Mean Scores and Standard Deviations for FKK Chance Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>FKK Chance Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	24.69 (7.79)	25.63 (7.60)	24.19 (5.94)
Low APP	29.00 (10.00)	20.33 (1.53)	17.67 (2.08)
Medium APP	26.20 (10.71)	18.00 (7.25)	22.20 (6.38)
High APP	25.33 (8.51)	23.33 (7.10)	19.00 (5.20)

Table 89
Mean Scores and Standard Deviations for FKK External Locus of Control Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>FKK External Locus of Control Scores</u>			
<u>APP Severity Level</u>	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	51.38 (16.47)	52.50 (13.29)	50.13 (12.12)
Low APP	57.33 (3.58)	42.00 (5.29)	43.33 (9.71)
Medium APP	52.40 (18.15)	39.80 (15.43)	44.80 (14.72)
High APP	51.33 (9.07)	52.33 (14.15)	45.00 (1.00)

Table 90
Mean Scores and Standard Deviations for IPSAQ Internal Positive Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>IPSAQ Internal Positive Scores</u>			
<u>APP Severity Level</u>	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	8.50 (2.83)	7.37 (3.29)	7.00 (3.89)
Low APP	8.00 (4.36)	11.33 (1.53)	12.00 (1.00)
Medium APP	8.00 (1.41)	9.50 (1.92)	8.00 (.82)
High APP	6.50 (.71)	9.50 (.71)	8.00 (1.41)

Table 91
Mean Scores and Standard Deviations for IPSAQ Personal Positive Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>IPSAQ Personal Positive Scores</u>			
<u>APP Severity Level</u>	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	4.87 (2.36)	3.75 (1.49)	2.75 (2.82)
Low APP	2.00 (1.73)	4.00 (2.00)	2.00 (.00)
Medium APP	4.50 (1.73)	3.00 (1.41)	3.00 (2.45)
High APP	4.00 (.00)	3.50 (.71)	3.50 (.71)

Table 92
Mean Scores and Standard Deviations for IPSAQ Situational Positive Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>IPSAQ Situational Positive Scores</u>			
<u>APP Severity Level</u>	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	2.63 (1.51)	4.88 (2.42)	5.38 (2.62)
Low APP	3.67 (1.53)	.67 (.58)	2.00 (1.00)
Medium APP	2.00 (1.63)	3.50 (2.08)	5.00 (2.58)
High APP	5.50 (.71)	3.00 (.00)	4.50 (2.12)

Table 93
Mean Scores and Standard Deviations for IPSAQ Internal Negative Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>IPSAQ Internal Negative Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>APP Severity Level</u>	M (SD)	M (SD)	M (SD)
No APP	5.63 (4.21)	6.50 (4.72)	4.00 (3.38)
Low APP	5.00 (1.00)	3.33 (4.16)	9.33 (4.73)
Medium APP	7.25 (2.63)	8.00 (.82)	7.25 (.50)
High APP	6.00 (2.83)	9.50 (.71)	8.50 (4.95)

Table 94
Mean Scores and Standard Deviations for IPSAQ Personal Negative Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>IPSAQ Personal Negative Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>APP Severity Level</u>	M (SD)	M (SD)	M (SD)
No APP	6.13 (4.19)	5.13 (3.68)	4.87 (3.60)
Low APP	4.00 (2.65)	5.00 (1.73)	2.67 (2.08)
Medium APP	4.00 (4.08)	4.00 (2.58)	5.00 (3.16)
High APP	5.00 (2.83)	5.00 (1.41)	5.00 (4.24)

Table 95
Mean Scores and Standard Deviations for IPSAQ Situational Negative Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>IPSAQ Situational Negative Scores</u>			
<u>APP Severity Level</u>	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	4.25 (2.12)	4.25 (2.05)	6.25 (1.83)
Low APP	4.00 (4.58)	7.33 (5.51)	4.00 (5.29)
Medium APP	3.75 (1.89)	4.00 (2.94)	3.75 (2.75)
High APP	5.00 (.00)	1.50 (.71)	2.50 (.71)

Table 96
Mean Scores and Standard Deviations for CST Social Support Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>CST Social Support Scores</u>			
<u>APP Severity Level</u>	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	32.75 (10.11)	37.17 (9.16)	40.67 (8.33)
Low APP	39.67 (11.24)	36.67 (3.51)	43.00 (7.21)
Medium APP	36.25 (15.90)	35.75 (16.22)	39.75 (13.15)
High APP	35.67 (5.51)	38.33 (5.13)	42.33 (12.70)

Table 97
Mean Scores and Standard Deviations for CST Self Controlling Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>CST Self Controlling Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	54.50 (5.14)	54.08 (7.10)	54.75 (4.83)
Low APP	49.33 (4.04)	46.33 (5.13)	51.67 (.58)
Medium APP	48.75 (5.68)	51.25 (14.10)	51.25 (9.61)
High APP	48.67 (8.15)	45.33 (4.04)	36.33 (13.58)

Table 98
Mean Scores and Standard Deviations for CST Escape Avoidance Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>CST Escape Avoidance Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	33.67 (5.28)	36.25 (6.08)	38.17 (3.74)
Low APP	34.67 (2.52)	37.33 (.58)	37.00 (3.00)
Medium APP	36.25 (7.14)	41.50 (5.97)	40.75 (5.32)
High APP	35.67 (7.10)	39.67 (5.03)	25.00 (19.93)

Table 99
Mean Scores and Standard Deviations for CST Planful Problem Solving Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity Level</u>	<u>CST Planful Problem Solving Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No APP	42.08 (8.85)	43.33 (5.03)	45.17 (6.24)
Low APP	43.00 (3.61)	43.67 (2.08)	41.33 (2.08)
Medium APP	40.25 (15.13)	33.75 (8.26)	40.00 (9.27)
High APP	41.33 (13.65)	41.67 (8.62)	41.67 (19.66)

Table 100
Service Use by IS Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>			
	IS Relabel	IS Awareness	IS Need for Treatment	IS Total Insight
	F (2,114)	F (2,114)	F (2, 114)	F (2, 114)
Main Effect				
Assessment Time	2.40	1.37	3.22*	2.37
Main Effect				
Service Use	.52	8.42**	2.77	5.47*
Interaction				
Assessment Time * Service Use	3.98*	2.52	1.96	3.26*

*p<.05, **p<.01, ***p<.001

Table 101
Mean Scores and Standard Deviations for IS Relabel Scale Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IS Relabel Scale Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	2.62 (1.08)	2.76 (.95)	1.97 (1.21)
Service Use	2.13 (1.07)	2.43 (.94)	2.43 (.77)

Table 102
Mean Scores and Standard Deviations for IS Awareness of Illness Scale Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IS Awareness of Illness Scale Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	2.66 (1.17)	2.83 (1.20)	2.21 (1.11)
Service Use	2.00 (1.30)	1.70 (1.29)	1.97 (1.22)

Table 103
Mean Scores and Standard Deviations for IS Need for Treatment Scale Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>IS Need for Treatment Scale Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Services Before Age 18</u>	M (SD)	M (SD)	M (SD)
No Service Use	2.52 (1.08)	2.47 (1.21)	1.83 (1.22)
Service Use	2.15 (1.18)	1.72 (1.32)	1.85 (1.22)

Table 104
Mean Scores and Standard Deviations for IS Total Insight Scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Services Before Age 18</u>	<u>IS Total Insight Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	7.79 (2.96)	8.05 (2.87)	6.00 (2.75)
Service Use	6.35 (2.87)	5.92 (2.88)	6.25 (2.48)

Table 105
APP Severity by IS Repeated Measures ANOVAs

<u>Variable</u>	<u>Measures</u>			
	IS Relabel F (6,110)	IS Awareness F (6,110)	IS Need for Treatment F (6, 110)	IS Total Insight F (6, 110)
Main Effect Assessment Time	.91	2.16	3.14*	1.98
Main Effect Service Use	.43	2.89*	.91	1.88
Interaction Assessment Time * Service Use	2.90**	2.48*	1.79	3.38**

*p<.05, **p<.01, ***p<.001

Table 106
Mean Scores and Standard Deviations for IS Relabel Scale Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity</u>	<u>IS Relabel Scale Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	2.62 (1.08)	2.76 (.95)	1.97 (1.21)
Low APP	3.25 (.96)	2.50 (1.00)	2.00 (.00)
Medium APP	1.63 (.92)	2.75 (.89)	2.63 (.52)
High APP	2.11 (1.02)	2.28 (.96)	2.44 (.92)

Table 107
Mean Scores and Standard Deviations for IS Awareness of Illness Scale Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity</u>	<u>IS Awareness of Illness Scale Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	2.66 (1.17)	2.83 (1.20)	2.21 (1.11)
Low APP	3.25 (.96)	1.50 (1.00)	1.50 (1.00)
Medium APP	1.50 (1.41)	2.13 (1.13)	1.75 (1.17)
High APP	2.28 (1.18)	1.56 (1.42)	2.17 (1.30)

Table 108
Mean Scores and Standard Deviations for IS Need for Treatment Scale Scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity</u>	<u>IS Need for Treatment Scale Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	2.52 (1.08)	2.47 (1.21)	1.83 (1.22)
Low APP	3.13 (1.03)	1.50 (1.00)	1.38 (1.11)
Medium APP	1.69 (.88)	2.00 (1.51)	1.94 (1.08)
High APP	2.14 (1.23)	1.64 (1.34)	1.92 (1.33)

Table 109
Mean Scores and Standard Deviations for IS Total Insight score at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>APP Severity</u>	<u>IS Total Insight Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	7.79 (2.96)	8.05 (2.87)	6.00 (2.75)
Low APP	9.63 (2.75)	5.50 (1.00)	4.88 (1.65)
Medium APP	4.81 (1.81)	7.13 (3.40)	6.31 (2.19)
High APP	6.31 (2.77)	5.47 (2.88)	6.53 (2.74)

Table 110
Service Use by NOSIE Subscales and Total Assets Repeated Measures ANOVAs

<u>Variable</u>	<u>NOSIE Subscales</u>						<u>NOSIE Total Assets</u>
	<u>Daily Schedule Competence</u>	<u>Social Interest</u>	<u>Neatness</u>	<u>Irritability</u>	<u>Psychoticism</u>	<u>Motor Retardation</u>	F (2, 188)
Time	26.86***	39.51***	20.89***	3.78*	7.47***	15.93***	21.68***
Service Use	2.54	.06	1.20	1.38	.87	1.50	1.97
Time * Service Use	.57	1.30	.33	.15	.96	.34	.21

*p<.05, **p<.01, ***p<.001

Table 111
*Means and Standard Deviations on NOSIE Subscales and Total Assets at Admission, 6
 Months, and 12 Months as a Function of Service Use Before Age 18*

<u>NOSIE Subscales</u>	<u>Admission</u>		<u>6 Months</u>		<u>12 Months</u>	
	<u>M</u>	<u>SD</u>	<u>M</u>	<u>SD</u>	<u>M</u>	<u>SD</u>
<u>Daily Schedule Competence</u>						
No Service	32.84	6.96	34.27	5.98	36.68	3.57
Service Use	31.44	6.76	31.89	7.01	35.34	5.52
<u>Social Interest</u>						
No Service	15.01	5.58	18.95	5.96	19.17	5.38
Service Use	15.14	6.65	18.41	6.44	20.40	6.56
<u>Neatness</u>						
No Service	23.05	5.41	24.00	5.41	25.79	4.25
Service Use	22.34	5.85	22.62	5.62	24.72	4.91
<u>Irritability</u>						
No Service	4.37	5.44	5.89	6.14	4.59	4.60
Service Use	5.85	6.59	7.00	6.38	5.43	5.28
<u>Psychoticism</u>						
No Service	1.55	2.90	2.21	3.56	2.03	2.96
Service Use	1.73	2.68	3.13	4.26	2.70	3.96
<u>Motor Retardation</u>						
No Service	6.43	4.10	6.00	3.63	4.18	2.92
Service Use	7.28	4.60	6.54	4.76	5.35	3.88
<u>NOSIE Total Assets</u>						
No Service	154.54	23.46	159.12	22.60	167.02	16.75
Service Use	148.47	25.70	152.21	26.50	162.71	23.39

Table 112
APP Severity Level by NOSIE Subscales and Total Assets Repeated Measures ANOVAs

<u>Variable</u>	<u>NOSIE Subscales</u>						<u>NOSIE</u>
	<u>Daily</u> <u>Schedule</u> <u>Competence</u>	<u>Social</u> <u>Interest</u>	<u>Neatness</u>	<u>Irritability</u>	<u>Psychoticism</u>	<u>Motor</u> <u>Retardation</u>	<u>Total</u> <u>Assets</u>
	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 184)
Time	23.23***	36.58***	19.85***	3.40*	8.07***	13.55***	22.23***
APP Severity Level	1.10	.73	.73	1.30	.36	1.63	1.48
Time * APP	.85	1.83	3.14**	.40	1.58	2.03	2.20*

*p<.05, **p<.01, ***p<.001

Table 113
Means and Standard Deviations on NOSIE Subscales and Total Assets at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

<u>NOSIE Subscales</u>	<u>Admission</u>		<u>6 Months</u>		<u>12 Months</u>	
	<u>M</u>	<u>SD</u>	<u>M</u>	<u>SD</u>	<u>M</u>	<u>SD</u>
<u>Daily Schedule Competence</u>						
No APP	32.84	6.96	34.27	5.98	36.68	3.57
Low APP	30.24	8.73	31.04	7.32	34.23	6.99
Med APP	30.83	6.36	33.07	6.36	36.27	3.50
High APP	32.75	5.50	31.60	7.48	35.42	5.74
<u>Social Interest</u>						
No APP	15.01	5.58	18.95	5.96	19.17	5.38
Low APP	13.16	6.20	16.11	5.45	19.64	7.09
Med APP	15.03	7.02	21.09	4.35	20.90	4.50
High APP	16.60	6.62	18.00	7.79	20.56	7.65
<u>Neatness</u>						
No APP	23.05	5.41	24.00	5.41	25.79	4.25
Low APP	20.42	6.71	22.06	5.68	24.36	5.47
Med APP	21.01	5.48	23.24	4.91	25.26	3.54
High APP	24.69	4.87	22.54	6.27	24.57	5.57
<u>Irritability</u>						
No APP	4.37	5.44	5.89	6.14	4.59	4.60
Low APP	6.75	7.46	7.90	8.64	5.30	4.70
Med APP	4.31	5.36	5.75	6.07	3.38	4.30
High APP	6.38	6.90	7.31	4.76	7.05	5.97
<u>Psychoticism</u>						
No APP	1.55	2.90	2.21	3.56	2.03	2.96
Low APP	1.74	2.58	3.83	5.50	2.39	4.32
Med APP	2.06	3.65	2.47	4.02	2.08	3.20
High APP	1.48	1.90	3.13	3.53	3.37	4.30
<u>Motor Retardation</u>						
No APP	6.43	4.10	6.00	3.63	4.18	2.92
Low APP	8.91	5.61	7.96	4.81	6.23	4.65
Med APP	8.39	4.70	5.56	3.83	4.93	2.65
High APP	5.31	2.92	6.28	5.31	5.05	4.16
<u>NOSIE Total Assets</u>						
No APP	154.54	23.46	159.12	22.60	167.02	16.75
Low APP	136.62	30.08	145.59	29.43	160.29	26.73
Med APP	148.09	23.25	159.65	18.94	168.03	12.23
High APP	157.05	21.68	151.26	28.98	160.42	27.39

Table 114
Service Use by BPRS Factors and BPRS Total Repeated Measures ANOVAs

<u>Variable</u>	<u>BPRS Factors</u>						<u>BPRS Total</u>
	<u>BPRS Psychotic Disorganization</u>	<u>BPRS Hallucination/ Delusions</u>	<u>BPRS Paranoia</u>	<u>BPRS Emotional Blunting</u>	<u>BPRS Anxiety/ Depression</u>	<u>BPRS Agitation / Elation</u>	
	F (2, 128)	F (2, 134)	F (2, 134)	F (2, 134)	F (2, 134)	F (2, 134)	F (2, 128)
Time	1.16	.82	.04	1.03	.50	.06	1.23
Service Use	1.20	.47	1.75	1.20	.64	.02	.23
Time * Service Use	3.17*	.85	.05	.19	.27	1.08	.86

*p<.05, **p<.01, ***p<.001

Table 115
APP Severity Level by BPRS Factors and BPRS Total Repeated Measures ANOVAs

<u>Variable</u>	<u>BPRS Factors</u>						<u>BPRS Total</u>
	<u>BPRS Psychotic Disorganization</u>	<u>BPRS Hallucination/ Delusions</u>	<u>BPRS Paranoia</u>	<u>BPRS Emotional Blunting</u>	<u>BPRS Anxiety/ Depression</u>	<u>BPRS Agitation / Elation</u>	
	F (6, 124)	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 184)	F (6, 124)
Time	.88	.52	.06	.75	.25	.07	.71
APP Severity Level	.81	.64	.79	1.53	2.22	.24	.50
Time * APP Severity	1.71	.79	.46	.45	.32	.67	.57

*p<.05, **p<.01, ***p<.001

Table 116
*Mean Scores and Standard Deviations for BPRS Total scores at Admission, 6 Months,
 and 12 Months as a Function of Service Use Before Age 18*

<u>Service Use</u>	<u>BPRS Total Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	45.75 (10.47)	46.34 (13.75)	45.32 (14.75)
Service Use	49.53 (16.32)	47.13 (12.52)	44.64 (13.22)

Table 117
*Mean Scores and Standard Deviations for BPRS Psychotic Disorganization Factor
 scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age
 18*

<u>Service Use</u>	<u>BPRS Psychotic Disorganization Factor Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	5.81 (2.13)	6.31 (2.69)	6.13 (2.91)
Service Use	7.45 (3.82)	6.67 (2.80)	6.12 (2.76)

Table 118
Mean Scores and Standard Deviations for BPRS Hallucinations/Delusions Factor scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Service Use</u>	<u>BPRS Hallucination/Delusions Factor Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	6.83 (4.20)	6.99 (4.31)	6.80 (5.23)
Service Use	6.96 (4.01)	6.15 (3.87)	5.76 (3.64)

Table 119
Mean Scores and Standard Deviations for BPRS Paranoia Factor scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Service Use</u>	<u>BPRS Paranoia Factor Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	8.92 (2.86)	8.88 (3.56)	8.68 (3.40)
Service Use	8.01 (3.58)	8.04 (2.95)	8.04 (3.27)

Table 120
Mean Scores and Standard Deviations for BPRS Emotional Blunting Factor scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>Service Use</u>	<u>BPRS Emotional Blunting Factor Scores</u>		
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
	M (SD)	M (SD)	M (SD)
No Service Use	5.99 (3.64)	5.59 (2.27)	5.63 (2.53)
Service Use	6.77 (3.62)	6.52 (3.69)	6.12 (3.39)

Table 121
Mean Scores and Standard Deviations for BPRS Anxiety/Depression Factor scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>BPRS Anxiety/Depression Factor Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Service Use</u>	M (SD)	M (SD)	M (SD)
No Service Use	8.55 (2.81)	8.36 (3.24)	8.42 (3.32)
Service Use	9.41 (4.21)	9.04 (4.55)	8.65 (3.65)

Table 122
Mean Scores and Standard Deviations for BPRS Agitation/Elation Factor scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

<u>BPRS Agitation/Elation Factor Scores</u>			
	<u>At Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>Service Use</u>	M (SD)	M (SD)	M (SD)
No Service Use	3.31 (1.18)	3.63 (2.38)	3.41 (1.78)
Service Use	3.58 (1.66)	3.28 (1.19)	3.36 (1.45)

Table 123
Mean Scores and Standard Deviations for BPRS Total and Factor scores at Admission, 6 Months, and 12 Months as a Function of APP Severity Level

	<u>APP</u> <u>Severity</u> <u>Level</u>	<u>Admission</u> <u>M (SD)</u>	<u>Six Months</u> <u>M (SD)</u>	<u>Twelve Months</u> <u>M (SD)</u>
BPRS				
BPRS Total	No APP	45.75 (10.47)	46.34 (13.75)	45.32 (14.75)
	Low APP	49.19 (14.37)	47.81 (9.52)	48.13 (14.62)
	Med APP	42.67 (9.05)	43.42(10.22)	41.33 (9.37)
	High APP	51.52 (18.39)	47.90 (14.19)	44.27 (13.83)
BPRS Psychotic Disorganization Factor	No APP	5.81 (2.13)	6.31 (2.69)	6.13 (2.91)
	Low APP	8.00 (3.89)	6.80 (2.35)	6.94 (3.42)
	Med APP	5.52 (1.52)	5.98 (2.48)	5.92 (2.25)
	High APP	7.77 (4.17)	6.80 (3.10)	5.87 (2.69)
BPRS Hallucination/Delusion Factor	No APP	6.83 (4.20)	6.99 (4.31)	6.80 (5.23)
	Low APP	7.25 (3.88)	8.31 (5.30)	7.13 (5.61)
	Med APP	6.17 (3.13)	4.58 (2.54)	5.67 (2.80)
	High APP	7.07 (4.37)	5.80 (3.40)	5.30 (2.99)
BPRS Paranoia Factor	No APP	8.92 (2.86)	8.88 (3.56)	8.68 (3.40)
	Low APP	8.68 (4.63)	7.90 (2.64)	9.22 (3.43)
	Med APP	7.50 (2.21)	8.57 (2.36)	6.62 (1.95)
	High APP	7.92 (3.59)	7.95 (3.27)	7.99 (3.44)
BPRS Emotional Blunting Factor	No APP	5.99 (3.64)	5.59 (2.27)	5.63 (2.53)
	Low APP	7.75 (2.98)	6.75 (2.65)	6.72 (2.83)
	Med APP	4.75 (1.41)	4.08 (2.33)	5.00 (2.86)
	High APP	6.96 (4.09)	7.08 (4.10)	6.21 (3.73)
BPRS Anxiety/Depression Factor	No APP	8.55 (2.81)	8.36 (3.24)	8.42 (3.32)
	Low APP	6.63 (1.73)	6.78 (2.37)	7.00 (1.87)
	Med APP	9.25 (4.29)	9.67 (3.78)	8.67 (3.92)
	High APP	10.41 (4.47)	9.67 (5.15)	9.22 (3.98)
BPRS Agitation/Elation Factor	No APP	3.31 (1.18)	3.63 (2.38)	3.41 (1.78)
	Low APP	3.13 (1.06)	3.16 (1.03)	2.94 (.73)
	Med APP	3.33 (.93)	3.08 (.97)	3.83 (1.63)
	High APP	3.80 (1.95)	3.37 (1.32)	3.39 (1.60)

*p<.05

Table 124
Mean Scores and Standard Deviations for BPRS item scores at Admission, 6 Months, and 12 Months as a Function of Service Use Before Age 18

		<u>Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>BPRS Items</u>		<u>M (SD)</u>	<u>M (SD)</u>	<u>M (SD)</u>
Somatic Concern	No Service	2.78 (1.66)	2.58 (1.68)	2.59 (1.72)
	Service Use	2.39 (1.71)	2.68 (1.54)	2.29 (1.49)
Anxiety	No Service	2.39 (.97)	2.45 (1.24)	2.16 (1.25)
	Service Use	2.54 (1.51)	2.53 (1.35)	2.37 (1.38)
Depression	No Service	2.09 (1.11)	2.02 (1.16)	1.95 (1.07)
	Service Use	2.39 (1.51)	2.20 (1.41)	2.16 (1.16)
Suicidality	No Service	1.16 (.37)	1.22 (.51)	1.09 (.30)
	Service Use	1.63 (1.26)	1.57 (1.03)	1.45 (1.23)
Guilt	No Service	1.81 (1.38)	1.77 (1.18)	2.02 (1.49)
	Service Use	2.07 (1.29)	2.13 (1.51)	2.07 (1.47)
Hostility	No Service	2.44 (1.36)	2.52 (1.45)	2.64 (1.42)
	Service Use	2.71 (1.63)	2.91 (1.76)	2.61 (1.71)
Elevated Mood	No Service	1.44 (.91)	1.56 (1.30)	1.33 (.90)
	Service Use	1.54 (1.22)	1.25 (.68)	1.45 (.98)
Grandiosity	No Service	2.14 (1.78)	2.14 (1.94)	2.39 (2.14)
	Service Use	1.66 (1.40)	1.76 (1.78)	1.63 (1.49)
Suspiciousness	No Service	3.03 (1.85)	2.97 (2.06)	2.53 (1.81)
	Service Use	2.28 (1.73)	2.21 (1.46)	2.17 (1.45)
Hallucinations	No Service	1.72 (1.46)	2.17 (1.85)	2.23 (1.84)
	Service Use	2.72 (2.10)	2.13 (1.68)	2.04 (1.60)
Unusual Thought Content	No Service	2.97 (2.10)	2.68 (1.66)	2.17 (1.91)
	Service Use	2.57 (1.99)	2.26 (1.84)	2.14 (1.69)
Bizarre Behavior	No Service	1.59 (1.34)	2.08 (1.31)	1.80 (1.32)
	Service Use	2.54 (1.76)	2.17 (1.45)	1.95 (1.55)
Self-Neglect	No Service	1.97 (.83)	2.56 (.93)	2.20 (1.05)
	Service Use	2.46 (1.11)	2.46 (.90)	1.95 (.96)
Disorientation	No Service	1.41 (.76)	1.41 (.76)	1.50 (1.02)
	Service Use	1.54 (1.07)	1.57 (1.13)	1.47 (1.12)
Conceptual Disorganization	No Service	1.89 (1.10)	1.86 (1.26)	1.91 (1.21)
	Service Use	2.14 (1.65)	1.89 (1.16)	1.69 (1.15)
Blunted Affect	No Service	2.50 (1.50)	2.56 (1.12)	2.64 (1.07)
	Service Use	2.78 (1.52)	2.97 (1.69)	2.82 (1.47)
Emotional Withdrawal	No Service	2.19 (1.40)	2.12 (1.30)	2.12 (1.18)
	Service Use	2.42 (1.30)	2.22 (1.38)	2.11 (1.18)
Motor Retardation	No Service	2.05 (1.19)	1.67 (.99)	1.67 (.88)
	Service Use	2.20 (1.27)	1.93 (1.16)	1.82 (1.06)
Tension	No Service	1.67 (.91)	1.52 (.91)	1.75 (1.04)
	Service Use	1.55 (.82)	1.51 (.87)	1.30 (.78)
Uncooperativeness	No Service	1.70 (1.09)	1.72 (1.14)	1.64 (1.30)
	Service Use	1.50 (.99)	1.49 (.80)	1.85 (1.16)
Excitement	No Service	1.48 (.82)	1.53 (1.22)	1.63 (1.19)
	Service Use	1.27 (.73)	1.20 (.60)	1.27 (.89)
Distractibility	No Service	1.72 (.92)	1.59 (1.16)	1.88 (1.28)
	Service Use	1.99 (1.42)	1.76 (1.23)	1.70 (1.10)
Motor hyperactivity	No Service	1.30 (.49)	1.45 (1.10)	1.53 (1.09)
	Service Use	1.20 (.64)	1.23 (.63)	1.18 (.65)
Mannerisms and Posturing	No Service	1.30 (.60)	1.07 (.37)	1.27 (.73)
	Service Use	1.26 (.65)	1.14 (.54)	1.31 (.94)

Table 125
*Mean Scores and Standard Deviations for BPRS item scores at Admission, 6 Months, and
 12 Months as a Function of APP Severity Level*

		<u>Admission</u>	<u>Six Months</u>	<u>Twelve Months</u>
<u>BPRS Items</u>	<u>APP Severity Level</u>	<u>M (SD)</u>	<u>M (SD)</u>	<u>M (SD)</u>
Somatic Concern	No APP	2.78 (1.66)	2.58 (1.68)	2.59 (1.72)
	Low APP	1.63 (.92)	2.56 (1.64)	2.62 (1.62)
	Med APP	2.67 (2.09)	3.33 (1.63)	1.42 (.80)
	High APP	2.58 (1.79)	2.56 (1.51)	2.40 (1.54)
Anxiety	No APP	2.39 (.97)	2.45 (1.24)	2.16 (1.25)
	Low APP	1.63 (.74)	2.06 (1.27)	2.31 (.88)
	Med APP	2.33 (1.97)	2.67 (.98)	2.50 (1.76)
	High APP	2.90 (1.49)	2.66 (1.47)	2.35 (1.47)
Depression	No APP	2.09 (1.11)	2.02 (1.16)	1.95 (1.07)
	Low APP	1.50 (.54)	1.50 (1.07)	1.63 (.74)
	Med APP	2.58 (1.56)	2.25 (1.41)	2.42 (1.39)
	High APP	2.65 (1.64)	2.42 (1.49)	2.27 (1.21)
Suicidality	No APP	1.16 (.37)	1.22 (.51)	1.09 (.30)
	Low APP	1.00 (.00)	1.19 (.53)	1.13 (.35)
	Med APP	1.83 (.98)	1.33 (.52)	1.00 (.00)
	High APP	1.79 (1.47)	1.75 (1.21)	1.67 (1.50)
Guilt	No APP	1.81 (1.38)	1.77 (1.18)	2.02 (1.49)
	Low APP	1.25 (.46)	1.13 (.35)	1.25 (.54)
	Med APP	1.75 (.99)	2.58 (1.74)	2.08 (1.11)
	High APP	2.42 (1.41)	2.35 (1.59)	2.33 (1.67)
Hostility	No APP	2.44 (1.36)	2.52 (1.45)	2.64 (1.42)
	Low APP	2.38 (1.51)	2.06 (1.43)	3.00 (2.25)
	Med APP	2.67 (1.37)	3.17 (1.72)	1.92 (1.20)
	High APP	2.83 (1.76)	3.13 (1.85)	2.65 (1.63)
Elevated Mood	No APP	1.44 (.91)	1.56 (1.30)	1.33 (.90)
	Low APP	1.25 (.46)	1.13 (.35)	1.25 (.71)
	Med APP	1.42 (.67)	1.25 (.42)	2.33 (1.63)
	High APP	1.67 (1.47)	1.29 (.81)	1.29 (.75)
Grandiosity	No APP	2.14 (1.78)	2.14 (1.94)	2.39 (2.14)
	Low APP	1.88 (1.73)	3.75 (2.66)	2.25 (2.32)
	Med APP	1.67 (1.63)	1.00 (.00)	1.67 (1.63)
	High APP	1.58 (1.27)	1.29 (1.08)	1.42 (1.09)
Suspiciousness	No APP	3.03 (1.85)	2.97 (2.06)	2.53 (1.81)
	Low APP	2.63 (2.07)	2.81 (2.45)	2.94 (1.78)
	Med APP	1.67 (.82)	2.42 (.97)	1.75 (1.41)
	High APP	2.31 (1.79)	1.96 (1.08)	2.02 (1.31)
Hallucinations	No APP	1.72 (1.46)	2.17 (1.85)	2.23 (1.84)
	Low APP	2.13 (2.10)	1.63 (1.41)	2.13 (1.81)
	Med APP	2.67 (2.25)	2.00 (1.27)	2.17 (2.04)
	High APP	2.94 (2.11)	2.33 (1.86)	1.98 (1.49)
Unusual Thought Content	No APP	2.97 (2.10)	2.68 (1.66)	2.17 (1.91)
	Low APP	3.25 (2.19)	2.94 (2.28)	2.75 (2.14)
	Med APP	1.83 (1.60)	1.58 (1.43)	1.83 (1.33)
	High APP	2.52 (2.01)	2.20 (1.76)	2.00 (1.62)
Bizarre Behavior	No APP	1.59 (1.34)	2.08 (1.31)	1.80 (1.32)
	Low APP	2.81 (1.89)	2.13 (1.53)	2.56 (1.99)
	Med APP	1.33 (.52)	1.92 (1.20)	1.83 (1.60)
	High APP	2.75 (1.84)	2.25 (1.53)	1.77 (1.40)
Self-Neglect	No APP	1.97 (.83)	2.56 (.93)	2.20 (1.05)
	Low APP	2.38 (1.06)	2.44 (.62)	2.25 (.93)
	Med APP	2.67 (.52)	2.42 (.92)	2.67 (1.03)
	High APP	2.44 (1.25)	2.48 (.99)	1.67 (.86)
Disorientation	No APP	1.41 (.76)	1.41 (.76)	1.50 (1.02)
	Low APP	2.63 (1.69)	2.56 (1.80)	2.19 (1.69)
	Med APP	1.00 (.00)	1.33 (.82)	1.17 (.41)
	High APP	1.30 (.64)	1.28 (.65)	1.30 (.93)

Conceptual	No APP	1.89 (1.10)	1.86 (1.26)	1.91 (1.21)
Disorganization	Low APP	2.75 (1.75)	2.38 (1.41)	2.13 (1.46)
	Med APP	1.50 (1.23)	1.33 (.82)	1.50 (1.23)
	High APP	2.09 (1.70)	1.87 (1.12)	1.59 (1.02)
Blunted Affect	No APP	2.50 (1.50)	2.56 (1.12)	2.64 (1.07)
	Low APP	3.25 (1.49)	2.94 (1.61)	3.25 (1.51)
	Med APP	1.67 (.82)	2.00 (1.10)	1.92 (1.11)
Emotional	High APP	2.91 (1.58)	3.24 (1.80)	2.91 (1.50)
	No APP	2.19 (1.40)	2.12 (1.30)	2.12 (1.18)
	Low APP	2.88 (1.13)	2.44 (.94)	2.38 (1.06)
Withdrawal	Med APP	1.83 (1.17)	1.50 (1.23)	2.08 (1.02)
	High APP	2.41 (1.37)	2.33 (1.53)	2.02 (1.28)
	No APP	2.05 (1.19)	1.67 (.94)	1.67 (.88)
Motor Retardation	Low APP	2.25 (1.04)	1.88 (.84)	1.75 (.89)
	Med APP	1.75 (.61)	1.17 (.41)	1.58 (.92)
	High APP	2.30 (1.47)	2.15 (1.31)	1.91 (1.17)
Tension	No APP	1.67 (.91)	1.52 (.91)	1.75 (1.04)
	Low APP	1.75 (1.04)	1.50 (1.07)	1.25 (.46)
	Med APP	1.67 (.82)	1.50 (.55)	1.17 (.41)
Uncooperativeness	High APP	1.46 (.75)	1.52 (.90)	1.35 (.94)
	No APP	1.70 (1.09)	1.72 (1.14)	1.64 (1.30)
	Low APP	2.00 (1.31)	1.63 (.52)	1.88 (1.13)
Excitement	Med APP	1.67 (1.03)	1.50 (.55)	1.67 (1.03)
	High APP	1.28 (.81)	1.43 (.95)	1.89 (1.24)
	No APP	1.48 (.82)	1.53 (1.22)	1.63 (1.19)
Distractibility	Low APP	1.25 (.46)	1.00 (.00)	1.31 (.70)
	Med APP	1.00 (.00)	1.17 (.41)	1.17 (.41)
	High APP	1.35 (.89)	1.28 (.72)	1.28 (1.05)
Motor hyperactivity	No APP	1.72 (.92)	1.59 (1.16)	1.88 (1.28)
	Low APP	1.88 (.84)	1.63 (.74)	1.81 (1.00)
	Med APP	1.50 (.84)	1.83 (1.60)	1.50 (.84)
Mannerisms and Posturing	High APP	2.15 (1.68)	1.78 (1.30)	1.72 (1.20)
	No APP	1.30 (.49)	1.45 (1.10)	1.53 (1.09)
	Low APP	1.38 (.74)	1.44 (1.05)	1.13 (.35)
Mannerisms and Posturing	Med APP	1.00 (.00)	1.17 (.41)	1.00 (.00)
	High APP	1.20 (.69)	1.17 (.49)	1.24 (.80)
	No APP	1.30 (.60)	1.07 (.37)	1.27 (.73)
Mannerisms and Posturing	Low APP	1.50 (.76)	1.13 (.35)	1.00 (.00)
	Med APP	1.00 (.00)	1.00 (.00)	1.00 (.00)
	High APP	1.25 (.69)	1.18 (.66)	1.50 (1.18)

*p<.05

Table 126
*Relationship Between Level of Discharge Location Restrictiveness
 and Service Use Before Age 18 (N=242)*

<u>Discharge Location Restrictiveness</u>	<u>Service Use</u>		<u>Total</u>
	<u>No Service Use</u>	<u>Service Use</u>	
1 ó Same or Higher Restrictiveness (LRC Transfer)	13	13	26
2 - Psychiatric Residential Rehabilitation	42	55	97
3 - Assisted Living	33	48	81
4 ó Independent Living/Living with Family	20	18	38
Total	108	134	242

Table 127
*Relationship Between Level of Discharge Location Restrictiveness
 and APP Severity Level (N=242)*

<u>Discharge Location Restrictiveness</u>	<u>No</u>	<u>APP Severity Level</u>			<u>Total</u>
		<u>Low</u>	<u>Med</u>	<u>High</u>	
1 ó Same or Higher Restrictiveness (LRC Transfer)	13	1	1	11	26
2 - Psychiatric Residential Rehabilitation	42	8	15	32	97
3 - Assisted Living	33	8	10	30	81
4 ó Independent Living/Living with Family	20	2	3	13	38
Total	108	19	29	86	242