

11-7-2015

# Reintroduction of Atlantic salmon in Lake Ontario: the Implications of Genetic Quality on Individual Fitness

Chantal Audet  
*University of Windsor*

Follow this and additional works at: <https://scholar.uwindsor.ca/etd>

---

## Recommended Citation

Audet, Chantal, "Reintroduction of Atlantic salmon in Lake Ontario: the Implications of Genetic Quality on Individual Fitness" (2015). *Electronic Theses and Dissertations*. 5519.  
<https://scholar.uwindsor.ca/etd/5519>

This online database contains the full-text of PhD dissertations and Masters' theses of University of Windsor students from 1954 forward. These documents are made available for personal study and research purposes only, in accordance with the Canadian Copyright Act and the Creative Commons license—CC BY-NC-ND (Attribution, Non-Commercial, No Derivative Works). Under this license, works must always be attributed to the copyright holder (original author), cannot be used for any commercial purposes, and may not be altered. Any other use would require the permission of the copyright holder. Students may inquire about withdrawing their dissertation and/or thesis from this database. For additional inquiries, please contact the repository administrator via email ([scholarship@uwindsor.ca](mailto:scholarship@uwindsor.ca)) or by telephone at 519-253-3000ext. 3208.

**Reintroduction of Atlantic salmon in Lake Ontario: the Implications of Genetic  
Quality on Individual Fitness**

By Chantal Lianne Audet

A Thesis

Submitted to the Faculty of Graduate Studies  
through Biological Sciences in Partial Fulfillment of the Requirements for  
the Degree of Master of Science at the  
University of Windsor

Windsor, Ontario, Canada

2015

©2015 Chantal Lianne Audet

Reintroduction of Atlantic salmon in Lake Ontario: the Implications of Genetic Quality  
on Individual Fitness

by

Chantal L Audet

**APPROVED BY:**

---

A Fisk

Great Lakes Institute for Environmental Research

---

D Higgs

Department of Biological Sciences

---

T Pitcher, Advisor

Department of Biological Sciences

17 August 2015

## **DECLARATION OF CO-AUTHORSHIP**

I hereby declare that this thesis incorporates material that is the result of joint research, as follows: my first and second data chapters were co-authored with my supervisor, Dr. Trevor Pitcher, and Dr. Chris Wilson. In each case, my collaborators provided valuable feedback, helped with the project design and statistical analysis, and provided editorial input during the writing of each manuscript; however, in both cases the primary contributions have all been made by the author. Chapter 2 has been prepared as a manuscript that will be submitted to Ecology of Freshwater Fish. Chapter 3 is in the process of being made into a manuscript that will be submitted to Reproductive Biology, respectively.

I am aware of the University of Windsor Senate Policy on Authorship and I certify that I have properly acknowledged the contribution of other researchers to my thesis, and have obtained written permission from my co-authors to include the above materials in my thesis.

I certify that, with the above qualification, this thesis, and the research to which it refers, is the product of my own work, completed during my registration as a graduate student at the University of Windsor.

I declare that, to the best of my knowledge, my thesis does not infringe upon anyone's copyright nor violate any proprietary rights and that any ideas, techniques, quotations, or any other material from the work of other people including in my thesis, published or otherwise, are fully acknowledged in accordance with the standard referencing practices. Furthermore, to the extent that I have included copyrighted material

that surpasses the bounds of fair dealing within the meaning of the Canada Copyright Act, I certify that I have obtained a written permission from the copyright owners to include such materials in my thesis.

I declare that this is a true copy of my thesis, including any final revisions, as approved by my thesis committee and the Graduate Studies office, and that this thesis has not been submitted for a higher degree in any other University of Institution.

## ABSTRACT

The genetic health of hatchery broodstock is important for the success of conservation management strategies. This thesis examined the genetic quality of the Atlantic salmon (*Salmo salar*) broodstock used in the Lake Ontario reintroduction effort by examining the potential for outbreeding depression in F<sub>1</sub> juvenile hybrids using a full-factorial breeding design. This study did not find evidence to support the occurrence of outbreeding depression in hybrids. This thesis also examined the relationship between genetic and gamete quality in a strain used in the Lake Ontario restoration effort. Although negative correlations were found between heterozygosity and sperm velocity and longevity, the correlations explained limited variance. Overall, there was little evidence to support the existence of correlations between gamete quality and heterozygosity in Atlantic salmon. This study is the first step into looking at potential implications of stocking multiple stains of Atlantic salmon and assessing the correlation between genetic and reproductive quality in their broodstock.

## ACKNOWLEDGEMENTS

I would like to thank my advisor, Dr. T. Pitcher for mentoring and helping me out with my research for the past two years. I would also like to thank my external collaborator, Dr. C. Wilson for his help and expertise that went into both my chapters. I would also like to give a special thank you to Bill Sloan, Scott Ferguson and Anne Kidd for providing technical expertise for my chapters. I would also like to thank Craig Black who started the project and Katelynn Johnson who helped make the crosses for my second Chapter. I would also like to thank Jessica Mayrand for helping with running the swimming trials. I also owe a special thank you to Jade Laycock for assisting me with both of my field seasons. I would also like to thank Chris Weaver, Brian Rosborough and the technicians at the Harwood Fish Culture Centre who were instrumental in providing me with the fish in order to accomplish my third Chapter. I would also like to thank my committee members, Dr. D. Higgs and Dr. A. Fisk for their feedback. Finally, I would also like to thank my lab colleagues Adriana Forest, Jason Lewis, Sarah Lehnert, Aleksa Zubic and Sumeet Bhardwaj for all of the assistance they provided me over these past two years.

## TABLE OF CONTENTS

|   |            |
|---|------------|
| <b>DECLARATION OF CO-AUTHORSHIP.....</b>  | <b>III</b> |
| <b>ABSTRACT.....</b>  | <b>V</b>   |
| <b>ACKNOWLEDGEMENTS .....</b>   | <b>VI</b>  |
| <b>LIST OF TABLES .....</b>   | <b>IX</b>  |
| <b>LIST OF FIGURES.....</b>   | <b>XII</b> |
| <b>LIST OF APPENDICES.....</b>  | <b>XIV</b> |
| <b>CHAPTER 1: GENERAL INTRODUCTION.....</b>   | <b>1</b>   |
| HYBRIDIZATION.....  | 1          |
| <i>OUTBREEDING DEPRESSION</i> .....   | 2          |
| <i>HETEROSIS</i> .....  | 3          |
| HETEROZYGOSITY FITNESS CORRELATIONS.....  | 5          |
| <i>GAMETE QUALITY TRAITS</i> .....  | 6          |
| ATLANTIC SALMON.....  | 7          |
| OVERVIEW OF THE THESIS .....  | 8          |
| REFERENCES.....   | 10         |
| <b>CHAPTER 2: EFFECTS OF INTRASPECIFIC HYBRIDIZATION BETWEEN TWO STRAINS OF ATLANTIC SALMON (<i>SALMO SALAR</i>) ON JUVENILE SURVIVAL AND FITNESS-RELATED TRAITS: IMPLICATIONS FOR RESTORATION ECOLOGY.....</b> | <b>18</b>  |
| SYNOPSIS .....  | 18         |
| INTRODUCTION.....   | 19         |
| METHODS.....  | 22         |
| <i>FULL FACTORIAL BREEDING DESIGN</i> .....   | 22         |
| <i>FERTILIZED EGG SURVIVAL</i> .....  | 22         |
| <i>REARING</i> .....  | 23         |
| <i>LENGTH, MASS AND FULTON'S CONDITION FACTOR</i> .....   | 24         |
| <i>SURVIVAL</i> .....   | 24         |
| <i>STATISTICAL ANALYSIS</i> .....   | 25         |
| RESULTS .....   | 26         |
| DISCUSSION.....   | 27         |
| ACKNOWLEDGEMENTS.....   | 34         |
| REFERENCES.....   | 35         |
| <b>CHAPTER 3: HETEROZYGOSITY AND GAMETE QUALITY TRAITS IN HATCHERY-REARED ATLANTIC SALMON (<i>SALMO SALAR</i>).....</b>   | <b>45</b>  |
| SYNOPSIS .....  | 45         |
| INTRODUCTION.....   | 46         |
| METHODS.....  | 50         |
| <i>BROODSTOCK</i> .....   | 50         |
| <i>MILT COLLECTION AND SPERM QUALITY ASSESSMENT</i> .....   | 50         |
| <i>EGG COLLECTION AND QUALITY ASSESSMENT</i> .....  | 53         |
| <i>HETEROZYGOSITY</i> .....   | 54         |
| <i>STATISTICAL ANALYSIS</i> .....   | 54         |



|  |            |
|--|------------|
| FEMALES.....                                   | 54         |
| MALES.....                                     | 55         |
| RESULTS.....                                   | 56         |
| FEMALES.....                                   | 56         |
| MALES.....                                     | 57         |
| DISCUSSION.....                                | 58         |
| ACKNOWLEDGEMENTS.....                          | 65         |
| REFERENCES.....                                | 66         |
| <b>CHAPTER 4: GENERAL DISCUSSION.....</b>      | <b>90</b>  |
| SUMMARY.....                                   | 90         |
| <i>CHAPTER TWO</i> .....                       | 90         |
| <i>CHANGES TO ENVIRONMENTAL SETTINGS</i> ..... | 91         |
| <i>F2, F3 AND BACKCROSSES</i> .....            | 92         |
| <i>LIFE HISTORY TRAITS</i> .....               | 93         |
| <i>CHAPTER THREE</i> .....                     | 94         |
| CONCLUSION.....                                | 95         |
| REFERENCES.....                                | 97         |
| APPENDICES.....                                | 101        |
| <b>VITA AUCTORIS.....</b>                      | <b>129</b> |

## LIST OF TABLES

|  |    |
|--|----|
| <b>Table 1.1</b> Approved populations of Atlantic salmon for Lake Ontario. The table represents a comparison between the known conditions of the current approved candidate populations and the speculated characteristics of the Lake Ontario population.....   | 17 |
| <b>Table 2.1</b> Mean percentage of survival of F <sub>1</sub> cross types of Atlantic salmon ( <i>Salmo salar</i> ) during four life stages with the significance (p) of cross type (“Cross”) as fixed effect as well as dam, sire and their interaction as random effects to the variance of the fitness characteristic (see methods for details)..... | 42 |
| <b>Table 2.2</b> Means (± standard error) of F <sub>1</sub> cross types of Atlantic salmon ( <i>Salmo salar</i> ) with the significance (p) of cross type (“Cross”) as fixed effect as well as dam, sire and their interaction as random effects to the phenotypic variance of the fitness characteristic (see methods for details).....                 | 43 |
| <b>Table 3.1</b> Summary of existing literature evaluating the occurrence of HFCs of gamete quality related traits. MLH indicates multilocus heterozygosity, SH indicates standardized heterozygosity, HL indicates homozygosity by weighed locus.....   | 74 |
| <b>Table 3.2</b> Hardy Weinberg equilibrium of the 19 loci used to calculate heterozygosity. The loci that are out of Hardy-Weinberg equilibrium are in bold. Annealing Temperature (°C), Na is the number of alleles, Ho is the observed heterozygosity, He is the expected heterozygosity.....   | 76 |
| <b>Table 3.3</b> Summary of the heterozygosity and gamete quality metrics for the hatchery-reared female LaHave Atlantic salmon ( <i>Salmo salar</i> ) (n=42) from the broodstock at the   |    |

Harwood Fish Culture Station. Mean, maximum value, minimum value and standard deviation as well as standard error of the population mean.....77

**Table 3.4** Summary of the heterozygosity and gamete quality metrics for the hatchery-reared male LaHave Atlantic salmon (*Salmo salar*) (n=38) from the broodstock at the Harwood Fish Culture Station. Mean, maximum value, minimum value and standard deviation as well as standard error of the population mean VCL stands for curvilinear velocity, VAP stands for average path velocity, VSL stands for progressive velocity..78

**Table 3.5** General linear regression between egg quality traits and multilocus heterozygosity for females. All values represent those in the full model (df=37) while controlling for size. Unstandardized b value, t value and p value for the multilocus heterozygosity final model as well as the difference in the adjusted  $r^2$  of the final model and the final model without heterozygosity.....79

**Table 3.6** Summary of the multivariate analysis of weight of each locus on fecundity. The adjusted R2 value given to each locus was calculated by subtracting the adjusted R2 value of the model without the locus in question from that of the full multivariate model....80

**Table 3.7** General linear regression between sperm quality traits and heterozygosity for males (n = 38). Unstandardized b value, t value and p value for the multilocus heterozygosity final model selected through AIC as well as the difference in the adjusted  $r^2$  of the final model and the final model without heterozygosity. Age was a covariate in all models with the exception of density. The size principal component was a covariate in all three velocity measures at 10 sec post activation, longevity and condition. Finally,

age:size interaction was a covariate in the models for condition and  
longevity.....81

**Table A1** Mean velocity in each portion of the swim flume. Section 6 (in bold) had a  
significantly lower velocity than the other sections.....117

## LIST OF FIGURES

- Figure 3.1** Rotational Line Crossing depiction of the parental generation and the following crosses. P represents the parental generation, F<sub>1</sub> represents the first generation of crosses and F<sub>2</sub> the second generation that are offspring of crosses. The solid lines represent the females used to produce the line. The dashed lines represent the males used to produce the line. (adapted from Kincaid 1977).....83
- Figure 3.2** Distribution of multilocus heterozygosity (MLH) for the male Atlantic salmon in this study (n=38). MLH was measured as the number of heterozygous loci divided by the total number of loci.....84
- Figure 3.3** Distribution of multilocus heterozygosity (MLH) for the female Atlantic salmon in this study (n=42). MLH was measured as the number of heterozygous loci divided by the total number of loci.....85
- Figure 3.4** Negative relationship between curvilinear velocity (VCL) at 5 seconds post activation and multilocus heterozygosity ( $b=-169.04$ ,  $t(35)=-2.075$ ,  $p=0.04$ , dashed line) and at 10 seconds post activation ( $b=-54.434$ ,  $t(34)=-1.674$ ,  $p=0.103$ , solid line).....86
- Figure 3.5** Negative relationship between average path velocity (VAP) at 5 seconds post activation and multilocus heterozygosity ( $b=-161.06$ ,  $t(35)=-2.071$ ,  $p=0.05$ , dashed line) and at 10 seconds post activation ( $b=-60.107$ ,  $t(34)=-1.80$ ,  $p=0.08$ , solid line).....87
- Figure 3.6** Negative relationship between progressive velocity (VSL) at 5 seconds post activation and multilocus heterozygosity ( $b=-124.55$ ,  $t(35)=-1.972$ ,  $p=0.057$ , dashed line) and at 10 seconds post activation ( $b=-56.037$ ,  $t(34)=-2.053$ ,  $p=0.048$ , solid line).....88

**Figure 3.7** Longevity Regression: Graph depicting the negative linear correlation ( $b=-22.37$ ,  $t(33)=-1.97$ ,  $p=0.057$ ) between longevity (i.e. time point when 95% of the sperm cells have died) and Multilocus heterozygosity.....89

**Figure A1** Graphical representation of the nine zones in which water velocity was measured in the Loligo Swim-30 swim flume used in the study.....119

**Figure A2** Graphical display of the morphological scores of the four different cross types (pure LaHave, pure Sebago, Sebago dam x LaHave sire and LaHave dam x Sebago sire) created using a Discriminant Function Analysis.....120

## LIST OF APPENDICES

|  |     |
|--|-----|
| <b>Appendix 1</b> Mean length, mass and Fulton’s condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in April 2013 with one unit of standard deviation.....    | 101 |
| <b>Appendix 2</b> Mean length, mass and Fulton’s condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in August 2013 with one unit of standard deviation.....   | 103 |
| <b>Appendix 3</b> Mean length, mass and Fulton’s condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in October 2013 with one unit of standard deviation.....  | 105 |
| <b>Appendix 4</b> Mean length, mass and Fulton’s condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in December 2013 with one unit of standard deviation..... | 107 |
| <b>Appendix 5</b> Mean length, mass and Fulton’s condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in February 2014 with one unit of standard deviation..... | 109 |

|   |     |
|---|-----|
| <b>Appendix 6</b> Morphometry and Swimming Ability of Intraspecific hybrid Atlantic salmon ( <i>Salmon salar</i> ) in the juvenile life stage.....  | 111 |
| <b>Appendix 7</b> Female Raw Data as presented per individual. Wet and dry weight is measured on a per egg basis. Fecundity was calculated by multiplying the number of eggs per litre produced by the female with the number of litres she produced. Fulton’s condition factor was measured as mass over length <sup>3</sup> . Multilocus heterozygosity was measured as a percentage of heterozygous loci. Egg diameter (mm) with one standard error, the coefficient of variation was calculated as the standard deviation/mean then converted to a percentage.....  | 121 |
| <b>Appendix 8</b> Male Raw Data All sperm quality values are the mean taken between all measured tracks. Fulton’s condition factor (g/cm <sup>3</sup> ), multilocus heterozygosity was the percentage of heterozygous loci per individual, average path, curvilinear and progressive velocity were measured as units of distance per second traveled, linearity is the straightness with which the cell travelled per unit of distance, motility is the percentage of cells in a recording that are motile, longevity is the point when which 95% of the cells died and density is the number of cells per 1ml of milt..... | 123 |
| <b>Appendix 9</b> Summary of the 19 tetra and dinucleotide loci used in the study originally derived from the studies O’Reilly et al. 1996, Paterson et al. 2004, King et al. 2005 and Olafsson et al.2010. Listed in the primer sequence as well as the number of base pairs. (For further information see referenced sources).....  | 125 |
| <b>Appendix 10</b> Written Permission from Co-Authors.....  | 127 |



## CHAPTER 1: GENERAL INTRODUCTION

Darwinian fitness or reproductive fitness is defined as the number of offspring that reach sexual maturity contributed by an individual (see Frankham et al. 2002). An individual's reproductive fitness includes attributes such as their survival, capability of siring offspring, etc. (Frankham et al. 2002). When predicting one's individual fitness, there are many genetic factors that can influence the quantitative trait, such as the level of relatedness or genomic similarity between the parental generation (Edmands 2002; reviewed in Frankham et al. 2002). Many are familiar with the negative impacts of mating individuals of close relatedness (i.e. inbreeding depression) and the negative effects of recessive deleterious alleles being phenotypically expressed as a result of increased homozygosity (Charlesworth and Charlesworth 1999). The opposite end of the spectrum (i.e. outbreeding, or causing increased heterozygosity) does not always result in higher fitness, and in certain cases can be just as negative of an influence on individual fitness as inbreeding (e.g. Edmands 1999; reviewed in Frankham et al. 2011). Known as outbreeding depression, it can arise from the hybridization of two reproductively segregated populations.

### **Hybridization**

The term hybrid is not isolated to the crossing of two different species, it is a term that can also be applied to describe the offspring of intraspecific crosses between individuals of two different populations that are distinguishable by at least one heritable characteristic (e.g. morphology, see Harrison 1990). Unlike interspecific hybridization, intraspecific hybridization often produces fertile individuals; however, the fitness of

offspring can still vary relative to that of their parental populations (Edmands et al. 1999, 2002, 2007; Frankham et al. 2002). When hybrid fitness is inferior relative to the mean of the parental populations it is known as outbreeding depression (or hybrid breakdown, Templeton 1986; Edmands 1999, 2007; Frankham et al. 2002), whereas when the fitness of hybrid offspring is superior relative to the mean fitness of both parental populations it is known as heterosis (or hybrid vigour, Frankham et al. 2002; Edmands and Timmerman 2003; Edmands 2007). Signs of outbreeding depression can include an overall decrease in fertility and survival (Harrison 1990), whereas evidence of heterosis can include an increase in biomass, fertility and growth rate in hybrid offspring compared to that of their parental populations (reviewed in Frankham et al. 2002). The mechanisms responsible for outbreeding depression and heterosis can be divided into extrinsic factors (i.e. factors under environmental influence) and intrinsic factors (factors influenced by the genotype) (Templeton 1986; Edmands and Timmerman 2003). Expressed differences in traits under selection will also vary depending on the underlying genetic architecture (number and effects of contributing genes) (Naish and Hard 2008).

### *Outbreeding Depression*

There are many potential causes that can contribute to outbreeding depression. The potential intrinsic causes behind outbreeding depression can be attributed to both additive and nonadditive genetic effects. One example of an intrinsic effect is underdominance (i.e. heterozygotes have inferior fitness to homozygotes) (Lynch 1991). Underdominance is a nonadditive genetic effect that may encourage outbreeding depression through epistatic interactions that heterozygotes have inherited that are not seen in homozygotes (Waser and Price 1983). Another potential cause of outbreeding

depression is the break-up of co-adapted gene complexes as a result of incompatibilities between the parental populations (Burton 1987; Lynch 1991; Edmands 1999; McGinnity et al. 2003). For example, Gilk et al. (2004) found that F<sub>2</sub> intraspecific Pink salmon hybrids (*Oncorhynchus gorbuscha*) had lower survival rates than the pure counterparts, suggesting it was the result of disrupted co-adapted gene complexes.

Outbreeding depression may also be the result of extrinsic causes such as the loss of local adaptations (Tymchuk et al. 2007). As a result of additive genetic effects, hybrids may have an intermediate phenotype compared to their parental populations, and therefore may have lost critical local adaptations as a result, causing them to be not suited for either parental population's environment (Templeton 1986; Edmands 2007; Tymchuk et al. 2007).

### *Heterosis*

As with outbreeding depression, the intrinsic causes behind heterosis can be attributed to both additive and nonadditive genetic effects. One of the potential nonadditive causes is referred to as overdominance (a.k.a. heterozygote advantage); the explanation that heterozygotes have superior fitness relative to homozygotes as a result of positive allelic interactions (Lynch 1991; Birchler et al. 2003; Edmands and Timmerman 2003). Another possible cause of heterosis is known as dominance; (i.e. when the recessive deleterious alleles of one parent are masked by superior dominant alleles of the other (Lynch et al. 1991; Birchler et al. 2003; Edmands and Timmerman 2003). A third potential cause of heterosis is the added genetic diversity that breeding separate populations provides. When parental populations are small, and highly inbred, the

addition of foreign genetic material may reverse inbreeding depression and result in the expression of hybrid vigour (Frankham et al. 2002; Crespel et al. 2013). An example of a population that experienced heterosis as a result of genetic diversity is the Florida panther (*Puma concolor coryi*) (Johnson et al. 2010). In the 1990's the population of *P. concolor coryi* in Florida was relegated to a few dozen mature individuals that were highly inbred (Johnson et al. 2010). Sexually mature individuals (n=8) from Texas were then transported to Florida in order to reproduce in the hopes of increasing genetic diversity and ultimately promoting hybrid vigour (Johnson et al. 2010). Over a decade after successful reproduction with the individuals from Texas, heterozygosity in the *P. concolor coryi* population increased in Florida-Texas hybrids; their survivorship was higher than that of the pure Florida crosses (Johnson et al. 2010).

In certain cases the added genetic divergence between parental populations leads to a mix of outbreeding depression and heterosis in different generations. For example, Edmands (1999) found that genetic divergence between distinct parental populations of the copepod species, *Trigriopus californicus*, was correlated with hybrid vigour in the F<sub>1</sub> generation and hybrid breakdown in the F<sub>2</sub> and backcross generations. This delayed manifestation of outbreeding depression is the result of the return of deleterious homozygotic combinations between alleles that were present in the parental populations (Templeton 1986; Edmands 2007). In this study, the reduced fitness caused by hybridization was even displayed in the F<sub>3</sub> generation with the variance in displayed fitness ranging from higher than parental fitness to lower than F<sub>2</sub> fitness (Edmands 1999).

## **Heterozygosity Fitness Correlations**

Heterozygosity fitness correlations (HFCs) are correlations between heterozygosity at at least one locus and measured fitness related traits (e.g. life history traits) (reviewed in Chapman et al. 2009; reviewed in Szulkin et al. 2010). As mentioned in previous paragraphs, although genetic diversity and therefore heterozygosity are generally assumed to have positive effects on fitness, in certain cases negative effects can be seen as well.

When heterozygosity was most commonly measured using allozymes, the prevailing hypothesis was that the correlation with fitness was a direct result of selection on the marker, otherwise known as the direct effect hypothesis (David 1998; Lynch and Walsh 1998; reviewed in Hansson and Westerberg 2002). Now that the majority of HFC studies are done with microsatellites, which are assumed to be neutral markers (Jarne and Lagoda 1996), there are two current prevailing hypotheses to explain observed correlations between fitness related traits and heterozygosity. The first is known as the general effects hypothesis which states; an individual's level of heterozygosity at the microsatellite level is reflective of their heterozygosity at the genome level (David 1998; Lynch and Walsh 1998; reviewed in Hansson and Westerberg 2002; Coltman and Slate 2003). The second is referred to as the local effects hypothesis which states that each marker reflects the heterozygosity of loci under selection in the chromosomal vicinity of the microsatellite due to linkage disequilibrium (David 1998; Lynch and Walsh 1998; reviewed in Hansson and Westerberg 2002). When HFCs are due to general effects, they are assumed to be positive in nature (Szulkin and David 2011). This is due to the fact that it is assumed that general effects are more common in populations with a high degree of

variance in the inbreeding coefficient, and if the heterozygosity at microsatellite markers is reflective of overall heterozygosity, the most heterozygous individuals will be the least inbred and therefore have higher fitness than their more homozygous counterparts (reviewed in Hansson and Westerberg 2002). However, HFC can be quadratic or negative in relation as well, either as a result of local effects (Olano-Marin 2011) or outbreeding depression (Neff 2004). Because life history traits such as gamete quality are particularly sensitive to selection, they make particularly good traits to measure when determining effects of inbreeding and outbreeding.

### *Gamete Quality Traits*

As both sperm and eggs are highly specialized cells, any alterations to their characteristics can affect fertilization rates as well as offspring survival (Gage et al. 2002; Srivastava and Brown 1991). Spermatozoa have many selected characteristics that optimize their chances at fertilizing the ova (Gage et al. 1995; Gage et al. 2002). Characteristics that have been shown to influence fertilization success and therefore the quality of the sperm include velocity (Gage et al. 2004), motility, longevity and density (Gage et al. 1995; reviewed in Snook 2005). Eggs also have characteristics which can influence the survival of the embryo. The volume and mass of the egg are indicative of the energy the embryo will receive until the exogenous feeding stage (Srivastava and Brown 1991), and studies have shown that individuals that emerge from larger eggs have higher survival than those that emerge from smaller eggs (Einum and Fleming 1999, 2000). Heterozygosity has been found to correlate with both sperm (Gage et al. 2006; Fitzpatrick and Evans 2009) and egg quality (Heath et al. 2002; Garcia-Navas et al. 2009;

Wetzel et al. 2012) in various taxa. However, the studies that have examined heterozygosity in relation to sperm and egg quality are limited in number.

### **Atlantic salmon**

Atlantic salmon (*Salmo salar*) are a post-glacial species with distribution across Europe as well as eastern Canada and the North-Eastern United States. Although the vast majority have an anadromous life cycle, migrating to the Atlantic Ocean in order to reach maturity, there are populations that have a completely freshwater life cycle (Ward 1932).

Atlantic salmon were native to Lake Ontario and were an important source of income to the economy of Upper Canada in the 18<sup>th</sup> century (Dunfield 1985). They formed what is believed to be a landlocked population (Dunfield 1985) until the late-19<sup>th</sup> century when they were extirpated as a result of anthropogenic activities that range from dam construction, overfishing and forestry (MacCrimmon 1977). Many of the dams that once impeded the upstream migration of Atlantic salmon have either been removed or fish ladders have been incorporated into others through massive restoration efforts in attempts to make conditions more favorable for Atlantic salmon (Stanfield and Jones 2003). In order to determine if Lake Ontario is currently more suitable than previous years for Atlantic salmon, since 1987, fry and fingerlings have been released annually into surrounding tributaries as a pilot study. It was determined that conditions were suitable for Atlantic salmon in Lake Ontario. Therefore, since 2005, the Ontario Ministry of Natural Resources and Forestry (MNR) and partners have implemented a stocking effort to try and reintroduce the species back into Lake Ontario with the use of 3 separate strains (i.e. LaHave, Sebago and Lac St-Jean). The LaHave strain originates from the

LaHave River in Nova-Scotia (MNRF 2006), and has been reared in captivity since 1989 (Dimond and Smitka 2005). Additional broodstocks were developed from Sebago Lake in Maine and Lac St-Jean in Quebec, and the stocking of these two strains into Lake Ontario tributaries was initiated in 2008 (MNRF 2009), and, as of that year, both populations have been stocked simultaneously. The Sebago Lake strain was chosen as a deliberate ecological contrast to the LaHave strain (Dimond and Smitka 2005) (see Table 1.1), as the wild population is landlocked (Ward 1932) and lake conditions are considered similar to those in Lake Ontario (Toivonen 1971). As the population's freshwater life cycle provides a suitability to a different environment compared to the LaHave strain, and it has been used for previous successful introductions in the state of Maine, it is considered a good candidate for the Lake Ontario reintroduction efforts. The Lac St-Jean strain is another landlocked strain that originates from a lake in central Quebec. The population is used in other stocking efforts and based on geographic proximity, it is speculated that the population is the most genetically similar to the historical population in Lake Ontario. Based on these characteristics, the population is considered another good candidate for Lake Ontario. However, as the survival in captive settings is not as good in the Lac St-Jean strain as it is in the Sebago and LaHave strains, it is not stocked in as high a quantity (MNRF 2009).

### **Overview of the thesis**

The objectives of my thesis were to evaluate fitness related traits in Atlantic salmon to determine the implications of genetic quality (i.e. the potential beneficial or detrimental effects of the genome on characteristics of phenotypic expression) on fitness related traits in the Atlantic salmon broodstock. The objective of Chapter 2 was to



evaluate fitness related traits in juvenile Atlantic salmon crosses between two of the three mentioned allopatric strains (i.e. Sebago and LaHave) in order to determine whether outbreeding depression or heterosis was occurring in the hybrid crosses using a 2 x 2 full factorial breeding design. As fitness of individuals in the hatchery where the stocks are reared is equally as important as those in the wild, Chapter 3 evaluates the occurrence of heterozygosity fitness correlations in the gamete quality of the LaHave strain (the most stocked strain). In order to do this, sperm and egg quality were measured and compared to multilocus heterozygosity using 19 microsatellite markers.

## References

Birchler JA, Auger DL, Riddle NC (2003) In Search of the Molecular Basis of Heterosis. *The Plant Cell* 15: 2236-2239

Burton RS (1987) Differentiation and Integration of the Genome in Populations of the Marine Copepod *Tigriopus californicus*. *Evolution* 41: 504-513

Crespel A, Audet C, Bernatchez L, Garant D (2013) Effects of rearing environmental and strain combination on hybrid vigour in brook trout. *N Am J Aquacult* 74: 188-198

Chapman JR, Nakagawa S, Coltman DW, Slates J, Sheldon BC (2009) A quantitative review of heterozygosity-fitness correlations in animal populations. *Mol Ecol* 18: 2746-2765

Charlesworth B, Charlesworth D (1999) The genetic basis of inbreeding depression. *Genet Res., Camb* 74: 329-340

Coltman DW, Slate J (2003) Microsatellite measures of inbreeding: a meta-analysis. *Evolution* 57: 971-983

David P (1998) Heterozygosity-fitness correlations: new perspectives on old problems. *Heredity* 80: 531-537

Dimond P, Smitka J (2005) Evaluation of Selected Strains of Atlantic Salmon as Potential Candidates for the Restoration of Lake Ontario. Trout Unlimited Canada Technical Report ON-12. Guelph, Ontario: Trout Unlimited Canada

Dunfield RW (1985) The Atlantic salmon in the History of North America. Fisheries and Oceans, Ottawa. pp.181

Einum S, Fleming IA (1999) Maternal effects of egg size in brown trout (*Salmo trutta*): norms of reaction to environmental quality. Proc R Soc Lond B 266:2095-2100

Einum S, Fleming IA (2000) Selection against late emergence and small offspring in atlantic salmon (*Salmo salar*). Evolution 54:628-639

Edmands S (1999) Hybrid vigour and outbreeding depression in interpopulation crosses spanning a wide range of divergence. Evolution 53: 1757-1768

Edmands S (2002) Does parental divergence predict reproductive compatibility? Trends Ecol Evol 17: 520-526

Edmands S, Timmerman CC (2003) Modeling factors affecting the severity of outbreeding depression. Conserv Biol 17: 883-892

Edmands S (2007) Between a rock and a hard place: evaluating the relative risks of inbreeding and outbreeding for conservation and management. Mol Ecol 16: 463-475

Fitzpatrick JL, Evans JP (2009) Reduced heterozygosity impairs sperm quality in endangered mammals. Biology Letters 5: 320-323

Frankham R, Ballou JD, Briscoe DA (2002) Introduction to Conservation Genetics. Cambridge University Press, UK.

Frankham R, Ballou JD, Briscoe DA (2010) Introduction to Conservation Genetics, The 2nd Ed. Cambridge, UK. pp.618

Frankham R, Ballou JD, Eldridge MDB, Lacy RC, Ralls K, Dudash MR and Fenster CB (2011) Predicting the Probability of Outbreeding Depression. *Conserv Biol* 25: 465-475

Gage MJG, Stockley P, Parker GA (1995) Effects of Alternative Male Mating Strategies on Characteristics of Sperm Production in the Atlantic Salmon (*Salmo salar*): Theoretical and Empirical Investigation. *Philosophical Transactions: Biological Sciences* 350: 391-399

Gage MJG, MacFarlane C, Yeates S, Shackleton R, Parker, GA (2002) Relationships between sperm morphometry and sperm motility in the Atlantic salmon. *Journal of Fish Biology* 61: 1528-1539

Gage MJG, Macfarlane CP, Yeates S, Ward RG, Searle JB, Parker GA (2004) Spermatozoal Traits and Sperm Competition in Atlantic Salmon: Relative Sperm Velocity Is the Primary Determinant of Fertilization Success. *Current Biology* 14: 44-47

Garcia-Navas V, Ortego J, Sanz JJ (2009) Heterozygosity-Based Assortive Mating in Blue-tits (*Cyanistes caeruleus*): Implications for the Evolution of Mate Choice. *Proceedings: Biological Sciences* 276:2931-2940

Gilk SE, Wang IA, Hoover CL, Smoker WW, Taylor SG, Gray AK, Gharrett AJ (2004) Outbreeding depression in hybrids between spatially separated pink salmon, *Oncorhynchus gorbuscha*, populations: marine survival, homing ability, and variability in family size. *Environ Biol Fishes* 69: 287-297

Granier S, Audet C, Bernatchez L (2011) Hybrid vigour and outbreeding depression between strains of young-of-the-year brook trout (*Salvelinus fontinalis*). *Can J Zool* 89: 190-198

Hansson B, Westerberg L (2002) On the correlation between heterozygosity and fitness in natural populations. *Mol Ecol* 11:2467-2474

Harrison RG (1990) Hybrid zones: windows on evolutionary processes. In: Futuyma, D. and Antonovics, J. (ed) *Oxford Surveys in Evolutionary Biology*. Oxford University Press, Oxford. pp:69-128

Heath DD, Bryden CA, Shrimpton JM, Iwama GK, Kelly J, Heath JW (2002) Relationships between heterozygosity, allelic distance ( $d_2$ ), and reproductive traits in chinook salmon, *Oncorhynchus tshawytscha*. *Can J Fish Aquat Sci* 59:77-84

Holm E, Mandrak NE, Burrige, ME (2009) *The ROM Field Guide to freshwater fishes of Ontario*. Royal Ontario Museum

Jarne P, Lagoda JL (1996) Microsatellites, from molecules to populations and back. *Trends Ecol Evol* 8: 285-288

Johnson WE, Onorato DP, Roelke ME, Land ED, Cunningham M, Belden RC, McBride R, Jansen D, Lotz M, Shindle D, Howard J, Wildt DE, Penfold LM, Hostetler JA, Oli MK, O'Brien SJ (2010) Genetic Restoration of the Florida Panther. *Science* 329:1641-1645

Lynch M (1991) The Genetic Interpretation of Inbreeding Depression and Outbreeding Depression. *Evolution* 45: 622-629

Lynch M, Walsh B (1998) *Genetics and Analysis of Quantitative Traits* Sinauer pp.980

MacCrimmon HR (1977) *Animals, man and change: alien and extinct wildlife of Ontario*. McClelland and Stewart, Toronto.

McGinnity P, Prodöhl P, Ferguson A, Hynes R, Maoiléidigh NO, Baker N, Cotter D, O’Hea B, Cooke D, Rogan G, Taggart J, Cross T (2003) Fitness Reduction and Potential Extinction of Wild Populations of Atlantic Salmon, *Salmo salar*, as a Result of Interactions with Escaped Farm Salmon. *Proceedings: Biological Sciences* 270: 2443-2450

Naish KA, Hard JJ (2008) Bridging the gap between the genotype and the phenotype: linking genetic variation, selection and adaptation in fishes. *Fish Fish* 9: 396-422

Neff BD (2004) Stabilizing selection on genomic divergence in a wild fish population. *PNAS* 101: 2381-2385

Olano-Marin J, Mueller JC, Kempenaers B (2011) Heterozygosity and survival in blue tits (*Cyanistes caeruleus*): contrasting effects of presumably functional and neutral loci. *Mol Ecol* 20: 4028-4041

Ontario Ministry of Natural Resources (2006) Lake Ontario Fish Communities and Fisheries: 2005 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.

Ontario Ministry of Natural Resources (2009) Lake Ontario Fish Communities and Fisheries: 2008 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.

Snook RR (2005) Sperm in competition: not playing by the numbers. *Trends Ecol Evol* 20: 46-53

- Srivastava RK, Brown JA (1991) The biochemical characteristics and hatching performance performance of cultured and wild Atlantic salmon (*Salmo salar*) eggs. Can J Zool 69: 2436-2441
- Stanfield L, Jones ML (2003) Factors Influencing Rearing Success of Atlantic Salmon Stocked as Fry and Parr in Lake Ontario Tributaries. N Am J Fish Sci 23:1175-1183
- Szulkin M, Bierne N, David P (2010) Heterozygosity-fitness correlations: a time for reappraisal. Evolution 64:1202-1217
- Szulkin M, David P (2011) Negative heterozygosity-fitness correlations observed with microsatellites located in functional areas of the genome. Mol Ecol 20: 3949-3952
- Templeton AR (1986) Coadaptation and outbreeding depression. In Soulé M (ed) Conservation biology: the science of scarcity and diversity. Sinauer, Sunderland, MA. pp. 105-116
- Toivonen J (1971) The fish fauna and limnology of large oligotrophic glacial lakes in Europe (about 1800 A.D.). J Fish Res Board Can 29: 629-637
- Tymchuk WE, Sundstrom LF, Devlin RH (2007) Growth and Survival Trade-offs and outbreeding depression in rainbow trout (*Oncorhynchus mykiss*). Evolution 61:1225-1237
- Van Burkirk J, Willi Y (2006) The Change in Quantitative Genetic Variation with Inbreeding. Evolution 60: 2428-2434
- Vladić TV, Afzelius BA, Bronnikov GE (2002) Sperm Quality as Reflected Through Morphology in Salmon Alternative Life Histories. Biol Reprod 66: 98-105

Ward HB (1932) The Origin of the Landlocked Habit in Salmon. Proc Nat Acad Sc 18: 569-580

Waser NM, Price MV (1983) Optimal and actual outcrossing in plants, and the nature of plant- pollinator interaction. In C. E. Jones and R. J. Little (eds.), Handbook of Experimental Pollination Biology. pp. 341-359

Wetzel DP, Stewart IRK, Westneat DF (2012) Heterozygosity predicts clutch and egg size but not plasticity in a house sparrow population with no evidence of inbreeding. Mol Ecol 21: 406-420



Table 1.1: Approved populations of Atlantic salmon for Lake Ontario. The table represents a comparison between the known conditions of the current 3 approved candidate populations and the speculated characteristics of the Lake Ontario population

|                          | Ontario<br>(extirpated) | LaHave (Nova-<br>Scotia) | Sebago (Maine) |
|--------------------------|-------------------------|--------------------------|----------------|
| Freshwater life<br>cycle | Yes                     | No                       | Yes            |
| Anadromous               | Yes                     | Yes                      | No             |
| Oligotrophic             | Yes                     | No                       | Yes            |
| Genetic<br>Similarity    | Yes                     | No                       | ?              |

## CHAPTER 2: EFFECTS OF INTRASPECIFIC HYBRIDIZATION BETWEEN TWO STRAINS OF ATLANTIC SALMON (*SALMO SALAR*) ON JUVENILE SURVIVAL AND FITNESS-RELATED TRAITS: IMPLICATIONS FOR RESTORATION ECOLOGY

### SYNOPSIS

Hybridization between species or divergent conspecific populations may result in hybrid offspring exhibiting either superior (heterosis) or inferior (outbreeding depression) fitness relative to their parental populations. As both heterosis and outbreeding depression have previously been demonstrated in salmonids, consequences of interbreeding between divergent populations may therefore be relevant to salmonid conservation programs and restoration efforts. Atlantic salmon (*Salmo salar*) were once native to Lake Ontario but were extirpated by the late-19<sup>th</sup> century as a result of anthropogenic causes. Multiple allopatric populations of Atlantic salmon are currently being stocked in an effort to re-establish a self-sustaining population in Lake Ontario. This study evaluated whether interbreeding between individuals from Sebago Lake (Maine) and the LaHave River (Nova-Scotia) will result in the expression of heterosis or outbreeding depression in juveniles. This was accomplished by generating full-factorial 2x2 mating crosses between the two strains and comparing multiple traits (growth and survival) associated with individual fitness between the four cross types (pure within-population breeding and reciprocal hybrids). Hybrid juveniles did not display any signs of outbreeding depression or heterosis. Despite these results, further studies on reproductive fitness and comparative fitness of backcross and second-generation hybrids are recommended to assess potential consequences for this and similar restoration efforts.

## **Introduction**

Although hybridization is often overlooked in the conservation and restoration management of endangered and extirpated populations, it is relevant, as it can be either a potent tool or significant challenge (Edmands 2007, Frankham et al. 2011). Unlike interspecific hybridization, intraspecific hybridization (hybridization within species) often produces viable and fertile individuals; however, the fitness of offspring can be higher (heterosis) or lower (outbreeding depression) than that of their parental populations. Evidence of heterosis can include an increase in biomass, fertility and growth rate in hybrid offspring compared to that of their parental populations (reviewed in Frankham et al. 2002), whereas signs of outbreeding depression can include an overall decrease in fertility and survival (Harrison 1990). As heterosis or neutral outcomes are not considered detrimental to conservation efforts, the focus of selecting candidate populations should be on the prevention of outbreeding depression. Outbreeding depression is considered a likely outcome when breeding populations with fixed chromosomal differences, if they have lived in different environments for over 20 generations and if they have not had any gene flow for over 500 years (Frankham et al. 2011). Therefore, in order to avoid the onset of outbreeding depression when more than one population is being used in reintroduction efforts, it is best to choose parental populations that have low genetic divergence from one another as well as similar local adaptations (Edmands 1999, 2002).

The mechanisms responsible for outbreeding depression can be divided into extrinsic factors (i.e. factors under environmental influence) and intrinsic factors (factors influenced by the genotype) (Templeton 1986; Edmands and Timmerman 2003). Expressed differences in traits under selection will also vary depending on their

underlying genetic architecture (number and effects of contributing genes) as well as the potential correlation with other traits (Naish and Hard 2008). An example of allopatric crosses which have resulted in outbreeding is, Gilk et al. (2004) which crossed allopatric populations of Pink salmon (*Oncorhynchus gorbuscha*) which resulted in lower return rates for the F<sub>1</sub> generation and lower survival for the F<sub>2</sub> generation. Other studies have similarly cautioned against interbreeding between wild and hatchery fish populations with the same population of origin as they may have divergent genetic traits. For example, Araki et al. (2007) saw a 37.5% decrease per generation in the reproductive success of captively reared steelhead trout (*O. mykiss*) due to underlying genetics. As the consequences of lower survival and overall fitness are a possible outcome of outbreeding depression, it is therefore a concern for any conservation or reintroduction effort that utilizes multiple populations.

Atlantic salmon (*Salmo salar*) were once native to Lake Ontario, but were extirpated in the late 19<sup>th</sup> century as a result of anthropogenic activities such as dam construction, land clearing, and overfishing (MacCrimmon 1977). Land restoration efforts as well as restoration efforts in many tributaries have made conditions more favorable for Atlantic salmon (Ontario Ministry of Natural Resources and Forestry (MNRF) 2003; Stanfield and Jones 2003). Restoration began in earnest in 2005, using an anadromous strain that originates from the LaHave River in Nova-Scotia (MNRF 2005). Additional broodstock was developed from Sebago Lake in Maine, and stocking into Lake Ontario tributaries was initiated in 2008 (MNRF 2008), and, as of this point, both populations have been stocked simultaneously. The Sebago Lake strain was chosen as a deliberate ecological contrast to the LaHave strain (Dimond and Smitka 2005), as the wild

population is landlocked (Ward 1932) and lake conditions are considered similar to those in Lake Ontario (Toivonen 1971). As the population's freshwater life cycle provides a suitability to a different environment compared to the LaHave strain, and it has been used for previous successful introductions in the state of Maine, it is considered a good candidate for the Lake Ontario reintroduction efforts. However, as the strains have been separated for over 500 years and living in different environmental conditions for over 20 generations, the manifestation of outbreeding depression is a possible outcome of breeding these two populations (see Frankham et al. 2011).

This study evaluated the implications of interbreeding (intraspecific hybridization) between these two Atlantic salmon strains for the Lake Ontario reintroduction effort, testing specifically for juvenile heterosis or outbreeding depression in a controlled hatchery environment that is consistent with current hatchery practices. As the manifestation of heterosis and outbreeding depression can also vary with life stages (Fraser et al. 2010; Granier et al. 2011; Crespel et al. 2013), the importance of examining the fitness of the juvenile stage is critical as there is high mortality in this stage; with up to 90% of juveniles dying by the smolt migration phase (Thorstad et al. 2011). The potential for outbreeding depression or heterosis was assessed by measuring survival and multiple fitness-related traits (length, mass, Fulton's condition, and growth rate) over time at ecologically relevant stages by crossing individuals from the Sebago and LaHave populations in order to create family blocks composed of both pure strains and their reciprocal hybrid siblings.

## **Materials and Methods**

### *Full-factorial breeding design*

To assess potential fitness effects resulting from intraspecific hybridization, reproductive adults from both the Sebago and LaHave strains maintained at the MNRFCodrington Fisheries Research Facility (44.18.05°N, 78.29.40°W) were selected haphazardly in order to create 20 distinct half-sibling family blocks . These 20 family blocks were created using a 2x2 breeding design (blocked factorial breeding design); using one male and female from each strain to produce half-sibling family blocks consisting of a pure Sebago cross (S/S), a pure LaHave cross (L/L) and their reciprocal hybrids (LaHave dam/Sebago sire (L/S) and Sebago dam/LaHave sire (S/L)). Each adult was used in only one 2x2 cross, resulting in 20 independent family blocks. The full factorial breeding design allows for the separate evaluation of intrinsic genetic factors and maternal effects (both additive and environmental) and paternal effects (Pitcher and Neff 2006, 2007; Neff et al. 2011). The eggs for the blocks were fertilized on two separate dates (November 22, 2012 (n=8 blocks) and December 4, 2012 (n=12 blocks)). After fertilization, the eggs from each separate cross were randomly allocated into the cells of two separate incubation stacks, each containing five trays with 16 cells per tray.

### *Fertilized Egg Survival*

Fertilization success assays for all family blocks took place between December 21, 2012 and January 10, 2013, and survival of the fertilized eggs was monitored three times a week from January 14, 2013 until the latest date of hatching (March 4, 2013). If the eggs changed from transparent or translucent to opaque, they were deemed dead. To

examine if the dead eggs had been fertilized pre-mortality, they were submersed in acetic acid (5%) (see Hoysak and Liley 2001); if the eggs turned white after exposure to the acetic acid, they were considered fertilized; eggs that remained clear were considered unfertilized. Only eggs that had been fertilized were considered in the survival comparisons.

### *Rearing*

Once the alevin had absorbed their yolk sacs and manual feeding began, up to 100 individuals (mean +/- s.e.:  $97 \pm 1.2$ ) from each full sibling cross were transported from the incubation trays and randomly allocated into separate 40 L family rearing tanks at the University of Windsor Great Lakes Fish and Research Centre in LaSalle, Ontario. The facility is equipped with a scaled down recirculation system to ensure that the water quality in all the tanks is similar, not unlike those found at the provincial Atlantic salmon hatchery Normandale. Water quality (dissolved oxygen, pH, and temperature) was examined daily to ensure families were being held at optimal water conditions. On April 29, 2013, each tank was manually thinned down haphazardly to 50 individuals by removing individuals present throughout the water tank, with the exception of two tanks which began with 16 and 13 fish due to low egg survival. During the first half of the rearing stage (March 2013 to August 2013), the tanks were thinned on three occasions in order to accommodate growth (April 29, June 21 and August 2013), and to keep densities at relatively consistent levels in order to limit density effects on the early growth of the fish during this critical growth period.

### *Length, Mass & Fulton's Condition Factor*

Length, mass and Fulton's condition factor (K) were measured a total of five times during juvenile development between April 2013 to February 2014; the measurements covered the significant life stages from button-up to smolting (fry, parr and smolt). The first measurement took place during the "button-up" stage (shortly after the yolk sac was absorbed), the second measurement took place during the fry stage, 20 fish per tank were haphazardly selected in order to be weighed and measured. In order to prevent bias, individuals were netted from all areas of the water column. The third, fourth and fifth measurements took place when they were parr, during later period of parr stage and during their smolting period (after they had lost their parr marks). Ten fish per cross type (i.e. pure LaHave cross, pure Sebago cross, Sebago dam x LaHave sire cross or LaHave dam x Sebago sire cross) within each family block were haphazardly selected to be weighed and measured. The first two measurements took place at two week intervals in order to account for the difference in fertilization dates. The fish were anaesthetized using MS-222, each individual's mass (+/- 0.001 g) was then taken using an electronic scale (Denver Instrument TP 323) and digital images (with a size standard) taken of each fish were analysed using Image J (<http://imagej.nih.gov/ij/>) in order to assess fork length. Fulton's condition factor was calculated as  $K = (W/L^3) * 10\ 000$  (Ricker 1975).

### *Survival*

The survival of the juveniles was analyzed during four time periods: egg stage (January 2013 - February 2013), fry stage (April 2013 - June 2013), early parr stage (October 2013 - December 2013), and late parr stage (December 2013 - February 2014)



by comparing the change in the number of offspring in each tank that had occurred over that period of time.

### *Statistical Analyses*

Survival was examined at four separate time points using a binomial generalized linear mixed effects model with LaPlace approximation using version 1.1-7 of the lme4 package (Bates et al. 2014) in R 3.1.1 (R Core Team 2014). Cross type was evaluated as a fixed effect in the model, whereas dam ID, sire ID and dam ID x sire ID interaction were evaluated as random factors. Due to biological significance, density, dam ID and sire ID were included in all final models, whereas position effects (i.e. stack, tray, cell) and dam ID x sire ID were included in the final model if deemed statistically significant. A likelihood ratio test fitted with maximum likelihood (ML) was used in order to generate p-values between a full model and a reduced model without the variable in question.

Analyses of length, mass and Fulton's condition factor data at the five time points were also completed using version 1.1-7 of the lme4 package in R to generate linear mixed effects models. Cross type and density were entered in the model as fixed effects, while dam ID, sire ID and dam ID x sire ID interaction were entered in the model as random factors. Due to biological significance, despite the results from the AIC, density, dam and sire effects were always included in the full model. The significance of cross type was assessed using an F-test with a Kenward-Roger degree of freedom estimation in the package pbkrtest (Halekoh and Højsgaard 2014) whereas, the p-values for the random factors were generated using a likelihood ratio test fitted with ML between a full model and a reduced model without the variable in question.

Of the 20 2x2 mating crosses that were established, 6 factorial crosses were discarded due to accidental mixing, missing measurements of certain tanks at one or more time points and equipment failure which lead to the mortality of one tank in a family block during the later portion of the rearing period. In order to better control for factors such as parental ID and uneven sampling of treatment groups that could affect the results, any such experimental error with one (or more) of the tanks in a full factorial cross, led to the discard of all tanks in that family, thereby excluding that family from any further statistical analysis.

## **Results**

Summary statistics of survival and the fitness-related traits for the remaining 14 family blocks are presented in Table 2.1. Cross type did not affect the survival of eggs ( $p=0.96$ ), fry ( $p=0.47$ ), early parr ( $p=0.25$ ) or late parr ( $p=0.90$ ) (Table 2.1). Dam ID effects had a large influence over the survival of eyed eggs and early parr, but were not significant for survival of other life stages (Table 2.1). Dam ID x sire ID interaction only had a significant influence over the survival of the fry stage (Table 2.1). Sire ID effects had no significant influence over the survival of the individuals during any of the life stages (Table 2.1).

Over the five intervals, cross type had no significant effect on any of the characteristics associated with fitness (Table 2.2). Dam ID effects were significant for length, mass and condition at early life stages, but decreased in importance at the parr and smolt life stages (Table 2.2). Sire ID effects also explained part of the variance of mass and length during the early measurements and condition during the later measurements

(Table 2.2). Interaction effects (dam ID x sire ID) contributed to the variance of the early condition measurements but little to the rest of the fitness related traits. Tank density had significant effects on length and mass at the majority of the measurement time points (Table 2.2).

## **Discussion**

The results of this study show no significant influence of hybridization on survival or any of the fitness related traits measured in several stages of ontogeny of juvenile Atlantic salmon. Understanding the potential effect of intraspecific hybridization of Atlantic salmon, particularly in the juvenile life stages, is critical to the successful reintroduction of the species back to Lake Ontario if the program intends to continue using more than one population simultaneously that have been separated for over a thousand generations, as it has the potential to cause complications for hybrid offspring.

Outbreeding depression has been identified as a significant concern for interbreeding between divergent populations (Edmands 2007, Frankham et al. 2011), and has previously been documented in hatchery-reared salmonids (e.g. Araki et al. 2007, Granier et al. 2011), the severity of which has varied across studies. However, the detection of outbreeding depression or heterosis can also be the result of the environmental surroundings. For example, using inbred crosses, Houde et al. (2011) detected heterosis or outbreeding depression in a limited number of their families of Atlantic salmon depending on rearing environment (due to the loss of local adaptations) (Houde et al. 2011). A subsequent study found that the strength of both inbreeding and outbreeding depression varied annually along with environmental quality (Rollinson et al.

2014). In years with poor environmental quality, maternal effects accounted for greater variance in juvenile fitness than years with closer to optimum environmental quality (Rollinson et al. 2014). Therefore, there are various reasons for why the crossing of allopatric populations did not result in the expression of outbreeding depression in the setting of this study.

The most likely reason for our lack of observation of outbreeding depression had to do with the fact that we were looking at the F<sub>1</sub> generation. When outbreeding depression is observed in the F<sub>1</sub> generation it is typically the result of the loss of local adaptations as the result of an intermediate phenotype (Lynch 1991; Edmands 2007). Intrinsic incompatibilities between parental populations (e.g. the break-up of co-adapted gene complexes) (Burton 1987; McGinnity et al. 2003; McClelland and Naish 2007; Tymchuk et al. 2007) typically result in outbreeding depression in the F<sub>2</sub> generation or later when the parental genomes are subject to recombination (Dobzhansky 1948; Lynch 1991; Edmands 1999; Birchler et al. 2003; Edmands and Timmerman 2003; McGinnity et al. 2003; McClelland et al. 2005; Tymchuk et al. 2007). Outbreeding depression as a result of the breakup of co-adapted gene complexes has been demonstrated in intraspecific salmonid hybrids, for example, Gilk et al. (2004) found reduced survival in F<sub>2</sub> intraspecific *Oncorhynchus gorbuscha* hybrids relative to their pure counterparts, suggesting it was the result of disrupted co-adapted gene complexes. Another example is with Atlantic salmon in McGinnity et al. (2003), F<sub>2</sub> hybrid crosses saw significantly higher egg mortality than backcrosses as well as other crosses by the same sire, it was therefore most likely outbreeding depression. If co-adapted gene complexes are disrupted

in the Sebago x LaHave hybrids, the effects of outbreeding depression would more likely be observed in the F<sub>2</sub> generation if it existed.

It is also possible that the environmental setting of the experiment made for a lack of outbreeding depression. Benign hatchery environments mostly test intrinsic factors (Tymchuk et al. 2007), that is the effect caused by the genotype. However, the phenotype is the product of an interaction between genotype and the environment, it is believed that extrinsic factors such as the loss of local adaptations have a stronger influence over outbreeding depression than do intrinsic factors such as the disruption of co-adapted gene complexes for species with many unique and highly local populations (Allendorf et al. 2001; Edmands and Timmerman 2003; reviewed in Garcia de Leaniz et al. 2007; Tymchuk et al. 2007; Vandersteen et al. 2012). In the F<sub>1</sub> generation, outbreeding depression is also more likely to be due to extrinsic factors than intrinsic ones (Lynch 1991, Edmands 2007). As the environmental surroundings will affect the capability for detecting both outbreeding depression and heterosis (Burton 1987; Edmands 2007; Tymchuk et al. 2007; Vandersteen et al. 2012; Crespel et al. 2013), it is therefore possible that any outbreeding depression resulting from the loss of local adaptations was negated by a relaxed environmental setting (Tymchuk et al. 2007). For example, in Tymchuk et al. (2007), F<sub>3</sub> hybrid rainbow trout displayed outbreeding depression related to growth and survival in certain environmental surroundings; as a result, the cause of the outbreeding depression is thought to be primarily the result of the loss of local adaptations that was simply expedited by intrinsic factors. In Vandersteen et al. (2012), it was found that survival of rainbow trout fry differed not only in the geographic area but also seasonally, as the survival of different genotypes varied between summer and winter.

Another study on Brook trout (*Salvelinus fontinalis*) by Crespel et al. (2013) found environmental interactions affected the presence of heterosis in certain hybrid crosses but not others, also suggesting that environmental influence has a varying effect on different genotypes. Therefore, in the case of this study, it is possible that if the juveniles had lost a local adaptation of importance, the outbreeding depression might have only been capable of being detected in more challenging natural settings. If one of the populations being used in the restoration effort is indigenous to the environment, the addition of a second phenotype may cause the offspring to be less suited for the environment compared to the local population, and may hinder the reintroduction effort rather than assist it. In such cases, it would be worth choosing populations with similar local adaptations in order to avoid an intermediate phenotype.

As this study examined the early life stage of Atlantic salmon, another potential reason for the lack of detection of outbreeding depression or heterosis are maternal effects. These environmental or genetic effects can influence the offspring's phenotype on a per subject basis in early life stages by overshadowing the influence of the offspring's genotype (reviewed by Wolf and Wade 2009). Therefore, the phenotypic expression of outbreeding depression can be outweighed by maternal effects during early life stages (Edmands 2007). Previous studies, have demonstrated hybrid fitness has been shown to vary as a result of maternal effects (McGinnity et al. 2003; Debes et al. 2013). McGinnity et al. (2003) found maternal effects affected the survival of egg stage as well as eyed egg stage in the early juvenile stage of farmed x wild Atlantic salmon hybrids, and Debes et al. (2013) found that the actual genotype of the hybrid wild-farm offspring had little effect on survival, whereas maternal effects accounted for almost all of the variance in

survival between hybrid types. As our study found significant dam ID effects (i.e. the effect of the inherited maternal genotype as well as maternal effects) on the egg survival and early size measurements, it is possible that maternal effects are masking interaction effects that may have otherwise lead to outbreeding depression later in life. For example, Heath et al. (1999) found that juvenile growth in Chinook salmon was predominantly the result of maternal effects, however, as the individuals matured the effect of any maternal effect on the offspring's phenotype decreased. Another study by Houde et al. (2015) on the Sebago and LaHave populations of Atlantic salmon supported such findings, with juvenile survival and fitness being primarily influenced by maternal environmental effects during the egg stage and that phenotypic influence was then primarily controlled for by nonadditive genetic effects in the fry stage. It is important to mention the egg survival in my study (~50%) was much lower than those typically seen in other studies involving Atlantic salmon (84%) (e.g. Taranger and Hansen 1993) or other salmonids (~90%) (e.g. Cho et al. 2002). It is possible that the low egg survival seen in this study was the result of environmental settings such as the water temperature. For example, Taranger et al. (1993) found that Atlantic salmon eggs that were in warm water conditions (13-14°C) had a 15.5% lower survival than those placed in cold water (5-7°C) and a 7.9% lower survival than the control (8-10°C). As our water temperature typically ranged from 9-12°C, it is possible that the warmer water temperatures were not favorable for the Atlantic salmon eggs. However, as the egg survival in our study were consistent with the egg survival in other studies using these broodstocks of Atlantic salmon (Houde et al. 2015) and that they were all exposed to the same water conditions throughout the experiment, it is assumed that the low egg survival did not influence the outcome of the results. Therefore, it would

demonstrate the importance of testing lifelong fitness of the hybrids to insure that heterosis or outbreeding depression is not being masked by maternal effects in the study.

Another possibility is that the neutral genetic distance between the parental populations (i.e. LaHave and Sebago) was not great enough to invoke any phenotypic expression of outbreeding depression or heterosis, as the occurrence and severity of both has previously been shown to be positively correlated with parental divergence (Edmands and Timmerman 2003; Edmands 2007; McClelland and Naish 2007; Fraser et al. 2010). Although the Sebago and LaHave populations have been isolated from one another for approximately a thousand generations, the neutral genetic distance between the two populations is relatively low (as measured by  $F_{ST}=0.038$ , He et al. 2015). It is therefore possible that the rate at which these populations of Atlantic salmon are diverging is not rapid enough to observe any form of heterosis or outbreeding depression. Makinen et al. (2014) detected low genetic divergence between domesticated populations of Atlantic salmon ( $F_{ST}<0.03$ ). Although factors such as the fact that Atlantic salmon have been under domestication for over 10 generations (Hutchings and Fraser 2008), as well as the polygenic nature of most selected traits may have led to the lack in findings of divergent artificial selection (Makinen et al. 2014), other studies have also suggested that selection in a hatchery environment may cause convergent selection on the same genes between populations of salmonids with 5 to 7 generations of selection and 4 generations respectively (Roberge et al. 2006; Sauvage et al. 2010). As both of the populations in this study have been domesticated for several generations, it is possible that since being in hatchery settings, their broodstocks have been under convergent natural selection for domesticated environments.



In summary, the simultaneous use of both the Sebago and LaHave populations does not appear to result in the expression of outbreeding depression or heterosis during the juvenile life stages of the  $F_1$  generation. Future research should investigate adult fecundity and reproductive fitness (gamete quality and survival) of  $F_1$  hybrids, as well as potential evidence of outbreeding depression in  $F_2$ , backcrosses and subsequent generations. As with any reintroduction which uses multiple populations simultaneously, hybridization when it results in outbreeding depression remains a potential hindrance for conservation efforts. Therefore, the use of multiple allopatric populations should be done with caution after research has gone into studying the full effects hybridization may have.

## **Acknowledgements**

This study was funded by the NSERC Strategic Project Grant (to TEP and CCW), NSERC Discovery Grant (to TEP), CFI (to TEP), the Ontario Ministry of Economic Development and Innovation (to TEP) as well as the Ontario Ministry of Natural Resources and Forestry (MNRF). We would like to thank Bill Sloan and Scott Ferguson from the MNRF Codrington Fisheries Research Facility for providing technical expertise, Craig Black for helping to initiate this research and Katelynn Johnson for helping to produce the crosses. We would also like to thank the Great Lakes Fish Research Centre in LaSalle for providing the logistical support in order to conduct the research as well as Aleksa Zubic, Sumeet Bhardwaj and Jade Laycock for helping with the maintenance or care of the fish during the rearing stage. Lastly, we would like to thank my colleagues and lab members Jason Lewis, Adriana Forest and Sarah Lehnert for their assistance and support during this project.

## References

- Allendorf FW, Thorgaard GH (1984) Tetraploidy and the evolution of salmonid fishes. In Turner B (ed) Evolutionary Genetics of Fishes. Plenum Press, New York. pp.1-53
- Araki H, Cooper B, Blouin MS (2007) Genetic Effects of Captive Breeding Cause a Rapid, Cumulative Fitness Decline in the Wild. *Science* 318: 100-103
- Bates D, Maechler M, Bolker B and Walker S (2014) lme4: Linear mixed-effects models using Eigen and S4. submitted to *Journal of Statistical Software*
- Birchler JA, Auger DL, Riddle NC (2003) In search of the molecular basis of heterosis. *Plant Cell* 15: 2236-2239
- Burton RS (1987) Differentiation and Integration of the Genome in Populations of the Marine Copepod *Tigriopus californicus*. *Evolution* 41: 504-513
- Cho GK, Heath JW, Heath DD (2002) Electroshocking Influences Chinook Salmon Egg Survival and Juvenile Physiology and Immunology. *T Am Fish Soc* 131:224-233
- Crespel A, Audet C, Bernatchez L, Garant D (2013) Effects of rearing environmental and strain combination on hybrid vigour in brook trout. *N Am J Aquacult* 74: 188-198
- Debes PV, Fraser DJ, McBride MC, Hutchings JA (2013) Multigenerational hybridisation and its consequences for maternal effects in Atlantic salmon. *Heredity* 111: 238-247
- Dimond P, Smitka J (2005) Evaluation of Selected Strains of Atlantic Salmon as Potential Candidates for the Restoration of Lake Ontario. Trout Unlimited Canada Technical Report ON-12. Guelph, Ontario: Trout Unlimited Canada

- Dobzhansky T (1948) Genetic of natural populations. XVIII. Experiments on chromosomes of *Drosophila pseudoobscura* from different geographical regions. Genetics 33:588-602
- Edmands S (1999) Hybrid vigour and outbreeding depression in interpopulation crosses spanning a wide range of divergence. Evolution 53: 1757-1768
- Edmands S (2002) Does parental divergence predict reproductive compatibility? Trends Ecol Evol 17: 520-526
- Edmands S, Timmerman CC (2003) Modeling factors affecting the severity of outbreeding depression. Conserv Biol 17: 883-892
- Edmands S (2007) Between a rock and a hard place: evaluating the relative risks of inbreeding and outbreeding for conservation and management. Mol Ecol 16: 463-475
- Frankham R, Ballou JD, Briscoe DA (2002) Introduction to Conservation Genetics. Cambridge University Press, UK.
- Frankham R, Ballou JD, Eldridge MDB, Lacy RC, Ralls K, Dudash MR and Fenster CB (2011) Predicting the Probability of Outbreeding Depression. Conserv Biol 25: 465-475
- Fraser DJ, Houde ALS, Debes PV, O'Reilly P, Eddington JD and Hutchings JA (2010) Consequences of farmed-wild hybridization across divergent wild populations and multiple traits in salmon. Ecol Appl 20: 935-953
- Garcia de Leaniz C, Fleming IA, Einum S, Verspoor E, Jordan WC, Consuegra S, Aubin-Horth N, Lajus D, Letcher BH, Youngson AF, Webb JH, Vøllestad LA, Villanueva B,

Ferguson A, Quinn TP (2007) A critical review of adaptive genetic variation in Atlantic salmon: implications for Conservation. *Biol Rev* 82: 173-211

Gilk SE, Wang IA, Hoover CL, Smoker WW, Taylor SG, Gray AK, Gharrett AJ (2004) Outbreeding depression in hybrids between spatially separated pink salmon, *Oncorhynchus gorbuscha*, populations: marine survival, homing ability, and variability in family size. *Environ Biol Fishes* 69: 287-297

Granier S, Audet C, Bernatchez L (2011) Hybrid vigour and outbreeding depression between strains of young-of-the-year brook trout (*Salvelinus fontinalis*). *Can J Zool* 89: 190-198

Halekoh U, Højsgaard S (2014) A Kenward-Roger Approximation and Parametric Bootstrap Methods for Tests in Linear Mixed Models - The R Package pbkrtest. *Journal of Statistical Software* 59: 1-30.

Harrison RG (1990) Hybrid zones: windows on evolutionary processes. In: Futuyma, D. and Antonovics, J. (ed) *Oxford Surveys in Evolutionary Biology*. Oxford University Press, Oxford. pp:69-128

He X, Wilson CC, Wellband KW, Houde ALS, Neff BD, Heath DD. 2015. Transcriptional profiling of two Atlantic salmon strains: implications for reintroduction into Lake Ontario. *Conserv Genet* 16: 277-287

Heath DD, Fox CW, Heath JW (1999) Maternal Effects on Offspring Size: Variation Through Early Development of Chinook Salmon. *Evolution* 53: 1605-1611

- Houde AL, Fraser DJ, O'Reilly P, Hutchings JA (2011) Relative risks of inbreeding and outbreeding depression in the wild in endangered salmon. *Evol Appl* 4: 634-647
- Houde ALS, Black CA, Wilson CC, Pitcher TE and Neff BD (2015) Genetic and maternal effects on juvenile survival and fitness-related traits in three populations of Atlantic salmon. *Can J Fish Aquat Sci* 72:751-758
- Hoysak DJ, Liley NR (2001) Fertilization dynamics in sockeye salmon and a comparison of sperm from alternative male phenotypes. *J Fish Biol* 58: 1286–1300
- Hutchings JA, DJ Fraser (2008) The nature of fisheries and farming-induced evolution. *Mol Ecol* 17: 294-313
- Lynch M (1991) The Genetic Interpretation of Inbreeding Depression and Outbreeding Depression. *Evolution* 45: 622-629
- MacCrimmon HR (1977) *Animals, man and change: alien and extinct wildlife of Ontario*. McClelland and Stewart, Toronto.
- Makinen H, Vasemagi A, McGinnity P, Cross T, Primmer C (2014) Population genomic analyses of early-phase Atlantic Salmon (*Salmo salar*) domestication/captive breeding. *Evol Appl* 8: 93-107
- McClelland EK, Myers JM, Hard JJ, Park LK, Naish KA (2005) Two generations of outbreeding in coho salmon (*Oncorhynchus kisutch*): effects on size and growth. *Can J Fish Aquat Sci* 62: 2538-2547

McClelland EK, Naish KA (2007) What is the fitness outcome of crossing unrelated fish populations? A meta-analysis and an evaluation of future research directions. *Conserv Genet* 8: 397-416

McGinnity P, Prodöhl P, Ferguson A, Hynes R, Maoiléidigh NO, Baker N, Cotter D, O’Hea B, Cooke D, Rogan G, Taggart J, Cross T (2003) Fitness Reduction and Potential Extinction of Wild Populations of Atlantic Salmon, *Salmo salar*, as a Result of Interactions with Escaped Farm Salmon. *Proceedings: Biological Sciences* 270: 2443-2450

Naish KA, Hard JJ (2008) Bridging the gap between the genotype and the phenotype: linking genetic variation, selection and adaptation in fishes. *Fish Fish* 9: 396-422

Neff BD, Garner SR, Pitcher TE (2011) Conservation and enhancement of wild fish populations: preserving genetic quality versus genetic diversity. *Can J Fish Aquat Sci.* 68: 1139-1154

Ontario Ministry of Natural Resources (2004) Lake Ontario Fish Communities and Fisheries: 2003 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.

Ontario Ministry of Natural Resources (2006) Lake Ontario Fish Communities and Fisheries: 2005 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.

Ontario Ministry of Natural Resources (2009) Lake Ontario Fish Communities and Fisheries: 2008 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.

Pitcher TE, Neff BD (2006) MHC class IIB alleles contribute to both additive and non-additive genetic effects on survival in Chinook salmon. *Mol Ecol* 15: 2357-2365

Pitcher TE, Neff BD (2007) Genetic quality and offspring performance in chinook salmon: Implications for supportive breeding. *Conserv Genet* 8: 607-616

Ricker WE (1975) Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 191: 1-382

Roberge C, Einum S, Guderley H, Bernatchez L (2006) Rapid parallel evolutionary changes of gene transcription profiles in farmed Atlantic salmon. *Mol Ecol* 15:9-20

Rollinson N, Keith DM, Houde ALS, Debes PV, McBride MC, Hutchings JA (2014) Risk Assessment of Inbreeding and Outbreeding Depression in a Captive-Breeding Program. *Conserv Biol* 28: 529-540

Sauvage C, Derôme N, Normandeau E, St.-Cyr J, Audet C, Bernatchez L (2010) Fast Transcriptional Responses to Domestication in the Brook Charr *Salvelinus fontinalis*. *Genetics* 185: 105-112

Stanfield L, Jones ML (2003) Factors Influencing Rearing Success of Atlantic Salmon Stocked as Fry and Parr in Lake Ontario Tributaries. *N Am J Fish Sci* 23:1175-1183

Taranger GL, Hansen T (1993) Ovulation and egg survival following exposure of Atlantic salmon, *Salmo salar* L., broodstock to different water temperatures. *Aquacult Fish Manage* 24:151-156



Templeton AR (1986) Coadaptation and outbreeding depression. In Soulé M (ed) Conservation biology: the science of scarcity and diversity. Sinauer, Sunderland, MA. pp. 105-116

Thorstad EB, Whoriskey F, Rikardsen AH, Aarestrup K (2010) Aquatic Nomads: The Life and Migrations of the Atlantic Salmon. In Atlantic Salmon Ecology, First Edition. Wiley-Blackwell Publishing. Hoboken, NJ. pp.1-23

Toivonen J (1971) The fish fauna and limnology of large oligotrophic glacial lakes in Europe (about 1800 A.D.). J Fish Res Board Can 29: 629-637

Tymchuk WE, Sundstrom LF, Devlin RH (2007) Growth and Survival Trade-offs and outbreeding depression in rainbow trout (*Oncorhynchus mykiss*). Evolution 61:1225-1237

Vandersteen W, Biro P, Harris L, Devlin R (2012) Introgression of domesticated alleles into a wild trout genotype and the impact on seasonal survival in natural lakes. Evol Appl 5: 76-88

Ward HB (1932) The Origin of the Landlocked Habit in Salmon. Proc Nat Acad Sc 18: 569-580

Wolf JB, Wade MJ (2009) What Are Maternal Effects (And What Are They Not)? Philosophical Transactions: Biological Sciences 364: 1107-1115

Table 2.1: Mean percentage of survival of F<sub>1</sub> cross types of Atlantic salmon (*Salmo salar*) during four life stages with the significance (p) of cross type (“Cross”) as fixed effect as well as dam, sire and their interaction as random effects to the variance of the fitness characteristic (see methods for details).

| Survival (%)     | Pure Crosses |       | Hybrids |       | Significance     |       |                  |       |
|------------------|--------------|-------|---------|-------|------------------|-------|------------------|-------|
|                  | LL           | SS    | LS      | SL    | Dam              | Sire  | Interaction      | Cross |
| Egg Stage        | 42.26        | 48.55 | 45.32   | 49.28 | <b>&lt;0.001</b> | >0.99 | >0.99            | 0.96  |
| Fry Stage        | 71.23        | 75.14 | 79.36   | 76.43 | 0.620            | 0.20  | <b>&lt;0.001</b> | 0.47  |
| Early Parr Stage | 94.98        | 93.44 | 95.89   | 98.57 | <b>0.02</b>      | >0.99 | >0.99            | 0.25  |
| Late Parr Stage  | 92.44        | 94.38 | 90.62   | 90.42 | >0.99            | 0.31  | >0.99            | 0.90  |

Significant fixed and random effects ( $p < 0.05$ ) are indicated in bold. Original egg number ( $p=0.44$ ) was used in the place of density during the egg stage. Stack, tray and cell effects were also kept in the model as they were significant ( $p<0.001$ ). The cells were kept as separate entities when calculating the mean survival %.

**Table 2.2** Means ( $\pm$  standard error) of F<sub>1</sub> cross types of Atlantic salmon (*Salmo salar*) with the significance (p) of cross type (“Cross”) as fixed effect as well as dam, sire and their interaction as random effects to the phenotypic variance of the fitness characteristic (see methods for details).

| Characteristic (units) | Pure Crosses      |                   | Hybrids          |                    | Significance (p) |        | Dam   | Sire | Interaction | Cross |
|------------------------|-------------------|-------------------|------------------|--------------------|------------------|--------|-------|------|-------------|-------|
|                        | LL                | SS                | LS               | SL                 |                  |        |       |      |             |       |
| Fork length (cm)       |                   |                   |                  |                    |                  |        |       |      |             |       |
| April 2013(button-up)  | 35.5 $\pm$ 0.230  | 35.9 $\pm$ 0.22   | 35.5 $\pm$ 0.23  | 35.8 $\pm$ 0.22    | <0.001           | <0.001 | >0.99 | 0.99 |             |       |
| August 2013(fry)       | 61.5 $\pm$ 0.473  | 62.9 $\pm$ 0.42   | 60.1 $\pm$ 0.41  | 60.6 $\pm$ 0.40    | <0.001           | <0.001 | >0.99 | 0.41 |             |       |
| October 2013(parr)     | 82.7 $\pm$ 1.13   | 84.1 $\pm$ 1.22   | 79.7 $\pm$ 0.98  | 79.5 $\pm$ 0.97    | >0.99            | >0.99  | >0.99 | 0.57 |             |       |
| December 2013(parr)    | 90.6 $\pm$ 1.39   | 92.8 $\pm$ 1.57   | 88.3 $\pm$ 1.29  | 88.2 $\pm$ 1.14    | >0.99            | >0.99  | >0.99 | 0.84 |             |       |
| February 2014(smolt)   | 104.7 $\pm$ 1.47  | 107.3 $\pm$ 1.80  | 104.0 $\pm$ 1.42 | 102.4 $\pm$ 1.34   | >0.99            | >0.99  | >0.99 | 0.67 |             |       |
| Mass (g)               |                   |                   |                  |                    |                  |        |       |      |             |       |
| April 2013(button-up)  | 0.46 $\pm$ 0.010  | 0.48 $\pm$ 0.010  | 0.47 $\pm$ 0.010 | 0.48 $\pm$ 0.00978 | <0.001           | 0.01   | >0.99 | 0.91 |             |       |
| August 2013(fry)       | 2.81 $\pm$ 0.065  | 2.96 $\pm$ 0.058  | 2.55 $\pm$ 0.048 | 2.63 $\pm$ 0.0502  | 0.002            | 0.004  | >0.99 | 0.09 |             |       |
| October 2013(parr)     | 6.02 $\pm$ 0.283  | 6.55 $\pm$ 0.298  | 5.39 $\pm$ 0.22  | 5.45 $\pm$ 0.218   | >0.99            | >0.99  | >0.99 | 0.59 |             |       |
| December 2013(parr)    | 9.14 $\pm$ 0.484  | 9.81 $\pm$ 0.523  | 8.40 $\pm$ 0.40  | 8.10 $\pm$ 0.362   | >0.99            | >0.99  | >0.99 | 0.93 |             |       |
| February 2014(smolt)   | 13.50 $\pm$ 0.596 | 14.64 $\pm$ 0.745 | 13.06 $\pm$ 0.53 | 12.19 $\pm$ 0.507  | >0.99            | >0.99  | >0.99 | 0.66 |             |       |

Condition ( $10\,000 \times \text{g}/\text{mm}^3$ )

|                       |                      |                     |                     |                     |                  |                  |                  |      |
|-----------------------|----------------------|---------------------|---------------------|---------------------|------------------|------------------|------------------|------|
| April 2013(button-up) | $0.099 \pm 0.000627$ | $0.101 \pm 0.00053$ | $0.100 \pm 0.00065$ | $0.100 \pm 0.00064$ | <b>0.02</b>      | >0.99            | <b>&lt;0.001</b> | 0.95 |
| August 2013(fry)      | $0.117 \pm 0.000942$ | $0.117 \pm 0.00074$ | $0.115 \pm 0.00081$ | $0.116 \pm 0.00101$ | 0.21             | 0.49             | <b>&lt;0.001</b> | 0.86 |
| October 2013(parr)    | $0.099 \pm 0.00102$  | $0.104 \pm 0.00095$ | $0.101 \pm 0.00083$ | $0.103 \pm 0.00088$ | <b>&lt;0.001</b> | <b>&lt;0.001</b> | >0.99            | 0.74 |
| December 2013(parr)   | $0.112 \pm 0.000628$ | $0.113 \pm 0.00060$ | $0.113 \pm 0.00050$ | $0.111 \pm 0.00062$ | 0.31             | <b>0.006</b>     | 0.19             | 0.30 |
| February 2014(smolt)  | $0.111 \pm 0.000702$ | $0.110 \pm 0.00073$ | $0.110 \pm 0.00070$ | $0.107 \pm 0.00055$ | <b>0.02</b>      | 0.10             | 0.22             | 0.23 |

---

Significant fixed and random effects ( $p < 0.05$ ) are indicated in bold. Density was  $p < 0.05$  for all measurements with the exception of fork length April 2013 ( $p=0.19$ ) and condition (August  $p=0.55$ , December 0.38 and February 0.06)

## CHAPTER 3: HETEROZYGOSITY AND GAMETE QUALITY TRAITS IN HATCHERY-REARED ATLANTIC SALMON (*SALMO SALAR*)

### SYNOPSIS

This chapter examined the relationship between gamete quality traits and multi-locus heterozygosity in male and female hatchery-reared Atlantic salmon (*Salmo salar*). Multi-locus heterozygosity (MLH; based on 19 polymorphic microsatellite markers) was correlated with sperm (including velocity, linearity, density, longevity and motility) and egg (including egg diameter, wet egg mass, dry egg mass, fecundity) quality metrics as well as Fulton's condition factor. For females, there was no significant relationship between MLH and any of the gamete quality metrics. For males, although significant negative correlations were found between MLH and sperm velocity traits and longevity traits, the correlation explained limited variance. Also, there was no significant correlation between sperm linearity, motility nor density and MLH. Neither sex displayed any significant correlation between Fulton's condition factor and MLH. Although negative correlations between certain sperm quality traits were found, overall there was little evidence to support the existence of heterozygosity fitness correlations between gamete quality traits potentially due to a lack of variability in the inbreeding coefficient or indicative of a lack of heterozygosity fitness correlations with gamete quality traits in Atlantic salmon.

## **Introduction**

Heterozygosity fitness correlations (HFC) are statistical correlations between an individual measure of heterozygosity (e.g. multi-locus heterozygosity (MLH)) and traits that are related to fitness (reviewed in Chapman et al. 2009; Szulkin et al. 2010), and they have become a widely utilized tool in evolutionary ecology (Mitton 1993; Coltman and Slate 2003; David 1998; reviewed in Chapman et al. 2009; reviewed in Szulkin et al. 2010). Most HFC studies are currently carried out with the use of polymorphic microsatellites, which are predominantly considered to be neutral markers (Jarne and Lagoda 1996). The occurrence of HFCs in neutral markers is explained by what are known as local and general effects. Local effects occur when a neutral marker reflects overdominance of loci under selection within their chromosomal vicinity as a result of linkage disequilibrium (LD) (David 1998; Lynch and Walsh 1998, reviewed in Hansson and Westerberg 2002). General effects occur as the result of the heterozygosity at the markers being reflective of the individual's overall heterozygosity, and as a result of being less inbred, heterozygotes will have a higher fitness than homozygotes as a result of being less inbred) (David 1998; Lynch and Walsh 1998; reviewed in Hansson and Westerberg 2002; Coltman and Slate 2003).

Based on the general effects hypothesis, theory suggests that if HFCs are the result of heterozygotes being less inbred than homozygotes, HFCs would be more prevalent in populations that have high variance in the values of inbreeding coefficients (David 1998; reviewed in Coltman and Slate 2003) due to a higher amount of additive genetic variance within such populations. However, an empirical review by Chapman et al. (2009) found no evidence of HFCs being more common in populations with high variance in

inbreeding coefficients, suggesting that certain studies were finding local effects rather than general ones or the presence of publication bias towards significant results in all populations. Traits that are more closely associated with fitness (e.g. life history traits), or those that are affected by multiple loci also usually show stronger associations in HFC studies compared to traits that are not strongly associated with fitness (Coltman and Slate 2003; reviewed Chapman et al. 2009; reviewed in Szulkin et al. 2010). As spermatogenesis and oogenesis are extremely sensitive to selection pressures, it is expected that gamete quality will be one of the first traits to reveal evidence of correlations with heterozygosity (Gage et al. 2006; Fitzpatrick et al. 2009). Although many studies have examined the relationship between heterozygosity and life history traits (reviewed in Chapman et al. 2009; reviewed in Szulkin et al. 2010), few studies have examined the relationship between heterozygosity and gamete quality (see Table 3.1 for a summary of studies).

There are several characteristics adapted by spermatozoa to increase chances of fertilization and are therefore representative of sperm quality such as motility, longevity (Gage et al. 1995), velocity (Gage et al. 2004) and density (reviewed in Snook 2005). Heterozygosity has been found to correlate positively with sperm quality within (Gage et al. 2006) and among (Fitzpatrick and Evans 2009) species. For example, Gage et al. (2006) found a negative correlation between mean heterozygosity and the production of abnormal sperm within and across wild rabbit populations (*Oryctolagus cuniculus*) using 29 microsatellite loci. The strongest correlations were found within the most fragmented populations, suggesting a possible relationship to inbreeding (Gage et al. 2006). Other studies have also found heterozygosity to be negatively correlated with abnormal sperm

percentage and positively associated with motile sperm in endangered mammals (Fitzpatrick and Evans 2009).

Just as with sperm, egg quality is also important for offspring fitness. Egg quality is important for survival as it is the only source of energy the embryo receives until it reaches the exogenous feeding stage (Pickova et al. 1997). Therefore, the quality such as the volume, mass and fecundity are important to an individual's overall fitness as they reflect the energy content the individual is allocating to offspring (Srivastava and Brown 1991). Heterozygosity has been found to correlate with egg quality and quantity in multiple studies across a variety of taxa (see Table 3.1 for summary of studies) (e.g. Heath et al. 2002; Garcia-Navas et al. 2009; Wetzel et al. 2012). For example, heterozygosity was positively correlated with clutch size in house sparrows (*Passer domesticus*) (Wetzel et al. 2012). Another study found correlation between heterozygosity and clutch size as well as egg pigmentation pattern in blue-tits (*Cyanistes caeruleus*) (Garcia-Navas et al. 2009).

Although gamete quality characteristics are well documented in Atlantic salmon (Gage et al. 1995; 1998; 2004; Lush et al. 2014), there is a lack of studies that have evaluated their correlation to heterozygosity. Atlantic salmon (*Salmo salar*) are an externally fertilizing fish species with a distribution across eastern North America (Holm et al. 2009). There are many characteristics adapted by gametes in Atlantic salmon in order to optimize their chances of fertilization success as well as hatching success. For example, sperm velocity has been shown to be an important characteristic for fertilization success (Gage et al. 1995, 2004). It is also well documented that individuals emerging from larger eggs have higher survival than individuals emerging from smaller eggs



(Einum and Fleming 2000), suggesting egg diameter and egg mass as important characteristics to embryo health and survival. As many salmonid populations, including Atlantic salmon, are currently being supported by hatcheries for restoration and reintroduction, the gamete quality of these hatchery reared fish and their offspring is crucial for future health of these populations. However, it has been demonstrated that the lack of selection that hatcheries pose cause divergence in the evolution of gamete quality selection (e.g. Heath et al. 2002). In salmonids, Coho salmon (*Oncorhynchus kisutch*) reared in hatchery settings displayed lower than average reproductive success in the wild than their wild counterparts (Thériault et al. 2011). Kekäläinen et al. (2013) found that sperm motility in Arctic charr was lower in hatchery populations than their wild counterparts. As there is currently a reintroduction effort taking place for Atlantic salmon in Lake Ontario, the gamete quality of the broodstock is an important characteristic for the success of the initiative.

The objective of this chapter is to examine and describe the existence of correlations between gamete quality and MLH through either local or general effects using a population of captively reared Atlantic salmon as a case study. The LaHave population is the most common strain currently used in Atlantic salmon stocking efforts in Lake Ontario and originates from the LaHave River in Nova-Scotia (Stanfield and Jones 2003), and has been bred in captivity for several generations using rotational line crosses (RLC) a breeding technique that exclusively breeds different year classes in order to avoid inbreeding. As this population is the main broodstock for reintroduction efforts around the province (MNRF 2009), it would be of use to know if genetic variance can explain chances at reproductive success. If HFCs do not exist in the LaHave broodstock

of Atlantic salmon, I would expect no correlation between the gamete quality traits and heterozygosity. However, if HFCs do exist in the gamete quality of the LaHave broodstock, than I would expect a positive correlation between the gamete quality traits and heterozygosity.

## **Methods**

### *Broodstock*

All Atlantic salmon used in this study were from the LaHave broodstock maintained at the Ontario Ministry of Natural Resources and Forestry (MNR) Harwood Fish Culture Station (44° 8'13.31"N 78°10'5.59"W). The broodstock was created using a breeding technique known as rotational line crosses (RLC) in which unrelated broodstock are crossed in order to create 3 new lines to be used in rotation (see Figure 3.1; for further information on RLC see Kincaid 1977). The female broodstock were composed of 3 year classes generated in 2007, 2008 and 2009 from a year class at MNR Normandale Fish Culture Station whose parents had been created from wild gametes. The male broodstock had two year classes created in 2007 and 2008 also generated from the same Normandale Fish Culture Station year class.

### *Milt Collection and Sperm Quality Assessment*

Males were anesthetized using MS-222 (~0.110 g/L), a weight ( $\pm 1.0$  g) and a fork length ( $\pm 1.0$  mm) were recorded and milt samples were collected from 38 individuals (aged 6 years (n=29) and 7 years (n=9)). In order to avoid contamination from water or mucus, the area extending from the individual's pelvic fins to the anal fin was thoroughly dried before the individual was stripped. Milt was then collected by gently applying

pressure to the abdomen of the individual and collected in 532 mL Whirl-Pak bags (Nasco, Newmarket, Ontario). The bags were then placed in a cooler until analysis 1-2 hours after collection.

In order to estimate sperm motility traits, sperm activity was recorded using a video camera (XC-ST50, Sony, Japan) mounted on a microscope (CX41 Olympus, Melville, New York) through a 10x objective (as per Pitcher et al. 2009). An aliquot of milt was placed on the edge of a chamber on a 2X-CEL glass slide (Hamilton Thorne, Massachusetts) and activated using 10  $\mu$ L of hatchery water maintained at 7°C using a Bionomic controller (model BC-110, 20/20 Technology Inc., Wilmington, North Carolina). The HTM-CEROS sperm analysis system was used to analyze the selected sperm traits (i.e. path velocity (VAP), curvilinear velocity (VCL), progressive velocity (VSL) Linearity (i.e. the degree of straightness per unit of distance traveled by a sperm cell), % motility (i.e. the percentage of motile spermatozoa in the recording over the number of total sperm cells in the recording) and longevity (i.e. the point in time during the recording in which 95 % of the cells were immobile)). The settings of the system were placed at the following: Image capture was set at 60 frames/sec, cell detection was set at a contrast of 13 and a minimum cell size of 3 pixels, the cell intensity defaults were set at 3-50, the progressive cell settings were set at a VAP of 26.0 and STR of 80.0 and the slow cell cut-off was set at 20.0-20.0. Percent motility was analyzed 5s post activation, whereas, velocity and linearity were analyzed at 5 seconds as well as 10 seconds post activation. Five and 10 seconds post-activation were used as the standard times because according to the findings of Liley et al. (2002) fertilization success declines after the 10 second time point. The reason percent motility was not analyzed at 10 seconds or 15

seconds post activation and velocity and linearity were not analyzed at 15 seconds post activation was due to the poor correlation coefficients ( $r^2 < 0.215$ ) for the different tracks taken by the same male, which would have increased error in the measuring of the metric. The mean value of all recordings per male was used as the value in statistical analysis. Two recordings of sperm activity were performed for every male (with the exception of 3 individuals of which only 1 recording was collected) and the recordings were analyzed (see below) and averaged to insure consistency between measured traits.

To estimate sperm density, an aliquot of 1.5  $\mu\text{L}$  of milt was pipetted into a mixture of 125 mL of gluteraldehyde and 500  $\mu\text{L}$  of Cortland's saline solution (7.25 g/L NaCl; 0.38 g/L KCl; 0.47 g/L  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ; 0.4 g/L  $\text{Na}_2\text{HPO}_4 \cdot \text{H}_2\text{O}$ ; 1.0 g/L  $\text{NaHCO}_3$ ; 0.22 g/L  $\text{MgCl}_2$ ; 1.0 g/L  $\text{C}_6\text{H}_{12}\text{O}_6$ ). The mixture was kept in a cooler until it was transported to a refrigerator (4°C) until analysis could be performed (within 24h). 10  $\mu\text{L}$  of the solution was pipetted onto a haemocytometer and left to sit for ~10 minutes to insure that all the sperm cells had settled. The haemocytometer was placed onto a microscope (CX41 Olympus, Melville, New York) and recorded using a negative contrast 20x objective. The cells were counted using a standardized procedure, where the cells in each of the 4 corner squares as well as the centre square was counted. The mean of the 5 counted squares was then multiplied by 25 to represent to total number of squares and then by 10. Then the value is multiplied by the volume of the diluted milt solution (625  $\mu\text{L}$ ) to obtain a value to the number of sperm cells estimated to be in 1mL of milt (as per Pitcher et al. 2009a).

### *Egg Collection and Quality Assessment*

Females were anaesthetized using MS-222 (~0.110 g/L), a weight ( $\pm 1.0$  g) and a fork length ( $\pm 1.0$  mm) were recorded and eggs were collected from 41 individuals (aged 5 years (n=9), 6 years (n=20), and 7 years (n=12)) Manual pressure was then placed on the individual's abdomen and eggs were then collected in 532 mL whirl-pak bags and placed in a cooler until wet weight was measured between 2-3 hours after collection.

To estimate the quantity of eggs that each female produced, the volume of eggs that each female produced was measured in litres, following which a count was done on a subsample and multiplied by the total volume to calculate the fecundity of each female. To estimate wet egg mass, eggs were placed into a strainer to remove ovarian fluid and between 20-71 eggs (MEAN  $\pm$  S.D. =  $38.64 \pm 10.64$ ) were placed in weigh boats and weighed with an electronic scale (Denver Instrument TP 323) in order to calculate the mass of each individual egg ( $\pm 0.001$  g) and a digital image taken. Three replicates were done per female. The egg diameter of 10 eggs was measured in each of the 3 replicates using Image J (<http://imagej.nih.gov/ij/>) to get an average diameter of each egg ( $\pm 0.1$  mm). 15 mL of eggs were then stored in falcon tubes and placed on dry ice until they were transported to a  $-80^{\circ}\text{C}$  freezer where they were stored until the dry weight was measured. In order to estimate dry egg mass, eggs were removed from the  $-80^{\circ}\text{C}$  freezer and left to thaw for ~2-3 hours before 10 eggs were counted and placed in aluminum foil weighing dishes. Three replicates were done per female. The eggs were then placed in a dry oven for 24h at  $55^{\circ}\text{C}$  (Einum and Fleming 2000), followed by a desiccator for 12 hours to insure that all moisture had been removed before weighing ( $\pm 0.0001$  g) with an electronic scale (Denver Instrument SI 234).

### *Heterozygosity*

The MLH of individual males and females was calculated using 19 different microsatellite loci, with a polymorphic range of 7-48 alleles (see Table 3.2 for further information on each locus and Figures 3.2 and 3.3 for the distribution of MLH in both males and females). Amplification was done for the following 6UM of forward and reverse primer 6 UM dNTP, 30 BSA ug/ml, 60 µL of Promega buffer, 1.5 U/µl of Taq DNA polymerase, 60 ng DNA and 94.5uL of autoclaved ddH<sub>2</sub>O. Primers were tested implementing the following protocol: 96°C for 3 min, 35 cycles of 95°C for 45sec, varying temperature for each locus for 1 min and 72°C for 1min, followed by 72°C for 10min and finally 4°C until use. Heterozygosity was measured using multi-locus heterozygosity (MLH; the total number of loci at which each individual was heterozygous and dividing it by the total number of loci analyzed).

In order to make sure that the alleles were in Hardy-Weinberg equilibrium (HWE), the expected and observed heterozygosity was calculated for all of the loci using GenAlEx (see Table 3.2). In males, Ssa197 and SsaA119 were not in HWE, and in females Ssa202 are not in HWE. As the number of loci not in HWE were minimal, they are not expected to affect the outcome of the results.

### *Statistical analysis*

#### *Females*

All statistics were performed in R version 3.2.1 (R Core Team 2015). A PCA was run on age, mass and length of the individuals as the three factors were highly correlated.

The first two principal components generated out of the PCA referred to from here on in the text as “SizeA” and “SizeB” were used in subsequent analysis.

In order to test for the presence of general effects (i.e. the effect influenced by an individual’s overall heterozygosity as reflected by the microsatellites), the dependent variables (i.e. fecundity, mean egg diameter, mean wet weight, mean dry weight and Fulton’s condition factor) were analyzed individually using a general linear regression (LM) with MLH, SizeA and SizeB as fixed factors in the model. Interaction factors were kept in the model if they provided a better fit to the model using the Akaike Information Criterion (AIC). As both SizeA and SizeB were biologically relevant, they were kept as fixed factors in all final models despite the AIC values. In order to assess whether any observed correlations were due to general or local effects, a univariate linear regression was run (i.e. using MLH) and a multivariate linear regression was run (i.e. using every individual locus as separate binary factors in the model). The results of the two models were then compared using an ANOVA to determine if either model explained significantly more variance.

### *Males*

A PCA was performed on mass and length in order to generate uncorrelated principal components to use in a subsequent analysis. The dependent variables (VAP, VCL, VSL, % motility, linearity (squared, in order to generate a normal distribution), density (log transformed), longevity, percent motility and Fulton’s condition factor) were analyzed using an LM with MLH, age and the first principal component generated from mass and length. Interaction terms were also included in the model if selected by AIC. In

order to assess whether any observed correlations were due to general or local effects, a univariate linear regression was run (i.e. using MLH) and a multivariate linear regression was run (i.e. using every individual locus as separate binary factors in the model). The results of the two models were then compared using an ANOVA to determine if either model explained significantly more variance. If the univariate model explains the most variance, than that is suggestive of general effects. However, if the multivariate method explains significantly more variance, than this is indicative of local effects (see Szulkin et al. 2010 for further information on the univariate vs. multivariate comparison).

### *Results*

#### *Females*

MLH did not correlate with fecundity ( $b=1802.20$ ,  $t(37)=1.24$ ,  $p=0.22$ ), wet egg mass ( $b=5.28$ ,  $t(37)=0.16$ ,  $p=0.87$ ), dry mass ( $b= -8.00$ ,  $t(36)= -0.56$ ,  $p=0.58$ ) or egg diameter ( $b=0.03$ ,  $t(37)= 0.082$ ,  $p=0.94$ ) (see Table 3.5). MLH also did not correlated with Fulton's condition factor ( $b=-0.004$ ,  $t(37)=-1.60$ ,  $p=0.12$ ).

Multivariate models did not significantly predict any more variance in the egg quality traits than the univariate models (wet weight  $F=1.56$ ,  $p=0.19$ ), (dry mass  $F=1.34$ ,  $p=0.28$ ), (diameter  $F=1.21$ ,  $p=0.36$ ) with the exception of fecundity ( $F=2.035$ ,  $p=0.08$ ). The loci SSsp2201, SSsp2215, SsaA124, SsaA119 were significantly different between homozygotes and heterozygotes, with homozygotes at alleles SSsp2201, SsaA124 and SsaA119 having higher fecundity than heterozygotes, while heterozygotes at locus SSsp2215 had higher fecundity than homozygotes. Fulton's condition factor also did not



show any increased correlation under the multivariate analysis ( $F=0.84$ ,  $p=0.65$ ) (see Table 3.6).

### *Males*

Curvilinear velocity at 5 seconds post activation was correlated with MLH explaining 7.9% of the variance ( $b=-169.04$ ,  $t(35)=-2.075$ ,  $p=0.04$ ) and the age of the individual ( $b=32.79$ ,  $t(35)=2.147$ ,  $p=0.04$ ) (see Table 3.7 & Fig. 3.4). VAP at 5 seconds post activation was also correlated with the individual's age ( $b=30.92$ ,  $t(35)=2.121$ ,  $p=0.04$ ) as well as the individual's MLH which explained 7.9% of the variance ( $b=-161.06$ ,  $t(35)=-2.071$ ,  $p=0.05$ ) (see Table 3.5, Fig. 3.5). VSL at 5 seconds post activation was also correlated with MLH explaining 7.4% of the variance ( $b=-124.55$ ,  $t(35)=-1.972$ ,  $p=0.057$ ) (see Fig. 3.6). At 10 seconds post activation, heterozygosity was still significantly correlated with VSL explaining 7.9% of the variance ( $b=-56.037$ ,  $t(34)=-2.053$ ,  $p=0.048$ ) but was no longer significantly correlated with VCL ( $b=-54.434$ ,  $t(34)=-1.674$ ,  $p=0.103$ ) nor VAP ( $b=-60.107$ ,  $t(34)=-1.80$ ,  $p=0.08$ ). However, age did still have a significant effect on both VCL and VAP ( $b=22.566$ ,  $t(34)=3.637$ ,  $p<0.001$ ) and ( $b=20.60$ ,  $t(34)=3.23$ ,  $p=0.003$ ) as well as a correlation with VSL ( $b=10.674$ ,  $t(35)=2.050$ ,  $p=0.048$ ) at 10 seconds post activation. Longevity also correlated with MLH which explained 6.4% of the variance ( $b=-22.37$ ,  $t(33)=-1.97$ ,  $p=0.057$ ) as well as age ( $b=5.486$ ,  $t(33)=2.540$ ,  $p=0.0160$ ) (see Figure 3.7).

Multi-locus heterozygosity did not significantly correlate with linearity at 5 seconds post activation ( $b=-7.079$ ,  $t(35)=-0.003$ ,  $p=0.997$ ) nor at 10 seconds post activation ( $b=-3544.30$ ,  $t(34)=-1.47$ ,  $p=0.149$ ). However age did have a significant

influence on both factors ( $b=-1117.609$ ,  $t(34)=-2159$ ,  $p=0.0378$ ) ( $b=-1028.60$ ,  $t(34)=-2.28$ ,  $p=0.0287$ ). MLH was not correlated with density ( $b=3.63$ ,  $t(35)=1.82$ ,  $p=0.0767$ ), but was correlated with the principal component generated by the length and weight ( $b=0.3323$ ,  $t(35)=2.809$ ,  $p=0.008$ ). Motility at 5 seconds post activation was not significantly correlated with MLH ( $b=-0.329$ ,  $t(34)=-1.01$ ,  $p=0.32$ ).

The condition of individuals was not significantly correlated with MLH ( $b=0.007$ ,  $t(33)=1.57$ ,  $p=0.126$ ). However, the size principal component ( $b=-0.007$ ,  $t(33)=-1.798$ ,  $p=0.08$ ) as well as the size:age interaction ( $b=0.00126$ ,  $t(33)=1.986$ ,  $p=0.055$ ) had influence on the variance in condition between males.

None of the multivariate methods that were examined explained any more variance in any of the fitness related traits that were measured VCL 5s ( $F=0.83$ ,  $p=0.66$ ), VCL 10s ( $F=0.80$ ,  $p=0.68$ ), VAP 5s ( $F=0.92$ ,  $p=0.57$ ), VAP 10s ( $F=0.90$ ,  $p=.59$ ), VSL 5s ( $F=1.27$ ,  $p=0.32$ ), VSL 10s ( $F=1.26$ ,  $p=0.33$ ), squared linearity 5s ( $F=1.00$ ,  $p=0.50$ ), squared linearity 10s ( $F=1.19$ ,  $p=0.37$ ), Motility 5s ( $F=0.80$ ,  $p=0.68$ ), longevity ( $F=1.79$ ,  $p=0.14$ ), density ( $F=0.48$ ,  $p=0.94$ ) and Fulton's condition factor ( $F=1.85$ ,  $p=0.12$ ), suggesting that any observed correlation was the result of general effects and not local effects.

## **Discussion**

In this study, MLH was found to correlate differently with distinct gamete quality traits between the sexes. For males, although negative correlations between MLH and velocity at 5 seconds post activation (VCL, VAP and VSL), 10 seconds post activation for VSL and longevity were found, the variance explained by the MLH ranged between

6.4% and 7.9% at maximum. Therefore, although the results were statistically significant, the correlation might not be highly biologically relevant. Since the multivariate model did not explain any additional variance to the univariate model, it is assumed that these negative correlations, if biologically relevant, were the result of general effects. In females, MLH was not found to correlate with any of the egg quality traits or Fulton's condition factor. Although this study showed no significant difference between the variance explained by the multivariate and univariate models on average, the multivariate model for fecundity did explain a variance of 11.4% compared to 0.39% of the variance in the univariate model, suggesting possible local effects with certain loci. As studies have found varying levels of HFCs with different reproductive traits as well as varying results between the sexes, there are several potential reasons for the variance in observations in the correlation between gamete quality of both sexes and MLH.

A potential reason for the finding of negative HFCs for velocity and longevity of spermatozoa is that they are not measures of fitness in the LaHave broodstock. Although studies have found that velocity (Gage et al. 2004) and longevity (Gage et al. 1995) increase fertilization success, it is possible that the increased fertilization success of these traits does not necessarily correlate with an increase in reproductive success. Although in the wild it is speculated that asynchronous release of gametes and the presence of multiple males will lead to factors such as longevity of sperm having a stronger effect on paternity, in the hatchery setting where the broodstock is reared, the RLC breeding tactic only uses one male's sperm per female. Therefore, sperm competition between males does not exist, resulting in factors such as longevity having a potentially negligible effect on paternity in this type of setting. Although studies have found velocity to be a key

determinant in fertilization success (Gage et al. 2004; Pitcher et al. 2009b), other studies have found little influence of sperm velocity on fertilization success or hatching success. For example, Linhart et al. (2005) found no correlation between the sperm velocity of Common carp (*Cyprinus carpio*) and fertilization success or hatching success. It is therefore possible that velocity does not correlate with reproductive success in this population.

Another potential reason for finding negative HFCs in the velocity and longevity of spermatozoa is outbreeding depression. Outbreeding depression has been found in the offspring of several separated populations (e.g. Gilk et al. 2004; Goldberg et al. 2005), and manifests as the fitness related traits (e.g. reproductive traits) being worse than either parental population (Frankham et al. 2002). Although outbreeding depression is better documented in cases of the crossing of separated populations, it can also manifest within populations (reviewed in Frankham et al. 2010; reviewed in Szulkin and David 2011). Optimum levels of outbreeding within a population do exist just the same as optimum levels of inbreeding, and the two levels are often not very divergent (Willi and Van Buskirk 2005). Under scenarios of outbreeding depression, HFCs are expected to be negative or quadratic (Marshall & Spalton 2000; Neff 2004a). For example, Neff (2004a) found that individual bluegill sunfish with intermediate levels of genetic divergence had better reproductive success than individuals that were highly outbred or highly inbred. As the Atlantic salmon in our study are bred haphazardly with rotational line crosses to insure minimal inbreeding and maximum genetic diversity is maintained, it is possible that with the low variation in MLH levels we are seeing individuals with intermediate heterozygosity (e.g. 70%) with higher fitness than those at the extreme end (e.g. 100%).

Therefore, if outbreeding depression has manifested in the LaHave population of Atlantic salmon, the negative relation between velocity and heterozygosity as well as longevity and heterozygosity would be expected. If our sample had individuals with lower amounts of heterozygosity it is possible that we would have also seen a quadratic relationship rather than a negative one. It is also, perhaps, this lack of variance in the data which lead to no observable correlation between MLH and the measured egg quality traits in the females.

The presence of HFCs in non-inbred populations may also be reflective of heterozygotes at one particular locus displaying higher fitness than homozygotes at that particular locus rather than general effects (Pogson and Zouros 1994; Thelen and Allendorf 2001). For example, Wetzel et al. (2012) found a relation between clutch size and individual MLH in sparrows, however, the relation between egg volume and MLH displayed a more complicated relationship that was indicative that there were multiple loci influencing the outcome variable (Wetzel et al. 2012). Although Heath et al. (2002) found significant general effects with fecundity in Chinook salmon, locus *Omy207* was a significant local influence in all measured female reproductive traits measured in the study (i.e. fecundity, egg size and egg survival). Although for the most part, my study showed no overall significant evidence of local effects, locus *SSsp2201* explained over 8% of the variance in the fecundity model, possibly indicating local effects for this reproductive trait. Therefore, it is possible that local effects are more important for reproductive traits in the LaHave broodstock, which is why no general correlations were found with females.

Life history traits are also known to be plastic (reviewed in Nylin and Gotthard 1998)). Hatchery environments have relaxed selection, which may lead to adaptations that are not suited for natural environments (Lynch and O'Hely 2001; Wedekind 2002; Heath et al. 2003; Thériault et al. 2011; Kekäläinen et al. 2013). Because food sources in hatcheries are not limited once fish reach the exogenous feeding stage, there is no directional selection to favour individuals to expend the energy to produce larger eggs as would be required in the wild (e.g., Srivastava and Brown 1991; Lush et al. 2014). Instead, evolution in hatcheries often will drive egg selection towards smaller yet more numerous eggs instead of fewer and larger eggs (Einum and Fleming 1999; Heath et al. 2003). As the individuals used in this study are the third generation of captively reared individuals, there is no directional selection to lead individuals possessing higher degrees of genetic diversity (i.e. heterozygosity) to have adapted eggs of higher quality (i.e. larger eggs) than individuals that are more homozygous. Certain studies have suggested that heterozygotes have increased homeostasis in varying environments for maintaining a positive phenotype in traits with high plasticity (Marshall and Jain 1968; Gillespie & Turelli 1989), and, although the LaHave broodstock used in this study was the third generation in a line produced from wild gametes in 2007 (personal communication), three generations of hatchery environment rearing may have been enough to influence the selection of traits. Makinen et al. (2014) found that after 5 generations of hatchery environment, convergent evolution between strains of Atlantic salmon took place. If three generations was enough for convergent evolution to start taking place, it would possibly lend explanation to our lack of observing HFCs in female gametes.

Another potential reason for the lack of correlations between heterozygosity and the female gamete quality traits is low additive variance. Traits closely related to fitness (e.g. reproductive traits and survival) have been under strong natural selection for long periods of time and will therefore have seen most of the additive variance fixed as a result (Lynch and Walsh 1998; reviewed in Van Buskirk and Willi 2006). Therefore, variance for these traits will be primarily non-additive variance (Lynch and Walsh 1998; reviewed in Van Buskirk and Willi 2006). Under these conditions, additive genetic variance increases with inbreeding due to the addition of 'low quality' additive variance that is reintroduced in the population once inbreeding occurs (reviewed in Van Buskirk and Willi 2006). In Gage et al. (2006), there was variance in the inbreeding coefficients between the populations of rabbits that were used in the study, whereas in this study the LaHave population has no inbreeding as a result of RLC employed by the hatchery. Therefore, the needed additive genetic variance in order to detect HFCs may not have been present. The fact that the variance was so low due to a lack of inbreeding meant that HFCs (i.e. fecundity, egg diameter, egg mass) could not have been easily detected in many of the gamete quality traits we measured unless the relationship was strong.

There are also several methods for which to measure genetic diversity (e.g. mean  $d^2$  Coulson et al. 1998) Standardized heterozygosity (SH) (Coltman et al. 1999), Interrelatedness (IR)). A meta-analysis, by Chapman et al. (2009), suggested that the results obtained from SH, IR and MLH are all strongly correlated and will lead to similar results in most studies. However, many studies that have observed HFCs in reproductive traits have used mean  $d^2$  (Heath et al. 2002; Neff 2004a; Manias et al. 2014). For example, although Manias et al. (2014) found a positive correlation between clutch size

and heterozygosity as well as egg volume and heterozygosity, the two traits only correlated when heterozygosity was measured as per  $d^2$  and none of the other metrics that were used (i.e. IR, HL, MLH). Although mean  $d^2$  can be a more powerful measure of genetic variation, the measure is more correlated with genetic deviance over long periods of time (e.g. speciation) (Coulson et al. 1998; reviewed in Coltman and Slate 2003; Neff 2004b). As a result, mean  $d^2$  would not have been an appropriate alternative measure of genetic variance for our study.

Finally, as the strength and consistency of HFCs has been questioned in several meta-analysis reviews (e.g. Chapman et al. 2009; Szulkin et al. 2010) it is possible that this study simply did not have the power necessary in order to detect HFCs. For example, in Table 3.1 it is shown that many studies that have found statistically significant HFCs have sample sizes of  $n \geq 100$ . Since our sample size consisted of  $n=42$  females and  $n=38$  males, if HFCs of small or even moderate effect do exist, than we may have needed to double our sample size in order to detect it.

In conclusion, the majority of the correlations were non-significant, with the exception of the negative correlations I found between sperm velocity and longevity with heterozygosity as well as potential local effects associated with fecundity in females. However, it is worth mentioning that these statistically significant correlations explained little variance in the gamete quality traits. Therefore, it would appear that the heterozygosity present in the LaHave broodstock is currently at sufficient levels to not affect the quality of gametes they produce, and therefore should not affect their reproductive success.



## **Acknowledgements**

This study was funded by the NSERC Strategic Project Grant (to TEP and CCW), NSERC Discovery Grant (to TEP), CFI (to TEP), the Ontario Ministry of Economic Development and Innovation (to TEP) as well as the Ontario Ministry of Natural Resources and Forestry (MNRF). I would like to thank Chris Wilson and Anne Kidd from the MNRF and Trent University for providing the microsattellites as well as technical expertise, Brian Rosborough and Chris Weaver from the MNRF for providing the fish as well as logistical support in order to accomplish the study. I would also like to thank Katelynn Johnson, Jade Laycock and Jennifer Smith for providing help with the field work that went into this study. Lastly, I would like to thank my colleagues and lab members Jason Lewis, Adriana Forest and Sarah Lehnert for their assistance and support during this project.

## References

- Chapman JR, Nakagawa S, Coltman DW, Slaters J, Sheldon BC (2009) A quantitative review of heterozygosity-fitness correlations in animal populations. *Mol Ecol* 18: 2746-2765
- Coltman DW, Pilkington JG, Smith JA, Pemberton JM (1999) Parasite-mediated selection against inbred Soay sheep in a free-living island population. *Evolution* 53:1259-1267
- Coltman DW, Slate J (2003) Microsatellite measures of inbreeding: a meta-analysis. *Evolution* 57: 971-983
- Coulson TN, Pemberton JM, Albon SD, Beaumont M, Marshall C, Slate J, Guinness FE, Clutton-Brock TH (1998) Microsatellites reveal heterosis in red deer. *Proc R Soc Lond B* 265: 489-495
- David P (1998) Heterozygosity-fitness correlations: new perspectives on old problems. *Heredity* 80: 531-537
- Einum S, Fleming IA (1999) Maternal effects of egg size in brown trout (*Salmo trutta*): norms of reaction to environmental quality. *Proc R Soc Lond B* 266:2095-2100
- Einum S, Fleming IA (2000) Selection against late emergence and small offspring in Atlantic salmon (*Salmo salar*). *Evolution* 54:628-639
- Fitzpatrick JL, Evans JP (2009) Reduced heterozygosity impairs sperm quality in endangered mammals. *Biol Letters* 5: 320-323

Fitzpatrick JL, Montgomerie R, Desjardins JK, Stiver KA, Kolm N, Balshine S (2009) Female promiscuity promotes the evolution of faster sperm in cichlid fishes. *106*: 1129-1132

Foerster K, Delhey K, Johnsen A, Lifjeld JT, Kempenaers B (2003) Females increase offspring heterozygosity and fitness through extra-pair matings. *Nature* 425, 714-717

Frankham, R., J. D. Ballou, and D. A. Briscoe (2002) *Introduction to Conservation Genetics*. Cambridge University Press, UK.

Frankham R, Ballou JD, Briscoe DA (2010) *Introduction to Conservation Genetics, The Second Edition*. Cambridge, UK (p.618)

Gage MJG, Stockley P, Parker GA (1995) Effects of Alternative Male Mating Strategies on Characteristics of Sperm Production in the Atlantic Salmon (*Salmo salar*): Theoretical and Empirical Investigation. *Philosophical Transactions: Biological Sciences* 350: 391-399

Gage MJG, Stockley P, Parker GA (1998) Sperm morphometry in Atlantic salmon. *J Fish Biol* 53: 835-840

Gage MJG, Macfarlane CP, Yeates S, Ward RG, Searle JB, Parker GA (2004) Spermatozoal Traits and Sperm Competition in Atlantic Salmon: Relative Sperm Velocity Is the Primary Determinant of Fertilization Success. *Curr Biol* 14: 44-47

Gage MJG, SurrIDGE AK, Tomkins JL, Green E, Wiskin L, Bell DJ, Hewitt GM (2006) Reduced Heterozygosity Depresses Sperm Quality in Wild Rabbits, *Oryctolagus cuniculus*. *Curr Biol* 16:612-617

Garcia-Navas V, Ortego J, Sanz JJ (2009) Heterozygosity-Based Assortive Mating in Blue-tits (*Cyanistes caeruleus*): Implications for the Evolution of Mate Choice. PNAS 276:2931-2940

Gilk SE, Wang IA, Hoover CL, Smoker WW, Taylor SG, Gray AK, Gharrett AJ (2004) Outbreeding depression in hybrids between spatially separated pink salmon, *Oncorhynchus gorbuscha*, populations: marine survival, homing ability, and variability in family size. Environ Biol Fishes 69: 287-297

Gillespie JH, Turelli M (1989) Genotype-Environment Interactions and the Maintenance of Polygenic Variation. Genetics 121: 129-138

Goldberg TL, Grant EC, Inendino KR, Kassler TW, Claussen JE, Philipp DP (2005) Increased infectious disease susceptibility resulting from outbreeding depression. Conservation Biology, 19, 455–462

Hansson B, Westerberg L (2002) On the correlation between heterozygosity and fitness in natural populations. Mol Ecol 11:2467-2474

Heath DD, Bryden CA, Shrimpton JM, Iwama GK, Kelly J, Heath JW (2002) Relationships between heterozygosity, allelic distance ( $d^2$ ), and reproductive traits in chinook salmon, *Oncorhynchus tshawytscha*. Can J Fish Aquat Sci 59:77-84

Heath DD, Heath JW, Bryden CA, Johnson RM, Fox CW (2003) Rapid Evolution of Egg Size in Captive Salmon. Science 299:1738-1740

Holm E, Mandrak NE, Burrige, ME (2009) The ROM Field Guide to freshwater fishes of Ontario. Royal Ontario Museum

- Jarne P, Lagoda JL (1996) Microsatellites, from molecules to populations and back. Trends Ecol Evol 8: 285-288
- Johnson K, Butts IAE, Wilson CC, Pitcher TE (2013) Sperm Quality of Hatchery-Reared Lake Trout Throughout the Spawning Season. N AM J Aquacult 75:102-108
- Kekalainen J, Figenschou L, Janhunnen M, Kortet R, Peuhkuri N, Rudolfson G (2013) Hatchery selection may depress the number of motile sperm but intensify selection for their swimming velocity in the Arctic charr. Aquacult Int 21: 405-411
- Keller LF, Waller DM (2002) Inbreeding effects in wild populations. Trends Ecol Evol 17: 230-240
- Kincaid HL (1977) Rotational Line Crossing: An Approach to the Reduction of Inbreeding Accumulation in Trout Broodstocks. Prog Fish Cult 39: 179-181
- King TL, Eackles MS, Letcher BH (2005) Microsatellite DNA markers for the study of Atlantic salmon (*Salmo salar*) kinship, population structure, and mixed-fishery analyses. Mol Ecol Notes 5: 130-132
- Liley NR, Tamkee P, Tsai R, Hoysak DJ (2002) Fertilization dynamics in rainbow trout (*Oncorhynchus mykiss*): effect of male age, social experience, and sperm concentration and motility on in vitro fertilization. Can J Fish Aquat Sci 59: 144-152
- Linhart O, Rodina M, Gela D, Kocour M, Vandeputte M (2005) Spermatozoal competition in common carp (*Cyprinus carpio*): what is the primary determinant of competition success? Reproduction 130: 705-711

Lush L, Burt K, Hamoutene D, Camarillo-Sepulveda N, Perez-Casanova JC, Kenny S, Goulet P, Hinks R, Collier C (2014) Size and ATP Content of Unfertilized Eggs from Farmed and Wild Atlantic Salmon in Newfoundland. *N AM J Aquacult* 76: 138-142

Lynch M (1991) The Genetic Interpretation of Inbreeding Depression and Outbreeding Depression. *Evolution* 45: 622-629

Lynch M, O'Hely M (2001) Captive breeding and the genetic fitness of natural populations. *Conserv Genet* 2: 363-378

Lynch M, Walsh B (1998) *Genetics and Analysis of Quantitative Traits* Sinauer pp.980

Makinen H, Vasemagi A, McGinnity P, Cross T, Primmer C (2014) Population genomic analyses of early-phase Atlantic Salmon (*Salmo salar*) domestication/captive breeding. *Evol Appl* 8: 93-107

Marshall DR, Jain SK (1968) Phenotypic Plasticity of *Avena fatua* and *A. barbata*. *Amer Nat* 102:457-467

Marshall TC, Spalton JA (2000) Simultaneous inbreeding and outbreeding depression in reintroduced Arabian oryx. *Anim Conserv* 3: 241-248

Minias P, Minias A, Dziadek J (2014) Heterozygosity correlates with body size, nest site quality and productivity in a colonial waterbird, the whiskered tern (*Chlidonias hybrid*, Aves: *Sternidae*). *J Zoolog Syst Evol Res.* 53:133-139

Mitton JB (1993) Enzyme heterozygosity, metabolism, and developmental stability. *Genetica* 89: 47-65

- Neff BD (2004a) Stabilizing selection on genomic divergence in a wild fish population. PNAS 101: 2381-2385
- Neff BD (2004b) Mean d2 and Divergence Time: Transformations and Standardizations. Heredity 21:165-171
- Nylin S, Gotthard K (1998) Plasticity in Life-History Traits. Annu Rev Entomol 43: 63-83
- Olafsson K, Hjorleifsdottir S, Pampoulie C, Hreggvidsson GO, Gudjonsson S (2010) Novel set of multiplex assays (SalPrint15) for efficient analysis of 15 microsatellite loci of contemporary samples of the Atlantic salmon (*Salmo salar*) Mol Ecol Resources 10: 533-537
- Olano-Marin J, Mueller JC, Kempnaers B (2011) Heterozygosity and survival in blue tits (*Cyanistes caeruleus*): contrasting effects of presumably functional and neutral loci. Mol Ecol 20: 4028-4041
- Ontario Ministry of Natural Resources (2009) Lake Ontario Fish Communities and Fisheries: 2008 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.
- O'Reilly PT, Hamilton LC, McConnell SK, Wright JM (1996) Rapid analysis of genetic variation in Atlantic salmon (*Salmo salar*) by PCR multiplexing of dinucleotide and tetranucleotide microsatellites. Can J Fish Aquat Sci 53: 2292-2298

- Paterson S, Piertney SB, Knox D, Gilbey J, Verspoor E (2004) Characterization and PCR multiplexing of novel highly variable tetranucleotide Atlantic salmon (*Salmo salar* L.) microsatellites. *Mol Ecol Notes* 4:160-162
- Pickova J, Dutta PC, Larsson PO, Kiessling A (1997) Early embryonic cleavage pattern, hatching success, and egg-lipid fatty acid composition: comparison between two cod (*Gadus morhua*) stocks. *Can J Fish Aquat Sci* 54: 2410-2416
- Pitcher TE, Doucet SM, Beausoleil J, Hanley D (2009a) Secondary sexual characters and sperm traits in Coho Salmon (*Oncorhynchus kisutch*). *J Fish Biol* 74:1450–1461
- Pitcher TE, Beausoleil JJ, Abbott AA, Vandereerden JL (2009b) Sperm design and function in the redbside dace (*Clinostomus elongates*). *J Fish Biol* 75: 924-931
- Pogson GH, Zouros E (1994) Allozyme and RFLP Heterozygosities as Correlates of Growth Rate in the Scallop *Placopecten magellanicus*: A Test of the Associative Overdominance Hypothesis. *Genetics* 157:221-231
- Snook RR (2005) Sperm in competition: not playing by the numbers. *Trends Ecol Evol* 20: 46-53
- Srivastava RK, Brown JA (1991) The biochemical characteristics and hatching performance of cultured and wild Atlantic salmon (*Salmo salar*) eggs. *Can J Zool* 69:2436-2441
- Stanfield L, Jones ML (2003) Factors Influencing Rearing Success of Atlantic Salmon Stocked as Fry and Parr in Lake Ontario Tributaries. *N Am J Fish Sci* 23:1175-1183



Szulkin M, Bierne N, David P (2010) Heterozygosity-fitness correlations: a time for reappraisal. *Evolution* 64:1202-1217

Szulkin M, David P (2011) Negative heterozygosity-fitness correlations observed with microsatellites located in functional areas of the genome. *Mol Ecol* 20:3949-3952

Thelen GC, Allendorf FW (2001) Heterozygosity-fitness correlations in rainbow trout: effects of allozyme loci or associative overdominance? *Evolution* 55: 1180-1187

Thériault V, Moyer GR, Jackson LS, Blouin MS, Banks MA (2011) Reduced reproductive success of hatchery coho salmon in the wild: insights into most likely mechanisms. *Molecular Ecology* 20: 1860-1869

Van Burkirk J, Willi Y (2006) The Change in Quantitative Genetic Variation with Inbreeding. *Evolution* 60: 2428-2434

Wedekind C (2002) Sexual Selection and Life-History Decisions: Implications for Supportive Breeding and the Management of Captive Populations. *Conservation Biology* 16: 1204-1211

Wetzel DP, Stewart IRK, Westneat DF (2012) Heterozygosity predicts clutch and egg size but not plasticity in a house sparrow population with no evidence of inbreeding. *Mol Ecol* 21: 406-420

Table 3.1: Summary of existing literature evaluating the occurrence of HFCs of gamete quality related traits. MLH indicates multilocus heterozygosity, SH indicates standardized heterozygosity, HL indicates homozygosity by weighed locus, mean  $d^2$  is a measure of allelic divergence and IR stands for interrelatedness.

| Taxa  | Trait                          | Sample Size (n) | Number of loci | Measure of heterozygosity | Local or general effects | Direction | Source study             |
|---|--------------------------------|-----------------|----------------|---------------------------|--------------------------|-----------|--------------------------|
| <b>Chinook salmon (<i>O. tshawytscha</i>)</b> | -GSI                           | 100             | 7              | Mean $d^2$                | General                  | (+)       | Heath et al. 2002        |
|   | -fecundity                     | 100             |                | Mean $d^2$                | General & Local          | (+)       |                          |
|   | -egg volume                    |                 |                | Mean $d^2$                | Local                    | (+)       |                          |
|   | -hatching success              |                 |                | Mean $d^2$                | Local                    | (+)       |                          |
| <b>Rabbits</b>                                | -abnormal sperm                | 91              | 29             | MLH                       | General                  | (-)       | Gage et al. 2006         |
| <b>Blue-tits (<i>Cyanistes caeruleus</i>)</b> | -number of eggs sire by a male | 1496            | 79             | MLH                       | General                  | (+)       | Olano-Marin et al. 2011  |
|   | -number of recruits            |                 |                | MLH                       | General                  | (+)       |                          |
|   | -Clutch size                   |                 |                | MLH                       | General                  | (+)       |                          |
| <b>Blue-tits (<i>Cyanistes caeruleus</i>)</b> | -Female hatching success       | 1496            | 79             | SH                        | Local                    | (-)       | Olano-Marin et al. 2011  |
|   | -Female local recruit          |                 |                | SH                        | Local                    | (-)       |                          |
| <b>Blue-tits (<i>Cyanistes caeruleus</i>)</b> | -clutch size                   | 22              | 6              | MLH                       | General                  | (+)       | Foerster et al. 2003     |
|   | -Number of recruits per male   |                 |                | MLH                       | General                  | (+)       |                          |
| <b>Blue-tits (<i>Cyanistes caeruleus</i>)</b> | -clutch size                   | 269             | 14             | MLH                       | General                  | (+)       | Garcia-Navas et al. 2009 |
|   | -Eggshell pattern              |                 |                | MLH                       | General                  | (+)       |                          |

|  |  |      |    |                                       |         |            |                     |
|--|--|------|----|---------------------------------------|---------|------------|---------------------|
| <b>Bluegill sunfish</b><br>( <i>Lepomis macrochirus</i> )    | Reproductive success                           | 142? | 11 | Mean $d^2$                            | General | Quadratic  | Neff 2004a          |
| <b>European shag</b><br>( <i>Phalacrocorax aristotelis</i> ) | Reproductive success for females               | 85   | 7  | MLH<br>HL                             | General | (+)<br>(-) | Velando et al. 2015 |
|  | -Survival                                      |      |    | HL                                    |         | (-)        |                     |
| <b>Whiskered tern</b><br>( <i>Chlidonias hybrid</i> )        | -clutch size<br>-egg size<br>-hatching success | 40   | 8  | -MLH<br>- $d^2$<br>-IR<br>-inverse hl | General | (+)        | Minias et al. 2014  |
| <b>House sparrow</b><br>( <i>Passer domesticus</i> )         | -clutch size                                   | 791  | 21 | MLH                                   | General | (+)        | Wetzel et al. 2012  |
|  | -egg size                                      |      |    | MLH                                   | Local   | (+)        |                     |
|  | -nestling survival                             |      |    | MLH                                   |         | No Cor     |                     |
|  | -hatching success                              |      |    | MLH                                   |         | No Cor     |                     |

\*All results reported in the table are statistically significant. (+) represents a positive linear correlation. (-) represent a negative linear correlation. No Cor stands for no correlation.

Table 3.2: Hardy Weinberg equilibrium of the 19 loci used to calculate heterozygosity. The loci that are out of Hardy-Weinberg equilibrium are in bold. Na is the number of alleles, Ho is the observed heterozygosity, He is the expected heterozygosity.

| Locus           | Size range | Na | Males<br>n= 38 |              | Na | Females<br>n=43 |              | Source Paper         |
|-----------------|------------|----|----------------|--------------|----|-----------------|--------------|----------------------|
|                 |            |    | Ho             | He           |    | Ho              | He           |                      |
| <b>SSsp1605</b> | 227-267    | 8  | 0.833          | 0.808        | 10 | 0.881           | 0.813        | Paterson et al. 2004 |
| <b>SSsp2201</b> | 270-370    | 23 | 0.929          | 0.940        | 24 | 0.929           | 0.934        | Paterson et al. 2004 |
| <b>SSsp2215</b> | 145-193    | 13 | 0.857          | 0.880        | 13 | 0.929           | 0.901        | Paterson et al. 2004 |
| <b>Ssa197</b>   | 163-203    | 9  | <b>0.929</b>   | <b>0.844</b> | 11 | 0.905           | 0.842        | O'Reilly et al. 1996 |
| <b>Ssa202</b>   | 272-312    | 11 | 0.854          | 0.869        | 11 | <b>0.829</b>    | <b>0.867</b> | O'Reilly et al. 1996 |
| <b>SSspG7</b>   | 123-219    | 14 | 0.881          | 0.896        | 15 | 0.786           | 0.869        | Paterson et al. 2004 |
| <b>SSsp2213</b> | 152-204    | 11 | 0.881          | 0.851        | 9  | 0.857           | 0.801        | Paterson et al. 2004 |
| <b>SSsp2216</b> | 195-259    | 14 | 0.929          | 0.908        | 17 | 0.976           | 0.916        | Paterson et al. 2004 |
| <b>SsaA124</b>  | 178-204    | 6  | 0.714          | 0.726        | 6  | 0.800           | 0.734        | King et al. 2005     |
| <b>SsaD190</b>  | 234-378    | 23 | 0.927          | 0.924        | 19 | 0.973           | 0.911        | King et al. 2005     |
| <b>SsaF43</b>   | 106-144    | 7  | 0.881          | 0.791        | 8  | 0.675           | 0.810        | Olafsson et al. 2010 |
| <b>SSsp2210</b> | 110-152    | 3  | 0.167          | 0.154        | 4  | 0.381           | 0.344        | Paterson et al. 2004 |
| <b>SsaA119</b>  | 180-188    | 4  | <b>0.357</b>   | <b>0.514</b> | 3  | 0.488           | 0.488        | King et al. 2005     |
| <b>SsaD157</b>  | 308-452    | 22 | 0.878          | 0.922        | 25 | 0.927           | 0.920        | King et al. 2005     |
| <b>SsaD486</b>  | 174-208    | 8  | 0.548          | 0.607        | 6  | 0.750           | 0.630        | King et al. 2005     |
| <b>SsaD58</b>   | 300-412    | 21 | 0.738          | 0.871        | 22 | 0.780           | 0.911        | King et al. 2005     |
| <b>Ssa171</b>   | 222-276    | 14 | 0.951          | 0.896        | 14 | 0.949           | 0.896        | O'Reilly et al. 1996 |
| <b>SsaA86</b>   | 174-220    | 9  | 0.707          | 0.673        | 10 | 0.825           | 0.744        | King et al. 2005     |
| <b>SsaD144</b>  | 172-284    | 20 | 0.976          | 0.933        | 20 | 0.974           | 0.920        | King et al. 2005     |

Table 3.3: Summary of the heterozygosity and gamete quality metrics for the hatchery-reared female LaHave Atlantic salmon (*Salmo salar*) (n=42) from the broodstock at the Harwood Fish Culture Station. Mean, maximum value, minimum value and standard deviation as well as standard error of the population mean.

| <b>Trait</b>  | <b>Mean</b> | <b>Maximum</b> | <b>Minimum</b> | <b>S.E.</b> |
|---|-------------|----------------|----------------|-------------|
| <b>Heterozygosity index (%)</b>                     | 0.821       | 1.00           | 0.5625         | 0.0175      |
| <b>Fecundity (# of eggs)</b>                        | 5679.039    | 8654           | 1653.3         | 284.47      |
| <b>Mean egg diameter (mm)</b>                       | 6.27        | 7.025          | 5.52           | 0.0482      |
| <b>Mean wet weight (mg)</b>                         | 167.85      | 220.82         | 111.32         | 3.674       |
| <b>Mean dry weight (mg)*</b>                        | 59.70       | 76.87          | 38.56          | 1.565       |
| <b>Fulton's condition factor (g/cm<sup>3</sup>)</b> | 0.01632     | 0.02022        | 0.00944        | 0.0003      |
| <b>Mass (g)</b>                                     | 4568.3      | 7927           | 1335           | 224.89      |
| <b>Fork length (inches)</b>                         | 25.49       | 31             | 18             | 0.435       |

\*N=41 for the dry egg weight

Table 3.4: Summary of the heterozygosity and gamete quality metrics for the hatchery-reared male LaHave Atlantic salmon (*Salmo salar*) (n=38) from the broodstock at the Harwood Fish Culture Station. Mean, maximum value, minimum value and standard deviation as well as standard error of the population mean. VCL stands for curvilinear velocity, VAP stands for average path velocity, VSL stands for progressive velocity.

| <b>Trait</b>   | <b>Mean</b> | <b>Maximum</b> | <b>Minimum</b> | <b>S.E.</b> |
|--|-------------|----------------|----------------|-------------|
| <b>Heterozygosity (%)</b>  | 0.787       | 0.895          | 0.632          | 0.0131      |
| <b>VCL (<math>\mu\text{m/s}</math>) at 5sec post activation</b>      | 114.00      | 190.7          | 44.5           | 7.00        |
| <b>VCL (<math>\mu\text{m/s}</math>) at 10 sec post activation</b>    | 69.79       | 110.05         | 35.5           | 3.00        |
| <b>VAP (<math>\mu\text{m/s}</math>) at 5 sec</b>                     | 105.03      | 181.35         | 40.5           | 6.67        |
| <b>VAP (<math>\mu\text{m/s}</math>) at 10 sec</b>                    | 63.82       | 104.85         | 30.1           | 3.02        |
| <b>VSL (<math>\mu\text{m/s}</math>) at 5 sec</b>                     | 81.38       | 160.4          | 37.1           | 5.21        |
| <b>VSL (<math>\mu\text{m/s}</math>) at 10 sec</b>                    | 52.73       | 97.35          | 27.7           | 2.34        |
| <b>Linearity (degree/<math>\mu\text{m}</math>) at 5 sec</b>          | 72.84       | 87             | 45             | 1.66        |
| <b>Linearity (degree/<math>\mu\text{m}</math>) at 10 sec</b>         | 76.99       | 91.5           | 50.5           | 1.41        |
| <b>Motility (%)</b>  | 60.67       | 91.02          | 32.59          | 2.61        |
| <b>Density (cells/ml)</b>  | 2961349     | 16250000       | 187500         | 628164.1    |
| <b>Longevity (s)</b>   | 26.47       | 40.75          | 14.25          | 1.02        |
| <b>Mass (g)</b>  | 4087.3      | 10409          | 1310           | 368.49      |
| <b>Fork Length (inches)</b>  | 25.87       | 48             | 18             | 0.999       |
| <b>Fulton's condition factor (<math>\text{m}/\text{cm}^3</math>)</b> | 0.0138      | 0.018029       | 0.00274        | 0.00044     |

Table 3.5: General linear regression between egg quality traits and multilocus heterozygosity for females. All values represent those in the full model (df=37) while controlling for size. Unstandardized b value, t value and p value for the multilocus heterozygosity final model as well as the difference in the adjusted  $r^2$  of the final model and the final model without heterozygosity.

| <b>Traits</b>                    | <b>Unstandardized b values</b> | <b>t values</b> | <b>P values</b> | <b>Difference in Adjusted R<sup>2</sup></b> |
|----------------------------------|--------------------------------|-----------------|-----------------|---|
| <b>Fecundity</b>                 | 1802.20                        | 1.235           | 0.22            | 0.0039                                      |
| <b>Wet Egg Mass</b>              | 5.28                           | 0.160           | 0.87            | -0.0225                                     |
| <b>Dry Egg Mass</b>              | -8.00                          | 0.560           | 0.58            | -0.0169                                     |
| <b>Egg Diameter</b>              | 0.03                           | 0.082           | 0.94            | -0.0184                                     |
| <b>Fulton's condition factor</b> | -0.005                         | 1.602           | 0.12            | 0.0320                                      |

Table 3.6: Summary of the multivariate analysis of weight of each locus on fecundity. The adjusted R<sup>2</sup> value given to each locus was calculated by subtracting the adjusted R<sup>2</sup> value of the model without the locus in question from that of the full multivariate model.

| <b>Locus</b>    | <b>Adjusted R<sup>2</sup></b> |
|-----------------|-------------------------------|
| <b>SSsp1605</b> | 0.017                         |
| <b>SSsp2201</b> | 0.080                         |
| <b>SSsp2215</b> | 0.024                         |
| <b>Ssa197</b>   | -0.011                        |
| <b>Ssa202</b>   | -0.012                        |
| <b>SSspG7</b>   | -0.0086                       |
| <b>SSsp2213</b> | -0.0002                       |
| <b>SSsp2216</b> | -0.011                        |
| <b>SsaA124</b>  | 0.006                         |
| <b>SsaD190</b>  | 0.030                         |
| <b>SsaF43</b>   | -0.011                        |
| <b>SSsp2210</b> | 0.0008                        |
| <b>SsaA119</b>  | 0.027                         |
| <b>SsaD157</b>  | 0.00                          |
| <b>SsaD486</b>  | 0.005                         |
| <b>SsaD58</b>   | 0.005                         |
| <b>Ssa171</b>   | 0.011                         |
| <b>SsaA86</b>   | 0.022                         |
| <b>SsaD144</b>  | -0.0029                       |



Table 3.7: General linear regression between sperm quality traits and heterogosity for males (n = 38). Unstandardized b value, t value and p value for the multilocus heterozygosity final model selected through AIC as well as the difference in the adjusted R<sup>2</sup> of the final model and the final model without heterozygosity. Age was a covariate in all models with the exception of density. The size principal component was a covariate in all three velocity measures at 10 sec post activation, longevity and condition. Finally, age:size interaction was a covariate in the models for condition and longevity.

| <b>Traits</b>           | <b>Unstandardized b values</b> | <b>T values</b> | <b>P values</b> | <b>Difference in Adjusted R<sup>2</sup></b> |
|-------------------------|--------------------------------|-----------------|-----------------|---|
| <b>VCL 5s</b>           | -169.04                        | -2.075          | 0.04            | 0.079                                       |
| <b>VCL 10s</b>          | -54.434                        | -1.674          | 0.103           | 0.038                                       |
| <b>VAP 5s</b>           | -161.06                        | -2.071          | 0.046           | 0.078                                       |
| <b>VAP 10s</b>          | -60.107                        | -1.796          | 0.081           | 0.049                                       |
| <b>VSL 5s</b>           | -124.55                        | -1.972          | 0.057           | 0.074                                       |
| <b>VSL 10s</b>          | -56.037                        | -2.053          | 0.048           | 0.079                                       |
| <b>Linearity 5s</b>     | -7.079                         | -0.003          | 0.99            | -0.026                                      |
| <b>Linearity 10s</b>    | -3544.3                        | -1.474          | 0.15            | 0.028                                       |
| <b>Motility 5s</b>      | -0.3294                        | -1.014          | 0.32            | 0.0008                                      |
| <b>Longevity</b>        | -22.366                        | -1.974          | 0.057           | 0.064                                       |
| <b>Density</b>          | 3.631                          | 1.824           | 0.077           | 0.052                                       |
| <b>Fulton's</b>         | 0.007                          | 1.566           | 0.13            | 0.034                                       |
| <b>Condition factor</b> |                                |                 |                 |   |

## Figure Captions

Figure 3.1 Rotational Line Crossing depiction of the parental generation and the following crosses. P represents the parental generation, F<sub>1</sub> represents the first generation of crosses and F<sub>2</sub> the second generation that are offspring of crosses. The solid lines represent the females used to produce the line. The dashed lines represent the males used to produce the line. (adapted from Kincaid 1977)

Figure 3.2 Distribution of multilocus heterozygosity (MLH) for the male Atlantic salmon in this study (n=38). MLH was measured as the number of heterozygous loci divided by the total number of loci.


Figure 3.3 Distribution of multilocus heterozygosity (MLH) for the female Atlantic salmon in this study (n=42). MLH was measured as the number of heterozygous loci divided by the total number of loci.

Figure 3.4 Negative relationship between curvilinear velocity (VCL) at 5 seconds post activation and multilocus heterozygosity ( $b=-169.04$ ,  $t(35)=-2.075$ ,  $p=0.04$ , dashed line) and at 10 seconds post activation ( $b=-54.434$ ,  $t(34)=-1.674$ ,  $p=0.103$ , solid line).


Figure 3.5 Negative relationship between average path velocity (VAP) at 5 seconds post activation and multilocus heterozygosity ( $b=-161.06$ ,  $t(35)=-2.071$ ,  $p=0.05$ , dashed line) and at 10 seconds post activation ( $b=-60.107$ ,  $t(34)=-1.80$ ,  $p=0.08$ , solid line).

Figure 3.6 Negative relationship between progressive velocity (VSL) at 5 seconds post activation and multilocus heterozygosity ( $b=-124.55$ ,  $t(35)=-1.972$ ,  $p=0.057$ , dashed line) and at 10 seconds post activation ( $b=-56.037$ ,  $t(34)=-2.053$ ,  $p=0.048$ , solid line).


**Figure 3.1**




**Figure 3.2**




**Figure 3.3**




**Figure 3.4**



**Figure 3.5**




**Figure 3.6**





**Figure 3.7**



## CHAPTER 4: GENERAL DISCUSSION

### Summary

In this thesis, I examined the relationship between genetic quality and phenotypic expression of fitness related traits (e.g. survival, morphology and gamete quality) in Atlantic salmon and how it may impact the reintroduction efforts to Lake Ontario. The purpose of this chapter is to summarize my findings of both Chapter 2 and Chapter 3, address the caveats that have come up in previous chapters and make suggestions regarding the future direction studies should take in order to further the research that was done in this thesis.

### Chapter 2

The manifestation of outbreeding depression becomes increasingly likely when crossing populations that have been separated for >500 years or have lived in different environments for 20 generations or more (Frankham et al. 2011). As the reintroduction effort to Lake Ontario is simultaneously stocking populations that have been segregated since the last glacial period (Ward 1932; MNRF 2009) outbreeding depression is a potential outcome. This chapter examined the likelihood of outbreeding depression manifesting in the juvenile life phase of F<sub>1</sub> hybrids between the Sebago and LaHave strains of Atlantic salmon by using a 2 x 2 full factorial design in order to compare fitness related traits between the pure strains and their reciprocal hybrids while controlling for maternal and paternal effects. The results of this study showed no significant influence of hybridization on survival or any of the fitness related traits measured in juvenile Atlantic salmon.

### *Changes to Environmental Settings*

As my study only tested for the occurrence of outbreeding depression under intrinsic factors (Tymchuk et al. 2007), it is possible that the environmental setting of the experiment made for a lack of outbreeding depression. As phenotype is the result of an interaction between genotype and the environment, a change in extrinsic factors such as environment can alter the individual fitness of *S. salar* as much as intrinsic factors depending on whether or not they are suited for their surroundings (Garcia de Leaniz et al. 2007). Isolated populations of species that display specific local adaptations for small geographic areas, such as salmonids, are severely affected by the loss of these local adaptations (Templeton 1986; Tymchuk et al. 2007; Vandersteen et al. 2012). In fact, it is believed that the loss of local adaptations has a stronger influence over outbreeding depression than do intrinsic factors such as the disruption of co-adapted gene complexes for these local populations (Allendorf et al. 2001; Edmands and Timmerman 2003; Tymchuk et al. 2007). Therefore, if outbreeding depression were to result from the loss of local adaptations, it may be masked by a relaxed environmental settings such as the one used in this study (Tymchuk et al. 2007; Crespel et al. 2013). A genotype that performs perfectly well in a hatchery setting may perform differently in the parental environment (Tymchuk et al. 2007; Crespel et al. 2013). Several examples of this are documented in salmonids. In Tymchuk et al. (2007), F<sub>3</sub> hybrids *O. mykiss* displayed signs of outbreeding depression in relation to growth and survival only under certain environmental conditions, suggesting that the outbreeding depression was the result of extrinsic factors simply expedited by intrinsic factors. The strength of environmental influences may also vary depending on genotype. Different genotypes of *O. mykiss* had varying survival depending

on geographic area as well as seasonality (Vandersteen et al. 2012). Crespel et al. (2013) found a similar result with Brook trout (*Salvelinus fontinalis*) in which environmental interactions affected certain hybrid crosses but not others. Therefore, in the case of this study, it is possible that if the juveniles had lost a local adaptation of importance, any resulting outbreeding depression might not have been possible to detect in a hatchery setting. Therefore, I recommend that future studies examining the effect of outbreeding, on  $F_1$  generations specifically, should conduct the study in both parental environments if possible in order to detect outbreeding depression caused by extrinsic factors. In the event where the individuals are being transplanted to a novel environment which is not similar to either parental environments, it would perhaps be of equal value to also test the individuals in multiple settings of the new environment during various life stages in order to assess if they would suffer from signs of outbreeding depression in the intended environment. In the case of future studies examining Atlantic salmon in Lake Ontario, testing the individuals in multiple tributaries around the lake to insure that they are not going to suffer outbreeding depression in relation to the environment would be recommended.

#### *F<sub>2</sub>, F<sub>3</sub> and backcrosses*

As outbreeding depression as a result of intrinsic factors is most likely to manifest in the  $F_2$  generation or later when the original parental genomes on the same chromosome are subject to recombination (Dobzhansky 1948; Lynch 1991; Edmands 1999 McClelland et al. 2005; Tymchuk et al. 2007), it is possible that outbreeding depression could still manifest in the  $F_2$  generation of the Sebago and LaHave crosses of Atlantic salmon. If the  $F_1$  generation displays no evidence of outbreeding depression, they will be equally as

likely to reproduce as their pure strain counterparts resulting in F<sub>2</sub> generation hybrids and backcrosses. Outbreeding depression resulting from the breakup of co-adapted gene complexes has been suggested in studies on intraspecific salmonid hybrids. For example, McGinnity et al. (2003) found that F<sub>2</sub> hybrid *S. salar* had higher egg mortality than backcrosses as well as crosses by the same sire, suggesting that the outbreeding depression was the result of intrinsic incompatibilities. Another study, Edmands (1999), demonstrated that outbreeding depression can last until the F<sub>3</sub> generation, future studies looking at outbreeding depression as a result of hybridization should conduct the study to the F<sub>3</sub> generation (with the inclusion of backcrosses) in order to determine that there is no breakdown of co-adapted gene complexes that may lead to outbreeding depression.

#### *Life History Traits and fitness beyond the juvenile life stage*

Certain traits (e.g. life history traits) are reported to be more closely linked to inbreeding depression than morphological traits (DeRose and Roff 1999; originally noted by Falconer 1989) they may be more correlated with outbreeding depression than morphological traits as well. For example the loss of local adaptations has been shown to influence behaviour and homing ability (Gilk et al. 2004). For example, *O. mykiss* x *O. clarki* hybrids migrated earlier than non-hybridized *O. clarki* populations due to altering reactions to photoperiod (Corsi et al. 2012). As migration affects spawning time and success, life history traits may be a better indicator of outbreeding depression over an individual's lifespan rather than early growth traits (McClelland et al. 2005). Therefore, any future studies should evaluate the lifelong fitness of individuals if possible rather than just the juvenile stage.

## Chapter 3

Heterozygosity fitness correlations (HFC) are correlations between fitness related traits and a measure of heterozygosity that can occur in life history traits as well as morphological traits (reviewed in Chapman et al. 2009; reviewed in Szulkin et al. 2010) and they can manifest through general or local effects (David 1998, Lynch and Walsh 1998). As heterozygosity is thought to be correlated with genetic diversity and a lack of inbreeding, HFCs are generally expected to be positive in nature and can be indicative of inbreeding depression. It is important to know the correlation of heterozygosity and gamete quality in populations. The LaHave population is the most common for stocking efforts in Lake Ontario, Chapter 3 examined the correlation between multilocus heterozygosity and sperm and egg quality within the broodstock which was accomplished using 19 microsatellite markers. As heterozygosity is indicative of genetic diversity, it was predicted that heterozygosity and the gamete quality traits would be positively correlated. However, I found a negative correlation between sperm velocity as well as sperm longevity and multilocus heterozygosity.

The results found in the study are possibly indicative of outbreeding depression in the LaHave strain of Atlantic salmon. As all of the individuals in this study were > 50% heterozygous, it is possible that the individuals closer to the central position of heterozygosity (i.e. 50%) have the highest degree of fitness while those at either extreme (i.e. 0% or 100%) have the lowest degree of fitness. For example, Neff (2004) found a quadratic relationship between reproductive performance and heterozygosity with intermediate levels displaying the highest level of reproductive success while individuals at either extreme (i.e. inbred and outbred individuals) displayed the lowest. As, our

population had little variance in the inbreeding coefficient, it is possible that if there was more variability in the multilocus heterozygosity in our population that the relationship found between velocity/longevity and multilocus heterozygosity would have been a quadratic one (Neff 2004). Therefore, for future studies researching HFCs in hatchery populations, the experimental design should include individuals inbred and outbred in order to obtain a full spectrum of inbreeding coefficients in order to better assess the direction of any observed correlation.

Although no general effects were found between multilocus heterozygosity and the egg quality traits, local effects may be present between locus SSsp2201 and fecundity. To further the studies around the local effects, I recommend using a linkage map (i.e. a chromosomal map displaying the relative positions of loci) in order to analyze any selected loci in the chromosomal vicinity using existing genome maps out of Europe. Although questions surrounding the usage of European Atlantic salmon maps for North-American Atlantic salmon have arisen (Lubieniecki et al. 2010), a recent study has found that although chromosomal changes do exist, as a whole on a finer scale, the genome remains largely conservative and can therefore be used in reference to North-American populations of Atlantic salmon (Brenna-Hansen et al. 2012).

## **Conclusion**

Conservation management rarely takes into account outbreeding depression when selecting candidate populations for reintroduction or when breeding individuals in the hatchery. This is partially the result of most effort going into avoiding inbreeding and the consequences associated with it. This thesis examined outbreeding depression and how it

may play a role in the reintroduction of Atlantic salmon on two fronts; first through looking at hybridization and second through looking at the gamete quality of the hatchery broodstock. This study found no evidence to support the occurrence of outbreeding depression as a result of hybridization, however negative correlations between sperm quality and heterozygosity was observed in the LaHave broodstock. It is imperative that conservation management responsible for the broodstock of fish that are going to be stocked consider outbreeding depression as just as severe a risk as inbreeding depression and construct methods to avoid its consequences.



## References

Allendorf FW, Thorgaard GH (1984) Tetraploidy and the evolution of salmonid fishes. In Turner B (ed) Evolutionary Genetics of Fishes. Plenum Press, New York. pp.1-53

Brenna-Hansen S, Li J, Kent M, Boulding E, Dominik S, Davidson W, Lien S (2012) Chromosomal Differences between European and North American Atlantic salmon Discovered by Linkage Mapping and Supported by Fluorescence in situ Hybridization Analysis. BMC Genomics 13:432

Chapman JR, Nakagawa S, Coltman DW, Slaters J, Sheldon BC (2009) A quantitative review of heterozygosity-fitness correlations in animal populations. Mol Ecol 18: 2746-2765

Crespel A, Audet C, Bernatchez L, Garant D (2013) Effects of rearing environmental and strain combination on hybrid vigour in brook trout. N Am J Aquacult 74: 188-198

Corsi MP, Eby LA, Barfoot CA (2012) Hybridization with rainbow trout alters life history traits of native western cutthroat trout. Can J Fish Aquat Sci 70: 895-904

David P (1998) Heterozygosity-fitness correlations: new perspectives on old problems. Heredity 80: 531–537

DeRose MA, Roff DA (1999) A comparison of inbreeding depression in life-history and morphological traits in animals. Evolution, 53: 1288–1292

Dobzhansky T (1948) Genetic of natural populations. XVIII. Experiments on chromosomes of *Drosophila pseudoobscura* from different geographical regions. Genetics 33:588-602

- Edmands S (1999) Hybrid vigour and outbreeding depression in interpopulation crosses spanning a wide range of divergence. *Evolution* 53: 1757-1768
- Edmands S, Timmerman CC (2003) Modeling factors affecting the severity of outbreeding depression. *Conserv Biol* 17: 883-892
- Falconer DS (1989) An introduction to quantitative genetics 3<sup>rd</sup> ed. Wiley, London
- Frankham R, Ballou JD, Eldridge MDB, Lacy RC, Ralls K, Dudash MR and Fenster CB (2011) Predicting the Probability of Outbreeding Depression. *Conserv Biol* 25: 465-475
- Garcia de Leaniz C, Fleming IA, Einum S, Verspoor E, Jordan WC, Consuegra S, Aubin-Horth N, Lajus D, Letcher BH, Youngson AF, Webb JH, Vøllestad LA, Villanueva B, Ferguson A, Quinn TP (2007) A critical review of adaptive genetic variation in Atlantic salmon: implications for Conservation. *Biol Rev* 82: 173-211
- Gilk SE, Wang IA, Hoover CL, Smoker WW, Taylor SG, Gray AK, Gharrett AJ (2004) Outbreeding depression in hybrids between spatially separated pink salmon, *Oncorhynchus gorbuscha*, populations: marine survival, homing ability, and variability in family size. *Environ Biol Fishes* 69: 287-297
- Lubieniecki KP, Jones SL, Davidson EA, Park J, Koop BF, Walker S, Davidson WS (2010) comparative Genomic Analysis of Atlantic Salmon, *Salmo Salar*, from Europe and North America. *GMC Genetics* 11:105
- Lynch M (1991) The Genetic Interpretation of Inbreeding Depression and Outbreeding Depression. *Evolution* 45: 622-629

Lynch M, Walsh B (1998) Genetics and Analysis of Quantitative Traits. Sinauer & Associates Inc, Sunderland, Massachusetts. pp.980

McClelland EK, Myers JM, Hard JJ, Park LK, Naish KA (2005) Two generations of outbreeding in coho salmon (*Oncorhynchus kisutch*): effects on size and growth. Can J Fish Aquat Sci 62: 2538-2547

McGinnity P, Prodöhl P, Ferguson A, Hynes R, Maoléidigh NO, Baker N, Cotter D, O'Hea B, Cooke D, Rogan G, Taggart J, Cross T (2003) Fitness Reduction and Potential Extinction of Wild Populations of Atlantic Salmon, *Salmo salar*, as a Result of Interactions with Escaped Farm Salmon. Proceedings: Biological Sciences 270: 2443-2450

Neff BD (2004) Stabilizing selection on genomic divergence in a wild fish population. PNAS 101: 2381-2385

Ontario Ministry of Natural Resources (2009) Lake Ontario Fish Communities and Fisheries: 2008 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada

Szulkin M, Bierne N, David P (2010) Heterozygosity-fitness correlations: a time for reappraisal. Evolution 64:1202-1217

Templeton AR (1986) Coadaptation and outbreeding depression. In Soulé M (ed) Conservation biology: the science of scarcity and diversity. Sinauer, Sunderland, MA. pp. 105-116

Tymchuk WE, Sundstrom LF, Devlin RH (2007) Growth and Survival Trade-offs and outbreeding depression in rainbow trout (*Oncorhynchus mykiss*). *Evolution* 61:1225-1237

Vandersteen W, Biro P, Harris L, Devlin R (2012) Introgression of domesticated alleles into a wild trout genotype and the impact on seasonal survival in natural lakes. *Evol Appl* 5: 76-88

Ward HB (1932) The Origin of the Landlocked Habit in Salmon. *PNAS* 18: 569-580

APPENDICES

Appendix 1: Mean length, mass and Fulton's condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in April 2013 with one unit of standard deviation

| <b>Family Block</b> | <b>Dam</b> | <b>Sire</b> | <b>Hybrid Type</b> | <b>Mean Length (mm)</b> | <b>Mean Mass (g)</b> | <b>Mean Condition (g/mm<sup>3</sup>*10000)</b> |
|---------------------|------------|-------------|--------------------|-------------------------|----------------------|--|
| 1                   | L6         | L10         | LL                 | 33.07±3.30              | 0.37±0.11            | 0.1010±0.0171                                  |
| 1                   | S15        | S27         | SS                 | 35.08±2.39              | 0.43±0.10            | 0.0966±0.0074                                  |
| 1                   | L6         | S27         | LS                 | 34.09±2.37              | 0.37±0.09            | 0.0911±0.0048                                  |
| 1                   | S15        | L10         | SL                 | 35.56±2.69              | 0.45±0.12            | 0.0982±0.0061                                  |
| 2                   | L4         | L14         | LL                 | 35.97±2.15              | 0.47±0.10            | 0.0988±0.0077                                  |
| 2                   | S13        | S35         | SS                 | 33.30±1.82              | 0.36±0.06            | 0.0966±0.0046                                  |
| 2                   | L4         | S35         | LS                 | 33.09±3.36              | 0.36±0.12            | 0.0944±0.0094                                  |
| 2                   | S13        | L14         | SL                 | 34.05±2.65              | 0.40±0.10            | 0.0976±0.0060                                  |
| 4                   | L1         | L19         | LL                 | 34.27±2.94              | 0.41±0.12            | 0.0988±0.0051                                  |
| 4                   | S10        | S36         | SS                 | 35.78±3.14              | 0.48±0.12            | 0.1017±0.0067                                  |
| 4                   | L1         | S36         | LS                 | 34.57±2.83              | 0.41±0.12            | 0.0960±0.0072                                  |
| 4                   | S10        | L19         | SL                 | 35.12±3.03              | 0.44±0.14            | 0.0977±0.0101                                  |
| 7                   | L8         | L16         | LL                 | 32.61±2.99              | 0.34±0.10            | 0.0932±0.0077                                  |
| 7                   | S11        | S29         | SS                 | 33.66±2.38              | 0.39±0.09            | 0.0993±0.0073                                  |
| 7                   | L8         | S29         | LS                 | 32.62±2.93              | 0.33±0.09            | 0.0929±0.0056                                  |
| 7                   | S11        | L16         | SL                 | 33.90±2.93              | 0.39±0.12            | 0.0969±0.0104                                  |
| 8                   | L7         | L20         | LL                 | 34.67±3.56              | 0.39±0.12            | 0.0889±0.0086                                  |
| 8                   | S5         | S33         | SS                 | 33.79±2.06              | 0.39±0.08            | 0.0985±0.0985                                  |
| 8                   | L7         | S33         | LS                 | 33.94±2.75              | 0.37±0.10            | 0.0928±0.0928                                  |
| 8                   | S5         | L20         | SL                 | 35.84±1.79              | 0.46±0.08            | 0.0986±0.0986                                  |
| 9                   | L118       | L100        | LL                 | 38.05±4.53              | 0.52±0.21            | 0.0905±0.0905                                  |
| 9                   | S116       | S100        | SS                 | 36.90±3.64              | 0.52±0.16            | 0.0995±0.0995                                  |
| 9                   | L118       | S100        | LS                 | 36.33±3.69              | 0.52±0.17            | 0.1044±0.0117                                  |
| 9                   | S116       | L100        | SL                 | 36.53±3.88              | 0.47±0.15            | 0.0931±0.0078                                  |
| 10                  | L119       | L101        | LL                 | 35.43±3.48              | 0.50±0.17            | 0.1074±0.0063                                  |
| 10                  | S117       | S101        | SS                 | 37.47±4.94              | 0.54±0.21            | 0.0971±0.0052                                  |
| 10                  | L119       | S101        | LS                 | 36.64±3.13              | 0.58±0.16            | 0.1143±0.0079                                  |
| 10                  | S117       | L101        | SL                 | 35.49±3.20              | 0.49±0.13            | 0.1078±0.0081                                  |
| 11                  | L120       | L102        | LL                 | 40.03±1.25              | 0.73±0.10            | 0.1132±0.0096                                  |
| 11                  | S118       | S102        | SS                 | 35.73±2.66              | 0.51±0.13            | 0.1105±0.0056                                  |
| 11                  | L120       | S102        | LS                 | 38.78±3.73              | 0.56±0.19            | 0.0919±0.0079                                  |
| 11                  | S118       | L102        | SL                 | 38.37±3.28              | 0.58±0.15            | 0.1018±0.0152                                  |
| 12                  | L121       | L103        | LL                 | 36.01±3.87              | 0.52±0.17            | 0.1072±0.0078                                  |

|           |      |      |    |            |           |               |
|-----------|------|------|----|------------|-----------|---------------|
| <b>12</b> | S119 | S103 | SS | 36.71±3.43 | 0.48±0.13 | 0.0950±0.0068 |
| <b>12</b> | L121 | S103 | LS | 37.40±2.98 | 0.58±0.14 | 0.1085±0.0060 |
| <b>12</b> | S119 | L103 | SL | 36.12±3.84 | 0.45±0.15 | 0.0915±0.0055 |
| <b>13</b> | L122 | L104 | LL | 34.82±4.10 | 0.45±0.17 | 0.1010±0.0093 |
| <b>13</b> | S120 | S104 | SS | 35.08±3.54 | 0.43±0.14 | 0.0950±0.0072 |
| <b>13</b> | L122 | S104 | LS | 37.09±3.63 | 0.57±0.18 | 0.1080±0.0078 |
| <b>13</b> | S120 | L104 | SL | 32.92±3.67 | 0.38±0.16 | 0.0997±0.0100 |
| <b>14</b> | L123 | L106 | LL | 34.81±2.98 | 0.43±0.11 | 0.0991±0.0081 |
| <b>14</b> | S121 | S105 | SS | 36.17±3.64 | 0.51±0.15 | 0.1057±0.0076 |
| <b>14</b> | L123 | S105 | LS | 31.98±3.01 | 0.36±0.12 | 0.1043±0.0109 |
| <b>14</b> | S121 | L106 | SL | 36.84±2.67 | 0.54±0.12 | 0.1051±0.0083 |
| <b>16</b> | L125 | L108 | LL | 33.82±3.19 | 0.43±0.13 | 0.1065±0.0083 |
| <b>16</b> | S124 | S107 | SS | 34.66±3.41 | 0.42±0.14 | 0.0963±0.0066 |
| <b>16</b> | L125 | S107 | LS | 35.82±3.01 | 0.48±0.13 | 0.1009±0.0059 |
| <b>16</b> | S124 | L108 | SL | 33.92±2.94 | 0.40±0.10 | 0.0997±0.0148 |
| <b>17</b> | L126 | L109 | LL | 37.54±3.49 | 0.51±0.14 | 0.0946±0.0053 |
| <b>17</b> | S125 | S108 | SS | 39.50±3.28 | 0.68±0.18 | 0.1069±0.0072 |
| <b>17</b> | L126 | S108 | LS | 37.50±2.38 | 0.57±0.12 | 0.1062±0.0059 |
| <b>17</b> | S125 | L109 | SL | 39.56±2.87 | 0.71±0.17 | 0.1115±0.0073 |
| <b>20</b> | L131 | L113 | LL | 38.63±1.76 | 0.58±0.07 | 0.0999±0.0052 |
| <b>20</b> | S128 | S113 | SS | 38.61±2.88 | 0.66±0.02 | 0.1121±0.0074 |
| <b>20</b> | L131 | S113 | LS | 41.39±2.33 | 0.72±0.12 | 0.1007±0.0055 |
| <b>20</b> | S128 | L113 | SL | 36.67±4.44 | 0.56±0.22 | 0.1061±0.0094 |

Appendix 2: Mean length, mass and Fulton's condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in August 2013 with one unit of standard deviation

| <b>Family Block</b> | <b>Dam</b> | <b>Sire</b> | <b>Hybrid Type</b> | <b>Mean Length (mm)</b> | <b>Mean Mass (g)</b> | <b>Mean Condition (g/mm<sup>3</sup>*10000)</b> |
|---------------------|------------|-------------|--------------------|-------------------------|----------------------|--|
| 1                   | L6         | L10         | LL                 | 60.96±5.93              | 3.05±0.74            | 0.1356±0.0226                                  |
| 1                   | S15        | S27         | SS                 | 58.43±6.18              | 2.77±0.82            | 0.1355±0.0186                                  |
| 1                   | L6         | S27         | LS                 | 60.22±7.13              | 2.50±0.73            | 0.1123±0.0089                                  |
| 1                   | S15        | L10         | SL                 | 56.71±4.93              | 2.49±0.81            | 0.1338±0.0246                                  |
| 2                   | L4         | L14         | LL                 | 65.12±4.90              | 3.27±0.73            | 0.1178±0.0155                                  |
| 2                   | S13        | S35         | SS                 | 61.44±4.39              | 2.91±0.74            | 0.1231±0.0119                                  |
| 2                   | L4         | S35         | LS                 | 56.96±6.92              | 2.58±0.85            | 0.1362±0.0167                                  |
| 2                   | S13        | L14         | SL                 | 57.28±5.58              | 2.57±0.89            | 0.1324±0.0173                                  |
| 4                   | L1         | L19         | LL                 | 59.11±6.97              | 2.62±0.94            | 0.1227±0.0148                                  |
| 4                   | S10        | S36         | SS                 | 58.90±6.24              | 2.76±0.90            | 0.1315±0.0144                                  |
| 4                   | L1         | S36         | LS                 | 55.17±5.92              | 2.28±0.65            | 0.1349±0.0237                                  |
| 4                   | S10        | L19         | SL                 | 61.93±5.44              | 3.18±0.80            | 0.1321±0.0141                                  |
| 7                   | L8         | L16         | LL                 | 54.53±7.17              | 2.19±0.78            | 0.1336±0.0223                                  |
| 7                   | S11        | S29         | SS                 | 58.96±5.02              | 2.54±0.65            | 0.1221±0.0100                                  |
| 7                   | L8         | S29         | LS                 | 54.15±3.96              | 1.98±0.43            | 0.1231±0.0104                                  |
| 7                   | S11        | L16         | SL                 | 53.28±6.85              | 2.08±0.87            | 0.1323±0.0135                                  |
| 8                   | L7         | L20         | LL                 | 56.70±6.04              | 2.21±0.56            | 0.1204±0.0170                                  |
| 8                   | S5         | S33         | SS                 | 59.03±3.87              | 2.53±0.60            | 0.1209±0.0074                                  |
| 8                   | L7         | S33         | LS                 | 53.87±5.74              | 2.00±0.64            | 0.1250±0.0152                                  |
| 8                   | S5         | L20         | SL                 | 57.57±4.85              | 2.46±0.66            | 0.1272±0.0153                                  |
| 9                   | L118       | L100        | LL                 | 67.28±6.70              | 3.89±1.52            | 0.1223±0.0105                                  |
| 9                   | S116       | S100        | SS                 | 66.48±7.45              | 3.25±1.09            | 0.1069±0.0116                                  |
| 9                   | L118       | S100        | LS                 | 62.86±4.81              | 3.00±0.67            | 0.1191±0.0084                                  |
| 9                   | S116       | L100        | SL                 | 63.94±6.00              | 2.85±0.83            | 0.1065±0.0073                                  |
| 10                  | L119       | L101        | LL                 | 63.08±4.61              | 2.89±0.74            | 0.1128±0.0070                                  |
| 10                  | S117       | S101        | SS                 | 67.37±5.28              | 3.54±0.82            | 0.1141±0.0111                                  |
| 10                  | L119       | S101        | LS                 | 63.14±5.43              | 2.90±0.77            | 0.1127±0.0053                                  |
| 10                  | S117       | L101        | SL                 | 62.76±5.17              | 2.93±0.82            | 0.1158±0.0082                                  |
| 11                  | L120       | L102        | LL                 | 71.85±10.45             | 3.91±1.15            | 0.1049±0.0141                                  |
| 11                  | S118       | S102        | SS                 | 62.86±4.61              | 2.81±0.65            | 0.1112±0.0053                                  |
| 11                  | L120       | S102        | LS                 | 63.77±5.22              | 2.78±0.68            | 0.1060±0.0118                                  |
| 11                  | S118       | L102        | SL                 | 65.26±3.20              | 2.71±0.53            | 0.0964±0.0075                                  |
| 12                  | L121       | L103        | LL                 | 65.12±5.78              | 3.11±0.79            | 0.1100±0.0060                                  |
| 12                  | S119       | S103        | SS                 | 62.68±6.03              | 3.00±0.91            | 0.1187±0.0061                                  |
| 12                  | L121       | S103        | LS                 | 63.71±4.48              | 2.77±0.58            | 0.1058±0.0085                                  |

|           |      |      |    |            |           |               |
|-----------|------|------|----|------------|-----------|---------------|
| <b>12</b> | S119 | L103 | SL | 59.53±5.52 | 2.53±0.80 | 0.1165±0.0071 |
| <b>13</b> | L122 | L104 | LL | 60.07±9.93 | 2.47±1.38 | 0.1063±0.0075 |
| <b>13</b> | S120 | S104 | SS | 62.00±7.15 | 3.01±1.20 | 0.1208±0.0096 |
| <b>13</b> | L122 | S104 | LS | 61.18±5.75 | 2.57±0.76 | 0.1100±0.0175 |
| <b>13</b> | S120 | L104 | SL | 54.00±5.30 | 2.07±0.55 | 0.1303±0.0194 |
| <b>14</b> | L123 | L106 | LL | 61.03±5.39 | 2.71±0.69 | 0.1168±0.0082 |
| <b>14</b> | S121 | S105 | SS | 64.78±6.02 | 3.03±0.95 | 0.1085±0.0080 |
| <b>14</b> | L123 | S105 | LS | 57.59±5.88 | 2.29±0.70 | 0.1172±0.0110 |
| <b>14</b> | S121 | L106 | SL | 64.22±5.39 | 2.92±0.71 | 0.1087±0.0082 |
| <b>16</b> | L125 | L108 | LL | 52.20±7.18 | 2.02±0.86 | 0.1382±0.0187 |
| <b>16</b> | S124 | S107 | SS | 62.59±7.01 | 2.85±0.97 | 0.1132±0.0079 |
| <b>16</b> | L125 | S107 | LS | 62.44±5.63 | 2.57±0.78 | 0.1034±0.0091 |
| <b>16</b> | S124 | L108 | SL | 57.26±6.77 | 2.15±0.80 | 0.1113±0.0116 |
| <b>17</b> | L126 | L109 | LL | 61.28±8.01 | 2.40±0.89 | 0.1011±0.0102 |
| <b>17</b> | S125 | S108 | SS | 65.87±5.52 | 3.42±0.73 | 0.1185±0.0065 |
| <b>17</b> | L126 | S108 | LS | 59.52±5.93 | 2.50±0.67 | 0.1163±0.0100 |
| <b>17</b> | S125 | L109 | SL | 64.66±6.22 | 2.63±0.69 | 0.0995±0.0322 |
| <b>20</b> | L131 | L113 | LL | 66.21±4.33 | 3.15±0.62 | 0.1072±0.0038 |
| <b>20</b> | S128 | S113 | SS | 68.87±4.25 | 3.42±0.68 | 0.1035±0.0060 |
| <b>20</b> | L131 | S113 | LS | 69.28±5.82 | 3.77±1.04 | 0.1115±0.0019 |
| <b>20</b> | S128 | L113 | SL | 62.06±5.50 | 2.79±0.81 | 0.1139±0.0091 |



Appendix 3: Mean length, mass and Fulton's condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in October 2013 with one unit of standard deviation

| <b>Family Block</b> | <b>Dam</b> | <b>Sire</b> | <b>Hybrid Type</b> | <b>Mean Length (mm)</b> | <b>Mean Mass (g)</b> | <b>Mean Condition (g/mm<sup>3</sup>*10000)</b> |
|---------------------|------------|-------------|--------------------|-------------------------|----------------------|--|
| 1                   | L6         | L10         | LL                 | 90.98±11.64             | 7.83±3.46            | 0.1005±0.0167                                  |
| 1                   | S15        | S27         | SS                 | 81.75±11.11             | 6.38±2.41            | 0.1113±0.0045                                  |
| 1                   | L6         | S27         | LS                 | 84.31±9.41              | 5.79±1.90            | 0.0934±0.0023                                  |
| 1                   | S15        | L10         | SL                 | 79.22±8.69              | 5.71±1.84            | 0.1107±0.0072                                  |
| 2                   | L4         | L14         | LL                 | 83.40±8.51              | 6.36±1.96            | 0.1070±0.0035                                  |
| 2                   | S13        | S35         | SS                 | 96.89±11.64             | 8.87±2.88            | 0.0951±0.0036                                  |
| 2                   | L4         | S35         | LS                 | 74.06±8.01              | 4.50±1.42            | 0.1073±0.0027                                  |
| 2                   | S13        | L14         | SL                 | 80.77±7.59              | 5.27±1.60            | 0.0970±0.0045                                  |
| 4                   | L1         | L19         | LL                 | 83.99±12.94             | 6.38±2.94            | 0.1009±0.0044                                  |
| 4                   | S10        | S36         | SS                 | 97.39±14.99             | 10.80±4.81           | 0.1104±0.0055                                  |
| 4                   | L1         | S36         | LS                 | 88.81±14.35             | 7.57±3.63            | 0.1012±0.0035                                  |
| 4                   | S10        | L19         | SL                 | 90.04±11.72             | 7.38±2.99            | 0.0955±0.0050                                  |
| 7                   | L8         | L16         | LL                 | 72.35±7.94              | 4.42±1.59            | 0.1120±0.0064                                  |
| 7                   | S11        | S29         | SS                 | 83.65±14.42             | 7.13±3.94            | 0.1123±0.0042                                  |
| 7                   | L8         | S29         | LS                 | 74.57±15.86             | 4.79±3.81            | 0.1002±0.0047                                  |
| 7                   | S11        | L16         | SL                 | 80.08±13.81             | 5.48±3.15            | 0.0977±0.0049                                  |
| 8                   | L7         | L20         | LL                 | 79.42±12.53             | 5.00±2.76            | 0.0933±0.0089                                  |
| 8                   | S5         | S33         | SS                 | 87.94±12.53             | 6.83±2.97            | 0.0946±0.0035                                  |
| 8                   | L7         | S33         | LS                 | 86.277±10.01            | 6.37±2.51            | 0.0953±0.0030                                  |
| 8                   | S5         | L20         | SL                 | 83.34±6.60              | 5.08±1.43            | 0.0857±0.0055                                  |
| 9                   | L118       | L100        | LL                 | 91.29±13.10             | 7.88±4.23            | 0.0967±0.0065                                  |
| 9                   | S116       | S100        | SS                 | 75.01±9.87              | 4.95±2.49            | 0.1101±0.0068                                  |
| 9                   | L118       | S100        | LS                 | 77.61±7.13              | 4.31±0.99            | 0.0909±0.0054                                  |
| 9                   | S116       | L100        | SL                 | 79.01±11.57             | 5.71±2.56            | 0.1092±0.0032                                  |
| 10                  | L119       | L101        | LL                 | 85.03±8.04              | 5.83±1.88            | 0.0918±0.0046                                  |
| 10                  | S117       | S101        | SS                 | 87.39±7.77              | 7.53±1.95            | 0.1110±0.0058                                  |
| 10                  | L119       | S101        | LS                 | 79.77±10.18             | 5.40±2.29            | 0.1001±0.0064                                  |
| 10                  | S117       | L101        | SL                 | 97.48±16.79             | 11.03±4.78           | 0.1150±0.0078                                  |
| 11                  | L120       | L102        | LL                 | 91.72±16.57             | 7.47±4.60            | 0.0896±0.0102                                  |
| 11                  | S118       | S102        | SS                 | 79.12±8.11              | 5.83±1.76            | 0.1146±0.0050                                  |
| 11                  | L120       | S102        | LS                 | 79.21±5.20              | 4.71±1.11            | 0.0930±0.0063                                  |
| 11                  | S118       | L102        | SL                 | 72.12±6.90              | 4.28±1.21            | 0.1117±0.0058                                  |
| 12                  | L121       | L103        | LL                 | 83.84±9.62              | 6.95±2.48            | 0.1135±0.0071                                  |
| 12                  | S119       | S103        | SS                 | 87.46±16.50             | 6.90±4.51            | 0.0915±0.0073                                  |
| 12                  | L121       | S103        | LS                 | 82.66±10.45             | 5.73±2.08            | 0.0985±0.0161                                  |

---

|           |      |      |    |              |            |               |
|-----------|------|------|----|--------------|------------|---------------|
| <b>12</b> | S119 | L103 | SL | 78.31±12.16  | 5.94±2.84  | 0.1153±0.0029 |
| <b>13</b> | L122 | L104 | LL | 77.23±16.00  | 5.92±4.61  | 0.1130±0.0061 |
| <b>13</b> | S120 | S104 | SS | 84.18±11.46  | 6.18±2.82  | 0.0976±0.0043 |
| <b>13</b> | L122 | S104 | LS | 75.57±13.31  | 5.32±3.30  | 0.1114±0.0054 |
| <b>13</b> | S120 | L104 | SL | 79.80±5.45   | 4.96±1.05  | 0.0963±0.0034 |
| <b>14</b> | L123 | L106 | LL | 82.67±11.29  | 5.65±2.90  | 0.0940±0.0051 |
| <b>14</b> | S121 | S105 | SS | 78.88±10.82  | 5.58±2.62  | 0.1074±0.0038 |
| <b>14</b> | L123 | S105 | LS | 76.93±11.93  | 5.35±2.60  | 0.1100±0.0048 |
| <b>14</b> | S121 | L106 | SL | 76.99±5.39   | 5.09±1.03  | 0.1102±0.0031 |
| <b>16</b> | L125 | L108 | LL | 83.80±19.15  | 6.27±4.97  | 0.0914±0.0056 |
| <b>16</b> | S124 | S107 | SS | 101.11±26.26 | 10.86±7.76 | 0.0955±0.0002 |
| <b>16</b> | L125 | S107 | LS | 80.54±11.57  | 5.28±2.63  | 0.0953±0.0063 |
| <b>16</b> | S124 | L108 | SL | 75.89±13.79  | 4.76±3.15  | 0.0982±0.0042 |
| <b>17</b> | L126 | L109 | LL | 82.99±9.93   | 5.36±2.29  | 0.0890±0.0057 |
| <b>17</b> | S125 | S108 | SS | 87.12±10.82  | 6.20±2.54  | 0.0895±0.0026 |
| <b>17</b> | L126 | S108 | LS | 79.14±8.65   | 5.70±1.96  | 0.1105±0.0054 |
| <b>17</b> | S125 | L109 | SL | 79.13±12.07  | 5.16±2.51  | 0.0978±0.0089 |
| <b>20</b> | L131 | L113 | LL | 79.76±8.86   | 4.60±1.54  | 0.0873±0.0053 |
| <b>20</b> | S128 | S113 | SS | 80.76±7.74   | 5.78±1.66  | 0.1075±0.0054 |
| <b>20</b> | L131 | S113 | LS | 92.93±22.71  | 9.88±7.42  | 0.1096±0.0009 |
| <b>20</b> | S128 | L113 | SL | 74.75±8.92   | 4.57±1.72  | 0.1057±0.0049 |

---

Appendix 4: Mean length, mass and Fulton's condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in December 2013 with one unit of standard deviation

| <b>Family Block</b> | <b>Dam</b> | <b>Sire</b> | <b>Hybrid Type</b> | <b>Mean Length (mm)</b> | <b>Mean Mass (g)</b> | <b>Mean Condition (g/mm<sup>3</sup>*10000)</b> |
|---------------------|------------|-------------|--------------------|-------------------------|----------------------|--|
| 1                   | L6         | L10         | LL                 | 96.91±12.22             | 11.63±5.46           | 0.1219±0.0170                                  |
| 1                   | S15        | S27         | SS                 | 89.53±14.77             | 8.77±3.93            | 0.1147±0.0074                                  |
| 1                   | L6         | S27         | LS                 | 88.06±11.82             | 8.12±3.49            | 0.1135±0.0043                                  |
| 1                   | S15        | L10         | SL                 | 89.94±7.48              | 8.45±2.29            | 0.1138±0.0043                                  |
| 2                   | L4         | L14         | LL                 | 87.46±10.78             | 7.85±2.83            | 0.1142±0.0059                                  |
| 2                   | S13        | S35         | SS                 | 108.92±7.89             | 14.97±3.52           | 0.1143±0.0036                                  |
| 2                   | L4         | S35         | LS                 | 86.24±14.27             | 7.78±3.90            | 0.1137±0.0039                                  |
| 2                   | S13        | L14         | SL                 | 88.40±8.70              | 7.89±2.21            | 0.1118±0.0053                                  |
| 4                   | L1         | L19         | LL                 | 92.68±15.78             | 9.49±4.84            | 0.1104±0.0051                                  |
| 4                   | S10        | S36         | SS                 | 110.87±18.79            | 16.42±7.84           | 0.1127±0.0054                                  |
| 4                   | L1         | S36         | LS                 | 97.52±16.38             | 11.26±5.65           | 0.1139±0.0049                                  |
| 4                   | S10        | L19         | SL                 | 96.35±15.40             | 10.60±5.32           | 0.1099±0.0052                                  |
| 7                   | L8         | L16         | LL                 | 85.58±18.75             | 8.03±6.30            | 0.1120±0.0057                                  |
| 7                   | S11        | S29         | SS                 | 94.06±15.03             | 10.65±5.17           | 0.1198±0.0037                                  |
| 7                   | L8         | S29         | LS                 | 82.49±19.36             | 7.60±6.05            | 0.1181±0.0054                                  |
| 7                   | S11        | L16         | SL                 | 88.25±16.18             | 8.60±5.31            | 0.1141±0.0032                                  |
| 8                   | L7         | L20         | LL                 | 85.82±15.71             | 8.02±4.83            | 0.1157±0.0063                                  |
| 8                   | S5         | S33         | SS                 | 90.13±16.34             | 9.36±4.85            | 0.1177±0.0051                                  |
| 8                   | L7         | S33         | LS                 | 93.80±11.53             | 9.43±3.49            | 0.1101±0.0058                                  |
| 8                   | S5         | L20         | SL                 | 86.75±8.21              | 7.28±2.10            | 0.1090±0.0053                                  |
| 9                   | L118       | L100        | LL                 | 96.12±15.77             | 11.01±6.43           | 0.1143±0.0053                                  |
| 9                   | S116       | S100        | SS                 | 87.740±17.39            | 8.47±5.54            | 0.1123±0.0044                                  |
| 9                   | L118       | S100        | LS                 | 81.220±9.18             | 6.58±2.04            | 0.1188±0.0052                                  |
| 9                   | S116       | L100        | SL                 | 89.49±15.04             | 8.51±4.10            | 0.1118±0.0049                                  |
| 10                  | L119       | L101        | LL                 | 92.82±10.68             | 9.76±3.63            | 0.1172±0.0042                                  |
| 10                  | S117       | S101        | SS                 | 97.99±16.40             | 10.41±5.34           | 0.1044±0.0023                                  |
| 10                  | L119       | S101        | LS                 | 86.71±14.69             | 8.01±4.39            | 0.1126±0.0034                                  |
| 10                  | S117       | L101        | SL                 | 111.84±23.39            | 18.01±10.98          | 0.1163±0.0058                                  |
| 11                  | L120       | L102        | LL                 | 100.50±23.19            | 12.89±9.00           | 0.1151±0.0076                                  |
| 11                  | S118       | S102        | SS                 | 92.06±10.57             | 9.52±3.33            | 0.1177±0.0038                                  |
| 11                  | L120       | S102        | LS                 | 89.66±14.30             | 8.97±4.22            | 0.1176±0.0067                                  |
| 11                  | S118       | L102        | SL                 | 84.20±7.54              | 6.92±1.98            | 0.1129±0.0052                                  |
| 12                  | L121       | L103        | LL                 | 93.95±13.91             | 9.70±4.54            | 0.1100±0.0049                                  |
| 12                  | S119       | S103        | SS                 | 93.89±24.64             | 11.31±9.39           | 0.1138±0.0077                                  |
| 12                  | L121       | S103        | LS                 | 91.24±11.54             | 8.78±3.39            | 0.1112±0.0053                                  |
| 12                  | S119       | L103        | SL                 | 94.96±17.21             | 10.41±5.44           | 0.1122±0.0051                                  |

---

|           |      |      |    |              |             |               |
|-----------|------|------|----|--------------|-------------|---------------|
| <b>13</b> | L122 | L104 | LL | 85.84±21.77  | 8.57±8.13   | 0.1125±0.0050 |
| <b>13</b> | S120 | S104 | SS | 91.08±15.56  | 9.14±5.13   | 0.1113±0.0031 |
| <b>13</b> | L122 | S104 | LS | 84.56±13.45  | 7.11±3.52   | 0.1094±0.0034 |
| <b>13</b> | S120 | L104 | SL | 89.13±7.78   | 8.12±2.06   | 0.1124±0.0040 |
| <b>14</b> | L123 | L106 | LL | 88.63±14.41  | 8.50±5.23   | 0.1124±0.0033 |
| <b>14</b> | S121 | S105 | SS | 90.16±17.12  | 8.84±5.00   | 0.1107±0.0054 |
| <b>14</b> | L123 | S105 | LS | 86.59±17.42  | 7.96±5.03   | 0.1110±0.0046 |
| <b>14</b> | S121 | L106 | SL | 88.63±10.61  | 8.00±3.21   | 0.1107±0.0043 |
| <b>16</b> | L125 | L108 | LL | 92.96±21.78  | 10.12±8.11  | 0.1090±0.0045 |
| <b>16</b> | S124 | S107 | SS | 110.57±31.83 | 16.89±13.27 | 0.1107±0.0011 |
| <b>16</b> | L125 | S107 | LS | 86.85±15.13  | 7.87±4.86   | 0.1110±0.0051 |
| <b>16</b> | S124 | L108 | SL | 76.21±8.41   | 5.13±1.77   | 0.1118±0.0028 |
| <b>17</b> | L126 | L109 | LL | 91.24±12.83  | 8.66±3.91   | 0.1075±0.0059 |
| <b>17</b> | S125 | S108 | SS | 91.93±12.65  | 8.78±3.60   | 0.1075±0.0045 |
| <b>17</b> | L126 | S108 | LS | 91.50±12.03  | 8.73±3.28   | 0.1090±0.0045 |
| <b>17</b> | S125 | L109 | SL | 82.11±11.29  | 6.15±2.43   | 0.1103±0.0164 |
| <b>20</b> | L131 | L113 | LL | 89.31±9.75   | 7.91±2.65   | 0.1072±0.0048 |
| <b>20</b> | S128 | S113 | SS | 90.52±9.62   | 8.38±2.64   | 0.1100±0.0042 |
| <b>20</b> | L131 | S113 | LS | 103.70±30.01 | 14.65±12.92 | 0.1112±0.0022 |
| <b>20</b> | S128 | L113 | SL | 86.52±11.40  | 7.16±2.75   | 0.1059±0.0037 |

---

Appendix 5: Mean length, mass and Fulton's condition factor of the cross types (pure LaHave LL, pure Sebago SS, dam Sebago and LaHave sire SL and LaHave dam and Sebago sire LS) of Atlantic salmon within each family block as measured in February 2014 with one unit of standard deviation

| <b>Family Block</b> | <b>Dam</b> | <b>Sire</b> | <b>Hybrid Type</b> | <b>Mean Length (mm)</b> | <b>Mean Mass (g)</b> | <b>Mean Condition (g/mm<sup>3</sup>*10000)</b> |
|---------------------|------------|-------------|--------------------|-------------------------|----------------------|--|
| 1                   | L6         | L10         | LL                 | 111.15±2.91             | 15.82±2.90           | 0.1153±0.0215                                  |
| 1                   | S15        | S27         | SS                 | 103.44±13.77            | 12.75±4.50           | 0.1108±0.0062                                  |
| 1                   | L6         | S27         | LS                 | 107.48±16.27            | 13.84±6.55           | 0.1057±0.0041                                  |
| 1                   | S15        | L10         | SL                 | 100.97±12.24            | 11.36±4.18           | 0.1062±0.0043                                  |
| 2                   | L4         | L14         | LL                 | 110.48±14.24            | 15.08±5.25           | 0.1088±0.0063                                  |
| 2                   | S13        | S35         | SS                 | 125.35±7.92             | 20.83±3.78           | 0.1050±0.0026                                  |
| 2                   | L4         | S35         | LS                 | 102.16±16.64            | 11.99±5.88           | 0.1057±0.0081                                  |
| 2                   | S13        | L14         | SL                 | 103.98±8.85             | 11.79±2.24           | 0.1043±0.0078                                  |
| 4                   | L1         | L19         | LL                 | 110.17±15.74            | 14.83±5.83           | 0.1065±0.0046                                  |
| 4                   | S10        | S36         | SS                 | 128.44±19.03            | 25.12±10.48          | 0.1132±0.0031                                  |
| 4                   | L1         | S36         | LS                 | 113.74±17.16            | 17.23±7.50           | 0.1119±0.0059                                  |
| 4                   | S10        | L19         | SL                 | 110.56±16.04            | 15.17±6.81           | 0.1059±0.0053                                  |
| 7                   | L8         | L16         | LL                 | 97.00±14.65             | 11.02±4.50           | 0.1140±0.0075                                  |
| 7                   | S11        | S29         | SS                 | 114.26±11.49            | 17.52±5.28           | 0.1148±0.0094                                  |
| 7                   | L8         | S29         | LS                 | 96.84±13.76             | 11.63±4.73           | 0.1216±0.0076                                  |
| 7                   | S11        | L16         | SL                 | 103.97±19.35            | 13.69±7.35           | 0.1135±0.0066                                  |
| 8                   | L7         | L20         | LL                 | 104.79±17.58            | 14.01±7.07           | 0.1141±0.0069                                  |
| 8                   | S5         | S33         | SS                 | 103.81±18.73            | 13.90±6.55           | 0.1163±0.0068                                  |
| 8                   | L7         | S33         | LS                 | 111.38±11.89            | 15.13±4.62           | 0.1068±0.0052                                  |
| 8                   | S5         | L20         | SL                 | 98.51±4.50              | 10.36±1.29           | 0.1081±0.0045                                  |
| 9                   | L118       | L100        | LL                 | 105.74±14.28            | 14.15±5.85           | 0.1147±0.0062                                  |
| 9                   | S116       | S100        | SS                 | 100.51±21.46            | 12.84±7.95           | 0.1155±0.0096                                  |
| 9                   | L118       | S100        | LS                 | 97.70±9.725             | 10.77±2.65           | 0.1144±0.0071                                  |
| 9                   | S116       | L100        | SL                 | 101.89±16.08            | 12.07±5.33           | 0.1089±0.0075                                  |
| 10                  | L119       | L101        | LL                 | 107.45±12.60            | 14.63±5.15           | 0.1143±0.0042                                  |
| 10                  | S117       | S101        | SS                 | 120.16±19.90            | 18.96±9.14           | 0.1040±0.0015                                  |
| 10                  | L119       | S101        | LS                 | 104.01±16.00            | 13.12±5.62           | 0.1102±0.0043                                  |
| 10                  | S117       | L101        | SL                 | 125.60±19.33            | 23.86±10.72          | 0.1162±0.0002                                  |
| 11                  | L120       | L102        | LL                 | 119.18±35.79            | 22.58±18.13          | 0.1177±0.0004                                  |
| 11                  | S118       | S102        | SS                 | 107.43±10.69            | 14.60±4.51           | 0.1144±0.0039                                  |
| 11                  | L120       | S102        | LS                 | 105.65±13.36            | 13.73±4.68           | 0.1130±0.0072                                  |
| 11                  | S118       | L102        | SL                 | 96.62±12.65             | 10.02±3.58           | 0.1069±0.0040                                  |
| 12                  | L121       | L103        | LL                 | 106.62±16.39            | 13.68±6.72           | 0.1057±0.0047                                  |
| 12                  | S119       | S103        | SS                 | 104.48±24.98            | 14.73±11.68          | 0.1145±0.0067                                  |
| 12                  | L121       | S103        | LS                 | 105.99±11.82            | 13.00±4.43           | 0.1061±0.0089                                  |
| 12                  | S119       | L103        | SL                 | 111.78±19.06            | 16.32±7.49           | 0.1103±0.0065                                  |

|           |      |      |    |              |             |               |
|-----------|------|------|----|--------------|-------------|---------------|
| <b>13</b> | L122 | L104 | LL | 95.51±15.65  | 10.70±5.65  | 0.1136±0.0058 |
| <b>13</b> | S120 | S104 | SS | 105.37±17.86 | 13.64±7.23  | 0.1079±0.0043 |
| <b>13</b> | L122 | S104 | LS | 104.40±21.12 | 13.38±8.05  | 0.1059±0.0026 |
| <b>13</b> | S120 | L104 | SL | 105.35±13.76 | 13.10±4.70  | 0.1084±0.0047 |
| <b>14</b> | L123 | L106 | LL | 104.02±17.41 | 13.09±7.73  | 0.1078±0.0031 |
| <b>14</b> | S121 | S105 | SS | 104.26±18.26 | 12.82±6.85  | 0.1045±0.0039 |
| <b>14</b> | L123 | S105 | LS | 99.96±19.47  | 11.75±6.97  | 0.1078±0.0059 |
| <b>14</b> | S121 | L106 | SL | 104.62±10.26 | 12.48±4.20  | 0.1062±0.0043 |
| <b>16</b> | L125 | L108 | LL | 106.02±22.51 | 14.50±10.31 | 0.1084±0.0049 |
| <b>16</b> | S124 | S107 | SS | 128.39±33.13 | 26.27±18.64 | 0.1130±0.0003 |
| <b>16</b> | L125 | S107 | LS | 102.38±17.79 | 12.31±6.63  | 0.1069±0.0070 |
| <b>16</b> | S124 | L108 | SL | 96.79±20.99  | 11.36±8.43  | 0.1111±0.0049 |
| <b>17</b> | L126 | L109 | LL | 103.50±14.14 | 12.69±5.61  | 0.1086±0.0069 |
| <b>17</b> | S125 | S108 | SS | 107.99±9.61  | 13.63±3.10  | 0.1067±0.0041 |
| <b>17</b> | L126 | S108 | LS | 104.55±14.24 | 13.41±5.61  | 0.1117±0.0067 |
| <b>17</b> | S125 | L109 | SL | 101.78±15.86 | 11.65±5.72  | 0.1044±0.0047 |
| <b>20</b> | L131 | L113 | LL | 101.10±12.18 | 11.32±4.04  | 0.1056±0.0048 |
| <b>20</b> | S128 | S113 | SS | 101.68±10.39 | 11.43±3.46  | 0.1059±0.0034 |
| <b>20</b> | L131 | S113 | LS | 119.55±3.18  | 18.34±0.65  | 0.1077±0.012  |
| <b>20</b> | S128 | L113 | SL | 95.35±11.32  | 9.13±3.34   | 0.1021±0.004  |

## APPENDIX 6: MORPHOMETRY AND SWIMMING ABILITY OF INTRASPECIFIC HYBRID ATLANTIC SALMON (SALMON SALAR) IN THE JUVENILE LIFE STAGE

### **Introduction**

Hybridization is the crossing of individuals from segregated populations and can lead to a phenomena called outbreeding depression (Frankham et al. 2002). When outbreeding depression occurs in the first generation, it is usually the result of an intermediate phenotype, where the hybrid offspring has lost local adaptations of importance and is no longer suitable for either parental environment (Lynch 1991; Edmands 2007). Salmonids are a highly philopatric species that often have local adaptations. Local adaptations between allopatric populations of salmonids often result in distinct morphological differences (Taylor 1991). These morphological differences can be explained by abiotic factors in the environment such as depth, temperature, DO, etc. (Turan 2000), and can influence swimming ability. Atlantic salmon have previously been shown to exhibit environmentally induced morphology with more fusiform shape in environments with faster flow (Taylor 1986; Drinan et al. 2012). Faster flowing water usually results in more streamline bodies in order to decrease swimming costs (Boily and Magnan 2002). As the reintroduction effort of Atlantic salmon to Lake Ontario is introducing multiple allopatric populations simultaneously (i.e. LaHave and Sebago) (Dimond and Smitka 2005, MNRF 2009), it is possible that morphological differences between the populations will result in an intermediate phenotype in juveniles and therefore affect their swimming ability. The objective of this appendix is to summarize and compare the swimming performance as well as the morphometric characteristics of the reciprocal LaHave x Sebago hybrids compared to the pure strains in order to determine if outbreeding depression resulting from the loss of local adaptations is a potential occurrence in the juvenile life phase.

## Methods

### *Swimming ability & Morphometrics*

Swim performance was measured between December 18, 2013 and January 20 2014. Five fish per cross type in each family block (20 fish per family block) had their swimming ability tested using a swim flume of 32 L (Swim-30, Loligo Systems, Tjele, Denmark). The individual was acclimated at 0.28 m/s current for 3 minutes, and, once the acclimation period was over, the swim flume was increased to 0.53 m/s and the trial began. Every 2 minutes the current was increased by 0.31 m/s until the fish was no longer able to swim or the maximum current of 1.63 m/s was reached. These recordings will be used to calculate the critical swimming speed ( $U_{crit}$ ); an important measurement that is used to represent a fish's maximum swimming ability in short durations (Fisher et al. 2005). Critical swimming speed ( $U_{crit}$ ) will be calculated as:  $U_{crit} = U_f + U_s \cdot (T_f / T_s)$ , where  $U_f$  is the speed of the last completed interval,  $U_s$  is the interval speed increment,  $T_f$  is the time spent during the last interval and  $T_s$  is the length of the interval (Fisher et al. 2005).

An area with lower current was detected in the swim flume (see Figure A1, Table A1), if a fish remained in that area they were disqualified from the data analysis. The sire identity in family block 5 was unknown and therefore individuals from family block 5 were disqualified as a result. There were technical difficulties with the camera resulting in data missing for family blocks 8 and 19 from the morphometric analysis, however the swimming data was still used.

Immediately after the swimming trials, the fish were anaesthetized using MS-222, their weight was taken and pictures of the right side of their body were taken using a Canon (Power Shot A570 IS) digital camera. Morphometrics of each individual will be assessed using tpsDig2



software which uses 18 landmarks on the fish's body in order to assess and compare overall body shapes as well as caudal peduncal width and depth between hybrid groups (see Colborne et al. 2011) and partial warp scores generated using CoordGen8. This data will be used in combination with the swim flume data in order to determine if hybrid and pure strain fish differ in terms of body shape and ultimately swimming ability.

## Statistical analysis

### *Swimming Ability*

The Ucrit was run in an ANCOVA against cross type with mass as a covariate.

### *Morphometry*

The partial warp scores that were generated through CoordGen8 were then run through a Discriminant Function Analysis in R 3.2.1 using the MASS package (version 7.3-43). The scores generated were then analyzed using an ANOVA with cross type as the independent variable and the morphology scores as the dependent variable. This was then visually examined using thin plate splines created in tpsRegr. Any significant results were then analysed using a Tukey's post hoc test.

## **Results**

### *Swimming Ability*

The ANCOVA demonstrated that there was no significant difference between Ucrit of the four cross types ( $F=1.84$ ,  $p=0.37$ ) out of the 324 individuals used in the study.

## *Morphometry*

Out of the 32 partial warp scores of  $n=323$  individuals that were generated and put through the DFA, three significant discriminant factors were produced. A one way ANOVA displayed significant results in body morphology explained by cross type with the first discriminant factor ( $F=21.93$ ,  $p<0.001$ ). A Tukey post hoc comparison revealed significant difference between SL and LL ( $p<0.001$ ), SS and LL ( $p<0.001$ ), SL and LS ( $p=0.007$ ) as well as SS and LS ( $p<0.001$ ). The second discriminant factor also displayed a significant difference between cross types ( $F=13.09$ ,  $p<0.001$ ). A Tukey's post hoc analysis revealed differences between the LS and LL ( $p<0.001$ ), SL and LS ( $p<0.001$ ) and SS and SL ( $p=0.003$ ). There was also a significant difference between cross types in the third discriminant factor ( $F=12.84$ ,  $p<0.001$ ) with a Tukey post hoc revealing differences between LS and LL ( $p<0.001$ ), SL and LL ( $p<0.001$ ) and SS and LS ( $p<0.001$ ) and SS and SL ( $p<0.001$ ).

## **Conclusion**

In conclusion, all groups differed over various morphological traits. The first discriminant factor differed most strongly between the two pure strains with the reciprocal hybrids being intermediate. The difference with the second discriminant factor seemed to be between groups with different sires. This further supports Chapter 2 in demonstrating the effects of hybridization in the  $F_1$  generation as primarily being the loss of local adaptations through intermediate phenotypes, and may also provide some evidence of dam strain and sire strain effects affecting juvenile morphology.

## **Acknowledgments**

This study was funded by the NSERC Strategic Project Grant (to TEP and CCW), NSERC Discovery Grant (to TEP), CFI (to TEP), the Ontario Ministry of Economic Development and Innovation (to TEP) as well as the Ontario Ministry of Natural Resources and Forestry (MNRF). We would like to thank Bill Sloan and Scott Ferguson from the MNRF Codrington Fisheries Research Facility for providing technical expertise, Craig Black for helping to initiate this research and Katelynn Johnson for helping to produce the crosses. We would also like to thank the Great Lakes Fish Research Centre in LaSalle for providing the logistical support in order to conduct the research as well as Jessica Mayrand and Jade Laycock for helping with the swim trials. Lastly, we would like to thank my colleagues and lab members Jason Lewis, Adriana Forest and Sarah Lehnert for their assistance and support during this project.

## **References**

- Boily P, Magnan P (2002) Relationship between individual variation in morphological characters and swimming costs in brook charr (*Salvelinus fontinalis*) and yellow perch (*Perca flavescens*). *The Journal of Experimental Biology* 205: 1031-1036
- Colborne SF, Bellemare MC, Peres-Neto PR, Neff BD (2011) Morphological and swim performance variation among reproductive tactics of bluegill sunfish (*Lepomis macrochirus*). *C J Fish Aquat Sci* 68 :1802-1810
- Dimond P, Smitka J (2005) Evaluation of Selected Strains of Atlantic Salmon as Potential Candidates for the Restoration of Lake Ontario. Trout Unlimited Canada Technical Report ON-12. Guelph, Ontario: Trout Unlimited Canada

Drinan TJ, McGinnity P, Coughlan JP, Cross TF, Harrison SSC (2012) Morphological variability of Atlantic salmon *Salmo salar* and brown trout *Salmo trutta* in different river environments.

Ecology of Freshwater Fish 21:420-432

Edmands S (2007) Between a rock and a hard place: evaluating the relative risks of inbreeding and outbreeding for conservation and management. Mol Ecol 16: 463-475

Frankham R, Ballou JD, and Briscoe DA (2002) Introduction to Conservation Genetics. Cambridge University Press, UK.

Lynch M (1991) The Genetic Interpretation of Inbreeding Depression and Outbreeding Depression. Evolution 45: 622-629

Ontario Ministry of Natural Resources (2009) Lake Ontario Fish Communities and Fisheries: 2008 Annual Report of the Lake Ontario Management Unit. Ontario Ministry of Natural Resources, Picton, Ontario, Canada.

Ricker WE (1975) Computation and interpretation of biological statistics of fish populations. Bulletin of the Fisheries Research Board of Canada 191: 1-382

Taylor EB (1986) Differences in morphology between wild and hatchery populations of juvenile Coho salmon. The Progressive Fish-Culturist 48: 171–176.

Taylor EB (1991) A review of local adaptation in Salmonidae, with particular reference to Pacific and Atlantic salmon. Aquaculture 98: 185–207

Turan C (2000) Otolith shape and meristic analysis of Herring (*Clupea harengus*) in the northeast Atlantic. Archive of Fishery and Marine Research 48: 283–295

Table A1: Mean Velocity in each portion of the swim flume. Section 6 (in bold) had a significantly lower velocity than the other sections.

| Section  | Hz | Mean Velocity   |
|----------|----|-----------------|
| 1        | 10 | 0.275842        |
| 1        | 15 | 0.415576        |
| 1        | 20 | 0.562296        |
| 2        | 10 | 0.271184        |
| 2        | 15 | 0.408589        |
| 2        | 20 | 0.556473        |
| 3        | 10 | 0.257211        |
| 3        | 15 | 0.386464        |
| 3        | 20 | 0.508731        |
| 4        | 10 | 0.273513        |
| 4        | 15 | 0.413247        |
| 4        | 20 | 0.555309        |
| 5        | 10 | 0.271184        |
| 5        | 15 | 0.421398        |
| 5        | 20 | 0.557638        |
| <b>6</b> | 10 | <b>0.242073</b> |
| <b>6</b> | 15 | <b>0.36434</b>  |
| <b>6</b> | 20 | <b>0.47962</b>  |
| 7        | 10 | 0.267691        |
| 7        | 15 | 0.410918        |
| 7        | 20 | 0.549487        |
| 8        | 10 | 0.264198        |
| 8        | 15 | 0.410918        |
| 8        | 20 | 0.550651        |
| 9        | 10 | 0.230429        |
| 9        | 15 | 0.355024        |
| 9        | 20 | 0.473798        |

## Figure Captions

Figure A1: Graphical representation of the nine zones in which water velocity was measured in the Loligo Swim-30 swim flume used in the study.

Figure A2: Graphical display of the morphological scores of the four different cross types (pure LaHave, pure Sebago, Sebago dam x LaHave sire and LaHave dam x Sebago sire) created using a Discriminant Function Analysis

**Figure A1**





Figure A2





Appendix 7 Female Raw Data as presented per individual. Wet and dry weight is measured on a per egg basis. Fecundity was calculated by multiplying the number of eggs per litre produced by the female with the number of litres she produced. Fulton's condition factor was measured as mass over length<sup>3</sup>. Multilocus heterozygosity was measured as a percentage of heterozygous loci. Egg diameter (mm) with one standard error, the coefficient of variation was calculated as the standard deviation/mean then converted to a percentage.

| Female ID | Age (years) | Fork Length (inches) | Weight (g) | Fulton's Condition factor in g/cm <sup>3</sup> | MLH % | Mean Egg diameter (mm) ± 1 S.E. | Mean Wet Egg weight (mg)±1 C.V. | Mean dry egg weight (mg) ± 1 C.V. | Fecundity |
|-----------|-------------|----------------------|------------|--|-------|---------------------------------|---------------------------------|-----------------------------------|-----------|
| LHF1      | 5           | 26                   | 4300       | 0.0149   | 0.842 | 6.04±0.052                      | 162.57±2.529                    | 64.123±9.942                      | 6978      |
| LHF2      | 5           | 26.5                 | 5315       | 0.0174   | 1.000 | 6.40±0.050                      | 178.22±1.245                    | 76.81±9.077                       | 6328.4    |
| LHF3      | 5           | 22                   | 3022       | 0.0173   | 0.611 | 5.98±0.062                      | 152.14±1.779                    | 53.283±25.221                     | 3857.7    |
| LHF4      | 5           | 25                   | 3975       | 0.0155   | 0.895 | 5.88±0.035                      | 139.35±1.750                    | 45.093±4.584                      | 5511      |
| LHF5      | 5           | 24                   | 2140       | 0.0095   | 1.000 | 5.52±0.059                      | 111.32±3.134                    | 38.563±4.988                      | 6062.1    |
| LHF6      | 5           | 25                   | 3014       | 0.0118   | 0.684 | 6.26±0.051                      | 163.38±1.794                    | 75.325±13.095                     | 3894.4    |
| LHF7      | 5           | 24                   | 2869       | 0.0127   | 0.944 | 6.01±0.054                      | 138.16±1.775                    | 51.233±7.490                      | 4070.5    |
| LHF9      | 5           | 26                   | 4909       | 0.0170   | 0.947 | 6.11±0.093                      | 160.44±4.665                    | 54.004±5.663                      | 8275.2    |
| LHF10     | 5           | 18                   | 1335       | 0.0140   | 0.944 | 5.80±0.036                      | 139.17±2.657                    | 51.77±6.631                       | 1653.3    |
| LHF11     | 7           | 29.5                 | 6185       | 0.0147   | 0.895 | 6.09±0.060                      | 149.15±2.606                    | 53.803±10.679                     | 8266.5    |
| LHF13     | 7           | 27                   | 6210       | 0.0193   | 0.563 | 6.63±0.094                      | 205.59±3.993                    | 74.56±10.547                      | 4426.65   |
| LHF14     | 7           | 22                   | 3201       | 0.0184   | 0.842 | 6.07±0.065                      | 158.01±2.918                    | 67.043±2.576                      | 2413.8    |
| LHF15     | 7           | 25                   | 4490       | 0.0175   | 0.737 | 6.50±0.111                      | 160.26±5.628                    | 43.855±11.432                     | 4327      |
| LHF16     | 7           | 26                   | 5220       | 0.0181   | 0.842 | 7.03±0.093                      | 220.82±5.201                    | 69.273±2.669                      | 6034.5    |
| LHF17     | 7           | 25                   | 4290       | 0.0168   | 0.895 | 6.40±0.037                      | 166.65±1.663                    | 55.883±4.759                      | 5020.4    |
| LHF18     | 7           | 28.5                 | 6065       | 0.0160   | 0.842 | 6.56±0.046                      | 177.93±1.876                    | 76.873±11.603                     | 6274.15   |
| LHF19     | 7           | 26                   | 5010       | 0.0174   | 1.000 | 6.65±0.041                      | 196.65±5.521                    | 71.67±3.659                       | 4502.4    |
| LHF20     | 7           | 27                   | 6211       | 0.0193   | 0.737 | 6.33±0.038                      | 166.46±7.289                    | 54.05±5.867                       | 6723.6    |
| LHF21     | 7           | 29                   | 6540       | 0.0164   | 0.769 | 6.64±0.043                      | 190.12±2.843                    | 72.203±1.036                      | 8254.4    |
| LHF23     | 7           | 29                   | 6100       | 0.0153   | 0.842 | 6.38±0.043                      | 177.65±3.585                    | 60.693±4.823                      | 6815.2    |
| LHF24     | 7           | 22                   | 3495       | 0.0200   | 0.684 | 6.26±0.054                      | 168.05±1.397                    | 56.403±3.546                      | 4868      |
| LHF25     | 7           | 31                   | 7927       | 0.0162   | 0.632 | 6.38±0.052                      | 161.98±1.720                    | 61.13±1.0642                      | 8654      |
| LHF26     | 7           | 24                   | 4010       | 0.0177   | 0.889 | 6.68±0.054                      | 216.70±13.324                   | NA                                | 4219.2    |

|              |   |      |      |        |       |            |              |                |        |
|--------------|---|------|------|--------|-------|------------|--------------|----------------|--------|
| <b>LHF27</b> | 7 | 27   | 5114 | 0.0159 | 0.842 | 6.34±0.053 | 166.46±3.900 | 66.767±12.3830 | 5192.4 |
| <b>LHF28</b> | 7 | 27   | 5960 | 0.0185 | 0.684 | 6.08±0.060 | 137.94±2.010 | 50.417±5.380   | 7302.4 |
| <b>LHF29</b> | 7 | 23   | 3359 | 0.0169 | 0.632 | 6.26±0.040 | 174.59±3.034 | 64.688±2.714   | 3620.7 |
| <b>LHF30</b> | 7 | 27   | 5065 | 0.0157 | 0.895 | 5.99±0.058 | 151.59±4.529 | 49.927±3.255   | 7164.3 |
| <b>LHF31</b> | 7 | 26   | 4760 | 0.0165 | 0.790 | 6.37±0.047 | 155.80±1.503 | 54.893±3.312   | 4502.4 |
| <b>LHF32</b> | 7 | 30   | 6290 | 0.0142 | 0.790 | 6.13±0.059 | 154.65±3.615 | 57.73±6.643    | 6206.4 |
| <b>LHF33</b> | 6 | 27   | 4727 | 0.0147 | 0.842 | 6.48±0.054 | 186.10±2.538 | 72.837±10.759  | 6389.6 |
| <b>LHF34</b> | 6 | 24   | 4580 | 0.0202 | 0.790 | 5.98±0.038 | 154.79±1.244 | 55.553±3.740   | 7240.8 |
| <b>LHF35</b> | 6 | 26   | 4593 | 0.0160 | 0.842 | 6.39±0.043 | 196.91±2.220 | 66.181±4.576   | 6490.5 |
| <b>LHF36</b> | 6 | 26   | 5130 | 0.0178 | 0.790 | 6.46±0.063 | 185.46±4.293 | 63.705±2.298   | 7240.8 |
| <b>LHF37</b> | 6 | 26   | 3828 | 0.0133 | 0.947 | 5.91±0.057 | 150.03±3.867 | 55.24±0.205    | 7164.3 |
| <b>LHF38</b> | 6 | 24   | 3690 | 0.0163 | 0.737 | 6.32±0.057 | 166.34±3.694 | 54.231±3.180   | 4759.7 |
| <b>LHF39</b> | 6 | 28   | 5609 | 0.0156 | 0.947 | 6.39±0.039 | 182.26±1.883 | 58.463±8.014   | 7302   |
| <b>LHF40</b> | 6 | 28   | 6020 | 0.0167 | 0.790 | 6.66±0.039 | 211.26±2.416 | 76.207±6.727   | 8046   |
| <b>LHF41</b> | 6 | 20   | 2125 | 0.0162 | 0.895 | 6.09±0.060 | 163.09±0.326 | 55.527±9.803   | 2326   |
| <b>LHF42</b> | 6 | 20   | 2030 | 0.0155 | 0.684 | 5.72±0.044 | 130.60±4.664 | 47.32±14.319   | 1957.5 |
| <b>LHF43</b> | 6 | 22.5 | 3387 | 0.0182 | 0.842 | 6.09±0.059 | 168.61±1.321 | 51.673±3.710   | 5689.2 |
| <b>LHF44</b> | 6 | 26   | 5200 | 0.0181 | 0.889 | 6.75±0.102 | 184.55±NA    | 58.965±4.665   | 6815.2 |

Appendix 8 Male Raw Data All sperm quality values are the mean taken between all measured tracks. Fulton's condition factor ( $\text{g}/\text{cm}^3$ ), multilocus heterozygosity was the percentage of heterozygous loci per individual, average path, curvilinear and progressive velocity were measured as units of distance per second traveled, linearity is the straightness with which the cell travelled per unit of distance, motility is the percentage of cells in a recording that are motile, longevity is the point when which 95% of the cells died and density is the number of cells per 1ml of milt.

| Male ID | Age (years) | Fork Length (inches) | Weight (g) | Fulton's condition factor ( $\text{g}/\text{cm}^3$ ) | MLH % | VAP 5s ( $\mu\text{m}/\text{s}$ ) | VAP 10s ( $\mu\text{m}/\text{s}$ ) | VCL 5s ( $\mu\text{m}/\text{s}$ ) | VCL 10s ( $\mu\text{m}/\text{s}$ ) | VSL 5s ( $\mu\text{m}/\text{s}$ ) | VSL 10s ( $\mu\text{m}/\text{s}$ ) | Lin 5s ( $^\circ/\mu\text{m}$ ) | Lin 10s ( $^\circ/\mu\text{m}$ ) | %Motility | Longevity (s) | Density |
|---------|-------------|----------------------|------------|--|-------|-----------------------------------|------------------------------------|-----------------------------------|------------------------------------|-----------------------------------|------------------------------------|---------------------------------|----------------------------------|-----------|---------------|---------|
| LH1     | 7           | 23                   | 3128       | 0.0157   | 0.895 | 122.95                            | 78.4                               | 134.3                             | 85.65                              | 78.15                             | 50.85                              | 57.5                            | 60.5                             | 0.56423   | 31            | 6656250 |
| LH2     | 7           | 22                   | 2517       | 0.0144   | 0.789 | 142.3                             | 94.35                              | 159.3                             | 105.25                             | 76.8                              | 51.3                               | 49                              | 50.5                             | 0.78190   | 36.5          | 1437500 |
| LH4     | 7           | 28                   | 5594       | 0.0156   | 0.737 | 110.1                             | 75.1                               | 124.3                             | 83                                 | 54.5                              | 44.6                               | 45                              | 57                               | 0.5749    | 25.5          | 593750  |
| LH5     | 7           | 30                   | 7430       | 0.0168   | 0.778 | 150.65                            | 90.3                               | 161.05                            | 95.95                              | 113.75                            | 64.25                              | 69                              | 66.5                             | 0.6302    | 34.25         | 812500  |
| LH6     | 7           | 32                   | 7858       | 0.0146   | 0.737 | 135.05                            | 64.85                              | 149.25                            | 69.45                              | 79.85                             | 51.3                               | 53.5                            | 75                               | 0.71421   | 32.5          | 343750  |
| LH7     | 7           | 19                   | 1722       | 0.0153   | 0.789 | 178                               | 103                                | 181.55                            | 105.45                             | 150.2                             | 97.35                              | 83                              | 91.5                             | 0.74125   | 31.5          | 3906250 |
| 123 LH8 | 7           | 48                   | 4979       | 0.0027   | 0.778 | 181.35                            | 93.1                               | 190.7                             | 98.15                              | 160.4                             | 86.2                               | 83                              | 89                               | 0.86124   | 40.75         | 1625000 |
| LH9     | 7           | 18                   | 1638       | 0.0171   | 0.842 | 59.2                              | 50.25                              | 64.75                             | 56.2                               | 48.7                              | 45                                 | 76.5                            | 78.5                             | 0.47099   | 21.5          | 1500000 |
| LH10    | 7           | 34                   | 7467       | 0.0116   | 0.842 | 61.85                             | 47.8                               | 69                                | 66.1                               | 50.8                              | 41.5                               | 77.5                            | 68                               | 0.65492   | 21            | 187500  |
| LH11    | 6           | 22                   | 2406       | 0.0138   | 0.684 | 136.1                             | 58.05                              | 143.9                             | 65.15                              | 119.75                            | 52.6                               | 83.5                            | 85                               | 0.91022   | 25            | 1375000 |
| LH12    | 6           | 18                   | 1515       | 0.0159   | 0.842 | 49.8                              | 40.5                               | 56.65                             | 46.45                              | 39.95                             | 34.7                               | 73.5                            | 76                               | 0.72655   | 22.25         | 2656250 |
| LH13    | 6           | 19                   | 1388       | 0.0123   | 0.684 | 140.85                            | 76.55                              | 149.95                            | 80.25                              | 122.2                             | 68.75                              | 79.5                            | 85.5                             | 0.52394   | 36.5          | 2750000 |
| LH14    | 6           | 24                   | 2417       | 0.0107   | 0.737 | 157.3                             | 104.85                             | 170.15                            | 110.05                             | 104.1                             | 74.6                               | 60                              | 68                               | 0.79600   | 33.75         | 2250000 |
| LH16    | 6           | 22                   | 2171       | 0.0124   | 0.789 | 83.4                              | 60.75                              | 91.35                             | 68.35                              | 64.05                             | 50.75                              | 71.5                            | 76.5                             | 0.58461   | 22            | 1125000 |
| LH18    | 6           | 30                   | 5747       | 0.0130   | 0.684 | 109.85                            | 67.4                               | 115.5                             | 70.9                               | 78.05                             | 55.05                              | 70                              | 78.5                             | 0.39814   | 20.25         | 343750  |
| LH19    | 6           | 20                   | 1707       | 0.0130   | 0.737 | 69.7                              | 60.2                               | 78.3                              | 67.2                               | 52.7                              | 50.6                               | 69                              | 77                               | 0.32768   | 23.5          | 1343750 |
| LH20    | 6           | 22                   | 3002       | 0.0172   | 0.895 | 61.75                             | 50.8                               | 68                                | 56.15                              | 50.95                             | 44.05                              | 75.5                            | 79                               | 0.50049   | 25.75         | 687500  |
| LH21    | 6           | 30                   | 6509       | 0.0147   | 0.895 | 83.3                              | 60.4                               | 89.15                             | 64.35                              | 70.85                             | 44.1                               | 78.5                            | 70                               | 0.38421   | 21            | 4406250 |
| LH22    | 6           | 24                   | 3665       | 0.0162   | 0.842 | 90.15                             | 56.15                              | 95.6                              | 61.4                               | 80.05                             | 50.05                              | 83.5                            | 80                               | 0.32591   | 20.25         | 843750  |
| LH23    | 6           | 22                   | 2215       | 0.0127   | 0.895 | 122.5                             | 87.4                               | 131.75                            | 95.15                              | 103.9                             | 72.8                               | 77                              | 77                               | 0.36927   | 22.5          | 2687500 |
| LH24    | 6           | 23                   | 2504       | 0.0126   | 0.684 | 118.7                             | 83.3                               | 134.3                             | 91.25                              | 84.25                             | 62.5                               | 63.5                            | 69.5                             | 0.39023   | 29.75         | 2718750 |
| LH27    | 6           | 23                   | 3272       | 0.0164   | 0.789 | 156.95                            | 61.35                              | 164.95                            | 63.8                               | 125.8                             | 57.15                              | 75                              | 89                               | 0.75598   | 22            | 3250000 |
| LH28    | 6           | 26                   | 3856       | 0.0134   | 0.789 | 43.25                             | 30.1                               | 47.4                              | 35.5                               | 38.35                             | 27.7                               | 82                              | 79                               | 0.65891   | 14.25         | 1531250 |

|             |   |    |       |        |       |        |       |        |       |        |       |      |      |         |       |          |
|-------------|---|----|-------|--------|-------|--------|-------|--------|-------|--------|-------|------|------|---------|-------|----------|
| <b>LH29</b> | 6 | 26 | 4489  | 0.0156 | 0.737 | 181.2  | 82.8  | 188.6  | 85.95 | 131    | 72.45 | 68.5 | 84.5 | 0.81442 | 36.25 | 7937500  |
| <b>LH30</b> | 6 | 28 | 2845  | 0.0079 | 0.842 | 87.05  | 61.55 | 97.45  | 67.15 | 68.15  | 49.55 | 72.5 | 76   | 0.67428 | 19.75 | 16250000 |
| <b>LH31</b> | 6 | 26 | 3825  | 0.0133 | 0.789 | 91.3   | 60.95 | 98.1   | 64.55 | 65.55  | 50.4  | 67.5 | 79   | 0.73670 | 27    | 3750000  |
| <b>LH32</b> | 6 | 30 | 5768  | 0.0130 | 0.789 | 40.5   | 42.85 | 44.5   | 47    | 37.05  | 37.95 | 83.5 | 82.5 | 0.51968 | 20.5  | 1125000  |
| <b>LH33</b> | 6 | 30 | 5388  | 0.0122 | 0.632 | 84.05  | 63.4  | 90.3   | 68.5  | 77.05  | 56.45 | 86.5 | 81   | 0.52173 | 30.75 | 343750   |
| <b>LH34</b> | 6 | 30 | 5685  | 0.0128 | 0.737 | 84.4   | 52.9  | 90.55  | 58.2  | 77.25  | 46.8  | 85   | 81   | 0.5643  | 22.25 | 1906250  |
| <b>LH35</b> | 6 | 21 | 2109  | 0.0139 | 0.632 | 114.2  | 64.85 | 121.95 | 69.95 | 97.25  | 57.45 | 79.5 | 83   | 0.53106 | 30    | 437500   |
| <b>LH36</b> | 6 | 31 | 6546  | 0.0134 | 0.684 | 48.7   | 39.1  | 56.2   | 46.1  | 43.3   | 37.5  | 79   | 82   | 0.63636 | 19    | 500000   |
| <b>LH37</b> | 6 | 32 | 7567  | 0.0141 | 0.895 | 56.45  | 39    | 62.3   | 48.1  | 47     | 34.6  | 77   | 71.5 | 0.46666 | 18.75 | 1937500  |
| <b>LH38</b> | 6 | 18 | 1310  | 0.0137 | 0.737 | 135.9  | 65.9  | 157.5  | 70.9  | 105.4  | 60.8  | 66   | 84   | 0.73437 | 33    | 15906250 |
| <b>LH39</b> | 6 | 36 | 10409 | 0.0136 | 0.895 | 96.45  | 57.1  | 103.85 | 61.6  | 69.2   | 43.75 | 69   | 73.5 | 0.63146 | 23.5  | 2031250  |
| <b>LH40</b> | 6 | 18 | 1723  | 0.0180 | 0.895 | 127.6  | 51.9  | 143.6  | 58.5  | 93     | 46.45 | 63.5 | 80   | 0.67729 | 30.25 | 11500000 |
| <b>LH41</b> | 6 | 22 | 2717  | 0.0156 | 0.895 | 89.6   | 57.25 | 100.4  | 64.3  | 69.3   | 47.3  | 71   | 73.5 | 0.79129 | 27.5  | 812500   |
| <b>LH42</b> | 6 | 27 | 4036  | 0.0125 | 0.733 | 141.75 | 55    | 154.3  | 59    | 118.95 | 50.9  | 76.5 | 85.5 | 0.77419 | 31.75 | 2625000  |
| <b>LH43</b> | 6 | 29 | 6192  | 0.0155 | 0.895 | 47     | 35.8  | 51.3   | 41.2  | 44.9   | 33.6  | 87   | 82   | 0.33333 | 22.5  | 437500   |

Appendix 9 : Summary of the 19 tetra and dinucleotide loci used in the study originally derived from the studies O'Reilly et al. 1996, Paterson et al. 2004, King et al. 2005 and Olafsson et al. 2010. Listed in the primer sequence as well as the number of base pairs. (For further information see referenced sources).

| Locus    | Primer sequence 5'-3'   | Base Pairs | GenBank accession number | Authors              |
|----------|---|------------|--------------------------|----------------------|
| AY081812 | F: CGCAATGGAAGTCAGTGGACTGG<br>R: CTGATTTAGCTTTTTAGTGCCCAATGC          | 438bp      | AY081812.1               | Paterson et al. 2004 |
| AY081807 | F: TTTAGATGGTGGGATACTGGGAGGC<br>R: CGGGAGCCCCATAACCCTACTAATAAC        | 520bp      | AY081807.1               | Paterson et al. 2004 |
| AY081810 | F: ACTAGCCAGGTGTCCTGCCGGTC<br>R: AGGGTCAGTCAGTCACACCATGCAC            | 461bp      | AY081810.1               | Paterson et al. 2004 |
| SSU43694 | (a) GGG TTG AGT AGG GAG GCT TG<br>(b) TGG CAG GGA TTT GAC ATA AC*     | 473bp      | U43694.1                 | O'Reilly et al. 1996 |
| SSU43695 | (a) CTT GGA ATA TCT AGA ATA TGG C<br>(b) TTC ATG TGT TAA TGT TGC GTG* | 344bp      | U43695.1                 | O'Reilly et al. 1996 |
| AY081813 | F: CTTGGTCCCGTTCTTACGACAACC<br>R: TGCACGCTGCTTGGTCCTTG                | 613bp      | AY081813.2               | Paterson et al. 2004 |
| AY081809 | F: ATGTGGAGGTCAACTAACCAGCGTG<br>R: CATCAATCACAGAGTGAGGCACTCG          | 520bp      | AY081809.1               | Paterson et al. 2004 |
| AY081811 | F: GGCCCAGACAGATAAAACAAACACGC<br>R: GCCAACAGCAGCATCTACACCCAG          | 520bp      | AY081811.1               | Paterson et al. 2004 |
| AF525202 | F: CTCCTGCACCTGACTTCTATTC<br>R: ACAGGCTATCACAGAACAGTTG                | 410bp      | AF525202.1               | King et al. 2005     |
| AF525206 | F: GGCATTGGAGGTAAGGACAC<br>R: CCAGACCACTGAACTTCTCATC                  | 406bp      | AF525206.1               | King et al. 2005     |
| SsaF43   | Missing   | 136bp      | NA                       | Olafsson et al. 2010 |
| AY081808 | F: AAGTATTCATGCACACACATTCCTGC<br>R: CAAGACCCTTTTTCCAATGGGATTC         | 349bp      | AY081808.1               | Paterson et al. 2004 |
| AF525201 | F: TCTGGAAGTTTCCCTACTTCTG<br>R: TCTTTAACTGTTGCCTTAACGAC               | 571bp      | AF525201.1               | King et al. 2005     |
| AF525204 | F: ATCGAAATGGAACCTTTTGAATG  | 451bp      | AF525204.1               | King et al. 2005     |

---

|          |   |       |            |                      |
|----------|---|-------|------------|----------------------|
|          | R: GCTTAGGGCTGAGAGAGGAATAC  |       |            |                      |
| AF525208 | F: TCGCTGTGTATCAGTATTTTGG<br>R: ACTCGGATAAACTCACAGGTC                 | 544bp | AF525208.1 | King et al. 2005     |
| AF525210 | F: TAGAGTTTGTCTCTGGCTTTG<br>R: AGACCCTAGGACTGGCTACTG                  | 550bp | AF525210.1 | King et al. 2005     |
| SSU43693 | (a) TTA TTA TCC AAA GGG GTC AAA A<br>(b) GAG GTC GCT GGG GTT TAC TAT* | 436bp | U43693.1   | O'Reilly et al. 1996 |
| AF525200 | F: TCTCCAGTGGTTCTAGATGAG<br>R: GGAGCTAAACTTCAAAGCACAG                 | 522bp | AF525200.1 | King et al. 2005     |
| AF525203 | F: TTGTGAAGGGGCTGACTAAC<br>R: TCAATTGTTGGGTGCACATAG                   | 439bp | AF525203.1 | King et al. 2005     |

---

## APPENDIX 10: WRITTEN PERMISSION FROM CO-AUTHORS

Wilson, Chris (MNRF) <[chris.wilson@ontario.ca](mailto:chris.wilson@ontario.ca)>

Aug 18 (11  
days ago)

to Trevor, me

Agreed – please use this as my permission to use both the hybrid and heterozygosity data for your thesis.

Cheers,

Chris.

---

Chris Wilson  
Aquatic Research and Monitoring Section  
Ontario Ministry of Natural Resources and Forestry  
Trent University  
2140 East Bank Drive,  
Peterborough, ON K9J 7B8  
Canada  
[\(705\) 755-2260](tel:7057552260) (office)  
[\(705\) 755-2261](tel:7057552261) (lab)  
[\(705\) 755-1559](tel:7057551559) (fax)

**From:** Trevor Pitcher [mailto:[tpitcher@uwindsor.ca](mailto:tpitcher@uwindsor.ca)]

**Sent:** August-18-15 3:30 PM

**To:** Chantal Audet

**Cc:** Wilson, Chris (MNRF)

**Subject:** Re: Written Approval to Deposit Thesis

Chantal,

Consider this email my permission to use the data collected for chapter 2 and 3 in your thesis.

Best,

Trevor

---

Trevor E. Pitcher, PhD

Associate Professor

Department of Biological Sciences & Great Lakes Institute for Environmental Research

University of Windsor

Windsor, Ontario

Canada, N9B 3P4

Phone: [519-253-3000](tel:519-253-3000) ext. 2710

Email: [tpitcher@uwindsor.ca](mailto:tpitcher@uwindsor.ca)

Web: [www.uwindsor.ca/pitcher](http://www.uwindsor.ca/pitcher)



## VITA AUCTORIS

Name: Chantal Lianne Audet

Place of Birth: Sudbury, Ontario

Year of Birth: 1989

EDUCATION: École Secondaire du Sacré-Cœur

Sudbury, Ontario

2003-2007

Université Laurentienne/ Laurentian University

Sudbury, Ontario

2007-2012

B.Sc. Honours in Integrated Biology

University of Windsor

Windsor, Ontario

2013-2015

M.Sc. Biological Sciences