

Exact and Perturbed Friedmann-Lemaître Cosmologies

by

Paul Ullrich

A thesis
presented to the University of Waterloo
in fulfilment of the
thesis requirement for the degree of
Master of Mathematics
in
Applied Mathematics

Waterloo, Ontario, Canada, 2007

© Paul Ullrich 2007

I hereby declare that I am the sole author of this thesis. This is a true copy of the thesis, including any required final revisions, as accepted by my examiners.

I understand that my thesis may be made electronically available to the public.

Abstract

In this thesis we first apply the 1+3 covariant description of general relativity to analyze n -fluid Friedmann-Lemaître (FL) cosmologies; that is, homogeneous and isotropic cosmologies whose matter-energy content consists of n non-interacting fluids. We are motivated to study FL models of this type as observations suggest the physical universe is closely described by a FL model with a matter content consisting of radiation, dust and a cosmological constant. Secondly, we use the 1 + 3 covariant description to analyse scalar, vector and tensor perturbations of FL cosmologies containing a perfect fluid and a cosmological constant. In particular, we provide a thorough discussion of the behaviour of perturbations in the physically interesting cases of a dust or radiation background.

Acknowledgements

First and foremost, I would like to thank Dr. John Wainwright for his patience, guidance and support throughout the preparation of this thesis. I also would like to thank Dr. Achim Kempf and Dr. C. G. Hewitt for their time and comments.

Contents

1	Introduction	1
1.1	Cosmological Models	1
1.2	Geometrical Background	3
1.2.1	Projection and Differentiation	4
1.2.2	Kinematic Quantities	5
1.2.3	Decomposition of the Energy-Momentum Tensor	7
1.2.4	The Weyl Curvature Tensor	7
1.2.5	Spatial Curvature	8
1.2.6	Evolution and Constraint Equations	10
1.3	Overview of the Thesis	11
2	The Friedmann-Lemaître Cosmologies	13
2.1	Properties of the RW Metric	13
2.2	Characterizations of the FL Cosmologies	17
2.2.1	The Kinematic Characterization	17
2.2.2	The Weyl Characterization	18
3	Dynamics of n-Fluid FL Cosmologies	20
3.1	Behaviour and Classification of n -Fluid FL Cosmologies	20
3.1.1	Basic Evolution Equations	21
3.1.2	n -Fluid FL Cosmologies	23

3.1.3	Hubble-Normalized Scalars	26
3.1.4	A Compact State Space for n -fluid FL Cosmologies	27
3.1.5	Invariant Sets and Equilibrium Points in the Compact State Space	32
3.1.6	Classification of n -Fluid FL Cosmologies	33
3.1.7	Asymptotic Behaviour of Solutions	35
3.2	Parameterization of FL Models	43
3.2.1	The Intrinsic Parameters	43
3.2.2	The Observational Parameters	45
3.2.3	Conserved Quantities	47
3.3	2-Fluid FL Cosmologies - Qualitative Analysis	49
3.3.1	General Features	49
3.3.2	FL Cosmologies with Radiation and Dust (RDC -universes)	51
3.3.3	FL Cosmologies with Dust and Cosmological Constant	53
3.4	3-Fluid FL Cosmologies - Qualitative Analysis	56
3.4.1	General Features	56
3.4.2	A Model of the Physical Universe	59
3.5	Explicit FL Cosmologies	63
3.5.1	Single Fluid FL Cosmologies	64
3.5.2	FL Vacuum Cosmologies	66
3.5.3	Flat FL with a Single Fluid and Λ	67
3.5.4	FL Cosmologies with Radiation and Dust (RDC -Universes)	68
3.5.5	FL Cosmologies with Radiation and Λ (RCA -Universes)	69
3.5.6	Explicit Solutions in Terms of Observational Parameters	71
3.6	Discussion	72
4	Perturbations of FL Models	75
4.1	Historical Development	75
4.1.1	Metric Approach	75
4.1.2	Geometrical Approach	77
4.2	The Linearized Einstein Field Equations	78
4.2.1	The Linearization Process	79
4.2.2	The Linearized Evolution and Constraint Equations	80

4.3	Harmonic Decomposition	83
4.4	Scalar Perturbations	84
4.4.1	The Governing DE	84
4.4.2	Kinematic Quantities and Weyl Curvature	86
4.5	Vector Perturbations	87
4.5.1	The Governing DE	87
4.5.2	Kinematic Quantities and Weyl Curvature	88
4.6	Tensor Perturbations	89
4.6.1	The Governing DE	89
4.6.2	Kinematic Quantities and Weyl Curvature	90
4.7	Dynamics of the Linear Perturbations: General Features	91
4.7.1	Conformal Time and the Particle Horizon	91
4.7.2	Governing DEs Using Conformal Time	93
4.7.3	General Solutions for Scalar and Tensor Perturbations with $\Lambda = 0$	96
4.7.4	Functional Dependence of First-Order Quantities	98
4.8	Perturbations of FL Models with Pressure-Free Matter	101
4.8.1	The General Solution in Integral Form for Scalar Perturbations	101
4.8.2	Scalar Perturbations with $\Lambda = 0$	103
4.8.3	Scalar Perturbations with $\Lambda > 0, K = 0$	105
4.8.4	Tensor Perturbations for $\Lambda = 0$	106
4.9	Perturbations of Radiation-Filled FL Models	108
4.10	Perturbations in Asymptotic Epochs of FL Models	112
4.10.1	The Flat FL Asymptotic Epoch	114
4.10.2	The Milne Asymptotic Epoch	117
4.10.3	The de Sitter Asymptotic Epoch	119
4.11	Discussion	122
A	Evolution and Constraint Equations	124
B	Spatial Curvature	128
B.1	The Generalized Gauss Equation	128
B.2	The Spatial Gradient of the 3-Ricci Scalar	129

C	Solutions of the Friedmann Equation	131
C.1	Solutions with a Stiff Fluid	132
C.2	Existence-Uniqueness for the Friedmann Equation	134
D	Conformal Time	135
E	Covariant Differential Identities	138
F	Covariant Harmonics	141
F.1	Scalar Harmonics	142
F.2	Vector Harmonics	143
F.3	Tensor Harmonics	145
G	The 3-Cotton-York Tensor	147
G.1	Evolution Equation for the 3-Cotton-York Tensor	148
G.2	Tensor Perturbations	149
H	Bessel and Associated Legendre Functions	150
H.1	Normal Forms of Bessel and Associated Legendre DEs	150
H.2	Special Solutions of the Governing DEs	151
H.3	Resonant Solutions of the Legendre DE	153
H.4	Asymptotic Behaviour	154
	Bibliography	155

List of Figures

3.1	The 2-fluid FL state space for RDC -universes	52
3.2	The 2-fluid FL state space for DCA -universes	54
3.3	Thumbnail plots depicting behaviour of the length scale in FL cosmologies	55
3.4	The 3-fluid FL state space for radiation, dust and cosmological constant.	60
3.5	3-fluid FL state space cross-section along the expanding flat FL manifold	60
3.6	Past and future evolution of the $RDCA$ model of the physical universe	62
4.1	Conformal time and the particle horizon.	93
4.2	Scalar perturbation modes for dust and spatial curvature.	105
4.3	Scalar perturbation modes for dust and cosmological constant.	107
4.4	Tensor modes of the Hubble-normalized electric Weyl tensor in open FL with dust	109
4.5	Tensor modes of the Hubble-normalized electric Weyl tensor in open FL with radiation	111

List of Tables

3.1	The past and future attractors of generic n -fluid FL cosmologies.	42
3.2	Asymptotic behaviour of Hubble-normalized quantities in generic n -fluid FL cosmologies.	42
4.1	Functional dependence of first-order quantities	100
4.2	Perturbations of flat FL with $k\eta \ll 1$	115
4.3	Perturbations of flat FL, with $k\eta \gg 1$	116
4.4	Perturbations of Milne	118
4.5	Perturbations of de Sitter ($k\tilde{\eta} \ll 1$)	120
4.6	Perturbations of de Sitter ($k\tilde{\eta} \gg 1$)	121

INTRODUCTION

1.1 Cosmological Models

The goal of cosmology is to describe the large-scale structure, origins and evolution of the universe as a whole. Within the framework of general relativity, a *cosmological model*, denoted $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ consists of a differential manifold \mathcal{M} , a Lorentzian metric \mathbf{g} (which describes the geometry of the space-time) and a *unit timelike vector field* \mathbf{u} ,

$$g_{ij}u^i u^j = -1, \quad (1.1)$$

that defines the world-lines of a family of fundamental observers. In cosmology one represents the matter content as a smooth distribution (a continuum) whose particles represent galaxies in the present epoch. The time-like vector field \mathbf{u} is the 4-velocity of the particles in this continuum. One member of the congruence of time-like curves defined by \mathbf{u} , namely the world-line of our galaxy, plays a particularly important role, since it is the position from which we make observations.

The evolution of the universe is assumed to be governed by the *Einstein field equations (EFEs)*,

$$R_{ab} - \frac{1}{2}Rg_{ab} = T_{ab}, \quad (1.2)$$

where R_{ab} is the *Ricci tensor*, R is the *Ricci scalar* ($R = g^{ab}R_{ab}$) and T_{ab} is the *energy-momentum tensor* of the distribution of matter and energy in the universe. Here and throughout we will use geometrized units so that $c = 1$ and $8\pi G = 1$, where c is the speed of light in vacuum and G is the gravitational constant.

There are two fundamental assumptions that are used to construct models of the universe:

I. *The universe is isotropic about our galaxy.* More precisely, at any event on the world-line of our galaxy, all directions orthogonal to the world-line are physically and geometrically equivalent, when the universe is viewed on a sufficiently large scale.

A variety of cosmological observations (*e.g.* number counts of distant galaxies and peculiar velocity surveys) provide support for assumption I in the present epoch. The strongest support is provided by the temperature of the cosmic microwave background radiation (CMBR) which is isotropic to within 10^{-5} (*i.e.* $\frac{\Delta T}{T} \sim 10^{-5}$).

II. *The Copernican Principle.* This principle states that our galaxy is not in a preferred position in the universe. Unlike assumption I, this assumption is philosophical, rather than observational.

Assumptions I and II together imply that the universe satisfies the so-called *Cosmological Principle*, namely

- i) the universe is isotropic about all fundamental world-lines, and
- ii) the universe is spatially homogeneous, *i.e.* all physical and geometric quantities do not vary in space.

It is a well-known consequence of assumptions I and II (we give a concise proof in Chapter 2; see Theorem 2.1.1) that the metric g_{ab} can be written in the form of the *Robertson-Walker (RW) metric*,¹

$$ds^2 = -dt^2 + \ell^2(t)\tilde{g}_{\alpha\beta}(x^\lambda)dx^\alpha dx^\beta, \quad (1.3)$$

where $\tilde{g}_{\alpha\beta}$ is a 3-metric of constant curvature. The preferred timelike vector field \mathbf{u} is given by

$$\mathbf{u} = \frac{\partial}{\partial t}. \quad (1.4)$$

The Robertson-Walker metric provides the basis for the Friedmann-Lemaître (FL) cosmological models that play a central role in modern cosmology. A *Friedmann-Lemaître cosmology*² is a cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ that satisfies the Einstein field equations (1.2) with a suitable energy-momentum tensor T_{ab} , with metric \mathbf{g} and fundamental congruence \mathbf{u} having the Robertson-Walker form (1.3) and (1.4).

In this thesis we study FL cosmologies and their perturbations within a general mathematical framework. In section 1.2 we provide the necessary geometrical background for this study.

¹Here we use Greek indices to denote spatial indices. They can take on the values 1, 2 or 3.

²The first explicit FL cosmological models were first discovered by Friedmann in 1929 (for some historical details, see Ellis (1989)).

1.2 Geometrical Background

In this thesis we will make use of the so-called 1 + 3 *covariant description of general relativity*³ which has been developed for use in spacetimes in which there is a preferred timelike congruence \mathbf{u} , as in the case of a cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$. The “1+3” refers to the fact that one performs a “time + space” decomposition relative to \mathbf{u} by projecting tensors and tensorial equations parallel to \mathbf{u} and orthogonal to \mathbf{u} . The second aspect of the 1 + 3 description is to write a tensor as a sum of algebraically simpler parts, *i.e.* to give an algebraic decomposition. For example, an arbitrary rank 2 contravariant tensor can be written as the sum of a symmetric tracefree tensor, an anti-symmetric tensor and the trace multiplied by the metric tensor.

The first step is to introduce the basic variables in the 1 + 3 description, which are defined by applying projection relative to \mathbf{u} and algebraically decomposing the covariant derivative $\nabla_b u_a$, the energy-momentum tensor T_{ab} and the Riemann-Christoffel tensor as follows:

- i) The *covariant derivative of the fundamental 4-velocity*, $\nabla_b u_a$, describes the kinematics of the continuum described by the vector field \mathbf{u} . The so-called *kinematic quantities* are defined by writing $\nabla_b u_a$ as a sum of simpler parts and projecting relative to \mathbf{u} . We define these quantities in section 1.2.2.
- ii) The *energy-momentum tensor* T_{ab} describes the matter and energy content of the universe and so gives rise to a set of quantities known as *matter variables* when decomposed relative to \mathbf{u} . We discuss this decomposition in section 1.2.3.
- iii) The *Weyl curvature tensor* C_{abcd} , which is defined as the tracefree part of the Riemann curvature tensor, is decomposed relative to \mathbf{u} into two tracefree symmetric tensors, the so-called⁴ electric part E_{ab} and magnetic part H_{ab} . We discuss this decomposition in section 1.2.4.

The second step is to derive evolution and constraint equations for the basic variables by using the Ricci identities and Bianchi identities in conjunction with the Einstein field equations. We will discuss this process in section 1.2.6.

³See, for example, Ellis (1973), Wainwright and Ellis (1997), p1-30, or Ellis and van Elst (1998), Chapter 2.

⁴There is a formal analogy with the decomposition of the electromagnetic field tensor F_{ab} into an electric field E_a and a magnetic field H_a .

1.2.1 Projection and Differentiation

Projection parallel to \mathbf{u} is achieved by simply contracting a vector or tensor with u^a . Projection into the 3-space orthogonal to \mathbf{u} is defined by the *projection tensor*

$$h_{ab} = g_{ab} + u_a u_b, \quad (1.5)$$

which satisfies the identities

$$h_a^c h_c^b = h_a^b, \quad h_a^b u_b = 0, \quad h_a^a = 3.$$

Corresponding to these two types of projection, two derivatives are defined.

The overdot $\dot{}$ denotes differentiation along the fundamental congruence, defined for scalars as

$$\dot{f} = f_{,a} u^a, \quad (1.6)$$

and for tensors as

$$\dot{T}_{ab\dots c} = u^d \nabla_d T_{ab\dots c}, \quad (1.7)$$

where ∇_a denotes covariant differentiation.

Following Maartens (1997), we use angle brackets to denote the orthogonal projections of vectors, the orthogonally projected symmetric trace-free part of tensors, and their time derivatives, *i.e.*

$$X^{(a)} = h^a_b X^b, \quad T^{(ab)} = \left[h^a_c h^b_d - \frac{1}{3} h^{ab} h_{cd} \right] T^{cd}, \quad (1.8)$$

$$\dot{X}^{(a)} = h^a_b \dot{X}^b, \quad \dot{T}^{(ab)} = \left[h^a_c h^b_d - \frac{1}{3} h^{ab} h_{cd} \right] \dot{T}^{cd}. \quad (1.9)$$

Further, we will use the projection symbol \perp to identify orthogonal projection over all tensor indices, *i.e.*

$$(T^{ab}_{cd})_{\perp} = h^a_p h^b_q h^r_c h^s_d T^{pq}_{rs}. \quad (1.10)$$

The *spatial covariant derivative* $\hat{\nabla}_a$ is defined by projecting all free indices of the covariant derivative (compare Ellis (1973), Ellis and Bruni (1989)). For example,

$$\hat{\nabla}_a f = h^b_a \nabla_b f, \quad \hat{\nabla}_a T^{bc}_{de} = h^p_a h^b_q h^r_c h^s_d h^t_e \nabla_p T^{qr}_{st}. \quad (1.11)$$

The covariant derivative of a scalar can then be decomposed into a derivative along the fundamental congruence \mathbf{u} and a spatial covariant derivative via

$$\nabla_a f = \hat{\nabla}_a f - u_a \dot{f}. \quad (1.12)$$

The totally skew pseudotensor is defined by

$$\eta^{abcd} = \eta^{[abcd]}, \quad \eta^{0123} = (-g)^{-1/2}, \quad (1.13)$$

where

$$g = \det(g_{ab}). \quad (1.14)$$

The *projected permutation tensor* is then defined by (compare Maartens et al. (1995)),

$$\epsilon_{abc} = \eta_{abcd} u^d. \quad (1.15)$$

and is used to define the *spatial curl of a vector* by

$$\text{curl } X_a = \epsilon_{abc} \hat{\nabla}^b X^c. \quad (1.16)$$

and the *spatial curl of a rank 2 tensor* by

$$\text{curl } T_{ab} = \epsilon_{cd(a} \hat{\nabla}^c T_{b)}^d. \quad (1.17)$$

1.2.2 Kinematic Quantities

We now focus on the decomposition of the covariant derivative of the fundamental 4-velocity. Following Ellis (1973, p8-11), we decompose the covariant derivative of a unit timelike vector field \mathbf{u} as

$$\nabla_b u_a = \sigma_{ab} + \omega_{ab} + H h_{ab} - \dot{u}_a u_b, \quad (1.18)$$

with σ_{ab} symmetric and trace-free, ω_{ab} antisymmetric, and

$$\sigma_{ab} u^a = 0, \quad \omega_{ab} u^a = 0, \quad \dot{u}_a u^a = 0. \quad (1.19)$$

It follows from (1.18) and (1.19) that

$$H = \frac{1}{3} \nabla_a u^a, \quad (1.20)$$

$$\dot{u}_a = u^b \nabla_b u_a, \quad (1.21)$$

$$\sigma_{ab} = \nabla_{(a} u_{b)} - H h_{ab} + \dot{u}_{(a} u_{b)}, \quad (1.22)$$

$$\omega_{ab} = \nabla_{[b} u_{a]} + \dot{u}_{[a} u_{b]}. \quad (1.23)$$

Here, H is the Hubble scalar⁵, \dot{u}_a is the acceleration vector, σ_{ab} is the rate of shear tensor and ω_{ab} is the vorticity tensor. In practice we will usually replace ω_{ab} by the vorticity vector, defined by

$$\omega^a = \frac{1}{2}\eta^{abcd}u_b\omega_{cd} \quad (1.24)$$

which satisfies $u^a\omega_a = 0$. We can write the vorticity tensor in terms of the vorticity vector using the relation

$$\omega^{ab} = \eta^{abcd}\omega_c u_d. \quad (1.25)$$

The magnitude of the rate of shear tensor and the vorticity tensor are defined by

$$\sigma^2 = \frac{1}{2}\sigma_{ab}\sigma^{ab}, \quad \omega^2 = \omega_a\omega^a.$$

We now state two useful propositions concerning an *irrotational* vector field u^a , i.e. $\omega_a = 0$.

Proposition 1.2.1 *Given a vector field \mathbf{u} , there exist scalar fields $f(x^a)$ and $t(x^a)$ such that⁶ $u_a = -ft_{,a}$ if and only if $\omega_a = 0$.*

Proof: Given $u_a = -ft_{,a}$, it follows by differentiating that $u_{[a,b}u_{c]} = 0$ and that $\omega_a = 0$, by (1.24). The converse is a special case of Frobenius' Theorem, which is usually stated in terms of differential forms (see for example Flanders (1989), see the theorem on page 94). \square

Comment: The significance of this result is that if $\omega = 0$, then the 3-spaces orthogonal to u^a at each point form spacelike hypersurfaces given by $t = \text{constant}$, i.e. there is a one-parameter family of spacelike hypersurfaces orthogonal to the timelike congruence defined by \mathbf{u} .

If we can further impose the condition of zero acceleration ($\dot{u}_a = 0$), we obtain the following result.

Proposition 1.2.2 $\omega_a = 0, \dot{u}_a = 0 \iff u_a = -t_{,a}$, for some scalar field $t(x^a)$.

⁵The rate of expansion scalar Θ , defined by $\Theta = 3H$, is often used in the expansion (1.18).

⁶The minus sign ensures that $\dot{t} > 0$, i.e. t increases into the future along the fundamental congruence.

Proof: Given $\omega_a = \dot{u}_a = 0$, equation (1.23) implies $u_{[a;b]} = u_{[a,b]} = 0$. It then follows from Proposition 1.2.1 that $f_{,[a}t,b]} = 0$. This in turn implies that $f = g(t)$ and hence that one can redefine t to set $f = 1$. \square

In this case, the scalar t that labels the hypersurfaces is clock time along the normal congruence.

1.2.3 Decomposition of the Energy-Momentum Tensor

The energy-momentum tensor T_{ab} is a symmetric tensor that specifies the total matter-energy content of a cosmological model. We use the standard decomposition of the energy-momentum tensor with respect to a timelike vector field \mathbf{u} (see Ellis (1973), p7):

$$T_{ab} = \mu u_a u_b + q_a u_b + q_b u_a + p h_{ab} + \pi_{ab}, \quad (1.26)$$

where

$$q_a u^a = 0, \quad \pi_{ab} u^b = 0, \quad \pi_a^a = 0, \quad \pi_{ab} = \pi_{ba}. \quad (1.27)$$

In this equation, μ is the *total energy density* measured by an observer moving with fluid 4-velocity u^a , q_a is the *energy flux relative to u^a* (representing heat conduction and diffusion), p is the *isotropic pressure* and π_{ab} is the *trace-free anisotropic pressure* (due to processes such as viscosity).

A *perfect fluid* with 4-velocity \mathbf{u} is defined by

$$q_a = 0, \quad \pi_{ab} = 0, \quad (1.28)$$

in which case the energy-momentum tensor (1.26) simplifies to

$$T_{ab} = \mu u_a u_b + p(g_{ab} + u_a u_b). \quad (1.29)$$

1.2.4 The Weyl Curvature Tensor

The *Weyl conformal curvature tensor* (or simply the *Weyl tensor*) is defined in terms of the Riemann tensor and the Ricci tensor according to

$$C_{abcd} = R_{abcd} - \frac{1}{2}(g_{ac}R_{bd} + g_{bd}R_{ac} - g_{bc}R_{ad} - g_{ad}R_{bc}) + \frac{1}{6}R(g_{ac}g_{bd} - g_{ad}g_{bc}). \quad (1.30)$$

(see, for example, Ellis 1973, p7). We note that the Weyl curvature tensor has all the symmetry properties of the Riemann tensor and is in addition trace-free ($C^a{}_{bad} = 0$).

It is helpful to view (1.30) as a decomposition of the Riemann tensor R_{abcd} , with its 20 independent components, into a tracefree part C_{abcd} and the trace R_{ab} :

$$R_{abcd} \longleftrightarrow \{C_{abcd}, R_{ab}\}, \quad (1.31)$$

each of which have 10 independent components. The Ricci tensor is determined at each point by the energy-momentum tensor through the field equations (1.2). On the other hand, as we shall see later, the Weyl tensor governs the propagation of changes in the gravitational field, *i.e.* gravitational waves.

Within the 1 + 3 formalism the Weyl tensor can be decomposed further into an “electric part” E_{ab} and a “magnetic part” H_{ab} by projecting C_{abcd} and its dual orthogonal to \mathbf{u} :

$$E_{ac} = C_{abcd}u^b u^d, \quad H_{ac} = \frac{1}{2}\eta_{ab}{}^{ef} C_{efcd}u^b u^d. \quad (1.32)$$

We note that E_{ab} and H_{ab} are symmetric, trace-free and orthogonal to \mathbf{u} :

$$E_{ab}u^b = 0, \quad H_{ab}u^b = 0. \quad (1.33)$$

In addition, this decomposition describes the Weyl tensor completely since⁷

$$E_{ab} = 0 = H_{ab} \iff C_{abcd} = 0. \quad (1.34)$$

Note that through (1.30) E_{ab} , H_{ab} and R_{ab} provide a full description of the space-time curvature as described by the Riemann tensor.

1.2.5 Spatial Curvature

If \mathbf{u} is irrotational, we have seen that there exists a family of hypersurfaces (*i.e.* 3-dimensional manifolds) orthogonal to \mathbf{u} . In this situation the projection tensor h_{ab} can be viewed as the 3-metric induced on these hypersurfaces and the spatial covariant derivative $\hat{\nabla}_a$ can be viewed as the covariant derivative determined by this 3-metric.

Analogous to the spacetime Ricci identities, the 3-space Ricci identities are given by

$$\hat{\nabla}_c \hat{\nabla}_d X_a - \hat{\nabla}_d \hat{\nabla}_c X_a = {}^{(3)}R_{abcd} X^b, \quad (1.35)$$

⁷The Weyl tensor can, in fact, be written in terms of E_{ab} and H_{ab} (see, for example, Ellis and van Elst (1998) eq. (22)).

where ${}^{(3)}R_{abcd}$ is the *spatial Riemann tensor* and X_a is any vector field in the hypersurfaces, *i.e.* $X_a u^a = 0$. It can be shown by applying (1.35) to an arbitrary vector field \mathbf{X} that is orthogonal to \mathbf{u} , expanding the left hand side using (1.11) and (1.5) and introducing the spacetime Riemann tensor via

$$\nabla_c \nabla_d X_a - \nabla_d \nabla_c X_a = R_{abcd} X^b, \quad (1.36)$$

that ${}^{(3)}R_{abcd}$ is related to R_{abcd} via

$${}^{(3)}R_{abcd} = (R_{abcd})_{\perp} - \Theta_{ac} \Theta_{bd} + \Theta_{ad} \Theta_{bc}, \quad (1.37)$$

where Θ_{ab} is the expansion tensor of the congruence \mathbf{u} , defined by

$$\Theta_{ab} = \sigma_{ab} + H h_{ab}. \quad (1.38)$$

Equation (1.37) is the well-known *Gauss equation* (see, for example, Ellis (1973), p33-4).

The *spatial Ricci tensor* and *spatial Ricci scalar* are then defined in the usual manner:

$${}^{(3)}R_{ab} = h^{cd} {}^{(3)}R_{acbd}, \quad {}^{(3)}R = h^{ab} {}^{(3)}R_{ab}. \quad (1.39)$$

Further, the *trace-free spatial Ricci tensor* is defined as

$${}^{(3)}S_{ab} = {}^{(3)}R_{ab} - \frac{1}{3} {}^{(3)}R h_{ab}. \quad (1.40)$$

We can obtain an expression for the spatial Ricci scalar and trace-free spatial Ricci tensor in terms of the spacetime quantities from (B.5), incorporating the Weyl tensor through (1.30) and (1.32):

$${}^{(3)}R = R + 2R_{bd} u^b u^d - 6H^2 + 2\sigma^2, \quad (1.41)$$

$${}^{(3)}S_{ab} = E_{ab} + \frac{1}{2} R_{\langle ab \rangle} - H \sigma_{ab} + \sigma^c_{\langle a} \sigma_{b \rangle c}. \quad (1.42)$$

Then using the field equations (1.2) in the form (A.2)-(A.5) to eliminate R and R_{ab} , we obtain

$${}^{(3)}R = -6H^2 + 2\mu + 2\sigma^2, \quad (1.43)$$

$${}^{(3)}S_{ab} = E_{ab} + \frac{1}{2} \pi_{ab} - H \sigma_{ab} + \sigma^c_{\langle a} \sigma_{b \rangle c}. \quad (1.44)$$

We conclude this section with a brief digression on spaces of constant curvature. A differentiable manifold \mathcal{M} of dimension n with metric g is said to be a *space of constant curvature* if and only if

$$R_{abcd} = C(g_{ac}g_{bd} - g_{ad}g_{bc}). \quad (1.45)$$

It follows by contraction that

$$R = n(n-1)C, \quad \text{and} \quad R_{ab} - \frac{1}{n}Rg_{ab} = 0, \quad (1.46)$$

i.e. the tracefree Ricci tensor $S_{ab} \equiv R_{ab} - \frac{1}{n}Rg_{ab}$ is zero. In addition it follows from the twice contracted Bianchi identities that if $n > 2$ then R is constant (see proposition 1.2.3). If $n = 3$, the Riemann tensor is completely determined by the Ricci tensor according to⁸

$$R_{abcd} = 4h_{[d[b}S_{a]c]} + \frac{1}{3}Rg_{a[c}g_{b]d}. \quad (1.47)$$

It follows that if $n = 3$ then $(\mathcal{M}, \mathbf{g})$ is a space of constant curvature if and only if $S_{ab} = 0$. Within the framework of the 1 + 3 covariant formalism this result can be stated as follows.

Proposition 1.2.3 *Let \mathbf{u} be an irrotational timelike vector field. The hypersurfaces orthogonal to \mathbf{u} are spaces of constant curvature if and only if*

$${}^{(3)}S_{ab} = 0. \quad (1.48)$$

Further, if this condition holds, then

$$\hat{\nabla}_a {}^{(3)}R = 0. \quad (1.49)$$

1.2.6 Evolution and Constraint Equations

In the 1 + 3 covariant formalism, the information contained in the EFEs is displayed in an indirect manner by using the *Ricci identities* applied to the fundamental 4-velocity \mathbf{u} ,

$$\nabla_c \nabla_d u_a - \nabla_d \nabla_c u_a = R_{abcd} u^b, \quad (1.50)$$

and the *Bianchi identities*,

$$\nabla_{[a} R_{bc]de} = 0. \quad (1.51)$$

The latter can be written in divergence form by contracting once and then twice and using (1.30) to introduce the Weyl tensor:⁹

$$\nabla_d C_{abc}{}^d = -\nabla_{[a} R^c{}_{b]} - \frac{1}{6} \delta^c_{[a} \nabla_{b]} R, \quad (1.52)$$

⁸See, for example, Ellis (1973) eq. (80).

⁹Also see Wainwright and Ellis (1997), p27.

and

$$\nabla_b(R_a^b - \frac{1}{2}R\delta_a^b) = 0. \quad (1.53)$$

The Riemann tensor in (1.50) is likewise expressed in terms of the Weyl tensor and Ricci tensor using (1.30). In all these identities, the Ricci tensor is expressed in terms of the energy-momentum tensor (1.26) using the EFEs in the form

$$R_{ab} = T_{ab} - \frac{1}{2}g_{ab}T^c_c, \quad (1.54)$$

which introduces the fluid 4-velocity \mathbf{u} into (1.52) and (1.53). Equation (1.18) is used to replace all occurrences of $\nabla_b u_a$ by the kinematic quantities and (1.32) is used to replace C_{abcd} by E_{ab} and H_{ab} . Finally, one projects all three identities parallel to and orthogonal to \mathbf{u} . The result is a set of *evolution equations* giving the derivatives of H , σ_{ab} , ω_a , $E_{ab} + \frac{1}{2}\pi_{ab}$, H_{ab} , μ and q_a along the congruence \mathbf{u} , and a set of *constraint equations* that contain only the spatial covariant derivative $\hat{\nabla}_a$. We refer the reader to Appendix A for the complete list of equations that arise from this process.

1.3 Overview of the Thesis

The main goal of this thesis is twofold. Firstly, we analyze the dynamics of so-called n -fluid FL cosmologies, that is, homogeneous and isotropic cosmologies whose matter-energy content consists of n non-interacting fluids. Secondly, we give a unified coordinate-independent discussion of perturbations of FL cosmologies containing a perfect fluid and a cosmological constant. Some parts of this thesis are new, while other parts provide a unified presentation of known results for these cosmologies. Our treatment throughout is completely coordinate-independent since it is based on the so-called 1 + 3 covariant description of general relativity and in this respect it differs from much of the current literature.

In Chapter 1, we introduce the 1 + 3 covariant description of general relativity and in Appendix A we give the full system of evolution equations and constraint equations. In this sense the thesis is self-contained. However, a reader who has no prior knowledge of the 1 + 3 covariant description will encounter an initially steep learning curve. We assume the reader has a basic knowledge of general relativity and some familiarity with dynamical systems theory (see, for example, Wainwright and Ellis (1997), Chapter 4).

Chapter 2 plays a supporting role. In this chapter we describe the geometrical, *i.e.* coordinate independent properties of the Robertson-Walker metric, and give two characterizations of the Friedmann-Lemaître models that highlight their very special nature.

In Chapter 3, we give a complete qualitative description of the dynamics of n -fluid FL cosmologies. A central role is played by a new formulation of the evolution equations for these models as a dynamical system on a compact n -dimensional state space. The most important example is the 3-fluid model in which the three fluids are radiation, pressure-free matter and a cosmological constant, since it provides a model of the physical universe. Our discussion of this model is one of the highlights of this chapter.

In Chapter 4, we give a unified derivation of the governing equations for scalar, vector and tensor perturbations of FL cosmologies containing a perfect fluid and a cosmological constant, using the so-called geometrical approach pioneered by Ellis et al (1989). We then analyze the behaviour of the perturbations in various regimes using both exact and approximate solutions. In particular we determine how the perturbations affect the so-called observational quantities that determine the extent to which a perturbed model deviates from an exact FL model.

The Appendices contain a wealth of additional material relating to perturbations of FL universes, not all of which is needed for the purposes of this thesis.

THE FRIEDMANN-LEMAÎTRE COSMOLOGIES

In this chapter we first derive the geometrical (*i.e.* coordinate independent) properties of the Robertson-Walker metric within the framework of the 1 + 3 description of general relativity. We then give two characterizations of the Friedmann-Lemaître cosmologies, firstly using the kinematic quantities and secondly using the Weyl tensor.

2.1 Properties of the RW Metric

In this section we show that the high symmetry of the RW line-element leads to strong restrictions on the kinematic quantities of the fundamental congruence, on the Weyl tensor and on the Ricci tensor. We then derive the RW line-element assuming that the Cosmological Principle holds.

Proposition 2.1.1 *The kinematic quantities of the fundamental congruence (1.4) associated with the Robertson-Walker metric (1.3) satisfy*

$$\sigma_{ab} = 0, \quad \omega_a = 0, \quad \dot{u}_a = 0, \quad \hat{\nabla}_a H = 0. \quad (2.1)$$

Proof. For the RW line-element (1.3), the covariant derivative

$$\nabla_b u_a = u_{a,b} - \Gamma_{ab}^c u_c \quad (2.2)$$

simplifies to

$$\nabla_b u_a = \frac{1}{2} g_{ab,0}. \quad (2.3)$$

Furthermore, it follows from the RW metric (1.3) that

$$g_{\alpha\beta,0} = 2 \frac{\dot{\ell}}{\ell} g_{\alpha\beta}, \quad \text{and} \quad g_{a0,0} = 0. \quad (2.4)$$

Since the projection tensor h_{ab} has components $h_{\alpha\beta} = g_{\alpha\beta}$, $h_{a0} = 0$, it follows that

$$\nabla_b u_a = \frac{\dot{\ell}}{\ell} h_{ab}. \quad (2.5)$$

The decomposition (1.18) immediately implies $\sigma_{ab} = 0$, $\omega_{ab} = 0$, $\dot{u}_a = 0$ and

$$H = \frac{\dot{\ell}}{\ell}. \quad (2.6)$$

Thus, H is purely a function of t , which implies $\hat{\nabla}_a H = 0$. \square

We now consider the restrictions on the spatial curvature associated with the Robertson-Walker metric.

Proposition 2.1.2 *The spatial curvature of the RW metric satisfies*

$${}^{(3)}S_{ab} = 0, \quad \hat{\nabla}_a {}^{(3)}R = 0. \quad (2.7)$$

Proof: Since the hypersurfaces orthogonal to \mathbf{u} are spaces of constant curvature, the result follows immediately from proposition 1.2.3. \square

We now show that the Weyl tensor of the Robertson-Walker metric is zero.

Proposition 2.1.3 *The Robertson-Walker metric (1.3) has zero Weyl tensor:*

$$E_{ab} = 0, \quad H_{ab} = 0 \quad (2.8)$$

Proof: We make use of the contracted Ricci identities in the form¹

$$\dot{\sigma}_{\langle ab \rangle} = -2H\sigma_{ab} + \nabla_{\langle a}\dot{u}_{b \rangle} + \dot{u}_{\langle a}\dot{u}_{b \rangle} - \sigma_{\langle a}{}^c\sigma_{b \rangle c} - \omega_{\langle a}\omega_{b \rangle} - (E_{ab} - \frac{1}{2}R_{\langle ab \rangle}), \quad (2.9)$$

$$0 = H_{ab} - 2\dot{u}_{\langle a}\omega_{b \rangle} - \hat{\nabla}_{\langle a}\omega_{b \rangle} - \text{curl}(\sigma_{ab}). \quad (2.10)$$

We note that $H_{ab} = 0$ is an immediate consequence of (2.10), on applying proposition 2.1.1. If we apply proposition 2.1.1 to (2.9) and 2.1.2 to (1.42), we obtain

$$E_{ab} - \frac{1}{2}R_{\langle ab \rangle} = 0, \quad E_{ab} + \frac{1}{2}R_{\langle ab \rangle} = 0, \quad (2.11)$$

which implies $E_{ab} = 0$. \square

Finally, we derive the restrictions satisfied by the Ricci tensor of the Robertson-Walker metric (1.3).

Proposition 2.1.4 *The Ricci tensor of the Robertson-Walker metric (1.3) satisfies*

$$R_{ab}u^a h^b{}_c = 0, \quad R_{\langle ab \rangle} = 0. \quad (2.12)$$

Proof: We note that $R_{\langle ab \rangle} = 0$ follows immediately from (2.11). In order to show $R_{ab}u^a h^b{}_c = 0$, we make use of the contracted Ricci identities in the form²

$$0 = \hat{\nabla}^b\sigma_{ab} - 2\hat{\nabla}_a H - \text{curl}(\omega_a) - 2\epsilon_{abc}\dot{u}^b\omega^c + R_{bc}u^b h^c{}_a, \quad (2.13)$$

Then on applying proposition 2.1.1 to (2.13), we obtain $R_{ab}u^a h^b{}_c = 0$. \square

We conclude this section by showing that the Cosmological Principle, as stated in section 1.1, leads to the RW metric.

Proposition 2.1.5 *If the fundamental congruence of a cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ satisfies*

$$\sigma_{ab} = 0, \quad \omega_a = 0, \quad \dot{u}_a = 0, \quad \hat{\nabla}_a H = 0, \quad (2.14)$$

¹These equations are an intermediate step in deriving (A.9) and (A.13), and follow from the Ricci identities, without use of the field equations. In fact, on applying the field equations (1.2) in the form (A.2)-(A.5) to (2.9) and (2.10), we obtain (A.9) and (A.13).

²This equation is an intermediate step in deriving (A.11), and follow from the Ricci identities, without use of the field equations. In fact, on applying the field equations (1.2) in the form (A.2)-(A.5) to (2.13), we obtain (A.11).

and the hypersurfaces orthogonal to \mathbf{u} satisfy

$${}^{(3)}S_{ab} = 0, \quad (2.15)$$

then \mathbf{g} and \mathbf{u} can be written in the RW form (1.3) and (1.4).

Proof: By proposition 1.2.2, the fundamental 4-velocity \mathbf{u} can be written as

$$u_a = -t_{,a}, \quad (2.16)$$

where t is a scalar field. We then introduce *co-moving coordinates* x^α , i.e. coordinates x^α that are constant along the world lines of the fundamental congruence. Then the metric and 4-velocity assume the form

$$ds^2 = -dt^2 + g_{\alpha\beta}(t, x^\lambda)dx^\alpha dx^\beta, \quad \mathbf{u} = \frac{\partial}{\partial t}. \quad (2.17)$$

Next, using the metric (2.17), it can be quickly shown that

$$\nabla_b u_a = \frac{1}{2}g_{ab,0}. \quad (2.18)$$

This equation, in conjunction with (1.5), (1.22), (2.14) and (2.16), leads to

$$g_{\alpha\beta,0} = 2H(t)g_{\alpha\beta}. \quad (2.19)$$

We introduce the length scale function $\ell(t)$ via

$$H(t) = \frac{\dot{\ell}}{\ell}, \quad (2.20)$$

and thus can integrate (2.19) to obtain

$$g_{\alpha\beta}(t, x^\lambda) = \ell(t)^2 \tilde{g}_{\alpha\beta}(x^\lambda). \quad (2.21)$$

Since we have that ${}^{(3)}S_{ab} = 0$, it follows from proposition 1.2.3 that $\tilde{g}_{\alpha\beta}$ describes a 3-space of constant curvature. Thus the metric (2.17) assumes the Robertson-Walker form (1.3). \square

Theorem 2.1.1 (Derivation of the RW Metric) *If a cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ satisfies the Cosmological Principle relative to \mathbf{u} , then \mathbf{g} and \mathbf{u} can be written in the RW form (1.3) and (1.4).*

Proof: The Cosmological Principle implies that any geometrically defined vector orthogonal to \mathbf{u} must be zero, since otherwise it would define a preferred direction orthogonal to \mathbf{u} . In particular, we have $\omega_a = \dot{u}_a = \hat{\nabla}_a H = 0$. Likewise, any geometrically defined rank two symmetric tracefree tensor X_{ab} that is orthogonal to \mathbf{u} ($X_{ab}u^b = 0$) must be zero, since otherwise its eigenvectors would define a preferred direction orthogonal to \mathbf{u} . In particular we have $\sigma_{ab} = {}^{(3)}S_{ab} = 0$. The desired result thus follows from proposition 2.1.5. \square

2.2 Characterizations of the FL Cosmologies

In this section we present two characterizations of FL cosmologies in terms of the kinematic quantities and Weyl curvature.

2.2.1 The Kinematic Characterization

We now present the *kinematic characterization of the Friedmann-Lemaître models*.³

Theorem 2.2.1 *A cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ which satisfies the Einstein field equations with perfect fluid source is a Friedmann-Lemaître cosmology if and only if the kinematic quantities of the fluid 4-velocity \mathbf{u} satisfy*

$$\sigma_{ab} = 0, \quad \omega_a = 0, \quad \dot{u}_a = 0. \quad (2.22)$$

Proof.⁴ If we assume the model is a FL cosmology, the kinematic restrictions (2.22) follow immediately from proposition 2.1.1.

Conversely, assume that the restrictions (2.22) hold. Since the source is a perfect fluid with 4-velocity \mathbf{u} , equation (1.28) implies $\pi_{ab} = 0$ and $q_a = 0$. Then the conditions (2.22) in conjunction with (A.9) imply $E_{ab} = 0$, which in turn implies ${}^{(3)}S_{ab} = 0$ via (1.44). The conditions (2.22) and $q_a = 0$ in conjunction with (A.11) imply $\hat{\nabla}_a H = 0$. The desired result then follows from proposition 2.1.5. \square

³One can replace the assumption $\dot{u}_a = 0$ in Theorem 2.2.1 by the assumption that the perfect fluid has a linear equation of state $p = (\gamma - 1)\mu$. However, the proof requires a more detailed analysis and is beyond the scope of this thesis (see instead Collins and Wainwright (1983)).

⁴Krasinski (1997) gives the outline of a proof, referring to other sources for some parts (see page 11).

2.2.2 The Weyl Characterization

We now present a characterization of the FL models using the Weyl tensor (see section 1.2.4).

Proposition 2.2.1 *If a cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ satisfies the Einstein field equations with perfect fluid source having 4-velocity \mathbf{u} and barotropic equation of state $p = p(\mu)$ with $\mu + p \neq 0$, and has zero Weyl tensor,*

$$E_{ab} = 0, \quad H_{ab} = 0, \quad (2.23)$$

then the kinematic quantities satisfy

$$\sigma_{ab} = 0, \quad \omega_{ab} = 0, \quad \dot{u}_a = 0. \quad (2.24)$$

Proof. On account of (2.23), equations (A.18), (A.20) and (A.21) simplify to

$$(\mu + p)\sigma_{ab} = 0, \quad \hat{\nabla}_a \mu = 0, \quad (\mu + p)\omega_a = 0. \quad (2.25)$$

Since $(\mu + p) \neq 0$, (2.25) implies $\sigma_{ab} = 0$ and $\omega_a = 0$. We now make use of the chain rule to write

$$\hat{\nabla}_a p = \frac{dp}{d\mu} \hat{\nabla}_a \mu = 0. \quad (2.26)$$

Then using the perfect fluid condition, *i.e.* $\pi_{ab} = 0$ and $q_a = 0$, it follows from (A.15) and (2.26) that $(\mu + p)\dot{u}_a = 0$, and hence $\dot{u}_a = 0$, which completes the proof. \square

We can now present the main result of this section, which we shall refer to as the *Weyl characterization of the Friedmann-Lemaître models*.

Theorem 2.2.2 *A cosmological model $(\mathcal{M}, \mathbf{g}, \mathbf{u})$, which satisfies the Einstein field equations with perfect fluid source having 4-velocity \mathbf{u} and barotropic equation of state $p = p(\mu)$ with $\mu + p \neq 0$, is a Friedmann-Lemaître model if and only if the Weyl tensor is zero:*

$$E_{ab} = 0, \quad H_{ab} = 0.$$

Proof. This result follows immediately from proposition 2.1.3 and theorem 2.2.1 in conjunction with proposition 2.2.1. \square

We now present one final result which restricts the gradients of the physical and geometrical scalars in an FL cosmology.

Proposition 2.2.2 *A Friedmann-Lemaître cosmology satisfies*

$$\hat{\nabla}_a \mu = \hat{\nabla}_a p = \hat{\nabla}_a H = \hat{\nabla}_a {}^{(3)}R = 0. \quad (2.27)$$

Proof: We note that $\hat{\nabla}_a H = \hat{\nabla}_a {}^{(3)}R = 0$ follows immediately from propositions 2.1.1 and 2.1.2. Further, it follows from (A.4), (A.5) and proposition 2.1.4 that $q_a = 0$ and $\pi_{ab} = 0$. Hence, (A.15) and (A.20) along with theorems 2.2.1 and 2.2.2 implies $\hat{\nabla}_a \mu = \hat{\nabla}_a p = 0$. \square

DYNAMICS OF n -FLUID FL COSMOLOGIES

The main goal of this chapter is a description of the dynamics of n -fluid FL cosmologies. In section 3.1, using the results from chapter 1, we derive the basic evolution equations for n -fluid FL cosmologies and describe their behaviour using dynamical systems techniques. In section 3.2, we describe the degrees of freedom of the n -fluid FL cosmologies in two ways, firstly, using intrinsic parameters and secondly, using observational parameters. In sections 3.3 and 3.4, we apply the general analysis to the case of 2-fluid and 3-fluid FL cosmologies. Finally, in section 3.5 we give a unified presentation of the known solutions of the EFEs for single fluid and 2-fluid FL cosmologies, and relate them to the state space analysis in section 3.3.

3.1 Behaviour and Classification of n -Fluid FL Cosmologies

In this section we derive the evolution equations for n -fluid FL cosmologies without introducing local coordinates and a line-element, by specializing the general evolution and constraint equations (appendix A) using the restrictions derived in chapter 2. We formulate the evolution equations as ODEs for a set of dimensionless variables in a compact state, which enables us to use dynamical systems methods to describe the evolution of n -fluid FL cosmologies qualitatively.

3.1.1 Basic Evolution Equations

We know from theorems 2.2.1 and 2.2.2 and proposition 2.1.4 that FL models have the following properties:

- i) the kinematic quantities of the fundamental congruence \mathbf{u} satisfy

$$\sigma_{ab} = 0, \quad \omega_a = 0, \quad \dot{u}_a = 0, \quad (3.1)$$

- ii) the Weyl curvature is zero:

$$E_{ab} = 0, \quad H_{ab} = 0, \quad (3.2)$$

and

- iii) the stress-energy tensor satisfies

$$\pi_{ab} = 0, \quad q_a = 0, \quad (3.3)$$

i.e. it has the form of a perfect fluid.

The remaining non-zero physical and geometric variables that appear in the general evolution and constraint equations in Appendix A (A.8)-(A.21) are the Hubble scalar H , the energy density μ and the pressure p of the matter content. Propositions 2.1.1 and 2.1.4 show that these scalars have zero spatial gradient

$$h_a^b H_{;b} = 0, \quad h_a^b \mu_{;b} = 0, \quad h_a^b p_{;b} = 0, \quad (3.4)$$

and hence are purely functions of clock time t along the fundamental congruence. With the above restrictions, the general evolution and constraint equations reduce to two evolution equations for μ and H , namely (A.8) and (A.14), which read

$$\dot{\mu} = -3H(\mu + p), \quad (3.5)$$

$$\dot{H} = -H^2 - \frac{1}{6}(\mu + 3p), \quad (3.6)$$

where the overdot denotes differentiation with respect to t . The spatial geometry of the model is determined by the Gauss equation in the form (1.43) and (1.44). It follows that the tracefree Ricci tensor is zero,

$${}^{(3)}S_{ab} = 0, \quad (3.7)$$

and that the spatial Ricci scalar is determined by H and μ according to

$${}^{(3)}R = -6H^2 + 2\mu. \quad (3.8)$$

In order to obtain the standard evolution equations, we introduce the length scale function $\ell(t)$, which is defined by H , up to a constant multiple, according to

$$H = \frac{\dot{\ell}}{\ell}. \quad (3.9)$$

This definition is consistent with the length scale defined by the RW metric (1.3).

To proceed we need the following result:

Proposition 3.1.1 *In a Friedmann-Lemaître model the quantity*

$${}^{(3)}R(t)\ell(t)^2 \quad (3.10)$$

is constant in time.

Proof. A straightforward calculation using (3.5)-(3.9) yields $({}^{(3)}R\ell^2)^\bullet = 0$. \square

Since ℓ is only defined by (3.9) up to a multiplicative constant, we may choose it to satisfy¹

$$\frac{1}{6} {}^{(3)}R\ell^2 = K = \pm 1, \text{ or } 0, \quad (3.11)$$

where K is the *spatial curvature parameter*. The value for K indicates whether ${}^{(3)}R$ is positive, negative or zero. Models where $K = -1, 0$ and $+1$ are then referred to, respectively, as *open FL models*, *flat FL models* and *closed FL models*.

It is important to note that if $K = \pm 1$, then $\ell(t)$ is completely determined via eq (3.11), by the spatial geometry, represented by ${}^{(3)}R$. On the other hand, if $K = 0$, equation (3.11) does not restrict $\ell(t)$, which means that *if $K = 0$ then $\ell(t)$ can be rescaled with a multiplicative constant*.

We now present the standard form of the evolution equations for FL models (see, for example, Wainwright and Ellis 1997, p52). Using (3.9) and (3.11), equations (3.5), (3.6) and (3.8) are rewritten as the *conservation equation*,

$$\dot{\mu} = -3\frac{\dot{\ell}}{\ell}(\mu + p), \quad (3.12)$$

¹The factor of $\frac{1}{6}$ is chosen for compatibility with the RW metric.

the Raychaudhuri equation,

$$\frac{\ddot{\ell}}{\ell} = -\frac{1}{6}(\mu + 3p), \quad (3.13)$$

and the Friedmann equation,

$$3\dot{\ell}^2 = \mu\ell^2 - 3K. \quad (3.14)$$

Note that (3.13) is a consequence of (3.12) and (3.14) provided that $\dot{\ell} \neq 0$, as follows by differentiating (3.14).

3.1.2 n -Fluid FL Cosmologies

In this chapter, we consider FL cosmological models whose matter content consists of n fluids with the same four-velocity \mathbf{u} . The total energy density and pressure are equal to a sum of the energy density and pressure of the individual components:

$$\mu = \sum_{i=1}^n \mu_i, \quad p = \sum_{i=1}^n p_i. \quad (3.15)$$

The fluids are assumed to be non-interacting, which means that each fluid component individually satisfies the conservation equation (3.12), *i.e.*

$$\dot{\mu}_i = -3\frac{\dot{\ell}}{\ell}(\mu_i + p_i). \quad (3.16)$$

We further assume that each fluid obeys a linear equation of state with equation of state parameters γ_i ($i = 1, \dots, n$), *i.e.*

$$p_i = (\gamma_i - 1)\mu_i, \quad (3.17)$$

with

$$0 \leq \gamma_i \leq 2. \quad (3.18)$$

The most important physical values are $\gamma_i = 1$ for pressure-free matter and $\gamma_i = \frac{4}{3}$ for radiation. The value $\gamma_i = 0$ can be used to represent a cosmological constant and $\gamma_i = 2$ is sometimes considered to correspond to a so-called stiff fluid. For definiteness we order the equation of state parameters according to

$$\gamma_1 > \gamma_2 > \dots > \gamma_n, \quad (3.19)$$

and assume that $\gamma_1 \geq 1$, *i.e.* at least one matter component is matter with a non-negative pressure. The conservation equations (3.16) then assume the form

$$\dot{\mu}_i = -3\frac{\dot{\ell}}{\ell}\gamma_i\mu_i. \quad (3.20)$$

As well, the Friedmann equation (3.14) and Raychaudhuri equation (3.13) assume the form

$$\dot{\ell}^2 = \sum_{i=1}^n \frac{1}{3} \mu_i \ell^2 - K, \quad (3.21)$$

and

$$\frac{\ddot{\ell}}{\dot{\ell}} = -\frac{1}{6} \sum_{i=1}^n (3\gamma_i - 2) \mu_i. \quad (3.22)$$

In summary, we will use the term *n-fluid FL cosmology* to refer to an FL cosmological model whose matter content consists of n non-interacting fluids with the same 4-velocity \mathbf{u} and equations of state given by (3.17)-(3.19). Following Jantzen and Ugla (1992), we will use abbreviations such as *RDC-universe* and *RCA-universe* to identify n -fluid FL cosmologies, where R denotes radiation, D denotes dust, C denotes curvature and Λ denotes a cosmological constant.

We have shown that the state of an n -fluid FL cosmology as a function of time t is determined by the functions $(\ell, \mu_1, \dots, \mu_n)$ which satisfy the system of ODEs (3.20) and (3.21). The state of an n -fluid cosmology can equivalently be described by the variables (H, μ_1, \dots, μ_n) , which satisfy the evolution equations

$$\dot{\mu}_i = -3H\gamma_i\mu_i, \quad (3.23)$$

$$\dot{H} = -H^2 - \frac{1}{6} \sum_{i=1}^n (3\gamma_i - 2) \mu_i. \quad (3.24)$$

The first of these is a restatement of (3.20) using the definition (3.9) of H . The second follows from (3.6) on using (3.15) and (3.17). Further, upon applying (3.9), the Friedmann equation (3.21) gives an additional constraint that must be satisfied, namely

$$H^2 = \frac{1}{3} \sum_{i=1}^n \mu_i - \frac{K}{\ell^2}. \quad (3.25)$$

We note that this equation explicitly determines ℓ in terms of H and μ when $K \neq 0$.

At this stage, we present four simple and well-known solutions of equations (3.20)-(3.22) that play an important role when we formulate the evolution equations (3.23)-(3.24) for the n -fluid FL cosmologies as a dynamical system.

1. *Flat FL universe* ($K = 0, 0 < \gamma \leq 2$). For a single fluid, the length scale, energy density and pressure are

$$\ell = \ell_0 \left(\frac{t}{t_0} \right)^{\frac{2}{3\gamma}}, \quad \mu = \frac{4}{3\gamma^2 t^2}, \quad p = (\gamma - 1)\mu. \quad (3.26)$$

In the case of dust ($\gamma = 1$), this solution is the *Einstein-de Sitter universe* (Einstein and de Sitter (1932)).

2. *de Sitter universe* ($K = 0$). The de Sitter universe can be thought of as the special case of the flat FL universe when the single fluid is a cosmological constant (*i.e.* when $\gamma = 0$). The length scale, energy density and pressure are

$$\ell = \ell_0 \exp\left(\sqrt{\frac{\Lambda}{3}}t\right), \quad \mu = \Lambda, \quad p = -\Lambda. \quad (3.27)$$

3. *Milne universe* ($K = -1$). The length scale, energy density and pressure are

$$\ell = t, \quad \mu = 0, \quad p = 0. \quad (3.28)$$

The Milne universe can be thought of as the open FL universe with zero matter content.

4. *Einstein static universe* ($K = +1$). This solution arises when we require that the length scale is constant. The length scale, total energy density and total pressure are

$$\ell = \ell_0, \quad \mu = 3\ell_0^{-2}, \quad p = -\ell_0^{-2}, \quad (3.29)$$

where $\ell_0 > 0$ is constant.² The matter-energy content can be interpreted as two, or more generally, n non-interacting fluids. For example, for a 2-fluid model, the energy densities are given by³

$$\mu_1 = \left(\frac{2 - 3\gamma_2}{\gamma_1 - \gamma_2}\right)\ell_0^{-2}, \quad \mu_2 = \left(\frac{3\gamma_1 - 2}{\gamma_1 - \gamma_2}\right)\ell_0^{-2}, \quad (3.30)$$

where $0 \leq \gamma_2 < \frac{2}{3} < \gamma_1 \leq 2$. Einstein's choice was $\gamma_1 = 1$ (pressure-free matter) and $\gamma_2 = 0$ (a cosmological constant) so that $\mu_1 = 2\ell_0^{-2}$, $p_1 = 0$ and $\mu_2 = \ell_0^{-2} = \Lambda$, $p_2 = -\Lambda$, where Λ is the cosmological constant.

²The values of μ and p follow (3.13) and (3.14).

³These expressions are obtained by solving the linear system

$$\mu_1 + \mu_2 = 3\ell^2, \quad (3\gamma_1 - 2)\mu_1 + (3\gamma_2 - 2)\mu_2 = 0,$$

which follows from (3.21) and (3.22) with $\ell = \ell_0$ and $n = 2$.

3.1.3 Hubble-Normalized Scalars

In this section we introduce so-called *Hubble-normalized scalars*, which form a set of dimensionless variables that, together with the Hubble scalar H , describe the state of n -fluid FL cosmologies. These dimensionless scalars also play an important role in relating the models to observations (see section 3.2.2).

Firstly, the *density parameters* Ω_i , which describe the dynamical effect of the matter densities, are defined by

$$\Omega_i = \frac{\mu_i}{3H^2}. \quad (3.31)$$

The *total density parameter* Ω , which describes the dynamical effect of the total matter content, is defined by

$$\Omega = \frac{\mu}{3H^2}. \quad (3.32)$$

It follows from (3.15) and (3.31) that

$$\Omega = \sum_{i=1}^n \Omega_i. \quad (3.33)$$

Lastly, the dynamical effect of the spatial curvature is described by Ω_k , defined by

$$\Omega_k = -\frac{{}^{(3)}R}{6H^2}. \quad (3.34)$$

It follows from (3.11) that

$$\Omega_k = -\frac{K}{H^2 \ell^2}. \quad (3.35)$$

In terms of the parameters Ω and Ω_k , the Friedmann equation (3.21) assumes the simple form

$$\Omega + \Omega_k = 1, \quad (3.36)$$

which may be further expanded in terms of Ω_i as

$$\sum_{i=1}^n \Omega_i + \Omega_k = 1. \quad (3.37)$$

We note that (3.36) implies that Ω determines the spatial curvature as follows:

$$\Omega < 1 \iff \text{open FL}, \quad (3.38)$$

$$\Omega = 1 \iff \text{flat FL}, \quad (3.39)$$

$$\Omega > 1 \iff \text{closed FL}. \quad (3.40)$$

There is an additional dimensionless variable that plays a fundamental role, namely the *deceleration parameter* q , defined by

$$q = -\frac{\ddot{\ell}}{(\dot{\ell})^2}. \quad (3.41)$$

Clearly, if $q < 0$ then the expansion is accelerating ($\ddot{\ell} > 0$) and if $q > 0$ then the expansion is decelerating ($\ddot{\ell} < 0$). In an n -fluid FL model, q can be expressed in terms of the density parameters by using the Raychaudhuri equation (3.22) with (3.9) and (3.31). We write

$$q = \frac{1}{2} \sum_{i=1}^n (3\gamma_i - 2)\Omega_i. \quad (3.42)$$

The quantities Ω_i , Ω , Ω_k and q are dimensionless, whereas H has dimensions of $(length)^{-1}$. Together, these variables are fundamental quantities in the FL models and more general models (see, for example, Wainwright and Ellis (1997), section 5.2 and chapters 12 and 13).

The evolution equations for Ω_i and Ω_k can be obtained by differentiating (3.31) and (3.35) and applying (3.14), (3.23), (3.24) and (3.42). On performing this procedure, we obtain

$$\frac{d\Omega_i}{dt} = (2q - (3\gamma_i - 2))\Omega_i H, \quad \text{and} \quad \frac{d\Omega_k}{dt} = 2q\Omega_k H. \quad (3.43)$$

3.1.4 A Compact State Space for n -fluid FL Cosmologies

The one drawback of using H -normalized variables is that they cannot give a unified description of models that have both an expanding ($H > 0$) and a contracting ($H < 0$) epoch. In particular, when $H = 0$ the density parameters (3.31) and (3.34) are undefined (*i.e.* $\Omega_i \rightarrow +\infty$ as $H \rightarrow 0$). In order to circumvent this difficulty, we introduce a set of variables which are bounded throughout the evolution of the model.

For an n -fluid FL cosmology, with total energy density

$$\mu = \sum_{i=1}^n \mu_i > 0, \quad (3.44)$$

we define the dimensionless *matter variables*, which describe the relative significance of the different matter components:

$$\chi_i = \frac{\mu_i}{\mu}, \quad \mu_i \geq 0. \quad (3.45)$$

Then by (3.44) we have

$$\sum_{i=1}^n \chi_i = 1, \quad 0 \leq \chi_i \leq 1, \quad (3.46)$$

i.e. the χ_i are bounded.

Following Wainwright (1996, p124-5), we now introduce a *curvature parameter* $\tilde{\Omega}$ so as to compactify the state space:

$$\tilde{\Omega} = \arctan \left(\frac{\sqrt{3}H}{\sqrt{\mu}} \right). \quad (3.47)$$

It follows that $\tilde{\Omega}$ satisfies

$$-\frac{\pi}{2} < \tilde{\Omega} < \frac{\pi}{2}, \quad (3.48)$$

and that

$$\text{sign}(\tilde{\Omega}) = \text{sign}(H). \quad (3.49)$$

This variable thus describes the state of expansion of a model. In addition, it describes the spatial curvature, *i.e.* the deviation from flatness, as a consequence of (3.32) and (3.38)-(3.40). In particular, it follows from (3.39) and (3.32) that $\tilde{\Omega} = \frac{\pi}{4}$ gives the flat FL models.

We now derive evolution equations for $\tilde{\Omega}$ and the χ_i . It is convenient to introduce a quantity \tilde{q} that plays the role of the deceleration parameter q , but unlike q , is bounded when $H = 0$. Motivated by (3.42), we define

$$\tilde{q} = \frac{1}{2} \sum_{i=1}^n (3\gamma_i - 2)\chi_i. \quad (3.50)$$

Together, (3.19) and (3.50) imply that \tilde{q} is restricted by

$$\frac{1}{2}(3\gamma_n - 2) \leq \tilde{q} \leq \frac{1}{2}(3\gamma_1 - 2). \quad (3.51)$$

It follows from (3.15), (3.23), (3.45), (3.46) and (3.50) that the total energy density satisfies

$$\frac{d\mu}{dt} = -2(\tilde{q} + 1)H\mu. \quad (3.52)$$

The evolution equation for χ_i then follows from (3.23), (3.45), (3.50) and (3.52) and is given by

$$\frac{d\chi_i}{dt} = [2\tilde{q} - (3\gamma_i - 2)]H\chi_i. \quad (3.53)$$

Upon noting that (3.47) implies

$$H = \sqrt{\frac{\mu}{3}} \tan \tilde{\Omega}, \quad (3.54)$$

we can rewrite (3.53) as

$$\frac{d\chi_i}{dt} = \sqrt{\frac{\mu}{3}} \frac{1}{\cos \tilde{\Omega}} [2\tilde{q} - (3\gamma_i - 2)](\sin \tilde{\Omega})\chi_i. \quad (3.55)$$

The form of (3.55) suggests defining a new time variable $\tilde{\tau}$ by

$$\frac{dt}{d\tilde{\tau}} = \sqrt{\frac{3}{\mu}} \cos \tilde{\Omega}. \quad (3.56)$$

It is essential to note that this change of variable is well-defined, since $\cos \tilde{\Omega} > 0$ on account of (3.48). In terms of $\tilde{\tau}$, (3.55) reads

$$\frac{d\chi_i}{d\tilde{\tau}} = [2\tilde{q} - (3\gamma_i - 2)](\sin \tilde{\Omega})\chi_i. \quad (3.57)$$

In order to obtain the evolution equation for $\tilde{\Omega}$ we need the evolution equations for μ and H . First, it follows from (3.52) and (3.56) that

$$\frac{d\mu}{d\tilde{\tau}} = -2\sqrt{3\mu}H(\tilde{q} + 1) \cos \tilde{\Omega}. \quad (3.58)$$

Secondly, equations (3.24), (3.45), (3.50), (3.54) and (3.56) lead to

$$\frac{dH}{d\tilde{\tau}} = -\sqrt{\frac{\mu}{3}}(\tan^2 \tilde{\Omega} + \tilde{q}) \cos \tilde{\Omega}. \quad (3.59)$$

We differentiate (3.54) with respect to $\tilde{\tau}$ and use (3.58). After simplifying using (3.54), we obtain

$$\frac{d\tilde{\Omega}}{d\tilde{\tau}} = -\tilde{q} \cos 2\tilde{\Omega} \cos \tilde{\Omega}. \quad (3.60)$$

Equations (3.57) and (3.60) are the desired evolution equations. The state space is the bounded n -dimensional subset of \mathbb{R}^{n+1} described by $(\tilde{\Omega}, \chi_1, \dots, \chi_n)$ and subject to the restrictions (3.46) and (3.48). The state space is not compact due to the strict inequality in (3.48) and so we seek to compactify this space, since then we can apply theorems from the theory of dynamical systems.

In order to compactify the space, we simply observe that the right hand sides of the ODEs (3.57) and (3.60) are well-defined, in fact analytic, at the points $\tilde{\Omega} = \pm\frac{\pi}{2}$, which form part of the boundary of the state space. We can thus extend the state space to include the points with $\tilde{\Omega} = \pm\frac{\pi}{2}$, thereby compactifying it. Since

$$\lim_{|\tilde{\Omega}| \rightarrow \frac{\pi}{2}^-} \mu = 0, \quad (3.61)$$

as follows from (3.54), we interpret the sets $\tilde{\Omega} = \pm\frac{\pi}{2}$ as representing FL cosmologies with $\mu = 0$. It follows from (3.14) that the set $\tilde{\Omega} = \pm\frac{\pi}{2}$ represents the Milne universe, given by (3.28).

State Space Representation of n -Fluid FL Cosmologies**State Space:**

$$(\tilde{\Omega}, \chi_1, \dots, \chi_n) \in \mathbb{R}^{n+1}$$

$$\sum_{i=1}^n \chi_i = 1, \quad 0 \leq \chi_i \leq 1, \quad -\frac{\pi}{2} \leq \tilde{\Omega} \leq \frac{\pi}{2}. \quad (3.62)$$

Evolution Equations:

$$\frac{d\tilde{\Omega}}{d\tilde{\tau}} = -\tilde{q} \cos 2\tilde{\Omega} \cos \tilde{\Omega}, \quad (3.63)$$

$$\frac{d\chi_i}{d\tilde{\tau}} = [2\tilde{q} - (3\gamma_i - 2)] (\sin \tilde{\Omega}) \chi_i, \quad (3.64)$$

with

$$\tilde{q} = \frac{1}{2} \sum_{i=1}^n (3\gamma_i - 2) \chi_i. \quad (3.65)$$

Comments:

- i) The ODEs (3.63) and (3.64) describe the essential dynamics of n -fluid FL cosmologies in terms of the dimensionless variables $(\tilde{\Omega}, \chi_1, \dots, \chi_n)$ and the dimensionless time $\tilde{\tau}$. Since the state space, as defined by (3.62) is compact, *the solutions of the ODEs are defined for all $\tilde{\tau} \in \mathbb{R}$* , and as a result the ODEs define a dynamical system on the state space (see, for example, Wainwright and Ellis (1997), p87, Corollary 4.1). The evolution of an n -fluid FL cosmology is thus described by an orbit of the dynamical system (a solution curve of the ODEs). As a consequence, we are guaranteed the existence of a past attractor \mathcal{A}^- of the ODEs (*i.e.* an invariant set to which all orbits, except possibly a set of measure zero, are past asymptotic to as $\tilde{\tau} \rightarrow -\infty$) and a future attractor \mathcal{A}^+ (*i.e.* as a past attractor, except as $\tilde{\tau} \rightarrow +\infty$). The past attractor describes the asymptotic regime at early times and the future attractor describes the asymptotic regime at late times.
- ii) This description of the evolution of n -fluid FL cosmologies as a dynamical system on a compact state space is new. It generalizes the formulation given by Wainwright (1996) for the case $n = 2$.
- iii) The physical state of an n -fluid FL cosmology is determined by the variables (H, μ_1, \dots, μ_n) ,

which are related to the dimensionless variables via the total density μ , according to

$$H = \sqrt{\frac{\mu}{3}} \tan \tilde{\Omega}, \quad \mu_i = \mu \chi_i, \quad (3.66)$$

(see (3.54) and (3.45)). On using (3.66) to eliminate H , (3.58) assumes the form

$$\frac{d\mu}{d\tilde{\tau}} = -2(\tilde{q} + 1)(\sin \tilde{\Omega})\mu. \quad (3.67)$$

Then (3.67) determines μ as an integral in terms of $\tilde{\Omega}(\tilde{\tau})$ and $\chi_i(\tilde{\tau})$, up to an arbitrary constant. This freedom implies that *each orbit determines a 1-parameter family of cosmological models*, whose physical variables are related by a multiplicative constant. Since an n -dimensional dynamical system has an $(n - 1)$ -parameter family of orbits, it follows that *the class of n -fluid FL cosmologies is labelled by n parameters*. We will specify these parameters explicitly in section 3.2.

- iv) Clock time t is related to $\tilde{\tau}$ via the ODE (3.56) and can thus be expressed as an integral involving $\tilde{\Omega}(\tilde{\tau})$ and $\mu(\tilde{\tau})$, the latter determined from (3.67). Clock time will take on values in an interval

$$t_i < t < t_f, \quad (3.68)$$

where t_i is finite or $-\infty$ and t_f is finite or $+\infty$, depending on the model.

- v) The dimensionless variables $\tilde{\Omega}$, χ_i and \tilde{q} are closely related to the Hubble-normalized scalars Ω_i and q when $H \neq 0$. From (3.32) and (3.47) we have

$$\tan^2 \tilde{\Omega} = \frac{1}{\Omega}, \quad -\frac{\pi}{2} < \tilde{\Omega} < \frac{\pi}{2}, \quad \tilde{\Omega} \neq 0. \quad (3.69)$$

Similarly, from (3.31), (3.32) and (3.45) we have

$$\chi_i = \frac{\Omega_i}{\Omega}. \quad (3.70)$$

Finally, from (3.42), (3.50) and (3.70) we have

$$\tilde{q} = \frac{q}{\Omega}. \quad (3.71)$$

3.1.5 Invariant Sets and Equilibrium Points in the Compact State Space

Our goal in the rest of this section is to describe the qualitative behaviour of the orbits in the compact state space, thereby understanding the dynamics of n -fluid FL cosmologies. Since the state space is compact, it follows that every orbit has a past and future attractor (see, for example, Wainwright and Ellis (1997), p99-100).

In general the attractors of a dynamical system are complicated invariant sets. But in the present situation we will show that they are determined by the equilibrium points of the dynamical system (section 3.1.7), which we now present.

The form of the ODEs (3.63) and (3.64) determines three physically important invariant sets, given as follows:

- 1) *Flat FL set.* This set is the $n - 1$ dimensional subset given by $\tilde{\Omega} = \pm \frac{\pi}{4}$ (which corresponds to $\Omega = 1$, $\Omega_k = 0$ on account of (3.69) and (3.36)). The orbits in this set describe flat n -fluid FL cosmologies, expanding if $\tilde{\Omega} = \frac{\pi}{4}$ and contracting if $\tilde{\Omega} = -\frac{\pi}{4}$.
- 2) *Milne set.* This set is the $n - 1$ dimensional subset given by $\tilde{\Omega} = \pm \frac{\pi}{2}$ (which corresponds to $\Omega = 0$ and $\Omega_k = 1$ on account of (3.69) and (3.36)). The orbits in this set describe the Milne vacuum solution.
- 3) *m -fluid sets ($1 \leq m < n$).* These sets are determined by setting $n - m$ of the χ_i variables to be zero, with the m remaining χ_i variables being non-zero. The orbits in these sets describe m -fluid FL cosmologies.

The equilibrium points are determined by setting the right hand side of equations (3.63) and (3.64) to zero. If $\tilde{\Omega} \neq 0$, then (3.64) and (3.65) implies $\chi_i = 1$, for one value of i , $\chi_j = 0$, for $j \neq i$, and $\tilde{q} = \frac{1}{2}(3\gamma_i - 2) \neq 0$. Then (3.63) implies $\cos \tilde{\Omega} = 0$ (*i.e.* $\tilde{\Omega} = \pm \frac{\pi}{2}$) or $\cos 2\tilde{\Omega} = 0$ (*i.e.* $\tilde{\Omega} = \pm \frac{\pi}{4}$). If $\tilde{\Omega} = 0$ then (3.63) implies $\tilde{q} = 0$, with \tilde{q} given by (3.65). In this way, we obtain the following sets of equilibrium points:

- 1) *Flat FL points (F_i^\pm)* are given by

$$\tilde{\Omega} = \pm \pi/4, \quad \chi_i = 1, \quad \chi_j = 0 \text{ if } j \neq i. \quad (3.72)$$

These equilibrium points lie in the intersection of the flat FL set and the various 1-fluid sets.

2) *Milne points* (M_i^\pm) are given by

$$\tilde{\Omega} = \pm \frac{\pi}{2}, \quad \chi_i = 1, \chi_j = 0 \text{ if } j \neq i. \quad (3.73)$$

These equilibrium points lie in the intersection of the Milne set and the various 1-fluid sets.

3) *Einstein static points* (E) are given by

$$\tilde{\Omega} = 0, \quad \sum_{i=1}^n (3\gamma_i - 2)\chi_i = 0, \quad (3.74)$$

and so constitute an $(n - 2)$ -dimensional set of equilibrium points. These equilibrium points correspond to the n -fluid interpretation of the Einstein static model (3.29) and only exist when $\gamma_n < 2/3$.

3.1.6 Classification of n -Fluid FL Cosmologies

In this section, we will prove that the n -fluid FL cosmologies that have an epoch of expansion (*i.e.* $H > 0$ in some time interval) form three qualitatively different generic⁴ subclasses, defined as follows:

- 1) *Ever-expanding models* satisfy $H > 0$ for all t .
- 2) *Recollapsing models* satisfy $H(t_*) = 0$, $H > 0$ for $t < t_*$ and $H < 0$ for $t > t_*$ for some t_* .
- 3) *Bouncing models* satisfy $H(t_*) = 0$, $H < 0$ for $t < t_*$ and $H > 0$ for $t > t_*$ for some t_* .

The variables $\tilde{\Omega}$ and \tilde{q} are particularly important in this analysis. We begin with the following proposition on the behaviour of \tilde{q} .

Proposition 3.1.2 *In any n -fluid FL cosmology with $n > 1$, \tilde{q} is strictly decreasing (increasing) in an epoch of expansion (contraction). Further, if $n = 1$ then \tilde{q} is constant.*

⁴Invariant sets of dimension n .

Proof: By differentiating (3.50) with respect to $\tilde{\tau}$ and using (3.57), we obtain⁵

$$\frac{d\tilde{q}}{d\tilde{\tau}} = \frac{1}{2} \left[4\tilde{q}^2 - \sum_{i=1}^n (3\gamma_i - 2)^2 \chi_i \right] \sin \tilde{\Omega}. \quad (3.75)$$

By the Cauchy-Schwarz inequality and (3.46) we have

$$4\tilde{q}^2 = \left[\sum_{i=1}^n (3\gamma_i - 2) \chi_i \right]^2 \leq \left[\sum_{i=1}^n (3\gamma_i - 2)^2 \chi_i \right] \left[\sum_{i=1}^n \chi_i \right] = \sum_{i=1}^n (3\gamma_i - 2)^2 \chi_i, \quad (3.76)$$

with equality holding if and only if $n = 1$.⁶

On recalling (3.49), it follows that

$$\text{sign} \left(\frac{d\tilde{q}}{d\tilde{\tau}} \right) = -\text{sign}(H), \quad (3.77)$$

which establishes the result. \square

This proposition leads to a second result, presented here:

Proposition 3.1.3 *For any non-static n -fluid FL cosmology, $\tilde{\Omega}(\tilde{\tau}_*) = 0$ is satisfied for at most one $\tilde{\tau}_*$, and in this situation $\frac{d\tilde{\Omega}}{d\tilde{\tau}}(\tilde{\tau}_*) \neq 0$.*

Proof: We first show, by contradiction, that $\tilde{\Omega}(\tilde{\tau}_*) = 0$ is satisfied for at most one $\tilde{\tau}_*$. Suppose $\tilde{\Omega}(\tilde{\tau}_1) = 0 = \tilde{\Omega}(\tilde{\tau}_2)$ and $\tilde{\Omega} \neq 0$ for $\tilde{\tau} \in (\tilde{\tau}_1, \tilde{\tau}_2)$. Without loss of generality, we assume that $\tilde{\Omega} > 0$ for $\tilde{\tau} \in (\tilde{\tau}_1, \tilde{\tau}_2)$. Then by proposition 3.1.2 it follows that $\tilde{q}(\tilde{\tau}_2) \leq \tilde{q}(\tilde{\tau}_1)$ since \tilde{q} is strictly decreasing whenever $\tilde{\Omega} > 0$. Further, by (3.60), we have that $\tilde{\Omega}$ satisfies

$$\tilde{\Omega}'(\tilde{\tau}_1) = -\tilde{q}(\tilde{\tau}_1), \quad \text{and} \quad \tilde{\Omega}'(\tilde{\tau}_2) = -\tilde{q}(\tilde{\tau}_2), \quad (3.78)$$

⁵This equation is equivalent to (14.15) in Wainwright and Ellis (1997), since $\tilde{q} = \frac{1}{2}(3w + 1)$. The proposition is equivalent to $c_s^2 - w > 0$, if $n > 1$.

⁶Equality holds in (3.76) if and only if there exists c such that

$$(3\gamma_i - 2)\sqrt{\chi_i} = c\sqrt{\chi_i}, \quad i = 1, 2, \dots, n,$$

which is possible if and only if $n = 1$.

where the prime (\prime) denotes differentiation with respect to $\tilde{\tau}$. But $\tilde{\Omega}(\tilde{\tau}) > 0$ for $\tilde{\tau} \in (\tilde{\tau}_1, \tilde{\tau}_2)$ implies that $\tilde{\Omega}'(\tilde{\tau}_1) > 0$ (i.e. $\tilde{q}(\tilde{\tau}_1) < 0$) and $\tilde{\Omega}'(\tilde{\tau}_2) < 0$ (i.e. $\tilde{q}(\tilde{\tau}_2) > 0$) and so $\tilde{q}(\tilde{\tau}_2) > \tilde{q}(\tilde{\tau}_1)$, which is a contradiction.

Finally, it follows from (3.63) that if $\tilde{\Omega} = 0$, $\frac{d\tilde{\Omega}}{d\tilde{\tau}} = 0$ is satisfied only when $\tilde{q} = 0$, which is disallowed for non-static models. \square

The main result of this section now follows immediately.

Theorem 3.1.1 *There are exactly three generic classes of n -fluid FL cosmologies ($n > 1$) that have an expanding epoch, namely ever-expanding models, recollapsing models and bouncing models. Bouncing models occur if and only if $\gamma_n < \frac{2}{3}$.*

Proof: Since $H = 0$ if and only if $\tilde{\Omega} = 0$ on account of (3.47), proposition 3.1.3 states that the Hubble scalar can be zero at most once, and that H changes sign if it becomes zero. Thus models with more than one expanding epoch are excluded, which implies that the only possible classes are those given. If $\gamma_n > \frac{2}{3}$, it follows from (3.19) and (3.50) that $\tilde{q} > \frac{1}{2}(3\gamma_n - 2) > 0$, and hence from (3.63) that bouncing models are disallowed. \square

To the best of the author's knowledge, the classification theorem is new for $n > 2$, although it is a well known result for $n = 2$, when the matter content is a perfect fluid ($\gamma_1 > \frac{2}{3}$) and a cosmological constant ($\gamma_2 = 0$) (see Rindler (1977), p234-8).

3.1.7 Asymptotic Behaviour of Solutions

In this section we determine the asymptotic behaviour of the three generic classes of n -fluid FL cosmologies identified in the previous section. We will first describe the behaviour of the length scale $\ell(t)$ as a function of clock time t in each of the three generic classes of FL cosmologies, and then use these results to determine the past attractor and future attractor.

Important in this analysis is the set of orbits asymptotic to an Einstein static solution. In the following proposition, we demonstrate that orbits of this type are non-generic.

Proposition 3.1.4 *The set of orbits in the compact state space that are asymptotic to an Einstein static equilibrium point has dimension $n - 1$.*

Proof: We can write the evolution equations (3.57) and (3.60) as a dynamical system of the form

$$\frac{dX_i}{d\tilde{\tau}} = F_i(X_0, X_1, \dots, X_n), \quad i = 0, \dots, n, \quad (3.79)$$

on \mathbb{R}^{n+1} , where $X_0 = \tilde{\Omega}$ and $X_j = \chi_j$ ($j = 1, \dots, n$). The physical state space is the hyperplane given by (3.62), and hence is n dimensional. Consider the Jacobian matrix $J_{ij} = \frac{\partial F_j}{\partial X_i}$. Upon evaluating J_{ij} on the Einstein static set, given by (3.74), we obtain

$$J_{00} = 0, \quad J_{0i} = -\frac{1}{2}(3\gamma_i - 2), \quad J_{i0} = -(3\gamma_i - 2)\chi_{i,E}, \quad J_{ij} = 0, \quad (3.80)$$

where $i, j = 1, \dots, n$ and $\chi_{i,E}$ denotes the value of χ_i at the Einstein equilibrium point. The Jacobian matrix, when restricted to the physical state space, has two non-zero eigenvalues

$$\lambda_{\pm} = \pm \sqrt{\frac{1}{2} \sum_{i=1}^n (3\gamma_i - 2)^2 \chi_i} > 0, \quad (3.81)$$

and $n - 2$ zero eigenvalues.⁷ It follows⁸ that each Einstein static equilibrium point has a 1-dimensional stable manifold and a 1-dimensional unstable manifold. Thus the set of orbits future asymptotic to the Einstein static set has dimension exactly one larger than the dimension of the Einstein static manifold, *i.e.* dimension $n - 1$. \square

In the next proposition, we establish the behaviour of the length scale ℓ as a function of time $\tilde{\tau}$, for ever-expanding models.

Proposition 3.1.5 *If $\tilde{\Omega} > 0$ for all $\tilde{\tau} \in \mathbb{R}$ and $\tilde{\Omega} \not\rightarrow 0$ as $\tilde{\tau} \rightarrow \pm\infty$, then*

$$\lim_{\tilde{\tau} \rightarrow -\infty} \ell(\tilde{\tau}) = -\infty, \quad \text{and} \quad \lim_{\tilde{\tau} \rightarrow +\infty} \ell(\tilde{\tau}) = +\infty. \quad (3.82)$$

Proof: The length scale ℓ defines a time variable τ according to

$$\ell = \ell_0 e^{\tau}. \quad (3.83)$$

⁷The zero eigenvalues reflect the fact that the set of Einstein static equilibrium points has dimension $n - 2$.

⁸Using the Stable/Unstable Manifold Theorem. See, for example, Perko (1996, p107).

It follows from (3.9) and (3.83) that

$$\frac{dt}{d\tau} = \frac{1}{H}. \quad (3.84)$$

Then, using (3.54), (3.56) and (3.84), and the chain rule, we obtain

$$\frac{d\tau}{d\tilde{\tau}} = \sin \tilde{\Omega}. \quad (3.85)$$

Since $\tilde{\Omega} > 0$ for all $\tilde{\tau} \in \mathbb{R}$ and $\tilde{\Omega} \not\rightarrow 0$ as $\tilde{\tau} \rightarrow \pm\infty$, (3.85) then implies

$$\lim_{\tilde{\tau} \rightarrow -\infty} \tau(\tilde{\tau}) = -\infty, \quad \text{and} \quad \lim_{\tilde{\tau} \rightarrow +\infty} \tau(\tilde{\tau}) = +\infty. \quad (3.86)$$

This completes the proof, on applying (3.83). \square

Proposition 3.1.6 *For a recollapsing model, $\tilde{q}(\tilde{\tau}) > 0$ for all $\tilde{\tau}$, and for a bouncing model, $\tilde{q}(\tilde{\tau}) < 0$ for all $\tilde{\tau}$.*

Proof: For a recollapsing model, it follows from the definition that there exists $\tilde{\tau}_*$ so that $\tilde{\Omega}(\tilde{\tau}_*) = 0$ and that $\tilde{\Omega}(\tilde{\tau})$ changes sign from positive to negative at $\tilde{\tau}_*$, which implies $\frac{d\tilde{\Omega}}{d\tilde{\tau}}(\tilde{\tau}_*) < 0$ (strict inequality follows from Proposition 3.1.3). The evolution equation (3.63) now implies $q(\tilde{\tau}_*) > 0$. By Proposition 3.1.2, \tilde{q} has a global minimum at $\tilde{\tau}_*$, i.e. $\tilde{q}(\tilde{\tau}) \geq \tilde{q}(\tilde{\tau}_*)$ for all $\tilde{\tau}$, which gives the desired result. The proof for bouncing models is similar. \square

We now determine the asymptotic behaviour of clock time t (i.e. as $\tilde{\tau} \rightarrow \pm\infty$). Let

$$t_i = \lim_{\tilde{\tau} \rightarrow -\infty} t(\tilde{\tau}), \quad t_f = \lim_{\tilde{\tau} \rightarrow +\infty} t(\tilde{\tau}), \quad (3.87)$$

where the dependence of t on $\tilde{\tau}$ is determined via equation (3.56). Then t_i and t_f are finite or infinite depending on the type of universe and during the evolution $t_i < t < t_f$. Clock time is related to the length scale via the Friedmann equation. In particular, for any n -fluid FL cosmology, we can integrate (3.20) to obtain⁹

$$\mu_i \ell^2 = 3 (\lambda_i \ell)^{-3\gamma_i + 2}. \quad (3.88)$$

⁹See p46.

where λ_i is an integration constant whose inverse can be interpreted as the characteristic length scale of fluid i . On substituting (3.88) into the Friedmann equation (3.21), we obtain

$$\left(\frac{d\ell}{dt}\right)^2 = \sum_{i=1}^n (\lambda_i \ell)^{-3\gamma_i+2} - K. \quad (3.89)$$

Theorem 3.1.2 *The following statements are true:*

- i) *For an ever-expanding n -fluid FL model that is not asymptotic to the Einstein static universe, t_i is finite, $t_f = \infty$ and the length scale satisfies $\ell(t_i) = 0$ for some t_i and $\ell \rightarrow +\infty$ as $t \rightarrow +\infty$.*
- ii) *For a recollapsing n -fluid FL model, t_i and t_f are finite, $\ell(t_i) = 0$ and $\ell(t_f) = 0$.*
- iii) *For a bouncing n -fluid FL model, t_i and t_f are infinite and $\ell(t)$ satisfies $\ell(t) \rightarrow +\infty$ as $t \rightarrow \pm\infty$.*

Proof: Case i) In this case, we will make use of the following elementary results from calculus:

- (I) If $f(a) = O(a^r)$ as $a \rightarrow 0$, and $r > -1$, then $\lim_{a \rightarrow 0} \int_a^1 f(\tilde{a}) d\tilde{a} < \infty$, *i.e.* the integral converges.
- (II) If $f(a) \in O(a^r)$, as $a \rightarrow \infty$, and $r \geq -1$ then $\lim_{a \rightarrow \infty} \int_1^a f(\tilde{a}) d\tilde{a} = \infty$, *i.e.* the integral diverges.

Recall that ever-expanding models satisfy $H(\tilde{\tau}) > 0 \forall \tilde{\tau}$. Thus by Proposition 3.1.5, solutions which are not asymptotic to the Einstein static solution satisfy $\ell \rightarrow 0$ as $\tilde{\tau} \rightarrow -\infty$ and $\ell \rightarrow \infty$ as $\tilde{\tau} \rightarrow \infty$. Let

$$f(\ell) = \frac{1}{\sqrt{\sum_{i=1}^n (\lambda_i \ell)^{-3\gamma_i+2} - K}}. \quad (3.90)$$

For an ever-expanding model, this function is defined and continuous (for $0 < a < \infty$) since $H > 0$ implies that the denominator is strictly positive (see (3.89)). Hence, we can integrate (3.89) to obtain

$$t(\ell) = t(1) - \int_{\ell}^1 f(\tilde{\ell}) d\tilde{\ell}, \quad \text{for } 0 < \ell < 1. \quad (3.91)$$

Further, the integrand satisfies

$$f(\ell) = O\left(\ell^{\frac{3\gamma_1-2}{2}}\right), \quad \text{as } \ell \rightarrow 0. \quad (3.92)$$

Since $\gamma_1 > \frac{2}{3}$ it follows from result (I) that there exists some finite time t_i , defined by $t_i = \lim_{\ell \rightarrow 0} t(\ell)$, such that $\ell(t_i) = 0$.

Similarly, we can integrate (3.89) to obtain

$$t(\ell) = t(1) + \int_1^\ell f(\tilde{\ell}) d\tilde{\ell}, \quad \text{for } \ell > 1. \quad (3.93)$$

Further, the integrand satisfies

$$f(a) = O\left(\ell^{\frac{3\gamma_n-2}{2}}\right), \quad \text{as } \ell \rightarrow \infty. \quad (3.94)$$

Since $\gamma_n \geq 0$ it follows from result (II) $t \rightarrow \infty$ as $\ell \rightarrow \infty$.

For the following two cases we make use of the fact that \tilde{q} and $\ddot{\ell}$ are related by

$$\frac{\ddot{\ell}}{\ell} = -\frac{1}{3}\mu\tilde{q}, \quad (3.95)$$

as follows from (3.22), (3.45) and (3.50).

Case ii) For a recollapsing model, Proposition 3.1.6 implies $\tilde{q}(\tilde{\tau}) > 0$ for all $\tilde{\tau} \in \mathbb{R}$. Then (3.95) implies that $\ell(t)$ is concave down on its domain and so we can conclude there exists t_i and t_f such that $\ell(t_i) = 0$ and $\ell(t_f) = 0$.

Case iii) For a bouncing model, Proposition 3.1.6 implies $\tilde{q}(\tilde{\tau}) < 0$ for all $\tilde{\tau} \in \mathbb{R}$. Then (3.95) implies that $\ell(t)$ is concave up on its domain and hence $\ell \rightarrow \infty$ as $\tilde{\tau} \rightarrow \pm\infty$. Since $\text{sign}(\dot{\ell}) = \text{sign}(H)$, it follows that ℓ is minimal at $\tilde{\tau}_*$. If we choose $\ell_{min} = \ell(\tilde{\tau}_*)$, we have that $\ell(\tilde{\tau}) \geq \ell_{min} > 0$ for all $\tilde{\tau} \in \mathbb{R}$.

We must now show that ℓ does not go to infinity in finite time. Choose ℓ_0 so that $H > 0$ for $\ell \geq \ell_0$. Then $f(\ell)$ in (3.90) is defined and continuous for $\ell \geq 1$. We can integrate (3.89) from $\ell = 1$ to obtain (3.93). Since the integrand satisfies

$$f(a) = O\left(\ell^{\frac{3\gamma_n-2}{2}}\right), \quad \text{as } a \rightarrow \infty, \quad (3.96)$$

and $\gamma_n \geq 0$, it follows from result (II) that (3.93) diverges, and so these models satisfy $\ell \rightarrow \infty$ as $t \rightarrow \infty$. By symmetry, we have that $\ell \rightarrow \infty$ as $t \rightarrow -\infty$. \square

The past and future attractors of the dynamical system on the compact state space can now be determined using theorem 3.1.2.

Theorem 3.1.3 *The past and future attractors in the state space of n -fluid FL cosmologies with an expanding epoch are*

$$\mathcal{A}^- = F_1^+ \cup F_n^-, \text{ and } \mathcal{A}^+ = F_1^- \cup F_n^+, \text{ for } \gamma_n < \frac{2}{3}, \quad (3.97)$$

and

$$\mathcal{A}^- = F_1^+, \text{ and } \mathcal{A}^+ = F_1^- \cup M_n^+ \text{ for } \gamma_n > \frac{2}{3}. \quad (3.98)$$

Proof: We begin by obtaining algebraic expressions for χ_i and $\tilde{\Omega}$. From (3.88) we have

$$\mu_i \ell^2 = 3 (\lambda_i \ell)^{-3\gamma_i+2}. \quad (3.99)$$

Then (3.45), (3.88) and (3.99) lead to

$$\chi_i = \frac{(\lambda_i \ell)^{-3\gamma_i+2}}{\sum_{j=1}^n (\lambda_j \ell)^{-3\gamma_j+2}} \quad (3.100)$$

Using (3.47), we write the Friedmann equation (3.25) in the form

$$1 - \tan^2 \tilde{\Omega} = \frac{3K}{\mu \ell^2}. \quad (3.101)$$

Then using (3.45) and (3.99) we have

$$1 - \tan^2 \tilde{\Omega} = \frac{K}{\sum_{j=1}^n (\lambda_j \ell)^{-3\gamma_j+2}}. \quad (3.102)$$

To determine the limits of (3.100) and (3.102) as $\ell \rightarrow 0$, we write

$$\chi_1 = \frac{\lambda_1^{-3\gamma_1+2}}{\sum_{j=1}^n \lambda_j^{-3\gamma_j+2} \ell^{3(\gamma_1-\gamma_j)}}. \quad (3.103)$$

and, using (3.44), (3.99) and (3.100),

$$1 - \tan^2 \tilde{\Omega} = K \chi_1 (\lambda_1 \ell)^{3\gamma_1-2}. \quad (3.104)$$

It follows that

$$\lim_{\ell \rightarrow 0^+} \chi_1 = 1, \quad \lim_{\ell \rightarrow 0^+} \tilde{\Omega} = \pm \frac{\pi}{4}, \quad (3.105)$$

since $\gamma_1 - \gamma_j > 0$ and $\gamma_1 > \frac{2}{3}$.

Similarly, by writing

$$\chi_n = \frac{\lambda_n^{-3\gamma_n+2}}{\sum_{j=1}^n \lambda_j^{-3\gamma_j+2} \ell^{3(\gamma_n-\gamma_j)}}, \quad (3.106)$$

and

$$1 - \tan^2 \tilde{\Omega} = K \chi_n (\lambda_n \ell)^{3\gamma_n - 2}, \quad (3.107)$$

we obtain

$$\lim_{\ell \rightarrow +\infty} \chi_n = 1, \quad \lim_{\ell \rightarrow +\infty} \tilde{\Omega} = \begin{cases} 0, & \text{if } \gamma_n < \frac{2}{3} \\ \pm \frac{\pi}{2}, & \text{if } \gamma_n > \frac{2}{3} \text{ and } K = -1. \end{cases} \quad (3.108)$$

It now follows from Theorem 3.1.2 and equations (3.72) and (3.73) that orbits describing models in the three generic classes approach the equilibrium points given in table 3.1, as $\tilde{\tau} \rightarrow -\infty$ and as $\tilde{\tau} \rightarrow +\infty$.

□

Comment: Since the orbits describing models in the three generic classes approach the equilibrium points in table 3.1 as $\tilde{\tau} \rightarrow \pm\infty$, it follows that the density parameters Ω_i and Ω_k approach their values at the equilibrium points, which we give in table 3.2.

<u>Model Type</u>	<u>$\tilde{\tau} \rightarrow -\infty$</u>	<u>$\tilde{\tau} \rightarrow +\infty$</u>
Ever-expanding models		
$\gamma_n < \frac{2}{3}$	F_1^+	F_n^+
$\gamma_n > \frac{2}{3}, K = -1$	F_1^+	M_n^+
Recollapsing models	F_1^+	F_1^-
Bouncing models ($\gamma_n < \frac{2}{3}$ only)	F_n^-	F_n^+

TABLE 3.1: The past and future attractors of generic n -fluid FL cosmologies.

<u>Equilibrium Point</u>	<u>Behaviour</u>
F_1^\pm	$\Omega_i \rightarrow 0$ for $i \neq 1$, $\Omega_1 \rightarrow 1$, $\Omega_k \rightarrow 0$,
F_n^\pm	$\Omega_i \rightarrow 0$ for $i \neq n$, $\Omega_n \rightarrow 1$, $\Omega_k \rightarrow 0$,
M_i^\pm	$\Omega_i \rightarrow 0$ for $i = 1, \dots, n$, $\Omega_k \rightarrow 1$.

TABLE 3.2: Asymptotic behaviour of Hubble-normalized quantities in generic n -fluid FL cosmologies.

3.2 Parameterization of FL Models

In this section we introduce the n essential parameters that label the family of n -fluid FL cosmologies, namely a conformal parameter λ and $n - 1$ dimensionless mass parameters m_1, \dots, m_{n-1} , that we shall refer to as *intrinsic parameters*. In section 3.2.2, we introduce an equivalent set of parameters, namely the observational parameters $\Omega_{i,0}$ and H_0 and show how they determine the intrinsic parameters. Finally, we show that the evolution equations (3.57) and (3.60) admit $n - 1$ independent conserved quantities, which characterize the orbits in the n -dimensional state space. In particular, we will show that the values of the conserved quantities are determined in terms of the intrinsic parameters.

3.2.1 The Intrinsic Parameters

We now integrate the conservation equation and demonstrate that the constants of integration of this equation can be used to parameterize n -fluid FL cosmologies. As we have seen previously in (3.88), on integrating (3.20), one obtains

$$\mu_i \ell^2 = 3 (\lambda_i \ell)^{-3\gamma_i+2}. \quad (3.109)$$

The constants of integration λ_i have dimension $(\text{length})^{-1}$ and so each can be thought of as defining a characteristic length scale for each fluid via λ_i^{-1} . When (3.109) is substituted into (3.21), there results a first order ODE for $\ell(t)$:

$$\left(\frac{d\ell}{dt}\right)^2 = \sum_{i=1}^n (\lambda_i \ell)^{-3\gamma_i+2} - K. \quad (3.110)$$

In order to define a dimensionless length scale and time variable for n -fluid FL cosmologies, one needs to choose a preferred λ_i that will play the role of an overall conformal factor. Typically we will choose λ to be λ_n , which is a natural choice if $\gamma_n = 0$. In this case, it follows from (3.109) that

$$\lambda^2 = \frac{1}{3}\Lambda. \quad (3.111)$$

The dimensionless length L and time T are then defined as

$$L = \lambda \ell, \quad \text{and} \quad T = \lambda t. \quad (3.112)$$

Equation (3.109) suggests defining a set of dimensionless parameters m_1, \dots, m_{n-1} via¹⁰

$$m_i = \left(\frac{\lambda_i}{\lambda}\right)^{-3\gamma_i+2}, \quad (3.113)$$

¹⁰For $i = n$, this equation states that $m_n = 1$.

so that

$$\mu_i = 3\lambda^2 m_i L^{-3\gamma_i}. \quad (3.114)$$

In dimensionless form, the Friedmann equation (3.110) then reads

$$\left(\frac{dL}{dT}\right)^2 = \sum_{i=1}^n m_i L^{-3\gamma_i+2} - K. \quad (3.115)$$

For any choice of parameters $m_i > 0$, $i = 1, \dots, n-1$, along with the conditions

$$L(0) = 0, \quad \text{and} \quad \frac{dL}{dT} > 0, \quad (3.116)$$

the DE (3.115) has unique solution (as shown in section C.2), given by

$$L = L(T, m_1, \dots, m_{n-1}). \quad (3.117)$$

It then follows from (3.112) that $\ell(t)$ can be written as

$$\ell(t) = \lambda^{-1} L(\lambda t, m_1, \dots, m_{n-1}). \quad (3.118)$$

The line element then takes the form

$$ds^2 = \lambda^{-2} (-dT^2 + L^2 d\Sigma^2), \quad (3.119)$$

where L is given by (3.117) and $d\Sigma^2$ is a metric of constant curvature.¹¹ From (3.9), (3.112) and (3.115) we obtain

$$H^2 = \lambda^2 \left(\sum_{i=1}^n m_i L^{-3\gamma_i} - K L^{-2} \right). \quad (3.120)$$

Further, using (3.31), (3.114) and (3.120) the density parameters Ω_i are written as

$$\Omega_i = \frac{m_i L^{-3\gamma_i+2}}{\sum_{j=1}^n m_j L^{-3\gamma_j+2} - K}. \quad (3.121)$$

In summary, equations (3.114) and (3.118) show that the n -fluid FL cosmologies form an n -parameter family, labelled by a conformal factor λ and $n-1$ mass parameters m_1, \dots, m_{n-1} . We will refer to these parameters as the *intrinsic parameters*.

¹¹Equations (3.114) and (3.119) illustrate the well-known scale-invariance of Einstein's field equations: if g_{ij} is a solution, then so is $\tilde{g}_{ij} = \lambda^{-2} g_{ij}$ for any constant λ , with the matter terms scaling appropriately, *e.g.* $\tilde{\mu} = \lambda^2 \mu$.

3.2.2 The Observational Parameters

In this section we introduce the observational parameters associated with n -fluid FL cosmologies, based on the current epoch t_0 , which represents the age of the universe. We denote the values of H , q , Ω_i and Ω_k at t_0 by

$$H_0 = H(t_0), \quad q_0 = q(t_0), \quad \Omega_{i,0} = \Omega_i(t_0), \quad \Omega_{k,0} = \Omega_k(t_0),$$

where H_0 is the *Hubble constant*. Then, the set of constants

$$\{t_0, H_0, q_0, \Omega_{A,0}, \Omega_{k,0}\}, \quad A = 1, \dots, n, \quad (3.122)$$

are collectively referred to as the *observational parameters* (see, for example, Wainwright and Ellis (1997), p55-8).

In addition, we require the parameter ℓ_0 , which denotes the length scale at the present time, *i.e.* $\ell_0 = \ell(t_0)$. This quantity is not an observational parameter, but if $K \neq 0$, ℓ_0 can be determined in terms of observational parameters using (3.35) and (3.36), *i.e.* via

$$\ell_0^2 = \frac{-K}{H_0^2 \Omega_{k,0}} = \frac{-K}{H_0^2 (1 - \sum_{i=1}^n \Omega_{i,0})}. \quad (3.123)$$

In the case of zero curvature ($K = 0$), ℓ_0 can be chosen arbitrarily in order to fix the multiplicative factor that determines ℓ (also see section 3.1.1).

We now present the observational formulation of the n -fluid FL cosmologies. In this formulation, the dimensionless length scale and dimensionless time variable are given by

$$a = \frac{\ell}{\ell_0}, \quad \text{and} \quad T = H_0 t. \quad (3.124)$$

From (3.88) we have

$$\mu_i = 3H_0^2 \Omega_{i,0} a^{-3\gamma_i}. \quad (3.125)$$

Using (3.124) and (3.125), the Friedmann equation (3.14) assumes the form

$$\left(\frac{da}{dT}\right)^2 = \sum_{i=1}^n \Omega_{i,0} a^{-3\gamma_i+2} + \Omega_{k,0}. \quad (3.126)$$

We use uniqueness of the solution to (3.126) under some appropriate initial conditions (see section C.2) to conclude that in an expanding epoch the length scale is given by

$$\ell = \ell_0 a(H_0 t, \Omega_{1,0}, \dots, \Omega_{n,0}). \quad (3.127)$$

It follows from (3.9), (3.124) and (3.126) that the Hubble scalar assumes the form¹²

$$H^2 = H_0^2 \left[\sum_{i=1}^n \Omega_{i,0} a^{-3\gamma_i} + \Omega_{k,0} a^{-2} \right], \quad (3.128)$$

and further that the density parameters (3.31) and (3.45) assume the form

$$\Omega_i = \frac{\Omega_{i,0} a^{-3\gamma_i+2}}{\sum_{j=1}^n \Omega_{j,0} a^{-3\gamma_j+2} + \Omega_{k,0}}, \quad (3.129)$$

and

$$\chi_i = \frac{\Omega_{i,0} a^{-3\gamma_i+2}}{\sum_{j=1}^n \Omega_{j,0} a^{-3\gamma_j+2}}. \quad (3.130)$$

The parameters $\Omega_{k,0}$, q_0 , t_0 in (3.122) can be related to the parameters $\{\Omega_{i,0}, H_0\}$, as follows: First, $\Omega_{k,0}$ is determined in terms of $\Omega_{i,0}$ via the Friedmann equation (3.37):

$$\Omega_{k,0} = 1 - \sum_{i=1}^n \Omega_{i,0}. \quad (3.131)$$

Second, q_0 is determined in terms of $\Omega_{i,0}$ via (3.42):

$$q_0 = \frac{1}{2} \sum_{i=1}^n (3\gamma_i - 2) \Omega_{i,0}. \quad (3.132)$$

Third, knowing H_0 , we can obtain t_0 by integrating (3.126) as

$$t_0 H_0 = \int_0^1 \frac{1}{\sqrt{F(a, \Omega_{i,0})}} da, \quad (3.133)$$

where

$$F(y, \Omega_{i,0}) = 1 + \sum_{i=1}^n \Omega_{i,0} (a^{2-3\gamma_i} - 1), \quad (3.134)$$

and $\Omega_{k,0}$ has been eliminated using (3.131). As a consequence, *the only independent observational parameters are the $n + 1$ quantities defined by*

$$\{\Omega_{i,0}, H_0\}. \quad (3.135)$$

¹²An expression of this form is given by Peacock (1999), eq. (3.8).

It is useful to note that the normalized scale factor can be expressed in terms of the redshift z , defined by

$$1 + z = \frac{\ell_0}{\ell} = \frac{1}{a}, \quad (3.136)$$

(see Wainwright and Ellis (1997), eq. (2.41)). One can thus use (3.128) and (3.129) to express H and Ω_i in terms of z .

We now present the link between the observational parameters and the intrinsic parameters. On substituting $\mu_i = 3H^2\Omega_i$ (see (3.31)) into (3.109) at $t = t_0$, we obtain

$$\lambda_i^{-1} = \ell_0^{-1}(\Omega_{i,0}H_0^2\ell_0^2)^{\frac{1}{3\gamma_i-2}}. \quad (3.137)$$

3.2.3 Conserved Quantities

We now show that the evolution equations (3.57) and (3.60) admit $n - 1$ conserved quantities.

From (3.45) and (3.114) we have that

$$\chi_i = \frac{3\lambda^2}{\mu} m_i L^{-3\gamma_i}. \quad (3.138)$$

We now eliminate λ^2/μ and L by taking products of powers of three distinct matter variables (we require $n \geq 3$). Indeed, it follows from (3.138) that

$$\chi_i^{\gamma_j-\gamma_k} \chi_j^{\gamma_k-\gamma_i} \chi_k^{\gamma_i-\gamma_j} = m_i^{\gamma_j-\gamma_k} m_j^{\gamma_k-\gamma_i} m_k^{\gamma_i-\gamma_j}. \quad (3.139)$$

In the case of $K \neq 0$, we also obtain a conserved quantity describing the curvature of a model. We use (3.9) and (3.47) to rewrite the Friedmann equation (3.14) as

$$|1 - \tan^2 \tilde{\Omega}| = \frac{3}{\mu\ell^2}. \quad (3.140)$$

Then from (3.138) and (3.140), we have

$$\chi_i^{3\gamma_j-2} \chi_j^{-3\gamma_i+2} |1 - \tan^2 \tilde{\Omega}|^{3(\gamma_i-\gamma_j)} = m_i^{3\gamma_j-2} m_j^{-3\gamma_i+2}. \quad (3.141)$$

Since there can be only $n - 1$ independent conserved quantities, we must fix some of the values of i , j and k in (3.139) and (3.141) to obtain a minimal independent set. We choose $i = 1$, $k = n$ in (3.139) (with $2 \leq j \leq n - 1$) and $i = 1$ and $j = n$ in (3.141) and so define¹³

$$\mathcal{M}_j(\chi_1, \dots, \chi_n) \equiv \chi_1^{\gamma_j-\gamma_n} \chi_j^{\gamma_n-\gamma_1} \chi_n^{\gamma_1-\gamma_j}, \quad (3.142)$$

¹³Observe that if we choose $j = 1$ or $j = n$ in (3.142) we simply obtain $\mathcal{M}_1 = 1$ and $\mathcal{M}_n = 1$.

and

$$\mathcal{K}(\chi_1, \chi_n, \tilde{\Omega}) \equiv \chi_1^{\frac{3\gamma_n-2}{3(\gamma_1-\gamma_n)}} \chi_n^{-\frac{(3\gamma_1-2)}{3(\gamma_1-\gamma_n)}} (1 - \tan^2 \tilde{\Omega}). \quad (3.143)$$

Upon recalling that $m_n = 1$, the conserved quantities are then determined by the intrinsic parameters m_1, \dots, m_{n-1} according to

$$\mathcal{M}_j(\chi_1, \dots, \chi_n) = m_1^{\gamma_j-\gamma_n} m_j^{\gamma_n-\gamma_1}, \quad j = 2, \dots, n-1, \quad (3.144)$$

and

$$\mathcal{K}(\chi_1, \chi_n, \tilde{\Omega}) = K m_1^{\frac{3\gamma_n-2}{3(\gamma_1-\gamma_n)}}. \quad (3.145)$$

Comment: We have derived these conserved quantities algebraically, and it follows immediately from (3.144) and (3.145) that

$$\frac{d\mathcal{M}_j}{d\tilde{\tau}} = 0, \quad \frac{d\mathcal{K}}{d\tilde{\tau}} = 0. \quad (3.146)$$

It involves a lengthy calculation to verify (3.146) using the ODEs (3.63) and (3.64).

If $K = 0$ (i.e. $\tilde{\Omega} = \frac{\pi}{4}$) then $\mathcal{K} = 0$. As described in section 3.2.2, we can use the freedom to rescale ℓ_0 to set

$$\lambda_1 = \lambda_n, \quad (3.147)$$

which implies $m_1 = 1$. Then (3.144) gives

$$\mathcal{M}_j = m_j^{\gamma_n-\gamma_1}, \quad j = 2, \dots, n-1. \quad (3.148)$$

The conserved quantities (3.142) and (3.143) can also be expressed in terms of the density parameters, as follows:

$$\mathcal{M}_j(\Omega_1, \dots, \Omega_n) = \Omega_1^{\gamma_j-\gamma_n} \Omega_j^{\gamma_n-\gamma_1} \Omega_n^{\gamma_1-\gamma_j}, \quad j = 2, \dots, n-1, \quad (3.149)$$

$$\mathcal{K}(\Omega_1, \Omega_n, \Omega_k) = -\Omega_1^{\frac{3\gamma_n-2}{3(\gamma_1-\gamma_n)}} \Omega_n^{-\frac{3\gamma_1-2}{3(\gamma_1-\gamma_n)}} \Omega_k, \quad (3.150)$$

The conserved quantities are thus related to the observational parameters by

$$\mathcal{M}_j(\Omega_1, \dots, \Omega_n) = \Omega_{1,0}^{(\gamma_j-\gamma_n)} \Omega_{j,0}^{(\gamma_n-\gamma_1)} \Omega_{n,0}^{(\gamma_1-\gamma_j)}, \quad (3.151)$$

and

$$\mathcal{K}(\Omega_1, \Omega_n, \Omega_k) = -\Omega_{1,0}^{\frac{3\gamma_n-2}{3(\gamma_1-\gamma_n)}} \Omega_{n,0}^{-\frac{(3\gamma_1-2)}{3(\gamma_1-\gamma_n)}} \Omega_{k,0}, \quad (3.152)$$

upon evaluating (3.149) and (3.150) at $t = t_0$. We note that conserved quantities of this type have been previously discovered, for example, by Ehlers and Rindler (1989) for *RDC* Λ FL universes, and more generally by Lake (2006) for *n*-fluid FL universes, using a different method.

3.3 2-Fluid FL Cosmologies - Qualitative Analysis

We now use the results of sections 3.1 and 3.2 to give a qualitative analysis of the dynamics of 2-fluid FL cosmologies. The results of this section, in particular figures 3.1-3.3, are contained in Wainwright (1996), although the use of the conserved quantity \mathcal{K} is new.

3.3.1 General Features

The state space is two-dimensional, being described by the variables $(\tilde{\Omega}, \chi_1, \chi_2)$ subject to the restrictions (3.46). We use (3.46) to eliminate χ_1 via

$$\chi_1 = 1 - \chi_2. \quad (3.153)$$

The state space is then the rectangle in the $(\tilde{\Omega}, \chi_2)$ -plane given by

$$-\pi/2 \leq \tilde{\Omega} \leq \pi/2, \quad \text{and} \quad 0 \leq \chi_2 \leq 1. \quad (3.154)$$

In accordance with theorem 3.1.1, there are two qualitatively different classes of 2-fluid FL cosmologies, depending on whether $\gamma_2 > \frac{2}{3}$ or $\gamma_2 < \frac{2}{3}$ (recall that γ_1 and γ_2 are assumed to satisfy $0 \leq \gamma_2 < \gamma_1 \leq 2$). We will discuss these two classes in sections 3.3.2 and 3.3.3 using the two most important examples from a physical point of view:¹⁴

- i) Universes with radiation and dust (*RDC*-universes, $\gamma_1 = \frac{4}{3}, \gamma_2 = 1$),
- ii) Universes with dust and cosmological constant (*DC* Λ -universes, $\gamma_1 = 1, \gamma_2 = 0$).

An important difference between these two cases is that if $\gamma_2 < \frac{2}{3}$ the Einstein static universe, represented by an equilibrium point E , is included in the family of models. The point E is given by¹⁵

$$\chi_{2,E} = \frac{3\gamma_1 - 2}{3(\gamma_1 - \gamma_2)}, \quad \tilde{\Omega} = 0, \quad K = +1, \quad (3.155)$$

(see equation (3.74)). A second related difference is the behaviour of the compact deceleration parameter \tilde{q} as given by (3.65). If $\gamma_2 > \frac{2}{3}$ then \tilde{q} is positive on the state space (3.62) and all models are decelerating. If

¹⁴Other choices are qualitatively similar

¹⁵The restriction $\gamma_2 < \frac{2}{3}$ ensures that $\chi_{2,E} < 1$, as required.

$\gamma_2 < \frac{2}{3}$ then \tilde{q} is negative on a subset of the state space. Figure 3.2 shows that the orbits of ever-expanding models eventually enter this subset, and as a result the models are accelerating at late times.

The portraits of the orbits are drawn by first sketching the various invariant sets and equilibrium points, as given in section 3.1.5, in particular the boundary of the state space (3.154) and the flat FL invariant set $\tilde{\Omega} = \pm \frac{\pi}{4}$. In the case $\gamma_2 < \frac{2}{3}$, the orbits that are past and future asymptotic to the Einstein static point E should be drawn. At this stage, the orbits that have been drawn represent exceptional models and form a “skeleton” for the state space. The orbits of typical models can now be drawn by noting the past and future attractors as given in theorem 3.1.3. In drawing the portraits it is helpful to note that the evolution equations (3.63) and (3.64) are invariant under the interchange

$$(\tilde{\Omega}, \tilde{\tau}) \rightarrow (-\tilde{\Omega}, -\tilde{\tau}), \quad (3.156)$$

i.e. changing the sign of $\tilde{\Omega}$ reverses the direction of time. In particular, the invariant set $-\frac{\pi}{2} \leq \tilde{\Omega} \leq -\frac{\pi}{4}$ represents cosmologies that evolve in the same way as those with $\frac{\pi}{4} \leq \tilde{\Omega} \leq \frac{\pi}{2}$, with the difference that the direction of time is reversed. These models are contracting throughout their evolution ($\tilde{\Omega} < 0$ implies $H < 0$) and hence are not potential models of the real universe.

In practice the state space can be sketched numerically¹⁶ using the conserved quantity \mathcal{K} that is defined by (3.143) for the family of 2-fluid cosmologies. On substituting (3.153) into (3.155) and choosing $n = 2$, we obtain

$$\mathcal{K}(\tilde{\Omega}, \chi_2) = (1 - \chi_2)^{\frac{3\gamma_2 - 2}{3(\gamma_1 - \gamma_2)}} \chi_2^{-\frac{(3\gamma_1 - 2)}{3(\gamma_1 - \gamma_2)}} (1 - \tan^2 \tilde{\Omega}). \quad (3.157)$$

The orbits are given by

$$\mathcal{K}(\tilde{\Omega}, \chi_2) = \text{constant}. \quad (3.158)$$

The value of \mathcal{K} at E is denoted \mathcal{K}_E and is given by

$$\mathcal{K}_E = (1 - \chi_{2,E})^{\frac{3\gamma_2 - 2}{3(\gamma_1 - \gamma_2)}} (\chi_{2,E})^{-\frac{(3\gamma_1 - 2)}{3(\gamma_1 - \gamma_2)}}. \quad (3.159)$$

It follows that the equation

$$\mathcal{K}(\tilde{\Omega}, \chi_2) = \mathcal{K}_E \quad (3.160)$$

describes the orbits that are past and future asymptotic to E , *i.e.* it describes the stable and unstable manifolds of E .

The value of \mathcal{K} determines whether a model is ever-expanding or not, as shown in the following table.

¹⁶For example, using the `plots[implicitplot]` command in Maple.

<u>Restriction on \mathcal{K}</u>	<u>Qualitative Behaviour</u>
<u>Case $\gamma_2 > \frac{2}{3}$</u>	
$-\infty < \mathcal{K} \leq 0$	Ever-expanding
$0 < \mathcal{K} < +\infty$	Recollapsing
<u>Case $\gamma_2 < \frac{2}{3}$</u>	
$-\infty < \mathcal{K} < \mathcal{K}_E$	Ever-expanding
$\mathcal{K}_E < \mathcal{K} < +\infty$	Recollapsing or bouncing

Further, the sign of \mathcal{K} determines whether the model is open, flat or closed. We note that all generic classes of FL models are covered by the given values of \mathcal{K} .

In the case $\gamma_2 > \frac{2}{3}$ the critical value of \mathcal{K} is $\mathcal{K}_{crit} = 0$, indicating that the flat FL models play an exceptional role as regards to future evolution. In the case $\gamma_2 < \frac{2}{3}$, $\mathcal{K}_{crit} = \mathcal{K}_E$ and the exceptional models are those that are past or future asymptotic to the Einstein static universe. This distinction is clearly indicated by comparing figure 3.1 (*RDC*-universes) with figure 3.2 (*DCA*-universes).

We note that the conserved quantity \mathcal{K} can also be expressed in terms of the density parameters. It follows from (3.150) with $n = 2$ that

$$\mathcal{K}(\Omega_1, \Omega_2, \Omega_k) = -\Omega_1^{\frac{3\gamma_2-2}{3(\gamma_1-\gamma_2)}} \Omega_2^{-\frac{(3\gamma_1-2)}{3(\gamma_1-\gamma_2)}} \Omega_k, \quad (3.161)$$

with

$$\Omega_k = 1 - \Omega_1 - \Omega_2. \quad (3.162)$$

3.3.2 FL Cosmologies with Radiation and Dust (*RDC*-universes)

We now consider 2-fluid FL cosmologies containing radiation ($\gamma_1 = \frac{4}{3}$) and dust ($\gamma_2 = 1$). The portrait of the orbits, shown in figure 3.1 is drawn as described in the previous subsection.

For the *RDC*-universes, the expression (3.157) for the conserved quantity \mathcal{K} becomes

$$\mathcal{K} = \frac{\chi_r}{\chi_d^2} (1 - \tan^2 \tilde{\Omega}). \quad (3.163)$$

Equivalently, the expression (3.161) gives \mathcal{K} in terms of the density parameters as

$$\mathcal{K} = -\frac{\Omega_r \Omega_k}{\Omega_d^2}. \quad (3.164)$$




FIGURE 3.1: The $(\tilde{\Omega}, \chi_2)$ state space for 2-fluid FL cosmologies containing radiation ($\gamma_1 = \frac{4}{3}$) and dust ($\gamma_2 = 1$), showing regions of expansion ($H > 0$, i.e. $\tilde{\Omega} > 0$) and regions of contraction ($H < 0$, i.e. $\tilde{\Omega} < 0$).

The value of \mathcal{K} is infinite or zero on the skeleton of the state space:

$$\begin{aligned} \mathcal{K} &= +\infty && \text{if } \chi_d = 0 \text{ and } |\tilde{\Omega}| < \frac{\pi}{4}, \\ \mathcal{K} &= 0 && \text{if } \chi_d = 1 \text{ or } |\tilde{\Omega}| = \frac{\pi}{4}, \\ \mathcal{K} &= -\infty && \text{if } \tilde{\Omega} = \pm \frac{\pi}{2}, \text{ or } \chi_d = 0 \text{ and } |\tilde{\Omega}| > \frac{\pi}{4}. \end{aligned}$$

Note that \mathcal{K} is indeterminate at the past attractor $\mathcal{A}^- = F_r^+$ ($\chi_d = 0, \tilde{\Omega} = \frac{\pi}{4}$) and at the future attractor $\mathcal{A}^+ = F_r^- \cup M_d^+$ ($\chi_d = 0, \tilde{\Omega} = \frac{\pi}{4}; \chi_d = 1, \tilde{\Omega} = \frac{\pi}{2}$). This indeterminacy arises since infinitely many orbits meet at \mathcal{A}^- and \mathcal{A}^+ .

Discussion: Typical models differ qualitatively depending on the value of \mathcal{K} , which determines where the orbit lies in the state space. Firstly, for ever-expanding models if the orbit passes close to M_r^+ , the dust component will never be significant dynamically (i.e. $\Omega_d \ll 1$) and this will correspond to \mathcal{K} being

large and negative. On the other hand if the orbit passes close to F_d^+ then Ω_d will attain a maximum value close to 1, corresponding to \mathcal{K} being negative and close to zero. Secondly, recollapsing models differ quantitatively by how close the orbit comes to F_d^\pm , *i.e.* by the maximum value of χ_d , which in turn will depend on \mathcal{K} . One can determine the dependence on \mathcal{K} explicitly, as follows:

i) For ever-expanding models, we obtain

$$\Omega_d|_{max} = \frac{1}{1 + 2\sqrt{-\mathcal{K}}}, \quad \mathcal{K} < 0. \quad (3.165)$$

ii) For recollapsing models an expression for $\chi_r|_{min}$ and $\chi_d|_{max}$ can be obtained by setting $\tilde{\Omega} = 0$ in (3.163) and solving the resulting second degree polynomial. We obtain

$$\chi_d|_{max} = \frac{2}{1 + \sqrt{1 + 4\mathcal{K}}}, \quad \chi_r|_{min} = 1 - \chi_d|_{max}, \quad \mathcal{K} > 0. \quad (3.166)$$

3.3.3 FL Cosmologies with Dust and Cosmological Constant

We now consider 2-fluid FL cosmologies containing radiation ($\gamma_1 = \frac{4}{3}$) and dust ($\gamma_2 = 1$). The portrait of the orbits, shown in figure 3.2 is drawn as described in section 3.3.1.

For the $DC\Lambda$ -universes, the expression (3.157) for the conserved quantity \mathcal{K} becomes

$$\mathcal{K}(\tilde{\Omega}, \chi_\Lambda) = \chi_d^{-2/3} \chi_\Lambda^{-1/3} (1 - \tan^2 \tilde{\Omega}). \quad (3.167)$$

Equivalently, the expression (3.161) gives \mathcal{K} in terms of the density parameters as

$$\mathcal{K} = -\Omega_d^{-2/3} \Omega_\Lambda^{-1/3} \Omega_k. \quad (3.168)$$

The critical value of χ_Λ , *i.e.* the value of χ_Λ at the Einstein static equilibrium point, is given by (3.155) as

$$\chi_{\Lambda,E} = \frac{1}{3}. \quad (3.169)$$

It then follows from (3.159) that the critical value of \mathcal{K} is

$$\mathcal{K}_E = \left(\frac{4}{27}\right)^{-1/3}. \quad (3.170)$$

The value of \mathcal{K} is infinite or zero on the skeleton of the state space:

$$\begin{aligned} \mathcal{K} &= +\infty && \text{if } |\tilde{\Omega}| < \frac{\pi}{4} \text{ and } \chi_\Lambda = 0 \text{ or } 1, \\ \mathcal{K} &= 0 && \text{if } \tilde{\Omega} = \pm \frac{\pi}{4}, \\ \mathcal{K} &= -\infty && \text{if } \tilde{\Omega} = \pm \frac{\pi}{2} \text{ or } |\tilde{\Omega}| > \frac{\pi}{4} \text{ and } \chi_\Lambda = 0 \text{ or } 1. \end{aligned}$$

Note that \mathcal{K} is indeterminate at the past attractor $\mathcal{A}^- = F_d^+$ ($\chi_\Lambda = 0, \tilde{\Omega} = \frac{\pi}{4}$) and at the future attractor $\mathcal{A}^+ = deS^+ \cup F_d^-$ ($\chi_\Lambda = 1, \tilde{\Omega} = \frac{\pi}{4}; \chi_\Lambda = 0, \tilde{\Omega} = -\frac{\pi}{4}$).




FIGURE 3.2: The $(\tilde{\Omega}, \chi_2)$ state space for 2-fluid FL cosmologies containing dust ($\gamma_1 = 1$) and cosmological constant ($\gamma_2 = 0$), showing regions of expansion ($H > 0$, *i.e.* $\tilde{\Omega} > 0$), regions of contraction ($H < 0$, *i.e.* $\tilde{\Omega} < 0$), regions of acceleration ($\tilde{q} < 0$) and regions of deceleration ($\tilde{q} > 0$) in relation to the Einstein static solution E , which satisfies $H = 0$ and $\tilde{q} = 0$.




FIGURE 3.3: The $(\tilde{\Omega}, \chi)$ representation of the state space for the FL models with matter, a cosmological constant and spatial curvature. The thumbnail plots depict the length scale $\ell(t)$ of FL models corresponding to orbits in each region of the state space and to each equilibrium point. Cases (1), (2) and (3) represent the three generic families of FL models, *i.e.* their orbits fill an open subset of the state space (2-parameter families). They represent, respectively, ever-expanding models, recollapsing models and bouncing models. On the other hand, cases (4), (5), (6a) and (6b) correspond to a single orbit in the state space (a 1-parameter family), while the cases in the third row represent the unique solutions corresponding to the equilibrium points.

Discussion: Firstly, ever-expanding models differ qualitatively depending on the magnitude and the sign of the curvature scalar Ω_k . If the orbit stays close to flat FL ($\tilde{\Omega} = \frac{\pi}{4}$), the curvature will never be significant dynamically (*i.e.* $\Omega_k \ll 1$) and this will correspond to \mathcal{K} being close to zero. Secondly, recollapsing models (and bouncing models) differ qualitatively by how close the orbit comes to the Einstein static model, *i.e.* by the maximal (or minimal) value of χ_Λ . Since $\chi_\Lambda < \chi_{\Lambda,E}$ in recollapsing models (see (3.155)), the cosmological constant will never be the dominant fluid. Similarly, since $\chi_\Lambda > \chi_{\Lambda,E}$ in bouncing models, there will be no dust dominant epoch, *i.e.* $\chi_d \approx 1$. One can determine the dependence on \mathcal{K} explicitly, as follows:

i) For ever-expanding models, we obtain

$$|\Omega_k|_{max} = \frac{|\mathcal{K}|}{\mathcal{K}_E - \mathcal{K}}, \quad \mathcal{K} < \mathcal{K}_E. \quad (3.171)$$

ii) For recollapsing and bouncing models, an expression for $\chi_\Lambda|_{max}$ and $\chi_\Lambda|_{min}$ can be obtained by setting $\tilde{\Omega} = 0$ in (3.167) and solving the resulting third degree polynomial.

3.4 3-Fluid FL Cosmologies - Qualitative Analysis

We now use the results of sections 3.1-3.3 to give a qualitative analysis of the dynamics of 3-fluid FL cosmologies in which the matter content is radiation, dust and a cosmological constant ($\Lambda > 0$) (labelling the χ_i as χ_r , χ_d and χ_Λ). The analysis given in this section is new.

3.4.1 General Features

The state space is three-dimensional, being described by the variables $(\tilde{\Omega}, \chi_r, \chi_d, \chi_\Lambda)$ subject to the restriction (3.46). We use (3.46) to eliminate χ_r via

$$\chi_r = 1 - \chi_d - \chi_\Lambda. \quad (3.172)$$

The physical state space is then a solid triangular prism in \mathbb{R}^3 given by

$$-\frac{\pi}{2} \leq \tilde{\Omega} \leq \frac{\pi}{2}, \quad 0 \leq \chi_d, \chi_\Lambda \leq 1, \quad \chi_d + \chi_\Lambda \leq 1. \quad (3.173)$$

We can construct a ‘‘skeleton’’ for the state space using various invariant subsets, as shown in figure 3.4. The choice of $\chi_i = 0$ for any one fluid gives a 2-fluid FL cosmology, identified as follows:

$$\begin{aligned}\chi_\Lambda = 0 & \quad RDC\text{-universes,} \\ \chi_d = 0 & \quad RC\Lambda\text{-universes,} \\ \chi_r = 0 & \quad DC\Lambda\text{-universes.}\end{aligned}$$

Further, the orbits in the planes $\tilde{\Omega} = \pm \frac{\pi}{4}$ represent the flat 3-fluid universes.

The Einstein static universe is represented by a line segment of equilibrium points denoted $E_r E_d$, which is the intersection of the planes $\tilde{\Omega} = 0$ and $\tilde{q} = 0$. The stable manifold \mathcal{S}^+ of the set of Einstein static points is a surface through the line segment $E_r E_d$ terminating at deS^- when $\chi_\Lambda = 1$ and at the line segment $F_r^+ F_d^+$ when $\chi_\Lambda = 0$. This surface can be described by an equation of the form

$$\tilde{\Omega} = f(\chi_d, \chi_\Lambda), \quad (3.174)$$

with χ_d and χ_Λ restricted as in (3.173). Because of the symmetry $(\tilde{\Omega}, \tilde{\tau}) \rightarrow (-\tilde{\Omega}, -\tilde{\tau})$ (see (3.156)), the unstable manifold \mathcal{S}^- is the surface

$$\tilde{\Omega} = -f(\chi_d, \chi_\Lambda). \quad (3.175)$$

We know from theorem 3.1.1 that there are three generic classes of $RDC\Lambda$ FL universes. Their orbits are contained in invariant subsets bounded by the surfaces \mathcal{S}^\pm , as follows:

- Ever-expanding universes $\tilde{\Omega} > |f(\chi_d, \chi_\Lambda)|$ (the subset to the right of \mathcal{S}^+ and \mathcal{S}^-),
- Recollapsing universes (the subset between \mathcal{S}^+ and \mathcal{S}^- and below the plane $\tilde{q} = 0$),
- Bouncing universes (the subset between \mathcal{S}^+ and \mathcal{S}^- and above the plane $\tilde{q} = 0$).

The planes $\tilde{\Omega} = 0$ and $\tilde{q} = 0$, although not invariant sets, play an important role in describing the dynamics. The plane $\tilde{\Omega} = 0$ divides the state space into a region of expansion ($\tilde{\Omega} > 0$) and a region of contraction ($\tilde{\Omega} < 0$). The variable \tilde{q} is given by (3.65) which here specializes to

$$\tilde{q} = \chi_r + \frac{1}{2}\chi_d - \chi_\Lambda. \quad (3.176)$$

On using (3.172) to eliminate χ_r , we find that $\tilde{q} = 0$ on the plane

$$\chi_d + 4\chi_\Lambda = 2. \quad (3.177)$$

Equation (3.177) then divides the state space into a region of acceleration ($\tilde{q} < 0$) and a region of deceleration ($\tilde{q} > 0$) as follows:

$$\begin{aligned} \text{Region of acceleration} \quad & \chi_\Lambda > \frac{1}{4}(2 - \chi_d), \\ \text{Region of deceleration} \quad & \chi_\Lambda < \frac{1}{4}(2 - \chi_d). \end{aligned}$$

It follows from the general discussion in section 3.2.3 that there are two conserved quantities for the $RDCA$ FL universes, defined by

$$\mathcal{M} = \left(\frac{\chi_r^3 \chi_\Lambda}{\chi_d^4} \right)^{1/3}, \quad \text{and} \quad \mathcal{K} = \frac{1 - \tan^2 \tilde{\Omega}}{(\chi_r \chi_\Lambda)^{1/2}}. \quad (3.178)$$

Using (3.172) we can eliminate χ_r from these equations and hence obtain \mathcal{M} and \mathcal{K} in terms of $\tilde{\Omega}$, χ_d and χ_Λ . The values of \mathcal{M} and \mathcal{K} at the Einstein static equilibrium points E are denoted \mathcal{M}_E and \mathcal{K}_E and are obtained on substituting (3.177) into (3.178). We have

$$\mathcal{M}_E = \left[\frac{(3\chi_\Lambda - 1)^3 \chi_\Lambda}{16(1 - 2\chi_\Lambda)^4} \right]^{1/3}, \quad 0 \leq \mathcal{M}_E < \infty, \quad (3.179)$$

$$\mathcal{K}_E = [(3\chi_\Lambda - 1)\chi_\Lambda]^{-1/2}, \quad 2 \leq \mathcal{K}_E < \infty. \quad (3.180)$$

The orbits of typical models are described by the equations

$$\mathcal{K}(\tilde{\Omega}, \chi_d, \chi_\Lambda) = \text{constant}, \quad \text{and} \quad \mathcal{M}(\chi_d, \chi_\Lambda) = \text{constant}, \quad (3.181)$$

i.e. the orbits are the intersections of these two families of surfaces in the state space. As a special case the flat universes satisfy $\mathcal{K} = 0$ (*i.e.* $\tilde{\Omega} = \frac{\pi}{4}$) and then the equation

$$\mathcal{M}(\tilde{\Omega}, \chi_d, \chi_\Lambda) = \text{constant} \quad (3.182)$$

describes the orbits in the invariant set $\tilde{\Omega} = \frac{\pi}{4}$ and can be used to numerically sketch them, as in figure 3.5.

Discussion: Knowing the past attractor $\mathcal{A}^- = F_r^+$ and the future attractor $\mathcal{A}^+ = F_r^- \cup deS^+$ one can visualize the orbits of typical universes. For example, the orbits of ever-expanding models join F_r^+ to deS^+ . The models will differ qualitatively depending on whether they come close to F_d^+ ($\Omega_d \approx 1$ in a neighbourhood of F_d^+) and whether they come close to the Milne set ($\tilde{\Omega} = \frac{\pi}{2}$) or to the surfaces \mathcal{S}^\pm , in which case there will be an epoch in which spatial curvature is dominant (*i.e.* $|\Omega_k| \approx 1$). One can determine the points in state space at which $|\Omega_k|$ and χ_d attain a maximum value, as follows. If $K \neq 0$, it follows from (3.36) and (3.69) that $\tilde{\Omega}$ and $|\Omega_k|$ are simultaneously extremal. Since in an ever-expanding

model $\tilde{\Omega}$ is extremal when $\tilde{q} = 0$, we conclude that $|\Omega_k|$ is extremal on the plane given by (3.177). The matter variable χ_d will attain a maximum when $\chi'_d = 0$, which implies $\tilde{q} = \frac{1}{2}$, as follows from (3.64) and (3.65). After substituting (3.172) to eliminate the χ_r term in (3.176), we conclude that χ_d will attain a maximum on the plane

$$\chi_d + 4\chi_\Lambda = 1. \quad (3.183)$$

One can then determine the dependence of $\chi_d|_{max}$ on \mathcal{M} explicitly from (3.172), (3.178) and (3.183), as follows:

$$\chi_d|_{max} = \frac{3^{3/4}}{4\mathcal{M}^{3/4} + 3^{3/4}}. \quad (3.184)$$

3.4.2 A Model of the Physical Universe

In this section we discuss the *RDCA* FL universes as viable models for the large scale dynamical behaviour of the physical universe. Current observations of distant galaxies, type I_a supernovae and the cosmic microwave background radiation support the following conclusions:

- i) The universe is expanding in the present epoch ($H_0 > 0$).
- ii) The universe is accelerating in the present epoch ($q_0 < 0$).
- iii) The early universe was radiation-dominated (hot big-bang).
- iv) Pressure-free matter (baryonic matter and dark matter) is dynamically significant in the present epoch.
- v) The spatial geometry is close to flatness in the present epoch.

These conclusions suggest modelling the physical universe as an *RDCA* FL cosmology, with observational parameters

$$\Omega_{r,0}, \quad \Omega_{d,0}, \quad \Omega_{k,0}, \quad \Omega_{\Lambda,0}, \quad q_0, \quad H_0, \quad \text{and} \quad t_0. \quad (3.185)$$

Equations (3.131)-(3.133) limit the number of independent quantities in this set to four, as expected for an *RDCA* model. Values for these parameters have been determined using observations, such as high redshift galaxy surveys (for example, by 2dF and SDSS; see Spergel et al. (2006) for detailed references) and analyses of the power spectrum of the cosmic microwave background (for example, by WMAP). We




FIGURE 3.4: The skeleton of the state space for $RDCA$ cosmologies, using $\tilde{\Omega}$, χ_d and χ_Λ as variables, with $\chi_r = 1 - \chi_d - \chi_\Lambda$.




FIGURE 3.5: The invariant set $\tilde{\Omega} = \frac{\pi}{4}$ in the state space for the $RDCA$ universes, describing the expanding flat FL models.

refer the reader to Spergel et al. (2006) and Tegmark et al. (2004) for an analysis of recent observational data, leading to the following values:

$$H_0 = 71.0 \pm 4.0 \text{ km s}^{-1} \text{ Mpc}^{-1}, \quad (3.186)$$

$$\Omega_{r,0} = (4.9 \pm 0.5) \times 10^{-5}, \quad (3.187)$$

$$\Omega_{d,0} = 0.27 \pm 0.03, \quad (3.188)$$

$$\Omega_{\Lambda,0} = 0.73 \pm 0.04, \quad (3.189)$$

$$\Omega_{k,0} = -0.010 \pm 0.014. \quad (3.190)$$

Using the observational values (3.186)-(3.190), one can apply (3.63)-(3.65) and (3.69)-(3.71) to numerically obtain the past and future behaviour of all matter quantities χ_i and the curvature indicator Ω . We present the results of one such numerical simulation in figure 3.6, along with computed epochs of note.

The conserved quantities in the $RDCA$ universe, as given by (3.178), can be expressed in terms of the density parameters as

$$\mathcal{M} = \left(\frac{\Omega_r^3 \Omega_\Lambda}{\Omega_d^4} \right)^{\frac{1}{3}}, \quad (3.191)$$

and

$$\mathcal{K} = \Omega_r^{-\frac{1}{2}} \Omega_\Lambda^{-\frac{1}{2}} \Omega_k, \quad (3.192)$$


(see (3.149) and (3.150)). Then using (3.187)-(3.190), we obtain

$$\mathcal{M} = (2.63 \pm 0.07) \times 10^{-3}, \quad (3.193)$$

and

$$\mathcal{K} = -1.72 \pm 1.91. \quad (3.194)$$

The values of these conserved quantities, which were fixed at the end of the inflationary epoch, determine a small subset of $RDCA$ universes that can potentially describe the real universe. The value of \mathcal{M} ensures the occurrence of a matter-dominated epoch that is sufficiently long, but not too long (see equation (3.184)). The value of \mathcal{K} ensures that Ω_k is close to 0 in the past and also into the future.



Epoch	Redshift (z)	Temperature (T)
A Big-bang nucleosynthesis	3.71×10^8	$1.01 \times 10^9 K$
B Matter-radiation equality	5380	9700K
C Last Scattering	1100	3010K
D Maximal matter density	13.5	39.6K
E Maximal spatial curvature	0.773	4.83K
F Matter- Λ equality	0.374	3.50K
G Present day	0.0	2.73K

FIGURE 3.6: Past and future evolution of the $RDC\Lambda$ model of the physical universe, depicting the matter variables (top), Hubble-normalized energy density (bottom) and epochs of primary interest.

3.5 Explicit FL Cosmologies

In this section we give a unified description of the known solutions of the Friedmann DE that can be expressed in terms of elementary functions, for 1-fluid and 2-fluid FL cosmologies. In these cases, the Friedmann DE can be transformed to the form

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}}\right)^2 = Q(\mathcal{L}, m, K), \quad (3.195)$$

where Q is a *quadratic function* in \mathcal{L} , m is a dimensionless mass parameter and K is the curvature indicator. Here \mathcal{L} is a function of the length scale factor ℓ and the variable \mathcal{T} is either proportional to conformal time η (in the case $\Lambda = 0$) or proportional to clock time t (in the case $\Lambda > 0$).

Recall that the n -fluid Friedmann equation can be written as a DE of the form (3.110), reproduced here for convenience:

$$\left(\frac{d\ell}{dt}\right)^2 = \sum_{i=1}^n (\lambda_i \ell)^{-3\gamma_i+2} - K. \quad (3.196)$$

We are primarily interested in expanding models, which begin at a big-bang singularity, *i.e.* the length scale satisfies $\ell(t_s) = 0$ at some time t_s , and $\frac{d\ell}{dt} > 0$ in some interval. Since the DE (3.196) is autonomous, it is invariant under the change $t \rightarrow t + C$. We will use this freedom to set $t_s = 0$, thereby fixing the constant of integration that arises in solving (3.196). We will thus solve (3.196) subject to the requirement $\frac{d\ell}{dt} > 0$ and the initial condition

$$\ell(0) = 0. \quad (3.197)$$

Using conformal time (D.1), equation (3.196) becomes a first order ODE for $\ell(\eta)$:

$$\frac{1}{\ell^2} \left(\frac{d\ell}{d\eta}\right)^2 = \sum_{i=1}^n (\lambda_i \ell)^{-3\gamma_i+2} - K. \quad (3.198)$$

The solution $\ell(\eta)$ can in principle be expressed in terms of clock time t using the equation (D.1). As with (3.196), when solving (3.198) for $\ell(\eta)$, we will impose the requirement $\frac{d\ell}{d\eta} > 0$ and the initial condition

$$\ell(0) = 0. \quad (3.199)$$

In the following sections we consider five classes of models in which either (3.196) or (3.198) can be transformed into (3.195). The solution of (3.195) subject to the requirement $\frac{d\mathcal{L}}{d\mathcal{T}} > 0$ and the initial condition $\mathcal{L}(0) = 0$ is given in Appendix C. In each case, we give the solution for $\ell(t)$ or $\ell(\eta)$ and note that the corresponding expressions for the energy densities and pressure are given by equation (3.15), (3.17) and (3.114).

3.5.1 Single Fluid FL Cosmologies

In this case we consider solutions to the Friedmann equation with $n = 1$, $\frac{2}{3} < \gamma \leq 2$ and arbitrary curvature. We choose the conformal factor to be $\lambda = \lambda_1$. The Friedmann DE (3.198) assumes the form

$$\frac{1}{\ell^2} \left(\frac{d\ell}{d\eta} \right)^2 = (\lambda\ell)^{-3\gamma+2} - K. \quad (3.200)$$

We perform the change of variable

$$\mathcal{L} = (\lambda\ell)^{1/\beta}, \quad \mathcal{T} = \frac{\eta}{\beta}, \quad (3.201)$$

with

$$\beta = \frac{2}{3\gamma - 2}, \quad (3.202)$$

so as to obtain

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}} \right)^2 = 1 - K\mathcal{L}^2. \quad (3.203)$$

Then from (C.3) and (C.4), the solution is

$$\mathcal{L} = S_K(\mathcal{T}), \quad (3.204)$$

where

$$S_K(\mathcal{T}) = \begin{cases} \sinh(\mathcal{T}), & \text{open FL } (K = -1), \\ \mathcal{T}, & \text{flat FL } (K = 0), \\ \sin(\mathcal{T}), & \text{closed FL } (K = +1). \end{cases} \quad (3.205)$$

Thus,

$$\ell(\eta) = \lambda^{-1} \left[S_K \left(\frac{\eta}{\beta} \right) \right]^\beta. \quad (3.206)$$

Using (3.114), the energy density is given by

$$\mu = 3\lambda^2 L^{-3\gamma}, \quad L = \lambda\ell. \quad (3.207)$$

This solution, for arbitrary γ , was first given by Harrison (1967) (see equations (152), (18) and (32) in Harrison (1967), for open, flat and closed models, respectively). Specific cases were discovered earlier, however. Friedmann (1922) gave this solution in the case of dust and positive curvature (*i.e.* $\gamma = 1$ and $K = +1$) and Tolman (1931) gave this solution in the case of radiation and positive curvature (*i.e.* $\gamma = \frac{4}{3}$ and $K = +1$).

These solutions describe two qualitatively different classes of models: ever-expanding models if $K = -1$ or 0 , and recollapsing models if $K = +1$. In this case, it follows from (3.205) and (3.206) that recollapse (*i.e.* $\ell \rightarrow 0$ as $\eta \rightarrow \eta_f$) will occur at

$$\eta_f = \beta\pi. \quad (3.208)$$

For flat FL ($K = 0$) we can obtain $\ell = \ell(t)$ explicitly by integrating (3.196) with $n = 1$ and $K = 0$. The result is

$$\ell(t) = \lambda^{-1} \left(\frac{3\gamma}{2} \lambda t \right)^{\frac{2}{3\gamma}}. \quad (3.209)$$

The parameter λ is not essential, since by rescaling ℓ it can be assigned any value. Equation (3.209) thus represents the unique flat FL universe with a single fluid.¹⁷

In the case of radiation ($\gamma = \frac{4}{3}$, $\beta = 1$) one can integrate (D.1) to obtain,

$$t = \begin{cases} \lambda^{-1}(\cosh \eta - 1), & \text{open FL } (K = -1), \\ \lambda^{-1}(1 - \cos \eta), & \text{closed FL } (K = +1). \end{cases} \quad (3.210)$$

Substituting back into (3.204) and rearranging yields

$$\ell(t) = \lambda^{-1} [2\lambda t - K(\lambda t)^2]^{1/2}. \quad (3.211)$$

We note that the second term here represents the contribution to the solution from spatial curvature, *i.e.* the drift from flat FL. If $K = -1$, the limit $\lambda \rightarrow \infty$, gives the Milne solution (given in section 3.1.2, equation (3.28)).

In the case of dust ($\gamma = 1$, $\beta = 2$) one can integrate (D.1) in closed form, but cannot invert the expression in terms of elementary functions. In this case t and η are related by

$$t = \begin{cases} \frac{1}{2}\lambda^{-1}(\sinh \eta - \eta), & \text{open FL } (K = -1), \\ \frac{1}{2}\lambda^{-1}(\eta - \sin \eta), & \text{closed FL } (K = +1). \end{cases} \quad (3.212)$$

¹⁷That (3.209) does define a unique solution is confirmed by the fact that the physical quantities H and μ do not depend on λ , being given by

$$H = \frac{2}{3\gamma} t^{-1}, \quad \text{and} \quad \mu = \frac{4}{3\gamma^2} t^{-2},$$

on using (3.9), (3.32) and the fact that $\Omega = 1$.

3.5.2 FL Vacuum Cosmologies

In this case we solve the Friedmann equation where the only fluid component is a cosmological constant, *i.e.* $n = 1$, $\gamma_1 = 0$. We choose the conformal factor to be $\lambda = \lambda_1$, which leads to

$$\lambda = \sqrt{\frac{1}{3}\Lambda}, \quad (3.213)$$

(see equation (3.111)). The Friedmann equation (3.196) in this case is

$$\left(\frac{d\ell}{dt}\right)^2 = (\lambda\ell)^2 - K. \quad (3.214)$$

We perform the change of variable

$$\mathcal{L} = \lambda\ell, \quad \mathcal{T} = \lambda t, \quad (3.215)$$

so as to obtain

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}}\right)^2 = \mathcal{L}^2 - K. \quad (3.216)$$

This DE can only be solved subject to $\mathcal{L}(0) = 0$ and $\frac{d\mathcal{L}}{d\mathcal{T}} > 0$ in the case of open FL, with the solution given by (C.3). For the case of non-negative curvature, we instead require that $\mathcal{L}(0) = 1$ and that \mathcal{L} is increasing for all $\mathcal{T} > 0$. It follows that (3.216) has a unique solution in each case, leading to

$$\ell(t) = \begin{cases} \lambda^{-1} \sinh(\lambda t), & \text{open FL } (K = -1), \\ \lambda^{-1} \exp(\lambda t), & \text{flat FL } (K = 0), \\ \lambda^{-1} \cosh(\lambda t), & \text{closed FL } (K = +1), \end{cases} \quad (3.217)$$

where λ is given by (3.213). The energy density is constant and given by

$$\mu = 3\lambda^2 = \Lambda. \quad (3.218)$$

These solutions were first given by Robertson (1933), de Sitter (1917) and Lanczos (1922) in the case of negative, zero and positive curvature, respectively. They correspond to the orbits $M^+ \rightarrow deS^+$ for $K = -1$, deS^+ for $K = 0$ and $deS^- \rightarrow deS^+$ for $K = +1$, along the 1-fluid cosmological constant manifold in figure 3.3. In the case of flat FL, (3.217) gives the *de Sitter* solution, described in section 3.1.2, equation (3.27).

Often, in the case of the de Sitter solution, it is preferable to use conformal time instead of clock time. In particular, when $K = 0$ this choice of time variable is defined on an interval $(-\infty, \eta_f)$ (see Appendix D). Upon defining

$$\tilde{\eta} = \eta_f - \eta, \quad (3.219)$$

the length scale (3.217) for a flat background is simply

$$\ell(\eta) = \frac{1}{\lambda \tilde{\eta}}. \quad (3.220)$$

3.5.3 Flat FL with a Single Fluid and Λ

In this case we solve the Friedmann equation with $n = 2$, $\gamma_1 = \gamma$ arbitrary, $\gamma_2 = 0$ and zero curvature (*i.e.* $K = 0$). We choose the conformal factor to be $\lambda = \lambda_2$, which leads to (3.213). As described in section 3.2.2, we can use the freedom to scale ℓ to set $\lambda_1 = \lambda$. The Friedmann equation (3.196) then takes the form

$$\left(\frac{d\ell}{dt}\right)^2 = (\lambda\ell)^{-3\gamma+2} + (\lambda\ell)^2. \quad (3.221)$$

We apply the change of variable

$$\mathcal{L} = (\lambda\ell)^{\frac{3}{2}\gamma}, \quad \mathcal{T} = \frac{3}{2}\gamma\lambda t \quad (3.222)$$

so as to write (3.196) as

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}}\right)^2 = 1 + \mathcal{L}^2. \quad (3.223)$$

Then using (C.3), we obtain

$$\mathcal{L} = \sinh \mathcal{T}, \quad (3.224)$$

and so

$$\ell(t) = \lambda^{-1} \left[\sinh \left(\frac{3\gamma}{2} \lambda t \right) \right]^{\frac{2}{3\gamma}}. \quad (3.225)$$

Using (3.114), the energy density is given by

$$\mu = \mu_f + \mu_\Lambda = 3\lambda^2 L^{-3\gamma} + \Lambda, \quad L = \lambda\ell. \quad (3.226)$$

These solutions were first given by Harrison (1967) in equation (23). They represent ever-expanding models and correspond to the orbit $F^+ \rightarrow deS^+$ along the $K = 0$ invariant set, as in figure 3.3.

3.5.4 FL Cosmologies with Radiation and Dust (*RDC-Universes*)

In this case we solve the Friedmann equation with $n = 2$, $\gamma_1 = \frac{4}{3}$ and $\gamma_2 = 1$. We choose $\lambda = \lambda_2$ as the conformal parameter. Then, by (3.113), the single mass parameter is given by

$$m = m_r = \left(\frac{\lambda_r}{\lambda}\right)^{-2}. \quad (3.227)$$

The Friedmann equation (3.198) assumes the form

$$\frac{1}{\ell^2} \left(\frac{d\ell}{d\eta}\right)^2 = m(\lambda\ell)^{-2} + (\lambda\ell)^{-1} - K. \quad (3.228)$$

We apply the change of variable

$$\mathcal{L} = \frac{1}{\sqrt{m}}(\lambda\ell), \quad \mathcal{T} = \eta, \quad (3.229)$$

which gives

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}}\right)^2 = 1 + \frac{1}{\sqrt{m}}\mathcal{L} - K\mathcal{L}^2. \quad (3.230)$$

Using (C.3) and (C.4), the solution is

$$\mathcal{L} = S_K(\mathcal{T}) + \frac{1}{\sqrt{m}}S_K\left(\frac{1}{2}\mathcal{T}\right)^2, \quad (3.231)$$

and so

$$\ell(\eta) = \lambda^{-1} \left(\sqrt{m}S_K(\eta) + S_K\left(\frac{1}{2}\eta\right)^2 \right), \quad (3.232)$$

where S_K is given by (3.205). Using (3.114), the energy density is given by

$$\mu = \mu_r + \mu_d = 3\lambda^2 m L^{-4} + 3\lambda^2 L^{-3}, \quad L = \lambda\ell. \quad (3.233)$$

In this case, one can integrate (D.1) so as to obtain an expression for the clock time $t = t(\eta)$, but cannot invert this expression to obtain $\ell = \ell(t)$:

$$t = \begin{cases} \lambda^{-1} \left(\sqrt{m}(\cosh \eta - 1) + \frac{1}{2}(\sinh \eta - \eta) \right), & \text{open } (K = -1), \\ \lambda^{-1} \left(\frac{1}{2}\sqrt{m}\eta^2 + \frac{1}{12}\eta^3 \right), & \text{flat } (K = 0), \\ \lambda^{-1} \left(\sqrt{m}(1 - \cos \eta) + \frac{1}{2}(\eta - \sin \eta) \right), & \text{closed } (K = +1). \end{cases} \quad (3.234)$$

The solution (3.232) contains the one-fluid dust and radiation solutions given in (3.206) as special cases. One can see this by writing (3.232) in terms of parameters $\lambda_r = \lambda_1$ and $\lambda_d = \lambda_2$,

$$\ell(\eta) = \lambda_r^{-1}S_K(\eta) + \lambda_d^{-1}S_K\left(\frac{1}{2}\eta\right)^2, \quad (3.235)$$

and letting $\lambda_r \rightarrow \infty$ or $\lambda_d \rightarrow \infty$.

These solutions were first given¹⁸ by Chernin (1966). They describe all orbits in the state space for *RDC*-universes with $H > 0$ in some epoch (see figure 3.1) and hence describe two qualitatively different classes of models: ever-expanding models if $K = -1$ (the orbits $F_1^+ \rightarrow M^+$) or $K = 0$ (the orbit $F_1^+ \rightarrow F_2^+$) and recollapsing models if $K = +1$ (the orbits $F_1^+ \rightarrow F_1^-$). The dimensionless parameter m determines key physical properties of the models. First, it determines the epoch of matter-radiation equality ($\mu_r = \mu_d$) according to

$$L_{eq} = m, \quad (3.236)$$

as follows from (3.233). Second, for recollapsing models, m determines the time of recollapse, according to

$$\eta_f = 2(\pi - \arctan(2\sqrt{m})), \quad (3.237)$$

as follows from (3.232).

3.5.5 FL Cosmologies with Radiation and Λ (*RCA*-Universes)

In this case we solve the Friedmann equation with $n = 2$, $\gamma_1 = \frac{4}{3}$, $\gamma_2 = 0$ and arbitrary curvature. We choose the conformal factor to be $\lambda = \lambda_2$ so that (3.213) holds. Then, by (3.113), the single mass parameter is given by

$$m = m_1 = \left(\frac{\lambda_1}{\lambda}\right)^{-2}. \quad (3.238)$$

The Friedmann equation (3.196) then takes the form

$$\left(\frac{d\ell}{dt}\right)^2 = m(\lambda\ell)^{-2} + (\lambda\ell)^2 - K. \quad (3.239)$$

Upon making a change of variable according to

$$\mathcal{L} = \frac{1}{\sqrt{m}}(\lambda\ell)^2, \quad \mathcal{T} = 2\lambda t, \quad (3.240)$$

we obtain

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}}\right)^2 = 1 - \frac{1}{\sqrt{m}}K\mathcal{L} + \mathcal{L}^2. \quad (3.241)$$

¹⁸Harrison reports that in the case $K = +1$ this solution was given by Lemaître (1927), but we have not verified this reference.

Using (C.3), the solution is

$$\mathcal{L} = \sinh \mathcal{T} - \frac{1}{\sqrt{m}} K \sinh^2\left(\frac{1}{2}\mathcal{T}\right), \quad (3.242)$$

and so

$$\ell(t) = \lambda^{-1} \left[\sqrt{m} \sinh(2\lambda t) - K \sinh^2(\lambda t) \right]^{1/2}. \quad (3.243)$$

Using (3.114), the energy density is given by

$$\mu = \mu_r + \mu_\Lambda = 3\lambda^2 m L^{-4} + \Lambda, \quad L = \lambda \ell. \quad (3.244)$$

The solution (3.243) contains the $C\Lambda$ vacuum solutions given in (3.217) and the one-fluid radiation solution given in (3.211) as special cases. One can see this by writing (3.243) in terms of parameters $\lambda_r = \lambda_1$ and $\lambda_\Lambda = \lambda_2$,

$$\ell(\eta) = \left[\lambda_r^{-1} \lambda_\Lambda^{-1} \sinh(2\lambda_\Lambda t) - K \lambda_\Lambda^{-2} \sinh^2(\lambda_\Lambda t) \right]^{1/2}, \quad (3.245)$$

and letting $\lambda_r \rightarrow \infty$ or $\lambda_\Lambda \rightarrow 0$. Also, in the case of zero curvature ($K = 0$), (3.243) reduces to (3.225) with $\gamma = \frac{4}{3}$.

This solution was first given by Harrison (1967) (see equations (59) and (38), for the case of negative ($K = -1$) and positive ($K = +1$) curvature, respectively). It describes all orbits in the state space for $RC\Lambda$ -universes with $H > 0$ in some epoch (see figure 3.2) and an initial singularity. Hence, this solution describes both ever-expanding models and recollapsing models. The parameter m determines key physical properties of the models. First, it determines the epoch of radiation-cosmological constant equality ($\mu_r = \mu_\Lambda$) according to

$$L_{eq} = m^{\frac{1}{4}}, \quad (3.246)$$

as follows from (3.213) and (3.244). Second, for recollapsing models, m determines the time of recollapse according to

$$t_f = \lambda^{-1} \operatorname{arctanh}(2\sqrt{m}). \quad (3.247)$$

This result follows upon rewriting (3.243) using standard hyperbolic identities as

$$\ell(t) = \lambda^{-1} \left[\frac{1}{2} \sinh(2\lambda t) (2\sqrt{m} - K \tanh(\lambda t)) \right]^{1/2}. \quad (3.248)$$

We note that at the critical value of m , denoted m^* and given by $m^* = \frac{1}{4}$, the solution (3.243) simplifies to

$$\ell(t) = \lambda^{-1} \left[\frac{1}{2} (1 - \exp(-2\lambda t)) \right]^{\frac{1}{2}}. \quad (3.249)$$

This solution is future asymptotic to the Einstein static solution and corresponds to the orbit $F^+ \rightarrow E$ in figure 3.2.

3.5.6 Explicit Solutions in Terms of Observational Parameters

In sections 3.5.1-3.5.5, we have presented the explicit solutions using intrinsic parameters, namely, a conformal factor λ , the curvature indicator K and, for the 2-fluid solutions with non-zero curvature, a dimensionless mass parameter m , which describes the relative significance of the two fluids.

In order to link with observations it is desirable to write the solutions in terms of the observational parameters. In order to do so we need to use the formula that relates the intrinsic parameters λ_i to the observational parameters, derived in section 3.2.2. In the case of $K \neq 0$, we can use equations (3.123) and (3.137) to write

$$\lambda_i^{-2} = \ell_0^2 \left(\frac{\Omega_{i,0}}{|\Omega_{k,0}|} \right)^{\frac{2}{3\gamma_i-2}}. \quad (3.250)$$

One can think of (3.250) as relating λ_i^{-1} , the length scale determined by the i^{th} fluid, to ℓ_0 , the length scale determined by the spatial curvature. In particular, we will need the following special cases of (3.250):

$$\lambda_r^{-2} = \ell_0^2 \left(\frac{\Omega_{r,0}}{|\Omega_{k,0}|} \right), \quad \text{Radiation } (\gamma_r = \frac{4}{3}), \quad (3.251)$$

$$\lambda_d^{-2} = \ell_0^2 \left(\frac{\Omega_{d,0}}{|\Omega_{k,0}|} \right)^2, \quad \text{Dust } (\gamma_d = 1), \quad (3.252)$$

Note that the Friedmann equation (3.36), evaluated at t_0 gives

$$\Omega_{k,0} = 1 - \Omega_0, \quad (3.253)$$

where Ω_0 is the total density parameter evaluated at t_0 . In the case of a cosmological constant (*i.e.* $\gamma = 0$), we can obtain an expression for λ_Λ directly from (3.137), as follows:

$$\lambda_\Lambda^{-2} = \frac{1}{\Omega_{\Lambda,0} H_0^2} = \frac{3}{\Lambda}, \quad (\gamma_\Lambda = 0). \quad (3.254)$$

In the case of $K = 0$ and $n > 1$, we use the freedom to scale ℓ in order to set λ_1 equal to λ_n . Then, as a result of (3.137), the length scale ℓ_0 is determined explicitly in terms of $\Omega_{i,0}$ and H_0 according to

$$(\ell_0 H_0)^{6(\gamma_1 - \gamma_n)} = \Omega_{1,0}^{3\gamma_n - 2} \Omega_{n,0}^{-(3\gamma_1 - 2)}. \quad (3.255)$$

The intrinsic parameters λ_i are then obtained via (3.137).

There is no work to be done in writing the densities in observational form, since they are given in general by equation (3.125), which we repeat for convenience:

$$\mu_i = 3H_0^2 \Omega_{i,0} a^{3\gamma_i}, \quad (3.256)$$

where $a = \ell/\ell_0$.

In order to illustrate this process we give the length scale function for the *RDC*-universes in terms of the observational parameters. We first rewrite $\ell(\eta)$ in (3.232) in terms of λ_r and λ_d :

$$\ell(\eta) = \lambda_r^{-1} S_K(\eta) + \lambda_d^{-1} S_K(\frac{1}{2}\eta)^2. \quad (3.257)$$

Then (3.251) and (3.252) immediately give

$$\ell(t) = \ell_0 \left[\left(\frac{\Omega_{r,0}}{|1 - \Omega_0|} \right)^{1/2} S_K(\eta) + \left(\frac{\Omega_{d,0}}{|1 - \Omega_0|} \right) S_K(\frac{1}{2}\eta)^2 \right], \quad (3.258)$$

where $\Omega_0 = \Omega_{r,0} + \Omega_{d,0}$. Then $\Omega_{r,0} = 0$ gives the *DC*-universe and $\Omega_{d,0} = 0$ gives the *RC*-universe. Peacock (1999) refers to the terms

$$\frac{\Omega_{d,0}}{2|1 - \Omega_0|}, \quad \text{and} \quad \frac{\Omega_{r,0}}{2|1 - \Omega_0|}$$

as the ‘‘dimensionless masses’’ (p79), when discussing *RDC*-universes. Note that these quantities are closely related to those given in (3.251) and (3.252), upon applying (3.253). These quantities also appear in the *DC*- and *RC*-universes as given by Coles and Lucchin (1995) (see p39, eq (2.4.2) for dust in terms of η and p41, eq (2.5.2) for radiation in terms of t).

3.6 Discussion

We summarize the principal features of the dynamics of *n*-fluid FL universes.

- i) *n*-fluid FL universes have at most one epoch of expansion ($H > 0$) and at most one epoch of acceleration ($q < 0$, or equivalently, $w < -\frac{1}{3}$). These results depend crucially on the assumption that the fluids are non-interacting. We refer to Tolman (1934), p402 and p429-431, and to Clifton and Barrow (2007) for a discussion of models in which this assumption is not made.
- ii) On approach to a singularity (either initial or final, $\ell \rightarrow 0$) the 1-fluid is dominant (*i.e.* $\chi_1 \rightarrow 1$) and the matter content dominates the spatial curvature ($\Omega \rightarrow 1$, $\Omega_k \rightarrow 0$). In a typical ever-expanding model ($\ell \rightarrow \infty$) the *n*-fluid is dominant ($\chi_n \rightarrow 1$), and the matter content dominates the spatial curvature if and only if $\gamma_n < \frac{2}{3}$.
- iii) The *n*-fluid FL cosmologies admit $n - 1$ conserved quantities formed from the density parameters $\Omega_1, \dots, \Omega_n$. In particular, they admit one curvature quantity and $n - 2$ matter quantities.

We now comment on an important difference between n -fluid FL cosmologies with $n > 2$ and 2-fluid cosmologies. For a 2-fluid FL model, there are two essential parameters, a conformal parameter λ and a mass parameter m_1 , or equivalently, two observational parameters $\Omega_{1,0}$ and $\Omega_{2,0}$. There is also one conserved quantity \mathcal{K} that keeps track of the spatial curvature. For a 3-fluid FL model there is an additional mass parameter m_2 , and hence three observational parameters $\Omega_{1,0}$, $\Omega_{2,0}$ and $\Omega_{3,0}$. There is also a second conserved quantity \mathcal{M} . The new dynamical feature that arises in generalizing from a 2-fluid model to a 3-fluid model relates to the intermediate dynamics. In an ever-expanding 3-fluid model the additional degree of freedom is the maximum value attained by χ_2 , which indicates the extent to which fluid-2 becomes dominant during the evolution (recall that fluid-1 is dominant at early times ($\chi_1 \rightarrow 1$) and fluid-3 is dominant at late times ($\chi_3 \rightarrow 1$)). In an $RDC\Lambda$ -model of the physical universe, the maximum value of χ_{dust} is close to unity, which ensures that there is an extended matter-dominated epoch.

The dynamical systems analysis of the n -fluid FL universes leads to an interesting conclusion regarding the behaviour of the spatial curvature, as follows. *For any $\epsilon > 0$, there is an open set of n -fluid FL universes with $\gamma_n < \frac{2}{3}$ whose spatial curvature scalar Ω_k satisfies $|\Omega_k| < \epsilon$ throughout the evolution.* In other words, if $\gamma_n < \frac{2}{3}$, in particular, if fluid n is a cosmological constant, the total matter-energy content “controls” the spatial curvature throughout the evolution. This result is a consequence of the fact that generic ever-expanding models satisfy $\Omega_k \rightarrow 0$ as $\ell \rightarrow 0$ and since $\gamma_n < \frac{2}{3}$, $\Omega_k \rightarrow 0$ as $\ell \rightarrow +\infty$. It follows that $|\Omega_k|$ attains a maximum value ϵ which depends on the orbit in question. By restricting to orbits that lie within a sufficiently small neighbourhood of the flat FL invariant set $\Omega_k = 0$, the value ϵ can be made arbitrarily small. The above result has been found using different methods by Lake (2005), for $DC\Lambda$ FL universes. He indicates that the result will also hold for $RDC\Lambda$ universes (see also Lake (2006)). Lake comments on the significance of this result in connection with the so-called flatness problem (Lake (2005)).

One of the main results of this chapter is the analysis of the $RDC\Lambda$ FL universes and the comparisons with observations. Discussion of these models dates back to the 1930's. Our analysis is novel in that it gives new insight into the dynamics of these models, by representing the evolution as orbits in a compact state space. We now give a brief survey of previous work on these models.

Tolman (1934) gives the Friedmann equation for closed $RDC\Lambda$ universes (equation (106.3), p408) and states that de Sitter has studied the integration of this equation (de Sitter (1930), (1931), not readily available). He gives the special solutions of this DE that represent universes that are past or future asymptotic to the Einstein static universe, in the form $t = f(\ell)$ [equations (161.8) and (161.4)].

Coquereaux and Grossman (1982) give a detailed qualitative discussion of the Friedmann equation for closed $RDC\Lambda$ -universes using elliptic and Weierstrass functions. Debrowski and Stelmach (1986) extend the analysis to open $RDC\Lambda$ -universes. They give an explicit solution in terms of elementary functions when the parameters are suitably restricted.

The work that is closest to ours is that of Ehlers and Rinder (1989), who give a dynamical systems analysis of $RDC\Lambda$ -universes, using Hubble-normalized variables. They give the explicit expressions for the density parameters as functions of the scale factor [equation (19)] and also show the existence of two conserved quantities [equation (29)]. Their state space representation is inevitably incomplete since the Hubble-normalized variables are undefined at the instant of maximum expansion in recollapsing and bouncing models.

PERTURBATIONS OF FL MODELS

The high degree of isotropy of the cosmic microwave background (CMB), when combined with the Copernican Principle, provides strong support for the belief that the large scale structure of the observable universe is very well described by the Friedmann-Lemaître universes, which are exactly isotropic and spatially homogeneous. The real universe is, of course, not exactly isotropic and spatially homogeneous, since there is complex structure associated with the observed distribution of galaxies, the overall expansion may not be exactly isotropic, and there may be primordial gravitational waves. But the current belief of most cosmologists is that the deviations from an exact FL cosmological model are sufficiently small that they can be described by considering linear perturbations of the FL models.

4.1 Historical Development

In this section we briefly discuss the history of the theory of linear perturbations of FL cosmologies. There are two main approaches, which we shall refer to as the *metric approach* and the *geometrical approach*.

4.1.1 Metric Approach

The metric approach to perturbations of the FL models was introduced by Lifshitz (1946) and subsequently discussed in greater detail by Lifshitz and Khalatnikov (1963). In the metric approach, one distinguishes a *background spacetime*, which is a FL model, and a *perturbed spacetime*, which represents the physical universe. The metric for the perturbed spacetime is written in the form

$$g_{ab} = g_{(0)ab} + \delta g_{ab}, \quad (4.1)$$

where $g_{(0)ab}$ is the metric of the FL model and δg_{ab} is a small perturbation, *i.e.* $|\delta g_{ab}| \ll |g_{(0)ab}|$. The fluid 4-velocity and energy density in the perturbed spacetime are likewise written in the form

$$u^i = u_{(0)}^i + \delta u^i, \quad (4.2)$$

$$\mu = \mu_{(0)} + \delta \mu. \quad (4.3)$$

The density contrast is defined by

$$\delta = \frac{\delta \mu}{\mu_{(0)}}. \quad (4.4)$$

The problem of gauge

In defining a perturbation, one is effectively establishing a one-to-one correspondence between points in the physical spacetime and points in the background spacetime. However, this correspondence is not unique: one may make an infinitesimal change in the correspondence keeping the background coordinates fixed. Such a transformation is called a *gauge transformation*. A quantity that is unchanged by an infinitesimal gauge transformation is said to be *gauge-invariant*. We recommend Ellis and Bruni (1989) for a detailed explanation of gauge-invariance.

The problem of gauge invariance has plagued the study of linear perturbations since the pioneering work of Lifshitz (1946), resulting in authors publishing contradictory predictions on the behaviour of perturbations of FL cosmologies. The reason is that the metric perturbation δg_{ab} , the velocity perturbation δu^i and the density contrast δ are not gauge-invariant quantities, and hence their time evolution can depend on the choice of gauge.

A result due to Stewart and Walker (1974) gives a useful criteria for identifying gauge-invariant quantities:

Theorem 4.1.1 (Stewart-Walker Lemma) *Let $T_{(0)}$ be a tensor field on a background spacetime and let $T = T_{(0)} + \Delta T$ be the corresponding tensor on a perturbed spacetime. If $T_{(0)} = 0$ then T is gauge-invariant.*

In 1980 Bardeen reformulated the metric approach using gauge-invariant variables. He did not make use of the Stewart-Walker Lemma in choosing the gauge-invariant variables. Instead, he wrote out the transformation laws for the perturbations δg_{ab} , δu^a and δ under an infinitesimal gauge transformation

and then, by inspection, formed linear combinations of the perturbations and their derivatives that were invariant.

This approach placed the analysis of perturbations of Friedmann-Lemaître models on a sounder mathematical foundation and has been used in much subsequent research.¹ Nevertheless, the metric approach still possesses several shortcomings:

- i) First, it is not clear what physical quantities Bardeen's gauge-invariant quantities correspond to. Indeed, a gauge-invariant quantity can have different physical interpretations in different gauges. The most important ambiguity concerns the density contrast δ , as defined by (4.4). Bardeen defines two gauge-invariant quantities in terms of δ , denoted by ϵ_m and ϵ_g . The first equals the density contrast δ when one uses the so-called co-moving gauge, while the second equals δ in the so-called Newtonian gauge (see Bardeen 1980, p22).
- ii) Second, as emphasized by Hawking (1966), the metric tensor is not a physically significant quantity since one cannot measure it directly.

4.1.2 Geometrical Approach

The geometrical approach has its origins in a paper by Hawking (1966). Instead of using the components of a perturbed metric as basic variables, he proposed using the evolution equations for the kinematic quantities, the Weyl curvature and the matter density to study how perturbations evolve. Although this paper initiated a significant new approach to studying linear perturbations of FL, the analysis of density perturbations was flawed and the paper had little impact. Subsequently, motivated by Hawking's paper, Lyth and Mukherjee (1988) used the evolution equations for μ and H to study density perturbations in an FL cosmology, although they did not do so in a fully gauge-invariant way. The decisive step was taken by Ellis & Bruni (1989), when they proposed using the spatial gradient of the matter density ($\hat{\nabla}_a \mu$) as the basic variable to describe density perturbations. Unlike the density contrast δ , the spatial gradient is a gauge-invariant quantity by the Stewart-Walker lemma, since it is zero in any FL model (see proposition 2.1.4). The first comprehensive discussion of the geometrical approach was given by Bruni, Dunsby and

¹We refer to Mukhanov (2005) as a recent text that uses a simplified version of the Bardeen approach, restricting considerations to perturbations of flat FL (see chapter 7).

Ellis (1992)². They also related the Bardeen approach to the geometrical approach, and gave clear physical interpretations of his gauge-invariant variables.

In this chapter we give an introduction to the geometrical approach to linear perturbations of FL models.

4.2 The Linearized Einstein Field Equations

The geometrical approach is based on the conviction that one should use variables that have a direct physical or geometric meaning, and are both coordinate-independent and gauge-invariant. One thus uses tensorial quantities that are identically zero in an FL universe, and hence gauge-invariant, on account of the Stewart-Walker lemma.

We shall see that a complete set of tensorial quantities is the following:

- i) The density and Hubble gradients: $\hat{\nabla}_a \mu, \hat{\nabla}_a H$
- ii) The fluid kinematic quantities: σ_{ab}, ω_a
- iii) The Weyl curvature: E_{ab}, H_{ab}

These variables have two additional advantages:

- a) they directly describe the deviation of the physical universe from an idealized FL universe, and
- b) they are, in principle, observable, when normalized to be dimensionless using a power of H . We refer in particular to the fundamental paper of Sachs and Wolfe (1967), who analyzed the observations of distant galaxies from this point of view, and Maartens et al. (1995b), who used the cosmic microwave background observations to bound these quantities.

In the geometrical approach one thus replaces the EFEs for the metric tensor components by the equivalent system of evolution equations and constraints, satisfied by the tensorial quantities i)-iii), as given in Appendix A.

For simplicity we will assume the matter content consists of a perfect fluid with barotropic equation of state $p = p(\mu)$ and a cosmological constant Λ . We will use the standard notation,

$$w = \frac{p}{\mu}, \quad c_s^2 = \frac{dp}{d\mu}, \quad (4.5)$$

²The name ‘‘geometrical approach’’ was introduced in this reference. This approach is also referred to as ‘‘the gauge-invariant and covariant (GIC) approach’’.

where c_s is the speed of sound.

4.2.1 The Linearization Process

In order to derive the governing equations for linear perturbations of a FL cosmology, we must distinguish between quantities which are non-zero in the background (referred to as *zero-order quantities*) and those which are zero in the background (referred to as *first-order quantities*). For a FL background, the zero-order quantities are

$$\mu, \quad w, \quad c_s^2 \quad \text{and} \quad H. \quad (4.6)$$

It follows from propositions 2.1.1-2.1.4 that the quantities

$$\hat{\nabla}_a \mu, \quad \hat{\nabla}_a H, \quad \sigma_{ab}, \quad \omega_a, \quad E_{ab}, \quad H_{ab}, \quad (4.7)$$

are first order. For sake of brevity, we define

$$X_a = \hat{\nabla}_a \mu, \quad (4.8)$$

and

$$Z_a = 3\hat{\nabla}_a H. \quad (4.9)$$

We note that the acceleration \dot{u}_a is not an independent quantity, since it can be expressed in terms of the density gradient X_a via (A.17), on using (4.5) and (4.8):

$$\dot{u}_a = -\frac{c_s^2}{(1+w)\mu} X_a. \quad (4.10)$$

The evolution equations for the zero-order quantities are simply the evolution equations for a FL cosmology with a single fluid and cosmological constant, given by (3.5) and (3.6), with μ replaced by $\mu + \Lambda$ and p by $p - \Lambda$ to make the cosmological constant explicit. We also include the evolution equation for w , which follows from (4.12) and (4.5).

Zero-Order Evolution Equations

$$\dot{H} = -H^2 - \frac{1}{6}(1+3w)\mu + \frac{1}{3}\Lambda, \quad (4.11)$$

$$\dot{\mu} = -3H(1+w)\mu, \quad (4.12)$$

$$\dot{w} = -3H(1+w)(c_s^2 - w). \quad (4.13)$$

Note: The Friedmann equation (3.8) plays an auxiliary role in that if $K \neq 0$, it serves to express $\ell(t)$ in terms of the zero-order quantities. We use (3.11) to write (3.8) as follows:

$$H^2 = \frac{1}{3}(\mu + \Lambda) - \frac{K}{\ell^2}. \quad (4.14)$$

The evolution equations for the first order quantities are obtained from the general system of evolution equations and constraints in Appendix A, by a process of linearization. There are three aspects to the *linearization procedure*:

- L_1) Drop all products of the first order quantities (4.7) and their spatial derivatives in the general non-linear evolution equations.
- L_2) The resulting equations form a coupled system of linear PDEs for the first order quantities, whose coefficients depend on the zero order quantities. For example, there are terms such as $H X_a$, $H E_{ab}$ and $\mu \sigma_{ab}$. The linearization process involves replacing the zeroth order quantity by its value in the background FL model in each of these products.
- L_3) The resulting equations also contain the projected spatial covariant derivative operator $\hat{\nabla}_a$ acting on first order quantities. The linearization process involves replacing $\hat{\nabla}_a$, which is defined on the physical spacetime, by the corresponding operator on the background FL spacetime.

4.2.2 The Linearized Evolution and Constraint Equations

We now derive the governing equations for linearized perturbations by applying the linearization procedure L_1 to the general system of evolution equations and constraint equations in Appendix A.

One aspect of the linearization process, namely dealing with the terms $\hat{\nabla}_{\langle a} \dot{u}_{b \rangle}$, $\text{curl}(\dot{u}_a)$ and $\hat{\nabla}_a(\hat{\nabla}^b \dot{u}_b)$ involving the spatial derivative of \dot{u}_a , requires particular attention. First, it follows from (4.10) and L_1 that³

$$\hat{\nabla}_{\langle a} \dot{u}_{b \rangle} = \frac{-c_s^2}{(1+w)\mu} \hat{\nabla}_{\langle a} X_{b \rangle}. \quad (4.15)$$

³Since $p = p(\mu)$, we can use (4.5) to write

$$\hat{\nabla}_a \frac{c_s^2}{(1+w)\mu} = F(\mu) \hat{\nabla}_a \mu = F(\mu) X_a,$$

where $F(\mu)$ is a function of μ whose specific form is unimportant. Then the term $F(\mu) X_a X_b$ is dropped.

Second, it follows in a similar manner from (4.10), (4.8) and L_1 that

$$\text{curl}(\dot{u}_a) = \frac{-c_s^2}{(1+w)\mu} \text{curl}(\hat{\nabla}_a \mu). \quad (4.16)$$

Applying (E.25) with $f = \mu$ and using (A.16) gives

$$\text{curl}(\dot{u}_a) = 6c_s^2 H \omega_a. \quad (4.17)$$

Finally, it again follows from (4.10) and repeated application of L_1 that

$$\hat{\nabla}_a(\hat{\nabla}^b \dot{u}_b) = \frac{-c_s^2}{(1+w)\mu} \hat{\nabla}_a(\hat{\nabla}^2 \mu). \quad (4.18)$$

Then the commutation property (E.18) with $f = \mu$ and the evolution equation (A.16) for μ leads to

$$\hat{\nabla}_a(\hat{\nabla}^b \dot{u}_b) = \frac{-c_s^2}{(1+w)\mu} \left(\hat{\nabla}^2 - \frac{2K}{\ell^2} \right) X_a - 6c_s^2 H \text{curl}(\omega_a). \quad (4.19)$$

The evolution equations for the first order quantities are now obtained from (A.9) (A.10), (A.18), (A.19), (A.25) and (A.26) on applying the linearization procedure L_1 and using (4.15), (4.17) and (4.19) to eliminate terms involving spatial derivatives of \dot{u}_a .

First-Order Evolution Equations

$$\dot{X}_{\langle a} = -4H X_a - (1+w)\mu Z_a, \quad (4.20)$$

$$\dot{Z}_{\langle a} = -3H Z_a - \frac{1}{2} X_a - \frac{c_s^2}{(1+w)\mu} \left(\hat{\nabla}^2 + \frac{K}{\ell^2} \right) X_a - 6c_s^2 H \text{curl}(\omega_a), \quad (4.21)$$

$$\dot{\sigma}_{\langle ab} = -2H \sigma_{ab} - \frac{c_s^2}{(1+w)\mu} \hat{\nabla}_{\langle a} X_{b\rangle} - E_{ab}, \quad (4.22)$$

$$\dot{\omega}_{\langle a} = (3c_s^2 - 2)H \omega_a, \quad (4.23)$$

$$\dot{E}_{\langle ab} = -3H E_{ab} + \text{curl}(H_{ab}) - \frac{1}{2}(1+w)\mu \sigma_{ab}, \quad (4.24)$$

$$\dot{H}_{\langle ab} = -3H H_{ab} - \text{curl}(E_{ab}). \quad (4.25)$$

Similarly, constraints on the first-order quantities are obtained on applying the linearization procedure L_1 to (A.11), (A.12), (A.13), (A.20) and (A.21).

First-Order Constraint Equations

$$\hat{\nabla}^b \sigma_{ab} = \frac{2}{3} Z_a + \text{curl}(\omega_a), \quad (4.26)$$

$$\hat{\nabla}^a \omega_a = 0, \quad (4.27)$$

$$\hat{\nabla}_{\langle a} \omega_{b \rangle} = H_{ab} - \text{curl}(\sigma_{ab}), \quad (4.28)$$

$$\hat{\nabla}^b E_{ab} = \frac{1}{3} X_a, \quad (4.29)$$

$$\hat{\nabla}^b H_{ab} = (1 + w) \mu \omega_a. \quad (4.30)$$

There are two additional constraints involving the curls of the spatial gradients X_a and Z_a . Indeed, using (4.10), (4.16) and (4.17) leads to

$$\text{curl}(X_a) = 6(1 + w) \mu H \omega_a. \quad (4.31)$$

This equation is in fact a consequence of the main system of evolution equations and constraints and hence we list it as an auxiliary equation. There is an analogous equation for $\text{curl}(Z_a)$ which we shall not need.

Comments:

- i) The evolution equations (4.20)-(4.25) and constraints (4.26)-(4.30) form a closed system of linear first order PDEs for the first order quantities (4.7). The coefficients μ , w , c_s^2 and H are zero-order quantities and are evaluated in the background FL model. Note that if $K \neq 0$ the term K/ℓ^2 in (4.21) is given in terms of zero order quantities by the constraint (4.14).
- ii) It can be shown that the system of evolution equations and constraints is consistent: that is, the time derivatives of each constraint equation (4.26)-(4.30) are identically satisfied as a consequence of the other equations. As a result, once a set of initial data is chosen to satisfy the constraint equations, these equations will hold at all later times (see Bruni et al. (1992), p42).
- iii) Equations (4.20)-(4.25) and (4.26)-(4.30) form the basis for the geometrical approach to the analysis of linear perturbations of FL models with a barotropic perfect fluid and cosmological constant as source. They can be extracted from Bruni et al. (1992) by specializing their more general equations

to the case of a barotropic perfect fluid (set q_a , π_{ab} and \mathcal{E} to zero in equations (50)-(54), (57)-(62)). Most authors, however, do not make use of the full system of evolution and constraint equations.

4.3 Harmonic Decomposition

Before proceeding, we must discuss the fundamental role of spatial harmonics in the geometrical theory. The harmonic decomposition is analogous to a Fourier decomposition of a quantity that is defined along the background 3-spaces. Hence, harmonics are defined so as to be constant along flow lines (*i.e.*, independent of proper time) and orthogonal to the fluid 4-velocity \mathbf{u} . We refer the reader to Appendix F for a complete discussion of spatial harmonics in the geometrical approach.

The key idea behind the harmonic decomposition is that a scalar, vector or tracefree symmetric rank 2 tensor orthogonal to u^a can be written as a linear combination of scalar, vector and tensor harmonics with purely time-dependent coefficients. In the literature it is tacitly assumed that the harmonics are complete in some appropriate space of tensor fields.

The expansions are as follows:

$$T = \sum_k T_{(0,k)} Q^{(0)}, \quad (4.32)$$

$$T_a = \sum_k T_{(0,k)} Q_a^{(0)} + \sum_k T_{(1,k)} Q_a^{(1)}, \quad (4.33)$$

$$T_{ab} = \sum_k T_{(0,k)} Q_{ab}^{(0)} + \sum_k T_{(1,k)} Q_{ab}^{(1)} + \sum_k T_{(2,k)} Q_{ab}^{(2)}. \quad (4.34)$$

This decomposition has two consequences: Firstly, it decouples the temporal and spatial dependence of each field, since all coefficients are purely functions of time and harmonics are constant along flow lines, *i.e.* $T_{(d,k)} = T_{(d,k)}(t)$ and $\dot{Q} = 0$ for all harmonics Q (see Appendix F). Second, in the framework of linear perturbation theory, it removes any co-dependence between scalar, vector and tensor harmonics. As in Bruni et al. (1992, p51), the summation in (4.32)-(4.34) can be over a discrete set or an integral over a continuously varying index.

We note that the scalar decomposition (4.32) is analogous to the 3-dimensional Fourier transform for scalar fields. The vector decomposition (4.33) is similarly analogous to the Fourier transform, except applied to the curl-free (scalar) component and the divergence-free (vector) component of the vector field. The tensor decomposition (4.34) separates the curl-free divergent (scalar) component, the remaining divergent (vector) component and the divergence-free (tensor) component of the vector field.

Using this decomposition, it is natural to consider the kinematic quantities and Weyl tensor components of the real universe to be the fundamental description of the linear perturbations (since they are each gauge-invariant, each of these components can be expanded using the harmonic basis functions).

The decoupling of the perturbation types is extremely important in the linearized theory since it allows us to consider scalar, vector and tensor perturbations independently. For each perturbation type, we will proceed as follows: First, we first define a *basic variable*. Second, we use the linearized evolution equations (4.11)-(4.30) to derive a second order differential equation which describes the evolution of the basic variable, which we shall call the *governing DE*. Finally, we use the linearized evolution and constraint equations to express all kinematic quantities and Weyl tensor components in terms of the basic variable, hence providing a complete solution to the linearized equations for each type of perturbation.

4.4 Scalar Perturbations

We first focus on *scalar perturbations*, which are defined as solutions of the linearized equations (4.11)-(4.30) that can be expanded in terms of the scalar harmonics $Q^{(0)}$, $Q_a^{(0)}$ and $Q_{ab}^{(0)}$, defined by equations (F.3), (F.5) and (F.6). Following Ellis et al. (1989), we choose the basic quantity for scalar perturbations to be the fractional density gradient \mathcal{D}_a , defined as

$$\mathcal{D}_a = \frac{\ell \hat{\nabla}_a \mu}{\mu}, \quad (4.35)$$

or equivalently, using (4.8),

$$\mathcal{D}_a = \frac{\ell}{\mu} X_a. \quad (4.36)$$

We also rescale Z_a according to $\mathcal{Z}_a = \ell Z_a$, *i.e.*

$$\mathcal{Z}_a = 3\ell \hat{\nabla}_a H. \quad (4.37)$$

4.4.1 The Governing DE

In this section we derive the governing DE for scalar perturbations, following Ellis et al. (1990) (also see Wainwright and Ellis (1997, p290-3)). Since we are considering scalar harmonics, we can write

$$\mathcal{D}_a = \sum_k \mathcal{D}_{(k)} Q_a^{(0,k)}. \quad (4.38)$$

Since (4.38) and (F.8) imply $\text{curl}(\mathcal{D}_a) = 0$, and hence $\text{curl}(X_a) = 0$, (4.31) gives

$$\omega_a = 0. \quad (4.39)$$

It follows that equations (4.20) and (4.21) decouple from the remaining linearized evolution equations. Expressing these equations in terms of the rescaled variables \mathcal{D}_a and \mathcal{Z}_a leads to

$$\dot{\mathcal{D}}_{\langle a} = 3wH\mathcal{D}_a - (1+w)\mu\mathcal{Z}_a, \quad (4.40)$$

$$\dot{\mathcal{Z}}_{\langle a} = -2H\mathcal{Z}_a - \frac{1}{2}\mu\mathcal{D}_a - \frac{c_s^2}{(1+w)\mu} \left(\hat{\nabla}^2 + \frac{K}{\ell^2} \right) \mathcal{D}_a. \quad (4.41)$$

These equations can be combined to give a second order evolution equation for \mathcal{D}_a . On differentiating (4.40) and using (4.11)-(4.13), (4.14) and (4.41), we obtain

$$\ddot{\mathcal{D}}_{\langle a} + \mathcal{A}H\dot{\mathcal{D}}_{\langle a} + \mathcal{B}H^2\mathcal{D}_a - c_s^2 \left(\hat{\nabla}^2 - \frac{2K}{\ell^2} \right) \mathcal{D}_a = 0, \quad (4.42)$$

where

$$\mathcal{A} = 2 - 3w - 3(w - c_s^2), \quad (4.43)$$

and

$$\mathcal{B} = -\frac{3}{2}(1-w)(1+3w)\Omega - 6w\Omega_\Lambda + 3(w - c_s^2) \left(\frac{K}{H^2\ell^2} - 3 \right). \quad (4.44)$$

We now expand each term in terms of scalar harmonics. The derivatives of \mathcal{D}_a along the fundamental congruence are obtained by differentiating (4.38) and noting $\dot{Q}_a^{(0,k)} = 0$ (see (F.10)), as follows:

$$\dot{\mathcal{D}}_{\langle a} = \sum_k \dot{\mathcal{D}}_{\langle k} Q_a^{(0,k)}, \quad \ddot{\mathcal{D}}_{\langle a} = \sum_k \ddot{\mathcal{D}}_{\langle k} Q_a^{(0,k)}. \quad (4.45)$$

In order to remove the Laplacian term from the evolution equation (4.42), we use the identity (F.13), which can be written as

$$\left(\hat{\nabla}^2 - \frac{2K}{\ell^2} \right) Q_a^{(0)} = -\frac{k^2}{\ell^2} Q_a^{(0)}. \quad (4.46)$$

On applying (4.45) and (4.46) to (4.42), we obtain

$$\ddot{\mathcal{D}}_{\langle k} + \mathcal{A}H\dot{\mathcal{D}}_{\langle k} + \left(\mathcal{B} + \frac{c_s^2 k^2}{H^2 \ell^2} \right) H^2 \mathcal{D}_{\langle k} = 0, \quad (4.47)$$

where \mathcal{A} and \mathcal{B} are given by (4.43) and (4.44). This is the governing equation for scalar perturbations, in terms of clock time t . This equation is equivalent⁴ to (73) in Bruni et al. (1992), upon specializing to the case of a perfect fluid, and (214) in Ellis and van Elst (1998), in the case of $\Lambda = 0$.

4.4.2 Kinematic Quantities and Weyl Curvature

We now write the kinematic quantities and Weyl curvature in terms of the density gradient coefficients $\mathcal{D}_{(k)}$ and the scalar harmonics $Q^{(0)}$. Since we are considering scalar harmonics, \mathcal{Z}_a and σ_{ab} can be expanded as

$$\mathcal{Z}_a = \sum_k \mathcal{Z}_{(k)} Q_a^{(0,k)}, \quad \sigma_{ab} = \sum_k \sigma_{(k)} Q_{ab}^{(0,k)}. \quad (4.48)$$

We can use (4.40) to express $\mathcal{Z}_{(k)}$ in terms of $\mathcal{D}_{(k)}$ and $\dot{\mathcal{D}}_{(k)}$, and since $\omega_a = 0$ it follows from (4.26) and (F.16) that

$$\mathcal{Z}_{(k)} = \frac{(k^2 - 3K)}{k} \sigma_{(k)}. \quad (4.49)$$

Next, it follows from (4.48) and (F.8) that $\text{curl}(\sigma_{ab}) = 0$. Hence, using (4.28) we have that

$$H_{ab} = 0. \quad (4.50)$$

Expanding E_{ab} in terms of scalar harmonics via

$$E_{ab} = \sum_k E_{(k)} Q_{ab}^{(0,k)}, \quad (4.51)$$

and using (F.16) leads to

$$\ell \hat{\nabla}^b E_{ab} = \sum_k \frac{2}{3k} (k^2 - 3K) E_{(k)} Q_a^{(0,k)}. \quad (4.52)$$

The coefficients $E_{(k)}$ can then be expressed in terms of $\mathcal{D}_{(k)}$ via the constraint (4.29). We thus obtain the following expressions for the harmonic coefficients of the non-zero Hubble-normalized first-order quantities:

⁴Bruni et al. (1992) make use of the scalar quantity $\Delta = \hat{\nabla}^a \mathcal{D}_a$ instead of $\mathcal{D}_{(k)}$, defined in this manner so as to restrict considerations to scalar perturbations without it being necessary to resort to a harmonic expansion. This choice of scalar quantity automatically removes all vector harmonic components from \mathcal{D}_a since vector harmonics are divergence-free by definition (F.22).

$$\mathcal{Z}_{(k)} = -\frac{1}{(1+w)} \left[-3wH\mathcal{D}_{(k)} + \dot{\mathcal{D}}_{(k)} \right], \quad (4.53)$$

$$\sigma_{(k)} = -\frac{k}{k^2 - 3K} \mathcal{Z}_{(k)}, \quad (4.54)$$

$$E_{(k)} = \frac{k}{2(k^2 - 3K)} \mu \mathcal{D}_{(k)}. \quad (4.55)$$

At this stage, we have satisfied all of the linearized evolution equations and constraint equations except for (4.22) and (4.24). We can now verify that D_a , σ_{ab} and E_{ab} , as given by (4.38), (4.48), (4.51) and (4.53)-(4.55) satisfy (4.22) and (4.24) (via Maple). We have thus established the following result:

Proposition 4.4.1 *For any solution $\mathcal{D}_{(k)}$ of the governing DE (4.47), the first order quantities given by (4.38), (4.48), (4.51) and (4.53)-(4.55) satisfy the complete set of linearized evolution and constraint equations (4.20)-(4.30).*

4.5 Vector Perturbations

In this section we focus on *vector perturbations*, defined as solutions of the linearized equations (4.20)-(4.30) that can be expanded in terms of the vector harmonics $Q_a^{(1)}$ and $Q_{ab}^{(1)}$. Vector perturbations are closely associated with vorticity, and so we choose the basic quantity for vector perturbations to be the co-moving vorticity vector, defined by

$$\mathcal{W}_a = \ell \omega_a. \quad (4.56)$$

4.5.1 The Governing DE

In this section, we derive the governing DE for vector perturbations. On differentiating (4.56) along the fundamental congruence and using (4.23), we obtain

$$\dot{\mathcal{W}}_{\langle a} - (3c_s^2 - 1)H\mathcal{W}_a = 0. \quad (4.57)$$

We expand (4.57) in terms of the associated vector harmonics $P_a^{(1,k)}$ as

$$\mathcal{W}_a = \sum_k \mathcal{W}_{(k)} P_a^{(1,k)}. \quad (4.58)$$

On substituting (4.58) into (4.57), we obtain

$$\dot{\mathcal{W}}_{(k)} - (3c_s^2 - 1)H\mathcal{W}_{(k)} = 0. \quad (4.59)$$

This equation is the governing DE for vector perturbations, in terms of clock time t .

4.5.2 Kinematic Quantities and Weyl Curvature

We now write the kinematic quantities and Weyl curvature in terms of the co-moving vorticity coefficients $\mathcal{W}_{(k)}$ and vector harmonics $Q^{(1)}$. By the definition of vector perturbations, all quantities can be expanded in terms of vector harmonics and associated vector harmonics, as follows:

$$\begin{aligned} \mathcal{D}_a &= \sum_k \mathcal{D}_{(k)} Q_a^{(1,k)}, & \mathcal{Z}_a &= \sum_k \mathcal{Z}_{(k)} Q_a^{(1,k)}, & \sigma_{ab} &= \sum_k \sigma_{(k)} Q_a^{(1,k)}, \\ E_{ab} &= \sum_k E_{(k)} Q_{ab}^{(1,k)}, & H_{ab} &= \sum_k H_{(k)} P_{ab}^{(1,k)}. \end{aligned} \quad (4.60)$$

From (4.31), (4.36) and (F.23), we can express $\mathcal{D}_{(k)}$ in terms of $\mathcal{W}_{(k)}$, and hence obtain $\mathcal{Z}_{(k)}$ in terms of $\mathcal{W}_{(k)}$ from (4.20), (4.36) and (4.37). Further, using (F.34), the divergences of σ_{ab} , E_{ab} and H_{ab} can be written as follows:

$$\ell \hat{\nabla}^b \sigma_{ab} = \sum_k \frac{1}{2k} (k^2 - 2K) \sigma_{(k)} Q_a, \quad (4.61)$$

$$\ell \hat{\nabla}^b E_{ab} = \sum_k \frac{1}{2k} (k^2 - 2K) E_{(k)} Q_a, \quad (4.62)$$

$$\ell \hat{\nabla}^b H_{ab} = \sum_k \frac{1}{2k} (k^2 - 2K) H_{(k)} Q_a. \quad (4.63)$$

We can now use (4.26), (4.29) and (4.30) to determine the coefficients $\sigma_{(k)}$, $E_{(k)}$ and $H_{(k)}$ in terms of $\mathcal{W}_{(k)}$. We thus obtain the following expressions for the harmonic coefficients of the first-order quantities:

$$\mathcal{D}_{(k)} = 6(1+w)H\ell\mathcal{W}_{(k)}, \quad (4.64)$$

$$\mathcal{Z}_{(k)} = -3\ell^{-1} [2K - (1+w)\mu\ell^2] \mathcal{W}_{(k)}, \quad (4.65)$$

$$\sigma_{(k)} = \frac{2k}{k^2 - 2K} \ell^{-1} [(k^2 - 2K) + 2(1+w)\mu_0\ell^2] \mathcal{W}_{(k)}, \quad (4.66)$$

$$E_{(k)} = \frac{4k}{k^2 - 2K} (1+w)\mu H\ell\mathcal{W}_{(k)}, \quad (4.67)$$

$$H_{(k)} = \frac{2k}{k^2 - 2K} (1+w)\mu\mathcal{W}_{(k)}. \quad (4.68)$$

At this stage, we have satisfied all of the linearized evolution equations and constraint equations except for (4.22), (4.24), (4.25) and (4.28). We can now verify that \mathcal{D}_a , \mathcal{Z}_a , σ_{ab} , ω_a , E_{ab} and H_{ab} , as given by (4.60) and (4.64)-(4.68) satisfy (4.22), (4.24), (4.25) and (4.28) (via Maple). We have thus established the following result:

Proposition 4.5.1 *For any solution $\mathcal{W}_{(k)}$ of the governing DE (4.59), the first order quantities given by (4.60) and (4.64)-(4.68) satisfy the complete set of linearized evolution and constraint equations (4.20)-(4.30).*

4.6 Tensor Perturbations

In this section we focus on *tensor perturbations*, defined as solutions of the linearized equations (4.11)-(4.30) that can be expanded in terms of the tensor harmonics $Q_{ab}^{(2)}$. Tensor perturbations are interpreted as gravitational waves, described by a coupling of the electric and magnetic Weyl curvature.

As regards to the choice of basic variable, we shall see that for tensor perturbations the only non-zero first order quantities are σ_{ab} , E_{ab} and H_{ab} . On the grounds of mathematical simplicity we choose the comoving shear tensor as the basic variable, denoted \mathcal{X}_{ab} and defined by

$$\mathcal{X}_{ab} = \ell\sigma_{ab}. \quad (4.69)$$

4.6.1 The Governing DE

In this section, we derive the governing DE for tensor perturbations.⁵ It is a consequence of applying tensor harmonics that rank two tensors have zero spatial divergence, *i.e.*

$$\hat{\nabla}^b \sigma_{ab} = 0, \quad \hat{\nabla}^b E_{ab} = 0, \quad \hat{\nabla}^b H_{ab} = 0. \quad (4.70)$$

It then follows from the constraints (4.26), (4.29) and (4.30) that all vector quantities are zero, *i.e.*

$$\mathcal{D}_a = 0, \quad \mathcal{Z}_a = 0, \quad \text{and} \quad \omega_a = 0. \quad (4.71)$$

Using (4.71), the shear evolution equation (4.22) can be rewritten in terms of \mathcal{X}_{ab} as

$$\dot{\mathcal{X}}_{(ab)} = -H\mathcal{X}_{ab} - \ell E_{ab}. \quad (4.72)$$

⁵A similar derivation was given by Dunsby et al. (1997).

We now use (4.24) to calculate $(\ell E_{ab})'_{\perp}$, noting that $\dot{\ell} = H\ell$. We use the constraint (4.28) to eliminate H_{ab} and then apply (E.31) to the ‘‘curl curl \mathcal{X}_{ab} ’’ term. The result is

$$(\ell E_{ab})'_{\perp} = -2H(\ell E_{ab}) + \left(-\frac{1}{2}(1+w)\mu + \frac{3K}{\ell^2} \right) \mathcal{X}_{ab} - \hat{\nabla}^2 \mathcal{X}_{ab}. \quad (4.73)$$

We can now combine these equations to give a second order DE for \mathcal{X}_{ab} (differentiate (4.72) using (4.11) and (4.73), and then use (4.72) to eliminate E_{ab} from the resulting equation). After expressing the coefficients of \mathcal{X}_{ab} in terms of the density parameters Ω and Ω_{Λ} (see 3.31), we obtain⁶

$$\ddot{\mathcal{X}}_{(ab)} + 3H\dot{\mathcal{X}}_{(ab)} + \left[-(3w+1)\Omega + 2\Omega_{\Lambda} + \frac{2K}{H^2\ell^2} \right] H^2 \mathcal{X}_{ab} - \hat{\nabla}^2 \mathcal{X}_{ab} = 0. \quad (4.74)$$

We now expand \mathcal{X}_{ab} in terms of tensor harmonics according to

$$\mathcal{X}_{ab} = \sum_k \mathcal{X}_{(k)} Q_{ab}^{(2,k)}. \quad (4.75)$$

Substituting (4.75) into (4.74) and using (F.38) yields

$$\ddot{\mathcal{X}}_{(k)} + 3H\dot{\mathcal{X}}_{(k)} + \left[-(3w+1)\Omega + 2\Omega_{\Lambda} + \frac{k^2 + 2K}{H^2\ell^2} \right] H^2 \mathcal{X}_{(k)} = 0. \quad (4.76)$$

This is the governing equation for tensor perturbations, in terms of clock time t .

4.6.2 Kinematic Quantities and Weyl Curvature

We now write the kinematic quantities and Weyl curvature in terms of the dimensionless shear coefficients $\mathcal{X}_{(k)}$, tensor harmonics $Q_{ab}^{(2)}$ and associated tensor harmonics $P_{ab}^{(2)}$. Since (4.70) confirms that σ_{ab} , E_{ab} and H_{ab} can be expanded in terms of tensor harmonics, we can write

$$\sigma_{ab} = \sum_k \sigma_{(k)} Q_{ab}^{(2,k)}, \quad E_{ab} = \sum_k E_{(k)} Q_{ab}^{(2,k)}, \quad H_{ab} = \sum_k H_{(k)} P_{ab}^{(2,k)}, \quad (4.77)$$

⁶It should be noted that the evolution equation (22) given in Dunsby et al. (1997, p1219) contains two errors. The original equation should instead be written as

$$\Delta\sigma_{ab} + \frac{5}{3}\Theta\dot{\sigma}_{ab} + \left[\frac{1}{9}\Theta^2 - \frac{1}{6}\mu(9\gamma - 10) \right] \sigma_{ab} = 0,$$

where $\Delta\sigma_{ab} = \ddot{\sigma}_{\perp ab} - \hat{\nabla}^2\sigma_{ab}$, so as to be consistent with the remainder of that paper. This equation is then equivalent to (4.74) for $\Lambda = 0$ on recalling that $\mathcal{X}_{ab} = \ell\sigma_{ab}$ and $\Theta = 3H$.

where ⁷

$$P_{ab} = \ell \text{curl}(Q_{ab}). \quad (4.78)$$

Then using (4.69), (4.22) and (4.28) we can write σ_{ab} , E_{ab} and H_{ab} in terms of $\mathcal{X}_{(k)}$. We thus obtain the following expressions for the harmonic coefficients of the non-zero first-order quantities:

$$\sigma_{(k)} = \ell^{-1} \mathcal{X}_{(k)}, \quad (4.79)$$

$$E_{(k)} = -\ell^{-1} \left(\dot{\mathcal{X}}_{(k)} + H \mathcal{X}_{(k)} \right), \quad (4.80)$$

$$H_{(k)} = \ell^{-2} \mathcal{X}_{(k)}. \quad (4.81)$$

At this stage, we have satisfied all of the linearized evolution equations and constraint equations except for (4.24) and (4.25). We can now verify that σ_{ab} , E_{ab} and H_{ab} , as given by (4.77) satisfy (4.24) and (4.25) (via Maple). We have thus established the following result:

Proposition 4.6.1 *For any solution $\mathcal{X}_{(k)}$ of the governing DE (4.76), the first order quantities given by (4.77) and (4.79)-(4.81) satisfy the complete set of linearized evolution and constraint equations (4.20)-(4.30).*

4.7 Dynamics of the Linear Perturbations: General Features

In this section we discuss the dynamics of the linear perturbations. We first discuss the importance of conformal time in analyzing perturbations of FL and rewrite the governing DEs for each perturbation type in terms of this time variable. Finally, we examine the dependence of the first-order quantities on the solutions to the governing DEs.

4.7.1 Conformal Time and the Particle Horizon

It is a consequence of propositions 4.4.1, 4.5.1 and 4.6.1 that the spatial gradients, kinematic quantities and Weyl tensor components can be determined directly from the set of solutions of the governing DEs. As a consequence, we turn our attention to the behaviour of solutions to these equations.

⁷See definition in section F.3.

We make the choice of conformal time η as the independent quantity, defined in terms of the length scale ℓ and clock time t by

$$\frac{d\eta}{dt} = \frac{1}{\ell}. \quad (4.82)$$

Under this choice of time variable, the Robertson-Walker line element (1.3) can be written in the form

$$ds^2 = \ell^2(\eta)(-d\eta^2 + \tilde{g}_{\alpha\beta}dx^\alpha dx^\beta), \quad (4.83)$$

where $\tilde{g}_{\alpha\beta}$ is a 3-metric of constant curvature.

Conformal time is directly linked to the notion of the *particle horizon*, defined as the boundary of the part of the universe that is visible to us. In terms of conformal time, the distance to the particle horizon (see figure 4.1) is given by

$$d_H = \ell(\eta)\eta. \quad (4.84)$$

Since the wavelength λ for a perturbation of mode k at some fixed time η is given by

$$\lambda(\eta) = \left(\frac{2\pi}{k}\right)\ell(\eta), \quad (4.85)$$

then from (4.84) we obtain the key relation

$$\frac{\lambda}{d_H} = \frac{2\pi}{k\eta}. \quad (4.86)$$

Hence, if $k\eta \ll 1$ the wavelength of the perturbation mode is large compared to the distance to the particle horizon (since $\lambda \gg d_H$). In this situation it is customary to say that the *perturbation is outside the horizon*. Conversely, if $k\eta \gg 1$ the wavelength of the perturbation mode is small compared to the distance to the horizon (since $\lambda \ll d_H$). Similarly, in this situation it is customary to say that the *perturbation is inside the horizon*.

The range of the conformal time variable varies depending on the characteristics of the FL model, as shown in Appendix D. A summary of the behaviour of conformal time and clock time for each of the generic classes of FL model with an initial singularity is given below:

<u>FL Model</u>	<u>Range of η</u>	<u>Range of t</u>
Ever-expanding models, $\Lambda = 0$	$0 < \eta < +\infty$	$0 < t < +\infty$
Ever-expanding models, $\Lambda \neq 0$	$0 < \eta < \eta_f$	$0 < t < +\infty$
Recollapsing models, Λ arbitrary	$0 < \eta < \eta_f$	$0 < t < t_f$




FIGURE 4.1: The relation between the distance to the particle horizon, denoted $d_H(\eta)$ and the conformal time parameter η .

4.7.2 Governing DEs Using Conformal Time

We now present the governing DEs for scalar, vector and tensor perturbations in terms of conformal time and specialized to a γ -law equation of state, *i.e.* $p = (\gamma - 1)\mu$. It follows from (4.5) that

$$w = c_s^2 = \gamma - 1. \quad (4.87)$$

The three governing DEs, given by (4.47), (4.59) and (4.76) in terms of clock time t , assume the following form in terms of conformal time. The change of variable can be performed simply by using these identities:

$$\ell \frac{d}{dt} = \frac{d}{d\eta}, \quad \ell^2 \frac{d^2}{dt^2} = \frac{d^2}{d\eta^2} - H\ell \frac{d}{d\eta}, \quad (4.88)$$

which follow from (4.82).

Governing DEs for Perturbations using Conformal Time**Scalar**

$$\mathcal{D}''_{(k)} + (4 - 3\gamma)H\ell\mathcal{D}'_{(k)} + \left(-\frac{3}{2}(2 - \gamma)(3\gamma - 2)\Omega - 6(\gamma - 1)\Omega_\Lambda + \frac{c_s^2 k^2}{H^2 \ell^2}\right) H^2 \ell^2 \mathcal{D}_{(k)} = 0. \quad (4.89)$$

Vector

$$\mathcal{W}'_{(k)} - (3\gamma - 4)H\ell\mathcal{W}_{(k)} = 0. \quad (4.90)$$

Tensor

$$\mathcal{X}''_{(k)} + 2H\ell\mathcal{X}'_{(k)} + \left(- (3\gamma - 2)\Omega + 2\Omega_\Lambda + \frac{k^2 + 2K}{H^2 \ell^2}\right) H^2 \ell^2 \mathcal{X}_{(k)} = 0. \quad (4.91)$$

In the case of vector perturbations, the governing DE (4.90) can be integrated directly, on applying⁸

$$H = \frac{\ell'}{\ell^2}. \quad (4.92)$$

We then obtain

$$\mathcal{W}_{(k)} = C(\lambda\ell)^{3\gamma-4}. \quad (4.93)$$

In the case of scalar and tensor perturbations, the second-order governing DEs can be transformed into normal form by making a change of dependent variable. In both cases the resulting DE⁹ has the following form, in terms of a constant frequency ω and a potential $U(\eta)$:

$$Y''_{(k)} + [\omega^2 - U(\eta)] Y_{(k)} = 0. \quad (4.94)$$

The wave number k enters only into the frequency, which also depends on the curvature index K , and, in the case of scalar perturbations, on the equation of state. The potential $U(\eta)$ depends on the background FL model (H , ℓ , Ω and Ω_Λ). The specific DEs, given in the table below, are obtained by making the indicated change of variable in the governing DEs (4.89) and (4.91). In order to simplify the subsequent analysis we use the notation

$$\beta = \frac{2}{3\gamma - 2}, \quad \text{or, equivalently,} \quad \beta = \frac{2}{3w + 1}, \quad (4.95)$$

⁸This expression is obtained from (3.9) on making the change of variable (4.88).

⁹To the best of the author's knowledge, this similarity between tensor and scalar perturbations has not been noted before.

as in Bardeen (1980).

Normal Form of Governing DEs for Scalar and Tensor Perturbations

Scalar

$$\text{Change of variable: } \mathcal{D}_{(k)} = (\lambda\ell)^{\frac{1}{\beta}-1} \hat{\mathcal{D}}_{(k)}, \quad (4.96)$$

$$\text{Governing DE: } \hat{\mathcal{D}}_{(k)}'' + [\omega^2 - U(\eta)] \hat{\mathcal{D}}_{(k)} = 0, \quad (4.97)$$

with

$$\omega^2 = c_s^2 k^2 + \frac{1}{\beta^2} (\beta - 1)^2 K, \quad (4.98)$$

$$U(\eta) = \frac{1}{\beta^2} (1 + \beta) (\beta\Omega + \Omega_\Lambda) H^2 \ell^2. \quad (4.99)$$

Tensor

$$\text{Change of variable: } \mathcal{X}_{(k)} = (\lambda\ell)^{-1} \hat{\mathcal{X}}_{(k)}, \quad (4.100)$$

$$\text{Governing DE: } \hat{\mathcal{X}}_{(k)}'' + [\omega^2 - U(\eta)] \hat{\mathcal{X}}_{(k)} = 0. \quad (4.101)$$

with

$$\omega^2 = k^2 + 3K, \quad (4.102)$$

$$U(\eta) = \frac{1}{\beta} (1 + \beta) \Omega H^2 \ell^2. \quad (4.103)$$

Note: Here λ is the constant conformal parameter introduced in section 3.2.1, having dimension $(length)^{-1}$.

Remark: Writing the governing DEs in normal form has several advantages. Firstly, it gives insight into the qualitative behaviour of the solutions. In particular, if $\omega^2 \gg U(\eta)$ the solutions will be approximated by oscillatory solutions of frequency ω , while if $\omega^2 \ll U(\eta)$, the solutions will be approximated by power

law solutions. Secondly, if the background FL model has $\Lambda = 0$, the general solution can be written in terms of Bessel or Legendre functions, as we now show.

4.7.3 General Solutions for Scalar and Tensor Perturbations with $\Lambda = 0$

In the case $\Lambda = 0$, the potential $U(\eta)$ in the governing DEs is given by

$$U(\eta) = \frac{1}{\beta}(1 + \beta)\Omega H^2 \ell^2 \quad (4.104)$$

for both scalar and tensor perturbations (see equations (4.99) and (4.103)). The background solution is (3.206):

$$\ell(\eta) = \lambda^{-1} [S_K(T)]^\beta, \quad T = \frac{\eta}{\beta}, \quad (4.105)$$

where

$$S_K(T) = (T, \sinh T, \sin T), \quad (4.106)$$

for $K = (0, -1, +1)$. On noting that $\Omega H^2 \ell^2 = \frac{1}{3}\mu \ell^2$, it follows from (3.207) that

$$\Omega H^2 \ell^2 = (\lambda \ell)^{-\frac{2}{\beta}}. \quad (4.107)$$

Thus, using (4.105) we obtain

$$\Omega H^2 \ell^2 = [S_K(T)]^{-2}. \quad (4.108)$$

Observe now that the governing DEs in normal form (equations (4.97) and (4.101)) coincide with the Bessel DE or the Legendre DE, as given in Appendix H (see equations (H.1)-(H.6)), with the parameters determined as follows:

<u>Curvature</u>	<u>DE Type</u>	<u>Parameters</u>	
$K = 0$	Bessel DE	$\mu = \beta + \frac{1}{2}$	$a = \omega$
$K = -1$	Legendre (Toroidal) DE	$\mu = \beta + \frac{1}{2}$	$\nu = -\frac{1}{2} + i\beta\omega$
$K = +1$	Legendre (Conical) DE	$\mu = \beta + \frac{1}{2}$	$\nu = -\frac{1}{2} + \beta\omega$

The value of ω , as given by (4.98) and (4.102) distinguishes between scalar and tensor perturbations. The solutions of the three DEs are given by (H.2), (H.4) and (H.6), in terms of Bessel and Legendre functions. The resulting expressions for $\mathcal{D}_{(k)}$ and $\mathcal{X}_{(k)}$, obtained from (4.96), (4.100) and (4.105) are listed in the following table

General Solution of the Governing DEs for Scalar and Tensor Perturbations When $\Lambda = 0$ **Flat FL ($K = 0$)**

$$\left. \begin{array}{l} \text{Scalar} \quad \mathcal{D}_{(k)}(\eta) = \eta^{\frac{3}{2}-\beta} \\ \text{Tensor} \quad \mathcal{X}_{(k)}(\eta) = \eta^{\frac{1}{2}-\beta} \end{array} \right\} \left[C_+ J_{\beta+\frac{1}{2}}(a\eta) + C_- Y_{\beta+\frac{1}{2}}(a\eta) \right], \quad (4.109)$$

where

$$a = (c_s k, k) \quad (4.110)$$

for scalar and tensor perturbations, respectively.

Open FL ($K = -1$)

$$\left. \begin{array}{l} \text{Scalar} \quad \mathcal{D}_{(k)}(\eta) = \left(\sinh \frac{\eta}{\beta} \right)^{\frac{3}{2}-\beta} \\ \text{Tensor} \quad \mathcal{X}_{(k)}(\eta) = \left(\sinh \frac{\eta}{\beta} \right)^{\frac{1}{2}-\beta} \end{array} \right\} \left[C_+ Q_\nu^\mu \left(\cosh \frac{\eta}{\beta} \right) + C_- P_\nu^\mu \left(\cosh \frac{\eta}{\beta} \right) \right], \quad (4.111)$$

where

$$\mu = \beta + \frac{1}{2}, \quad \nu + \frac{1}{2} = i\beta\omega, \quad (4.112)$$

and the frequency ω is given by (4.98) and (4.102) respectively.

Closed FL ($K = +1$)

Replace hyperbolic functions by trigonometric functions in the solutions (4.111) for open FL, and use

$$\nu + \frac{1}{2} = \beta\omega. \quad (4.113)$$

Remark: The physically important cases, namely when the matter content is dust (*i.e.* $\gamma = 1, \beta = 2$) or radiation (*i.e.* $\gamma = \frac{4}{3}, \beta = 1$) require comment. For dust, the scalar solution is not valid since $c_s^2 = \gamma - 1 = 0$. We will derive the solution in this case in section 4.8. The tensor solution is valid, however, and the relevant Bessel and Legendre functions are in fact elementary and are given in section

4.8.4. For radiation, both the scalar and tensor solutions are elementary. We will give them in section 4.9.

One can use asymptotic approximations of the Bessel and Legendre functions (Appendix H) to obtain approximate solutions in restricted regimes.

4.7.4 Functional Dependence of First-Order Quantities

We are interested in describing to what extent a perturbed FL cosmology deviates from an exact FL cosmology.

It is known that cosmological observations place bounds on the first order quantities, specifically on dimensionless quantities formed by dividing the first order quantities by an appropriate power of the Hubble scalar. The specific quantities are

$$\text{Kinematic quantities: } \frac{\sigma_{ab}}{H}, \frac{\omega_a}{H}, \quad (4.114)$$

$$\text{Spatial gradients: } \frac{\hat{\nabla}_a \mu}{H\mu}, \frac{\hat{\nabla}_a H}{H^2}, \quad (4.115)$$

$$\text{Weyl tensor: } \frac{E_{ab}}{H^2}, \frac{H_{ab}}{H^2}. \quad (4.116)$$

The first analysis of this type was due to Sachs and Wolfe (1967) who showed that observations of distant galaxies could, in principle, place bounds on these quantities. More recently, Maartens et al (1995a,b) have shown that the high isotropy of the CMBR ($\frac{\Delta T}{T} \approx 10^{-5}$) places strong bounds on these quantities.¹⁰

It is natural to use these Hubble-normalized quantities as a measure of how close the physical model is to an idealized FL model, and we will thus calculate their time dependence for the three perturbation types. A growing mode is an indication of an instability in the background FL model.

We now briefly mention the connection between the geometrical approach and Bardeen's gauge-invariant metric approach to cosmological perturbations (Bardeen (1980)). In Bardeen's approach, the basic variables are gauge-invariant linear combinations of the metric perturbations. However, it turns out that Bardeen's metric potentials Φ_H , Ψ and $H_T^{(2)}$ have the same time dependence as certain expressions

¹⁰The analysis is complicated and subject to assumptions being imposed on the time derivatives of the multipoles of the CMBR.

formed from the basic variables in the geometrical approach. It can be shown that¹¹

$$\Phi_H \sim \Omega H^2 \ell^2 \mathcal{D}_{(k)} \quad (\text{scalar}), \quad (4.117)$$

$$\Psi \sim \Omega H^2 \ell^2 \mathcal{W}_{(k)} \quad (\text{vector}), \quad (4.118)$$

$$H_T^{(2)} \sim 2H\ell \mathcal{X}_{(k)} + \mathcal{X}'_{(k)} \quad (\text{tensor}). \quad (4.119)$$

We refer the reader to Bruni et al. (1992, p45-9) for an overview of the relation between the geometrical framework and Bardeen's metric approach.

Using the results in sections 4.4.2, 4.5.2 and 4.6.2 we can obtain the dependence of the Hubble-normalized quantities (4.114)-(4.116) on the basic variables $\mathcal{D}_{(k)}$, $\mathcal{W}_{(k)}$ and $\mathcal{X}_{(k)}$ in terms of conformal time. In order to enable comparisons to be made, we also include Bardeen's metric potentials Φ_H (4.117)-(4.119). These results are provided in table 4.1.

¹¹This result relies on comparing the governing DEs for $\mathcal{D}_{(k)}$, $\mathcal{W}_{(k)}$ and $\mathcal{X}_{(k)}$ with the governing DEs defined in Bardeen's approach (Bardeen equation (4.9), (4.13) and (4.14)) and noting that the paired quantities $(\mathcal{D}_{(k)}, \epsilon_m)$, $(\mathcal{W}_{(k)}, V_c)$ and $(\mathcal{X}_{(k)}, \frac{d}{d\eta} H_T^{(2)})$ satisfy the same evolutions equation. Equations (4.117)-(4.118) then follow from Bardeen (4.3) and (4.12).

Functional dependence of first-order quantities.

Quantity	Scalar Perturbations	Vector Perturbations	Tensor Perturbations
$\frac{\sigma_{ab}}{H} \sim$	$-3(\gamma - 1)\mathcal{D}_{(k)} + \frac{\mathcal{D}'_{(k)}}{H\ell}$	$\left[\frac{k^2 - 2K}{H^2\ell^2} + 6\gamma\Omega \right] H\ell\mathcal{W}_{(k)}$	$\frac{\mathcal{X}_{(k)}}{H\ell}$
$\frac{\omega_a}{H} \sim$	0	$\frac{\mathcal{W}_{(k)}}{H\ell}$	0
$\frac{\hat{\nabla}_a\mu}{H\mu} \sim$	$\frac{\mathcal{D}_{(k)}}{H\ell}$	$\mathcal{W}_{(k)}$	0
$\frac{\hat{\nabla}_a H}{H^2} \sim$	$\frac{1}{H\ell} \left[3(\gamma - 1)\mathcal{D}_{(k)} - \frac{\mathcal{D}'_{(k)}}{H\ell} \right]$	$\left[\frac{2K}{H^2\ell^2} - 3\gamma\Omega \right] \mathcal{W}_{(k)}$	0
$\frac{E_{ab}}{H^2} \sim$	$\Omega\mathcal{D}_{(k)}$	$\Omega H\ell\mathcal{W}_{(k)}$	$\frac{\mathcal{X}_{(k)}}{H\ell} + \frac{\mathcal{X}'_{(k)}}{H^2\ell^2}$
$\frac{H_{ab}}{H^2} \sim$	0	$\Omega\mathcal{W}_{(k)}$	$\frac{\mathcal{X}_{(k)}}{H^2\ell^2}$
$\Phi_H \sim$	$\Omega H^2\ell^2\mathcal{D}_{(k)}$	-	-
$\Psi \sim$	-	$\Omega H^2\ell^2\mathcal{W}_{(k)}$	-
$H_T^{(2)} \sim$	-	-	$2H\ell\mathcal{X}_{(k)} + \mathcal{X}'_{(k)}$
$\frac{\hat{\nabla}_a^{(3)}R}{H^3} \sim$	$\left[\frac{3}{2}\gamma\Omega - 3(\gamma - 1) \right] \frac{\mathcal{D}_{(k)}}{H\ell} + \frac{\mathcal{D}'_{(k)}}{H^2\ell^2}$	$\frac{K}{H^2\ell^2}\mathcal{W}_{(k)}$	0
$\frac{{}^{(3)}C_{ab}}{H^3} \sim$	0	0	$2\frac{\mathcal{X}_{(k)}}{H^2\ell^2} + \frac{\mathcal{X}'_{(k)}}{H^3\ell^3}$

TABLE 4.1: Expressions for the first order quantities in terms of the basic variable for each perturbation type.

4.8 Perturbations of FL Models with Pressure-Free Matter

For the case where the background FL model contains dust and possibly a cosmological constant, the analysis of scalar perturbations simplifies significantly since one can avoid performing the harmonic decomposition. The evolution equation (4.42) for the fractional density gradient \mathcal{D}_a simplifies to

$$\ddot{\mathcal{D}}_{\langle a} + 2H\dot{\mathcal{D}}_{\langle a} - \frac{3}{2}\Omega H^2 \mathcal{D}_a = 0. \quad (4.120)$$

Since (4.10) implies $\dot{u}_a = 0$ for a dust background, a short calculation using (1.5) and (1.18) reveals $\dot{h}_{ab} = 0$, and hence that

$$\dot{\mathcal{D}}_{\langle a} = \dot{\mathcal{D}}_a, \quad \text{and} \quad \ddot{\mathcal{D}}_{\langle a} = \ddot{\mathcal{D}}_a. \quad (4.121)$$

Namely, we use the fact that the fluid 4-velocity is a geodesic in the case of dust. Thus, we can drop the spatial projections from (4.120) and so obtain

$$\ddot{\mathcal{D}}_a + 2H\dot{\mathcal{D}}_a - \frac{3}{2}\Omega H^2 \mathcal{D}_a = 0. \quad (4.122)$$

Although it follows that \mathcal{D}_a does not depend on the perturbation mode k , a harmonic decomposition is required in order to obtain expressions for the shear and electric Weyl tensor (see, for example, (4.54) and (4.55)).

4.8.1 The General Solution in Integral Form for Scalar Perturbations

A second simplification is that the evolution equation (4.122) has a first integral, which we derive as follows, using the spatial gradients \mathcal{D}_a , \mathcal{Z}_a and \mathcal{R}_a .

In the case of dust, the evolution equations, given in (4.40), (4.41) and (B.13) simplify significantly, and can be written as

$$\dot{\mathcal{D}}_a = -\mathcal{Z}_a, \quad (4.123)$$

$$(\ell^2 \mathcal{Z}_a)^\cdot = -\frac{1}{2}\mu\ell^2 \mathcal{D}_a, \quad (4.124)$$

$$(\ell^2 \mathcal{R}_a)^\cdot = -4K \mathcal{Z}_a. \quad (4.125)$$

Note that these DEs are not independent, since the 3 spatial gradients are linearly dependent. This result can be seen upon taking the spatial gradient of (1.43) and linearizing so as to obtain

$$\mathcal{R}_a = -4H\mathcal{Z}_a + 2\mu\mathcal{D}_a. \quad (4.126)$$

Observe that the quantity \tilde{C}_a , defined by¹²

$$\tilde{C}_a = \ell^2 \mathcal{R}_a - 4K \mathcal{D}_a. \quad (4.127)$$

satisfies

$$\frac{d}{dt} \tilde{C}_a = 0. \quad (4.128)$$

We can rewrite (4.127) in terms of $\mathcal{D}_a \dot{\mathcal{D}}_a$ using (4.123), (4.126), (3.31) and (3.35), so as to obtain

$$\tilde{C}_a = 4\ell^2 (H \dot{\mathcal{D}}_a + (\frac{3}{2}\Omega + \Omega_k) H^2 \mathcal{D}_a). \quad (4.129)$$

This equation is in fact a first integral of the evolution equation (4.122), since (4.122) is a consequence of (4.123) and (4.124).

On noting that, in the case of dust, (4.11) can be rewritten as

$$\dot{H} = -(\frac{3}{2}\Omega + \Omega_k) H^2, \quad (4.130)$$

we can further simplify (4.129) to read

$$\frac{d}{dt} \left(\frac{\mathcal{D}_a}{H} \right) = \frac{\tilde{C}_a}{4H^2 \ell^2}. \quad (4.131)$$

The solution of (4.131) is thus

$$\mathcal{D}_a(t) = C_a^{(+)} H \int_0^t \frac{d\tilde{t}}{H^2 \ell^2} + C_a^{(-)} \frac{H}{\lambda}, \quad (4.132)$$

for time-independent vectors $C_a^{(-)}$ and $C_a^{(+)}$, with $C_a^{(+)} = \tilde{C}_a$. For simplicity, we define

$$\mathcal{D}^{(+)}(t) = H \int_0^t \frac{d\tilde{t}}{H^2 \ell^2}, \quad \mathcal{D}^{(-)}(t) = \frac{H}{\lambda}, \quad (4.133)$$

and so write (4.132) as

$$\mathcal{D}_a(t) = C_a^{(+)} \mathcal{D}^{(+)}(t) + C_a^{(-)} \mathcal{D}^{(-)}(t). \quad (4.134)$$

In terms of conformal time η , the growing mode $\mathcal{D}^{(+)}(\eta)$, as given by (4.133), reads

$$\mathcal{D}^{(+)}(\eta) = H \int_0^\eta \frac{d\tilde{\eta}}{H^2 \ell}. \quad (4.135)$$

¹²Also see Bruni et al. (1992), eq (38).

Equation (4.135) can often be used to gain insight into the asymptotic behaviour of dust perturbations. If $H \rightarrow 0$ or $H \rightarrow \infty$ as $\eta \rightarrow \eta^*$ (where $\eta^* = \pm\infty$ is allowed), we can rewrite $\mathcal{D}^{(+)}$ as

$$\mathcal{D}^{(+)} \sim \frac{\left(\int_0^\eta \frac{d\tilde{\eta}}{H^2 \ell} \right)}{\left(\frac{1}{H} \right)}. \quad (4.136)$$

On applying L'Hôpital's rule, in the limit as $\eta \rightarrow \eta^*$, we obtain

$$\mathcal{D}^{(+)} \sim -\frac{1}{H' \ell}. \quad (4.137)$$

Then (4.130) gives an expression for the asymptotic behaviour of the growing mode in terms of zero-order quantities:

$$\mathcal{D}^{(+)}(\eta) \sim \frac{1}{\frac{3}{2}\Omega H^2 \ell^2 - K}, \quad \text{as } \eta \rightarrow \eta^*. \quad (4.138)$$

Although the general solution (4.133) cannot always be obtained in closed form, one can analyze the asymptotic behaviour of these solutions. We are primarily interested in three asymptotic epochs, namely flat FL ($\eta \approx 0$), Milne ($\eta \gg 1$, $\Omega_k \approx 1$) and de Sitter ($\eta \approx \eta_f$, $\Omega_\Lambda \approx 1$). The general behaviour of perturbations within these epochs shall be discussed in detail in section 4.10.

4.8.2 Scalar Perturbations with $\Lambda = 0$

We now use the general solution (4.135) to obtain the solution for a dust background with $\Lambda = 0$. The length scale (4.105), when specialized to the case of dust, is given by

$$\lambda \ell = (S_K(\eta/2))^2, \quad (4.139)$$

and the Hubble parameter by $H = \ell' / \ell^2$ and (4.139),

$$H \ell = \frac{C_K(\eta/2)}{S_K(\eta/2)}, \quad (4.140)$$

where $C_K(T)$ is defined analogous to $S_K(T)$ to be

$$C_K(T) = (1, \cosh T, \cos T), \quad (4.141)$$

for $K = (0, -1, +1)$.

We now consider the case of $K \neq 0$ separately. In this case, the growing mode can be determined from (4.135), (4.139) and (4.140), on integrating

$$\mathcal{D}^{(+)}(\eta) = \frac{H}{\lambda} \int \frac{S_K(\eta/2)^4}{C_K(\eta/2)^2} d\eta. \quad (4.142)$$

This integral is elementary and can be expressed in terms of H and ℓ after a short calculation:

$$\mathcal{D}^{(+)} = \frac{1}{\lambda\ell} \left(2 - \frac{3}{2}\eta H\ell \right) + H^2\ell^2. \quad (4.143)$$

It follows from (4.107) with $\beta = 2$ that

$$\Omega H^2\ell^2 = \frac{1}{\lambda\ell}, \quad (4.144)$$

which allows us to rewrite the Friedmann equation (3.36) as

$$H^2\ell^2 = -K + \frac{1}{\lambda\ell}. \quad (4.145)$$

We can now eliminate the $H^2\ell^2$ term in (4.143), obtaining

$$\mathcal{D}^{(+)} = \frac{3}{\lambda\ell} \left(1 - \frac{1}{2}\eta H\ell \right) - K. \quad (4.146)$$

The solution of (4.122) for $K \neq 0$ is then

$$\mathcal{D}_a = C_a^{(+)} \left[\frac{3}{\lambda\ell} \left(1 - \frac{1}{2}\eta H\ell \right) - K \right] + C_a^{(-)} \left(\frac{H}{\lambda} \right), \quad (4.147)$$

where H and ℓ are given by (4.139) and (4.140).

In the case of $K = 0$, the integral (4.142) evaluates to

$$\mathcal{D}^{(+)} = \frac{1}{10}\eta^2, \quad (4.148)$$

where the coefficient $\frac{1}{10}$ is chosen so as to be asymptotic to the growing mode solution for $K \neq 0$ when $\eta \rightarrow 0$. The solution for a flat FL background is then

$$\mathcal{D}_a = C_a^{(+)} \left(\frac{1}{10}\eta^2 \right) + C_a^{(-)} (8\eta^{-3}). \quad (4.149)$$

We depict the growing and decaying mode solutions in figure 4.2.

We note that the solution (4.147) can also be derived from the governing DE (4.122), upon applying the general solutions (H.2), (H.34) and (H.35).




FIGURE 4.2: A depiction of the qualitative behaviour of the growing and decaying mode solutions of (4.122) when $\Lambda = 0$. From left to right, the plots depict negative ($K = -1$), flat ($K = 0$) and positive ($K = +1$) curvature.

4.8.3 Scalar Perturbations with $\Lambda > 0$, $K = 0$

In the case $\Lambda > 0$ with flat spatial geometry ($K = 0$), the general solution (4.133) can be given in terms of toroidal Legendre functions with clock time as the time variable, as follows. We transform the governing DE (4.122) to normal form by means of the change of variable

$$\mathcal{D}_a = (\lambda\ell)^{-1}\hat{\mathcal{D}}_a. \quad (4.150)$$

The result is

$$\hat{\mathcal{D}}_a'' - H^2\hat{\mathcal{D}}_a = 0, \quad \hat{\mathcal{D}}_a = (\lambda\ell)^{-1}\mathcal{D}_a. \quad (4.151)$$

The length scale of the background solution is given by (3.225). We calculate H using $H = \dot{\ell}/\ell$. The results are

$$\ell = \lambda^{-1}(\sinh T)^{2/3}, \quad H = \lambda \coth T, \quad (4.152)$$

where

$$T = \frac{3}{2}\lambda t, \quad \text{and} \quad \lambda = \sqrt{\frac{\Lambda}{3}}. \quad (4.153)$$

We substitute (4.152) into (4.151) and use T as the independent variable. After applying a standard identity, we obtain

$$\frac{d^2}{dT^2}\hat{\mathcal{D}}_a - \frac{4}{9}\left[1 + \frac{1}{\sinh^2 T}\right]\hat{\mathcal{D}}_a = 0, \quad \mathcal{D}_a = \left[\sinh\left(\frac{1}{2}\sqrt{3\Lambda}t\right)\right]^{-2/3}\hat{\mathcal{D}}_a. \quad (4.154)$$

This DE is the toroidal Legendre DE (H.3) with $\mu = \frac{5}{6}$ and $\nu = \frac{1}{6}$ whose solution is given by (H.4). Thus, we have

$$\mathcal{D}_a(t) = (\sinh T)^{-\frac{1}{6}} \left[C_a^{(+)} Q_{1/6}^{5/6}(\cosh T) + C_a^{(-)} P_{1/6}^{5/6}(\cosh T) \right]. \quad (4.155)$$

The second term of this solution is in fact an elementary function due to the identity¹³

$$(\sinh T)^{-\frac{1}{6}} P_{1/6}^{5/6}(\cosh T) = A \coth(T), \quad (4.156)$$

for A constant. It follows from (4.152) that (4.155) can be rewritten as

$$\mathcal{D}_a(t) = C_a^{(+)} (\sinh T)^{-\frac{1}{6}} Q_{1/6}^{5/6}(\cosh T) + C_a^{(-)} \frac{H}{\lambda}, \quad (4.157)$$

where T is given by (4.153). Observe that the second term corresponds to the second term in the general solution (4.132). However, the growing mode in (4.157) does not correspond to the growing mode given in the general solution (4.132), since it diverges as $t \rightarrow 0^+$ (see (H.39)). Thus the growing mode $\mathcal{D}^{(+)}(t)$ in (4.132) must be a linear combination of the two solutions in (4.157):

$$\mathcal{D}^{(+)} = H \int_0^t \frac{dt}{H^2 \ell^2} = a_1 (\sinh T)^{-\frac{1}{6}} Q_{1/6}^{5/6}(\cosh T) + a_2 \frac{H}{\lambda}. \quad (4.158)$$

The growing mode defined in this way will satisfy

$$\lim_{t \rightarrow 0} \mathcal{D}^{(+)} = 0. \quad (4.159)$$

We depict the growing and decaying mode solutions in figure 4.3.

4.8.4 Tensor Perturbations for $\Lambda = 0$

The solution for $\mathcal{X}_{(k)}(\eta)$ is obtained in terms of Bessel and Legendre functions by choosing $\beta = 2$ in the general solution (4.109), (4.111) and (4.113). Since the index has the value $\mu = \frac{5}{2}$, these solutions are elementary functions, given by (H.20)-(H.25). Upon comparing the functional dependence in each case, we obtain the following unified form of the solution:

$$\mathcal{X}_{(k)}(\eta) = \frac{1}{\lambda \ell} [c_+ \mathcal{X}_{(k)}^+(\eta) + c_- \mathcal{X}_{(k)}^-(\eta)], \quad (4.160)$$

¹³This result is obtained from Abramowitz and Stegun (1970), equations (8.5.1) and (8.6.16).




FIGURE 4.3: A depiction of the qualitative behaviour of the growing and decaying modes in an FL cosmology for density perturbations with dust and cosmological constant.

where

$$\mathcal{X}_{(k)}^+(\eta) = (4n^2 - K - 3H^2\ell^2) \sin(n\eta) + 6nH\ell \cos(n\eta), \quad (4.161)$$

$$\mathcal{X}_{(k)}^-(\eta) = (4n^2 - K - 3H^2\ell^2) \cos(n\eta) - 6nH\ell \sin(n\eta), \quad (4.162)$$

and $n^2 = k^2 + 3K$. Here, the coefficients c_+ and c_- are constants that depend on k .

Tensor perturbations describe the propagation of gravitational waves, which are represented by the electric and magnetic Weyl tensor, E_{ab} and H_{ab} . We specialize the solution (4.160) to $K = -1$ and use (3.206) and $H = \ell'/\ell^2$ to determine ℓ and H . On applying (4.80) and (4.81), we can determine the asymptotic behaviour of E_{ab} and H_{ab} in three asymptotic regimes: The first regime ($\eta \ll 1$ and $k\eta \ll 1$) describes long wavelength perturbations that occur outside the particle horizon close to the singularity; the second regime ($\eta \ll 1$ and $k\eta \gg 1$) describes perturbations within the particle horizon close to the singularity; lastly, the long term behaviour of the perturbation, when spatial curvature is significant, is described in the third regime ($\eta \gg 1$).

	$\eta \ll 1$ and $k\eta \ll 1$	$\eta \ll 1$ and $k\eta \gg 1$	$\eta \gg 1$
E_{ab}/H^2	$c_+\eta^2 + c_-\eta^{-3}$	$A \cos(n\eta + \phi)$	$Ae^{-\eta} \cos(n\eta + \phi)$
H_{ab}/H^2	$c_+\eta^3 + c_-\eta^{-2}$	$A \sin(n\eta + \phi)$	$Ae^{-\eta} \sin(n\eta + \phi)$

Here c_{\pm} , A and ϕ are arbitrary constants that depend on k . In the first regime we can see that the

tensor perturbations do not represent gravitational waves since they have a power law dependence on η . We depict the behaviour of $E_{(k)}/H^2$ in figure 4.4, for three given choices of n .

4.9 Perturbations of Radiation-Filled FL Models

We now derive the general solution of the governing DEs for a radiation fluid with zero cosmological constant ($\Lambda = 0$) in terms of elementary functions.

The solution for $\mathcal{D}_{(k)}(\eta)$ is obtained by choosing $\beta = 1$ in the general solution (4.109), (4.111) and (4.113). Since the index has the value $\mu = \frac{3}{2}$, these solutions are elementary functions, given by (H.14)-(H.19). Upon comparing the functional dependence in each case, we obtain the following unified form of the solution:

$$\mathcal{D}_{(k)}(\eta) = c_+ \mathcal{D}^{(+)}(\eta) + c_- \mathcal{D}^{(-)}(\eta), \quad (4.163)$$

where

$$\mathcal{D}^{(+)}(\eta) = H\ell \sin(c_s k \eta) - c_s k \cos(c_s k \eta), \quad (4.164)$$

$$\mathcal{D}^{(-)}(\eta) = H\ell \cos(c_s k \eta) + c_s k \sin(c_s k \eta), \quad (4.165)$$

and

$$H\ell = \begin{cases} \coth \eta & \text{Open FL } (K = -1), \\ \eta^{-1} & \text{Flat FL } (K = 0), \\ \cot \eta & \text{Closed FL } (K = +1). \end{cases} \quad (4.166)$$

The coefficients c_+ and c_- are constants that depend on k .

The solution for $\mathcal{X}_{(k)}(\eta)$ is obtained by choosing $\beta = 1$ in the general solution (4.109), (4.111) and (4.113). Since the index has the value $\mu = \frac{3}{2}$, these solutions are elementary functions, given by (H.14)-(H.19). Upon comparing the functional dependence in each case, we obtain the following unified form of the solution:

$$\mathcal{X}_{(k)}(\eta) = \frac{1}{\lambda\ell} [c_+ \mathcal{X}_{(k)}^+(\eta) + c_- \mathcal{X}_{(k)}^-(\eta)], \quad (4.167)$$

where

$$\mathcal{X}_{(k)}^+(\eta) = H\ell \sin(n\eta) - n \cos(n\eta), \quad (4.168)$$

$$\mathcal{X}_{(k)}^-(\eta) = H\ell \cos(n\eta) + n \sin(n\eta), \quad (4.169)$$

$$(4.170)$$




FIGURE 4.4: Tensor modes of the Hubble-normalized electric Weyl tensor $E_{(k)}/H^2$ for an open FL background ($K = -1$) with dust. From top to bottom, the eigenmodes in each plot are $n = 10$, $n = 100$ and $n = 1000$.

and $n^2 = k^2 + 3K$. The coefficients c_+ and c_- are constants that depend on k .

As with perturbations of a dust background, we specialize the solution (4.163) to $K = -1$ and use (3.206) and $H = \ell'/\ell^2$ to determine ℓ and H . On applying (4.80) and (4.80), we can determine the asymptotic behaviour of E_{ab} and H_{ab} in three asymptotic regimes.

	$\eta \ll 1$ and $k\eta \ll 1$	$\eta \ll 1$ and $k\eta \gg 1$	$\eta \gg 1$
E_{ab}/H^2	$c_+\eta^2 + c_-\eta^{-1}$	$A\eta \cos(n\eta + \phi)$	$Ae^{-\eta} \cos(n\eta + \phi)$
H_{ab}/H^2	$c_+\eta^3 + c_-$	$A\eta \sin(n\eta + \phi)$	$Ae^{-\eta} \sin(n\eta + \phi)$

Here c_{\pm} , A and ϕ are arbitrary constants that depend on k . We depict the behaviour of $E_{(k)}/H^2$ in figure 4.5, for three given choices of n .




FIGURE 4.5: Tensor modes of the Hubble-normalized electric Weyl tensor $E_{(k)}/H^2$ for an open FL background ($K = -1$) with radiation. From top to bottom, the eigenmodes in each plot are $n = 10$, $n = 100$ and $n = 1000$.

4.10 Perturbations in Asymptotic Epochs of FL Models

In this section we give approximate solutions of the governing ODEs for scalar and tensor perturbations in three asymptotic epochs for FL cosmologies. We also give the behaviour of the spatial gradients, kinematic quantities and Weyl tensor components for these approximate solutions. Further, to facilitate comparisons, we give the Bardeen (1980) gauge-invariant metric potentials for the approximate solutions.

The three asymptotic epochs are as follows:

Flat FL. The flat FL background is described by

$$\Omega = 1, \quad \Omega_\Lambda = 0, \quad \Omega_k = 0, \quad \lambda\ell = \left(\frac{\eta}{\beta}\right)^\beta, \quad H\ell = \frac{\beta}{\eta}. \quad (4.171)$$

This solution approximates the dynamics of any FL cosmology with an initial singularity with cosmological constant and arbitrary curvature at early times, since these models satisfy

$$\lim_{\eta \rightarrow 0} (\Omega, \Omega_\Lambda, \Omega_k) = (1, 0, 0). \quad (4.172)$$

(see table 3.2).

Milne. The Milne background is described by

$$\Omega = 0, \quad \Omega_\Lambda = 0, \quad \Omega_k = 1, \quad \lambda\ell = e^\eta, \quad H\ell = 1. \quad (4.173)$$

This solution approximates the dynamics of any open FL cosmology with zero cosmological constant at late times, since these models satisfy

$$\lim_{\eta \rightarrow \infty} (\Omega, \Omega_\Lambda, \Omega_k) = (0, 0, 1). \quad (4.174)$$

(see table 3.2).

de Sitter. The de Sitter background is described by

$$\Omega = 0, \quad \Omega_\Lambda = 1, \quad \Omega_k = 0, \quad \lambda\ell = \frac{1}{\tilde{\eta}}, \quad H\ell = \frac{1}{\tilde{\eta}}, \quad (4.175)$$

where $\tilde{\eta}$ is defined by (3.219):

$$\tilde{\eta} = \eta - \eta_f. \quad (4.176)$$

This solution approximates the dynamics of any ever-expanding FL cosmology with cosmological constant and arbitrary curvature at late times, since these models satisfy

$$\lim_{\eta \rightarrow \eta_f} (\Omega, \Omega_\Lambda, \Omega_k) = (0, 1, 0). \quad (4.177)$$

(see table 3.2).

Remark: The approximate solution for the case of flat FL with $k\eta \ll 1$ (Table 4.2) have been given by Goode (1989). To the best of the author's knowledge, the other results are new.

4.10.1 The Flat FL Asymptotic Epoch

The solutions of the governing equations in the flat FL background are obtained by substituting (4.171) into (4.109) and (4.93) and are given by

$$\mathcal{D}_{(k)}(\eta) = \eta^{\frac{3}{2}-\beta} \left[c_+ J_{\beta+\frac{1}{2}}(c_s k \eta) + c_- Y_{\beta+\frac{1}{2}}(c_s k \eta) \right], \quad (4.178)$$

$$\mathcal{W}_{(k)}(\eta) = C \eta^{2(1-\beta)}, \quad (4.179)$$

$$\mathcal{X}_{(k)}(\eta) = \eta^{\frac{1}{2}-\beta} \left[c_+ J_{\beta+\frac{1}{2}}(k \eta) + c_- Y_{\beta+\frac{1}{2}}(k \eta) \right]. \quad (4.180)$$

In the expressions for $\mathcal{D}_{(k)}(\eta)$ and $\mathcal{X}_{(k)}(\eta)$ we can identify two asymptotic regimes, namely $k\eta \ll 1$ and $k\eta \gg 1$. The restriction $k\eta \ll 1$ means that the perturbation mode corresponding to k is outside the particle horizon, while $k\eta \gg 1$ means the perturbation mode is inside the particle horizon (see section 4.7.1). The approximate solutions subject to these restrictions can be obtained from the asymptotic expressions for the Bessel functions in Appendix H.4.

These approximate solutions will also be valid for an open or closed FL background, for the following reason. If $K = \pm 1$ and $k\eta \ll 1$, it follows that η must satisfy $\eta \ll 1$ since $k > 1$ – see (F.11), (F.32) and (F.43). It thus follows from (4.172) that

$$\frac{\Omega_k}{\Omega} \ll 1, \quad (4.181)$$

i.e. the spatial curvature is dynamically negligible in the regime $k\eta \ll 1$.

The approximate solutions and the resulting time dependence of the first order quantities are given in table 4.2 ($k\eta \ll 1$) and table 4.3 ($k\eta \gg 1$).

Perturbations of FL models subject to $k\eta \ll 1$ and $\Omega_k/\Omega_m \ll 1$.

Quantity		Scalar Perturbations	Vector Perturbations	Tensor Perturbations
$\mathcal{D}_{(k)}$	\sim	$\{\eta^2, \eta^{1-2\beta}\}$	-	-
$\mathcal{W}_{(k)}$	\sim	-	$\{\eta^{2-2\beta}\}$	-
$\mathcal{X}_{(k)}$	\sim	-	-	$\{\eta, \eta^{-2\beta}\}$
<hr/>				
$\frac{\sigma_{ab}}{H}$	\sim	$\{\eta^2, \eta^{1-2\beta}\}$	$\{\eta^{1-2\beta}\}$	$\{\eta^2, \eta^{1-2\beta}\}$
$\frac{\omega_a}{H}$	\sim	0	$\{\eta^{3-2\beta}\}$	0
$\frac{\hat{\nabla}_a \mu}{H\mu}$	\sim	$\{\eta^3, \eta^{2-2\beta}\}$	$\{\eta^{2-2\beta}\}$	0
$\frac{\hat{\nabla}_a H}{H^2}$	\sim	$\{\eta^3, \eta^{2-2\beta}\}$	$\{\eta^{2-2\beta}\}$	0
$\frac{\hat{\nabla}_a^{(3)} R}{H^3}$	\sim	$\{\eta^3, \eta^{4-2\beta}\}^\dagger$	0	0
$\frac{E_{ab}}{H^2}$	\sim	$\{\eta^2, \eta^{1-2\beta}\}$	$\{\eta^{1-2\beta}\}$	$\{\eta^2, \eta^{1-2\beta}\}$
$\frac{H_{ab}}{H^2}$	\sim	0	$\{\eta^{2-2\beta}\}$	$\{\eta^3, \eta^{2-2\beta}\}$
<hr/>				
Φ_H	\sim	$\{1, \eta^{-1-2\beta}\}$	-	-
Ψ	\sim	-	$\{\eta^{-2\beta}\}$	-
$H_T^{(2)}$	\sim	-	-	$\{1, \eta^{1-2\beta}\}^\ddagger$

TABLE 4.2: This table describes perturbations of any FL model subject to $k\eta \ll 1$ and $\Omega_k/\Omega_m \ll 1$. For open and closed FL models the second restriction requires that $\eta \ll 1$, which is necessarily satisfied on account of the first restriction, since $k > 1$.

[†] The second mode in this case is obtained directly from the exact solution and is zero in the case of dust.

[‡] The second mode in this case cannot be obtained from the asymptotic behaviour of $\mathcal{X}_{(k)}$.

Perturbations of FL models subject to $k\eta \gg 1$ and $\Omega_k/\Omega_m \ll 1$.

Quantity		Scalar Perturbations ($\gamma \neq 1$) [†]	Vector Perturbations	Tensor Perturbations
$\mathcal{D}_{(k)}$	\sim	$\eta^{1-\beta} \sin(c_s k\eta + \phi)$	-	-
$\mathcal{W}_{(k)}$	\sim	-	$\eta^{2-2\beta}$	-
$\mathcal{X}_{(k)}$	\sim	-	-	$\eta^{-\beta} \sin(k\eta + \phi)$
$\frac{\sigma_{ab}}{H}$	\sim	$\eta^{2-\beta} \cos(c_s k\eta + \phi)$	$\eta^{3-2\beta}$	$\eta^{1-\beta} \sin(k\eta + \phi)$
$\frac{\omega_a}{H}$	\sim	0	$\eta^{3-2\beta}$	0
$\frac{\hat{\nabla}_a \mu}{H\mu}$	\sim	$\eta^{2-\beta} \sin(c_s k\eta + \phi)$	$\eta^{2-2\beta}$	0
$\frac{\hat{\nabla}_a H}{H^2}$	\sim	$\eta^{3-\beta} \cos(c_s k\eta + \phi)$	$\eta^{2-2\beta}$	0
$\frac{\hat{\nabla}_a^{(3)} R}{H^3}$	\sim	$\eta^{3-\beta} \sin(c_s k\eta + \tilde{\phi})$	0	0
$\frac{E_{ab}}{H^2}$	\sim	$\eta^{1-\beta} \sin(c_s k\eta + \phi)$	$\eta^{1-2\beta}$	$\eta^{2-\beta} \cos(k\eta + \phi)$
$\frac{H_{ab}}{H^2}$	\sim	0	$\eta^{2-2\beta}$	$\eta^{2-\beta} \sin(k\eta + \phi)$
Φ_H	\sim	$\eta^{-1-\beta} \sin(c_s k\eta + \phi)$	-	-
Ψ	\sim	-	$\eta^{-2\beta}$	-
$H_T^{(2)}$	\sim	-	-	$\eta^{-\beta} \cos(k\eta + \phi)$

TABLE 4.3: This table describes perturbations of flat FL with $k\eta \gg 1$, and open and closed FL with $k\eta \gg 1$ and $\eta \ll 1$. For open and closed FL models the second restriction requires that $\frac{\Omega_k}{\Omega} \ll 1$. The first restriction then requires that k is sufficiently large.

[†] For the case of $\gamma = 1$ see table 4.2.

4.10.2 The Milne Asymptotic Epoch

The governing DEs in the Milne background are obtained by substituting (4.173) into (4.97), (4.90) and (4.101) and are given by

$$\hat{\mathcal{D}}''_{(k)} + \omega^2 \hat{\mathcal{D}}_{(k)} = 0, \quad \mathcal{D}_{(k)} = e^{-(1-\frac{1}{\beta})\eta} \hat{\mathcal{D}}_{(k)}, \quad (4.182)$$

$$\mathcal{W}'_{(k)} + \frac{2}{\beta}(\beta - 1)\mathcal{W}_{(k)} = 0, \quad (4.183)$$

$$\hat{\mathcal{X}}''_{(k)} + n^2 \hat{\mathcal{X}}_{(k)} = 0, \quad \mathcal{X}_{(k)} = e^{-\eta} \hat{\mathcal{X}}_{(k)}. \quad (4.184)$$

where

$$\omega^2 = c_s^2 k^2 - (1 - \frac{1}{\beta})^2 \quad (4.185)$$

holds for scalar perturbations and

$$n^2 = k^2 - 3 \quad (4.186)$$

holds for tensor perturbations. For scalar perturbations there are two cases, namely $\omega^2 > 0$ and $\omega^2 \leq 0$. The second case includes dust (*i.e.* $c_s^2 = \gamma - 1 = 0$, $\beta = 2$), while the first case includes radiation (*i.e.* $\gamma = \frac{4}{3}$, $\beta = 1$). In addition, the condition $\omega^2 > 0$ is satisfied for all perfect fluids other than dust (*i.e.* $1 < \gamma < 2$) provided that the wave number k is sufficiently large, *i.e.* $k > k_{crit}$. We will restrict our considerations to the cases $\omega^2 > 0$ and dust.

The solutions are

$$\mathcal{D}_{(k)}(\eta) = \begin{cases} Ae^{-(1-\frac{1}{\beta})\eta} \sin(\omega\eta + \phi), & \text{if } \gamma > 1, k > k_{crit}, \\ c_+ + c_- e^{-\eta}, & \text{if } \gamma = 1, \end{cases} \quad (4.187)$$

$$\mathcal{W}_{(k)}(\eta) = C e^{-2(1-\frac{1}{\beta})\eta}, \quad (4.188)$$

$$\mathcal{X}_{(k)}(\eta) = Ae^{-\eta} \sin(n\eta + \phi), \quad (4.189)$$

where A is an arbitrary constant amplitude.

In order to calculate the approximate expression for E_{ab} we need an approximate expression for Ω subject to $\eta \gg 1$. A simple calculation using (3.206) and (3.207) yields the approximate asymptotic behaviour

$$\Omega = \exp\left(-\frac{2}{\beta}\eta\right) \ll 1. \quad (4.190)$$

The approximate solutions and the resulting time dependence of the first order quantities are given in table 4.4.

Perturbations of Milne.

Quantity	Scalar Perturbations		Vector Perturbations	Tensor Perturbations
	$\gamma = 1$	$\gamma > 1, k^2 > k_{crit}^2$		
$\mathcal{D}_{(k)}$	$\sim \{1, e^{-\eta}\}$	$e^{-(1-\frac{1}{\beta})\eta} \sin(\omega\eta + \phi)$	-	-
$\mathcal{W}_{(k)}$	-	-	$e^{-2(1-\frac{1}{\beta})\eta}$	-
$\mathcal{X}_{(k)}$	-	-	-	$e^{-\eta} \sin(n\eta + \phi)$
$\frac{\sigma_{ab}}{H}$	$\sim \{\eta e^{-\eta}, e^{-\eta}\}^\dagger$	$e^{-(1-\frac{1}{\beta})\eta} \sin(\omega\eta + \tilde{\phi})$	$e^{-2(1-\frac{1}{\beta})\eta}$	$e^{-\eta} \sin(n\eta + \phi)$
$\frac{\omega_a}{H}$	~ 0	0	$e^{-2(1-\frac{1}{\beta})\eta}$	0
$\frac{\hat{\nabla}_a \mu}{H\mu}$	$\sim \{1, e^{-\eta}\}$	$e^{-(1-\frac{1}{\beta})\eta} \sin(\omega\eta + \phi)$	$e^{-2(1-\frac{1}{\beta})\eta}$	0
$\frac{\hat{\nabla}_a H}{H^2}$	$\sim \{\eta e^{-\eta}, e^{-\eta}\}^\dagger$	$e^{-(1-\frac{1}{\beta})\eta} \sin(\omega\eta + \tilde{\phi})$	$e^{-2(1-\frac{1}{\beta})\eta}$	0
$\frac{\hat{\nabla}_a^{(3)} R}{H^3}$	$\sim \{e^{-\eta}, e^{-\eta}\}$	$e^{-(1-\frac{1}{\beta})\eta} \sin(\omega\eta + \tilde{\phi})$	$e^{-2(1-\frac{1}{\beta})\eta}$	0
$\frac{E_{ab}}{H^2}$	$\sim \{e^{-\eta}, e^{-2\eta}\}$	$e^{-(1+\frac{1}{\beta})\eta} \sin(\omega\eta + \phi)$	$e^{-2\eta}$	$e^{-\eta} \cos(n\eta + \phi)$
$\frac{H_{ab}}{H^2}$	~ 0	0	$e^{-2\eta}$	$e^{-\eta} \sin(n\eta + \phi)$
Φ_H	$\sim \{e^{-\eta}, e^{-2\eta}\}$	$e^{-(1+\frac{1}{\beta})\eta} \sin(\omega\eta + \phi)$	-	-
Ψ	-	-	$e^{-2\eta}$	-
$H_T^{(2)}$	-	-	-	$e^{-\eta} \sin(n\eta + \phi)$

TABLE 4.4: This table describes perturbations of open FL models with $\eta \gg 1$ and $\Omega_k/\Omega_m \gg 1$ and hence $k\eta \gg 1$.

[†] In the case of $\gamma = 1$ the growing mode is obtained from the growing mode for dust in an open FL background, given by (4.146).

4.10.3 The de Sitter Asymptotic Epoch

The governing DEs in the flat de Sitter background are obtained by substituting (4.175) into (4.97), (4.90) and (4.101) and are given by

$$\hat{\mathcal{D}}''_{(k)} + \left[c_s^2 k^2 - \frac{1}{\beta} \left(1 + \frac{1}{\beta} \right) \frac{1}{\tilde{\eta}^2} \right] \hat{\mathcal{D}}_{(k)} = 0, \quad \mathcal{D}_{(k)} = \tilde{\eta}^{1-\frac{1}{\beta}} \hat{\mathcal{D}}_{(k)}, \quad (4.191)$$

$$\mathcal{W}'_{(k)} + 2 \left(\frac{1}{\beta} - 1 \right) \frac{1}{\tilde{\eta}} \mathcal{W}_{(k)} = 0, \quad (4.192)$$

$$\hat{\mathcal{X}}''_{(k)} + n^2 \hat{\mathcal{X}}_{(k)} = 0, \quad \mathcal{X}_{(k)} = \tilde{\eta} \hat{\mathcal{X}}_{(k)}, \quad (4.193)$$

where the prime (') denotes differentiation with respect to $\tilde{\eta}$ and $n^2 = k^2 + 3$ – see (F.43). The solutions to these equations are given by¹⁴

$$\mathcal{D}_{(k)}(\eta) = \begin{cases} \tilde{\eta}^{\frac{3}{2}-\frac{1}{\beta}} \left[c_- J_{\frac{1}{2}+\frac{1}{\beta}}(c_s k \tilde{\eta}) + c_+ Y_{\frac{1}{2}+\frac{1}{\beta}}(c_s k \tilde{\eta}) \right], & \text{if } \gamma \neq 1 \\ c_- \tilde{\eta}^2 + c_+, & \text{if } \gamma = 1 \end{cases} \quad (4.194)$$

$$\mathcal{W}_{(k)}(\eta) = c \tilde{\eta}^{2-\frac{2}{\beta}}, \quad (4.195)$$

$$\mathcal{X}_{(k)}(\eta) = c_- \tilde{\eta} \sin(n\tilde{\eta}) + c_+ \tilde{\eta} \cos(n\tilde{\eta}). \quad (4.196)$$

As in the case of the flat FL background, we can distinguish two regimes, depending on the wave number, *i.e.* $k\tilde{\eta} \ll 1$ and $k\tilde{\eta} \gg 1$. These results are valid also for $K = \pm 1$, but require that $\Omega_k/\Omega_\Lambda \ll 1$, which holds for $\tilde{\eta} \ll 1$ on account of (4.175).

In order to calculate the approximate expression for E_{ab} we need an approximate expression for Ω subject to $\tilde{\eta} \ll 1$. A simple calculation using (3.225) and (3.226) yields the approximate asymptotic behaviour

$$\Omega \sim \tilde{\eta}^{3\gamma}. \quad (4.197)$$

The approximate solutions and the resulting time dependence of the first order quantities are given in table 4.5 ($k\tilde{\eta} \ll 1$) and table 4.6 ($k\tilde{\eta} \gg 1$).

¹⁴Observe that (4.191) is Bessel's DE (H.1) in normal form with $a = c_s k$ and $\mu^2 - \frac{1}{4} = \frac{1}{\beta} \left(1 + \frac{1}{\beta} \right)$, *i.e.* $\mu = \frac{1}{2} + \frac{1}{\beta}$.

Perturbations of de Sitter subject to $k\tilde{\eta} \ll 1$ and $\Omega_k/\Omega_\Lambda \ll 1$.

<u>Quantity</u>		<u>Scalar Perturbations</u>	<u>Vector Perturbations</u>	<u>Tensor Perturbations</u>
$\mathcal{D}_{(k)}$	\sim	$\{\tilde{\eta}^{1-\frac{2}{\beta}}, \tilde{\eta}^2\}$	-	-
$\mathcal{W}_{(k)}$	\sim	-	$\tilde{\eta}^{2-\frac{2}{\beta}}$	-
$\mathcal{X}_{(k)}$	\sim	-	-	$\{\tilde{\eta}, \tilde{\eta}^2\}$
$\frac{\sigma_{ab}}{H}$	\sim	$\{\tilde{\eta}^{3-\frac{2}{\beta}}, \tilde{\eta}^2\}^\dagger$	$\tilde{\eta}^{3-\frac{2}{\beta}}$	$\{\tilde{\eta}^2, \tilde{\eta}^3\}$
$\frac{\omega_a}{H}$	\sim	0	$\tilde{\eta}^{3-\frac{2}{\beta}}$	0
$\frac{\hat{\nabla}_a \mu}{H\mu}$	\sim	$\{\tilde{\eta}^{2-\frac{2}{\beta}}, \tilde{\eta}^3\}$	$\tilde{\eta}^{2-\frac{2}{\beta}}$	0
$\frac{\hat{\nabla}_a H}{H^2}$	\sim	$\{\tilde{\eta}^{4-\frac{2}{\beta}}, \tilde{\eta}^3\}^\dagger$	$\tilde{\eta}^4$	0
$\frac{\hat{\nabla}_a^{(3)} R}{H^3}$	\sim	$\{\tilde{\eta}^{4-\frac{2}{\beta}}, \tilde{\eta}^3\}$	0	0
$\frac{E_{ab}}{H^2}$	\sim	$\{\tilde{\eta}^3, \tilde{\eta}^{2(2+\frac{1}{\beta})}\}$	$\tilde{\eta}^3$	$\{\tilde{\eta}^4, \tilde{\eta}^3\}$
$\frac{H_{ab}}{H^2}$	\sim	0	$\tilde{\eta}^4$	$\{\tilde{\eta}^3, \tilde{\eta}^4\}$
Φ_H	\sim	$\{\tilde{\eta}, \tilde{\eta}^{3\gamma}\}$	-	-
Ψ	\sim	-	$\tilde{\eta}^2$	-
$H_T^{(2)}$	\sim	-	-	$\{1, \tilde{\eta}^3\}$

TABLE 4.5: This table describes perturbations of any FL model that is future asymptotic to de Sitter, subject to $k\tilde{\eta} \ll 1$.

[†] Due to cancellation of higher order terms, the growing mode must be obtained from the exact solution (4.194).

Perturbations of de Sitter subject to $k\tilde{\eta} \gg 1$ and $\Omega_k/\Omega_\Lambda \ll 1$.

Quantity	Scalar Perturbations ($\gamma \neq 1$)	Tensor Perturbations
$\mathcal{D}_{(k)} \sim$	$\tilde{\eta}^{1-\frac{1}{\beta}} \sin(c_s k \tilde{\eta} + \phi)$	-
$\mathcal{X}_{(k)} \sim$	-	$\tilde{\eta} \sin(n\tilde{\eta} + \phi)$
$\frac{\sigma_{ab}}{H} \sim$	$\tilde{\eta}^{2-\frac{1}{\beta}} \cos(c_s k \tilde{\eta} + \phi)$	$\tilde{\eta}^2 \sin(n\tilde{\eta} + \phi)$
$\frac{\omega_a}{H} \sim$	0	0
$\frac{\hat{\nabla}_a \mu}{H\mu} \sim$	$\tilde{\eta}^{2-\frac{1}{\beta}} \sin(c_s k \tilde{\eta} + \phi)$	0
$\frac{\hat{\nabla}_a H}{H^2} \sim$	$\tilde{\eta}^{3-\frac{1}{\beta}} \cos(c_s k \tilde{\eta} + \phi)$	0
$\frac{\hat{\nabla}_a^{(3)} R}{H^3} \sim$	$\tilde{\eta}^{3-\frac{1}{\beta}} \cos(c_s k \tilde{\eta} + \phi)$	0
$\frac{E_{ab}}{H^2} \sim$	$\tilde{\eta}^{3+\frac{1}{\beta}} \sin(c_s k \tilde{\eta} + \phi)$	$\tilde{\eta}^3 \cos(n\tilde{\eta} + \phi)$
$\frac{H_{ab}}{H^2} \sim$	0	$\tilde{\eta}^3 \sin(n\tilde{\eta} + \phi)$
$\Phi_H \sim$	$\tilde{\eta}^{1+\frac{1}{\beta}} \sin(c_s k \tilde{\eta} + \phi)$	-
$H_T^{(2)} \sim$	-	$\tilde{\eta} \cos(n\tilde{\eta} + \phi)$

TABLE 4.6: This table describes perturbations of any FL model that is future asymptotic to de Sitter, with $k\tilde{\eta} \gg 1$ and $\tilde{\eta} \ll 1$. In the case of scalar perturbations with $\gamma = 1$ or vector perturbations, the behaviour is identical to that given in table 4.5.

4.11 Discussion

We now summarize the principal features of linear perturbations of FL cosmologies.

- i) For each perturbation type the complete system of linearized evolution and constraint equations is reduced to a single ODE for the harmonic modes, together with expressions for the Hubble-normalized quantities (4.114)-(4.116). Solutions of the governing DE for vector perturbations can always be determined explicitly in terms of ℓ according to (4.93).
- ii) When written in normal form using conformal time, the governing DEs for scalar and tensor perturbations are essentially the same if $\Lambda = 0$, *i.e.* they are governed by the same potential U with a different frequency ω , but differ significantly if $\Lambda > 0$.
- iii) We have given a unified derivation of the known explicit solutions of the governing DEs for scalar and tensor perturbations in (4.109)-(4.113). Solutions are given in terms of Bessel functions if $\Lambda = 0$ and $K = 0$ with γ arbitrary. Solutions are given in terms of Legendre functions if $\Lambda = 0$ and $K = \pm 1$ with γ arbitrary. In the case of dust ($\gamma = 1$) and radiation ($\gamma = \frac{4}{3}$), solutions for all perturbation modes can be written in terms of elementary functions.

We now give a brief overview of the behaviour of scalar and tensor perturbations when the matter content is dust or radiation and possibly with a cosmological constant.

Scalar perturbations, as described by the spatial density gradient \mathcal{D}_a , have a growing mode and a decaying mode in the regime $k\eta \ll 1$. For *pressure-free matter* these two modes persist for all η . The growing mode is bounded in the Milne regime, *i.e.* if $\Lambda = 0$ and $K = -1$, and in the de Sitter regime, *i.e.* if $\Lambda > 0$.

For *radiation* (or more generally if the barotropic index satisfies $w > 1$), scalar perturbations are wavelike in the regime $c_s k\eta \gg 1$ (interpreted as sound waves with speed of propagation c_s). The spatial density gradient is wavelike with constant amplitude in the Milne regime (unbounded if $w > \frac{1}{3}$) and in the de Sitter regime (with $k\tilde{\eta} \ll 1$) it has a growing and unbounded mode and a decaying mode (more generally this occurs if $w > 0$).

Tensor perturbations, as described by the electric and magnetic Weyl curvature, have a growing mode and a decaying mode in the regime $k\eta \ll 1$. In the regime $k\eta \gg 1$ where $\Omega_k/\Omega_m \ll 1$ the behaviour is sinusoidal, with power-law amplitude.

In tables 4.2-4.6 we summarize the time-dependence of the Hubble-normalized kinematic quantities, Hubble-normalized Weyl tensor components and metric potentials in three asymptotic epochs for FL cosmologies. Tables 4.2 and 4.3 show clearly that the flat FL model is unstable, both with respect to long wavelength and short wavelength perturbations. This instability is not revealed by the gauge invariant metric potentials. Tables 4.4 - 4.6 show that spatial curvature has a stabilizing effect, as does the presence of a cosmological constant.

EVOLUTION AND CONSTRAINT EQUATIONS

As discussed in section 1.2.6, when the Ricci identities (1.50) and Bianchi identities (1.51) are applied to the Einstein field equations (1.2), we obtain a set of evolution and constraint equations for the kinematic quantities and Weyl tensor components. For an outline of the derivation of these equations see Ellis (1973, p24-30) and Ellis and van Elst (1998, p10-12).

Projected Field Equations. The Einstein field equations (1.2) can be rewritten as

$$R_{ab} = T_{ab} - \frac{1}{2}Tg_{ab}. \quad (\text{A.1})$$

We contract (A.1) with h_{ab} and u_a and use (1.26) in order to obtain the following relations:

$$R = \mu - 3p, \quad (\text{A.2})$$

$$R_{ab}u^a u^b = \frac{1}{2}(\mu + 3p), \quad (\text{A.3})$$

$$R_{ab}u^a h^b_c = -q_c, \quad (\text{A.4})$$

$$R_{(ab)} = \pi_{ab}, \quad (\text{A.5})$$

(see Ellis (1973), p24). Using (1.30) we also obtain two important relations relating the Riemann curvature tensor and Weyl tensor components:

$$R_{acbd}u^c u^d = E_{ab} - \frac{1}{2}R_{(ab)} + \frac{1}{3}h_{ab}R_{cd}u^c u^d, \quad (\text{A.6})$$

$$\frac{1}{2}\eta_{ac}{}^{ef}R_{efbd}u^c u^d = H_{ab} + \frac{1}{2}\eta_{abcd}R^c{}_e u^d u^e. \quad (\text{A.7})$$

Ricci Identities

$$\dot{H} = -H^2 + \frac{1}{3}(\hat{\nabla}_a \dot{u}^a + \dot{u}_a \dot{u}^a - 2\sigma^2 + 2\omega^2) - \frac{1}{6}(\mu + 3p), \quad (\text{A.8})$$

$$\dot{\sigma}_{\langle ab \rangle} = -2H\sigma_{ab} + \hat{\nabla}_{\langle a} \dot{u}_{b \rangle} + \dot{u}_{\langle a} \dot{u}_{b \rangle} - \sigma_{\langle a}^c \sigma_{b \rangle c} - \omega_{\langle a} \omega_{b \rangle} - (E_{ab} - \frac{1}{2}\pi_{ab}), \quad (\text{A.9})$$

$$\dot{\omega}_{\langle a \rangle} = -2H\omega_a + \sigma_{ab}\omega^b - \frac{1}{2}\text{curl}(\dot{u}_a), \quad (\text{A.10})$$

$$0 = \hat{\nabla}^b \sigma_{ab} - 2\hat{\nabla}_a H - \text{curl}(\omega_a) - 2\epsilon_{abc}\dot{u}^b \omega^c + q_a, \quad (\text{A.11})$$

$$0 = \hat{\nabla}^a \omega_a - \dot{u}^a \omega_a, \quad (\text{A.12})$$

$$0 = H_{ab} - 2\dot{u}_{\langle a} \omega_{b \rangle} - \hat{\nabla}_{\langle a} \omega_{b \rangle} - \text{curl}(\sigma_{ab}). \quad (\text{A.13})$$

The twice-contracted Bianchi identities (1.53) lead to the following evolution equations:

$$\dot{\mu} = -3H(\mu + p) - \hat{\nabla}_a q^a - 2\dot{u}_a q^a - \sigma_a^b \pi_b^a, \quad (\text{A.14})$$

$$\begin{aligned} \dot{q}_{\langle a \rangle} &= -4Hq_a - \hat{\nabla}_a p - (\mu + p)\dot{u}_a - \hat{\nabla}^b \pi_{ab} \\ &\quad - \dot{u}^b \pi_{ab} - \sigma^{ab} q_b + \epsilon_{abc} \omega^b q^c. \end{aligned} \quad (\text{A.15})$$

We are primarily interested in the case of a perfect fluid, where these equations reduce to

Twice-Contracted Bianchi Identities (Perfect Fluid Source)

$$\dot{\mu} = -3H(\mu + p), \quad (\text{A.16})$$

$$0 = (\mu + p)\dot{u}_a + \hat{\nabla}_a p. \quad (\text{A.17})$$

For simplicity, the equations obtained from the full Bianchi identities (1.51) are presented here in the case of a perfect fluid source.

Bianchi Identities (Perfect Fluid Source)

$$\begin{aligned}\dot{E}_{\langle ab \rangle} &= -3HE_{ab} + 3\sigma_{\langle a}{}^c E_{b \rangle c} - \frac{1}{2}(\mu + p)\sigma_{ab} + \text{curl}(H_{ab}) \\ &\quad + \epsilon_{ef(a} (2\dot{u}^e H_{b)}^f - \omega^e E_{b)}^f),\end{aligned}\tag{A.18}$$

$$\begin{aligned}\dot{H}_{\langle ab \rangle} &= -3HH_{ab} + 3\sigma_{\langle a}{}^c H_{b \rangle c} - \text{curl}(E_{ab}) \\ &\quad - \epsilon_{ef(a} (2\dot{u}^e E_{b)}^f + \omega^e H_{b)}^f),\end{aligned}\tag{A.19}$$

$$0 = \hat{\nabla}^b E_{ab} + 3\omega^b H_{ab} + \epsilon_{abc}\sigma^{be} H_e^c - \frac{1}{3}\hat{\nabla}_a \mu,\tag{A.20}$$

$$0 = \hat{\nabla}^b H_{ab} - 3\omega^b E_{ab} - \epsilon_{abc}\sigma^{be} E_e^c - (\mu + p)\omega_a.\tag{A.21}$$

Spatial Gradients of Zero-Order Quantities. In order to derive the governing equations for linear perturbations of FL, it is necessary to derive evolution equations for the spatial gradients of the matter density μ and the Hubble parameter H , defined by (4.8) and (4.9). By propositions 2.1.4 and 2.1.1, these quantities are zero in the background and hence gauge-invariant by the Stewart-Walker lemma.

We make use of the fact that the Raychaudhuri equation (A.8) can be rewritten as

$$\dot{H} = \frac{1}{3}\mathcal{T} - \frac{1}{2}(\mu + p),\tag{A.22}$$

where

$$\mathcal{T} = -3H^2 + \mu + \Lambda + \hat{\nabla}_a \dot{u}^a + \dot{u}^a \dot{u}_a - 2\sigma^2 + 2\omega^2.\tag{A.23}$$

In order to proceed, we require an identity which permits the interchange of the $\dot{}$ derivative and the spatial derivative $\hat{\nabla}_a$ when applied to a scalar field f . A straightforward calculation using (1.18) yields

$$(\hat{\nabla}_a f)^\cdot{}_{\perp} = -H\hat{\nabla}_a f + (\hat{\nabla}_a + \dot{u}_a)\dot{f} - (\sigma_a^b + \omega_a^b)\hat{\nabla}_b f,\tag{A.24}$$

for any scalar f (also see equation (92) in Ellis et al. (1990)).

We now derive evolution equations for X_a and Z_a , following Ellis and van Elst (1998, p56). Substituting $f = \mu$ into (A.24) and using (4.8), (4.9), (A.16) and (A.17) yields the evolution equation for X_a . If we instead choose $f = H$ in (A.24) and use (4.8), (4.9), (A.22), (A.16) and (A.17) we obtain the evolution equation for Z_a . The results of these derivations are given as follows:

Spatial Gradients of Zero-Order Quantities

$$\dot{X}_{(a)} = -4H X_a - (\sigma_a^b + \omega_a^b) X_b - (1 + w)\mu Z_a, \quad (\text{A.25})$$

$$\begin{aligned} \dot{Z}_{(a)} = & -3H Z_a - (\sigma_a^b + \omega_a^b) Z_b - \frac{1}{2} X_a \\ & + \mathcal{T} \dot{u}_a + \hat{\nabla}_a (\hat{\nabla}^b \dot{u}_b + \dot{u}^a \dot{u}_a - 2\sigma^2 + 2\omega^2). \end{aligned} \quad (\text{A.26})$$

SPATIAL CURVATURE

In this appendix we introduce the 3-curvature when the fundamental congruence has non-zero vorticity. This analysis leads to a generalization of the Gauss equation (1.37), the 3-Ricci tensor (1.39) and 3-Ricci scalar.

B.1 The Generalized Gauss Equation

We note that the projected covariant derivative $\hat{\nabla}_a$, defined in (1.11), is a non-commutative operator on the space-time manifold when vorticity is present, *i.e.* $\omega_a \neq 0$. After a short calculation using (1.5), (1.11) and (1.18), we obtain

$$\hat{\nabla}_a \hat{\nabla}_b f - \hat{\nabla}_b \hat{\nabla}_a f = -2\omega_{ab} \dot{f}. \quad (\text{B.1})$$

We now derive a similar identity for tensors with one or more index.

On projecting (1.18) orthogonal to u^a and applying (1.11), we obtain

$$\hat{\nabla}_b u_a = k_{ab}, \quad (\text{B.2})$$

where k_{ab} is defined by

$$k_{ab} = \omega_{ab} + \sigma_{ab} + Hh_{ab}, \quad (\text{B.3})$$

(see Ellis et al. (1990), eq. (76)).

Upon performing a similar computation as in (B.1) and using (1.50), we obtain the expression

$$\hat{\nabla}_a \hat{\nabla}_b X_c - \hat{\nabla}_b \hat{\nabla}_a X_c = -2\omega_{ab} \dot{X}_{\langle c} + {}^{(3)}R_{abcd} X^d, \quad (\text{B.4})$$

where ${}^{(3)}R_{abcd}$ is defined by

$${}^{(3)}R_{abcd} = (R_{abcd})_{\perp} - k_{ac}k_{bd} + k_{ad}k_{bc}, \quad (\text{B.5})$$

(see Ellis et al. (1990), eq. (79)). We note that this definition reduces to the 3-Riemann tensor given in (1.37) when u^a is irrotational. A similar identity to (B.4) can be obtained for tensors with two or more indices, generalized according to

$$\hat{\nabla}_{[a}\hat{\nabla}_{b]}S_{cd\dots ef} = -\omega_{ab}\dot{S}_{\perp cd\dots ef} + \frac{1}{2}({}^{(3)}R_{scba}T_{d\dots ef}^s + \dots + {}^{(3)}R_{sfba}T_{cd\dots e}^s). \quad (\text{B.6})$$

On contracting (B.5), we can obtain an expression for the spatial Ricci scalar and trace-free spatial Ricci tensor in terms of the spacetime quantities from (B.5), incorporating the Weyl tensor through (1.30) and (1.32):

$${}^{(3)}R = R + 2R_{bd}u^b u^d - 6H^2 + 2\sigma^2 - 2\omega^2, \quad (\text{B.7})$$

$$\begin{aligned} {}^{(3)}S_{ab} &= E_{ab} + \frac{1}{2}(h_a^c h_b^d - \frac{1}{3}h^{cd}h_{ab})R_{cd} - H\sigma_{ab} - H\omega_{ab} \\ &\quad - \frac{2}{3}(\sigma^2 - \omega^2)h_{ab} + \sigma_a^c \sigma_{cb} + \omega_a^c \omega_{cb} + \omega_a^c \sigma_{cb} + \sigma_a^c \omega_{cb}. \end{aligned} \quad (\text{B.8})$$

Using the field equations (1.2) in the form (A.2)-(A.5) to eliminate R and R_{ab} , we obtain

$${}^{(3)}R = -6H^2 + 2\mu + 2\sigma^2 - 2\omega^2, \quad (\text{B.9})$$

$$\begin{aligned} {}^{(3)}S_{ab} &= E_{ab} + \frac{1}{2}\pi_{ab} - H\sigma_{ab} - H\omega_{ab} - \frac{2}{3}(\sigma^2 - \omega^2)h_{ab} \\ &\quad + \sigma_a^c \sigma_{cb} + \omega_a^c \omega_{cb} + \omega_a^c \sigma_{cb} + \sigma_a^c \omega_{cb}. \end{aligned} \quad (\text{B.10})$$

B.2 The Spatial Gradient of the 3-Ricci Scalar

The *spatial gradient of the 3-Ricci scalar* is defined by

$$\mathcal{R}_a = \ell \hat{\nabla}_a {}^{(3)}R. \quad (\text{B.11})$$

An evolution equation for \mathcal{R}_a is obtained in the same manner as for \mathcal{D}_a and \mathcal{Z}_a , defined in (4.35) and (4.37), respectively. If we choose $f = {}^{(3)}R$ in (A.24) and use (4.9), (A.22), (A.16) and (A.17) we obtain

$$\begin{aligned} \dot{\mathcal{R}}_{\langle a} &= -2H\mathcal{R}_a - (\sigma_a^b + \omega_a^b)\mathcal{R}_b - \frac{4}{3}\mathcal{T}\mathcal{Z}_a \\ &\quad - 4\ell\mathcal{T}\dot{u}_a + \ell\hat{\nabla}_a(\hat{\nabla}^b \dot{u}_b - 2\sigma^2 + 2\omega^2). \end{aligned} \quad (\text{B.12})$$

Further, we can obtain an auxiliary equation describing the evolution of \mathcal{R}_a from (B.12) and (4.19), presented here for completeness:

$$\dot{\mathcal{R}}_{\langle a} = -2H\mathcal{R}_a - \frac{4K}{\ell^2}\mathcal{Z}_a + 4H\frac{c_s^2}{(1+w)}\left(\hat{\nabla}^2 + \frac{K}{\ell^2}\right)\mathcal{D}_a + 24c_s^2H^2\ell\text{curl}(\omega_a). \quad (\text{B.13})$$

SOLUTIONS OF THE FRIEDMANN EQUATION

In this appendix we list the solutions of the three first order quadratic DEs that arise by transforming the Friedmann DE. In the first two cases, the solution is uniquely determined by requiring $\frac{dL}{dT} > 0$ and the initial condition

$$L(0) = 0. \tag{C.1}$$

In the third case we require that

$$L(0) = 1, \quad \text{and} \quad L'(T) > 0 \text{ for } T > 0. \tag{C.2}$$

	<u>Differential Equation</u>	<u>Solution</u>
Case 1	$\left(\frac{dL}{dT}\right)^2 = 1 + 2\alpha L + L^2,$	$L = \sinh T + 2\alpha \sinh^2(\frac{1}{2}T).$
Case 2	$\left(\frac{dL}{dT}\right)^2 = 1 + 2\alpha L - L^2,$	$L = \sin T + 2\alpha \sin^2(\frac{1}{2}T).$
Case 3	$\left(\frac{dL}{dT}\right)^2 = L^2 - 1,$	$L = \cosh T.$

There are four known cases where the Friedmann equation can be transformed into the quadratic form (3.195) via a change of variable:

- i) 1-fluid plus spatial curvature.
- ii) 2-fluid flat FL.
- iii) 2-fluid FL plus arbitrary curvature. The equation of state parameters must satisfy

$$\gamma_1 = \frac{2}{3} + 2\Delta\gamma, \quad \gamma_2 = \frac{2}{3} + \Delta\gamma, \quad (\text{C.6})$$

where $\Delta\gamma$ is constant.

- iv) 3-fluid FL. The equation of state parameters must satisfy

$$\gamma_1 > \gamma_2 > \gamma_3, \quad \gamma_1 - \gamma_2 = \gamma_2 - \gamma_3 = \Delta\gamma, \quad (\text{C.7})$$

where $\Delta\gamma$ is constant.

C.1 Solutions with a Stiff Fluid

There are three known cases where we can obtain a solution to the Friedmann equation where one fluid is a so-called stiff fluid, *i.e.* $\gamma_s = 2$. In this section we briefly present each of these solutions.

SD-Universes. We choose the conformal parameter to be $\lambda = \lambda_d$, which, on using (3.113), leads to a single mass parameter defined by

$$m_s = \left(\frac{\lambda_s}{\lambda} \right)^{-4}. \quad (\text{C.8})$$

In this case the Friedmann equation reads

$$\left(\frac{d\ell}{dt} \right)^2 = (\lambda_s \ell)^{-4} + (\lambda_d \ell)^{-1}. \quad (\text{C.9})$$

We apply the change of variable

$$\mathcal{L} = \frac{\lambda_s^4}{\lambda} \ell^3, \quad \mathcal{T} = 3 \frac{\lambda_s^2}{\lambda} t, \quad (\text{C.10})$$

obtaining

$$\left(\frac{d\mathcal{L}}{d\mathcal{T}} \right)^2 = 1 + \mathcal{L}. \quad (\text{C.11})$$

Subject to the IC $\mathcal{L}(0) = 0$, the solution is

$$\mathcal{L}(T) = \frac{1}{4}T^2 + T, \quad (\text{C.12})$$

or equivalently

$$\ell(t) = \lambda^{-1} \left[\frac{9}{4}(\lambda t)^2 + 3\sqrt{m_s}(\lambda t) \right]^{1/3}. \quad (\text{C.13})$$

SRC-Universes. We choose the conformal parameter to be $\lambda = \lambda_r$, which, on using (3.113), leads to a single mass parameter defined by

$$m_s = \left(\frac{\lambda_s}{\lambda} \right)^{-4}. \quad (\text{C.14})$$

In this case the Friedmann equation reads

$$\left(\frac{d\ell}{d\eta} \right)^2 = (\lambda_s \ell)^{-4} + (\lambda_r \ell)^{-2} - K. \quad (\text{C.15})$$

We apply the change of variable

$$\mathcal{L} = \lambda_s^2 \ell^2, \quad T = 2\eta, \quad (\text{C.16})$$

obtaining

$$\left(\frac{d\mathcal{L}}{dT} \right)^2 = 1 + \frac{1}{\sqrt{m_s}} \mathcal{L} - K \mathcal{L}^2. \quad (\text{C.17})$$

Subject to the IC $\mathcal{L}(0) = 0$, the solution is

$$\ell(t) = \lambda^{-1} \left[\sqrt{m_s} S_K(2\eta) + S_K(\eta)^2 \right]^{1/2}. \quad (\text{C.18})$$

SDA-Universes. We choose the conformal parameter to be $\lambda = \lambda_\Lambda$, which, on using (3.113), leads to two mass parameters defined by

$$m_s = \left(\frac{\lambda_s}{\lambda} \right)^{-4}, \quad m_d = \left(\frac{\lambda_d}{\lambda} \right)^{-1}. \quad (\text{C.19})$$

In this case the Friedmann equation reads

$$\left(\frac{d\ell}{d\eta} \right)^2 = (\lambda_s \ell)^{-4} + (\lambda_d \ell)^{-1} + (\lambda \ell)^2. \quad (\text{C.20})$$

We apply the change of variable

$$\mathcal{L} = \lambda_s^2 \lambda \ell^3, \quad T = 3\lambda t, \quad (\text{C.21})$$

obtaining

$$\left(\frac{d\mathcal{L}}{dT}\right)^2 = 1 + \frac{m_d}{\sqrt{m_s}}\mathcal{L} + \mathcal{L}^2. \quad (\text{C.22})$$

Subject to the IC $\mathcal{L}(0) = 0$, the solution is

$$\ell(t) = \lambda^{-1} \left[\sqrt{m_s} \sinh(3\lambda t) + m_d \sinh^2\left(\frac{3}{2}\lambda t\right) \right]^{1/3}. \quad (\text{C.23})$$

C.2 Existence-Uniqueness for the Friedmann Equation

In this section we prove existence-uniqueness of solutions to the Friedmann DE (3.115), given by

$$\left(\frac{dL}{dT}\right)^2 = \sum_{i=1}^n m_i L^{-3\gamma_i+2} - K, \quad (\text{C.24})$$

for any choice of parameters $m_i > 0$, $i = 1, \dots, n-1$ and subject to the conditions

$$L(0) = 0, \quad \text{and} \quad \frac{dL}{dT} > 0, \quad (\text{C.25})$$

We can rewrite (C.24) as

$$\left(\frac{dL}{dT}\right)^2 = L^{-(3\gamma_1-2)} \left[\sum_{i=1}^n m_i L^{3(\gamma_1-\gamma_i)} - KL^{3\gamma_1-2} \right]. \quad (\text{C.26})$$

Since $\gamma_1 > \gamma_2 > \dots > \gamma_n$ and $\gamma_1 > \frac{2}{3}$, this DE is of the form

$$\left(\frac{dL}{dT}\right)^2 = L^{-(3\gamma_1-2)} F(L), \quad (\text{C.27})$$

with

$$F(0) = m_1 > 0. \quad (\text{C.28})$$

Since $\frac{dL}{dT} > 0$, we can rewrite (C.27) as

$$\frac{dT}{dL} = \frac{L^{\frac{1}{2}(3\gamma_1-2)}}{F(L)^{1/2}}, \quad L \geq 0. \quad (\text{C.29})$$

The general solution to (C.29) is then, for all $m_i > 0$,

$$T = G(L) + C, \quad (\text{C.30})$$

where $G(L)$ is the antiderivative of the RHS of (C.29). The requirement that $T = 0$ when $L = 0$ fixes C , and hence gives a unique solution.

CONFORMAL TIME

The conformal time variable η is defined in terms of clock time t and the length scale ℓ via

$$\frac{d\eta}{dt} = \frac{1}{\ell}. \quad (\text{D.1})$$

Unlike clock time, conformal time is not necessarily unbounded into the future for ever-expanding models. In fact, the presence or absence of certain types of fluids determines the behaviour of the conformal time variable. We present the following theorem:

Theorem D.1 *Let $(\mathcal{M}, \mathbf{g}, \mathbf{u})$ be an ever-expanding n -fluid FL cosmology whose length scale satisfies $\ell(0) = 0$ and where each fluid satisfies a linear equation of state (3.17) with equation of state parameters γ_i ordered according to $\gamma_1 > \gamma_2 > \dots > \gamma_n$. It follows that η satisfies the following constraints:*

- (a) η is bounded below iff $\gamma_1 > \frac{2}{3}$.
- (b) η is bounded above iff $\gamma_2 < \frac{2}{3}$.

Proof. It follows from the definition of an ever-expanding model (see section 3.1.6) that

$$\frac{d\ell}{dt} > 0 \quad \forall t > 0. \quad (\text{D.2})$$

Using the chain rule we expand (D.1) so as to obtain

$$\frac{d\eta}{d\ell} = \left[\ell \frac{d\ell}{dt} \right]^{-1}. \quad (\text{D.3})$$

It then follows from (D.2) and (D.3) that η is a strictly increasing function of ℓ for $t > 0$.

Let η_0 and η_* be the value of conformal time at ℓ_0 and ℓ_* , respectively. We now integrate (D.3) over (η_0, η_*) so as to obtain

$$\eta_* - \eta_0 = F(\ell_*) = \int_{\ell_0}^{\ell_*} \left[\ell \left(\frac{d\ell}{dt} \right) \right]^{-1} d\ell. \quad (\text{D.4})$$

Since η is strictly increasing, we note that η_* is bounded above if and only if $F(\ell_*)$ is bounded as $\ell_* \rightarrow +\infty$. Similarly, we note that η_* is bounded below if and only if $F(\ell_*)$ is bounded as $\ell_* \rightarrow 0^+$. We now prove the two results separately:

(a) Assume that condition (a) is not met. It then follows from (3.196) that \exists a constant $C > 0$ such that $\frac{d\ell}{dt} < C \forall t < t_0$. The integral (D.4) then takes the form

$$\eta_* = \eta_0 - \int_0^{\ell_0} \left[\ell \left(\frac{d\ell}{dt} \right) \right]^{-1} d\ell < \eta_0 - \frac{1}{C} \int_0^{\ell_0} \frac{d\ell}{\ell}, \quad (\text{D.5})$$

which is unbounded.

If condition (a) is met it then follows from (3.196) that \exists constants $C > 0$ and $\epsilon > 0$ such that $\frac{d\ell}{dt} > C\ell^{-\epsilon} \forall t < t_0$. The integral (D.4) then takes the form

$$\eta_* = \eta_0 - \int_0^{\ell_0} \left[\ell \left(\frac{d\ell}{dt} \right) \right]^{-1} d\ell > \eta_0 - \frac{1}{C} \int_0^{\ell_0} \frac{d\ell}{\ell^{1-\epsilon}}, \quad (\text{D.6})$$

which is bounded.

(b) Assume that condition (b) is not met. It then follows from (3.196) that \exists a constant $C > 0$ such that $\frac{d\ell}{dt} < C \forall t > t_0$. The integral (D.4) then takes the form

$$\eta_* = \eta_0 + \lim_{\ell_* \rightarrow +\infty} \int_{\ell_0}^{\ell_*} \left[\ell \left(\frac{d\ell}{dt} \right) \right]^{-1} d\ell > \eta_0 + \lim_{\ell_* \rightarrow +\infty} \frac{1}{C} \int_{\ell_0}^{\ell_*} \frac{d\ell}{\ell}, \quad (\text{D.7})$$

which is unbounded.

If condition (b) is met it then follows from (3.196) that \exists constants $C > 0$ and $\epsilon > 0$ such that $\frac{d\ell}{dt} > C\ell^\epsilon \forall t > t_0$. The integral (D.4) then takes the form

$$\eta_* = \eta_0 + \lim_{\ell_* \rightarrow +\infty} \int_{\ell_0}^{\ell_*} \left[\ell \left(\frac{d\ell}{dt} \right) \right]^{-1} d\ell > \eta_0 + \lim_{\ell_* \rightarrow +\infty} \frac{1}{C} \int_{\ell_0}^{\ell_*} \frac{d\ell}{\ell^{1+\epsilon}}, \quad (\text{D.8})$$

which is bounded.

Thus, we conclude that the constraints hold, completing the proof. \square

We note that this theorem also holds in the case of closed FL as long as $\frac{d\ell}{dt} > C$, where C is a constant satisfying $C > 0$. This constraint prevents models which recollapse or whose length scale tend to a constant as $t \rightarrow +\infty$.

COVARIANT DIFFERENTIAL IDENTITIES

In this appendix, we present a list of useful differential identities involving the derivative operators used in this thesis. This list represents an extension of the list given in Ellis et al. (1990) Appendix A by incorporating identities involving the Laplacian and curl operators.

Zero-order 3-curvature quantities for FL models: Before proceeding, we must first obtain zero order expressions for the background curvature. It quickly follows from (3.11), (1.40) and (1.47) that, to zero order, the background 3-curvature quantities satisfy the following relationships:

$$\ell^2 {}^{(3)}R_{abcd} = K (h_{ac}h_{bd} - h_{ad}h_{bc}), \quad (\text{E.1})$$

$$\ell^2 {}^{(3)}R_{ac} = 2K h_{ac}, \quad (\text{E.2})$$

$$\ell^2 {}^{(3)}R = 6K. \quad (\text{E.3})$$

We now present a list of linearized identities, derived using (1.7), (1.11), (1.16)-(1.17), (B.1) and (B.6). In all of the following relations, we shall assume that $\hat{\nabla}_a f$, X_a and T_{ab} and all their derivatives are first order quantities. Further, we assume that T_{ab} is symmetric and trace-free.

First-order time derivatives: The following set of linearized identities are used when interchanging the derivative along the fundamental congruence (\cdot) with $\hat{\nabla}$, $\text{curl}(\cdot)$ or $\hat{\nabla}^2$:

$$\hat{\nabla}_a(\dot{f}) - (\hat{\nabla}_a f)^\cdot{}_\perp = H\hat{\nabla}_a f - \dot{f}\dot{u}_a, \quad (\text{E.4})$$

$$\hat{\nabla}_a(\dot{X}_b) - (\hat{\nabla}_a X_b)^\cdot{}_\perp = H\hat{\nabla}_a X_b, \quad (\text{E.5})$$

$$\hat{\nabla}_a(\dot{T}_{bc}) - (\hat{\nabla}_a T_{bc})^\cdot{}_\perp = H\hat{\nabla}_a T_{bc}. \quad (\text{E.6})$$

$$\ell\hat{\nabla}_a(\dot{f}) = (\ell\hat{\nabla}_a f)^\cdot{}_\perp - \ell\dot{f}\dot{u}_a, \quad (\text{E.7})$$

$$\ell\hat{\nabla}_a(\dot{X}_b) = (\ell\hat{\nabla}_a X_b)^\cdot{}_\perp, \quad (\text{E.8})$$

$$\ell\hat{\nabla}_a(\dot{T}_{bc}) = (\ell\hat{\nabla}_a T_{bc})^\cdot{}_\perp. \quad (\text{E.9})$$

$$\text{curl}(\dot{X}_a) - (\text{curl}(X_a))^\cdot = H\text{curl}(X_a), \quad (\text{E.10})$$

$$\text{curl}(\dot{T}_{ab}) - (\text{curl}(T_{ab}))^\cdot = H\text{curl}(T_{ab}). \quad (\text{E.11})$$

$$\ell\text{curl}(\dot{X}_a) = (\ell\text{curl}(X_a))^\cdot{}_\perp, \quad (\text{E.12})$$

$$\ell\text{curl}(\dot{T}_{ab}) = (\ell\text{curl}(T_{ab}))^\cdot{}_\perp. \quad (\text{E.13})$$

$$\hat{\nabla}^2(\dot{X}_a) = (\hat{\nabla}^2 X_a)^\cdot{}_\perp + 2H\hat{\nabla}^2 X_a, \quad (\text{E.14})$$

$$\hat{\nabla}^2(\dot{T}_{ab}) = (\hat{\nabla}^2 T_{ab})^\cdot{}_\perp + 2H\hat{\nabla}^2 T_{ab}. \quad (\text{E.15})$$

$$\ell^2\hat{\nabla}^2(\dot{X}_a) = (\ell^2\hat{\nabla}^2 X_a)^\cdot{}_\perp, \quad (\text{E.16})$$

$$\ell^2\hat{\nabla}^2(\dot{T}_{ab}) = (\ell^2\hat{\nabla}^2 T_{ab})^\cdot{}_\perp. \quad (\text{E.17})$$

Laplacian identities: The following linearized identities are used when interchanging the Laplacian operator $\hat{\nabla}^2$ with $\hat{\nabla}$ or $\text{curl}(\cdot)$:

$$\ell^2\hat{\nabla}^a\hat{\nabla}^2 f = \ell^2\hat{\nabla}^2\hat{\nabla}^a f - 2K\hat{\nabla}^a f - 2\dot{f}\text{curl}(\omega_a), \quad (\text{E.18})$$

$$\ell^2\hat{\nabla}^a\hat{\nabla}^2 X_b = \ell^2\hat{\nabla}^2\hat{\nabla}^a X_b - 2K[\hat{\nabla}^a X_b + \hat{\nabla}^b X_a - h_b^a\hat{\nabla}^c X_c], \quad (\text{E.19})$$

$$\ell^2\hat{\nabla}^a\hat{\nabla}^2 T_{bc} = \ell^2\hat{\nabla}^2\hat{\nabla}^a T_{bc} - 2K\left[\hat{\nabla}^a T_{bc} + \hat{\nabla}_b T_c^a + \hat{\nabla}_c T_b^a - h_b^a\hat{\nabla}^s T_{sc} - h_c^a\hat{\nabla}^s T_{bs}\right], \quad (\text{E.20})$$

$$\ell^2 \hat{\nabla}^a \hat{\nabla}^2 X_a = \ell^2 \hat{\nabla}^2 \hat{\nabla}^a X_a + 2K \hat{\nabla}^a X_a, \quad (\text{E.21})$$

$$\ell^2 \hat{\nabla}^b \hat{\nabla}^2 T_{ab} = \ell^2 \hat{\nabla}^2 \hat{\nabla}^b T_{ab} + 4K \hat{\nabla}^b T_{ab}, \quad (\text{E.22})$$

$$\hat{\nabla}^2 \text{curl} X_a = \text{curl}(\hat{\nabla}^2 X_a), \quad (\text{E.23})$$

$$\hat{\nabla}^2 \text{curl} T_{ab} = \text{curl}(\hat{\nabla}^2 T_{ab}). \quad (\text{E.24})$$

Curl identities: Finally, the following linearized identities are used when interchanging the curl operator $\text{curl}(\cdot)$ with $\hat{\nabla}$:

$$\text{curl}(\hat{\nabla}_a f) = -2\omega_a \dot{f}, \quad (\text{E.25})$$

$$\text{curl}(\hat{\nabla}_a X_b) = \hat{\nabla}_{(a} \text{curl}(X_b), \quad (\text{E.26})$$

$$\text{curl}(\hat{\nabla}_{(a} X_b) = \frac{1}{2} \hat{\nabla}_{(a} \text{curl}(X_b), \quad (\text{E.27})$$

$$\hat{\nabla}^a \text{curl}(X_a) = 0, \quad (\text{E.28})$$

$$\text{curl}(\hat{\nabla}^b T_{ab}) = 2\hat{\nabla}^b \text{curl}(T_{ab}), \quad (\text{E.29})$$

Further, we obtain the following linearized identities for the curl of a curl, in the linearized context:

$$\ell^2 \text{curl} \text{curl} X_a = -\ell^2 \hat{\nabla}^2 X_a + \ell^2 \hat{\nabla}_a \hat{\nabla}^b X_b + 2K X_a, \quad (\text{E.30})$$

$$\ell^2 \text{curl} \text{curl} T_{ab} = -\ell^2 \hat{\nabla}^2 T_{ab} + \frac{3}{2} \ell^2 \hat{\nabla}_{(a} \hat{\nabla}^c T_{b)c} + 3K T_{ab}, \quad (\text{E.31})$$

COVARIANT HARMONICS

In this appendix, we present an overview of general identities and results related to the covariant harmonics. We extend the results of Bruni et al. (1992) Appendix B by introducing the notion of associated vector and tensor harmonics.

As in Hawking (1966), we define the *spatial harmonics* $Q_{ab\dots c}^{(n)}$ as eigenfunctions of the spatial Laplace-Beltrami operator:

$$\hat{\nabla}^2 Q_{ab\dots c}^{(k)} + \frac{k^2}{\ell^2} Q_{ab\dots c}^{(k)} = 0, \quad (\text{F.1})$$

where k is known as the *wavenumber* of the harmonic. Further, using (F.1) we impose that the $Q^{(k)}$ quantities are constant in time:

$$\dot{Q}_{ab\dots c}^{(k)} = 0. \quad (\text{F.2})$$

See Bruni et al. (1992, Appendix B) for the relation between the metric harmonics Y and the covariant harmonics Q . Hereafter, as we are working in the geometrical formalism, the use of the term *harmonics* will refer to the covariant harmonics defined by (F.1) and (F.2). For the problem at hand, we only require scalar harmonics, vector harmonics and two-index tensor harmonics (which, for purposes of brevity will be referred to as tensor harmonics).

We briefly discuss notation for harmonics and quantities derived from harmonics. The 0 superscript of the harmonic quantity $Q^{(0,k)}$ denotes that it is a scalar harmonic. Similarly, we use 1 and 2 to denote vector and tensor harmonics, respectively. It is common in the literature to drop the wavenumber k when writing a harmonic quantity, so writing $Q^{(0,k)}$ as $Q^{(0)}$ instead. The wavenumber of the harmonic being used is instead implicitly defined by the equation where the harmonic appears. Further, here and elsewhere, we shall use Q to denote the whole set of harmonics and the standard quantities constructed from Q : we have

scalar harmonics $Q^{(0)}$, $Q_a^{(0)}$, $Q_{ab}^{(0)}$, vector harmonics $Q_a^{(1)}$, $Q_{ab}^{(1)}$ and tensor harmonics $Q_{ab}^{(2)}$.

We now give the definitions and properties of each of these three classes of harmonic types.

F.1 Scalar Harmonics

Scalar harmonics are defined as solutions of

$$\hat{\nabla}^2 Q = -\frac{k^2}{\ell^2} Q, \quad (\text{F.3})$$

that satisfy the time-independence constraint

$$\dot{Q} = 0. \quad (\text{F.4})$$

Each scalar harmonics Q defines a vector and a trace-free symmetric tensor via

$$Q_a = -\frac{\ell}{k} \hat{\nabla}_a Q, \quad (\text{F.5})$$

and

$$Q_{ab} = \frac{\ell^2}{k^2} \hat{\nabla}_b \hat{\nabla}_a Q + \frac{1}{3} h_{ab} Q. \quad (\text{F.6})$$

Since it follows from (F.4) and (B.1) that Q satisfies

$$\hat{\nabla}_{[a} \hat{\nabla}_{b]} Q = 0, \quad (\text{F.7})$$

it can be quickly shown that Q_a and Q_{ab} satisfy the curl-free constraint,

$$\text{curl}(Q_a) = 0, \quad \text{curl}(Q_{ab}) = 0, \quad (\text{F.8})$$

congruence-orthogonality,

$$u^a Q_a = 0, \quad u^a Q_{ab} = 0, \quad u^b Q_{ab} = 0, \quad (\text{F.9})$$

and time invariance,

$$\dot{Q}_a = 0, \quad \dot{Q}_{ab} = 0. \quad (\text{F.10})$$

We are restricted in our choice of k^2 to the eigenvalues determined by Lifshitz (1946), which in turn are determined by the spatial curvature according to

$$k^2 = n^2 - K \quad \begin{cases} n > 0 & \text{if } K = -1, 0, \\ n = 3, 4, \dots & \text{if } K = +1. \end{cases} \quad (\text{F.11})$$

We note that it is common to include $n = 2$ as a possibly harmonic mode in the case of $K = +1$, but it can be shown via the constraint equation (4.29) and identity (F.16) that this mode is unphysical. In fact, this mode is gauge-dependent and associated with the freedom of choice of the velocity field u^a .

Some important identities involving the vector quantity Q_a can be derived using the differential identities in Appendix E and (F.3)-(F.5):

$$\ell \hat{\nabla}_a Q^a = kQ, \quad (\text{F.12})$$

$$\ell^2 \hat{\nabla}^2 Q_a = -(k^2 - 2K)Q_a, \quad (\text{F.13})$$

$$\ell \hat{\nabla}_b Q_a = -k(Q_{ab} - \frac{1}{3}h_{ab}Q), \quad (\text{F.14})$$

$$\ell^2 \hat{\nabla}_{[a} \hat{\nabla}_{b]} Q_c = \frac{1}{2}K(h_{ac}Q_b - h_{bc}Q_a). \quad (\text{F.15})$$

Similarly, some important identities involving the tensor quantity Q_{ab} can be derived using the differential identities in appendix E and (F.3)-(F.6):

$$\ell \hat{\nabla}^b Q_{ab} = \frac{2}{3}k^{-1}(k^2 - 3K)Q_a, \quad (\text{F.16})$$

$$\ell^2 \hat{\nabla}^a \hat{\nabla}^b Q_{ab} = \frac{2}{3}(k^2 - 3K)Q, \quad (\text{F.17})$$

$$\ell^2 \hat{\nabla}_b \hat{\nabla}^c Q_{ac} = -\frac{2}{3}(k^2 - 3K)(Q_{ab} - \frac{1}{3}h_{ab}Q), \quad (\text{F.18})$$

$$\ell^2 \hat{\nabla}^2 Q_{ab} = -(k^2 - 6K)Q_{ab}, \quad (\text{F.19})$$

$$\ell^2 \hat{\nabla}_{[a} \hat{\nabla}_{b]} Q_{cd} = \frac{1}{2}K[(h_{ac}Q_{bd} - h_{bc}Q_{ad}) + (h_{ad}Q_{bc} - h_{bd}Q_{ac})]. \quad (\text{F.20})$$

F.2 Vector Harmonics

Vector harmonics are defined as solutions of

$$\hat{\nabla}^2 Q_a = -\frac{k^2}{\ell^2}Q_a, \quad (\text{F.21})$$

where Q_a is divergence-free and time-independent,

$$\hat{\nabla}^a Q_a = 0, \quad \dot{Q}_a = 0. \quad (\text{F.22})$$

Every vector harmonic Q_a defines an *associated vector harmonic* P_a via

$$P_a = \ell \text{curl}(Q_a), \quad (\text{F.23})$$

which can be shown to satisfy (F.21) and (F.22) using (E.23), (E.28) and (E.12). On taking the curl of (F.23) and applying (F.21) and (E.30), we obtain Q_a in terms of P_a ,

$$Q_a = (k^2 + 2K)^{-1} \ell \operatorname{curl}(P_a). \quad (\text{F.24})$$

The vector harmonics Q and associated vector harmonics P then define trace-free symmetric tensor quantities via

$$Q_{ab} = -\frac{\ell}{k} \hat{\nabla}_{(a} Q_{b)}, \quad (\text{F.25})$$

and

$$P_{ab} = -\frac{\ell}{k} \hat{\nabla}_{(a} P_{b)}. \quad (\text{F.26})$$

These tensor-valued vector harmonics are defined in order to satisfy zero second divergence,

$$\hat{\nabla}^a \hat{\nabla}^b Q_{ab} = 0, \quad \hat{\nabla}^a \hat{\nabla}^b P_{ab} = 0, \quad (\text{F.27})$$

congruence orthogonality,

$$u^a Q_{ab} = 0, \quad u^b Q_{ab} = 0, \quad u^a P_{ab} = 0, \quad u^b P_{ab} = 0, \quad (\text{F.28})$$

and time invariance,

$$\dot{Q}_{ab} = 0, \quad \dot{P}_{ab} = 0. \quad (\text{F.29})$$

The tensor-valued vector harmonics are related using (F.25), (F.26) and (E.24), as follows:

$$\ell \operatorname{curl}(P_{ab}) = \frac{1}{2}(k^2 + 2K)Q_{ab}, \quad (\text{F.30})$$

$$\ell \operatorname{curl}(Q_{ab}) = \frac{1}{2}P_{ab}. \quad (\text{F.31})$$

As with scalar harmonics, we are restricted in our choice of k^2 to the eigenvalues determined by Lifshitz (1946), according to

$$k^2 = n^2 - 2K \quad \begin{cases} n > 0 & \text{if } K = -1, 0 \\ n = 3, 4, \dots & \text{if } K = +1 \end{cases}. \quad (\text{F.32})$$

Again, it is common to include $n = 2$ as a possibly harmonic mode in the case of $K = +1$, but it can be shown via the constraint equation (A.21) that this mode is unphysical.

Using the differential identities in Appendix E and (F.21)-(F.23) we can derive the following identity for the vector harmonics Q_a and associated vector harmonics P_a (in this case, we may replace Q_a with P_a to obtain a second identity):

$$\ell^2 \hat{\nabla}_{[a} \hat{\nabla}_{b]} Q_c = \frac{1}{2} K (h_{ac} Q_b - h_{bc} Q_a). \quad (\text{F.33})$$

Some identities involving the tensor-valued vector harmonics Q_{ab} and associated tensor-valued vector harmonics P_{ab} can be derived using the differential identities in appendix E and (F.21)-(F.26). In each of the following identities, the pair (Q_{ab}, Q_a) may be replaced with (P_{ab}, P_a) :

$$\ell \hat{\nabla}^b Q_{ab} = \frac{1}{2} k^{-1} (k^2 - 2K) Q_a, \quad (\text{F.34})$$

$$\ell^2 \hat{\nabla}_{(a} \hat{\nabla}^c Q_{a)c} = -\frac{1}{2} (k^2 - 2K) Q_{ab}, \quad (\text{F.35})$$

$$\ell^2 \hat{\nabla}^2 Q_{ab} = -(k^2 - 4K) Q_{ab}, \quad (\text{F.36})$$

$$\ell^2 \hat{\nabla}_{[a} \hat{\nabla}_{b]} Q_{cd} = \frac{1}{2} K [(h_{ac} Q_{bd} - h_{bc} Q_{ad}) + (h_{ad} Q_{bc} - h_{bd} Q_{ac})]. \quad (\text{F.37})$$

F.3 Tensor Harmonics

Tensor harmonics are defined as solutions of

$$\hat{\nabla}^2 Q_{ab} = -\frac{k^2}{\ell^2} Q_{ab}, \quad (\text{F.38})$$

where Q_{ab} is divergence-free, trace-free and time independent:

$$\hat{\nabla}^b Q_{ab} = 0, \quad Q^a_a = 0, \quad \dot{Q}_{ab} = 0. \quad (\text{F.39})$$

As with vector harmonics, every tensor harmonic Q_{ab} defines an *associated tensor harmonic* P_{ab} via

$$P_{ab} = \ell \text{curl}(Q_{ab}). \quad (\text{F.40})$$

It can be shown that P_{ab} satisfies the definition of a tensor harmonic (F.38)-(F.39) using (E.24), (E.29) and (E.13). It then follows immediately from (F.38) that the quantities Q_{ab} and P_{ab} satisfy congruence orthogonality:

$$u^a Q_{ab} = 0, \quad u^b Q_{ab} = 0, \quad u^a P_{ab} = 0, \quad u^b P_{ab} = 0. \quad (\text{F.41})$$

Upon taking the curl of (F.40) and applying (F.38) and (E.31), we obtain Q_{ab} in terms of P_{ab} :

$$Q_{ab} = (k^2 + 3K)^{-1} \ell \text{curl}(P_{ab}). \quad (\text{F.42})$$

As with scalar and vector harmonics, we are restricted in our choice of k^2 to the eigenvalues determined by Lifshitz (1946):

$$k^2 = n^2 - 3K \quad \begin{cases} n > 0 & \text{if } K = -1, 0 \\ n = 3, 4, \dots & \text{if } K = +1 \end{cases} . \quad (\text{F.43})$$

We can obtain an important identity for Q_{ab} and P_{ab} using (B.6) and (E.1), as follows (the identity for P_{ab} is identical, except with Q_{ab} replaced with P_{ab}):

$$\ell^2 \hat{\nabla}_{[a} \hat{\nabla}_{b]} Q_{cd} = \frac{1}{2} K [(h_{ac} Q_{bd} - h_{bc} Q_{ad}) + (h_{ad} Q_{bc} - h_{bd} Q_{ac})] . \quad (\text{F.44})$$

THE 3-COTTON-YORK TENSOR

In this appendix we introduce the 3-Cotton-York tensor, which is useful in the analysis of tensor perturbations. Specifically, we are interested in using the 3-Cotton-York tensor in place of the co-moving shear \mathcal{X}_{ab} in developing a governing DE for tensor perturbations. It is of interest in this regard due to its relation with Bardeen's metric potential $H_T^{(2)}$. We will further show that the coefficients of the shear and Weyl tensor components can be rewritten in terms of $\mathcal{C}_{(k)}$. To the best of the author's knowledge, this analysis of tensor perturbations in terms of the 3-Cotton-York tensor is new.

In terms of the trace-free spatial Ricci tensor (1.40), the *3-Cotton-York tensor* is defined as (see, for example, van Elst and Uggla (1997, p2680-1))

$${}^{(3)}C_{ab} = \text{curl}({}^{(3)}S_{ab}), \quad (\text{G.1})$$

and hence is symmetric and tracefree. In the linearized context, we can use (B.10) and (4.28) to obtain an alternative first-order expression for ${}^{(3)}C_{ab}$ in terms of the Weyl tensor components, namely

$${}^{(3)}C_{ab} = -HH_{ab} + \text{curl}(E_{ab}). \quad (\text{G.2})$$

We note that an important property of this quantity is that it has zero spatial divergence in the linear theory. Indeed, upon taking the divergence of (G.2) and applying (4.31), (4.29), (4.30) and (E.29), we obtain, to first order,

$$\hat{\nabla}^b {}^{(3)}C_{ab} = 0. \quad (\text{G.3})$$

The evolution equation for ${}^{(3)}C_{ab}$ then follows upon taking the time derivative of (G.2) and applying (4.11), (4.24), (4.25) and (E.11), giving

$${}^{(3)}\dot{C}_{ab} = -3H {}^{(3)}C_{ab} + \frac{2K}{\ell^2} H_{ab} - \hat{\nabla}^2 H_{ab} + 2\hat{\nabla}_{\langle a} \hat{\nabla}^c H_{b\rangle c}. \quad (\text{G.4})$$

G.1 Evolution Equation for the 3-Cotton-York Tensor

Instead of choosing the co-moving shear to be the basic quantity for tensor perturbations (as in section 4.6), we can instead choose the basic quantity to be the *dimensionless 3-Cotton-York tensor*, defined by

$$\mathcal{C}_{ab} = \ell^3 {}^{(3)}\mathcal{C}_{ab}. \quad (\text{G.5})$$

Upon recalling from (G.3) that the 3-Cotton-York tensor has zero divergence, it quickly follows that \mathcal{C}_{ab} also has zero divergence, *i.e.*

$$\hat{\nabla}^b \mathcal{C}_{ab} = 0. \quad (\text{G.6})$$

Hence, \mathcal{C}_{ab} can be expanded purely in terms of associated tensor harmonics, according to

$$\mathcal{C}_{ab} = \sum_k \mathcal{C}_{(k)} P_{ab}^{(2)}, \quad (\text{G.7})$$

The derivatives of \mathcal{C}_{ab} along the fundamental congruence are obtained by differentiating (G.7), giving

$$\dot{\mathcal{C}}_{\langle ab \rangle} = \sum_k \dot{\mathcal{C}}_{(k)} P_{ab}^{(2,k)}, \quad \ddot{\mathcal{C}}_{\langle ab \rangle} = \sum_k \ddot{\mathcal{C}}_{(k)} P_{ab}^{(2,k)}. \quad (\text{G.8})$$

Upon differentiating (G.5) and using (3.9) and (G.4) we obtain

$$\dot{\mathcal{C}}_{ab} = \ell^3 \left(\frac{2K}{\ell^2} - \hat{\nabla}^2 \right) H_{ab}. \quad (\text{G.9})$$

Differentiating again and applying (G.5) and (E.17) yields

$$\ddot{\mathcal{C}}_{ab} + 3H\dot{\mathcal{C}}_{ab} + \left(\frac{2K}{\ell^2} - \hat{\nabla}^2 \right) \mathcal{C}_{ab} = 0. \quad (\text{G.10})$$

We now expand each term of this equation in terms of tensor harmonics using (G.7) and (G.8), applying (F.38) to remove the Laplacian term, obtaining

$$\ddot{\mathcal{C}}_{(k)} + 3H\dot{\mathcal{C}}_{(k)} + \frac{1}{\ell^2} (k^2 + 2K) \mathcal{C}_{(k)} = 0. \quad (\text{G.11})$$

This is then the second-order DE governing the evolution of the 3-Cotton-York tensor. In terms of conformal time, this DE reads

$$\mathcal{C}''_{(k)} + 2H\ell\mathcal{C}'_{(k)} + (k^2 + 2K)\mathcal{C}_{(k)} = 0, \quad (\text{G.12})$$

which is identical to Bardeen's DE for the metric potential $H_T^{(2)}$, given in Bardeen (1980) equation (4.14).

G.2 Tensor Perturbations

We now show the relation between (G.11) and the governing DE for tensor perturbations (4.76). We note that it follows by (G.3) that the 3-Cotton-York tensor is purely tensorial, *i.e.* the scalar and vector contributions to ${}^{(3)}C_{(k)}$ are exactly zero. Expanding ${}^{(3)}C_{(k)}$ in terms of associated tensor harmonics yields

$${}^{(3)}C_{ab} = \sum_k {}^{(3)}C_{(k)} P_{ab}^{(2)}. \quad (\text{G.13})$$

The coefficients ${}^{(3)}C_{(k)}$ can then be determined in terms of $\mathcal{X}_{(k)}$ and the tensor harmonics via (G.2) upon applying (4.77), (4.80), (4.81) and (F.40). We obtain

$${}^{(3)}C_{(k)} \sim -\frac{1}{H^2 \ell^2} \left(\dot{\mathcal{X}}_{(k)} + 2H \mathcal{X}_{(k)} \right). \quad (\text{G.14})$$

If we expand (G.9) in terms of tensor harmonics using (4.77), (F.38) and (G.8), we obtain

$$\dot{\mathcal{C}}_{(k)} = \ell (k^2 + 2K) H_{(k)}, \quad (\text{G.15})$$

and hence using (4.81) that

$$\mathcal{X}_{(k)} = \frac{1}{k^2 + 2K} \ell \dot{\mathcal{C}}_{(k)}. \quad (\text{G.16})$$

Now using (G.16), we can rewrite (4.79)-(4.81) as

$$\sigma_{(k)} = \frac{1}{k^2 + 2K} \dot{\mathcal{C}}_{(k)}, \quad (\text{G.17})$$

$$E_{(k)} = -\frac{1}{k^2 + 2K} \left[\ddot{\mathcal{C}}_{(k)} + 2H \dot{\mathcal{C}}_{(k)} \right], \quad (\text{G.18})$$

$$H_{(k)} = \frac{1}{k^2 + 2K} \ell^{-1} \dot{\mathcal{C}}_{(k)}. \quad (\text{G.19})$$

Thus, we obtain the coefficients of the shear and Weyl tensor components in terms of the coefficients of the 3-Cotton-York tensor $\mathcal{C}_{(k)}$.

BESSEL AND ASSOCIATED LEGENDRE FUNCTIONS

In this appendix we present solutions of the Bessel DE, toroidal Legendre DE and conical Legendre DE, which are used in solving the governing DEs for scalar and tensor perturbations. In section H.1 we present the normal forms of the Bessel and associated Legendre DEs and their general solutions in terms of Bessel and associated Legendre functions. As discussed in section H.2, these solutions can be written in terms of elementary functions in certain special cases. Further, resonant solutions of the Legendre DE appear in certain cases, as discussed in section H.3. Finally, the asymptotic behaviour of the general solutions is given in section H.4.

H.1 Normal Forms of Bessel and Associated Legendre DEs

In this section we present the normal forms of Bessel's DE, the toroidal Legendre DE and the conical Legendre DE along with their general solution in terms of Bessel functions and associated Legendre functions. Formally, the toroidal and conical Legendre functions are defined as the composition of the associated Legendre functions with \cosh and \cos , respectively.

Bessel's DE Normal Form:

$$\frac{d^2 X}{dT^2} + \left[a^2 - \frac{(\mu^2 - \frac{1}{4})}{T^2} \right] X = 0. \quad (\text{H.1})$$

Solution:

$$X(T) = T^{\frac{1}{2}} [c_1 J_\mu(aT) + c_2 Y_\mu(aT)]. \quad (\text{H.2})$$

Legendre DE (Toroidal) Normal Form:

$$\frac{d^2 X}{dT^2} + \left[-\left(\nu + \frac{1}{2}\right)^2 - \frac{(\mu^2 - \frac{1}{4})}{\sinh^2 T} \right] X = 0. \quad (\text{H.3})$$

Solution:

$$X(T) = (\sinh T)^{\frac{1}{2}} [c_1 P_\nu^\mu(\cosh T) + c_2 Q_\nu^\mu(\cosh T)]. \quad (\text{H.4})$$

Legendre DE (Conical) Normal Form:

$$\frac{d^2 X}{dT^2} + \left[\left(\nu + \frac{1}{2}\right)^2 - \frac{(\mu^2 - \frac{1}{4})}{\sin^2 T} \right] X = 0. \quad (\text{H.5})$$

Solution:

$$X(T) = (\sin T)^{\frac{1}{2}} [c_1 P_\nu^\mu(\cos T) + c_2 Q_\nu^\mu(\cos T)]. \quad (\text{H.6})$$

H.2 Special Solutions of the Governing DEs

For certain values of γ , μ and ν , the Bessel functions and associated Legendre P functions reduce to closed-form trigonometric expressions. Further, in each of these cases we can derive a second linearly independent solution \tilde{Q} of the associated Legendre DEs (H.3) and (H.5) that is defined so as to reduce to $J_\mu(bz)$ in the limit $z \rightarrow 0$. This solution is a preferred choice, since it emphasizes the relation between solutions for $K = \pm 1$ and solutions with $K = 0$ in the early time limit. In this section, we present the first few instances of this behaviour and its importance in modelling perturbations with a given fluid background.

For brevity, we shall make use of the notation

$$C(z) = \cos(z), \quad S(z) = \sin(z), \quad (\text{H.7})$$

and use the tilde \sim to denote dependence up to a multiplicative constant.

$\mu = \frac{1}{2}, \nu = -\frac{1}{2} + \sqrt{K}b$ Describes evolution of $(\mathcal{C}, \gamma = \frac{4}{3})$.

$$z^{1/2} \cdot Y_\mu(bz) \sim C(bz), \quad (\text{H.8})$$

$$z^{1/2} \cdot J_\mu(bz) \sim S(bz), \quad (\text{H.9})$$

$$(\sinh z)^{1/2} \cdot P_\nu^\mu(\cosh z) \sim C(bz), \quad (\text{H.10})$$

$$(\sinh z)^{1/2} \cdot \tilde{Q}_\nu^\mu(\cosh z) \sim S(bz), \quad (\text{H.11})$$

$$(\sin z)^{1/2} \cdot P_\nu^\mu(\cos z) \sim C(bz), \quad (\text{H.12})$$

$$(\sin z)^{1/2} \cdot \tilde{Q}_\nu^\mu(\cos z) \sim S(bz). \quad (\text{H.13})$$

$\mu = \frac{3}{2}, \nu = -\frac{1}{2} + \sqrt{K}b$ Describes evolution of $(\mathcal{C}, \gamma = 1), (\mathcal{X}, \gamma = \frac{4}{3}), (\mathcal{D}, \gamma = \frac{4}{3})$.

$$z^{1/2} \cdot Y_\mu(bz) \sim z^{-1}C(bz) + bS(bz), \quad (\text{H.14})$$

$$z^{1/2} \cdot J_\mu(bz) \sim -z^{-1}S(bz) + bC(bz), \quad (\text{H.15})$$

$$(\sinh z)^{1/2} \cdot P_\nu^\mu(\cosh z) \sim (\coth z)C(bz) + bS(bz), \quad (\text{H.16})$$

$$(\sinh z)^{1/2} \cdot \tilde{Q}_\nu^\mu(\cosh z) \sim -(\coth z)S(bz) + bC(bz), \quad (\text{H.17})$$

$$(\sin z)^{1/2} \cdot P_\nu^\mu(\cos z) \sim (\cot z)C(bz) + bS(bz), \quad (\text{H.18})$$

$$(\sin z)^{1/2} \cdot \tilde{Q}_\nu^\mu(\cos z) \sim -(\cot z)S(bz) + bC(bz). \quad (\text{H.19})$$

$\mu = \frac{5}{2}, \nu = -\frac{1}{2} + \sqrt{K}b$ Describes evolution of $(\mathcal{C}, \gamma = \frac{1}{3}), (\mathcal{X}, \gamma = 1)$.

$$z^{1/2} \cdot Y_\mu(bz) \sim (b^2 - 3z^{-2})C(bz) - 3bz^{-1}S(bz), \quad (\text{H.20})$$

$$z^{1/2} \cdot J_\mu(bz) \sim (b^2 - 3z^{-2})S(bz) + 3bz^{-1}C(bz), \quad (\text{H.21})$$

$$(\sinh z)^{1/2} \cdot P_\nu^\mu(\cosh z) \sim (b^2 - 3\coth^2 z + 1)C(bz) - 3b(\coth z)S(bz), \quad (\text{H.22})$$

$$(\sinh z)^{1/2} \cdot \tilde{Q}_\nu^\mu(\cosh z) \sim (b^2 - 3\coth^2 z + 1)S(bz) + 3b(\coth z)C(bz), \quad (\text{H.23})$$

$$(\sin z)^{1/2} \cdot P_\nu^\mu(\cos z) \sim (b^2 - 3\cot^2 z - 1)C(bz) - 3b(\cot z)S(bz), \quad (\text{H.24})$$

$$(\sin z)^{1/2} \cdot \tilde{Q}_\nu^\mu(\cos z) \sim (b^2 - 3\cot^2 z - 1)S(bz) + 3b(\cot z)C(bz). \quad (\text{H.25})$$

H.3 Resonant Solutions of the Legendre DE

For certain special cases, the associated Legendre functions P and Q are no longer linearly independent (see Abramowitz and Stegun p333, eq. (8.1.8)). One such example is the case of scalar perturbations with a dust background ($\gamma = 1$), where a second solution must be determined through alternative means. We now give the first five examples of resonance in the Legendre DE and give the second resonant solution of the Legendre DE for each case, denoted by $R_{\nu}^{\mu}(z)$. The toroidal solutions are given below. Conical solutions are obtained by replacing hyperbolic functions with trigonometric functions.

$$\mu = \frac{1}{2}, \nu = -\frac{1}{2}$$

$$(\sinh z)^{1/2} \cdot P_{-1/2}^{1/2}(\cosh z) \sim 1, \quad (\text{H.26})$$

$$(\sinh z)^{1/2} \cdot R_{-1/2}^{1/2}(\cosh z) \sim z. \quad (\text{H.27})$$

$$\mu = \frac{3}{2}, \nu = -\frac{1}{2}$$

$$(\sinh z)^{1/2} \cdot P_{-1/2}^{3/2}(\cosh z) \sim \coth z, \quad (\text{H.28})$$

$$(\sinh z)^{1/2} \cdot R_{-1/2}^{3/2}(\cosh z) \sim z \coth z - 1. \quad (\text{H.29})$$

$$\mu = \frac{5}{2}, \nu = -\frac{1}{2}$$

$$(\sinh z)^{1/2} \cdot P_{-1/2}^{5/2}(\cosh z) \sim 2(\coth z)^2 + (\sinh z)^{-2}, \quad (\text{H.30})$$

$$(\sinh z)^{1/2} \cdot R_{-1/2}^{5/2}(\cosh z) \sim z(2(\coth z)^2 + (\sinh z)^{-2}) - 3(\coth z). \quad (\text{H.31})$$

$$\mu = \frac{3}{2}, \nu = \frac{1}{2}, -\frac{3}{2}$$

$$(\sinh z)^{1/2} \cdot P_{-3/2}^{3/2}(\cosh z) \sim (\sinh z)^{-1}, \quad (\text{H.32})$$

$$(\sinh z)^{1/2} \cdot R_{-3/2}^{3/2}(\cosh z) \sim z(\sinh z)^{-1} - \cosh z. \quad (\text{H.33})$$

$$\mu = \frac{5}{2}, \nu = \frac{1}{2}, -\frac{3}{2}$$

$$(\sinh z)^{1/2} \cdot P_{-3/2}^{5/2}(\cosh z) \sim (\coth z)(\sinh z)^{-1}, \quad (\text{H.34})$$

$$(\sinh z)^{1/2} \cdot R_{-3/2}^{5/2}(\cosh z) \sim 3z(\coth z)(\sinh z)^{-1} - ((\cosh z)^2 + 2)(\sinh z)^{-1}. \quad (\text{H.35})$$

H.4 Asymptotic Behaviour

We now present the asymptotic behaviour of the Bessel functions and associated Legendre functions, both for early times and late times.

Bessel Functions

$$J_u(t \rightarrow 0) \sim \begin{cases} t^{|u|} & u \text{ integer} \\ t^u & u \text{ noninteger} \end{cases} \quad Y_u(t \rightarrow 0) \sim \begin{cases} t^{-|u|} & u \neq 0 \\ \ln(t) & u = 0 \end{cases} \quad (\text{H.36})$$

$$J_u(t \rightarrow +\infty) \sim t^{-1/2} \sin(t + \phi) \quad Y_u(t \rightarrow +\infty) \sim t^{-1/2} \cos(t + \phi) \quad (\text{H.37})$$

Associated Legendre Functions ($t \rightarrow 1$)

$$P_\nu^\mu(t \rightarrow 1) \sim \begin{cases} 1 & \mu = 0 \\ (t-1)^{|\mu|/2} & \mu \text{ integer} \\ (t-1)^{-|\mu|/2} & \mu \text{ noninteger} \end{cases} \quad (\text{H.38})$$

$$Q_\nu^\mu(t \rightarrow 1) \sim \begin{cases} \ln(t-1) & \mu = 0 \\ (t-1)^{-|\mu|/2} & \mu \text{ integer} \\ (t-1)^{-|\mu|/2} & \mu \text{ noninteger} \end{cases} \quad (\text{H.39})$$

Toroidal Legendre Functions ($t \rightarrow \infty$)

For the case

$$\nu = -\frac{1}{2} + ib, \quad b > 0, \quad (\text{H.40})$$

the asymptotic behaviour of the toroidal Legendre functions is given by

$$P_\nu^\mu(\cosh(t \rightarrow \infty)) \sim \exp(-t/2) \cos(bt + \phi), \quad (\text{H.41})$$

$$Q_\nu^\mu(\cosh(t \rightarrow \infty)) \sim \exp(-t/2) \sin(bt + \phi), \quad (\text{H.42})$$

as $t \rightarrow \infty$.

Bibliography

- [1] Abramowitz, M. and Stegun, I. (1970). *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th ed.* New York: Dover Publications, Inc.
- [2] Bardeen, J.M. (1980). Gauge Invariant Cosmological Perturbations, *Phys. Rev. D* **22**, p1882-905.
- [3] Bruni, M., Dunsby, P.K.S. and Ellis, G.F.R. (1992). Cosmological perturbations and the physical meaning of gauge-invariant variables, *Astrophys. J.* **395**, 34-53.
- [4] Clifton, T. and Barrow, J.D. (2007). The Ups and Downs of Cyclic Universes, *Phys. Rev. D* **75**.
- [5] Coles, P. and Lucchin, F. (1995). *Cosmology: the Origin and Evolution of Cosmic Structure.* John Wiley and Sons.
- [6] Collins, C.B. and Wainwright, J. (1983). The role of shear in general-relativistic cosmological and stellar models, *Phys. Rev. D* **27**, 1209-18.
- [7] Coquereaux, R. and Grossmann, A. (1982). Analytic Discussion of Spatially Closed Friedman Universes with Cosmological Constant and Radiation Pressure, *Ann. of Phys.* **143**, 296-356.
- [8] Dabrowski, M. and Stelmach, J. (1986). Analytic Solutions of Friedman Equation for Spatially Opened Universes with Cosmological Constant and Radiation Pressure, *Ann. of Phys.* **166**, 422-42.
- [9] Demiański, M., Zdzisław, A.G. and Woszczyńska, A. (2005). Evolution of density perturbations in a realistic universe, *Gen. Relativ. Gravit.* **37**, 2063-2082.

-
- [10] Dunsby, P.K.S., Bassett B.A. and Ellis, G.F.R. (1997). Covariant analysis of gravitational waves in a cosmological context, *Class. Quan. Grav.* **14**, 1215-22.
- [11] Ehlers, J. and Rindler, W. (1989). A phase-space representation of Friedmann-Lemaître universes containing both dust and radiation and the inevitability of a big bang, *Mon. Not. R. Astr. Soc.* **238**, 503-21.
- [12] Einstein, A. and de Sitter, W. (1932). On the relation between the expansion and the mean density of the universe, *Proc. Natl. Acad. Sci. USA* **18**, 213-14.
- [13] Ellis, G.F.R. (1973). Relativistic Cosmology. In *Cargèse Lectures in Physics Physics, Volume 6*, ed. E. Schatzmann. Gordon and Breach.
- [14] Ellis, G.F.R. (1989). *The Expanding Universe: A History of Cosmology from 1917 to 1960*, in D. Howard and J. Stachel, ed., *Einstein and the History of General Relativity*, p367-431. Boston: The Center for Einstein Studies.
- [15] Ellis, G.F.R. and Bruni, M. (1989). Covariant and gauge-invariant approach to cosmological density fluctuations, *Phys. Rev. D* **40**, 1804-18.
- [16] Ellis, G.F.R., Hwang, J. and Bruni, M. (1989). Covariant and gauge-independent perfect-fluid Robertson-Walker perturbations, *Phys. Rev. D* **40**, 1819-26.
- [17] Ellis, G.F.R., Bruni, M. and Hwang, J. (1990). Density-gradient-vorticity relation in perfect-fluid Robertson-Walker perturbations, *Phys. Rev. D* **42**, 1035-46.
- [18] Ellis, G.F.R. and van Elst, H. (1998). Cosmological Models. In *Cargèse Lectures 1998*.
- [19] Flanders, H. (1989). *Differential Forms with Applications to the Physical Sciences*. New York: Dover Publications, Inc.
- [20] Goode, S.W. (1983). Spatially inhomogeneous cosmologies and their relation with the Friedmann-Robertson-Walker cosmologies, *PhD Thesis* University of Waterloo.
- [21] Goode, S.W. (1989). Analysis of spatially inhomogeneous perturbations of the FRW cosmologies, *Phys. Rev. D* **39**, 2882-92.
-

-
- [22] Hall, G.S. and Pulham, J.R. (1996). *General Relativity*. Bristol: Scottish Universities Summer School in Physics.
- [23] Harrison, E.R. (1967). Classification of Uniform Cosmological Models, *Mon. Not. R. Astron. Soc.* **137**, 69.
- [24] Hawking, S.W. (1966). Perturbations of an expanding universe, *Astrophys. J.* **145**, 544-54.
- [25] Hawking, S.W. and Ellis, G.F.R. (1973). *The Large Scale Structure of Space-Time*. Cambridge: Cambridge University Press.
- [26] Jantzen, R.T. and Uggla, C. (1992). Structure of the generalized Friedmann problem, *Gen. Rel. Grav.* **24**, 59-85.
- [27] Lake, K. (2005). *The Flatness Problem and Λ* . *Phys. Rev. Lett.* **94**, 201102.
- [28] Lake, K. (2006). *Integration of the Friedmann equation for universes of arbitrary complexity*. Preprint, gr-qc/0603028.
- [29] Lifshitz, M. (1946). On the Gravitational Stability of the Expanding Universe, *J. Phys (Moscow)* **10**, 116.
- [30] Lifshitz, M. and Khalatnikov, I. (1963). Investigations in Relativistic Cosmology, *Adv. Phys.* **12**, 185-249.
- [31] Lyth, D.H. and Mukherjee, M. (1988). Fluid flow description of density irregularities in the universe, *Phys. Rev. D* **38**, 485-9.
- [32] Krasinski, A. (1997). *Inhomogeneous Cosmological Models*. Cambridge: Cambridge University Press.
- [33] Maartens, R., Ellis, G.F.R. and Stoeger, W.R. (1995a). Limits on anisotropy and inhomogeneity from the cosmic background radiation, *Phys. Rev. D* **51**, 1525-35.
- [34] Maartens, R., Ellis, G.F.R. and Stoeger, W.R. (1995b). Improved limits on anisotropy and inhomogeneity from the cosmic background radiation, *Phys. Rev. D* **51**, 5942-45.
- [35] Maartens, R. (1997). Linearization instability of gravity waves? *Phys. Rev. D* **55**, 463-7.
-

-
- [36] Mukhanov, V. (2005). *Physical Foundations of Cosmology*. Cambridge: Cambridge University Press.
- [37] Peacock, J.A. (1999). *Cosmological Physics*. Cambridge: Cambridge University Press.
- [38] Perko, L. (1996). *Differential Equations and Dynamical Systems, 2nd ed.* New York: Springer-Verlag.
- [39] Ratra, B. and Peebles P.J.E. (1995). Inflation in an open universe, *Phys. Rev. D* **52**, 1837-94.
- [40] Rindler, W. (1977). *Essential Relativity: Special, General, and Cosmological, revised 2nd ed.* New York: Springer-Verlag.
- [41] Sachs, R.K. and Wolfe, A.M. (1967). Perturbations of a cosmological model and angular variations of the microwave background, *Astrophys. J.* **147**, 73-90.
- [42] Spergel, D.N. et al. (2006). Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology. Preprint, astro-ph/0603449.
- [43] Stephani, H. et al. (2003). *Exact Solutions to Einstein's Field Equations 2nd ed.* Cambridge: Cambridge University Press.
- [44] Stewart, J.M. Perturbations of Friedmann-Robertson-Walker cosmological models, *Class. Quan. Grav* **7**, 1169-80.
- [45] Stewart, J.M. and Walker, M. (1974). Perturbations of spacetimes in general relativity, *Proc. Roy. Soc. London A* **341**, 49-74. [Chapter 14]
- [46] Tegmark, M. et al. (2004). Cosmological parameters from SDSS and WMAP, *Phys. Rev. D* **69**, 103501.
- [47] Tolman, R.C. (1934). *Relativity Thermodynamics and Cosmology*. Oxford: Clarendon Press.
- [48] van Elst, H. and Uggla, C. (1997). General Relativistic 1 + 3 Orthonormal Frame Approach, *Class. Quan. Grav* **14**, 267395.
- [49] Wainwright, J. (1996) *Relativistic Cosmology*, in Hall, G.S. and Pulham, J.R. General Relativity. Great Britain: Scottish Universities Summer School in Physics, p107-41.
-

-
- [50] Wainwright, J. and Ellis, G.F.R. (1997). *Dynamical Systems in Cosmology*. Cambridge: Cambridge University Press.
- [51] Wald, R. (1984). *General Relativity*. Chicago and London: The University of Chicago Press.
- [52] Weinberg, S. (1972). *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, New York: Wiley.
-