



2014

Negative Urgency, Pubertal Onset and the Longitudinal Prediction of Alcohol Consumption During the Transition from Preadolescence to Adolescence

Lauren Helena Boyle

University of Kentucky, laurenhelena@gmail.com

[Click here to let us know how access to this document benefits you.](#)

Recommended Citation

Boyle, Lauren Helena, "Negative Urgency, Pubertal Onset and the Longitudinal Prediction of Alcohol Consumption During the Transition from Preadolescence to Adolescence" (2014). *Theses and Dissertations--Psychology*. 50.
https://uknowledge.uky.edu/psychology_etds/50

This Master's Thesis is brought to you for free and open access by the Psychology at UKnowledge. It has been accepted for inclusion in Theses and Dissertations--Psychology by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

STUDENT AGREEMENT:

I represent that my thesis or dissertation and abstract are my original work. Proper attribution has been given to all outside sources. I understand that I am solely responsible for obtaining any needed copyright permissions. I have obtained needed written permission statement(s) from the owner(s) of each third-party copyrighted matter to be included in my work, allowing electronic distribution (if such use is not permitted by the fair use doctrine) which will be submitted to UKnowledge as Additional File.

I hereby grant to The University of Kentucky and its agents the irrevocable, non-exclusive, and royalty-free license to archive and make accessible my work in whole or in part in all forms of media, now or hereafter known. I agree that the document mentioned above may be made available immediately for worldwide access unless an embargo applies.

I retain all other ownership rights to the copyright of my work. I also retain the right to use in future works (such as articles or books) all or part of my work. I understand that I am free to register the copyright to my work.

REVIEW, APPROVAL AND ACCEPTANCE

The document mentioned above has been reviewed and accepted by the student's advisor, on behalf of the advisory committee, and by the Director of Graduate Studies (DGS), on behalf of the program; we verify that this is the final, approved version of the student's thesis including all changes required by the advisory committee. The undersigned agree to abide by the statements above.

Lauren Helena Boyle, Student

Dr. Gregory T. Smith, Major Professor

Dr. Gregory T. Smith, Director of Graduate Studies

NEGATIVE URGENCY, PUBERTAL ONSET AND
THE LONGITUDINAL PREDICTION OF ALCOHOL
CONSUMPTION DURING THE TRANSITION FROM
PREADOLESCENCE TO ADOLESCENCE

THESIS

A thesis submitted in partial fulfillment of the
requirements for the degree of Master of Science in the
College of Arts and Sciences
at the University of Kentucky

By

Lauren Helena Boyle

Lexington, Kentucky

Director: Dr. Gregory T. Smith, Professor of Psychology

Lexington, Kentucky

2014

Copyright© Lauren Helena Boyle 2014

ABSTRACT OF THESIS

NEGATIVE URGENCY, PUBERTAL ONSET AND THE LONGITUDINAL PREDICTION OF ALCOHOL CONSUMPTION DURING THE TRANSITION FROM PREADOLESCENCE TO ADOLESCENCE

Alcohol use in early adolescence is associated with numerous concurrent and future problems, including diagnosable alcohol use disorders. The trait of negative urgency, the tendency to act rashly when distressed, is an important predictor of alcohol-related dysfunction in youth and adults. The aim of this study was to test a model proposed by Cyders and Smith (2008) specifying a puberty-based developmental increase in negative urgency, which in turn predicts subsequent increases in early adolescent drinking. In a sample of 1,910 youth assessed semi-annually from spring of 5th grade through spring of 8th grade, we found support for this model. Pubertal onset was associated with both a mean increase and subsequent rises in negative urgency over time. Drinking frequency at any wave was predicted by prior wave assessments of drinking frequency, negative urgency, and pubertal onset. The slope of increase in drinking also increased as a function of pubertal onset. This model applied to negative urgency but not to other impulsivity-related traits. These findings highlight the importance of personality change in early adolescence as part of the risk matrix for early onset alcohol consumption.

KEYWORDS: Personality, Longitudinal, Drinking, Negative Urgency, Pubertal Onset

Lauren Helena Boyle

June 30, 2014

NEGATIVE URGENCY, PUBERTAL ONSET AND
THE LONGITUDINAL PREDICTION OF ALCOHOL
CONSUMPTION DURING THE TRANSITION FROM
PREADOLESCENCE TO ADOLESCENCE

By

Lauren Helena Boyle

Dr. Gregory T. Smith

Director of Thesis & Director of
Graduate Studies

June 30, 2014

DEDICATION

*To mom and dad. Thank you for giving me strength, direction,
and inspiration. (And maybe a few loans).*

*To Nirvana and Alice in Chains, whose enjoyably depressing music
was my constant companion while I worked.*

To the memory of Judson Eli Wickham. Enjoy your orange, Jud.

TABLE OF CONTENTS

| | |
|--------------------------------------------------------------------------------------|----|
| List of Tables | iv |
| List of Figures | v |
| Chapter One: Introduction | 1 |
| Early Adolescent Drinking | 2 |
| Negative Urgency and Other Impulsivity-Related Traits..... | 3 |
| Adolescent Increases in Negative Urgency and the Role of Pubertal Onset..... | 4 |
| The Current Study | 6 |
| Chapter Two: Methods..... | 7 |
| Participants..... | 7 |
| Measures..... | 7 |
| Demographic and background questionnaire | 7 |
| The Pubertal Development Scale | 7 |
| The UPPS-P-Child Version..... | 8 |
| The Drinking Styles Questionnaire | 8 |
| Procedure | 9 |
| Data collection | 9 |
| Data Analytic Method..... | 10 |
| Chapter Three: Results | 13 |
| Participant Retention | 13 |
| Possible Effects due to School Membership | 13 |
| Descriptive Statistics | 13 |
| Correlations Among Traits, Behaviors and Pubertal Status Across Waves..... | 13 |
| Model Tests..... | 14 |
| Change in negative urgency across early adolescence..... | 14 |
| Change in drinking behavior | 14 |
| Negative urgency predicting change in drinking | 15 |
| Tests of Specificity of the Relationship Between Pubertal Onset and Negative Urgency | 15 |
| Chapter Four: Discussion..... | 20 |
| References | 23 |
| Vita | 29 |

LIST OF TABLES

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Table 3.1, Descriptive Statistics for Pubertal Statuorientations, Drinker Status, and Five Personality Trait Contributors to Rash Action | 17 |
| Table 3.2, Bivariate Correlations Among Study Variables in Waves 1 and 7..... | 18 |

LIST OF FIGURES

| | |
|----------------------------------------------------------------------|----|
| Figure 2.1, Figural Representation of the Hypothesized Process | 12 |
|----------------------------------------------------------------------|----|

Chapter One: Introduction

The aim of this study was to evaluate a model of one aspect of risk for early adolescent drinking behavior that integrates both biological and personality processes. Specifically, we examined the relationships between the biological event of pubertal onset and subsequent changes in high-risk personality traits, and between these trait changes and subsequent shifts in drinking behavior. In brief, the rationale for this research is as follows. Early adolescent alcohol consumption is significantly associated with current and future dysfunction, including diagnostic status and engagement in other high-risk behaviors (DeWit, Adlaf, Offord, & Ogborne, 2000; Grant & Dawson, 1997; Gruber, DiClemente, Anderson, & Lodico, 1996; Guo, Collins, Hill, & Hawkins, 2000; Guttmanova et al., 2012; SAMHSA, 2012). Furthermore, the number of youth who drink, and the frequency with which they do so, increases in a somewhat linear fashion across the adolescent years (Chung et al., 2012). One aspect of risk for early adolescent drinking involves personality, particularly personality traits that are associated with rash or impulsive action (Gunn & Smith, 2010; Settles, Zapolski, & Smith, 2014; Stautz & Cooper, 2013). Among the best trait predictors of early adolescent drinking is the trait of negative urgency (the tendency to act rashly when extremely upset: Gunn & Smith, 2010; Stautz & Cooper, 2013). Cyders and Smith (2008) argued that pubertal onset is associated with mean and slope increases in negative urgency during early adolescence. This theoretical contention has not previously been tested.

The aim of this paper was to test the Cyders and Smith (2008) theoretical contention that pubertal onset is associated with increases in negative urgency, and further that changes in negative urgency are associated with subsequent changes in drinking behavior. To introduce this empirical investigation, we first briefly review research indicating that early adolescent alcohol consumption has deleterious effects on subsequent functioning and quality of life. After introducing the trait of negative urgency

and considering related traits, we review the basis for the hypothesis that pubertal onset leads to increases in negative urgency in particular. We then present our model relating changes in negative urgency to subsequent changes in drinking behavior, and introduce the specifics of the current investigation.

Early Adolescent Drinking

A small but significant portion of youth have already consumed more than a few sips of alcohol prior to age 12, with estimates that approximately 9% or 10% of 11-year-olds have done so (Donovan, 2007; Gunn & Smith, 2010). From age 11 through the rest of adolescence, the prevalence of use within the past year rises steadily to 67% by age 18 (Chung et al., 2012). This early use, and the trend of increasing use, is quite important clinically. For both boys and girls age 12, reports of having consumed alcohol just one or more days in the preceding year had sensitivity of 1.0 and specificity of .95 (girls) or .94 (boys) in the concurrent prediction of any past-year DSM IV alcohol use disorder symptom (Chung et al., 2012). The success of this indicator as a predictor of concurrent problems was not limited to age 12; it provided the best sensitivity and specificity through age 15 for both sexes.

Alcohol consumption during the middle school years is also associated with several other problem behaviors, including early onset marijuana use, early sexual intercourse, and low value on academic achievement (Jessor, 1987). In addition, early consumption is a significant predictor of diagnostic status and alcohol problems in later adolescence and adulthood (DeWit, Adlaf, Offord, & Ogborne, 2000; Grant & Dawson, 1997; Gruber et al., 1996; Guo et al., 2000; Guttmanova et al., 2012). Further, among adolescents in treatment for alcohol and substance use disorders, the majority report first using substances between the ages of 12 and 14 (SAMHSA, 2012). Alcohol consumption at this early age is therefore not only a good marker of current dysfunction, but also of risk for significant future dysfunction.

Negative Urgency and Other Impulsivity-Related Traits

Negative urgency is understood to reflect individual differences in the disposition to act in rash, impulsive ways when distressed (Cyders & Smith, 2008; Whiteside & Lynam, 2001). Urgency theory holds that individuals high in the trait are inclined to respond to distress with immediate action that is designed to distract from, or reduce, the distress (Cyders & Smith, 2008). Such actions are chosen because they provide the immediate benefit of relief from distress, not because they help meet long-term needs and goals. Often, such actions can be described as rash or impulsive. Examples of behaviors that serve the negative reinforcement of relief from distress that are predicted prospectively by negative urgency include alcohol consumption (Settles, Cyders, & Smith, 2010), smoking (Doran, Khoddam, Sanders, Schweizer, Trim, & Myers, 2013), and binge eating and purging (Fischer, Peterson, & McCarthy, 2013; Pearson, Combs, Zapolski, & Smith, 2012). Concerning alcohol consumption, meta-analyses of early adolescents highlight the predictive role of negative urgency in drinking behavior (Stautz & Cooper, 2013), and meta-analyses of late adolescents and adults find that negative urgency predicts heavy quantity of consumption and problem drinking behavior (Coskunpinar, Dir, & Cyders, 2013).

Negative urgency concerns a disposition toward emotion-driven rash action, and is thus different from most other impulsivity-related personality traits. For example, sensation seeking refers to a need to seek stimulation, not to a tendency to act when emotional. Similarly, neither lack of planning (acting without forethought) nor lack of perseverance (inability to maintain a focus on a task when faced with potential distraction) are emotion-based (Cyders & Smith, 2008; Whiteside & Lynam, 2001). Negative urgency shares very little variance with these other three traits (Cyders & Smith, 2007; Smith et al., 2007). Negative urgency is highly related to positive urgency, which is the tendency to act rashly when in an unusually positive mood; indeed, the two

traits can be considered facets of an overall urgency trait domain (Cyders & Smith, 2007). Negative and positive urgency, though highly correlated, play different predictive roles for some behaviors. Negative urgency predicts binge eating and purging, but positive urgency does not (Cyders et al., 2007; Pearson et al., 2012; Pearson, Zapolski, & Smith, 2014). More broadly, in prospective research, negative urgency predicts increases in negative mood-based rash action and positive urgency predicts increases in positive mood-based rash action (Cyders & Smith, 2010). Positive urgency does not appear to operate via a negative reinforcement mechanism.

Thus, among this set of five traits, negative urgency is one of two traits that is emotion-based and the only trait that (a) focuses on negative affect-driven rash action and (b) appears to facilitate actions that provide immediate negative reinforcement (in the form of distress relief). See Cyders and Smith (2007, 2008) and Whiteside and Lynam (2001) for the model specifying these five traits; Smith and Guller (in press) provide a discussion of other, similar models.

Adolescent Increases in Negative Urgency and the role of Pubertal Onset

There is a great deal of evidence that individual differences in personality are relatively stable over the life span (Caspi & Roberts, 2001; McCrae & Costa, 1994; Roberts & DelVecchio, 2000). However, it is also true that the greatest stability occurs beginning around age 30; prior to age 30, personality does change to a modest degree over time, particularly during periods of important developmental transitions (Bleidorn, 2012; Neyer & Asendorpf, 2001; Roberts, 2006). Bleidorn (2012) documented personality change over just six months in youth making the transition from high school to adult life. Roberts, Caspi, and Moffitt (2001) note that personality traits appear to be particularly malleable at younger ages, such as the transition into middle school from elementary school. This thesis is designed to test the theoretical position that the post-pubertal, early adolescent years are a period during which the trait of negative urgency

does change. Specifically, we believe there are two factors that contribute to increases in negative urgency during the early adolescent years.

The first factor is initial levels of negative urgency, which can be reliably and validly assessed among pre-adolescents (Gunn & Smith, 2010; Zapolski, Stairs, Settles, Combs, & Smith, 2010). Youth high in negative urgency tend to respond to distress with immediate action, designed to provide prompt negative reinforcement in the form of relief or distraction from their distress (Cyders & Smith, 2008). Each time an individual engages in such actions, which provide immediate relief, the chosen behavior is reinforced. Over time, negative urgency – based rash actions become increasingly likely. At the same time, on each such occasion the individual is not learning other, more adaptive, long-term responses to distress. Thus, rash actions become increasingly likely and problem-solving skills do not develop in the same way as they do in other youth. In this way, the trait is reinforced and becomes stronger over time (Pearson, Guller, Spillane, & Smith, 2011).

The second proposed influence on increases in negative urgency is the experience of pubertal onset. One characteristic of brain development following pubertal onset is that development of motivational and emotional subcortical connections occurs faster than development of connections that support prefrontal cortex (PFC) activity (Casey & Caudle, 2013). This imbalance in rate of development is important because certain aspects of PFC activity modulate emotional reactivity (Maxwell & Davidson, 2007). One result of this imbalance is that adolescents, compared to children and adults, rely more on motivational/emotional subcortical brain activity to guide behavior. Thus, when teens are experiencing intense emotion, they are more likely than children or adults to act in rash or impulsive ways (Casey & Caudle, 2013; Spear, 2001). Given the heightened negative affect associated with adolescence (Brooks-Gunn et al., 1994;

Compas et al., 1995; Nelson et al., 2002), post-pubertal teens appear to experience an increase in negative affect-driven rash action, i.e., an increase in negative urgency.

The Current Study

We tested the following model. Pubertal onset occurs at different times for different youth. When it occurs, it is expected to predict subsequent increases in both the mean and the rate of increase in negative urgency across early adolescence. Because negative urgency is a risk factor for early adolescent drinking, the mean and slope increases in this trait will predict subsequent mean and slope increases in adolescent drinking behavior. Our expectation was that this puberty-based risk process was specific to negative urgency and thus would not operate for positive urgency (because of the absence of negative reinforcement), sensation seeking, lack of planning, or lack of perseverance. To test this model, we studied all five traits, pubertal status, and drinking behavior in adolescents ($n = 1910$) longitudinally across the developmental period from the end of elementary school through the end of middle school.

Chapter Two: Methods

Participants

A total of 1910 children completed questionnaires in seven waves from the spring of 5th grade (the last year of elementary school in the participating school districts) through the spring of 8th grade (the last year of middle school), collected every six months. Participants in the study consisted of students from urban, rural, and suburban backgrounds, all from public school systems. The sample was approximately equally divided between boys ($n = 972$) and girls ($n = 938$). The breakdown of students by ethnicity was as follows: 61.6% European American, 17.0% African American, 6.9% Hispanic/Latino, 3% Asian American, and 11.5% of students reporting other ethnic backgrounds. We oversampled for African American and Hispanic children, and the sample represents those groups at a higher rate than is characteristic of the local population. The majority of the 5th graders, 66.8%, were 11 years of age; 22.8% were 10 years of age, 10% were age 12, and both 9 and 13-year-olds made up 0.2% of the participants.

Measures

Demographic and background questionnaire. This brief measure provided demographic information including age, grade level, gender, and ethnicity. Participants were asked to hand-write their current age and grade level, circle their sex and indicate which label(s) best described their ethnic background.

The Pubertal Development Scale (PDS, Petersen, Crockett, Richards, & Boxer, 1988). This self-report scale ascertained pubertal development status for participants, and is scored on a likert scale. Questions differed for boys and girls, and assessed information such as body hair, menstruation, and breast development, depending on subject gender. There were four questions for boys (“Do you have facial hair yet?”) and five questions for girls (“Have you begun to have your period?”).

Internal consistency ratings of the PDS show Cronbach's α 's ranging from .67 to .76 for 11-year-olds (Brooks-Gunn, Warren, Rosso, & Gargiulo, 1987). Inter-rater reliability estimates show good agreement with the gold standard – a physician administered assessment of pubertal status using Tanner ratings – with r values ranging from .61 to .67 (Brooks-Gunn et al., 1987; Coleman & Coleman, 2002). As is frequently practiced, we dichotomized pubertal status, with a mean score of 2.5 demarcating pre- and post-pubertal onset (Culbert, Burt, McGue, Iacono, & Klump, 2009).

The UPPS-P-Child Version (Zapolski et al., 2010). This self-report measure was used to measure the five impulsivity-related traits. The measure uses a four-point likert-type scale, with answer choices spanning from “not at all like me” to “very much like me.” Sample items and wave 1 internal consistency estimates for each scale are the following (for each scale, internal consistency estimates increased after wave 1). For negative urgency, a sample item is “When I am upset I often act without thinking” ($\alpha = .85$). For positive urgency, “When I am very happy, I tend to do things that may cause problems in my life” ($\alpha = .89$). For sensation seeking, “I like new, thrilling things, even if they are a little scary” ($\alpha = .79$). For lack of planning, “I tend to blurt out things without thinking” ($\alpha = .77$). For lack of perseverance, “Once I get going on something, I hate to stop,” which is reverse scored ($\alpha = .65$).

The Drinking Styles Questionnaire (Smith, McCarthy, & Goldman, 1995). This self-report measure was used to assess drinking behavior. We measured drinking frequency because it has been shown to be the best predictor of alcohol use disorder symptoms in youth this young (Chung et al., 2012). We used a single item, asking “Which of the following best describes how often you drink alcohol?” Response options are: “I have never had a drink of alcohol,” “I have only had 1, 2, 3, or 4 drinks of alcohol in my life,” “I only drink alcohol 3 or 4 times a year,” “I drink alcohol about once a month,” “I drink alcohol once or twice a week,” and “I drink alcohol almost daily.” The

single item measure has proven to be stable over time (Smith et al., 1995), and there is good evidence for its validity (Gunn & Smith, 2010; Settles, Fischer, Cyders, Combs, Gunn, & Smith, 2012).

Procedure

Data collection. At wave 1, the questionnaires were administered in 23 public elementary schools during school hours. In subsequent waves, the measures were administered in 15 public middle schools, again during school hours (in addition, children who had moved completed the measures by mail). We used a passive consent procedure. The families of each of 1,988 5th grade children in each of the 23 participating elementary schools were sent a letter introducing the study through the U.S. Mail. Families were asked to return an enclosed, stamped letter or call a phone number if they did not want their child to participate. At each wave of the study, participants completed active assent forms for their participation. Out of 1,988 5th graders in the participating schools, 1910 participated in the study (95.8%). A total of 78 students did not participate due to one of the following reasons: families declined participation for their child; students declined assent; or a variety of other reasons, such as language disabilities that precluded completing the questionnaires.

Data collections took place in the spring of 5th grade, and then in the fall and spring of 6th, 7th, and 8th grades, for a total of 7 data collection waves. Questionnaires were administered in classrooms, the school cafeteria, or the school library. Participants were assured that their responses would be kept strictly confidential: Neither teachers nor parents would have access to their responses. Researchers introduced the federal certificate of confidentiality to the participants and noted that the certificate legally protected the privacy of their responses. The procedure took 60 minutes or less.

Participants who moved during the course of the study were contacted and invited to continue to participate. Those who agreed did so by mail or by secure website.

Participants assessed during school hours were not paid for their participation, but those who had moved were paid \$30 per wave.

Data analytic method. During the first stage of analysis, we calculated the overall frequency of drinking behavior and pubertal onset at each wave. We also calculated the bivariate correlations among pubertal onset, drinking behavior and the five personality traits.

To test our theoretical model, we used a multilevel modeling approach in a series of steps. This method provided several advantages. First, we were able to model both within-person change over time and between-person variability in growth patterns. Pubertal onset occurs at different times for different youth, so we modeled it as a time-varying predictor. In the first step, we tested whether, within persons, negative urgency changed as a function of time, pubertal onset, and time since pubertal onset. Between persons, we tested whether negative urgency differed by sex. In the second step, we tested whether, within persons, pubertal onset was associated with increases in the mean and slope of drinking behavior. In the third step, we began with the model used in the first step and added drinking frequency measured both concurrently with negative urgency and one wave (6 months) later than negative urgency. This step enabled us to test our hypothesis that, within persons, change in negative urgency would predict a mean increase in subsequent drinking behavior (controlling for prior drinking behavior) as well as a slope increase in subsequent drinking behavior (again controlled for prior drinking behavior). We also tested whether sex moderated any of the predictive effects.

To determine if our model of personality and subsequent drinking change was specific to negative urgency, we ran the same set of models four additional times, replacing negative urgency in the model with positive urgency, sensation seeking, lack of planning, and lack of perseverance. We used Mplus (Muthén & Muthén, 1998-2010) for these analyses.

The process of modeling change as a function of a time-varying predictor can be depicted in equation form. We begin with an equation reflecting general, linear change in a trait as a function of time (for simplicity, leaving out additional predictors):

$$Y_{ij} = \pi_{0i} + \pi_{1i} \text{TIME}_{ij} + \varepsilon_{ij}, \quad (1)$$

where Y reflects person i 's trait score at time j ; π_{0i} reflects the intercept of the growth curve, π_{1i} reflects the slope of the growth curve, and ε reflects normally distributed error. In the above figure, I have depicted this initial slope as 0. When one adds pubertal onset (PO) as dichotomous,

$$Y_{ij} = \pi_{0i} + \pi_{1i} \text{TIME}_{ij} + \pi_{2i} \text{PO} + \varepsilon_{ij}, \quad (2)$$

where π_{2i} captures the magnitude of the shift when PO goes from 0 to 1. Because PO is treated as dichotomous, the resulting equation is identical to equation (1) when PO is zero. When PO is 1, the equation becomes

$$Y_{ij} = \pi_{0i} + \pi_{2i} + \pi_{1i} \text{TIME}_{ij} + \varepsilon_{ij}, \quad (2a)$$

indicating a higher intercept. To capture the hypothesized higher slope, one adds a second time variable, which counts time starting at PO

$$Y_{ij} = \pi_{0i} + \pi_{1i} \text{TIME}_{ij} + \pi_{2i} \text{PO} + \pi_{3i} \text{POTIME} + \varepsilon_{ij}, \quad (3)$$

where π_{3i} reflects the rate of increase as a function of POTIME (time beginning with PO, calculated as $\text{PO} \times \text{TIME}$). When PO is 0, POTIME is also 0, and the equation is again equation (1). When PO is 1, POTIME takes on nonzero values, and the equation becomes

$$Y_{ij} = \pi_{0i} + \pi_{2i} + \pi_{1i} \text{TIME}_{ij} + \pi_{3i} \text{POTIME} + \varepsilon_{ij}, \quad (3a)$$

reflecting both a higher intercept and change in Y as a function of two time variables, time plus time beginning with PO. The result can be thought of as two line segments: one before PO and one beginning with PO, as depicted in Figure 2.1. One can construct a parallel equation for drinking behavior (where the second time variable would be $\text{POTIME} + 1$).

Figure 2.1




Figure 2.1. Figural representation of the hypothesized sequential process by which pubertal onset (PO) leads to changes in negative urgency (NU), which in turn leads to changes in drinking behavior. The solid line represents negative urgency, and the left panel of the figure represents a mean increase in negative urgency with pubertal onset, followed by a slope increase that reflects increasing levels of negative urgency over time. In the right panel of the figure, the dotted line represents drinking behavior.

Chapter Three: Results

Participant Retention

Of the 1,910 participants, 96.5% participated at wave 1 and 82% participated at wave 7). At each wave, participating and non-participating participants did not differ on any study variables, so we inferred that data were missing at random. We therefore used the expectation maximization (EM) procedure to impute missing values, a procedure shown to more accurately approximate population data than traditional, alternative methods, such as case deletion or mean substitution (Enders, 2006).

Possible Effects due to School Membership

In order to determine whether there was significant covariance among the study variables due to participants attending the same school, we calculated intraclass coefficients for each variable (using elementary school membership, $n = 23$, as the nesting variable). Intraclass coefficients ranged from .03 (positive urgency) to .00. We therefore concluded that school membership was essentially unrelated to study variables.

Descriptive Statistics

The sample of 1,910 children includes 972 boys and 938 girls. Table 3.1 provides the following information: the percentage of youth who had experienced pubertal onset by each wave, drinking frequency by wave, and mean scores on each of the 5 impulsivity-related traits at each wave.

Correlations Among Traits, Behaviors and Pubertal Status Across Waves

Correlations among pubertal status, negative urgency, and drinking frequency across all 7 waves are presented in Table 3.2. All correlations involving pubertal status are point biserial correlations due to the dichotomous nature of the variable. Correlations involving all other variables are product-moment correlations. The cutoff for statistical significance was set to $p < .001$ because of the large sample size.

Model Tests

We first present results of the test of whether pubertal onset predicted mean and slope increases in negative urgency. We then provide results of the test of whether pubertal onset predicted mean and slope increases in drinking behavior. Next, we provide results of the test of whether negative urgency predicted subsequent mean and slope increases in drinking behavior. We then summarize the parallel tests we conducted for each of the other four personality traits. For each analysis described below, we tested whether any of the observed effects were moderated by sex. None were.

Change in negative urgency across early adolescence. We first tested whether there was, in fact, significant variability in change in negative urgency across the 7 waves. We thus examined between-person variability in the growth of negative urgency over time, applying both linear and quadratic growth models. There was significant variability in growth patterns of negative urgency over time ($p < .001$). The presence of this variability indicates the value of identifying determinants of that variability. In the current study, we identified pubertal onset as such a predictor. Mean levels of negative urgency did not increase as a function of time, but did increase with pubertal onset ($\beta = .06, p < .001$); this finding reflects a mean increase in negative urgency associated with the onset of puberty. Negative urgency also increased over time since pubertal onset ($\beta = .04, p < .001$), indicating an increase in the slope of growth in negative urgency following pubertal onset. Thus, our hypotheses were confirmed.

Change in drinking behavior. We identified significant variability around linear and quadratic growth curves for drinking behavior ($p < .001$). We found that drinking behavior increased as a function of time ($\beta = .21, p < .001$), pubertal onset ($\beta = .07, p < .001$) and time since pubertal onset ($\beta = .06, p < .001$). While existing research has shown that drinking behavior tends to increase over time during early adolescence

(Chung et al., 2012), the current findings indicate that the time-varying covariate of pubertal onset is associated with mean and slope increases in drinking frequency.

Negative urgency predicting change in drinking. We anticipated increases in negative urgency to concurrently and prospectively predict increases in drinking behavior. Negative urgency did predict increases in drinking behavior both concurrently ($\beta = .13, p < .001$) and prospectively ($\beta = .02, p < .001$). The prospective effect is above and beyond prediction from prior drinking behavior, which also predicted subsequent drinking ($\beta = .13, p < .001$), pubertal onset ($\beta = .04, p < .001$), and time since pubertal onset ($\beta = .13, p < .001$). Together, this set of findings indicates that drinking frequency in a given wave was predicted from negative urgency, drinking frequency, and pubertal onset all measured the prior wave. In addition, the rate of increase in drinking increased beginning the wave following pubertal onset. Each of these findings is consistent with our model.

Tests of Specificity of the Relationship Between Pubertal Onset and Negative Urgency

We next tested whether any of the other four impulsivity-related traits (positive urgency, sensation seeking, lack of planning, and lack of perseverance) underwent the same pattern of change in relation to pubertal onset. Our hypothesis was that the effects described above would be specific to negative urgency. For each of these traits, there was significant variability in trait growth over time ($p < .001$).

Positive urgency did not increase as a function of time, but did increase with pubertal onset ($\beta = .05, p < .001$), indicating a mean increase in the trait following pubertal onset. Positive urgency change was unrelated to time following pubertal onset. Unlike negative urgency, the slope of change in positive urgency did not become more positive following pubertal onset.

Sensation seeking increased as a function of time ($\beta = .09, p < .001$), and pubertal onset was associated with a positive mean increase in the trait ($\beta = .07, p < .001$). However, time since pubertal onset was significantly negatively related to change in sensation seeking ($\beta = -.06, p < .001$).

Mean trait levels of lack of planning increased as a function of time ($\beta = .18, p < .001$), did not have a mean change with pubertal onset, and time since pubertal onset was negatively associated with change in lack of planning ($\beta = -.06, p < .001$).

Lack of perseverance increased as a function of time ($\beta = .05, p < .001$) and decreased as a function of pubertal onset ($\beta = -.04, p < .001$); that is, the mean level of the trait declined with the onset of puberty. There was no association between pubertal time and lack of perseverance. Thus, none of the other four traits manifested both mean and slope increases following the onset of puberty.

Although none of the other four traits showed the pattern of change related to pubertal onset shown by negative urgency, it is of course important to investigate their roles in predicting drinking behavior. Each of the following prospective prediction findings included control for prior drinking. Positive urgency was positively associated with drinking concurrently ($\beta = .10, p < .001$) and prospectively ($\beta = .02, p < .001$). Sensation seeking was associated with concurrent changes in drinking ($\beta = .05, p < .001$), but did not predict future changes in drinking. Lack of planning was associated positively with changes in drinking both concurrently ($\beta = .06, p < .001$) and prospectively ($\beta = .04, p < .001$). Lack of perseverance was not associated with concurrent changes in drinking but did positively predict subsequent changes in drinking ($\beta = .05, p < .001$).

Table 3.1

Descriptive Statistics for Pubertal Statuorientations, Drinker Status, and Five Personality Trait Contributors to Rash Action

| Pubertal Status | Drinker Status | NU | PU | SS | LPI | LPs | |
|-----------------|----------------|--------------|---------------|---------------|---------------|---------------|---------------|
| | % | mean (sd) | mean (sd) | mean (sd) | mean (sd) | mean (sd) | |
| Wave 1 | 23.2 | 10.3 | 2.21 (.70) | 2.14 (.74) | 2.63 (.70) | 2.01 (.55) | 2.04 (.48) |
| Wave 2 | 32.1 | 11.4 | 2.11 (.69) | 2.04 (.76) | 2.63 (.74) | 2.10 (.58) | 2.09 (.52) |
| Wave 3 | 42.0 | 14.4 | 2.15 (.72) | 2.06 (.79) | 2.66 (.73) | 2.20 (.59) | 2.11 (.54) |
| Wave 4 | 50.5 | 17.1 | 2.19 (.71) | 2.06 (.78) | 2.70 (.71) | 2.14 (.57) | 2.03 (.53) |
| Wave 5 | 62.8 | 21.6 | 2.18 (.72) | 2.06 (.79) | 2.70 (.74) | 2.22 (.57) | 2.09 (.57) |
| Wave 6 | 62.8 | 31.0 | 2.19 (.69) | 2.06 (.77) | 2.76 (.71) | 2.18 (.52) | 2.04 (.52) |
| Wave 7 | 73.4 | 32.1 | 2.21 (.70) | 2.06 (.78) | 2.73 (.73) | 2.20 (.56) | 2.10 (.58) |

Note: NU: negative urgency; PU: positive urgency; SS: sensation seeking; LPI: lack of planning; LPs: lack of perseverance. *N* = 1910.

Table 3.2
Bivariate Correlations Among Study Variables in Waves 1 and 7

| | PUB-1 | PUB-7 | NU-1 | NU-7 | PU-1 | PU-7 | SS-1 | SS-7 | LPL-1 | LPL-7 | LPS-1 | LPS-7 | DR-1 | DR-7 |
|-------|-------|-------|--------|--------|-------|-------|--------|--------|-------|-------|-------|-------|-------|------|
| PUB-1 | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| PUB-7 | .112* | - | - | - | - | - | - | - | - | - | - | - | - | - |
| NU-1 | .113* | .035 | - | - | - | - | - | - | - | - | - | - | - | - |
| NU-7 | .110* | .090* | .349* | - | - | - | - | - | - | - | - | - | - | - |
| PU-1 | .126* | .023 | .627* | .336* | - | - | - | - | - | - | - | - | - | - |
| PU-7 | .122* | .054 | .281* | .673* | .354* | - | - | - | - | - | - | - | - | - |
| SS-1 | .076* | .095* | .269* | .148* | .316* | .145* | - | - | - | - | - | - | - | - |
| SS-7 | .022 | .086* | .075* | .330* | .120* | .289* | .443* | - | - | - | - | - | - | - |
| LPL-1 | .066* | .036 | .358* | .189* | .308* | .173* | .134* | .089* | - | - | - | - | - | - |
| LPL-7 | .061* | .017 | .193* | .175* | .175* | .202* | .072* | -.076* | .309* | - | - | - | - | - |
| LPS-1 | -.020 | -.024 | -.040* | .047 | -.039 | .041 | -.295* | -.138* | .425* | .169* | - | - | - | - |
| LPS-7 | .029 | -.020 | .065* | -.077* | .045 | -.046 | .069* | .349* | .158* | .657* | .265* | - | - | - |
| DR-1 | .153* | -.010 | .217* | .109* | .180* | .092* | .138* | .022 | .180* | .102* | .081* | .063* | - | - |
| DR-7 | .136* | .069* | .161* | .327* | .129* | .262* | .149* | .175* | .147* | .182* | .024 | .045 | .246* | - |

Note: PUB, pubertal onset; NU, negative urgency; PU, positive urgency; SS, sensation seeking; LPL, lack of planning; LPS, lack of perseverance; DR, drinking; each number following the variable represents the wave in which the data was collected (i.e., NU-1 = Negative Urgency, wave 1). Correlations with puberty are point-biserial correlations, and all others are Pearson correlations. * $p < .001$.

Chapter Four: Discussion

In this study, we found empirical support for a novel model that helps explain part of the well-known finding that alcohol use increases dramatically across the early adolescent years (Chung et al., 2012). Specifically, pubertal onset led to mean and slope increases in negative urgency; these trait changes in turn led to mean and slope increases in drinking behavior across early adolescence. That is, the wave at which pubertal onset was reported was associated with both mean and slope increases in negative urgency. Negative urgency, in turn, predicted the mean level of drinking frequency the following wave, as well as the slope of increase in drinking beginning that following wave. This prediction occurred in the context of the rigorous control of predicting drinking each wave from drinking the prior wave (6 months earlier). It thus appears that one part of the explanation for increased drinking during early adolescence may lie in the intersection between the biological event of pubertal onset and associated changes in the high risk personality trait of negative urgency.

This pattern of findings is consistent with the developmental model proposed by Cyders and Smith (2008). The experience of pubertal onset is known to be associated with increases in negative affect and in rash, risky behavior (Casey & Caudle, 2013; Compas et al., 1995). Cyders and Smith argued that these changes reflected a developmental increase in negative urgency. We found support for this contention as well as for the contention that the developmental increase in the trait helps account for subsequent increases in drinking behavior. This finding is quite important clinically, because early adolescent alcohol consumption is associated with a host of co-occurring and future problems (Chung et al., 2012; DeWit et al., 2000; Grant & Dawson, 1997; Jessor, 1987).

Cyders and Smith (2008) argued that, in a context in which negative affect occurs frequently, each time a person acts rashly to alleviate his or her distress, the rash action is negatively reinforced (due to distraction from the original source of distress). Therefore, over time, rash actions such as drinking become more likely. In addition, on each such occasion the person misses an opportunity to learn alternative, more adaptive ways of coping with distress. Thus, there is a combination of reinforcement for the disposition toward negative urgency and a lack of development of alternatives to negative urgency-based rash action. This process results in a profile of increasing risk over time.

In important ways, this phenomenon was specific to the trait of negative urgency. The slope of change in negative urgency became more positive over time following pubertal onset. This effect was not observed for any of the other four traits studied. In fact, pubertal onset was negatively associated with the slopes of sensation seeking and lack of planning. Pubertal onset was unrelated to the slopes of positive urgency and lack of perseverance. This specificity lends further support to our model.

It is important to appreciate that other traits also predicted drinking behavior. Positive urgency, sensation seeking, and lack of planning were each associated with concurrent drinking behavior. Positive urgency, lack of planning, and lack of perseverance predicted subsequent drinking behavior, controlling for prior drinking behavior. None of these three traits changed as a function of pubertal onset in the way that negative urgency did. Thus, although their roles in the developmental process appear to differ from that of negative urgency, they did predict subsequent drinking in this early adolescent sample. Their possible functional roles in relation to the development of early adolescent drinking merit further investigation.

None of the effects summarized above were moderated by sex; that is, we found no evidence that any of the identified processes differed for girls and boys. It is certainly

possible that a longer-term prospective study, investigating the emergence of diagnosable alcohol abuse and dependence, would reveal important sex differences. Nonetheless, it was striking that the same process appeared to operate for both sexes.

We studied the predictive roles of each of the five traits separately, running different multilevel models for each trait. We elected to do so for three reasons. First, the theoretical model driving this work concerned a specific puberty-based process for negative urgency, and we judged it important to test that process directly. Second, it did seem important to understand whether each trait, individually, changed as a function of time and pubertal status. Third, at this early stage of investigation of the intersection of developmental trait change and drinking risk, it seemed premature to investigate those parts of each trait that are independent of the other four traits, which is what would be involved in modeling change in, and prediction from, all five traits at once. In prior correlational work with both early adolescents and college students, the predictive roles of sensation seeking, lack of planning, and lack of perseverance were not present when those traits were controlled for negative and positive urgency (Cyders, Flory, Rainer, & Smith, 2009; Settles et al., 2014).

The findings of this study should be considered in the context of its limitations. We had some missing data. Although there was no indication that missing-ness biased the study's results, we cannot know with certainty whether any findings would differ had there been no attrition. We assessed each trait, pubertal status, and drinking behavior by questionnaire. Our method precluded detailed responses to questions about the survey and the kind of clarification that could result from an interview format. However, there is good evidence for the reliability and validity of each measure used. A strength of the study is that we studied youth across 7 waves, from the end of elementary school through the end of middle school. However, the absence of information about trait change and drinking behavior change during the high school years limits the conclusions

we can draw. Although we did model the effects of pubertal onset as a time-varying predictor, which allowed us to model that effect in a person-specific way, our data analytic plan did not allow for the possibility that different youth proceed along different developmental trajectories with respect to both traits and drinking behavior. Trajectory analyses, which are designed to determine the value of modeling distinct developmental groups, can provide a valuable addition to the findings of this study.

In one way, the model we tested is complex and elaborate. At the same time, we isolated and studied just one specific developmental process. Comprehensive models of risk will need to integrate the current findings with many other risk processes that have been identified in the literature. For example, low parental monitoring and responsiveness (Baumrind, 1985), greater parental drinking (Wong, Brower, Fitzgerald, & Zucker, 2004), parental alcoholism (Russell, 1990), children's sleep loss (Wong et al., 2004), children's engagement in externalizing behaviors (Clark, Parker, & Lynch, 1999; McGue, Iacono, Legrand, & Elkins, 2001), children's engagement in internalizing behaviors (Hussong, Jones, Stein, Baucom, & Boeding, 2011), and learned expectancies for reinforcement from alcohol (Smith, Goldman, Greenbaum, & Christiansen, 1995) all predict early drinking. It is likely that risk reflects developmental transactions among all of these variables, pubertal status, and others, both within and across time.

References

- Barbas, H. (2007). Specialized elements of orbitofrontal cortex in primates. *Annals of the New York Academy of Sciences*, 1121, 10-32. doi: 10.1196/annals.1401.015
- Baumrind, D. (1985). Familial antecedents of adolescent drug use: A developmental perspective. In Jones, C. L., and Battles, R. J. (Eds.), *Etiology of Drug Abuse: Implications for Prevention*. Rockville, MD: National Institute on Drug Abuse, NIDA Research Monograph 56.
- Bechara, A. (2005). Decision making, impulse control and loss of willpower to resist drugs: A neurocognitive perspective. *Nature Neuroscience*, 8(11), 1458-1463. doi: 10.1038/nn1584
- Bleidorn, W. (2012). Hitting the road to adulthood: Short-term personality development during a major life transition. *Personality and Social Psychology Bulletin*, 38(12), 1594-1608.
- Brooks-Gunn, J., Graber, J. A., & Paikoff, R. L. (1994). Studying links between hormones and negative affect: Models and measures. *Journal of Research on Adolescence*, 4(4), 469-486.
- Brooks-Gunn, J., Warren, M. P., Rosso, J., & Gargiulo, J. (1987). Validity of self-report measures of girls' pubertal status. *Child Development*, 58(3).
- Cardinal, R. N., Parkinson, J. A., Hall, J., & Everitt, B. J. (2002). Emotion and motivation: The role of the amygdala, ventral striatum, and prefrontal cortex. *Neuroscience & Biobehavioral Reviews*, 26(3), 321-352.
- Caspi, A., & Roberts, B. W. (2001). Personality development across the life course: The argument for change and continuity. *Psychological Inquiry*, 12(2), 49-66.
- Chung, T., Smith, G. T., Donovan, J. E., Windle, M., Faden, V. B., Chen, C. M., & Martin, C. S. (2012). Drinking frequency as a brief screen for adolescent alcohol problems. *Pediatrics*, 129(2), 205-212. doi: 10.1542/peds.2011-1828
- Clark, D. B., Parker, A. M., & Lynch, K. G. (1999). Psychopathology and substance-related problems during early adolescence: A survival analysis. *Journal of Clinical Child Psychology*, 28, 333-341.
- Coleman, L., & Coleman, J. (2002). The measurement of puberty: A review. *Journal of Adolescence*, 25(5), 535-550.
- Combs, J. L., Spillane, N. S., Caudill, L., Stark, B., & Smith, G. T. (2012). The acquired preparedness risk model applied to smoking in 5th grade children. *Addictive Behaviors*, 37(3), 331-334. doi: 10.1016/j.addbeh.2011.11.005
- Coskunpinar, A., Dir, A. L., & Cyders, M. A. (2013). Multidimensionality in impulsivity and alcohol use: A meta-analysis using the UPPS-P model of impulsivity. *Alcoholism: Clinical and Experimental Research*, 37, 1441-1450.

- Culbert, K. M., Burt, S. A., McGue, M., Iacono, W. G., & Klump, K. L. (2009). Puberty and the genetic diathesis of disordered eating attitudes and behaviors. *Journal of Abnormal Psychology, 118*(4), 788-796.
- Curran, P. J. (2003). Have multilevel models been structural equation models all along? *Multivariate Behavioral Research, 38*(4), 529-569.
- Cyders, M. A., & Smith, G. T. (2008). Emotion-based dispositions to rash action: Positive and negative urgency. *Psychological Bulletin, 134*(6), 807-828.
- Cyders, M. A., & Smith, G. T. (2010). Longitudinal validation of the urgency traits over the first year of college. *Journal of Personality Assessment, 92*, 63-69.
- Cyders, M. A., Smith, G. T., Spillane, N. S., Fischer, S., Annus, A. M., & Peterson, C. (2007). Integration of impulsivity and positive mood to predict risky behavior: development and validation of a measure of positive urgency. *Psychological Assessment, 19*(1), 107-118.
- DeWit, D. J., Adlaf, E. M., Offord, D. R., & Ogborne, A. C. (2000). Age at first alcohol use: A risk factor for the development of alcohol disorders. *American Journal of Psychiatry, 157*(5), 745-750.
- Donovan, J. E. (2007). Really underage drinkers: The epidemiology of children's alcohol use in the United States. *Prevention Science, 8*(3), 192-205. doi: 10.1007/s11121-007-0072-7
- Enders, C. (2006). A primer on the use of modern missing-data methods in psychosomatic medicine research. *Psychosomatic Medicine, 68*, 427-736.
- Fischer, S., Peterson, C. M., & McCarthy, D. (2013). A prospective test of the influence of negative urgency and expectancies on binge eating and purging. *Psychology of Addictive Behaviors, 27*(1), 294-300. doi: 10.1037/a0029323
- Grant, B. F., & Dawson, D. A. (1997). Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: results from the National Longitudinal Alcohol Epidemiologic Survey. *Journal of Substance Abuse, 9*, 103-110.
- Gruber, E., DiClemente, R. J., Anderson, M. M., & Lodico, M. (1996). Early drinking onset and its association with alcohol use and problem behavior in late adolescence. *Preventive Medicine, 25*(3), 293-300. doi: 10.1006/pmed.1996.0059
- Guller, L., Zapolski, T. C., & Smith, G. T. (2012). *Impulsivity traits and the longitudinal prediction of addictive behaviors during the transition from preadolescence to adolescence*. Manuscript submitted for publication.
- Gunn, R. L., & Smith, G. T. (2010). Risk factors for elementary school drinking: Pubertal status, personality, and alcohol expectancies concurrently predict fifth grade alcohol consumption. *Psychology of Addictive Behaviors, 24*(4), 617-627. doi: 10.1037/a0020334

- Guo, J., Collins, L. M., Hill, K. G., & Hawkins, J. D. (2000). Developmental pathways to alcohol abuse and dependence in young adulthood. *Journal of Studies on Alcohol*, 61(6), 799-808.
- Guttmanova, K., Hill, K. G., Bailey, J. A., Lee, J. O., Hartigan, L. A., Hawkins, J. D., & Catalano, R. F. (2012). Examining explanatory mechanisms of the effects of early alcohol use on young adult alcohol dependence. *J Stud Alcohol Drugs*, 73(3), 379-390.
- Hariri, A. R., Drabant, E. M., & Weinberger, D. R. (2006). Imaging genetics: Perspectives from studies of genetically driven variation in serotonin function and corticolimbic affective processing. *Biological Psychiatry*, 59(10), 888-897. doi: 10.1016/j.biopsych.2005.11.005
- Hussong, A.M., Jones, D.J., Stein, G.L., Baucom, D.H., & Boeding, S. (2011). An internalizing pathway to alcohol use and disorder. *Psychology of Addictive Behaviors*, 25, 390-404. Doi: 10.1037/a0024519
- Jackson, D. L. (2001). Sample Size and Number of Parameter Estimates in Maximum Likelihood Confirmatory Factor Analysis: A Monte Carlo Investigation. *Structural Equation Modeling: A Multidisciplinary Journal*, 8(2), 205-223.
- Jackson, D. L. (2003). Revisiting sample size and number of parameter estimates: some support for the N:q hypothesis. *Structural Equation Modeling: A Multidisciplinary Journal*, 10(1), 128-141. doi: 10.1207/s15328007sem1001_6
- Jessor, R. (1987). Problem-behavior theory, psychosocial development, and adolescent problem drinking. *British Journal of Addiction*, 82(4), 331-342.
- Kline, R. B. (1998). *Principles and practice of structural equation modeling*. New York: Guilford Press.
- Lewis, M. D., & Todd, R. M. (2007). The self-regulating brain: Cortical-subcortical feedback and the development of intelligent action. *Cognitive Development*, 22(4), 406-430.
- Luna, B., & Sweeney, J. A. (2004). The emergence of collaborative brain function: FMRI studies of the development of response inhibition. *Annals of the New York Academy of Sciences*, 1021, 296-309.
- Maggs, J. L., & Hurrelmann, K. (1995). *Health impairments in adolescence: The biopsychosocial "costs" of modern life-style*. Oxford, England: Walter De Gruyter.
- McCrae, R. R., & Costa, P. T. (1994). The stability of personality: Observations and evaluations. *Current Directions in Psychological Science*, 3(6), 173-175.
- McGue, M., Iacono, W. G., Legrand, L. N., & Elkins, I. Origins and consequences of age at first drink, part II: Familial risk and heritability. *Alcoholism: Clinical and Experimental Research*, 25, 1166-1175.

- Metcalfe, J., & Mischel, W. (1999). A hot/cool-system analysis of delay of gratification: Dynamics of willpower. *Psychological Review*, *106*(1), 3-19.
- Moffitt, T. E. (1993). Adolescence-limited and life-course-persistent antisocial behavior: A developmental taxonomy. *Psychological Review*, *100*(4), 674-701.
- Muthén, L. K., & Muthén, B. O. (1998-2010). *Mplus User's Guide* (Seventh ed.). Los Angeles, CA: Muthén & Muthén
- Nelson, C. A., Bloom, F. E., Cameron, J. L., Amaral, D., Dahl, R. E., & Pine, D. (2002). An integrative, multidisciplinary approach to the study of brain-behavior relations in the context of typical and atypical development. [Research Support, Non-U.S. Gov't]. *Development and Psychopathology*, *14*(3), 499-520.
- Neyer, F. J., & Asendorpf, J. B. (2001). Personality-relationship transaction in young adulthood. *Journal of Personality and Social Psychology*, *81*(6), 1190-1204.
- Pearson, C. M., Combs, J. L., Zapolski, T. C., & Smith, G. T. (2012). A longitudinal transactional risk model for early eating disorder onset. *Journal of Abnormal Psychology*, *121*(3), 707-718. doi: 10.1037/a0027567
- Pearson, C. M., Guller, L., Spillane, N. S., & Smith, G. T. (2011). A developmental model of addictive behavior: From impulsivity to compulsivity. In A. M. Columbus (Ed.), *Advances in Psychology Research* (Vol. 77). New York: Nova Science Publishers.
- Petersen, A., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*, *17*(2), 117-133. doi: 10.1007/bf01537962
- Raudenbush, S., & Liu, X. (2001). Effects of study duration, frequency of observation, and sample size on power in studies of group differences in polynomial change. *Psychological Methods*, *6*(4), 387-401.
- Roberts, B. W. (2006). Personality Development and Organizational Behavior. *Research in Organizational Behavior*, *27*, 1-40.
- Roberts, B. W., Caspi, A., & Moffitt, T. E. (2001). The kids are alright: Growth and stability in personality development from adolescence to adulthood. *Journal of Personality and Social Psychology*, *81*(4), 670-683.
- Roberts, B. W., & DelVecchio, W. F. (2000). The rank-order consistency of personality traits from childhood to old age: A quantitative review of longitudinal studies. *Psychological Bulletin*, *126*(1), 3.
- Russell, M. (1990). Prevalence of alcoholism among children of alcoholics. In Windle, M. (Ed.), *Children of Alcoholics: Critical Perspectives*, 9-38. New York: Guilford.
- Settles, R. E., & Smith, G. T. (2011). *Toward a truly developmental model of adolescent problem drinking*. Paper presented at the the annual meeting of the Research Society on Alcoholism, Atlanta, GA.

- Settles, R. E., Fischer, S., Cyders, M. A., Combs, J. L., Gunn, R. L., & Smith, G. T. (2012). Negative urgency: A personality predictor of externalizing behavior characterized by Neuroticism, low Conscientiousness, and Disagreeableness. *Journal of Abnormal Psychology, 121*, 160-172.
- Settles, R. F., Cyders, M. A., & Smith, G. T. (2010). Longitudinal validation of the acquired preparedness model of drinking risk. *Psychology of Addictive Behaviors, 24*(2), 198-208.
- Smith, G. T., Fischer, S., Cyders, M., Annus, A., Spillane, N., & McCarthy, D. (2007). On the validity and utility of discriminating among impulsivity-like traits. *Assessment, 14*(2), 155-170.
- Smith, G. T., Goldman, M. S., Greenbaum, P., & Christiansen, B. A. (1995). The expectancy for social facilitation from drinking: the divergent paths of high-expectancy and low-expectancy adolescents. *Journal of Abnormal Psychology, 104*, 32-40.
- Smith, G. T., McCarthy, D. M., & Goldman, M. S. (1995). Self-reported drinking and alcohol-related problems among early adolescents: Dimensionality and validity over 24 months. *Journal of Studies on Alcohol, 56*(4), 383-394.
- Stautz, K., & Cooper, A. (2013). Impulsivity-related personality traits and adolescent alcohol use: A meta-analytic review. *Clinical Psychology Review, 33*(4), 574-592.
- Steinberg, L. (2004). Risk taking in adolescence: What changes, and why? *Annals of the New York Academy of Sciences, 1021*, 51.
- Steinberg, L., Albert, D., Cauffman, E., Banich, M., Graham, S., & Woolard, J. (2008). Age differences in sensation seeking and impulsivity as indexed by behavior and self-report: Evidence for a dual systems model. *Developmental Psychology, 44*(6), 1764-1778.
- Substance Abuse and Mental Health Services Administration. (2012). *Results from the 2011 National Survey on Drug Use and Health: Summary of National Findings*. (NSDUH Series H-44, HHS Publication No. (SMA) 12-4713). Rockville, MD: Substance Abuse and Mental Health Services Administration. Retrieved from <http://www.samhsa.gov/data/nsduh/2k11results/nsduhresults2011>.
- Thapar, A., Cooper, M., Jefferies, R., & Stergiakouli, E. (2012). What causes attention deficit hyperactivity disorder? *Archives of Disease in Childhood, 97*(3), 260-265. doi: 10.1136/archdischild-2011-300482
- Whiteside, S., & Lynam, D. (2001). The Five Factor Model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences, 30*(4), 669-689.

Wong, M.M., Brower, K.J., Fitzgerald, H.E., & Zucker, R.A. (2004). Sleep problems in early childhood and early onset of alcohol and other drug use in adolescence. *Alcoholism: Clinical and Experimental Research*, 28, 578-587. doi: 10.1097/01.ALC.0000121651.75952.

Zapolski, T. C., Stairs, A. M., Settles, R. F., Combs, J. L., & Smith, G. T. (2010). The measurement of dispositions to rash action in children. *Assessment*, 17(1), 116-125.

LAUREN HELENA BOYLE

VITA
Department of Psychology
University of Kentucky

EDUCATION

University of California, Berkeley
B.A. in Psychology
Date of Completion: May 2005

HONORS & AWARDS

College of Letters & Science, Honors, UC Berkeley December 2001

PUBLICATIONS

Boyle, L.H., and Smith, G.T. (Forthcoming). "Spectrum model." In R. Cautin and S. Lilienfeld (Eds.), *The Encyclopedia of Clinical Psychology*. Malden, Oxford: John Wiley and Sons, Inc.

Guller, L., **Boyle, L.**, Spillane, N.S., Smith, G.T. (In press). Impulsive action and impulsive inaction: Toward an integrative theory of impulsivity. In M.C. Olmstead (Ed.), *Psychology of Impulsivity: New Research*. Hauppauge, NY: Nova Science Publishers.

PROFESSIONAL POSITIONS

| | |
|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 2013-2014 | Department of Psychology, University of Kentucky <i>Research Assistant</i> |
| 2013-2014 | Jesse G. Harris Psychological Services Center, University of Kentucky <i>Therapist</i> |
| 2013-2014 | Counseling Center, University of Kentucky <i>Student Therapist</i> |
| 2010-2012 | UC Davis Medical Investigation of Neurodevelopmental Disorders (M.I.N.D.) Institute, University of California, Davis <i>Junior Specialist/Study Coordinator</i> |
| 2010-2012 | Imaging Research Center, University of California, Davis <i>Research Assistant</i> |