

8-2009

Ridge preservation comparing the clinical and histologic healing of a mineralized particulate allograft with a nonporous PTFE membrane vs. mineralized particulate xenograft with a collagen plug membrane.

Jason A. Witonsky  
*University of Louisville*

Follow this and additional works at: <https://ir.library.louisville.edu/etd>

---

#### Recommended Citation

Witonsky, Jason A., "Ridge preservation comparing the clinical and histologic healing of a mineralized particulate allograft with a nonporous PTFE membrane vs. mineralized particulate xenograft with a collagen plug membrane." (2009). *Electronic Theses and Dissertations*. Paper 1585.

<https://doi.org/10.18297/etd/1585>

This Master's Thesis is brought to you for free and open access by ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. This title appears here courtesy of the author, who has retained all other copyrights. For more information, please contact [thinkir@louisville.edu](mailto:thinkir@louisville.edu).

**RIDGE PRESERVATION COMPARING THE CLINICAL AND HISTOLOGIC  
HEALING OF A MINERALIZED PARTICULATE ALLOGRAFT WITH A  
NONPOROUS PTFE MEMBRANE VS. MINERALIZED PARTICULATE  
XENOGRAFT WITH A COLLAGEN PLUG MEMBRANE**

By

Jason A. Witonsky  
DMD, University of Pennsylvania, 2006

A Thesis  
Submitted to the Faculty of the  
Graduate School of the University of Louisville  
in Partial Fulfillment of the Requirements  
for the Degree of

Master of Science

Program in Oral Biology  
School of Dentistry  
University of Louisville  
Louisville, Kentucky

August 2009

**RIDGE PRESERVATION COMPARING THE CLINICAL AND HISTOLOGIC  
HEALING OF A MINERALIZED PARTICULATE ALLOGRAFT WITH A  
NONPOROUS PTFE MEMBRANE VS. MINERALIZED PARTICULATE  
XENOGRAFT WITH A COLLAGEN PLUG MEMBRANE**

By

Jason A. Witonsky  
DMD, University of Pennsylvania, 2006

A Thesis Approved on

August 7, 2009

By the following Thesis Committee:

---

Thesis Director

---

---

## **DEDICATION**

This manuscript is dedicated to my parents for their love and support and to the pursuit of science. The love and support from my parents has been invaluable. It has made me a humble, hard working, caring, and skillful healthcare provider. Science has the ability to enlighten humanity; therefore my efforts go towards its pursuit.

## ACKNOWLEDGEMENTS

I would like to express my sincere gratitude to the following individuals:

Dr. Henry Greenwell, Program Director of Graduate Periodontics, for his invaluable guidance and mentoring in my training as a periodontist, his help in the preparation of this thesis and for the lifelong friend I have acquired.

Dr. Margaret Hill, Assistant Program Director, for her inspiration and commitment to excellence and her caring encouragement of all the residents.

Dr. Brian S. Shumway, Assistant Professor, thank you very much for all of your help. I appreciate all of your input into my thesis.

## ABSTRACT

### RIDGE PRESERVATION COMPARING THE CLINICAL AND HISTOLOGIC HEALING OF A MINERALIZED PARTICULATE ALLOGRAFT WITH A NONPOROUS PTFE MEMBRANE VS. MINERALIZED PARTICULATE XENOGRAFT WITH A COLLAGEN PLUG MEMBRANE

Jason Witonsky, DMD

August 7th, 2009

**Aim.** To compare two techniques of ridge preservation using a cancellous mineralized particulate xenograft plus a collagen plug to a cortical mineralized particulate allograft plus a PTFE membrane using ridge dimension data to assess the outcome.

**Methods.** Twenty-eight total patients were seen in the Graduate Periodontics Clinic at the University of Louisville School of Dentistry. Fourteen positive controls received a mineralized particulate xenograft (0.25 to 1.00 mm) covered by a collagen plug using a full-thickness flap technique (Plug group). Fourteen test patients received an intrasocket mineralized cortical particulate allograft (500 to 800  $\mu$ m) covered with a nonporous PTFE membrane also using a full thickness flap technique (PTFE group). Following tooth extraction, horizontal ridge dimensions were measured with a digital caliper and vertical ridge dimensions were measured from a stent. Each site was re-entered for implant placement at about 4 months. Prior to implant placement, a 2.7 X 6 mm trephine

core was obtained and preserved in formalin for histologic analysis.

**Results.** The mean horizontal ridge width at the crest of the Plug group decreased from  $8.6 \pm 1.0$  mm to  $7.3 \pm 1.0$  mm for a mean loss of  $-1.3 \pm 0.9$  mm ( $p < 0.05$ ) while the PTFE group decreased from  $7.9 \pm 1.5$  mm to  $6.8 \pm 1.4$  mm for a mean loss of  $-1.1 \pm 1.1$  mm ( $p < 0.05$ ). There were no statistically significance differences between the two groups ( $p > 0.05$ ). The mean mid-buccal vertical change for the Plug group was a loss of  $-0.1 \pm 1.6$  mm ( $p > 0.05$ ) vs. a gain of  $0.4 \pm 2.1$  mm ( $p > 0.05$ ) for the PTFE group. There were no statistically significant differences between groups for vertical change ( $p > 0.05$ ). The Plug group demonstrated  $28 \pm 20\%$  vital bone,  $37 \pm 16\%$  non-vital bone, and  $35 \pm 13\%$  trabecular space. The PTFE group demonstrated  $35 \pm 21\%$  vital bone,  $31 \pm 22\%$  non-vital bone, and  $34 \pm 10\%$  trabecular space. There were no statistically significant differences between groups ( $p > 0.05$ ).

**Conclusions.** Mean crestal ridge width was preserved for both the Plug and PTFE groups and there were no statistically significant differences between groups ( $p > 0.05$ ). There was a trend toward greater loss of mean mid-buccal ridge height for the Plug group, although there were no statistically significant differences between groups ( $p > 0.05$ ). The mean CEJ to osseous crest distance showed only a minimal loss of 0.7 mm or less, with no statistically significant differences between groups ( $p > 0.05$ ).

## TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS.....	iv
ABSTRACT.....	v
TABLE OF CONTENTS.....	vii
LIST OF TABLES.....	ix
LIST OF FIGURES.....	x
CHAPTER	
I. LITERATURE REVIEW	
Animal Extraction Socket Healing Sequence .....	1
Human Extraction Socket Healing Sequence .....	6
Alveolar Ridge Resorption following Tooth Extraction.....	8
Clinical Studies of Ridge Preservation .....	11
Polytetrafluoroethylene Technique Studies.....	16
Bio-Col Technique Studies.....	20
Histologic Evaluation of Ridge Preservation.....	24
Extraction Alone Studies .....	25
Allograft Studies .....	26
Xenograft Studies.....	28
Alloplast Studies .....	29
Summary of Histologic Findings.....	31
Summary of Literature Review.....	34



II.	MATERIALS AND METHODS	
	Study Design .....	39
	Patient Selection .....	41
	Post-Surgical Exclusion .....	41
	Pre-surgical Management.....	42
	Clinical measurements .....	42
	Surgical treatment.....	43
	Histologic analysis .....	44
	Statistical analysis .....	45
III.	RESULTS .....	46
IV.	DISCUSSION.....	54
V.	CONCLUSIONS.....	56
REFERENCES .....		59
APPENDICES.....		66
CURRICULUM VITAE.....		72

## LIST OF TABLES

TABLE	PAGE
1. Animal extraction socket healing 31 days (Clafin 1936).....	3
2. Animal extraction socket healing 180 days (Cardaropoli et al. 2003).....	3
3. Human extraction socket healing 56 days (Araujo et al. 2005a) .....	4
4. Human Extraction Socket Healing over 100 Days .....	6
5. Events In Extraction Socket Healing .....	8
6. Extraction Alone Studies Showing Change Alone .....	10
7. Extraction Alone Studies Showing Ridge Dimensions.....	11
8. Ridge Preservation Studies Showing Change Alone.....	23
9. Ridge Preservation Studies Showing Ridge Dimensions.....	24
10. Comparison of Histologic Data on Extraction Alone Studies.....	32
11. Comparison of Histologic Data on Ridge Preservation Studies .....	32
12. Root Dimensions at the Cervix by Tooth Types .....	36
13. Horizontal Ridge Width at the Crest for U of L Studies.....	38
14. U of L Studies by Tooth Type.....	38
15. Clinical Indices for Plug and PTFE Sites.....	49
16. Horizontal Ridge Width for Plug and PTFE Sites .....	50
17. Vertical Ridge Height Changes for Plug and PTFE Sites.....	51
18. CEJ to Osseous Crest Changes.....	52
19. Histologic Data for Plug and PTFE Sites.....	53

## LIST OF FIGURES

FIGURE	PAGE
1. Study Design.....	40
2a. Pre-op PTFE .....	57
2b. Cortical allograft.....	57
2c. PTFE membrane .....	57
2d. Pre-op 4-month re-entry.....	57
3a. Pre-op collagen plug .....	58
3b. Cancellous xenograft .....	58
3c. Collagen plug.....	58
3d. Pre-op 4-month re-entry.....	58

## **CHAPTER I**

### **LITERATURE REVIEW**

Tooth extraction has become a significant part of periodontal practice. The demand for dental implants as a tooth replacement requires clinicians to perform ridge preservation at the time of extraction to maintain post-extraction ridge width. This ensures a better site for future implant placement that satisfies both functional and esthetic requirements. Ridge preservation is an alternative to immediate implant placement when requirements of primary stability, adequate ridge dimensions or esthetics cannot be achieved. It is important to understand the events following tooth extraction that have been studied in both animals and humans in order to appreciate the most likely effects on ridge dimensions with or without a ridge preservation procedure.

#### **Animal Extraction Socket Healing Sequence**

The majority of the information about animal socket healing has been studied using the canine model. The earliest studies were completed in animals in the 1930's. Clafin (1936) provided data on the histologic healing of extraction sockets up to 31 days in dogs (Table 1). According to Clafin (1936), healing began with clot formation at day 1, followed by infiltration with osteoclasts at day 3, then bone formation around 5-7 days.

Epithelialization was complete over the clot around 7-9 days and complete socket fill occurred by 31 days. Despite complete socket fill, osteoclasts were still present, indicating that the healing was not complete at 31 days. Cardaropoli et al. (2003) extended the histologic analysis of the healing process of extraction sockets in beagle dogs to 180 days (Table 2). Both studies showed that the initial process after extraction was the formation of a blood clot at day 1. Subsequent to that, neovascularization played a significant role up to 14 days when new bone was formed along the socket walls. By day 30, in accord with Clafin, the socket was completely filled with bone. According to Cardaropoli et al. (2003), the bone at 30 days was immature. It was not until day 90 that this woven or immature bone had remodeled to become lamellar, mature bone. By day 180, the lamellar bone had undergone further remodeling and showed a slight decrease in mineralization due to the replacement of lamellar bone with bone marrow. Araujo et al. (2005a) examined histologic socket healing in the dog model using 12 sockets in 12 mongrel dogs over a period of 8 weeks (Table 3). At 1 week, the central portion of the socket was occupied by coagulum. At the apical portion, islands of newly formed woven bone were noted adjacent to the bundle bone. By 2 weeks, large amounts of newly formed bone were found in the apical and lateral portion of the socket. The surface of the woven bone was lined with densely packed osteoblasts and included a primitive bone marrow. By week 4, the crestal bone, which was completely composed of bundle bone, was lost. Apical to the crestal region, a multitude of osteoclasts were observed on the outer surfaces of the buccal and lingual walls. By week 8, the lingual wall had become wider than the buccal wall and was positioned 2 mm coronally to the buccal wall. A zone of mineralized tissue, which consisted of a mixture of woven and lamellar bone, had

formed between the buccal and lingual walls. This bridge of mineralized tissue traveled in an oblique direction. Two major findings from this study were: 1) the bundle bone began to disappear as early as 2 weeks post-extraction, and 2) the buccal wall undergoes a significantly greater amount of resorption than the lingual wall.

**Table 1**

**Animal Extraction Socket Healing 31 Days (Clafin 1936)**

<b>Time</b>	<b>Event</b>
Day 1	Blood clot formation
Day 3	Osteoclast appear at crest of bone and fibroblast emerge form socket walls
Day 5 to 7	First bone formation
Day 7 to 9	Epithelialization over clot completed
Day 11 to 15	New bone reaching the alveolar crest
Day 28 to 31	Socket filled with new bone, with osteoclasts still present

**Table 2**

**Animal Extraction Socket Healing 180 Days (Cardaropoli et al. 2003)**

<b>Time</b>	<b>Event</b>
Day 1	Blood clot formation comprising mostly of erythrocytes and platelets
Day 3	Lysis of erythrocytes and clot being replaced by vascularized tissue
Day 7	New blood vessel formation
Day 14	New bone formation on socket walls
Day 30	Socket filled with new bone
Day 90	Woven bone replaced by lamellar bone
Day 180	Some lamellar bone being replaced by bone marrow spaces

**Table 3****Animal Extraction Socket Healing 56 Days (Araujo et al. 2005a)**

<b>Time</b>	<b>Event</b>
Day 7 (1 week)	- internal portion of the socket occupied by coagulum - apical portion showed islands of newly formed woven bone adjacent to the bundle bone.
Day 14 (2 weeks)	- apical & lateral portions showed large amounts of newly formed woven bone - surface of the woven bone was lined with densely packed osteoblasts - primitive bone marrow.
Day 28 (4 weeks)	- at the crestal region, all bundle bone had been lost - crestal lamellar bone replaced with woven bone. - apical to the crestal region, a multitude of osteoclasts were observed on the outer surfaces of the buccal and lingual walls.
Day 56 (8 weeks)	- lingual wall wider than buccal wall - lingual wall positioned 2 mm coronal to buccal wall - zone of mineralized tissue which consist of a mixture of woven and lamellar bone had formed between the buccal and lingual walls traveling in an oblique direction.

Araujo et al. (2005b) also examined socket healing with placement of an immediate implant and Berglundh et al. (1994) studied the vascular supply around Branemark implants in beagle dogs. It was observed that the blood vessels of the peri-implant mucosa were terminal branches of larger vessels from the periosteum at the implant site. The peri-implant supracrestal connective tissue, in comparison to a tooth, was almost devoid of vascular supply, which could potentially influence healing. Carmagnola et al. (2000) examined the histologic healing around implants placed in sites previously grafted with mineralized cortical bovine xenograft (Bio-Oss). They created 16 surgical defects in 4 beagle dogs. Osseointegration failed to occur at the implant surfaces and a well-defined connective tissue capsule was present between implant surfaces. A deep vertical bone defect was frequently present along the lingual surface of the implant.

Botticelli et al. (2004) examined the effects of three different surgically created defect configurations on bone healing around implants. They found that the 4-wall defects fully resolved following implant placement. Two wall defects with the buccal and lingual plates intentionally removed showed incomplete healing. Botticelli et al. (2005), in a follow-up study, examined the effects of implant surface, implant position and the presence of combined horizontal and vertical residual peri-implant defects on osseointegration in Labrador dogs. After 4 months of healing, regardless of whether the implant was placed in a submerged or nonsubmerged position, a substantial amount of bone fill and a high degree of osseointegration was noted around roughened implants compared to machined implants. This result suggests that implant surface characteristics played an important role in the amount of hard tissue fill and level of osseointegration. Araujo et al. (2005b) studied the effects of immediate implant placement on the dimensional alterations of the alveolar ridge in beagle dogs. They compared sites that received an immediate implant to contralateral sites that received extraction alone over a period of 3 months. Results revealed that marked dimensional alterations, including decrease both in height and width of the ridge, had occurred in the extraction alone sites. The placement of an immediate implant decreased but did not prevent dimensional changes that occurred in the ridge. After 3 months of healing results were similar for both groups.



## Human Extraction Socket Healing Sequence

Various authors studied the extraction socket healing sequence in humans. Amler (1960), examined histologically a total of 75 human extraction sockets over a period of 100 days. In a study of 12 patients requiring extractions of all remaining maxillary teeth, Boyne (1966) examined the histological healing of one of the maxillary first premolar sockets over 23 days. Evian (1982) examined the histologic healing in 10 patients over a period of 16 weeks. Biopsies were taken at 4, 6, 8, 10, 12, and 16 weeks post-extraction. Taken together, these studies showed that the human healing sequence followed a similar pattern to the dog model (Table 4).

**Table 4**

### Human Extraction Socket Healing over 100 Days

<b>Time</b>	<b>Event</b>
Day 1	Blood clot formation
Day 2-3	Granulation tissue appears
Day 4	Contraction of the blood clot begins
Day 7-10	New bone formation
Day 14	1/3 socket filled
Day 20	Connective tissue replaces granulation tissue
Day 38	2/3 socket filled
Day 100	Radiopacity of socket was identical to surrounding bone

The first event in the healing sequence of both human and dogs was the formation of a blood clot at day 1 (Clafin 1936, Amler 1960). The first evidence of new bone formation

in dogs was seen around day 5 and along the lateral aspect of the socket by day 11 (Clafin 1936). In humans, the first evidence of new bone was not detected until day 7-10. Complete socket fill was observed around day 30 in dogs. This is in contrast to human studies where Boyne (1966) reported that only 1/3 of the socket was filled by day 14, and Amler noted that only 2/3 of the socket was filled at day 38. Mature, lamellar bone was seen in dogs at day 90 (Cardaropoli et al. 2003), and this was not evident until day 100 in humans (Amler 1960). Table 5 compares the socket healing sequence for the dog and human models.

**Table 5****Events In Extraction Socket Healing**

<b>Event</b>	<b>Time</b>	<b>Species</b>	<b>Study</b>
Blood Clot Formation	0 to 3 days	Dog	Clafin (1936)
	0 to 1 day	Human	Amler et al. (1960)
Fibroblast Proliferation	3 days	Dog	Clafin (1936)
	2 to 35 days	Human	Amler et al. (1960)
Osteoclast activity	3 to 31 days	Dog	Clafin (1936)
Osteoblast activity	5 to 31 days	Dog	Clafin (1936)
	7 days	Human	Amler et al. (1960)
	10 days	Human	Boyne (1966)
	28 days	Human	Evian et al. (1982)
First evidence of new bone	5 days	Dog	Clafin (1936)
	7-10 days	Human	Amler (1960)
Complete socket fill	30 days	Dogs	Clafin (1936)
1/3 socket fill	14 days	Human	Boyne (1966)
2/3 socket fill	38 days	Human	Amler (1960)
Mature bone present	90 days	Dog	Cardaropoli et al. (2003)
	100 days	Human	Amler (1960)

**Alveolar Ridge Resorption Following Tooth Extraction**

Loss of alveolar bone volume, both width and height, after tooth extraction is a inevitable outcome. The greatest amount of bone loss occurs within the first 2 years after tooth removal (Ashman 2000). Loss of alveolar ridge width and height can complicate placement of an endosseous dental implant since there must be adequate bone to completely surround the dental implant. Whether the residual ridge position is centered

compared to the original edentulous ridge, or it has shifted toward the lingual, is an important consideration. Ridge position can have a significant effect on implant placement, esthetics, and the subsequent occlusal relationship of the restored implant. Previous studies have reported that most ridge resorption occurs on the buccal, resulting in a shift of the center of the ridge toward the palatal/lingual, (Lekovic et al. 1997, Lekovic et al. 1998, Iasella et al. 2003). Pietrokovski and Massler (1967) evaluated 149 dental casts with one tooth missing. They found that the buccal surface of both the maxilla and the mandible resorb more than the lingual/palatal sides with a distinct shift of the center of the ridge to the palatal/lingual. The amount of facial resorption varied considerably between individual sites. Schropp et al. (2003) evaluated study casts from 46 patients with a single premolar or molar extraction over a 12-month period and found that most (2/3) resorption happened within the first 3 months. Yilmaz et al. (1998) examined study casts from 5 patients (10 sites) with a single maxillary incisor extraction that was followed for a 12-month period and noted a 17% decrease in ridge width. Barone et al. (2008) evaluated 40 patients (40 sites) in a non-molar extraction study that was followed for 7 months. He noted a decrease of 41.7% in ridge width. The amount of buccal-lingual ridge resorption after tooth extraction has been reported as 17-63% with the ridge height decreasing by 1 mm, (Lekovic et al. 1997, Lekovic et al. 1998, Yilmaz et al. 1998, Camargo et al. 2000, Schropp et al. 2003, Iasella et al. 2003, Barone et al. 2008). Data from these studies indicated that change in ridge width following tooth extraction varied substantially. Table 6 consists of a list of studies that examined the mean change in the horizontal and vertical ridge dimensions following tooth extraction alone. These resorptive changes in ridge dimension may preclude future implant

placement, or require additional surgical treatment to allow placement of functional, esthetic implants if ridge preservation is not performed at the time of extraction. Table 7 reports the ridge dimensions for the studies and percent change in ridge width.

**Table 6**

**Extraction Alone Studies Showing Change Alone**

<b>Extraction Alone Studies</b>				
<b>Study</b>	<b>Reentry Time (months)</b>	<b>Mean Horizontal Change mm</b>	<b>Percent Horizontal Change</b>	<b>Mean Vertical Change mm</b>
Lekovic et al. 1997	6	-4.43 ± 0.52	-62.9%	-0.88 ± 0.26
Lekovic et al. 1998	6	-4.59 ± 0.23	-61.3%	-1.50 ± 0.21
Yilmaz et al. 1998*	12	-0.75 ± 0.59	-17.0%	-1.35 ± 1.05
Camargo et al. 2000	6	-3.06 ± 2.41	-40.8%	-1.00 ± 2.25
Iasella et al. 2002	4-6	-2.63 ± 2.29	-28.6%	-0.90 ± 1.60
Schropp et al. 2003*	12	-6.1 ± 3.00	-50.8%	-0.20 ± 1.60
Barone et al. 2008	7	-4.5 ± 0.8	-41.7%	-3.60 ± 1.50
<b>Mean</b>	<b>7.6 ± 3.2</b>	<b>-3.7 ± 1.7</b>	<b>-43 ± 17</b>	<b>-1.2 ± 1.1</b>

\* = measured from study casts

**Table 7****Extraction Alone Studies Showing Ridge Dimensions**

<b>Study</b>	<b>Reentry Time (months)</b>	<b>Mean Initial Horiz</b>	<b>Mean Fin Horiz</b>	<b>Mean Horiz Change</b>	<b>% change</b>
Lekovic et al. 1997	6	7.0	2.6	-4.4	-63
Lekovic et al. 1998	6	7.5	2.9	-4.6	-61
Yilmaz et al. 1998*	12	4.7	3.9	-0.8	-17
Camargo et al. 2000	6	7.5	4.4	-3.1	-41
Iasella et al. 2002	4-6	9.1	6.4	-2.6	-29
Schropp et al. 2003*	12	12.0	5.9	-6.1	-51
Barone et al. 2008	7	10.8	6.3	-4.5	-42
<b>Mean</b>	<b>7.6 ± 3.2</b>	<b>8.4 ± 2.5</b>	<b>4.6 ± 1.6</b>	<b>-3.7 ± 1.7</b>	<b>-43 ± 17</b>

**Clinical Studies of Ridge Preservation**

With the emergence of dental implants, ridge preservation has become a frequent part of periodontal plastic and reconstructive surgery. The goal of ridge preservation is minimizing bone loss to preserve the maximum final, healed ridge dimensions. Osseous ridge preservation is done using a hard tissue graft. Without this procedure there may be inadequate ridge width to allow implant placement. Ashman (2000) noted that when an extraction takes place and ridge preservation is not utilized the site of extraction could lose 40% to 60% of bone height and width within 2 to 3 years and subsequent loss of 0.25% to 0.5% annually. Iasella (2003) reported as much as 4 mm loss of ridge width in extraction alone sites within 6 months. Using an atraumatic tooth extraction technique preserves osseous walls thereby improving the chances of osseous graft success. Garg (2001) discussed 5 steps he considered necessary for an atraumatic extraction: 1) do not

reflect the interdental papilla, especially in the esthetic zone; 2) focus on the actual process of tooth removal; 3) use elevators and forceps properly to reduce bony involvement and preserve bone contours; 4) section the tooth to help prevent bone loss; and 5) remove any soft tissue fragments or pathology. Horowitz (2005) added that use of a periosteal elevator is an important adjunct to atraumatic extractions. He says it is used to sever the periodontal ligament fibers, which enables the extraction to be accomplished with significantly less trauma. The greater the number of bony walls present following extraction the more likely the osseous graft will be successful. According to Garg (2001), the bone defect can be categorized into one of the following categories: five-walled, four-walled, three-walled, two-walled, or one-walled defects. Comparison studies have shown that intrasocket ridge preservation prevents most, but not all, ridge resorption. Several ridge preservation studies have used barrier membranes to attempt to improve quality and quantity of bone fill in extraction sites. Both resorbable and non-resorbable barrier membranes have been used; some studies used membranes alone, others used membranes in conjunction with intrasocket grafting materials. Lekovic et al. (1997) compared extraction alone to use of a non-resorbable barrier membrane alone (Gore-Tex®) and Lekovic et al. (1998) compared extraction alone to use of a resorbable barrier membrane alone (Resolut®). In both studies, the teeth included were anterior teeth or premolars. The teeth were atraumatically extracted, the membrane was placed and primary closure was obtained. Reentry was performed 6-months post-extraction. The results showed that both the non-resorbable (Gore-Tex®) and resorbable (Resolut®) barrier membranes provided comparable results. There was mean vertical resorption of 0.35 mm and a mean horizontal resorption of 1.53 mm (20%). Results from Lekovic et

al. (1997, 1998) reveal that the mean horizontal bone loss in the non-resorbable group (Gore-Tex®) was 1.73 mm, which was greater than the mean of 1.32 mm found in the resorbable membrane (Resolut®) group. The extraction alone control group lost a mean of 4.5 mm. The non-resorbable membrane sites had a mean of 3.70 mm (2.5-times) reduced horizontal loss than sites treated with extraction alone while the resorbable membrane sites had a mean of 3.27 mm (3.5-times) reduced horizontal loss. These two studies show that there is not much difference between use of a resorbable vs. a non-resorbable membrane for ridge preservation. Membrane use did, however, greatly decrease the amount of horizontal and vertical bone resorption when compared to extraction alone.

Yilmaz et al. (1998), using study models in a 16-patient, 27-socket study compared the use of bioactive glass (PerioGlas®) in fresh maxillary incisor extraction sites to extraction alone. Sites treated with bioactive glass (PerioGlas®) had a slight gain (0.2 mm) in ridge width, and minimal (0.1 mm) loss of ridge height over a period of 12 months. This was in contrast to the extraction alone group, which demonstrated a much greater loss of ridge width (0.75 mm), and ridge height (1.35 mm).

Camargo et al. (2000), in a 32 nonmolar site ridge preservation study with 6 month re-entry examined the use of bioactive glass (BioGran®) and calcium sulfate (Capset®) to extraction alone. They reported that the mixture of bioactive glass and calcium sulfate resulted in a mean loss of ridge width and height of 3.48 mm and 0.4 mm, respectively. In contrast, the extraction alone group showed slightly less loss in ridge width (3.06 mm), and a greater loss in ridge height (1.0 mm) over 6 months. Iasella et al. (2003) in a 4 to 6-month reentry study used 24 nonmolar sites and compared the use of



freeze-dried bone allograft (FDBA) with a membrane to extraction alone. After four to six months of healing, the sites grafted with FDBA gained 1.3 mm in ridge height and lost only 1.2 mm in ridge width, in comparison to the extraction alone group, which had twice the amount of ridge width loss (2.6 mm), and 0.9 mm ridge height loss.

Barone et al. (2008), examined corticocancellous porcine bone and a collagen membrane to extraction alone in a 40 nonmolar ridge preservation study with a 7 month re-entry. He reported that the corticocancellous porcine bone and collagen membrane group had a mean loss of ridge width and height of 2.5 mm and 0.7 mm, respectively. For the extraction alone group, he reported a mean loss of ridge width and height of 4.3 mm and 3.6 mm, respectively. In a 10 patient case series, Cardaropoli (2008) also studied corticocancellous porcine bone and a collagen membrane over 4 months. He reported a mean loss of 1.8 mm in ridge width after 4 months.

In addition to the extraction alone comparison studies, others have evaluated the effects of various graft materials used to preserve ridge dimensions. Nemcovsky and Serfaty (1996), in a 12-month, 23-patient, 23-socket study using non-resorbable hydroxyapatite (HA) crystals, showed a loss of ridge width of 0.6 mm and a loss of ridge height of 1.4 mm over 1 year. Simon et al. (2000) in a 4-month reentry study using particulate demineralized freeze-dried bone allograft (DFDBA) as an intrasocket and a buccal overlay graft along with a barrier membrane (Resolut XT®), reported an initial ridge width of  $6.2 \pm 0.2$  mm increasing to  $7.3 \pm 0.2$  mm for a gain of 1.1 mm. Zubillaga et al. (2003), in a 10-patient, 11-socket study compared the use of DFDBA (Regenafil®) and a resorbable barrier membrane (Resolut®) with or without fixation at four months. They reported that the mean change in ridge dimensions was a loss of 1.8 mm width, and

a gain of 1 mm height. Vance et al. (2004), in a 4-month nonmolar reentry study using 24 extraction sockets compared the use of anorganic bovine bone matrix (BioOss®) with a membrane to DFDBA plus mixture of calcium sulfate and carboxymethylcellulose (CalMatrix®). They demonstrated that both groups had a mean loss of 0.5 mm ridge width. The BioOss® group showed a gain in mean ridge height of 0.7 mm, while the CalMatrix® group showed a mean loss of 0.3 mm. Adams et al. (2005) compared two different ridge preservation techniques in nonmolar sites in a 4 month re-entry study. An intrasocket FDBA graft alone was compared to an intrasocket plus a buccal overlay (extrasocket) FDBA graft. The intrasocket alone group had a mean ridge width loss of 2 mm and no change in ridge height. In contrast, the overlay group showed a mean ridge width loss of 1.4 mm and a gain of 2.2 mm of ridge height. Brkovic et al. (2008) in a single case report evaluated an alveolar preservation technique involving placement of a cone of beta-tri-calcium phosphate (TCP) combined with type I collagen without the use of a barrier or flap. Nine months after tooth extraction, they reported no reduction in ridge height and no change in ridge width (12 mm). Neiva et al. (2008) in a 24 patient study over 4 months compared an anorganic bovine-derived hydroxyapatite matrix combined with a synthetic P-15 (Putty P15) and a bioabsorbable collagen wound dressing to a bioabsorbable wound dressing alone. Neiva reported a loss of 1.31 mm in ridge width and a gain of 0.15 mm in ridge height for the Putty P15 group. For the bioabsorbable collagen wound dressing alone, a loss of 1.43 mm for ridge width and a loss of 0.56 mm in ridge height was reported (Table 8,9).

## **Polytetrafluoroethylene Technique Studies**

Traditionally, porous non-resorbable membranes such as the expanded polytetrafluoroethylene (ePTFE) membrane and resorbable membranes had complications during guided bone regeneration (GBR) procedures. The ePTFE membrane had high rates of infection due to membrane exposure and its porous nature (Bartee 2001). Most resorbable membranes are type I collagen or type I-III collagen which have necessitated primary closure to prevent exposure to oral environment. Infections do not occur as frequently with resorbable membranes as with the ePTFE membrane; however, degradation of the membrane does occur with membrane exposure. To counter the problems of the ePTFE membrane and resorbable membranes, Bartee (2001) developed a dense polytetrafluoroethylene (PTFE) membrane.

According to Bartee (2001), the dense PTFE membrane offers 4 primary advantages over the ePTFE and resorbable membranes in extraction site reconstruction:

- 1) Due to the low porosity (< 0.3 micrometers), the dense membrane resists the incorporation of bacteria into its structure and can be left exposed in the mouth with a low risk of infection and subsequent graft loss. Exposure of the membrane does not compromise the underlying bone graft.
- 2) The ability of the membrane to remain exposed also reduces the need for the development of large flaps and vertical incisions to achieve primary closure. The nonresorbable polymer prevents premature degradation associated with exposure of resorbable membranes.
- 3) Conservation of soft tissue architecture is achieved since primary closure is not required. There is no loss of

vestibular depth, and the attached mucosa and interdental papilla can be preserved by using careful surgical technique. 4) The membrane does not allow ingrowth of the surrounding connective tissues, and removal is accomplished without anesthesia, surgery, or trauma to the adjacent tissues.

The procedure is indicated following the extraction of single or multiple teeth. Active infection is the only absolute contraindication. Barteo (2001) describes the PTFE ridge preservation technique as follows: 1) The first rule of ridge preservation is nontraumatic extraction. 2) Following root removal, sharp curettage should be carried out to remove remnants of periodontal ligament as well as any soft tissues such as periradicular cysts. These tissues may harbor pathogenic bacteria that may lead to postoperative complications. 3) Perforation of the socket cortical plate (decortication) is optional but may be helpful in establishing blood supply to the graft from the adjacent bone. 4) Using a periosteal elevator or syringe, the graft material is delivered to the extraction site and packed gently to the apex of the site. Overpacking is to be avoided because this only hinders revascularization of the site. 5) A section of membrane material is then cut to fit over the site extending 3 to 4 mm beyond the socket margins onto sound host bone. The membrane should be trimmed to maintain a 1.0 mm margin from adjacent tooth root to facilitate reattachment of the papilla to the interdental bone. The membrane should fit over the site and under the mucoperiosteal flap without wrinkling or buckling. 6) Suturing is accomplished with interrupted sutures at the interdental papillae and a single or horizontal mattress suture across the socket opening. The recommended suture material is 3-0 polyglycolide (Vicryl, Ethicon Inc, Somerville, NJ) or PTFE Monofilament. Excessive tension on the flaps should be avoided to

maximize blood flow within the flap and avoid necrosis of the flap margins. 7)  
Thorough irrigation of the site to remove all remaining graft particles is done. 8)  
Postoperatively, the patient should be observed at 1 week. At 2 weeks, sutures should be removed and the membrane cleaned if there is significant bacterial accumulation. 9)  
Membrane removal is done at either 3 or 4 weeks postoperatively, depending on the size of the defect and the condition of the walls. Removal is accomplished by grasping the membrane with forceps and gently removing it from the tissue bed. No anesthesia is required for this procedure, however topical anesthetic may be used. Upon removal, the graft material can usually be visualized, well consolidated in the osteoid matrix underlying the membrane. Re-epithelialization of the underlying tissue will occur over the next 7 to 10 days.

In a 4 patient case series, Bartee (1998) describes the histologic findings of PTFE ridge preservation and implant guided tissue regeneration (GTR). Two patients had ridge preservation alone and two patients had immediate implants with grafting. The patients received a graft paste consisting of 60% human freeze-dried demineralized bone (FDDB) granules (Dembone 300-500 micrometers and low-density, and 40% resorbable calcium phosphate (OsteoGraft/LD 300 OsteoGen). After adequate reflection of the mucoperiosteal flaps, the membrane was trimmed with sharp scissors and placed over the extraction site, extending 3 to 5 mm beyond the defect. Tissue samples were demineralized and stained with hematoxylin and eosin. New bone formation was clearly evident in all tissue sections. No areas of inflammatory infiltrate were noted. The bone graft particles were observed in various states of dissolution, resorption, or remodeling.

The bone was mature, dense lamellar bone. Overall, the regenerated tissue was well vascularized and there were no areas of fibrosis or chronic inflammation.

Horowitz (2005) reviewed the nonexpanded polytetrafluoroethylene (TefGen-FD) barrier in 2 case reports. The patients in this study had sites that were treated simply by protecting a blood clot with a removable, nonexpanded PTFE barrier over the extraction socket. This investigation evaluated the early (3 to 6 month range) healing and ingrowth of vital bone into an extraction socket and the maturation of overlying soft tissue. "To help promote GBR and protection of the healing socket while maintaining stability of the barrier, it is placed directly onto the outersurface of the bone and overlaps the facial and lingual walls of the socket". Proximal contouring of the material is performed to leave 1 mm of bone adjacent to the proximal teeth, so a blood supply can be reestablished from the alveolar bone to the papillae. No attempt is made to attain primary closure of the socket. Clinically, when an ePTFE membrane becomes exposed, bacteria penetrate the site and require a secondary surgery for removal. "Nonexpanded PTFE barriers are removed when one edge becomes exposed to the oral cavity, generally 3 to 6 weeks after insertion." Hoffmann et al. (2008) evaluated the non-porous, non-resorbable membrane in a retrospective private practice study consisting of 276 sockets in 276 subjects after 8 months of healing. He evaluated the dPTFE (Cytoplast Regentex GBR-200) membrane, a membrane made of high density polytetrafluoroethylene which does not require primary closure. The sockets were grouped as either single sockets or side-by-side sockets. His results for ridge height show that about 50% of the sockets had 0.5 mm of bone loss at the buccal site and 50% of the sockets had 1.0 mm at the buccal site. For ridge width, 50% of the sockets had 0.0 mm bone loss and 50% had 0.5 mm bone loss.

Histologic evaluation indicate that the newly formed tissue in the socket was mainly regular trabecular bone and typical cells indicating normal healing mechanisms were not impaired. He concluded that the use of dPTFE allows for the preservation of ridge width and height, however the treatment outcome is mainly limited by the architecture of the existing bony walls.

### **Bio-Col Technique Studies**

Sclar (2000) developed the 'Bio-Col' technique for preserving alveolar ridge anatomy following tooth removal in esthetic areas. He says the clinical goal of any ridge preservation technique should be to preserve both the hard and soft tissues following tooth removal, especially the interdental papillae, in such a way to optimize esthetics and function. "Maintaining a stable osteoconductive scaffold within the entire area of the socket that is slowly resorbed and eventually replaced by vital bone and isolating this scaffold from the deleterious effects of the oral environment during healing is an essential biologic consideration (Sclar 2000)."

The steps involved in this technique are as follows: 1) The tooth is extracted atraumatically without flap reflection. 2) Perforation of the socket walls in order to promote bleeding and enhance the invasion of osteoprogenitor cells. 3) The socket is grafted with deproteinized bovine bone mineral (Bio-Oss: Osteohealth Co, Shirley, NY), and isolated from the oral environment with absorbable collagen dressing (CollaPlug, Sulzer Calitek, Inc., Carlsbad, CA). 4) It is sealed with cyanoacrylate, an impervious tissue cement (Isodent: Ellman International, Hewlett, NY). This allows for guided bone

regeneration without the need for flap elevation and primary closure, thus preserving the surrounding soft-tissue volume (Sclar 2004). 5) An interim provisional restoration of ovate pontic design that replicates the contours of the tooth that was removed, supports the surrounding soft tissues and avoids implant loading when an immediate implant is placed.

The author's desire in developing the technique was to isolate the grafted socket and obtain a membrane effect without the elevation and advancement of large mucoperiosteal flaps that result in soft-tissue disfigurement and loss of volume at the site. When used in conjunction with immediate implant placement, the Bio-Col technique results in high rates of osseointegration (98.3% 58 sites follow-up time of 10-63 months) and excellent esthetics. Retrospective analysis also revealed a successful osseointegration rate of 94% for 248 sites treated with the Bio-Col in conjunction with delayed implant placement with a follow-up ranging from 6 to 73 months.

Sclar's selection of Bio-Oss as a bone graft material is based on Bio-Oss's osteoconductive properties. Bio-Oss consists of the mineral portion of bovine bone and provides the body with a matrix for bone cell migration. It is also integrated during the natural remodeling process of the human bone and slowly resorbed due to small crystallite size which is comparable to human bone ([www.Osteohealth.com](http://www.Osteohealth.com)).

Fowler and Whicker (2004) revealed a modification to the Bio-Col technique in a case report. They report the modification simplifies the procedure without compromising the esthetic result. The modifications consist of 4 changes to Sclar's description. First, the CollaPlug wound dressing is used in a significantly smaller quantity simply to cover the Bio-Oss graft, not layered to the level of the free gingival margin. On average only



the terminal 1/5 of the entire plug is utilized. Second, the horizontal mattress suture is eliminated. Suturing is only done if soft tissue trauma occurred (i.e. the interdental papilla is torn). Third, when a fixed provisional is utilized, the cyanoacrylate is not applied to the CollaPlug. Instead, the provisional is first temporarily cemented and the cyanoacrylate is placed at the gingival margin-pontic interface to “seal” this area. Finally, it is recommended the provisional be removed and modified between 3-6 weeks post-surgery.

As demonstrated by the aforementioned studies, despite the use of ridge preservation techniques to minimize the amount of bone resorption after an extraction, some loss of vertical and horizontal dimensions may still occur. On the other hand, if ridge preservation was not performed, a substantial decrease in the horizontal dimension of the ridge, ranging from 17-63% (0.75 to 6.1 mm) over 4-6 months can be anticipated, which may create enough deficiency of bone to preclude implant placement (Lekovic et al. 1997, Lekovic et al. 1998, Iasella et al. 2003, Schropp et al. 2003, Barone et al. 2008).

**Table 8**

**Ridge Preservation Studies Showing Change Alone**

Study	Reentry Time months	Treatment	Mean Horizontal Change mm	Percent Horizontal Change	Mean Vertical Change mm
Nemcovsky & Serfaty 1996	12	Nonresorbable HA crystals	-0.6 ± 0.66	N/A‡	-1.4 ± 0.50
Lekovic et al. 1997	6	ePTFE	-1.7 ± 0.56	-23.3%	-0.3 ± 0.26
Lekovic et al. 1998	6	Resolut	-1.3 ± 0.21	-17.6%	-0.4 ± 0.20
Yilmaz et al. 1998	6	PerioGlas cones	+0.2 ± 0.52	+3.6%	-0.1 ± 0.87
Camargo et al. 2000	6	BioGran Capset	-3.5 ± 2.68	-44.3%	-0.4 ± 3.18
Simon et al. 2000	4	DFDBA/ Resolut XT®	+1.1 ± NG*	+18%	-1.4 ± NG*
Iasella et al. 2003	4	FDBA/ BioMend	-1.2 ± 0.93	-13.0%	+1.3 ± 2.00
Zubillaga et al. 2003	4	Regenafil	-1.8 ± NG*	-16.8%	+1.0 ± NG*
Vance et al. 2004	4	BioOss/ BioGide	-0.5 ± 0.8	-5.2%	+0.7 ± 0.4
Vance et al. 2004	4	CalMatrix/ Capset	-0.5 ± 0.8	-5.6%	-0.3 ± 0.6
Barone et al. 2008	7	xenograft, collagen mem	-2.0 ± 0.9	-23.6%	-0.7 ± 1.4
Brkovic et al. 2008	9	B-TCP + coll	-1.4 ± 1.0	0.0%	0.0
Cardaropoli et al. 08	4	xenograft/coll membrane	-1.9 ± 1.7	-16.1%	NA
Neiva et al. 2008	4	P15/Collaplug	-1.3 ± 0.9	NA	+0.2 ± 1.8
Neiva et al. 2008	4	Collaplug	-1.4 ± 1.1	NA	-0.6 ± 1.0
<b>Mean ± sd</b>			<b>-1.1 ± 1.1</b>	<b>-12 ± 15</b>	<b>0.0 ± 0.8</b>

\* NG = not given in article

‡ = no baseline measurements reported, unable to determine percentage

**Table 9****Ridge Preservation Studies Showing Ridge Dimensions**

<b>Study</b>	<b>Reentry Time (months)</b>	<b>Mean Initial Horiz</b>	<b>Mean Fin Horiz</b>	<b>Mean Horiz Change</b>	<b>% change</b>
Nemcovsky & Serfaty 1996	12			-0.6	
Lekovic et al. 1997	6	7.3	5.6	-1.7	-23
Lekovic et al. 1998	6	7.4	6.1	-1.3	-18
Yilmaz et al. 1998	6	5.5	5.7	+0.2	+4
Camargo et al. 2000	6	7.9	4.4	-3.5	-44
Simon et al. 2000	4	6.2	7.3	+1.1	+18
Iasella et al. 2003	4	9.2	8.0	-1.2	-13
Zubillaga et al. 2003	4	10.7	8.9	-1.8	-17
Vance et al. 2004	4	8.9	8.4	-0.5	-6
Vance et al. 2004	4	9.7	9.2	-0.5	-5
Barone et al. 2008	7	10.6	8.1	-2.5	-24
Brkovic et al. 2008	9	12.0	12.0	0.0	0
Cardaropoli et al. 08	4	11.8	9.9	-1.9	-16
Neiva et al. 2008	4			-1.3	
Neiva et al. 2008	4			-1.4	
<b>Mean</b>	<b>5.6 ± 2.3</b>	<b>8.9 ± 2.1</b>	<b>7.8 ± 2.1</b>	<b>-1.1 ± 1.1</b>	<b>-12 ± 16</b>

**Histologic Evaluation of Ridge Preservation**

The goal of ridge preservation procedures is to prevent the collapse of the ridge by allowing the alveolar socket to fill in with as much bone volume as possible. The ideal bone grafting material will promote vital host bone to rapidly fill the socket and minimize the loss of ridge dimensions. It is very important to determine histologically how much

bone is present relative to the amount of trabecular space since this measure is a reflection of bone quality that may influence implant placement. A bone quality index has been described by Lekholm and Zarb (1985) which includes Type I bone being homogenous compact bone, Type II being a thick layer of compact bone surrounding a core of dense trabecular bone, Type III being a thin layer of cortical bone surrounding dense trabecular bone of favorable strength and Type IV being a thin layer of cortical bone surrounding a low-density trabecular bone. Type I bone is preferred for implant placement since it has the highest density of cortical bone and Type IV is the least preferred due to its very low density.

### **Extraction Alone Studies**

When extraction sockets are left alone and heal without any type of ridge preservation procedure the amount of vital bone present after 4-8 months of healing ranges from 26-54% with 46-67% of trabecular space (Iasella et al. 2003, Froum et al. 2002, Serino et al. 2003, Barone et al. 2008). In the canine model performing extraction alone in 9 sockets, Cardaropoli et al. (2005) reported only 15% vital bone and 85% trabecular space over 6 months. Histologic results from autogenous bone grafts have shown vital bone (osteocytes within the lacunae), non-vital bone (residual graft particles), vascular channels, osteoblasts and secondary osteon formation. Cement lines usually surround the non-vital bone, which joins the immature new bone with the non-vital bone chips (Becker et al. 1994, 1996, 1998)(Table 10).

## **Allograft Studies**

Allografts are usually available in one of two forms: mineralized particulate freeze-dried bone allograft (FDBA) and demineralized particulate freeze-dried bone allograft (DFDBA). FDBA provides an osteoconductive scaffold while DFDBA may provide osteoinductive proteins in addition to the osteoconductive scaffold (Mellonig et al. 1981, Mellonig 1991). The osteoinductive properties of DFDBA have been attributed to the presence of bone morphogenetic protein (BMP). Urist et al. (1971) isolated BMP from human cortical bone. He placed them in ectopic sites in athymic mice and found that they initiated bone formation. The demineralization process of allograft preparation releases BMP and allows osseous induction to occur. The donor age and health status can also affect the osteoinductive potential. Schwartz et al. (1996, 1998, 2000) found that there is a wide variation in the osteoinductive capabilities of commercial DFDBA from different bone banks. There was an age-dependent decrease in the new bone induction score as measured by histomorphometric analysis. Donors over the age of 50 showed significantly less induction ability, but there were no differences attributable to gender. Studies of demineralized freeze-dried bone allograft (DFDBA) used in ridge preservation procedures have reported conflicting results. Several studies have found that non-vital DFDBA particles are still present in biopsy cores. (Smukler et al. 1999, Froum et al. 2002)(Table 11). It has also been reported that DFDBA has osteoinductive properties and should induce bone growth, but in several histologic samples the DFDBA particles were encapsulated in fibrous connective tissue with no evidence of either osteoblastic or osteoclastic activity (Becker et al. 1994, 1996, 1998). If DFDBA particles do not provide any osteoinductive properties, it is believed they might interfere with normal bone

formation and may weaken the bone at the grafted site (Becker et al. 1994). The amount of non-vital graft particles remaining relative to the amount of vital bone may be an important factor. Several studies have reported that DFDBA particles do resorb and in some cases fully resorb leaving only vital bone (Vance et al. 2004). Histologic examination reveals that ridge preservation utilizing DFDBA has residual graft particles surrounded by intimately apposed woven and lamellar bone with distinct cement lines and a lack of fibrous encapsulation. Osteoblasts lined endosteal spaces and the new bone marrow exhibited a mild degree of fibrosis without signs of an inflammatory reaction (Brugnami et al. 1996, 1999, Smukler et al. 1999). Vance et al. (2004) examined 12 sockets grafted with a combination of DFDBA and an alloplastic putty consisting of calcium sulfate and carboxymethylcellulose (CalMatrix®) over 4 months. They reported 61% vital bone, 3% non-vital bone, and 36% trabecular space. The percentage of vital bone present after utilizing DFDBA in ridge preservation ranged from 35 to 60% with only 3-14% having non-vital bone (Smukler et al. 1999, Froum et al. 2002). Becker et al. (1996, 1998) reported more residual graft particles and fibrous encapsulation, which may be due to their failure to use an occlusive barrier membrane.

Freeze-dried bone allograft (FDBA), has also been used in ridge preservation procedures and showed a histologic result of 28% vital bone, 37% non-vital bone and 35% trabecular space over 4-6 months (Iasella et al. 2003). The residual FDBA particles were often surrounded by vital woven or lamellar bone, or were encapsulated in fibrous connective tissue. The residual graft material was higher than the amount with DFDBA, which may be due to the shorter healing period of 4-6 months vs. up to 48 months for DFDBA. Wang et al. (2008), grafted five patients with solvent preserved mineralized

particulate cancellous allograft (Puros). After 5 to 6 months they reported 69% vital bone 3.8% residual graft particles and 27% trabecular space. Comparison of the two grafting materials is difficult since the healing periods are different for each of the studies.

### **Xenograft Studies**

Xenografts, mostly anorganic bovine bone, have also been utilized in ridge preservation procedures with similar results to allografts. Generally, bone encircled and adhered to the grafted particles in a concentric and/or lamellar arrangement. Newly formed bone was observed, mostly in direct connection with the grafted particles (Artzi et al. 1998, 2001, Froum et al. 2004)(Table 11). Vance et al. (2004) showed that BioOss® had 26% vital bone with 16% non-vital bone and 58% trabecular space after 4 months of healing. This agrees with a 6-month study of 6 sockets grafted with BioOss® by Zitzmann et al. (1997, 2001) where they reported 27% vital bone, 30% non-vital bone, and 43% trabecular space. In contrast, Artzi et al. (2000) in a 9-month study, grafted 15 sockets in 15 patients using BioOss® and reported a much greater percentage of vital bone at 46%, along with 31% non-vital bone, and 23% trabecular space. Froum et al. (2004) in a 6 to 8 month study grafted 8 sockets with a nonresorbable anorganic bovine bone substitute (OsteoGraf R/N-300), 4 of which were combined with an ePTFE barrier, and the other 4 with Alloderm (ADM) as a barrier. In the OsteoGraf/ePTFE group, there was 18% vital bone, 21% non-vital bone, and 61% trabecular space. In the OsteoGraf/ADM® group, 42% vital bone, 13% non-vital bone, and 45% trabecular space. The difference in the amount of vital bone between the two groups could possibly

be attributed to the choice of barrier used. The vascular channels in the Alloderm may have provided better revascularization compared to the ePTFE barrier. Araujo et al. (2008), grafted one quadrant of fresh extractions sockets in mongrel dogs with Bio-Oss Collagen. After 3 months healing there was 27% bone marrow, 58% vital bone, and 12% residual particles of Bio-Oss Collagen. In a 40 patient study, Barone et al. (2008) compared grafting 20 sockets with OsteoBiol MP3 and a collagen membrane (OsteoBiol Evolution) to extraction alone over 7 months. In the OsteoBiol MP3/Evolution group, they reported 36% vital bone, 29% non-vital bone, and 37% connective tissue. The vital bone of 36% falls in the middle of xenograft histologic studies. The extraction alone group resulted in vital bone of 26% and 59% connective tissue. Neiva et al. (2008) reported on a 24 patient study examining a putty-form anorganic bovine-derived hydroxyapatite matrix combined with a synthetic cell-binding peptide P-15 (Putty P15) and a bioabsorbable collagen membrane to a bioabsorbable collagen dressing alone. He reported the Putty P15 having 29.92% vital bone, 65.25% bone marrow and 6.25% non-vital. The bioabsorbable group was reported to have 36.54% vital bone and 62.67% bone marrow.

### **Alloplast Studies**

Alloplastic materials such as bioactive glass, hydroxyapatite (HA) and calcium sulfate have been shown to have vital bone from 25 to 60% (MacNeill et al. 1999, Froum et al. 2002, 2004 Guarnieri et al. 2004, and Mangano et al. 2008). Alloplasts are well tolerated by the host and tend to be osteoconductive in nature rather than osteoinductive.



Guarnieri et al. (2004) in a 10 socket study utilizing medical grade calcium sulfate hemihydrate, found at 3 months that 100% of the graft had been resorbed and that there was 58% vital bone present throughout the preservation site. The site was also devoid of any inflammatory cells and connective tissue. The resorption time with calcium sulfate is much faster than the xenografts or the allografts previously mentioned. Mangano et al. (2008) on the other hand, discussed the slow rate of resorption of hydroxyapatite in a 20 year case report. Mangano et al (2008) reported using very dense HA with a mean size of 1 to 2 micrometers in a mandibular cuspid socket. After 20 years of follow up, the socket demonstrated 25% vital bone, 41% marrow space, and 38% residual HA particles. The author reports the slow rate of resorption due to an intimate binding between a patient's bone and HA particles. MacNeill et al. (1999) compared the osseous healing of 4 different alloplasts: hydroxyapatite (HA, OsteoGraf/P), bioactive glass #1 (BioGran® 300-360  $\mu\text{m}$ ), bioactive glass #2 (PerioGlas® 90-710  $\mu\text{m}$ ), and calcium sulfate (Capset®) with autogenous bone, in osteotomy sites surgically created in the rabbit tibia over 28 days. All graft sites showed evidence of new bone formation at one month with the Capset plus autogenous bone showing the greatest mean percentage of vital bone (58.8%) and PerioGlas® showing the least (40.4%), while the BioGran and OsteoGraf/P group both showed 41.8% vital bone. Froum et al. (2002) treated 19 human sockets with BioGran® and reported similar results with 59% vital bone, 6% non-vital bone, and 35% trabecular space over 6-8 months. Froum et al. (2004) treated 8 sockets with absorbable HA (OsteoGraf R/LD), 4 of which were combined with an ePTFE barrier, and the remaining 4 with an Alloderm® (ADM) barrier. After 6-8 months of healing, the HA/ADM group showed 35% vital bone, 4% non-vital bone, and 62% trabecular space,

while the HA/ePTFE group showed 28% vital bone, 12% non-vital bone, and 61% trabecular space (Table 11). In contrast, Luczyszyn et al. (2005) grafted 15 sockets in 11 patients using bioabsorbable HA (Algipore®) with an ADM barrier over 6 months. They reported only 1% vital bone, 42% non-vital bone, and 57% trabecular space. Serino et al. (2003), in a non-graft study, treated 34 sockets in 32 patients over 6 months with a bioabsorbable polylactide/polyglycolic acid sponge (Fisiograft®) to encourage vascular ingrowth. They reported 67% vital bone and 33% trabecular space. These results compare well to the results seen by Vance et al. (2004) with DFDBA and the calcium sulfate putty (CalMatrix®) and Guarnieri et al. (2004) with the medical grade calcium sulfate. In a single case report, Brkovic et al. (2008) evaluated beta-TCP with type I collagen (RTR Cone, Septodont, Saint-Maur-des-Fosses, France) and reported 62.6% vital bone, 21.1% marrow and 16.3% residual B-TCP graft. This is the highest percentage of vital bone reported of the alloplasts.

### **Summary of Histologic Findings**

When analyzing histologic findings the studies demonstrate that when ridge preservation procedures are performed with a variety of grafting materials, including allografts (DFDBA, FDBA), xenografts (anorganic bovine bone mineral), or alloplasts (hydroxyapatite, calcium sulfate, and polylactide/polyglycolic acid sponge), the percentage of vital and nonvital bone as well as trabecular space varies considerably. The percentage of vital bone ranged from 1-67%, the percentage of non-vital bone ranged from 0-42%, and the percentage of trabecular space ranged from 33-85%.

**Table 10****Comparison of Histologic Data on Extraction Alone studies**

Author/Yr	Species	Healing Months	% Vital Bone	% Trabecular Space
Froum et al. 2002	Human	6-8	32.4	67.6
Iasella et al. 2003	Human	4-6	54.0	46.0
Serino et al. 2003	Human	6	44.0	56.0
Barone et al. 2008	Human	7	26	59.0
<b>Mean ± sd</b>		<b>6 ± 1</b>	<b>39 ± 12</b>	<b>54 ± 14</b>

**Table 11****Comparison of Histologic Data on Ridge Preservation studies**

Author/Yr	Graft Material	Particle Size	Healing Months	% Vital Bone	% Non-Vital Bone	% Trabecular Space
<b>Allografts</b>						
Froum et al. 2002	DFDBA	250 to 500 $\mu\text{m}$	6-8	34.7	13.5	51.8
Iasella et al. 2003	FDBA	500-1000 $\mu\text{m}$	4-6	30.1	34.7	35.2
Vance et al. 2004	DFDBA/putty (CalMatrix®)	500-1000 $\mu\text{m}$	4	61.0	3.0	36.0
<b>Mean ± sd</b>				<b>41 ± 17</b>	<b>18 ± 17</b>	<b>38 ± 13</b>
<b>Xenografts</b>						
Artzi et al. 2000	BioOss®	250-1000 $\mu\text{m}$	9	46.3	30.8	42.6
Zitzmann et al. 2001	BioOss®	250-1000 $\mu\text{m}$	6	26.9	30.5	42.6
Froum et al. 2004	OsteoGraf R/N300 + ADM	250-420 $\mu\text{m}$	4	42.0	13.0	45.0
Froum et al. 2004	OsteoGraf R/N300 +ePTFE	250-420 $\mu\text{m}$	4	18.0	21.0	61.0

Vance et al. 2004	BioOss®	250-500 $\mu\text{m}$	4	26.0	16.0	54.0
Barone et al. 2008	OsteoBiol MP3 + OsteoBiol Evolution	600-1000 $\mu\text{m}$	7	35.5	29.2	36.6
Cardaropoli et al. 2008	OsteoBiol GenOs + OsteoBiol Evolution	250-1000 $\mu\text{m}$	4	NR	24.5	NR
Neiva et al. 2008	Putty P-15 + collaPlug	250-420 $\mu\text{m}$	4	29.9	6.3	65.3
<b>Mean</b>				<b>31 <math>\pm</math> 9</b>	<b>23 <math>\pm</math> 11</b>	<b>47 <math>\pm</math> 14</b>
<b>Alloplasts</b>						
Froum et al. 2002	Bioactive Glass (BioGran®)	300-355 $\mu\text{m}$	6-8	59.5	5.5	35.0
Froum et al. 2004	HA (OsteoGraf R/LD) + ADM	250-420 $\mu\text{m}$	4	35.0	4.0	62.0
Froum et al. 2004	HA (OsteoGraf R/LD) + ePTFE	250-420 $\mu\text{m}$	4	28.0	12.0	61.0
Luczyszyn et al. 2005	HA (Algipore®) + ADM	NA	6	1.0	42.0	57.0
Brkovic et al. 2008	B-TCP, Type 1 collagen	500-1000 $\mu\text{m}$	9	62.6	16.3	21.1
Mangano et al. 2008	dense HA	1 to 2 $\mu\text{m}$	240	25.4	38.1	41.3
<b>Mean</b>				<b>35 <math>\pm</math> 23</b>	<b>20 <math>\pm</math> 16</b>	<b>46 <math>\pm</math> 17</b>
<b>Membrane Alone</b>						
Luczyszyn et al. 2005	ADM	NA	6	46.0	0.0	54.0
<b>Collagen Filler Material</b>						
Serino et al. 2003	Poly lactide/ Polyglycolic acid sponge (Fisiograft®)	NA	6	67.0	0.0	33.0
Neiva et al. 2008	Collaplug	NA	4	36.5	0.0	62.7
<b>Mean</b>				<b>52 <math>\pm</math> 21</b>	<b>0 <math>\pm</math> 0</b>	<b>48 <math>\pm</math> 21</b>

\*NR= not reported in article

## Summary of Literature Review

Extraction alone studies have utilized both animal and human models. The healing sequence of an extraction socket begins with the formation of a blood clot around day 1, followed by neovascularization around day 3, and subsequent new bone formation starting at around 5-7 days (Clafin 1936, Cardaropoli et al. 2003, Amler 1960, Boyne 1966, Evian 1982). Complete socket fill was noted at day 30 in dogs (Clafin 1936), while only 2/3 of the socket was filled in humans at day 38 (Amler 1960). Mature, lamellar bone was seen as early as 90 days in dogs (Cardaropoli et al. 2003), and this was not present until day 100 in humans (Amler 1960).

Studies of the histologic healing of the extraction sockets have shown that without any type of ridge preservation procedure the amount of vital bone present after 4-8 months of healing ranges from 33-54% with 34-67% of trabecular space (Iasella et al. 2003, Froum et al. 2002, Serino et al. 2003). In contrast, in the canine model, Cardaropoli et al (2003) reported only 15% vital bone and 85% trabecular space after 6 months of healing. Significantly, Araujo et al. (2005), in an 8-week study using the canine model, reported that the bundle bone began to disappear as early as 2-weeks post-extraction, and that the buccal wall undergoes a greater amount of resorption than the lingual wall.

Histologic results from autogenous bone grafts have shown mostly vital bone (osteocytes within the lacunae). Studies using allografts (DFDBA, FDBA) for ridge preservation (Smukler et al. 1999, Froum et al. 2002, Vance et al. 2004, Iasella et al. 2003) have yielded variable results. Percentage of vital bone ranged from 30-61%,

% non-vital bone ranged from 3-35%, while percentage trabecular space ranged from 35-56%. This broad range of results could be attributed to the range in reentry times from 4-23 months. Ridge preservation studies using xenografts (BioOss®, OsteoGraf) showed similar results to allografts with a range of 18-46% of vital bone, 13-31% of non-vital bone, and 43-61% of trabecular space. A broader range of results was seen with studies using alloplasts (BioGran®, PerioGlas®, Algipore®, hydroxyapatite, calcium sulfate) with re-entry times from 1 to 8 months. From these studies, a range of 1-60% vital bone, 4-42% non-vital bone, and 35-57% trabecular space was reported. Lastly, (Serino et al. 2003), examined the use of a polylactide/polyglycolic acid sponge (Fisiograft®) for ridge preservation and they reported 67% vital bone, an absence of non-vital bone, and 33% trabecular space.

Alveolar ridge resorption has been reported as a common sequelae following tooth extraction. Loss of alveolar ridge width and height can be problematic if a dental implant is selected for tooth replacement. While the dimensions of the healed alveolar ridge determine the feasibility of placement of a dental implant, the immediate, post-extraction ridge dimensions may be predictive of the final outcome. In other words, both wide and narrow sockets will lose horizontal width. Thus, if a narrow socket is present initially, the final result may be too narrow to accommodate implant placement. Table 10 summarizes the root dimensions at the cervix as categorized by tooth types.

**Table 12****Root Dimensions at the Cervix by Tooth Types (Ash-Wheeler 6<sup>th</sup> Edition 1984, Woelfel 1990)**

Tooth Types	Bucco-lingual/palatal dimensions mm		Mesio-distal dimensions mm	
	Ash-Wheeler	Woelfel	Ash-Wheeler	Woelfel
Mandibular incisors				
Central	5.3	5.4	3.5	3.5
Lateral	5.8	5.8	4.0	3.8
Maxillary incisors				
Central	6.0	6.4	7.0	6.4
Lateral	5.0	5.8	5.0	4.7
Mandibular & Maxillary canines	7.0	Mx: 7.6 Mn: 7.5	5.5	Mx: 5.6 Mn: 5.2
Mandibular 1 <sup>st</sup> premolars	6.5	7.0	5.0	4.8
Mandibular 2 <sup>nd</sup> premolars	7.0	7.3	5.0	5.0
Maxillary premolars (1 <sup>st</sup> & 2 <sup>nd</sup> )	8.0	1 <sup>st</sup> : 8.2 2 <sup>nd</sup> : 8.1	5.0	1 <sup>st</sup> : 4.8 2 <sup>nd</sup> : 4.7
Mandibular 1 <sup>st</sup> molars	9.0	10.7	9.0	7.9
Mandibular 2 <sup>nd</sup> molars	9.0	10.7	8.0	7.6
Mandibular 3 <sup>rd</sup> molars	9.0	10.4	7.5	7.2
Maxillary 1 <sup>st</sup> molars	10.0	9.0	8.0	9.2
Maxillary 2 <sup>nd</sup> molars	10.0	8.8	7.0	9.1
Maxillary 3 <sup>rd</sup> molars	9.5	8.9	6.5	9.2

As is evident from Table 12, different tooth types possess different bucco-lingual/palatal and mesio-distal dimensions. In general, incisors are the smallest, while molars are the widest in dimension. As a result, ridge preservation becomes increasingly critical for the smaller tooth types (especially mandibular incisors) since even a small amount of horizontal ridge resorption can be detrimental.

Despite the use of a bone graft to preserve alveolar ridge dimensions, most studies have reported a net loss in horizontal and/or vertical ridge dimensions. Simon et al.

(2000) in a 4-month reentry study using particulate DFDBA as an intrasocket and a buccal overlay graft along with a barrier membrane (Resolut XT®); however, reported a mean net gain of approximately 1.1 mm of ridge width.

The goal of ridge preservation is to minimize the amount of ridge resorption after extraction. As was evident from the extraction alone studies reviewed (Lekovic et al. 1997, Lekovic et al. 1998, Yilmaz et al. 1998, Camargo et al. 2000, Iasella et al. 2002, Schropp et al. 2003), the change in ridge width following tooth extraction varies substantially, and this broad range (30-60%) may have a profound influence on the future tooth replacement options available.

The University of Louisville has studied ridge preservation since 2003 starting with Iasella. Since that time horizontal ridge width change has ranged from -0.5 to -2.0 mm with a mean of -1.1 mm. The percent change has ranged from -5 % to -21 % with a mean of -13 %. A possible cause of Vance's (2004) small amount of ridge loss could be due to the small amount of time the flap was open, as opposed to Adam's (2005) study which employed a longer surgical procedure (Table 13). Another factor in varying results is tooth type. According to the University of Louisville studies (Table 14), maxillary teeth compared to mandibular teeth and anterior teeth compared to posterior teeth have a greater percentage ridge width loss. Thus, results of a study could vary based on the distribution of teeth in the sample (Table 14). This study resulted in a mean horizontal ridge change of -1.3 mm (15%) for the Bio-Col technique and -1.1 mm (14%) for the PTFE technique.



**Table 13**  
**Horizontal Ridge Width at the Crest for U of L Studies**  
**Mean ± sd in mm**

	<b>Initial</b>	<b>Final</b>	<b>Change</b>	<b>% Change</b>
<b>Iasella 2003 FDBA</b>	9.2 ± 1.2	8.0 ± 1.4	-1.2 ± 0.9	-13
<b>Vance 2004 Calmatrix</b>	8.9 ± 1.8	8.4 ± 1.5	-0.5 ± 0.7	-6
<b>Vance 2004 BioGide/BioOss</b>	9.7 ± 1.1	9.2 ± 1.1	-0.5 ± 0.8	-5
<b>Adams 2005 Intra FDBA</b>	9.4 ± 1.2	7.4 ± 1.5	-2.0 ± 0.9*	-21
<b>Adams 2005 Overlay FDBA</b>	8.5 ± 1.0	7.1 ± 1.2	-1.4 ± 1.0*	-17
<b>Siu 2007 Flap</b>	8.5 ± 1.5	7.5 ± 1.5	-1.0 ± 1.1	-12
<b>Siu 2007 Flapless</b>	8.3 ± 1.3	7.0 ± 1.9	-1.3 ± 1.0	-16
<b>Witonsky 2009 BioCol</b>	8.6 ± 1.0	7.3 ± 1.0	-1.3 ± 0.9	-15
<b>Witonsky 2009 PTFE</b>	7.9 ± 1.5	6.8 ± 1.4	-1.1 ± 1.1	-14
<b>Mean</b>	8.8 ± 0.6	7.6 ± 0.8	-1.1 ± 0.5	-13 ± 5

\* = p < 0.05 between initial and 4-month values

**Table 14**  
**U of L Studies by Tooth Type**  
**Mean ± sd in mm**

	<b>n</b>	<b>Initial</b>	<b>Final</b>	<b>Change</b>	<b>% Change</b>
<b>Maxillary Incisor</b>	23	7.8 ± 1.0	6.3 ± 1.2	-1.5 ± 0.9	-19 ± 11
<b>Mandibular Incisor</b>	1	6.1	5.1	-1.0	-17
<b>Maxillary Canine</b>	4	8.8 ± 1.0	7.3 ± 1.9	-1.5 ± 1.0	-18 ± 14
<b>Mandibular Canine</b>	2	7.9 ± 2.6	8.1 ± 2.3	0.2 ± 0.2	+4 ± 4
<b>Maxillary Premolar</b>	69	9.3 ± 1.2	8.2 ± 1.4	-1.2 ± 1.0	-12 ± 11
<b>Mandibular Premolar</b>	15	8.0 ± 1.3	7.6 ± 1.4	-0.4 ± 0.9	-5 ± 11


## CHAPTER II

### METHODS

**Study design.** Twenty-eight patients were invited to participate in this randomized, controlled, single, blinded clinical trial. By random selection, using a coin toss technique, fourteen positive controls were selected to receive a mineralized cancellous particulate bone xenograft (Bio-Oss, Geistlich Pharma, Switzerland), which was covered by a collagen plug, (CollaPlug, Zimmer Dental, California) and used a full-thickness flap technique. Fourteen test patients received an intrasocket cortical mineralized particulate allograft (500 to 800  $\mu\text{m}$ ) (RegenerOss, Biomet 3i, Palm Beach, FL) which was covered with a nonporous PTFE membrane, and also used a full thickness flap technique. Each patient received a post-surgical regimen of 50 mg of doxycycline hyclate (Warner Chilcott Inc. Morris Planes, New Jersey) 1 tab qd for 2 weeks; 375 mg of naproxen sodium (Geneva Pharmaceuticals, Inc. Broomfield, CO) 1 tab q12h for 1 week; chlorhexidine 0.12% (Colgate Oral Pharmaceutical Canton, Massachusetts), twice daily, and analgesics as needed.

At 4-months post-surgery, a trephine core was taken from the grafted site immediately prior to implant placement and was submitted for histologic preparation using hematoxylin and eosin staining after all specimens were collected.

**Figure 1**



## **Patient Selection**

### **Inclusion criteria**

- 1) Have one non-molar tooth requiring extraction that will be replaced by a dental implant. The site must be bordered by at least one tooth.
- 2) Must be at least 18 years old.
- 3) Must sign an informed consent approved by the University of Louisville Human Studies Committee.

### **Exclusion Criteria**

- 1) Debilitating systemic diseases, or diseases that affect the periodontium.
- 2) Molar teeth.
- 3) Allergy to any material or medication used in the study.
- 4) Require prophylactic antibiotics.
- 5) Previous head and neck radiation therapy.
- 6) Chemotherapy in the previous 12 months.
- 7) Long term NSAID or steroid therapy.

### **Post-Surgical Exclusion**

Any site that is excluded after surgery will be reported. Sites were excluded if there was:

- 1) Loss of graft or barrier material.
- 2) Unanticipated healing complications that will adversely affect treatment results.

## **Presurgical Management**

Each patient received a diagnostic work-up including standardized periapical radiographs (Appendix D), study casts, clinical photographs, and a clinical examination to record attachment level, probing depth, recession, and mobility of teeth adjacent to the extracted sites. Customized triad occlusal stents were fabricated on the study casts to serve as fixed reference guides for the measurements (Appendix F).

Presurgical preparation included detailed oral hygiene instructions. Baseline data was collected just before the surgical phase of the treatment. Baseline data included:

## **Clinical Measurements**

- Plaque index: Silness and Loe 1964 (Appendix A).
- Gingival index: Loe 1967 – Gingival index (Appendix B).
- Bleeding on Probing Index: Dichotomous index (Appendix C).
- Gingival margin levels: Measured from CEJ to the gingival margin.
- Keratinized tissue: Measured from the gingival margin to the mucogingival junction
- Clinical attachment level: Measured from CEJ to the bottom of the clinical periodontal pocket.
- Clinical tooth mobility: Measured by using the modified Miller's Index.
- Horizontal Ridge width: A digital caliper was used to measure total ridge width to the nearest  $10^{-2}$  mm at the mid point of the alveolar crest and 5 mm apical to the crest, measured post-extraction and prior to implant placement.
- Vertical Change in alveolar crest: Measured post-extraction from the stent to alveolar crest minus re-entry stent to alveolar crest values.

- Radiographic examination: A customized stent was constructed using Triad® light cured resin (Appendix F) and a Rinn-XCP on the patient model (Appendix D) to ensure standardization of the projection.
- Clinical photographs.

### **Surgical treatment**

Patients were anesthetized with 2% lidocaine containing epinephrine in both 1:100,000 and 1:50,000 concentrations. A full-thickness, mucoperiosteal flap utilizing papilla preservation was elevated on the buccal and palatal/lingual with long releasing incisions up to the mucogingival junction to expose the alveolar ridge. Teeth were elevated and atraumatically extracted with periostomes, elevators, and forceps. The extraction socket was then curetted to remove all soft tissue. After flap reflection, the triad stent was used to obtain vertical bone height relative to the stent.

A digital caliper was applied to the ridge to measure the total alveolar ridge width at the mid-socket crest and 5 mm apical to the crest. In the PTFE group, the extraction socket was grafted with an intrasocket mineralized cortical particulate allograft composed of cortical chips 500 to 800  $\mu\text{m}$  (RegenerOss, Biomet 3i, Palm Beach, FL) then covered with a nonporous PTFE membrane. The PTFE membrane was shaped to extend 3 to 4 mm beyond the socket margins and 1.0 mm from the adjacent root. The Plug group received a mineralized cancellous particulate (0.25 to 1.00 mm) xenograft (Bio-Oss, Geistlich Pharma, Switzerland), covered by a collagen plug (CollaPlug, Zimmer Dental, California). The flaps were replaced and sutured with Cytoplast PTFE sutures (Osteogenics Biomedical Lubbock, Tx). Patients were given naproxen 375 mg (Geneva

Pharmaceuticals, Inc. Broomfield, CO), one tab q12h, doxycycline hyclate 50 mg (Warner Chilcott Inc. Morris Plains, New Jersey), 1 tab qd, chlorhexidine 0.12% (Colgate Oral Pharmaceutical Canton, Massachusetts), twice daily, and analgesics as needed.

Patients were seen for postoperative appointments at 2, 4, 8, and 12 weeks. Photographs were taken at each interval. In addition, at 4 weeks, patients in the PTFE group were seen for membrane removal. The membranes were removed atraumatically without flap reflection, when possible.

At 4 months, another standardized radiograph was taken. All baseline measurements were repeated. Patients were anesthetized with 2% lidocaine containing both 1:100,000 and 1:50,000 concentrations of epinephrine, and full-thickness mucoperiosteal flaps were elevated on the buccal and palatal/lingual. Papilla were again preserved and not included in the flap design. The acrylic stent was placed and measurements were obtained of vertical ridge height relative to the stent. The digital caliper was used to measure alveolar ridge width at the mid-buccal crestal sites and 5 mm apical to the crest. A blinded examiner performed all clinical measurements for both the initial and final data collection points.

**Histologic analysis.** A 2.7 X 6 mm trephine (H & H Company Ontario, California) was used with copious chilled irrigation to remove a trephine core from the experimental or control site. The osseous core was removed from the trephine using a periodontal probe that was placed into a window and elevated. The core was subsequently placed directly into a bottle of 10% buffered formalin for histologic preservation. The cores were decalcified and 12 to 15 step serial sections were taken from each

longitudinally sectioned core. The sections were stained with hematoxylin and eosin. This resulted in 10 slides per patient with at least 4 sections per slide. Vital and non-vital bone and trabecular space quantitation was performed using an American Optical microscope at 150X with a 10 X 10 ocular grid. For each patient 6 of 10 slides were assessed and for each slide at least 100 squares on the ocular grid were counted. A mean percentage of vital and non-vital bone and trabecular space was calculated for each patient.

An osteotomy site was then prepared with a surgical handpiece, using copious irrigation, and each patient received an endosseous root form dental implant. Flaps were replaced, and sutured with 4-0 silk sutures. Patients were again given naproxen 375 mg, doxycycline hyclate 50 mg and analgesics as needed.

**Statistical analysis.** For the statistical analysis, a t-test was used to evaluate the statistical significance of both the within and between group differences for both clinical and histologic data.



## CHAPTER III

### RESULTS

A total of 22 females and 6 males with a mean age of 54.9, ranging from 32 to 83, were enrolled. The Plug group consisted of 3 maxillary central incisor, 9 maxillary premolars, and 2 mandibular premolars. The PTFE group consisted of 1 maxillary lateral incisor, 9 maxillary premolars, and 4 mandibular premolars. There was 1 smoker in the Plug group and 2 smokers in the PTFE group. Smokers were excluded if they smoked more than 1/2 pack per day. Data from this study was derived from 14 patients, 7 per group, completed by Dr. Elliot Bermudez and the remaining 14 were completed by Dr. Jason Witonsky.

**Clinical Indices.** Plaque index, gingival index and bleeding on probing had low initial values and did not change significantly ( $p > 0.05$ , Table 14). There were no statistically significant differences between groups for any of the 3 clinical indices ( $p > 0.05$ ).

**Horizontal Ridge Width Changes.** The Plug group presented with a mean initial width at the crest of  $8.6 \pm 1.0$  mm, which changed to  $7.3 \pm 1.0$  mm at month 4 for a mean loss of  $1.3 \pm 0.9$  mm ( $p < 0.05$ , Table 15). The PTFE group had a mean initial width at the crest of  $7.9 \pm 1.5$  mm, which decreased to  $6.8 \pm 1.4$  mm for a mean loss of  $-1.1 \pm 1.1$  mm ( $p < 0.05$ ). There were no statistically significant differences between groups ( $p > 0.05$ ).

For the Plug group the mean initial width 5 mm apical to the crest was  $9.5 \pm 1.1$  mm, which decreased to  $8.0 \pm 1.2$  mm at month 4 for a mean loss of  $-1.5 \pm 1.0$  mm ( $p < 0.05$ ). The PTFE group had a mean initial width 5 mm apical to the crest of  $8.9 \pm 1.8$  mm, changed to  $7.4 \pm 1.4$  mm for mean loss of  $-1.5 \pm 1.3$  mm ( $p < 0.05$ ). There were no statistically significant differences between groups ( $p > 0.05$ ).

**Vertical mid-Buccal Ridge Height Changes.** Mid-buccal ridge height for the Plug group had a mean loss of  $-0.1 \pm 1.6$  mm ( $p > 0.05$ , Table 16). For the PTFE group, mid-buccal height had a mean gain of  $0.4 \pm 2.1$  mm ( $p > 0.05$ ). There were no statistically significance differences between groups ( $p > 0.05$ ).

**Vertical mid-Lingual Ridge Height Changes.** Mid-lingual ridge height for the Plug group had a mean loss of  $-0.3 \pm 1.3$  mm ( $p < 0.05$ , Table 16). For the PTFE group, mid-lingual height had a mean loss of  $-1.1 \pm 2.0$  mm ( $p < 0.05$ ). There were no statistically significance differences between groups ( $p > 0.05$ ).

**Vertical Mesial Ridge Height Changes.** Vertical mesial ridge height for the Plug group had a mean loss of  $-0.5 \pm 0.7$  mm ( $p < 0.05$ , table 16). For the PTFE group, mid-lingual height had a mean loss of  $-0.8 \pm 0.9$  mm ( $p < 0.05$ ). There were no statistically significance differences between groups ( $p > 0.05$ ).

**Vertical Distal Ridge Height Changes.** Vertical distal ridge height for the Plug group had a mean loss of  $-0.6 \pm 0.6$  mm ( $p > 0.05$ , table 16). For the PTFE group, mid-lingual height had a mean loss of  $-0.7 \pm 1.1$  mm ( $p > 0.05$ ). There were no statistically significance differences between groups ( $p > 0.05$ ).

**CEJ to Osseous Crest Changes.** The Plug group presented with a mean initial mesial CEJ to osseous crest distance of  $2.8 \pm 0.5$  mm, which increased to  $3.3 \pm 1.4$  mm at

month 4 for a mean loss of  $-0.5 \pm 1.0$  mm ( $p < 0.05$ , table 17). The PTFE group had a mean initial mesial CEJ to osseous crest distance of  $3.0 \pm 0.6$  mm, which increased to  $3.2 \pm 0.6$  mm for a mean loss of  $-0.2 \pm 0.9$  mm ( $p > 0.05$ ). There were no statistically significant differences between groups ( $p > 0.05$ ).

For the Plug group there was a mean initial distal CEJ to osseous crest distance of  $3.0 \pm 0.7$  mm, which increased to  $3.4 \pm 1.0$  mm at month 4 for a mean loss of  $-0.4 \pm 0.9$  mm ( $p > 0.05$ ). The PTFE group had a mean initial distal CEJ to osseous crest distance of  $2.9 \pm 0.9$  mm, which increased to  $3.6 \pm 1.2$  mm for a mean loss of  $-0.7 \pm 1.5$  mm ( $p < 0.05$ ). There were no statistically significant differences between groups ( $p > 0.5$ ).

**Histologic evaluation.** A high percentage of vital bone was found in both groups (Table 18). Histologic analysis revealed that Plug sites healed with  $28 \pm 20\%$  vital bone,  $37 \pm 16\%$  non-vital bone,  $35 \pm 13\%$  trabecular space. The PTFE sites healed with  $35 \pm 21\%$  vital bone,  $31 \pm 22\%$  non-vital bone,  $34 \pm 10\%$  trabecular space. There were no statistically significant differences between the two groups ( $p > 0.05$ ).

**Table 15**

**Clinical Indices for Plug and PTFE Sites**

**Mean  $\pm$  sd in index units**

		<b>Initial</b>	<b>Final</b>	<b>Change</b>
<b>Plaque</b>	Plug	0.1 $\pm$ 0.3	0.2 $\pm$ 0.2	0.1 $\pm$ 0.4
<b>Index</b>	PTFE	0.3 $\pm$ 0.4	0.3 $\pm$ 0.5	0.0 $\pm$ 0.7
<b>Gingival</b>	Plug	0.3 $\pm$ 0.4	0.3 $\pm$ 0.4	0.0 $\pm$ 0.4
<b>Index</b>	PTFE	0.4 $\pm$ 0.4	0.3 $\pm$ 0.3	-0.1 $\pm$ 0.5
<b>Bleeding on Probing</b>	Plug	0.2 $\pm$ 0.3	0.1 $\pm$ 0.1	-0.1 $\pm$ 0.3
	PTFE	0.1 $\pm$ 0.1	0.8 $\pm$ 2.5	0.7 $\pm$ 2.4

**Table 16**  
**Horizontal Ridge Width for Plug and PTFE Sites**  
**Mean ± sd in mm**

	<b>Initial</b>	<b>Final</b>	<b>Change</b>	<b>Range</b>
<b>Plug at Crest</b>	8.6 ± 1.0	7.3 ± 1.0	-1.3 ± 0.9*	-2.8 to +0.3
<b>PTFE at Crest</b>	7.9 ± 1.5	6.8 ± 1.4	-1.1 ± 1.1*	-2.5 to +1.2
<b>Plug at 5 mm</b>	9.5 ± 1.1	8.0 ± 1.2	-1.5 ± 1.0*	-3.0 to +0.7
<b>PTFE at 5 mm</b>	8.9 ± 1.8	7.4 ± 1.4	-1.5 ± 1.3*	-3.9 to +0.7

\* = p < 0.05 between initial and 4-month values

**Table 17**  
**Vertical Ridge Height Change for Plug and PTFE Sites**  
**Mean ± sd in mm**

<b>Location</b>	<b>Plug</b>	<b>PTFE</b>	<b>Plug</b>	<b>PTFE</b>
	Mean Change ± sd in mm		Range in mm	
<b>Mid-Buccal</b>	-0.1 ± 1.6	0.4 ± 2.1	-2.5 to 4.0	-2.0 to 4.0
<b>Mid-Lingual</b>	-0.3 ± 1.3	-1.1 ± 2.0*	-1.5 to 3.0	-5.0 to 2.5
<b>Mesial</b>	-0.5 ± 0.7*	-0.8 ± 0.9*	-2.0 to 0.3	-2.0 to 1.2
<b>Distal</b>	-0.6 ± 0.6*	-0.7 ± 1.1*	-1.8 to 0.3	-1.8 to 2.2

\* = p < 0.05 between initial and 4-month values

**Table 18**  
**CEJ to Osseous Crest Change at Adjacent Teeth**  
**Mean ± sd in mm**

	<b>n</b>	<b>Initial</b>	<b>Final</b>	<b>Change</b>
<b><u>Plug</u></b>				
<b>Mesial</b>	14	2.8 ± 0.5	3.3 ± 0.8	-0.5 ± 1.0*
<b>Distal</b>	13	3.0 ± 0.7	3.4 ± 1.0	-0.4 ± 0.9
<b><u>PTFE</u></b>				
<b>Mesial</b>	13	3.0 ± 0.6	3.2 ± 0.6	-0.2 ± 0.9
<b>Distal</b>	10	2.9 ± 0.9	3.6 ± 1.2	-0.7 ± 1.5*

\* = p < 0.05 between initial and 4-month values

**Table 19**  
**Histologic Data for PTFE and Collagen Plug Sites**

**Mean  $\pm$  sd**

<b>Group</b>	<b>Time</b>	<b>n</b>	<b>% Vital</b>	<b>% Non-vital</b>	<b>% Trabecular</b>
<b>Plug</b>	4 month	14	28 $\pm$ 20	37 $\pm$ 16	35 $\pm$ 13
<b>PTFE</b>	4 month	14	35 $\pm$ 21	31 $\pm$ 22	34 $\pm$ 10



## CHAPTER IV

### DISCUSSION

In this 4-month randomized, controlled, blinded clinical study of intrasocket ridge preservation in humans, the BioCol technique which utilizes a cancellous particulate xenograft (BioOss) plus a collagen plug (Plug group) was compared to a technique utilizing a mineralized particulate cortical allograft plus a nonporous polytetrafluoroethylene (PTFE) membrane (PTFE group).

In this study there were no statistically significant differences in the change in mean horizontal ridge width between groups. At the crest and 5 mm apical to the crest both groups showed significant loss ( $p < 0.05$ ), each losing -1.5 mm of mean ridge width.

Ridge preservation studies show less loss of mean ridge width when compared to treatment by extraction alone (Lekovic et al. 1997, Lekovic et al. 1998, Iasella et al. 2003). Extraction alone most often leads to extensive ridge resorption. In general, the longer the time period studied, the greater the ridge resorption reported (Lekovic et al. 1997, Lekovic et al. 1998, Schropp et al. 2003, Iasella et al. 2003). The ridge width dimension is compromised to a greater degree than ridge height, which is generally minimally affected. Ridge preservation does not totally eliminate loss of ridge width and most studies show that some loss still occurs. Previous studies have shown that

extraction alone leads to a mean loss of 43% ridge width versus 12% loss for ridge preservation (Tables 6, 7, 8, 9).

This study showed more loss of ridge dimension with a preservation procedure than 1 of the earlier studies at this institution, Vance et al. (2004), but less than 3 previous studies; Adams, Iasella et al. (2003), and Siu, (Table 16).

Both groups lost mean ridge height at all locations (mid-buccal, mid-lingual, mesial and distal) except the mid-buccal site for the PTFE group, which gained 0.4 mm. The PTFE group showed a statistically significant loss of -1.1 mm at the mid-lingual site and -0.8 on the mesial ( $p < 0.05$ ). The plug group showed a statistically significant loss of -0.5 mm on the mesial and -0.6 mm on the distal ( $p < 0.05$ ). None of these changes were statistically significant between groups ( $p > 0.05$ ).

This study evaluated loss of crestal width in extraction sites with at least one adjacent tooth. Twelve of 14 sites had 2 adjacent teeth. Loss of crestal width may be greater when there are no adjacent teeth, especially when all teeth in an arch are being removed. Thus, the means and ranges reported in this study may not be generalizable and should be limited in application to sites with adjacent teeth. Further study is warranted to document the resorptive response when an arch is edentulated.

The mean CEJ to osseous crest distance changed 1 mm or less for both the Plug and PTFE groups with no statistically significant differences between groups ( $p < 0.05$ ).

The PTFE group had more vital bone (35 vs 28%) and less non-vital bone (31 vs. 37%) than the Plug group. The xenograft used for the Plug group typically resorbs slowly and this results was not unexpected. Use of the allograft resulted in the presence of more vital bone in the area of implant placement. The significance of increased vital

bone for long term implant success or survival has not been established, however, most clinicians prefer to have increased vital bone.

Based on the results of this study, the change in ridge dimensions did not show any statistically significant differences between the Plug or PTFE ridge preservation techniques. Both treatments were effective in the preservation of horizontal and vertical ridge dimensions.

## **CHAPTER V**


### **CONCLUSIONS**

Within the limits of this study design and sample size it may be concluded that:

- 1) Mean crestal ridge width was preserved for both the Plug and PTFE groups and there were no statistically significant differences between groups ( $p > 0.05$ ).
- 2) There was a trend toward greater loss of mean mid-buccal ridge height for the Plug group, although there were no statistically significant differences between groups ( $p > 0.05$ ).
- 3) The mean CEJ to osseous crest distance showed only a minimal loss of 1 mm or less for both the Plug and PTFE groups and there were no statistically significant differences between groups ( $p > 0.05$ ).
- 4) The PTFE group had more vital bone and less non-vital bone than the Plug group, however, there were no statistically significant differences between groups ( $p > 0.05$ ).



Figure 2. a) Pre-op PTFE;



b) Cortical allograft;





Figure 2. c) PTFE membrane;



d) Pre-op 4-month re-entry.







Figure 3. a) Pre-op collagen plug;



b) Cancellous xenograft;





Figure 3. c) Collagen plug;



d) Pre-op 4-month re-entry..

## REFERENCES

- Adams DA. Ridge preservation comparing an intrasocket FDBA allograft alone to a combined intra-and extra-socket FDBA allograft both covered with acellular dermal matrix membrane. [Thesis]. 2005. Louisville, Kentucky. University of Louisville.
- Amler MH, Johnson PL, Salaman I. Histologic and histochemical investigation of human alveolar socket healing in undisturbed extraction wounds. *J Am Dent Assoc* 1960;6:132-144.
- Araujo MG, Lindhe J. Dimensional ridge alterations following tooth extraction. An experimental study in the dog. *J Clin Periodontol* 2005a;32(2):212-218.
- Araujo MG, Sukekava F, Wennstrom JL, Lindhe J. Ridge alterations following implant placement in fresh extraction sockets: an experimental study in the dog. *J Clin Periodontol* 2005b;32(6):645-652.
- Araujo M, Linder E, Wennstrom J, Lindhe J. The influence of bio-oss collagen on healing of an extraction socket: an experimental study in the dog *Int J Periodontics Restorative Dent* 2008;28(2):123-35.
- Artzi Z, Nemcovsky CE. The application of deproteinized bovine bone mineral for ridge preservation prior to implantation. Clinical and histological observations in a case report. *J Periodontol* 1998;69(9):1062-1067.
- Artzi Z, Tal H, Dayan D. Porous bovine bone mineral in healing of human extraction sockets: Part 1: histomorphometric evaluations at 9 months. *J Periodontol* 2000;71(6):1015-1023.
- Artzi Z, Tal H, Dayan D. Porous bovine bone mineral in healing of human extraction sockets: 2. Histochemical observations at 9 months. *J Periodontol* 2001;72(2):152-159.
- Ash MM, and Wheeler RC. *Wheeler's Atlas of Tooth Form*. Philadelphia: Saunders, 1984.

Ashman A. Postextraction ridge preservation using a synthetic alloplast. *Implant Dent* 2000;9(2):168-176.

Barone A, Aldini NN, Fini M, Giardino R, Guirado JL, Covani U. Xenograft versus extraction alone for ridge preservation after tooth removal: A clinical and histomorphometric study *J Periodontol* 2008;79:1370-1377.

Bartee BK. Evaluation of a new polytetrafluoroethylene guided tissue regeneration membrane in healing extraction sites *Compend Contin Educ Dent* 1998;19(12):1256-1264.

Bartee BK. Extraction site reconstruction for alveolar ridge preservation. Part 2: Membrane-assisted surgical technique. *J Oral Implantol* 2001;27(4):194-197.

Becker W, Becker BE, Caffesse R. A comparison of demineralized freeze-dried bone and autologous bone to induce bone formation in human extraction sockets. *J Periodontol* 1994;65(12):1128-1133.

Becker W, Urist M, Becker BE, Jackson W, Parry DA, Bartold M, Vincenzzi G, De Georges D, Niederwanger M. Clinical and histologic observations of sites implanted with intraoral autologous bone grafts or allografts. 15 human case reports. *J Periodontol* 1996;67(10):1025-1033.

Becker W, Clokie C, Sennerby L, Urist MR, Becker BE. Histologic findings after implantation and evaluation of different grafting materials and titanium micro screws into extraction sockets. Case Reports. *J Periodontol* 1998;69(4):414-421.

Berglundh T, Lindhe J, Jonsson K, Ericsson I. The topography of the vascular systems in the periodontal and peri-implant tissues in the dog. *J Clin Periodontol* 1994;21(3):189-193.

Botticelli D, Berglundh T, Lindhe J. Resolution of bone defects of varying dimension and configuration in the marginal portion of the peri-implant bone. An experimental study in the dog. *J Clin Periodontol* 2004;31(4):309-317.

Botticelli D, Berglundh T, Persson LG, Lindhe J. Bone regeneration at implants with turned or rough surfaces in self-contained defects. An experimental study in the dog. *J Clin Periodontol* 2005;32(5):448-455.

Boyne PJ. Osseous repair of the postextraction alveolus in man. *Oral Surg Oral Med Oral Pathol* 1966;21(6):805-813.



Brkovic BM, Prasad HS, Konandreas G, Milan R, Antunovic D, Sandor GK, Rohrer MD. Simple preservation of a maxillary extraction socket using beta-tricalcium phosphate with type I: Preliminary clinical and histomorphometric observations J Can Dent Assoc 2008;74(6):523-528.

Brugnami F, Then PR, Moroi H, Leone CW. Histologic evaluation of human extraction sockets treated with demineralized freeze-dried bone allograft (DFDBA) and cell occlusive membrane. J Periodontol 1996;67(7):821-825.

Brugnami F, Then PR, Moroi H, Kabani S, Leone CW. GBR in human extraction sockets and ridge defects prior to implant placement: Clinical results and histologic evidence of osteoblastic and osteoclastic activities in DFDBA. Int J Periodontics Restorative Dent 1999;19(3):259-268.

Camargo PM, Lekovic V, Weinlaender M, Klokkevold PR, Kenney EB, Dimitrijevic B, Nedic M, Jancovic S, Orsini M. Influence of bioactive glass on changes in alveolar process dimensions after exodontia. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2000;90(5):581-586.

Cardaropoli G, Araujo M, Lindhe J. Dynamics of bone tissue formation in tooth extraction sites. An experimental study in dogs. J Clin Periodontol 2003;30(9):809-818.

Cardaropoli G, Araujo MG, Hayacibara R, Sukekava F, Lindhe J. Healing of extraction sockets and surgically produced - augmented and non-augmented - defects in the alveolar ridge. An experimental study in the dog. J Clin Periodontol 2005;32(5):435-440.

Cardaropoli D, Cardaropoli G. Preservation of the postextraction alveolar ridge: A clinical and histologic study J Periodont Res 2008;28:469-477.

Carmagnola D, Berglundh T, Araujo M, Albrektsson T, Lindhe J. Bone healing around implants placed in a jaw defect augmented with Bio-Oss. An experimental study in dogs. J Clin Periodontol 2000;27(11):799-805.

Clafin R. Healing of disturbed and undisturbed extraction sites. J Am Dent Assoc 1936;23(6):945-959.

Evian C, Rosenberg E, Coslet J, Corn H. The osteogenic activity of bone removed from healing extraction sockets in humans. J Periodontol 1982;53(2):81-85.

Fowler EB, Whicker R. Modified approach to the bio-col ridge preservation technique: A case report J Contemp Dent Pract 2004;5(3):1-9.

Froum S, Cho S, Rosenberg E, Rohrer M, Tarnow D. Histologic Comparison of Healing Extraction Sockets Implanted With Bioactive Glass or Demineralized Freeze-Dried Bone Allograft: A Pilot Study. *J Periodontol* 2002;73(1):94-102.

Froum S, Cho SC, Elian N, Rosenberg E, Rohrer M, Tarnow D. Extraction sockets and Implantation of Hydroxyapatites with membrane barriers a histologic study. *Implant Dent* 2004;13(2):153-161.

Garg AK. Alveolar ridge preservation during and after surgical tooth removal. *Dent Implantol Update* 2001;11(8):57-62.

Guarnieri R, Pecora G, Fini M, Nicoli Aldini N, Giardino R, Orsini G, Piattelli A. Medical grade calcium sulfate hemihydrate in healing of human extraction sockets: Clinical and histological observations at 3 months. *J Periodontol* 2004;75(5):902-908.

Hoffmann O, Bartee BK, Beaumont C, Kasaj A, Deli G, Zafiroopoulos GG. Alveolar bone preservation in extraction sockets using non-resorbable dPTFE membrnes: A retrospective non-randomized study *J Periodontol* 2008;79:1355-1369.

Horowitz RA. Extraction environment enhancement: Critical evaluation of early socket healing in long-term barrier protected extraction sockets *Compend Contin Educ Dent* 2005;26(10):703-713.

Iasella JM, Greenwell H, Miller RL, Hill M, Drisko C, Bohra AA, Scheetz JP. Ridge preservation with freeze dried bone allograft and a collagen membrane compared to extraction alone for implant site development: A clinical and histologic study in humans. *J Periodontol* 2003;74(7):988-997.

Lekholm U, Zarb GA. Patient selection and preparation. In Branemark P-I, Zarb GA, Albrektsson T, editors: *Tissue integrated prostheses: osseointegration in clinical dentistry*. Quintessence Int 1985;199-209.

Lekovic V, Kenney EB, Weinlaender M, Han T, Klokkevold P, Nedic M, Orsini M. A bone regenerative approach to alveolar ridge maintenance following tooth extraction. Report of 10 cases. *J Periodontol* 1997;68(6):563-570.

Lekovic V, Camargo PM, Klokkevold PR, Weinlaender M, Kenney EB, Dimitrijevic B, Nedic M. Preservation of alveolar bone in extraction sockets using bioabsorbable membranes. *J Periodontol*. 1998;69(9):1044-1049.

Loe H. The gingival index, the plaque index and the retention index systems. *J Periodontol* 1967;38(Suppl):610-617.

Lucyszyn SM, Papalexiou V, Novaes AB Jr, Grisi MF, Souza SL, Taba M Jr. Acellular dermal matrix and hydroxyapatite in prevention of ridge deformities after tooth extraction. *Implant Dent* 2005;14(2):176-184.

MacNeill SR, Cobb CM, Rapley JW, Glaros AG, Spencer P. In vivo comparison of synthetic osseous graft materials. A preliminary study. *J Clin Periodontol* 1999;26(4):239-245.

Mangano C, Piatelli, A, Perrotti V, Iezzi G. Dense hydroxyapatite inserted into postextraction sockets: A histologic and histomorphometric 20-year case report *J Periodontol* 2008;79:929-933.

Mellonig J, Bowers G, Cotton W. Comparison of bone graft materials. Part II. New bone formation with autografts and allografts: A histological evaluation. *J Periodontol* 1981;52(6):297-302.

Mellonig J. Freeze-Dried bone allografts in periodontal reconstructive surgery. *Dent Clin North Am* 1991;35(3):505-520.

Neiva RF, Tsao YP, Eber R, Shotwell J, Billy E, Wang HL. Effects of a putty-form hydroxyapatite matrix combined with the synthetic cell-binding peptide p-15 on alveolar ridge preservation *J Periodontol* 2008;79:291-299.

Nemcovsky CE Serfaty V. Alveolar ridge preservation following extraction of maxillary anterior teeth. Report on 23 consecutive cases. *J Periodontol* 1996;67(4):390-395.

Pietrokovski J, Massler M. Alveolar ridge resorption following tooth extraction. *J Prost Dent* 1967;17(1):21-27.

Schropp L, Wenzel A, Kostopoulos L, Karring T. Bone healing and soft tissue contour changes following single-tooth extraction: A clinical and radiographic 12 month prospective study. *Int J Periodontics Restorative Dent* 2003;23(4):313-323.

Schwartz Z, Mellonig JT, Carnes DL Jr, De La Fontaine J, Cochran DL, Dean DD, Boyan BD. Ability of commercial demineralized freeze-dried bone allograft to induce new bone formation. *J Periodontol* 1996;67(9):918-926.

Schwartz Z, Somers A, Mellonig JT, Carnes DL, Dean DD, Cochran DL, Boyan BD. Ability of commercial demineralized freeze-dried bone allograft to induce new bone formation is dependent on donor age but not gender. *J Periodontol* 1998;69(4):470-478.

Schwartz Z, Weesner T, van Dijk S, Cochran DL, Mellonig JT, Lohmann CH, Carnes DL, Goldstein M, Dean DD, Boyan BD. Ability of deproteinized cancellous bovine bone to induce new bone formation. *J Periodontol* 2000;71(8):1258-1269.

Sclar A. Cosmetic soft-tissue enhancement for dental implants. *Alpha Omegan* 2000;93(3):38-46.

Sclar AG. Strategies for management of single tooth extraction sites in aesthetic implant therapy *J Oral Maxillofac Surg* 2004;62s:90-105.

Serino G, Biancu S, Iezzi G, Piatelli A. Ridge preservation following tooth extraction using a polylactide and polyglycolic sponge as space filler: a clinical and histological study in humans. *Clin Oral Implants Res* 2003;14(5):651-658.

Silness J, Loe H. Periodontal disease in pregnancy. II. Correlation between oral hygiene and periodontal conditions. *Acta Odontol Scand* 1964;22(1):121-135.

Simon BI, Von Hagen S, Deasy MJ, Faldu M, Resnansky D. Changes in alveolar bone height and width following ridge augmentation using bone graft and membranes. *J Periodontol* 2000;71(1):1774-1791.

Siu TL. Ridge preservation comparing a flap vs. flapless technique using a demineralized bone matrix allograft plus mineralized particulate allograft and covered with a calcium sulfate barrier. [Thesis]. 2007. Louisville, Kentucky. University of Louisville.

Smukler H, Landi L, Setayesh R. Histomorphometric evaluation of extraction sockets and deficient alveolar ridges treated with allograft and barrier membrane: A pilot study. *Int J Oral Maxillofac Implants* 1999;14(3):407-416.

Urist MR, Strates BS. Bone morphogenetic protein. *J Dent Res* 1971;50(9):1392-1406.

Vance GS, Greenwell H, Miller RL, Hill M, Johnston H, Scheetz JP. Comparison of an allograft in an experimental putty carrier and a bovine-derived xenograft used in ridge preservation: A clinical and histologic study in humans. *Int J Oral Maxillofac Implants* 2004;19(4):491-497.

Wang HL, Sao JP. Histologic evaluation of socket augmentation with mineralized human allograft *Int J Periodontics Restorative Dent* 2008;28(3):231-237.

Woelfel, JB. *Dental Anatomy: Its Relevance to Dentistry*. New York: Lippincott, 1990.

Yilmaz S, Efeoglu E, Kilic AR. Alveolar ridge reconstruction and/or preservation using root form bioglass cones *J Clin Periodontol* 1998;25(10):832-839.

Zitzmann NU, Naef R, Scharer P. Resorbable versus nonresorbable membranes in combination with Bio-Oss for guided bone regeneration. *Int J Oral Maxillofac Implants* 1997;12(3):844-852.

Zitzmann NU, Scharer P, Marinello CP, Schupbach P, Berglundh T. Alveolar ridge augmentation with Bio-Oss: A histologic study in humans. *Int J Periodontics Restorative Dent* 2001;21(3):289-296.

Zubillaga G, Von Hagen S, Simon BI, Deasy MJ. Changes in alveolar bone height and width following post-extraction ridge augmentation using a fixed bioabsorbable membrane and demineralized freeze-dried bone osteoinductive graft. *J Periodontol* 2003;74(7):965-975.

## Appendix A

### The Plaque Index

The plaque index of Silness and Loe (1964) was measured. Scores were as follows:

0 - No plaque

1 - A film of plaque adhering to the free gingival margin and adjacent area of the tooth.

The plaque may be seen in situ only after application of disclosing solution or by using the probe on the tooth surface.

2 - Moderate accumulation of soft deposits within the gingival pocket, or on the tooth and gingival margin, which can be seen with the naked eye.

3 - Abundance of soft matter within the gingival pocket and/or on the tooth and gingival margin.

Each gingival unit (buccal, lingual, mesiobuccal, distobuccal, mesiolingual, and distolingual) of the individual tooth was given a score from 0-3, called the plaque index for the area. The scores from the 6 areas of the tooth were added and divided by 6 to give the plaque index for the tooth.

## Appendix B

### Gingival Index

The gingival index of Loe (1967) was measured for the extracted tooth and any adjacent teeth. Scores were be recorded as follows:

0 = Normal gingiva.

1 = Mild inflammation - slight change in color slight edema, no bleeding on probing.

2 = Moderate inflammation - redness, edema, and glazing, bleeding on probing.

3 = Severe inflammation - marked redness and edema, ulceration and tendency to spontaneous bleeding.

Each gingival unit (mesiobuccal, buccal, distobuccal, distolingual, lingual, mesiolingual) of the tooth was given a score 0-3. The scores for each unit were added together and divided by 6 to give the gingival index for that tooth. The score of the test tooth and the two adjacent teeth were added and divided by 3 to give the gingival index for the test of control sites.

## Appendix C

### **Bleeding on Probing Index**

Dichotomous scoring was used for bleeding on probing:

0 = No bleeding;

1 = Bleeding on probing to the bottom of the pocket.



## Appendix D

### Standardized Radiographic technique

An occlusal stent was used to provide a stable foundation for the radiograph holder. A light cured resin material was placed on a Rinn radiograph holder and positioned to allow as near as possible paralleling technique. This material was light cured so that standardized radiographs can be compared. Radiographs were taken at baseline and 4 months.

## Appendix E

### Arithmetic determinations:

**Ridge width (Post-extraction)** = A digital caliper was used to measure total ridge width to the nearest  $10^{-2}$  mm at one point, mid socket, at the alveolar crest and 5 mm from the alveolar crest.

**Ridge width (4 month re-entry)** = Again, a digital caliper measured total ridge width to the nearest  $10^{-2}$  mm at one point, mid socket, at the alveolar crest and 5 mm from the alveolar crest.

**Change in alveolar crest height** = Initial: stent to alveolar crest minus re-entry stent to alveolar crest.

## Appendix F

### Stent fabrication

Rigid stents were made of 3 mm thick light cured resin material in order to provide reproducible measurements. The tooth to be extracted was ground off the model and the light cured resin material was pressed over a cast. Three channels were prepared on the labial and three on the palato/lingual aspect of the stent in which a North Carolina periodontal probe was placed so that mesial, mid and distal measurements could be made on the labial and palato/lingual aspects of the crestal bone. Additionally, two channels were also prepared on the occlusal portion of the stent to provide measures of mesial and distal occlusal ridge height. Holes were prepared with a high-speed hand-piece. In this way, reproducible probing spots and directions of probe insertions were possible.

## **CURRICULUM VITAE**

Jason Witonsky, DMD, MS

### **EDUCATION**

2006-2009 University of Louisville Graduate Periodontics  
Certificate in Periodontics

2006-2009 University of Louisville  
Master of Science in Oral Biology

2002-2006 University of Pennsylvania School of Dental Medicine  
Doctor Medicine Dentistry (D.M.D)

1997-2001 Emory University  
Bachelor of Science (B.S.)

### **LICENSURE**

June 2006-Present Commonwealth of Kentucky Dental License

Category II Laser certification 2007

DEA licensure granted 2006

ACLS certification 2007

**PROFESSIONAL ASSOCIATIONS**

American Dental Association member

American Academy of Periodontology member

Southern Academy of Periodontology member

Academy of Osseointegration member

**EXTERNSHIPS**

Nova Southeastern University, Graduate Periodontics, August 2005

Oral Medicine Rotation, University of Pennsylvania Medical Hospital, February 2006

Internal Medicine Rotation, University of Louisville Hospital, May 2007

Anesthesiology Rotation, University of Louisville Hospital, June 2006