

12-2013

Assessing Trichloromethane Formation and Control in Algal-Stimulated Waters Amended With Nitrogen and Phosphorus

Clinton Mash

University of Arkansas, Fayetteville

Follow this and additional works at: <http://scholarworks.uark.edu/etd>

 Part of the [Environmental Engineering Commons](#), and the [Fresh Water Studies Commons](#)

Recommended Citation

Mash, Clinton, "Assessing Trichloromethane Formation and Control in Algal-Stimulated Waters Amended With Nitrogen and Phosphorus" (2013). *Theses and Dissertations*. 985.

<http://scholarworks.uark.edu/etd/985>

This Thesis is brought to you for free and open access by ScholarWorks@UARK. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of ScholarWorks@UARK. For more information, please contact scholar@uark.edu, ccmiddle@uark.edu.

Assessing Trichloromethane Formation and Control
in Algal-Stimulated Waters Amended With Nitrogen and Phosphorus

Assessing Trichloromethane Formation and Control
in Algal-Stimulated Waters Amended With Nitrogen and Phosphorus

A thesis submitted in partial fulfillment
of the requirements for the degree of
Master of Science in Environmental Engineering

by

Clinton Mash
Arkansas State University – Beebe
Associate of Arts in Liberal Arts, 2008
University of Arkansas
Bachelor of Science in Physics, 2011

December 2013
University of Arkansas

This thesis is approved for recommendation to the Graduate Council.

Dr. Julian Fairey
Thesis Director

Dr. Wen Zhang
Committee Member

Dr. William Oliver III
Committee Member

Abstract

Nitrogen (N) and phosphorus (P) enrichments can stimulate algal growth in drinking water sources, which can cause increased production of disinfection byproduct (DBP) precursors. However, the effect of systematic N and P enrichments on DBP formation and control has not been adequately studied. In this work, we enriched samples from a drinking water source – sampled on April 5, May 30, and August 19, 2013 – with N and P to stimulate algal growth at N:P ratios covering almost five orders of magnitude (0.2-4,429). To simulate DBP-precursor removal processes at drinking water treatment plants (DWTPs), the samples were treated with ClO₂ followed by alum coagulation prior to free chlorine addition to assess the DBP formation potential (FP). Trichloromethane (TCM) was the predominant DBP formed and the TCMFP was the highest at intermediate N:P ratios (~10-50), which corresponded with the peak in algal biomass, as measured by chlorophyll-*a* (Chl-*a*). Algal biomass was P-limited throughout the study period, and co-limited by N for the August 19 sampling set. The differences in TCMFP between the raw and treated waters decreased with increasing P amendment, indicating that ClO₂ and alum coagulation became less effective for TCM precursor removal as algal biomass increased. This study highlights the impact of nutrient enrichments on TCM formation and control and has implications for nutrient management strategies related to source water protection and for DWTPs that use source waters increasingly enriched with N and P.

Acknowledgements

I would like to thank the following people for many reasons... a brief few are listed. Please note that order denotes no specific meaning.

- Dr. Julian Fairey and Dr. Wen Zhang: my advisors – in every meaning of the word.
- Dr. William “Lin” Oliver III and Dr. Ashley Pifer: as friends, teachers, and advisors, your contributions are extensive.
- Dr. Thad Scott and Dr. Byron Winston: your assistance in experimental setup, sample amendment, and nutrient analysis expanded the scope of this work markedly.
- Duc Thein “B” Do and David Meints: my colleagues and assistants, through many late nights and long discussions.
- Jenny Doyle, Erin Wright, and Marcus Ward: my colleagues, students, assistants, confidants, and teachers – as it should be.
- My family and friends: your patience, tolerance, and distraction have aided me more than just scholastically.

Table of Contents

Introduction.....	1
Materials and Methods.....	3
A. Sampling Location and Nutrient Enrichment Experiments.....	3
B. Water Quality Tests	5
C. Chlorine Dioxide Preparation	5
D. Alum Coagulation Jar Tests	6
E. Fluorescence Measurements.....	6
F. Disinfection Byproducts.....	7
Results and Discussion	8
A. Algal Biomass, Nutrient Concentrations, and N:P Ratios	8
B. Water Quality Tests	8
C. DBPFP Tests.....	10
Conclusions.....	13
References	35

List of Tables

Table 1 – Nitrogen and phosphorus doses and raw water quality data for April 5, 2013 sample collection.14

Table 2 – Nitrogen and phosphorus doses and water quality data of raw and treated waters for May 30, 2013 sample collection.15

Table 3 – Nitrogen and phosphorus doses and water quality data of raw and treated waters for August 19, 2013 sample collection.17

Table 4 – Excitation and Emission maxima of fluorescence-PARAFAC components.19

List of Figures

- Figure 1** – Chlorophyll-*a* (Chl-*a*) of the raw water samples as a function of the phosphorus (P) amendment gradient with constant nitrogen ($2,000 \mu\text{g L}^{-1}$) on a log-log basis. Lines represent the least squares best fit.20
- Figure 2** – Chlorophyll-*a* (Chl-*a*) of the raw water samples as a function of the nitrogen (N) amendment gradient with constant phosphorus ($200 \mu\text{g L}^{-1}$) on a semi-log basis. Lines represent the least squares best fit.21
- Figure 3** – Chlorophyll-*a* (Chl-*a*) of the raw water samples as a function of the molar N:P ratio of all samples on a log-log basis. Lines represent triplicate averages for the May 30 and August 19 sample collection. See Table 1 for details on N:P ratio.22
- Figure 4** – Trichloromethane formation potential (TCMFP) as a function of nitrogen (N) amendment for April 5 raw water. The P dose for all N-amended samples was $200 \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.23
- Figure 5** – Trichloromethane formation potential (TCMFP) as a function phosphorus (P) amendment for April 5 raw water. The N dose for all P-amended samples was $2,000 \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.24
- Figure 6** – Trichloromethane formation potential (TCMFP) as a function of nitrogen (N) amendment for May 30 raw and treated waters (ClO_2 dose of 1 mg L^{-1} as Cl_2 and alum dose of 40 mg L^{-1}). The P dose for all N-amended samples was $200 \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.25

Figure 7 – Trichloromethane formation potential (TCMFP) as a function of phosphorus (P) amendment for May 30 raw and treated waters (ClO_2 dose of 1 mg L^{-1} as Cl_2 and alum dose of 40 mg L^{-1}). The N dose for all P-amended samples was $2,000 \text{ } \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.26

Figure 8 – Trichloromethane formation potential (TCMFP) as a function of nitrogen (N) amendment for August 19 raw and treated waters (ClO_2 dose of 2 mg L^{-1} as Cl_2 and alum dose of 80 mg L^{-1}). The P dose for all N-amended samples was $200 \text{ } \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.27

Figure 9 – Trichloromethane formation potential (TCMFP) as a function of phosphorus (P) amendment for August 19 raw and treated waters (ClO_2 dose of 2 mg L^{-1} as Cl_2 and alum dose of 80 mg L^{-1}). The N dose for all P-amended samples was $2,000 \text{ } \mu\text{g L}^{-1}$. Lines represent triplicate averages except for the $\text{P} = 100 \text{ } \mu\text{g L}^{-1}$ dose, which was excluded. Filled markers represent blank samples without any nutrient amendment.28

Figure 10 – Trichloromethane formation potential (TCMFP) for the raw water samples amended with nitrogen (N) and phosphorus (P) for the April 5, May 30, and August 19 samples as a function of the log-molar N:P ratio, where N and P represent the applied doses. Lines represent triplicate averages for each sample collection. See Table 1 for details on N:P ratio.29

Figure 11 – Trichloromethane formation potential (TCMFP) for the raw water samples amended with nitrogen (N) and phosphorus (P) for the April 5, May 30, and August 19 samples as a function of chlorophyll-a (Chl-a). Lines represent the least squares best fit.30

Figure 12 – Correlations between trichloromethane formation potential (TCMFP) and DOC. Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. DOC is the dissolved organic carbon. Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.31

Figure 13 – Correlations between trichloromethane formation potential (TCMFP) and UV_{254} . Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. UV_{254} is the ultraviolet absorbance at 254 nm. Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.32

Figure 14 – Correlations between trichloromethane formation potential (TCMFP) and $I_{344/425}$. Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. $I_{344/425}$ is the fluorescence intensity at an excitation of 344 nm and an emission of

425 nm. Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.....33

Figure 15 – Correlations between trichloromethane formation potential (TCMFP) and C2 F_{MAX} . Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. C2 F_{MAX} is the maximum fluorescence intensity for PARAFAC Component 2 (see Table 4 for description of the fluorescence-PARAFAC components). Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.34

Introduction

Despite the discovery of disinfection byproducts (DBPs) in chlorinated waters almost four decades ago (Rook, 1976), DBP control at drinking water treatment plants (DWTPs) remains an ongoing challenge. DBPs are formed by reactions between disinfectants (*e.g.*, free chlorine and chlorine dioxide) and natural organic matter (NOM). While over 600 individual DBPs have been identified (Richardson et al., 2007), only 11 are regulated by the United States Environmental Protection Agency under the Stage 2 Disinfectants/DBP Rule – four trihalomethanes (THMs), five haloacetic acids, chlorite, and bromate.

DWTPs can draw from a two-pronged approach to curb formation of regulated DBPs: (1) increase NOM removal, by processes such as enhanced coagulation in which more coagulant is added than is necessary for turbidity removal (Krasner and Amy, 1995; Edzwald and Tobiason, 1999), and (2) switch primary and/or secondary disinfectants. One common primary disinfectant for DWTPs seeking to curb DBPs is chlorine dioxide (ClO_2), which can improve NOM coagulation (Cheng and Chi, 2003) and does not react with NOM to form THMs (Miltner, 1976). However, the use of ClO_2 necessitates the addition of a secondary disinfectant, like free chlorine, to maintain a residual throughout the distribution system. As such, DBPs can still form, but only after some NOM removal has occurred through the coagulation process. The drawbacks of chlorine dioxide addition are that it is reduced to chlorite (Korn et al., 2002; Karanfil et al., 2003), a regulated DBP that can be removed by the addition of ferrous salts, and that it may lyse algal cells and release intracellular organic matter, a potential source of DBP precursors (Plummer and Edzwald, 2002).

It has long been recognized that DBP formation is impacted by nutrient loadings to source waters. As urban and agricultural land use intensifies, nitrogen (N) and phosphorus (P)

enrichments can cause increases in algal biomass and productivity (Carpenter et al., 1998; Smith et al., 1999; Elser et al., 2007), decreasing the availability of pristine water supplies. Increased algal biomass and extracellular products (Myklestad, 1995) can react with disinfectants to form DBPs (Hoehn et al., 1980; Graham et al., 1998; Jack et al., 2002; Fang et al., 2010). In addition to elevated nutrients increasing algal biomass, the ratio of N:P can influence the type of algae growing in lakes (Schindler, 1974; Smith, 1983), which also has consequences for water quality. Eutrophic waters often have high algal productivity and lower N:P ratios (Downing and McCauley, 1992), which favor nitrogen-fixing cyanobacteria, and can deteriorate water quality through the production of toxins and taste-and-odor forming compounds (Huisman et al., 2005). On the other hand, oligotrophic lakes are often characterized by low productivity and high N:P ratios, conditions under which cyanobacteria are rare and diatoms typically dominate the phytoplankton community composition.

Despite these previous research efforts, comparatively little is known about DBP formation and control in waters enriched across environmentally relevant gradients of N and P. Such work is important to help guide nutrient management strategies and to assist DWTPs in adapting DBP control processes for increasingly impaired water sources. The research objective of this work was to assess the effect of algal growth driven by N and P enrichments on DBP formation and control. Source water was sampled in the spring and summer 2013 from Beaver Lake near a DWTP intake (Lowell, AR) and amended with N and P at various N:P ratios to stimulate biomass growth. To simulate DBP-precursor removal processes at DWTPs, these waters were subjected to ClO_2 oxidation and alum coagulation. After each treatment, the samples were filtered and various DBP-precursor surrogate parameters were measured (Pifer et al., in press). The raw and treated waters were chlorinated to assess the DBP formation potential

(DBPFP) as a function of N and P amendments, and correlations were sought between DBPFP and the various precursor surrogate parameters.

Materials and Methods

A. Sampling Location and Nutrient Enrichment Experiments

Source waters were collected from the transition zone of Beaver Lake Reservoir (Lowell, AR) near the Beaver Water District (BWD) DWTP intake structure and used as an algae seed culture. This reservoir provides drinking water and recreation opportunities for the Northwest Arkansas region. It has an average depth of 18-m and an average hydraulic retention time of 1.5 years. Trophic conditions range from eutrophic at the mouth of the White River to oligotrophic near the dam. The reservoir is also fed by Richland Creek, War Eagle Creek, and Brush Creek, and comprises a total hydraulic catchment area of 300,000-ha of largely forested (69%) and agricultural (26%) land (Sen et al., 2007).

Beaver Lake water was collected from a boat in the spring and summer of 2013 on April 5, May 30, and August 19. On each day, a 120-L composite sample was collected from across the photic zone and transported to the University of Arkansas for bioassay experiments. Samples were mixed and dispensed in 3-L aliquots into 4-L acid-washed plastic cubitainers. For each sampling date, a total of 36 cubitainers were used for a nutrient enrichment experiment. The nutrient enrichment bioassay experiment on each date was intended to create various nutrient-amendment rates and various N:P ratios. A P enrichment gradient of 0, 0.025, 0.05, 0.075, 0.1, and 0.2 mg L⁻¹ P as disodium hydrogen phosphate (Na₂HPO₄) along with 2 mg L⁻¹ nitrogen as potassium nitrate (KNO₃) was created to achieve 6 triplicate N:P ratios of ~4429, 442, 177, 89, 44, and 22 by moles, respectively. A separate N enrichment gradient of 0, 0.1, 0.25, 0.5, and 1 mg L⁻¹ N (as KNO₃) along with 0.2 mg L⁻¹ P (as Na₂HPO₄) was created to achieve 5 triplicate

molar N:P ratios of ~0.22, 1.1, 2.8, 5.5, and 11.1, respectively. As such, the combined N:P ratio gradient spanned almost five orders of magnitude, while the N and P enrichment gradients spanned more than one order of magnitude each.

After N and P amendment, samples were placed in a 30°C water bath under artificial lighting. Lights were controlled by a 12-hour on/off timer and measured to be 500 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ during illumination. Each cubitainer was opened to the atmosphere and shaken daily by hand to aid in aeration and minimize attached growth. Algal biomass was estimated daily as raw water fluorescence measurements using a Turner Design Trilogy fluorometer (Turner Designs, Sunnyvale, CA) at 880 nm. Once the samples had achieved their maximum biomass (~4 days), the cubitainers were shaken vigorously and 2-L were poured into prepared HDPE containers. These containers were stored in the dark at 4°C for DBPFP experiments. The remaining cubitainer volume was divided evenly for analyses of phytoplankton biomass and particulate nutrients. Aliquots were filtered onto Whatman glass fiber filters (GFFs) and stored frozen for measurement of phytoplankton biomass as extracted chlorophyll-*a* (Chl-*a*).

Chl-*a* was measured to estimate phytoplankton biomass according to Standard Methods 10200 H (Eaton et al., 2005), with modifications. One filter from each sample was protected from light and transferred to a 15 mL test tube containing 7 mL of 90% acetone solution. The samples were placed in a dark freezer for 24 hours to further enhance pigment extraction. In a dark room, 3 mL of each sample extract were then transferred into disposable test tubes and were analyzed using the Turner Design fluorometer at 880 nm. To adjust for the chlorophyll degradation product pheophytin, each sample was re-measured 90 seconds after addition of 0.1 mL of 0.1 N HCl.

B. Water Quality Tests

Laboratory glassware and plastic ware were prepared in accordance with previous work (Pifer et al., 2011). All stock chemicals used were ACS grade, and aqueous solutions were made with Milli-Q water (18.2 M Ω -cm) generated by a Millipore Integral 3 (Billerica, MA) water purification system. The pH and turbidity of the raw waters were measured using equipment and methods described previously (Pifer et al., 2011). Prior to measurement of dissolved organic carbon (DOC) and UV absorbance, samples were filtered through prepared 0.45- μ m nominal pore size polyethersulfone (PES) membranes. These filters were prepared by rinsing with 500-mL of Milli-Q water prior to use (Karanfil et al., 2003). The first 25-mL of filtered sample was wasted for each new filter, to minimize organic carbon adsorption. Filtered samples were then stored in 250-mL amber glass screw top bottles in the dark at 4°C. DOC analysis was performed on a Sievers 900 Portable Total Organic Carbon Analyzer (GE Analytical Instruments, Boulder, CO). UV absorbance scans from 225- to 600 nm were performed on a Shimadzu UV-Vis 2450 (Kyoto, Japan) spectrophotometer using a 1-cm path length low volume quartz cell.

C. Chlorine Dioxide Preparation

Chlorine dioxide was generated using methods described previously (Granderson et al., 2013). Before dosing, raw water samples were poured into prepared 1-L amber glass screw top bottles and placed in a water bath at 24°C. The stock chlorine dioxide concentration was measured by absorptivity at 360-nm after dilution with Milli-Q water, using an assumed molar absorptivity of 1,225 M⁻¹ cm⁻¹. The nutrient amended samples generated from source water collected on May 30, 2013 were dosed with chlorine dioxide at 1 mg L⁻¹, whereas the August 19 samples were dosed at 2 mg L⁻¹. After dosing, samples were capped headspace-free and placed in the dark at room temperature for 24 hours.

D. Alum Coagulation Jar Tests

After the chlorine dioxide dosing and hold time, 500-mL aliquots of each sample water were alum coagulated in square-bottom plastic jars equipped with 5-cm magnetic PTFE stir bars with ring-collared ends on an eight-position magnetic stir plate (Challenge Technology, Springdale, AR). Samples were mixed at 200 rpm to simulate rapid mix conditions prior to the simultaneous addition of alum (aluminum sulfate octadecahydrate) as a coagulant and sodium carbonate to aid in pH control. May 30 samples were dosed with 40 mg L⁻¹ alum and 25 mg L⁻¹ sodium carbonate, while August 19 samples were dosed with 80 mg L⁻¹ alum and 85 mg L⁻¹ sodium carbonate. After 30 seconds of rapid mix (~200 rpm), the jars were moved to an adjacent eight-position magnetic stir plate for flocculation at 40 rpm for 30 minutes. The samples were then allowed to settle quiescently for at least 30 minutes before decanting. The supernatant was characterized and filtered as described in the Water Quality Tests, then used for subsequent experiments as detailed in the remainder of this section.

E. Fluorescence Measurements

Fluorescence excitation-emission matrices (EEMs) were collected for every raw and treated water sample (244 EEMs). Excitation wavelengths ranged from 225- to 400 nm in 1 nm step sizes and emission data was collected from 270- to 600-nm in 1 nm step sizes, resulting in a total of 58,256 fluorescence intensity values, $I_{Ex/Em}$, per EEM. Scatter correction methods used were described previously (Zepp et al., 2004; Pifer et al., 2011). For the group of 244 EEMs, each $I_{Ex/Em}$ pair was regressed against the DBPFP data using an in-house MATLAB[®] code.

In addition to the pair-picking procedure, EEM data was modeled with PARAFAC analysis, following methods described previously (Pifer et al., 2011). Of the 244 EEM sample set, one sample was classified as an outlier and removed from the dataset based on high leverage

and apparent measurement error (Stedmon and Bro, 2008). A 5-component model was validated using split-halves analysis as detailed previously (Pifer et al., 2011), and fluorescence maximum (F_{MAX}) values from each component and EEM were used in DBPFP regression analyses.

F. Disinfection Byproducts

The DBPFP was measured following Standard Methods 5710 B (Eaton et al., 2005). Filtered samples were poured into 125-mL amber glass bottles and buffered with a phosphate solution to $\text{pH } 7.0 \pm 0.2$. Sodium hypochlorite stock solution was standardized following Standard Methods 4500-Cl B, and then diluted to a lower concentration (between 2- and 4 g L^{-1} as Cl_2) for dosing with a micropipette. The free chlorine dose required to achieve 7-day chlorine residuals of 3- to 5 mg L^{-1} as Cl_2 was estimated based on raw water DOC. Free chlorine doses were stair-stepped with nutrient loading and ranged from 9- to 22 mg L^{-1} as Cl_2 . After addition of free chlorine, samples were capped headspace-free and placed in the dark at room temperature. After seven days, the chlorine residual was measured. Standards of free chlorine were prepared and analyzed with DPD total chlorine reagent powder pillows (Hach Company) and a spectrophotometer (Shimadzu UV-vis 2450) to provide a measurement range of 1- to 7 mg L^{-1} as Cl_2 ($n = 5$, $R^2 = 0.99$, data not shown). An aliquot of sample was wasted before gently inverting the bottle three times, to minimize possible sample stratification. Precisely 5 mL of sample was pipetted into 5 mL of Milli-Q water for measurement of chlorine residual, to accommodate high residuals.

Precisely 30 mL of the remaining sample was withdrawn for DBPFP testing as described previously (Pifer and Fairey, 2012), with modifications. Two additional standard curve concentrations ($150 \text{ } \mu\text{g L}^{-1}$ and $200 \text{ } \mu\text{g L}^{-1}$) were added to encompass higher trichloromethane (TCM) yields. Blanks and check standards were analyzed every 18 injections for quality control

and 90% of check standards were within $\pm 20\%$ of the standard concentration, and all check standards were within $\pm 25\%$, which is considered to be acceptable based on EPA 551.1.

Results and Discussion

A. Algal Biomass, Nutrient Concentrations, and N:P Ratios

Algal biomass, measured as Chl-*a*, increased proportionally along the P enrichment gradient when N availability was high in experiments from all three months (Figure 1). Similarly, algal biomass increased along the N enrichment gradient when P availability was high in the August 19 experiment only (Figure 2). As a result, there was an obvious pattern in algal biomass along the experimental N:P gradient (Figure 3). For the May 30 and August 19 samples, algal biomass was greatest at intermediate N:P (~ 5 -50 by moles) and decreased substantially when the molar N:P ratio exceeded ~ 80 , indicating P-limiting conditions. These results indicate that P at least partially controlled algal biomass in Beaver Lake throughout the summer of 2013. Nitrogen exerted little control on algal biomass in spring, but partially controlled algal biomass in August (Figure 2). These results are consistent with previously reported patterns showing the seasonal transition between P- and N-limited algal growth in southern U.S. river impoundment reservoirs (Scott et al., 2008; Scott and Grantz, 2013).

B. Water Quality Tests

Raw water quality results for the April 5 sample collection are shown in Table 1. DOC increased with P dose from an average of 2.26- to 2.77 mg L⁻¹ as C, suggesting the increased algal biomass (Figure 1) augmented the DOC by release of extracellular organic matter. While UV₂₅₄ increased with P dose, the average SUVA decreased from 1.89- to 1.81 mg L⁻¹ m⁻¹, indicating the DOC produced was not enriched with aromatic carbon. This is a noteworthy result given the aromatic carbon fraction has been shown to be a significant source of THM precursors

(Weishaar et al., 2003). In contrast with the trends in P dose, DOC, UV_{254} , and SUVA did not change across the range of N doses. Taken together, these results suggest P-limited growth for the April 5 sampling set, which is consistent with the biomass data (Figures 1-3). The free chlorine residuals after 7 days (FC-7d) were between 4- and 7 $mg\ L^{-1}$ as Cl_2 , with no trends based on the N or P dose.

Raw and treated water quality results for the May 30 sample collection are shown in Table 2. Similar to the April results, raw water DOC increased with P dose from an average of 3.99- to 4.91 $mg\ L^{-1}$ as C and did not increase uniformly with N dose, indicating P-limited growth. For all twelve N and P doses, ClO_2 treatment increased the average DOC and decreased the average SUVA, suggesting algal cells were lysed by ClO_2 oxidation and released intracellular organic matter with relatively low aromatic carbon content, similar to previous results (Li et al., 2012). Subsequent alum coagulation decreased the average DOC below their corresponding raw waters in all 6 cases across the P gradient, but only in 3 of 5 cases across the N gradient. This indicates that DOC produced by N enrichment was more resistant to removal by alum coagulation. It is worth noting that the average FC-7d residuals in Table 2 were between 10- and 16 $mg\ L^{-1}$ as Cl_2 , above the target window of 3-5 $mg\ L^{-1}$ as Cl_2 for the DBPFP tests. Ongoing experiments in our laboratory suggest these higher residuals will enhance formation of chlorinated THMs at the expense of bromine-substituted species and haloacetonitriles.

Raw and treated water quality results for the August 19 sample collection are shown in Table 3. For the P-gradient, the raw water DOC ranged from 2.96- to 3.35 $mg\ L^{-1}$ as C, but in contrast to April and May samples only increased for the two highest P doses (100- and 200 $\mu g\ L^{-1}$). No discernible trends in average DOC were apparent across the N gradient, although Figure 2 indicates N was co-limiting for the August 19 samples. ClO_2 treatment increased the average

DOC and decreased the average SUVA, supporting the previous results (Table 2) that lysis of algal cells occurred and released DOC depleted in aromatic carbon. Subsequent alum coagulation decreased the average DOC relative to their corresponding raw waters for all 11 nutrient amended samples. The ranges of the average SUVA for raw, ClO₂-treated only, and ClO₂+alum coagulated waters were 1.54-1.70 mg L⁻¹ m⁻¹, 1.20-1.36 mg L⁻¹ m⁻¹, and 1.28-1.61 mg L⁻¹ m⁻¹. The modest increase in SUVA following alum coagulation of ClO₂-treated waters for all 11 samples was unexpected and suggests that alum coagulation preferentially removed the less aromatic DOC. FC-7d residuals ranged from 5- to 9 mg L⁻¹ as Cl₂, more inline with the target residual for the DBPFP tests (3-5 mg L⁻¹ as Cl₂) compared to the April samples (Table 2), but nevertheless relatively high, which, as stated previously, favors the formation of chlorinated THMs.

C. DBPFP Tests

As expected based on the high free chlorine residuals (Tables 1, 2, and 3) trichloromethane (TCM) was the predominant DBP formed, comprising 89-98% by mass of the total THMs (data not shown). Additionally, other DBPs quantified as part of EPA 551.1, such as dichloroacetonitrile, formed at relatively low concentrations (below 1.76 µg L⁻¹) and, as a result, further discussion is focused on TCM only. TCMFP results are presented in Figures 4-9, organized by sample month (April 5, May 30, and August 19) and nutrient gradient (N or P). The relatively high raw water TCMFP concentrations for the May 30 samples (approximately 50 µg L⁻¹ higher than the April 5 and August 19 samples) are likely due to the comparatively high FC-7d values (Tables 1, 2, and 3), rather than a greater abundance of TCM precursors. For the April 5 samples, the average TCMFP did not change across the N amendment (Figure 4), but increased 13% across the P amendment (from 90.0 to 102.8 µg L⁻¹, Figure 5). For the May 30 samples, the

average TCMFP in raw waters showed similar trends, with no increase across the N amendment (Figure 6), and an increase of 15% across the P amendment (from 165.7- to 195.1 $\mu\text{g L}^{-1}$, Figure 7). For the August 19 samples, by contrast, the average TCMFP in the raw waters increased 18% across N amendment (from 103.9- to 126.9 $\mu\text{g L}^{-1}$, Figure 8), and 9% across the P amendment (from 106.8- to 117.3 $\mu\text{g L}^{-1}$, Figure 9). For the raw water samples, TCMFP was greatest at intermediate values of the experimental N:P gradient (~ 10 -50, Figure 10), which corresponded with the greatest algal biomass across all experiments (Figure 3). Thus, TCMFP was positively correlated with algal biomass as Chl-*a* in all experiments, with the steepest and strongest relationship occurring for the May 30 samples (Figure 11).

Treatment of raw waters occurred for the samples collected on May 30 and August 19 only. The May 30 samples were treated with ClO_2 at 1 mg L^{-1} and an alum dose of 40 mg L^{-1} ; to achieve greater TCM precursor removal, both of these doses were doubled for the August 19 samples. Figure 5 shows that treatment with 1 mg L^{-1} ClO_2 increased the average TCMFP relative to the raw waters for the lowest two N amendments, and was similar to the raw waters for the higher N doses. Figure 6 shows this same dose of ClO_2 had little impact on TCMFP across the P amendment. This result indicates that the aromatic carbon depleted DOC released by ClO_2 treatment (Table 2 – DOC and SUVA) was not a significant source of TCM precursors. For August 19 samples, a ClO_2 dose of 2 mg L^{-1} decreased the average TCMFP by 20-30 $\mu\text{g L}^{-1}$ across the N amendments (Figure 8) and 22-47 $\mu\text{g L}^{-1}$ across the P amendments (Figure 9). Further, Figure 9 shows that the differences in TCMFP between the raw and ClO_2 treated samples decreased with increasing P amendment, presumably because the biomass produced (Figure 1) exerted a demand for ClO_2 , more so than directly contributing to the TCM precursor pool.

Alum coagulation following ClO_2 treatment lowered the average TCMFP, an expected result based on previous research (Granderson et al., 2013). The one exception to this trend occurred for the May 30 samples at an N amendment of $1000 \mu\text{g L}^{-1}$ (Figure 6), in which the average TCMFP values were similar for both treatments. Figure 7 shows that alum coagulation decreased the average TCMFP by $34\text{-}64 \mu\text{g L}^{-1}$ compared to ClO_2 -only, but the difference between treatments decreased as the P amendment increased. For the August 19 samples, alum coagulation decreased TCMFP by $10\text{-}20 \mu\text{g L}^{-1}$ relative to ClO_2 -only for both nutrient amendments (Figures 8-9). The implication of this result for DWTPs is that ClO_2 pre-oxidation and alum coagulation may be less effective for removal of TCM precursors as source waters become more nutrient enriched.

To further explain the TCMFP data, correlations were sought with known TCM precursor surrogate parameters (*e.g.*, UV_{254} , DOC, $I_{\text{Ex/Em}}$, and PARAFAC component F_{MAX} values). For this dataset, $I_{344/425}$ and F_{MAX} from Component 2 (Table 4) were the most strongly correlated fluorescence metrics ($I_{\text{Ex/Em}}$ correlation results not shown). Figures 12-15 show correlations between TCMFP and (i) DOC ($r^2 = 0.72$, Figure 12), (ii) UV_{254} ($r^2 = 0.88$, Figure 13), (iii) $I_{344/425}$ ($r^2 = 0.62$, Figure 14), and (iv) C2 F_{MAX} ($r^2 = 0.61$, Figure 15). A weaker correlation was found between TCMFP and SUVA ($r^2 = 0.57$, data not shown), an expected result given that SUVA is an intensive property. Data presented in Fig. 4 includes all samples and treatments except seven samples (out of 244) that were determined to be outliers – five of these samples had TCM concentrations that were 150% greater (*e.g.*, $300\text{-}700 \mu\text{g L}^{-1}$) than the highest value in the GC standard curve, one sample had no measurable FC-7d residual, and the other sample was determined to be an outlier during the PARAFAC modeling process. The comparatively strong TCMFP:DOC correlation ($r^2 = 0.72$, Figure 12) was unexpected because ClO_2 treatment

increased DOC (Tables 2 and 3) but decreased TCMFP (Figures 6-9). The high TCMFP:UV₂₅₄ correlation ($r^2 = 0.88$, Figure 13) is in agreement with prior research (Edzwald et al., 1985), supporting the contention that released DOC from nutrient stimulated biomass was both low in aromatic carbon and did not contribute significantly to the pool of TCM precursors. The comparatively weak correlations between TCMFP and the fluorescence metrics (Figures 14-15) were unexpected based on previous research (Pifer and Fairey, 2012; Granderson et al., 2013) and suggest that dissolved species present in the samples from the nutrient enrichments (*e.g.*, algal extrudates and intracellular organic matter) may have interfered with fluorescence measurements more so than UV₂₅₄.

Conclusions

The experiments presented here demonstrate that nutrient-driven increases in algal biomass reduced the effectiveness of two common DBP control measures, ClO₂ oxidation and alum coagulation. Algal biomass in nutrient amended waters was shown to be P-limited for the April 5, May 30, and August 19 sampling sets, with an N co-limitation for the August 19 samples. For the nutrient amended raw waters, algal biomass, measured as Chl-*a*, was a maximum at molar N:P ratios of ~10-50, which following chlorination corresponded to a measurable increase in the TCMFP. Oxidation of the sample waters with chlorine dioxide increased the DOC with aromatic-depleted compounds that were not significant TCM precursors. Across the experimental P-gradient, the differences in TCMFP between the raw and ClO₂+alum coagulated waters decreased with increasing P amendment, indicating the algal biomass exerted a demand for ClO₂ and alum. Results from this study can be used to guide nutrient management strategies for source water protection and can be used by DWTPs to assess the impact of N and P enrichments on TCM formation and control.

Table 1 – Nitrogen and phosphorus doses and raw water quality data for April 5, 2013 sample collection.

N Dose ($\mu\text{g L}^{-1}$)	P Dose ($\mu\text{g L}^{-1}$)	N:P (mol/mol)	DOC (mg L^{-1})	UV ₂₅₄ (m^{-1})	SUVA ($\text{mg L}^{-1} \text{m}^{-1}$)	FC Dose/FC-7d (mg L^{-1} as Cl ₂)
0	0	NA	2.31	4.3	1.86	9/5.22
2000	0	4429	2.26 ± 0.02	4.3 ± 0.1	1.89 ± 0.04	$9/5.59 \pm 0.13$
2000	10	442.3	2.37 ± 0.05	4.5 ± 0.1	1.89 ± 0.06	$10/6.02 \pm 0.04$
2000	25	176.9	2.44 ± 0.03	4.6 ± 0.0	1.89 ± 0.02	$11/6.34 \pm 0.16$
2000	50	88.5	2.50 ± 0.07	4.7 ± 0.1	1.87 ± 0.04	$12/6.64 \pm 0.24$
2000	100	44.2	2.56 ± 0.05	4.6 ± 0.2	1.81 ± 0.03	$12/6.59 \pm 0.11$
2000	200	22.1	2.77 ± 0.10	5.0 ± 0.0	1.81 ± 0.06	$13/6.85 \pm 0.17$
0	200	0.2	2.87 ± 0.07	5.0 ± 0.1	1.76 ± 0.03	$9/4.30 \pm 0.07$
100	200	1.1	2.83 ± 0.09	5.0 ± 0.1	1.77 ± 0.02	$10/5.00 \pm 0.13$
250	200	2.8	2.80 ± 0.05	5.0 ± 0.1	1.77 ± 0.01	$9/4.44 \pm 0.08$
500	200	5.5	2.82 ± 0.09	5.1 ± 0.1	1.80 ± 0.05	$12/6.09 \pm 0.24$
1000	200	11.1	2.87 ± 0.07	5.0 ± 0.1	1.75 ± 0.02	$13/6.48 \pm 0.32$

Values are averages \pm standard deviations.

N = Nitrogen added as KNO₃; P = Phosphorus added as Na₂HPO₄; DOC = Dissolved Organic Carbon; UV₂₅₄ = Ultraviolet Absorbance at 254 nm; SUVA = Specific UV₂₅₄ (UV₂₅₄/DOC); FC = free chlorine; FC-7d = free chlorine residual after 7-day hold time; N:P = molar nitrogen to phosphorus ratio based on amended doses, with the exception of two values (4429 and 0.2) which were calculated using the initial background concentrations; NA = not applicable.

Note: Free chlorine was dosed after all other reported measurements.

Table 2 – Nitrogen and phosphorus doses and water quality data of raw and treated waters for May 30, 2013 sample collection.

Sample Type	N Dose ($\mu\text{g L}^{-1}$)	P Dose ($\mu\text{g L}^{-1}$)	pH	Turbidity (NTU)	DOC (mg L^{-1})	UV ₂₅₄ (m^{-1})	SUVA ($\text{mg L}^{-1} \text{m}^{-1}$)	FC Dose/FC-7d ($\text{mg L}^{-1} \text{as Cl}_2$)
R	0	0	8.18	12.00	4.05	10.0	2.47	18/13.54
C	0	0	7.79	8.50	4.52	8.6	1.90	18/13.56
CA	0	0	NM	NM	3.31	4.6	1.39	18/15.66
R	2000	0	8.14 ± 0.02	9.23 ± 0.15	3.99 ± 0.06	9.5 ± 0.0	2.38 ± 0.03	$18/13.74 \pm 0.23$
C	2000	0	7.80 ± 0.03	8.70 ± 0.44	4.37 ± 0.05	8.6 ± 0.1	1.97 ± 0.02	$18/13.53 \pm 0.19$
CA	2000	0	NM	NM	3.02 ± 0.13	4.1 ± 0.1	1.37 ± 0.05	$18/15.69 \pm 0.45$
R	2000	10	9.07 ± 0.08	9.60 ± 0.00	4.08 ± 0.04	9.4 ± 0.1	2.30 ± 0.03	$19/14.00 \pm 0.33$
C	2000	10	8.22 ± 0.09	10.33 ± 0.29	4.56 ± 0.28	8.7 ± 0.0	1.91 ± 0.11	$19/14.49 \pm 0.50$
CA	2000	10	NM	NM	3.37 ± 0.15	4.4 ± 0.2	1.31 ± 0.02	$19/15.82 \pm 0.23$
R	2000	25	9.37 ± 0.08	9.67 ± 0.83	4.18 ± 0.08	9.5 ± 0.2	2.28 ± 0.05	$20/14.52 \pm 0.17$
C	2000	25	8.76 ± 0.12	10.83 ± 0.76	4.67 ± 0.10	9.1 ± 0.2	1.95 ± 0.01	$20/14.56 \pm 0.45$
CA	2000	25	NM	NM	3.66 ± 0.18	4.8 ± 0.2	1.31 ± 0.02	$20/15.97 \pm 0.26$
R	2000	50	9.84 ± 0.04	11.33 ± 0.58	4.32 ± 0.03	9.7 ± 0.2	2.24 ± 0.05	$21/14.60 \pm 0.64$
C	2000	50	9.44 ± 0.06	10.50 ± 0.87	4.89 ± 0.04	9.4 ± 0.1	1.93 ± 0.03	$21/14.15 \pm 0.49$
CA	2000	50	NM	NM	3.75 ± 0.04	6.2 ± 0.5	1.66 ± 0.12	$21/15.66 \pm 0.08$
R	2000	100	10.07 ± 0.04	11.00 ± 0.00	4.55 ± 0.15	10.1 ± 0.2	2.21 ± 0.03	$21/14.09 \pm 0.27$
C	2000	100	9.73 ± 0.02	11.67 ± 0.29	5.17 ± 0.12	9.7 ± 0.1	1.87 ± 0.03	$21/12.91 \pm 1.05$
CA	2000	100	NM	NM	4.56 ± 0.42	9.3 ± 0.4	2.05 ± 0.10	$21/15.38 \pm 0.25$
R	2000	200	10.26 ± 0.01	11.75 ± 0.35	4.91 ± 0.13	10.6 ± 0.2	2.15 ± 0.01	$22/13.93 \pm 0.07$
C	2000	200	9.78 ± 0.03	11.40 ± 5.09	6.79 ± 1.77	9.9 ± 0.1	1.50 ± 0.40	$22/12.36 \pm 0.72$
CA	2000	200	NM	NM	4.52 ± 0.45	8.9 ± 0.6	1.96 ± 0.06	$22/14.12 \pm 0.27$
R	0	200	10.11 ± 0.20	12.67 ± 0.58	4.66 ± 0.17	9.8 ± 0.4	2.11 ± 0.09	$18/11.54 \pm 0.25$

C	0	200	9.67 ± 0.17	7.13 ± 3.35	5.45 ± 0.38	9.5 ± 0.1	1.74 ± 0.12	18/11.18 ± 0.27
CA	0	200	NM	NM	5.75 ± 0.72	7.5 ± 1.2	1.32 ± 0.26	18/12.69 ± 0.80
R	100	200	10.19 ± 0.08	11.67 ± 0.58	6.58 ± 3.31	10.1 ± 0.4	1.75 ± 0.65	19/11.90 ± 1.85
C	100	200	9.78 ± 0.11	15.33 ± 2.08	7.20 ± 3.29	9.6 ± 0.1	1.50 ± 0.54	19/9.95 ± 1.78
CA	100	200	NM	NM	6.07 ± 2.67	7.8 ± 0.7	1.47 ± 0.61	19/12.27 ± 1.91
R	250	200	10.25 ± 0.10	12.00 ± 0.00	4.72 ± 0.09	10.1 ± 0.3	2.14 ± 0.05	20/12.53 ± 1.32
C	250	200	9.71 ± 0.08	12.33 ± 0.58	5.14 ± 0.03	9.6 ± 0.1	1.88 ± 0.02	20/11.89 ± 0.52
CA	250	200	NM	NM	4.12 ± 0.20	7.8 ± 0.6	1.90 ± 0.07	20/13.89 ± 0.26
R	500	200	10.28 ± 0.01	12.00 ± 0.00	4.66 ± 0.09	9.9 ± 0.2	2.13 ± 0.04	21/13.55 ± 0.18
C	500	200	9.82 ± 0.06	14.33 ± 1.15	5.21 ± 0.03	9.8 ± 0.1	1.87 ± 0.02	21/11.97 ± 0.27
CA	500	200	NM	NM	4.70 ± 0.18	9.6 ± 0.2	2.05 ± 0.12	21/13.46 ± 0.44
R	1000	200	10.29 ± 0.07	11.33 ± 0.58	4.98 ± 0.55	9.9 ± 0.1	2.01 ± 0.20	22/14.42 ± 0.68
C	1000	200	9.85 ± 0.04	13.33 ± 0.58	5.09 ± 0.02	9.7 ± 0.1	1.91 ± 0.01	22/12.97 ± 0.09
CA	1000	200	NM	NM	4.68 ± 0.43	10.1 ± 0.4	2.16 ± 0.19	22/13.33 ± 0.29

Values are averages ± standard deviations.

N = Nitrogen added as KNO₃; P = Phosphorus added as Na₂HPO₄; DOC = Dissolved Organic Carbon; UV₂₅₄ = Ultraviolet

Absorbance at 254 nm; SUVA = Specific UV₂₅₄ (UV₂₅₄/DOC); FC = free chlorine; FC-7d = free chlorine residual after 7-day hold time; C = Chlorine dioxide dosed at 1 mg L⁻¹ as Cl₂; CA = Chlorine dioxide dosed at 1 mg L⁻¹ as Cl₂ and Alum coagulation at 40 mg L⁻¹ as alum; R = nutrient amended raw water; NM = not measured

Note: Free chlorine was dosed after all other reported measurements.

Table 3 – Nitrogen and phosphorus doses and water quality data of raw and treated waters for August 19, 2013 sample collection.

Sample Type	N Dose ($\mu\text{g L}^{-1}$)	P Dose ($\mu\text{g L}^{-1}$)	pH	Turbidity (NTU)	DOC (mg L^{-1})	UV ₂₅₄ (m^{-1})	SUVA ($\text{mg L}^{-1} \text{m}^{-1}$)	FC Dose/FC-7d (mg L^{-1} as Cl ₂)
R	0	0	8.63	3.20	3.10	4.8	1.55	9/5.36
C	0	0	7.94	2.70	3.47	4.2	1.21	9/5.23
CA	0	0	8.23	0.90	3.23	3.3	1.02	9/6.11
R	0	0	8.83 ± 0.03	1.53 ± 0.06	3.09 ± 0.06	4.8 ± 0.0	1.56 ± 0.03	10/6.47 ± 0.04
C	0	0	8.12 ± 0.06	1.83 ± 0.12	3.18 ± 0.07	3.9 ± 0.1	1.24 ± 0.03	10/6.43 ± 0.07
CA	0	0	8.32 ± 0.04	0.43 ± 0.03	2.72 ± 0.07	3.5 ± 0.1	1.29 ± 0.05	10/7.01 ± 0.10
R	2000	0	8.94 ± 0.17	1.80 ± 0.26	3.09 ± 0.03	5.0 ± 0.1	1.61 ± 0.03	10/6.58 ± 0.15
C	2000	0	8.21 ± 0.25	1.80 ± 0.26	3.25 ± 0.03	4.0 ± 0.1	1.24 ± 0.01	10/6.23 ± 0.36
CA	2000	0	8.28 ± 0.03	0.42 ± 0.05	2.77 ± 0.03	3.6 ± 0.2	1.30 ± 0.06	10/7.11 ± 0.20
R	2000	10	8.92 ± 0.13	1.53 ± 0.15	3.10 ± 0.01	5.0 ± 0.0	1.61 ± 0.01	11/7.91 ± 0.51
C	2000	10	8.23 ± 0.12	1.77 ± 0.21	3.18 ± 0.05	3.9 ± 0.1	1.22 ± 0.00	11/7.41 ± 0.24
CA	2000	10	8.26 ± 0.01	0.40 ± 0.08	2.69 ± 0.14	3.6 ± 0.2	1.32 ± 0.04	11/8.51 ± 0.49
R	2000	25	9.25 ± 0.01	2.43 ± 0.23	3.06 ± 0.03	4.9 ± 0.1	1.61 ± 0.03	11/7.80 ± 0.34
C	2000	25	8.61 ± 0.05	2.13 ± 0.42	3.34 ± 0.02	4.3 ± 0.1	1.28 ± 0.01	11/7.25 ± 0.54
CA	2000	25	8.34 ± 0.03	0.83 ± 0.32	2.70 ± 0.05	3.8 ± 0.1	1.40 ± 0.00	11/8.11 ± 0.39
R	2000	50	9.36 ± 0.03	2.87 ± 0.57	2.96 ± 0.04	5.0 ± 0.1	1.70 ± 0.01	12/8.75 ± 0.26
C	2000	50	8.78 ± 0.05	3.20 ± 0.20	3.40 ± 0.06	4.4 ± 0.1	1.29 ± 0.02	12/8.25 ± 0.38
CA	2000	50	8.36 ± 0.01	0.83 ± 0.32	2.77 ± 0.06	3.8 ± 0.1	1.37 ± 0.05	12/9.33 ± 0.29
R	2000	100	9.55 ± 0.28	4.23 ± 0.75	3.24 ± 0.03	5.1 ± 0.1	1.58 ± 0.04	12/7.81 ± 0.88
C	2000	100	9.00 ± 0.28	4.53 ± 0.64	3.76 ± 0.51	4.7 ± 0.4	1.27 ± 0.09	12/7.25 ± 0.82
CA	2000	100	8.37 ± 0.03	0.77 ± 0.15	3.12 ± 0.39	4.2 ± 0.4	1.34 ± 0.11	12/8.21 ± 0.54
R	2000	200	9.80 ± 0.12	5.40 ± 0.53	3.35 ± 0.08	5.3 ± 0.1	1.57 ± 0.01	13/7.83 ± 0.27

C	2000	200	9.28 ± 0.16	5.57 ± 0.25	3.73 ± 0.05	5.1 ± 0.2	1.36 ± 0.03	13/7.71 ± 0.07
CA	2000	200	8.60 ± 0.11	1.27 ± 0.29	3.03 ± 0.08	4.9 ± 0.7	1.61 ± 0.20	13/8.81 ± 0.19
R	0	200	9.34 ± 0.01	2.23 ± 0.25	3.27 ± 0.03	5.2 ± 0.2	1.60 ± 0.03	10/6.96 ± 0.17
C	0	200	8.71 ± 0.04	2.30 ± 0.17	3.43 ± 0.07	4.1 ± 0.1	1.20 ± 0.02	10/6.26 ± 0.10
CA	0	200	8.43 ± 0.04	0.77 ± 0.21	2.88 ± 0.03	3.7 ± 0.1	1.28 ± 0.04	10/7.10 ± 0.16
R	100	200	9.56 ± 0.01	3.30 ± 0.62	3.17 ± 0.03	5.1 ± 0.1	1.62 ± 0.02	11/7.47 ± 0.10
C	100	200	9.06 ± 0.05	3.63 ± 0.15	3.72 ± 0.06	4.5 ± 0.1	1.22 ± 0.04	11/6.69 ± 0.11
CA	100	200	8.49 ± 0.03	0.87 ± 0.31	3.06 ± 0.08	4.3 ± 0.2	1.39 ± 0.02	11/7.63 ± 0.20
R	250	200	9.67 ± 0.02	3.53 ± 0.50	3.24 ± 0.02	5.1 ± 0.1	1.58 ± 0.01	12/8.35 ± 0.15
C	250	200	9.20 ± 0.01	4.07 ± 0.45	3.84 ± 0.05	4.7 ± 0.1	1.23 ± 0.00	12/7.10 ± 0.07
CA	250	200	8.52 ± 0.02	0.97 ± 0.21	3.12 ± 0.01	4.7 ± 0.1	1.50 ± 0.03	12/8.47 ± 0.05
R	500	200	9.70 ± 0.06	3.73 ± 0.06	3.30 ± 0.05	5.3 ± 0.1	1.59 ± 0.00	13/7.32 ± 0.29
C	500	200	9.14 ± 0.08	4.33 ± 0.31	3.78 ± 0.08	5.0 ± 0.1	1.32 ± 0.01	13/7.23 ± 0.10
CA	500	200	8.47 ± 0.02	1.03 ± 0.21	3.01 ± 0.05	4.5 ± 0.1	1.48 ± 0.01	13/8.41 ± 0.15
R	1000	200	9.76 ± 0.10	4.27 ± 0.64	3.33 ± 0.09	5.1 ± 0.1	1.54 ± 0.03	14/8.37 ± 0.23
C	1000	200	9.24 ± 0.05	4.67 ± 0.58	3.89 ± 0.05	5.2 ± 0.1	1.33 ± 0.01	14/7.95 ± 0.24
CA	1000	200	8.42 ± 0.05	1.00 ± 0.26	3.03 ± 0.05	4.8 ± 0.1	1.57 ± 0.03	14/9.09 ± 0.02

Values are averages ± standard deviations.

N = Nitrogen added as KNO₃; P = Phosphorus added as Na₂HPO₄; DOC = Dissolved Organic Carbon; UV₂₅₄ = Ultraviolet Absorbance at 254 nm; SUVA = Specific UV₂₅₄ (UV₂₅₄/DOC); FC = free chlorine; FC-7d = free chlorine residual after 7-day hold time; C = Chlorine dioxide dosed at 2 mg L⁻¹ as Cl₂; CA = Chlorine dioxide dosed at 2 mg L⁻¹ as Cl₂ and Alum coagulation at 80 mg L⁻¹ as alum; R = nutrient amended raw water.

Note: Free chlorine was dosed after all other reported measurements.

Table 4 – Excitation and Emission maxima of fluorescence-PARAFAC components.

Component	Excitation Maxima (nm)	Emission Maxima (nm)	r^2 (TCMFP: F_{MAX})
C1	235 (325, 386)	422 (476)	0.55
C2	337 (237)	375 (423)	0.61
C3	267 (367)	456	0.52
C4	226 (280)	355	0.18
C5	400 (370, 309)	490 (394)	0.47

Values in parentheses are secondary and tertiary maxima; r^2 values describe the linear correlations between trichloromethane formation potential (TCMFP) and the fluorescence maximum values (F_{MAX}) for each parallel factor (PARAFAC) component




Figure 1 – Chlorophyll-*a* (Chl-*a*) of the raw water samples as a function of the phosphorus (P) amendment gradient with constant nitrogen ($2,000 \mu\text{g L}^{-1}$) on a log-log basis. Lines represent the least squares best fit.




Figure 2 – Chlorophyll-*a* (Chl-*a*) of the raw water samples as a function of the nitrogen (N) amendment gradient with constant phosphorus ($200 \mu\text{g L}^{-1}$) on a semi-log basis. Lines represent the least squares best fit.




Figure 3 – Chlorophyll-*a* (Chl-*a*) of the raw water samples as a function of the molar N:P ratio of all samples on a log-log basis. Lines represent triplicate averages for the May 30 and August 19 sample collection. See Table 1 for details on N:P ratio.




Figure 4 – Trichloromethane formation potential (TCMFP) as a function of nitrogen (N) amendment for April 5 raw water. The P dose for all N-amended samples was $200 \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.




Figure 5 – Trichloromethane formation potential (TCMFP) as a function phosphorus (P) amendment for April 5 raw water. The N dose for all P-amended samples was 2,000 µg L⁻¹. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.




Figure 6 – Trichloromethane formation potential (TCMFP) as a function of nitrogen (N) amendment for May 30 raw and treated waters (ClO_2 dose of 1 mg L^{-1} as Cl_2 and alum dose of 40 mg L^{-1}). The P dose for all N-amended samples was $200 \text{ } \mu\text{g L}^{-1}$. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.




Figure 7 – Trichloromethane formation potential (TCMFP) as a function of phosphorus (P) amendment for May 30 raw and treated waters (ClO₂ dose of 1 mg L⁻¹ as Cl₂ and alum dose of 40 mg L⁻¹). The N dose for all P-amended samples was 2,000 µg L⁻¹. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.




Figure 8 – Trichloromethane formation potential (TCMFP) as a function of nitrogen (N) amendment for August 19 raw and treated waters (ClO₂ dose of 2 mg L⁻¹ as Cl₂ and alum dose of 80 mg L⁻¹). The P dose for all N-amended samples was 200 µg L⁻¹. Lines represent triplicate averages; filled markers represent blank samples without any nutrient amendment.




Figure 9 – Trichloromethane formation potential (TCMFP) as a function of phosphorus (P) amendment for August 19 raw and treated waters (ClO₂ dose of 2 mg L⁻¹ as Cl₂ and alum dose of 80 mg L⁻¹). The N dose for all P-amended samples was 2,000 µg L⁻¹. Lines represent triplicate averages except for the P = 100 µg L⁻¹ dose, which was excluded. Filled markers represent blank samples without any nutrient amendment.




Figure 10 – Trichloromethane formation potential (TCMFP) for the raw water samples amended with nitrogen (N) and phosphorus (P) for the April 5, May 30, and August 19 samples as a function of the log-molar N:P ratio, where N and P represent the applied doses. Lines represent triplicate averages for each sample collection. See Table 1 for details on N:P ratio.




Figure 11 – Trichloromethane formation potential (TCMFP) for the raw water samples amended with nitrogen (N) and phosphorus (P) for the April 5, May 30, and August 19 samples as a function of chlorophyll-a (Chl-a). Lines represent the least squares best fit.




Figure 12 – Correlations between trichloromethane formation potential (TCMFP) and DOC. Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. DOC is the dissolved organic carbon. Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBFP section.




Figure 13 – Correlations between trichloromethane formation potential (TCMFP) and UV₂₅₄. Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. UV₂₅₄ is the ultraviolet absorbance at 254 nm. Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.




Figure 14 – Correlations between trichloromethane formation potential (TCMFP) and $I_{344/425}$. Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. $I_{344/425}$ is the fluorescence intensity at an excitation of 344 nm and an emission of 425 nm. Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.




Figure 15 – Correlations between trichloromethane formation potential (TCMFP) and C2 F_{MAX} . Linear best-fit models (solid lines) were determined based on least-squares analyses of raw (R), chlorine dioxide treated (C), and chlorine dioxide treated and alum coagulated (CA) waters from the April 5, May 30, and August 19 sampling collections. Dashed lines encompass the upper and lower 95% prediction intervals for the linear models. C2 F_{MAX} is the maximum fluorescence intensity for PARAFAC Component 2 (see Table 4 for description of the fluorescence-PARAFAC components). Seven samples (out of 244) were excluded from this figure because they were determined to be outliers as described in the Results and Discussion – DBPFP section.

References

- Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N. and Smith, V. H., 1998. Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Applications* 8 (3), 559-568.
- Cheng, W. P. and Chi, F. H., 2003. Influence of eutrophication on the coagulation efficiency in reservoir water. *Chemosphere* 53 (7), 773-778.
- Downing, J. A. and McCauley, E., 1992. The nitrogen-phosphorus relationship in lakes. *Limnology and Oceanography* 37 (5), 936-945.
- Eaton, A. D., Clesceri, L. S., Rice, E. W. and Greenberg, A. E., Eds. (2005). Standard methods for the examination of water and wastewater. Washington, D.C., APHA-AWWA-WEF.
- Edzwald, J. K., Becker, W. C. and Wattier, K. L., 1985. Surrogate Parameters for Monitoring Organic-Matter and THM Precursors. *Journal American Water Works Association* 77 (4), 122-132.
- Edzwald, J. K. and Tobiasson, J. E., 1999. Enhanced coagulation: US requirements and a broader view. *Water Science and Technology* 40 (9), 63-70.
- Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H., Ngai, J. T., Seabloom, E. W., Shurin, J. B. and Smith, J. E., 2007. Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters* 10 (12), 1135-1142.
- Fang, J. Y., Yang, X., Ma, J., Shang, C. and Zhao, Q. A., 2010. Characterization of algal organic matter and formation of DBPs from chlor(am)ination. *Water Research* 44 (20), 5897-5906.
- Graham, N. J. D., Wardlaw, V. E., Perry, R. and Jiang, J. Q., 1998. The significance of algae as trihalomethane precursors. *Water Science and Technology* 37 (2), 83-89.
- Granderson, C. W., Pifer, A. D. and Fairey, J. L., 2013. An improved chloroform surrogate for chlorine dioxide- and alum-treated waters. *Journal American Water Works Association* 105 (3), 45-46.
- Hoehn, R. C., Barnes, D. B., Thompson, B. C., Randall, C. W., Grizzard, T. J. and Shaffer, P. T. B., 1980. Algae as sources of trihalomethane precursors. *Journal American Water Works Association* 72 (6), 344-350.

- Huisman, J., Matthijs, H. C. P. and Visser, P. M., Eds. (2005). Harmful Cyanobacteria. The Netherlands, Springer.
- Jack, J., Sellers, T. and Bukaveckas, P. A., 2002. Algal production and trihalomethane formation potential: an experimental assessment and inter-river comparison. *Canadian Journal of Fisheries and Aquatic Sciences* 59 (9), 1482-1491.
- Karanfil, T., Erdogan, I. and Schlautman, M. A., 2003. Selecting filter membranes for measuring DOC and UV254. *Journal American Water Works Association* 95 (3), 86-100.
- Korn, C., Andrews, R. C. and Escobar, M. D., 2002. Development of chlorine dioxide-related by-product models for drinking water treatment. *Water Research* 36 (1), 330-342.
- Krasner, S. W. and Amy, G., 1995. Jar-test evaluations of enhanced coagulation. *Journal American Water Works Association* 87 (10), 93-107.
- Li, L., Gao, N. Y., Deng, Y., Yao, J. J. and Zhang, K. J., 2012. Characterization of intracellular & extracellular algae organic matters (AOM) of *Microcystis aeruginosa* and formation of AOM-associated disinfection byproducts and odor & taste compounds. *Water Research* 46 (4), 1233-1240.
- Miltner, R. J., 1976. *The Effect of Chlorine Dioxide on Trihalomethanes in Drinking Water*, edition, University of Cincinnati.
- Myklestad, S. M., 1995. Release of extracellular products by phytoplankton with special emphasis on polysaccharides. *Science of the Total Environment* 165 (1-3), 155-164.
- Pifer, A. D., Cousins, S. L. and Fairey, J. L., in press. Assessing UV- and fluorescence-based metrics as disinfection byproduct precursor surrogate parameters in a water body influenced by a heavy rainfall event. *Journal of Water Supply: Research and Technology - AQUA*.
- Pifer, A. D. and Fairey, J. L., 2012. Improving on SUVA₂₅₄ using fluorescence-PARAFAC analysis and asymmetric flow-field flow fractionation for assessing disinfection byproduct formation and control. *Water Research* 46 (9), 2927-2936.
- Pifer, A. D., Miskin, D. R., Cousins, S. L. and Fairey, J. L., 2011. Coupling asymmetric flow-field flow fractionation and fluorescence parallel factor analysis reveals stratification of dissolved organic matter in a drinking water reservoir. *Journal of Chromatography A* 1218 (27), 4167-4178.

- Plummer, J. D. and Edzwald, J. K., 2002. Effects of chlorine and ozone on algal cell properties and removal of algae by coagulation. *Journal of Water Supply Research and Technology-Aqua* 51 (6), 307-318.
- Richardson, S. D., Plewa, M. J., Wagner, E. D., Schoeny, R. and DeMarini, D. M., 2007. Occurrence, genotoxicity, and carcinogenicity of regulated and emerging disinfection by-products in drinking water: A review and roadmap for research. *Mutation Research-Reviews in Mutation Research* 636 (1-3), 178-242.
- Rook, J. J., 1976. Haloforms in drinking water. *Journal American Water Works Association* 68 (3), 168-172.
- Schindler, D. W., 1974. Eutrophication and recovery in Experimental Lakes - Implications for lake management. *Science* 184 (4139), 897-899.
- Scott, J. T., Doyle, R. D., Prochnow, S. J. and White, J. D., 2008. Are watershed and lacustrine controls on planktonic N-2 fixation hierarchically structured? *Ecological Applications* 18 (3), 805-819.
- Scott, J. T. and Grantz, E. M., 2013. N-2 fixation exceeds internal nitrogen loading as a phytoplankton nutrient source in perpetually nitrogen-limited reservoirs. *Freshwater Science* 32 (3), 849-861.
- Sen, S., Haggard, B. E., Chaubey, I., Brye, K. R., Costello, T. A. and Matlock, M. D., 2007. Sediment phosphorus release at Beaver Reservoir, Northwest Arkansas, USA, 2002-2003: A preliminary investigation. *Water Air and Soil Pollution* 179 (1-4), 67-77.
- Smith, V. H., 1983. Low nitrogen to phosphorus ratios favor dominance by blue-green algae in lake phytoplankton. *Science* 221 (4611), 669-671.
- Smith, V. H., Tilman, G. D. and Nekola, J. C., 1999. Eutrophication: impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* 100 (1-3), 179-196.
- Stedmon, C. A. and Bro, R., 2008. Characterizing dissolved organic matter fluorescence with parallel factor analysis: a tutorial. *Limnology and Oceanography-Methods* 6, 572-579.
- Weishaar, J. L., Aiken, G. R., Bergamaschi, B. A., Fram, M. S., Fujii, R. and Mopper, K., 2003. Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Environmental Science & Technology* 37 (20), 4702-4708.

Zepp, R. G., Sheldon, W. M. and Moran, M. A., 2004. Dissolved organic fluorophores in southeastern US coastal waters: correction method for eliminating Rayleigh and Raman scattering peaks in excitation-emission matrices. *Marine Chemistry* 89 (1-4), 15-36.