

8-28-2012

Alkynal involved organocatalytic reactions

Xinshuai Zhang

Follow this and additional works at: https://digitalrepository.unm.edu/chem_etds

Recommended Citation

Zhang, Xinshuai. "Alkynal involved organocatalytic reactions." (2012). https://digitalrepository.unm.edu/chem_etds/27

This Dissertation is brought to you for free and open access by the Electronic Theses and Dissertations at UNM Digital Repository. It has been accepted for inclusion in Chemistry ETDs by an authorized administrator of UNM Digital Repository. For more information, please contact disc@unm.edu.

Xinshuai Zhang

Candidate

Chemistry and Chemical Biology

Department

This dissertation is approved, and it is acceptable in quality and form for publication:

Approved by the Dissertation Committee:

Wei Wang , Chairperson

Patrick S. Mariano

Hua Guo

Changjian Feng

ALKYNAL INVOLVED ORGANOCATALYTIC REACTIONS

by

XINSHUAI ZHANG

B.S., Chemistry, Shan'xi Normal University, P.R.China, 2001

M.S., Chemistry, Suzhou University, P.R. China, 2004

DISSERTATION

Submitted in Partial Fulfillment of the
Requirements for the Degree of

Doctor of Philosophy

Chemistry

The University of New Mexico
Albuquerque, New Mexico

July, 2012

ACKNOWLEDGMENTS

Foremost, I would like to express my greatest and sincerest gratitude to my research advisor - Professor Wei Wang. He always expressed his insight, creativity in knowledge and patience in guidance. With his sustained encouragement and guidance, I successfully solved the problems encountered in my research. He set a great example of successful research career in chemistry for me. What I learned from him is and will be invaluable treasure of my whole life.

Secondly, I wish to convey my sincere appreciation to all my committee members, Professor Patrick S. Mariano, Professor Hua Guo and Professor Changjian Feng for their precious time in reading and evaluating my dissertation. Especially, I want to express my deepest gratitude to Professor Patrick S. Mariano who gave me immense help in finding post-doctoral position.

I also would like to extend my thanks for current and ex-members in Professor Wei Wang's research group. They are my colleagues in lab and nice friends in life. I learn a lot from discussion research projects with them. My appreciation covered Dr. Shilei Zhang, Dr. Hexin Xie, Dr. Wei Jiang, Dr. Yinan Zhang, Mr. Xixi Song, Mr. Sihan Zhao, Ms. Xiaobei Chen, Ms. Yanting Cao, Mr. Chunliang Liu, Mr. Weimin Xuan, Mr. Aiguo Song, Mr. Wenhan He, Mr. Yuan Liu, Ms. Xichen Li. Among them, I want to point out my special thanks to Dr. Shilei Zhang. He helped me a lot in my research work

and shared his own research experience with me. I benefit a lot from his diligence and experience.

Finally, my conserved gratitude is left to my husband, Hua Huang who always strongly supports me during my Ph.D period. Without him, the life here will become gloomy and the research work should be even tough. Also, my special thanks go to my family: my parents and sister. Without their support, love and encouragement, none of these would be possible.

ALKYNAL INVOLVED ORGANOCATALYTIC REACTIONS

By

XINSHUAI ZHANG

B.S., Chemistry, Shan'xi Normal University, P.R.China, 2001

M.S., Chemistry, Suzhou University, P.R.China, 2004

Ph.D., Chemistry, University of New Mexico, 2012

ABSTRACT

During the past twenty years, vast progress has been achieved in the field of organocatalysis. One of the fruitful areas is iminium catalysis using α , β -unsaturated aldehydes as essential reactants. Enals involved enantioselective reactions (including single-step reactions and cascade/tandem reactions) represent an enormous and fruitful research branch. In contrast, the protocols for the corresponding alkynals, even in single-step conjugate addition reactions, are extremely rare.

Toward this end, my Ph. D research work mainly focuses on the development of alkynal involved organocatalytic reactions and expands the scope of organocatalysis by uncovering new activation modes. Unlike enals, ynals with sp hybridized C≡C bonds are much less studied and the related chemistries are much less understood. Within the context, firstly, we have developed an organocatalytic regio- and stereoselective Michael additions of 1*H*-1, 2, 3-triazole to alkynals which afford the trisubstituted alkenes. It is found that the reaction conditions are critical for the 1- vs 2-isomers. Furthermore, we have discovered several novel asymmetric oxa-Michael-Michael/aldol and aza-Michael-aldol(-aromatization) reaction sequences initiated by the Michael addition to alkynals. Notably, an unprecedented chiral iminium-allenamine cascade is disclosed for the first time. Significantly, these processes serve as efficient and straightforward entries to biologically significant chiral 4*H*-chromenes, polysubstituted quinolines and chiral 1,4-dihydroquinolines with excellent yield and enantioselectivity, as well as broad substrate compatibility. The proposed iminium-allenamine activation mode adds a new domain in the field of organocatalytic enantioselective cascade reactions. Finally, unprecedented organocatalytic 1,6-conjugated addition to ynals, an unmet challenging problem in organic synthesis, also has been realized.

TABLE OF CONTENTS

List of Schemes.....	X
List of Figures.....	xiii
List of Tables.....	xiv
List of Abbreviations.....	xvi
Chapter 1 Development of Organocatalysis	1
1.1 Introduction	1
1.2 Classification of Organocatalysts	3
1.2.1 Covalent Organocatalysts	4
1.2.1.1 Enamine Catalysts.....	4
1.2.1.2 Iminium Catalysts	8
1.2.1.3 SOMO Catalysis	13
1.2.1.4 Dienamine Catalysis	14
1.2.1.5 Oxidative Enamine Catalysis	16
1.2.2 Non-Covalent Organocatalysts	18
1.3 Summary and Outlook	23
1.4 Reference.....	24
Chapter 2 Organocatalytic Regio- and Stereoselective Michael Addition of 1<i>H</i>-1, 2, 3-Triazole to Alkynals.....	33
2.1 Background and Significance.....	33
2.2 Research Design.....	35
2.3 Introduction	35
2.4 Results and Discussion.....	36
2.5 Conclusions	44
2.6 Supporting Information.....	45
2.7 Reference.....	61
Chapter 3 Alkynal Involved Organocatalytic Cascade Reactions.....	64

3.1 Background and Significance.....	64
3.2 Research Design.....	67
3.3 Iminium–Allenamine Cascade Catalysis: One-Pot Access to Chiral 4 <i>H</i> -Chromenes by a Highly Enantioselective Michael–Michael Sequence ¹⁴	69
3.3.1 Introduction	69
3.3.2 Results and Discussion	70
3.3.3 Conclusion	75
3.3.4 Supporting Information	75
3.4 Development of Organocatalytic Asymmetric Cascade Double Michael Addition Reactions of 2-(2, 2-Bis-benzenesulfonyl-vinyl)-phenol to Alkynals	84
3.4.1 Introduction	84
3.4.2 Results and Discussion	86
3.4.3 Conclusion	91
3.4.4 Supporting Information	92
3.5 “One-Pot” Access to 4 <i>H</i> -Chromenes with Formation of a Chiral Quaternary Stereogenic Center by a Highly Enantioselective Iminium-allenamine Involved Oxa- Michael Aldol Cascade ²⁹	103
3.5.1 Introduction	103
3.5.2 Results and Discussion	106
3.5.3 Conclusion	111
3.5.4 Supporting Information	112
3.6 Efficient Synthesis of Polysubstituted Quinolines and Chiral 1,4-Dihydroquinolines by a Divergent Organocatalytic Cascade Approach. An Unexpected Effect of “ <i>N</i> ” Protecting Groups on Formed Products	127
3.6.1 Introduction	127
3.6.2 Results and Discussion	129
3.6.2.1 An Unexpected Organocatalytic aza-Michael-Aldol-Aromization Cascade	129
3.6.2.2 Organocatalytic Enantioselective Michael-aldol Cascade Reactions	137
3.6.3 Conclusion	143
3.6.4 Supporting Information	144

3.7 Reference.....	170
Chapter 4 Organocatalytic 1,6-Addition Reactions.....	184
4.1 Background and Significance.....	184
4.2 Research Design.....	186
4.3 Results and Discussion.....	187
4.4 Conclusions.....	195
4.5 Supporting Information.....	195
4.6 Reference.....	203

List of Schemes

Scheme 1.1 Pracejus' Enantioselective Ester Synthesis Catalyzed by <i>O</i> -acetylquinine....	2
Scheme 1.2 Proline Catalyzed Intramolecular Asymmetric Aldol Reaction (Hajos-Parrish-Eder-Sauer-Wiechert Reaction)	5
Scheme 1.3 Proline Catalyzed Intermolecular Aldol Reaction	5
Scheme 1.4 Other Examples of Enantioselective Enamine Catalysis	6
Scheme 1.5 Imidazolidinone Catalyzed Diels–Alder Reaction	9
Scheme 1.6 Other Examples of Enantioselective Iminium Catalysis	9
Scheme 1.7 Primary Amine Based Catalyst in Enamine / Iminium Catalysis	12
Scheme 1.8 Addition Reaction between Enamine Cation Radical and Olefine.....	14
Scheme 1.9 Representative SOMO Catalysis in Enantioselective α -Allylation of Aldehyde	14
Scheme 1.10 Formation of the Dienamine Intermediate in the Reaction between 2-Pentenal and the Chiral Amine Catalyst	15
Scheme 1.11 Organocatalytic γ -Amination of α , β -Unsaturated Aldehyde	15
Scheme 1.12 Other Examples of Dienamine Catalytic Transformations.....	16
Scheme 1.13 Oxidative Enamine Catalysis in Multiple Cascade Reactions.....	18
Scheme 1.14 Thiourea-Catalyzed Asymmetric Strecker Reaction	19
Scheme 1.15 Thiourea-Catalyzed Asymmetric Mannich Reaction	19
Scheme 1.16 Other Examples of Thiourea Catalyst as Double H-Bonding Catalyst	20
Scheme 1.17 Bicyclic Guanidine Catalyst for Asymmetric Strecker Reaction	21

Scheme 1.18 Michael Additions of Malonates to Nitroolefins Catalyzed by Takemoto Thiourea Catalyst I	22
Scheme 1.19 Michael Addition of Nitromethane to Chalcones Catalyzed by Cinchona Alkaloid Derived Bifunctional Thioureas Catalyst II	22
Scheme 1.20 Catalyst III Promoted Asymmetric Morita-Baylis-Hillman Reaction	23
Scheme 2.1 Examples of Catalytic Asymmetric Michael Additions to Alkynones or Acetylenic Esters	34
Scheme 2.2 Examples of Acetylenic Dienophiles in Asymmetric Diels-Alder Reaction	34
Scheme 3.1 the Iminium-Enamine Combination	65
Scheme 3.2 Orgacatalytic Michael-aldol Cascade for the Assembly of Optically Active Cyclohexanones	65
Scheme 3.3 Orgacatalytic Michael-Michael-aldol Triple Cascade Reaction.....	66
Scheme 3.4 Selected Examples of Michael Additions to Vinyl Sulfones.....	85
Scheme 3.5 Synthetic Utility of Adduct 5	91
Scheme 3.6 Alkynals Participating in Enantioselective Cascade Oxa-Michael-Michael and Aldol Reactions Promoted by a Chiral Secondary Amine Involving a Novel Iminium-allenamine Activation Mode.....	105
Scheme 3.7 Organocatalytic Michael-aldol Cascade Reactions Involving Chiral Allenamines	130
Scheme 3.8 Organocatalytic aza-Michael-aldol-aromatization Reaction	134
Scheme 4.1 Examples of Organocatalytic 1,6-Addition	185
Scheme 4.2 Proposal for Organocatalytic 1,6-Addition.....	187

Scheme 4.3 Organocatalytic 1,6-Addition between (<i>E</i>)-5-Phenylpent-2-en-4-ynal 1a and Thiophenol	188
Scheme 4.4 Proposed Mechanism for 1,6-Addition of Thiophenol to (<i>E</i>)-5-Phenylpent-2-en-4-ynal 1a	189
Scheme 4.5 Verification of the Structure of 1,6-Addition Product 2a	190
Scheme 4.6 1,6-Addition of Nitrogen Nucleophiles to (<i>E</i>)-5-Phenylpent-2-en-4-ynal 1a	193

List of Figures

Figure 1.1 Enamine Catalytic Cycle	8
Figure 1.2 Iminium Catalytic Cycle for Michael Addition and Cycloaddition	11
Figure 1.3 SOMO Catalysis via Single-Electron Oxidation of a Transiently Formed Enamine	14
Figure 1.4 Aminocatalysis: (1) Iminium Catalysis. (2) Oxidative Enamine Catalysis....	17
Figure 2.1 Secondary Amine Catalysts Screened	38
Figure 2.2 Other Secondary Amine Catalysts Screened	40
Figure 2.3 Tautomerism of Neutral 1 <i>H</i> -1, 2, 3-Triazole.....	42
Figure 2.4 Tautomerism of Triazolium Ion	42
Figure 3.1 Conventional Iminium-Enamine (1) and Novel Iminium-Allenamine (2) Activation Strategies in Organocatalytic Enantioselective Cascade Reactions.....	68
Figure 3.2 Selected Examples of 4 <i>H</i> -Chromenes as Substructures in Biologically Interesting Natural Products	70
Figure 3.3 Secondary Amine Catalysts Screened	71
Figure 3.4 the Single-Crystal X-ray Structure of Compound 3g , with Ellipsoids Set at 20% Probability.....	74
Figure 3.5 Secondary Amine Catalysts Screened	87
Figure 3.6 the Single-Crystal X-ray Structure of Compound 5a	90
Figure 3.7 X-ray Structure of 7h	111
Figure 3.8 Skeletons of Quinolines A and Hydroquinolines B-D	128
Figure 3.9 X-ray Crystal Structure of Compound 12g	143

List of Tables

Table 2.1 Optimization Reaction Conditions for the Organocatalytic Regio- and Stereocontrolled Michael Addition of 1a and 1 <i>H</i> -1, 2, 3-Triazole to Afford 2a^a	38
Table 2.2 Optimization Reaction Conditions for the Organocatalytic Regio- and Stereocontrolled Michael Addition of 1a and 1 <i>H</i> -1, 2, 3-Triazole to Afford 3a^a	41
Table 2.3 Scope of the Organocatalytic Regio- and Stereoselective Michael Addition of 1 <i>H</i> -1, 2, 3-Triazole to Substituted Phenyl-propynal 1a-m to Afford 2^a and 3^b Respectively.....	43
Table 3.1 Optimization Reaction Conditions for the Organocatalytic Asymmetric Cascade Oxa-Michael-Michael Reaction between Phenyl-propynal 1a and 2-(<i>E</i>)-(2-Nitrovinyl)-phenol 2a^a	71
Table 3.2 Catalyst III Promoted Asymmetric Cascade Oxa-Michael-Michael Reactions of Propynal 1 with 2-(<i>E</i>)-(2-Nitrovinyl)-phenol 2^a	73
Table 3.3 Optimization Reaction Conditions for the Organocatalytic Asymmetric Cascade Michael-Michael Reaction between Phenyl-propynal 1a and 2-(2, 2-Bis-benzenesulfonyl-vinyl)-phenol 4a^a	87
Table 3.4 Catalyst III Promoted Asymmetric Cascade Michael-Michael Reactions of Propynal 1 with 2-(2, 2-Bis-benzenesulfonyl-vinyl)-phenol 4^a	89
Table 3.5 Development and Optimization of Reaction Conditions of an Organocatalytic Enantioselective Cascade Oxa-Michael-aldol Cascade Reaction ^a	108
Table 3.6 Catalyst III Promoted Enantioselective Cascade Oxa-Michael-aldol Reactions of Alkynals 1 with Ethyl 2-(2-Hydroxyphenyl)-2-oxoacetates 6^a	110

Table 3.7 Exploration of Organocatalytic Aza-Michael-aldol-aromatization Cascade Reaction ^a	131
Table 3.8 Substrate Scope of I -Catalyzed “one-pot” Synthesis of Quinolines 10^a	136
Table 3.9 Optimization of Reaction Conditions for Organocatalytic Enantioselective Aza-Michael-aldol Cascade Reactions ^a	139
Table 3.10 Enantioselective Organocatalytic Cascade Aza-Michael-aldol Reactions between Ynals 1 and 2'-(Trifluoromethanesulfonyl)aminoketones 11^a	141
Table 4.1 Optimization of Reaction Condition of 1,6-Addition between (<i>E</i>)-5-Phenylpent-2-en-4-ynal 1a and Phenol ^a	191
Table 4.2 Scope of 1,6-Addition Process ^a	193

List of Abbreviations

$[\alpha]_D$	Specific Rotation at Wavelength of Sodium D Line
Ac	Acetyl Acetate
aq.	Aqueous
Ar	Aromatic
Bn	Benzyl
Boc	<i>t</i> -Butoxycarbonyl
bp	Boiling Point
<i>t</i> -Bu	<i>t</i> -Butyl
Bz	Benzoyl
<i>c</i>	Concentration
Cbz	Benzyloxycarbonyl
CDCl ₃	Deuterated Chloroform
δ	Chemical Shift
DCE	Dichloroethane
DCM	Dichloromethane
DMF	<i>N,N</i> -Dimethylformamide
DMSO	Dimethyl Sulfoxide
D ₂ O	Deuterated Water
EDG	Electron Donating Group

ee	Enantiomeric Excess
equiv.	equivalent
EtOAc	Ethyl Acetate
EWG	Electron Withdrawing Group
g	Gram
h	Hour
HOMO	Highest Occupied Molecular Orbital
HPLC	High Performance Liquid Chromatography
HRMS	High Resolution Mass Spectra
Hz	Hertz
<i>J</i>	Coupling Constant
λ	Wavelength
LUMO	Lowest Unoccupied Molecular Orbital
<i>m</i> -	Meta-
Me	Methyl
MeCN	Acetonitrile
Mg	Magnesium
mg	Milligram
min.	Minute
mL	Millimole

mp	Melting Point
Ms-	Methanesulfonyl
NMR	Nuclear Magnetic Resonance
<i>o</i> -	Ortho-
<i>p</i> -	Para-
Ph	Phenyl
PG	Protecting Group
ppm	Parts per Million
<i>i</i> -PrOH	Isopropyl Alcohol
rt	Room Temperature
TB(DM)S	<i>tert</i> -Butyldimethylsilyl
TBME	<i>tert</i> -Butylmethylether
TEA	Triethylamine
TES	Triethylsilyl
Tf	Trifluoromethanesulfonyl
TFA	Trifluoroacetic Acid
THF	Tetrahydrofuran
TLC	Thin-Layer Chromatography
TMS	Trimethylsilyl
Ts	<i>p</i> -Toluenesulfonyl

μL

Microliter

Chapter 1

Development of Organocatalysis


1.1 Introduction

We live in a world that new chemicals, materials and therapeutics come from synthesis, while many chemical processes heavily rely on catalysis. As reported, 90% of all commercially produced chemical products involve catalysts at some stages in the process of their manufacture¹. Particularly, the application of catalysis to the pharmaceutical industry has steadily increased over the past two decades due to the facts that 80% of small-molecule drugs approved by the FDA are chiral and syntheses using conventional asymmetric methods cannot meet the demand². Therefore, the development of new synthesis strategies becomes essential. Notably organocatalysis using pure small organic molecules has been an important player over the past 20 years in organic synthesis³.

Organocatalysis has a rich history. The first reported asymmetric organocatalytic reaction dated as early as 1912⁴. An enantioselective catalytic addition of HCN to benzaldehyde in the presence of alkaloids quinine and quinidine was realized although the optical activities of the resulting cyanohydrins were less than 10%. Subsequently, in 1960, Pracejus reported that the addition of methanol to phenylmethylketene using *O*-acetylquinine as catalyst again can get (-)- α -phenyl methylpropionate with synthetically

useful levels of enantioselectivity (74% ee)⁵ (Scheme 1.1). Further breakthroughs were achieved between 1970s and 1980s. The most famous example is the discovery of the L-proline promoted asymmetric Robinson annulation reaction which is also called Hajos-Parrish-Eder-Sauer-Wiechert reaction⁶. However, the catalytic potential of proline was ignored until 2000, the seminal works were published simultaneously by List, Barbas⁷ and MacMillan⁸. Since then, it has grown at an explosive pace over the past decade and the number of publications on the topic of organocatalysis has increased remarkably. Organocatalysis currently seems to be in the state of a “gold rush”⁹.

Scheme 1.1 Pracejus’ Enantioselective Ester Synthesis Catalyzed by *O*-acetylquinine



Compared with traditional approaches including transition metal catalysis and enzymatic transformations, organocatalysts have several advantages^{3b,10}. First, they are relatively inert toward moisture and oxygen, so experimental operations are simple and no demanding reaction conditions such as inert atmosphere, dry solvents are required in most cases. Second, they are relatively non-toxic and environmentally benign because of the absence of transition metals. This is particularly attractive for the preparation of pharmaceutical products, which do not tolerate toxic metal contamination. The increasing

focuses on the development of environmentally sustainable manufacturing processes also make organocatalysis as one of the most appealing research subjects. Third, organocatalysts are readily available and inexpensive because most of their precursors are directly obtained from naturally occurring optically pure molecules, such as amino acids and alkaloids. Finally, enzymes are usually substrate and reaction specific; however, one kind of organocatalyst usually can catalyze a variety of reactions with broad range of substrate scopes. Although the impact of transition metal catalysis on synthetic chemistry cannot be understated, the advent of organocatalysis indeed has brought the prospect of a complementary mode of catalysis^{3b}.

1.2 Classification of Organocatalysts

The field of organocatalysis has been a rapidly growing research area which evolved from its infancy through the “gold rush” stage over the last decade¹¹. Not only have the scopes and applications of enantioselective organocatalytic reactions been expanded dramatically, but also the fundamental mechanisms and catalytic modes underlying each type of organocatalytic reactions have been deeply comprehended. Obviously, it is relatively straightforward and reliable to apply the established activation modes as a platform to rationally design new enantioselective organocatalytic reactions and discover novel activation modes^{3b}. Encouraged by the significant value of catalytic mechanisms, herein, I will briefly summarize them in two categories.

Generally speaking, based on the typical interactions between an organocatalyst and substrates within the catalytic cycle, the vast majority of organocatalytic transformations can be divided into the two subgroups “covalent organocatalysis” and “non-covalent organocatalysis”. In the former case, covalent adducts are formed between the substrates and the catalyst. Whilst, in the latter case, non-covalent interactions such as hydrogen bonding or the formation of ion pairs play a part in the catalytic process⁹. It is worthy of point out that a bunch of combined catalytic strategies have emerged as a new branch in the field of organocatalyst very recently.

1.2.1 Covalent Organocatalysts


Aminocatalysts are probably the most commonly used organocatalysts in asymmetric transformations¹². Their properties are closely related with the particular reactivity of the nucleophilic nitrogen, whereby the nitrogen atom condenses with a substrate which is usually a carbonyl compound and give rise either to enamine or iminium intermediates. Besides that, SOMO (Singly Occupied Molecular Orbital) catalysis and dienamine catalysis are also fully investigated very recently. As to other types of covalent catalysts which are not closely related with my Ph. D research work such as planar-chiral heterocycles as nucleophilic catalysis, *N*-heterocyclic carbene catalyst will not be discussed in my dissertation.

1.2.1.1 Enamine Catalysts


In 1971, the proline-catalyzed intramolecular aldol reaction in the synthesis of the Wieland–Miescher ketone was reported simultaneously by Zoltan Hajos, David Parrish and Rudolf Weichert, Gerhard Sauer, Ulrich Eder (Scheme 1.2)⁶. The wonderful reaction

scenario was recognized by the synthetic community; however, the underlying catalytic mechanistic detail was not explored until more than 30 years later. In 2000, the great potential of activation mode – enamine catalysis was realized by the synthetic community along with the seminal work of intermolecular aldol reaction catalyzed by secondary amine via enamine by List and Barbars research group (Scheme 1.3)⁷. Since then, a number of new types of enamine catalysts¹³ have been identified and a broad range of asymmetric chemical transformations^{13, 14} with aldehydes and ketones via enamine catalysis have been achieved such as Mannich reactions¹⁵⁻¹⁷, Michael addition reactions¹⁸, α -functionalization of carbonyl compounds with a great many of electrophiles, namely, α -amination¹⁹⁻²⁰, α -oxygenation²¹, α -halogenation²²⁻²⁴, α -sulfenylation²⁵, α -selenylation²⁶⁻²⁷ and intramolecular α -alkylation of aldehydes²⁸ (Scheme 1.4).


Scheme 1.2 Proline Catalyzed Intramolecular Asymmetric Aldol Reaction (Hajos-Parrish-Eder-Sauer-Wiechert Reaction)




Scheme 1.3 Proline Catalyzed Intermolecular Aldol Reaction




Scheme 1.4 Other Examples of Enantioselective Enamine Catalysis





In the enamine catalytic cycle, the LUMO lowering effect results in dramatic increase in C-H acidity of the initial iminium ion **I** (Figure 1.1). Then the active enamine **II** species is reversibly generated. The resulting enamine **II** reacts with the electrophile via nucleophilic substitution or nucleophilic addition to form iminium ion **III** which is then hydrolyzed to afford the desired product¹³. In short, the essence of enamine catalysis is generation of the more nucleophilic enamine species whose HOMO energy is increased.

Figure 1.1 Enamine Catalytic Cycle




1.2.1.2 Iminium Catalysts


Although the iminium-catalysis strategy had never been documented until 2000, the experimental basis for this organocatalytic concept probably can be stretched back to several well-established methodological investigations. One of the well-known examples is the Knoevenagel condensation²⁹ mediated by primary or secondary amines. In 1931, Blanchard et al. suggested that an iminium ion species reversibly generated by an aldehyde and an amine is involved in the process³⁰. In 1953, Crowell and co-worker presented the kinetic evidence for imine/iminium intermediates in the Knoevenagel condensations³¹. The key point for this process is that only the iminium ion is of sufficient electronic deficiency to undergo the following nucleophilic attack which is closely related with LUMO-lowering activation^{11b}.


The iminium catalysis concept was first raised by MacMillan for the enantioselective Diels-Alder reaction between enal dienophiles and dienes in the presence of chiral imidazolidinone in 2000 (Scheme 1.5)³². Thereafter, the potential of the iminium catalysis was further explored. A myriad of transformations related with enals or enones, for example, [3+2] cycloaddition³³, [4+3]-cycloaddition³⁴, Friedel-Crafts alkylation³⁵, Mukaiyama Michael addition³⁶, Michael addition³⁷, hydrogenation³⁸, cyclopropanation³⁹ and epoxidation⁴⁰ have been extensively exploited (Scheme 1.6).

Scheme 1.5 Imidazolidinone Catalyzed Diels–Alder Reaction




Scheme 1.6 Other Examples of Enantioselective Iminium Catalysis



3. 


 $\text{N-Me-pyrrole} + \text{R-CH=CH-CHO} \xrightarrow[\text{THF-H}_2\text{O, -60~-30}^\circ\text{C, TFA}]{\text{Catalyst (20 mol\%)}}$

 $\text{R} = \text{Alkyl, -Aryl, -CO}_2\text{Me}$

 Yield 72-90%
 ee 87-93%
4. 


 $\text{Furan derivative} + \text{R-CH=CH-CHO} \xrightarrow[\text{CH}_2\text{Cl}_2\text{-H}_2\text{O, -70~-20}^\circ\text{C, DNBA}]{\text{Catalyst (20 mol\%)}}$

 $\text{R} = \text{Alkyl, -Aryl, -CO}_2\text{Me}$

 Yield 77-87%
 ee 84-99%
5. 


 $\text{R}^1\text{-CH=CH-CHO} + \text{R}^2\text{O}_2\text{C-CH}_2\text{-CO}_2\text{R}^2 \xrightarrow[\text{EtOH, 0}^\circ\text{C}]{\text{Catalyst (10 mol\%)}}$

 $\text{R}^1 = \text{Aryl}$ $\text{R}^2 = \text{Bn, Me}$

 Ar = 3,5-(CF₃)₂-Ph
 Yield 31-95%
 ee 86-95%
6. 


 $\text{R}^1\text{-CH=CH-CHO} + \text{EtO}_2\text{C-CH=C(Me)-CH}_2\text{-CO}_2\text{Et} \xrightarrow[\text{CHCl}_3, -30}^\circ\text{C, TFA}]{\text{Catalyst (20 mol\%)}}$

 $\text{R}^1, \text{R}^2 = \text{Alkyl, Aryl}$

 Yield 74-95%
 ee 90-97%
7. 

 $\text{R}^1\text{-CH=CH-CHO} + \text{Me}_2\text{S}^+\text{-CH}_2\text{-R}^2 \xrightarrow[\text{CHCl}_3, -10}^\circ\text{C}]{\text{Catalyst (20 mol\%)}}$

 $\text{R}^1 = \text{Alkyl, Ph}$ $\text{R}^2 = \text{COPh, CO}t\text{-Bu}$

 Yield 63-85%
 ee 89-96%
 dr 6:1~72:1
8. 


 $\text{R}^1\text{-CH=CH-CHO} + \text{R}^2\text{O}_2\text{C-CH}_2\text{-CO}_2\text{R}^2 \xrightarrow[\text{H}_2\text{O}_2, \text{CH}_2\text{Cl}_2, \text{r.t., 4h}]{\text{Catalyst (10 mol\%)}}$

 $\text{R}^1 = \text{Alkyl, Aryl}$ $\text{R}^2 = \text{H, Me}$

 Ar = 3,5-(CF₃)₂-Ph
 Yield 60->90%
 ee 75-98%
 dr 9:1~98:2

These reactions all share the same iminium activation mechanism. In the iminium catalytic cycle, the condensation of enals or enones with chiral amines results in the reversible formation of active iminium ions **IV** whose LUMO energy is lowered⁴¹. Accordingly, the iminium ions are more electrophilically subjected to attack by a variety of suitable nucleophiles or electron rich dienes. Subsequently, Michael addition results in an enamine **V** followed by iminium **VI** and cycloaddition results in an iminium ion intermediate **VII**. These intermediates (**VI** and **VII**) are then hydrolysed to give the desired product and regenerate chiral amine catalyst (Figure 1.2).

Figure 1.2 Iminium Catalytic Cycle for Michael Addition and Cycloaddition





In summary, the HOMO-raising enamine activation and the LUMO-lowering iminium activation are two divergent activation modes in organocatalysis despite the fundamental analogies in the structure of these catalysts and substrates. On the other hand, enamine and iminium catalysis are closely interrelated with each other which can be seen from their catalytic cycles. Enamine catalysis always results in the formation of iminium ion

and iminium catalysis proceeds via the enamine intermediate¹². The combinations of enamine and iminium catalysis in a cascade reaction leading to the formation of molecules of higher level complexity constitute a second milestone in the field of amine-based organocatalysis⁴².

Meanwhile, many new types of chiral amine catalysts have been identified. Remarkably, the diarylprolinol silyl ether catalysts have emerged as highly powerful, potentially general organocatalysts in promoting either enamine catalytic cycle or iminium catalytic cycle⁴³. In general, secondary amine is suitable for less hindered carbonyl compounds serving as either enamine or iminium catalysis. However, these catalysts have the inherent difficulties in generating congested covalent intermediates with sterically demanding substrates. Fortunately, the problem was overcome by less hindered primary amine based catalysts which were directly derived from natural cinchona alkaloids⁴⁴. And this strategy has been successfully demonstrated in some cases for activating challenging ketones⁴⁵ and α,β -disubstituted enals⁴⁶ (Scheme 1.7).

Scheme 1.7 Primary Amine Based Catalyst in Enamine / Iminium Catalysis





1.2.1.3 SOMO Catalysis

In 1992, K. Narasaka and co-workers reported the addition reactions of enamine cation radicals generated in the presence of stoichiometric amount of oxidant ammonium cerium (IV) nitrate (CAN) to olefins (Scheme 1.8)⁴⁷. About 15 years later, the MacMillan research group proposed a novel SOMO (single occupied molecular orbital) organocatalysis concept⁴⁸. SOMO catalysis is based on the idea that single-electron oxidation of a transiently produced, electron-rich enamine selectively generates a three π -electrons radical cation with a single occupied molecular orbital (Figure 1.3). This intermediate readily reacts with a variety of weak nucleophilic carbon-based “SOMOphiles” to afford enantioselective α -functionalized carbonyl-containing compounds (Scheme 1.9)⁴⁸. Although SOMO catalysis is one of the most recently discovered activation modes, it has already been extended to many asymmetric transformations⁴⁹⁻⁵¹ which are complementary to those via enamine catalysis.

Scheme 1.8 Addition Reaction between Enamine Cation Radical and Olefine






Figure 1.3 SOMO Catalysis via Single-Electron Oxidation of a Transiently Formed Enamine



Scheme 1.9 Representative SOMO Catalysis in Enantioselective α -Allylation of Aldehyde




1.2.1.4 Dienamine Catalysis


Enamine, iminium and SOMO catalysis have provided us feasible methods for stereoselective functionalization of carbonyl compounds at their α and/or β positions. In 2006, the seminal work about the corresponding γ -functionalization of carbonyl

compounds was reported by Jørgensen and co-workers who launch the unique dienamine organocatalyst⁵² concept for the first time. When they undertook ¹HNMR spectroscopic investigations in the attempt to characterize the expected electrophilic iminium-ion intermediate **A** formed by the reaction of 2-pentenal and chiral amine catalyst, contrary to their expectations, electron-rich dienamine species **B** was observed (Scheme 1.10)⁵². Inspired by the observation, Jørgensen demonstrated the first organocatalytic γ -amination of α,β -unsaturated aldehydes with high enantioselectivity by inverting the normal electrophilic alkene into electron-rich diene (Scheme 1.11)⁵².

Scheme 1.10 Formation of the Dienamine Intermediate in the Reaction between 2-Pentenal and the Chiral Amine Catalyst




Scheme 1.11 Organocatalytic γ -Amination of α,β -Unsaturated Aldehyde



Since the initial studies, the potential of dienamine catalysis was further exploited. Many dienamine catalytic transformations were designed to expand the scope of γ -

activation such as the intramolecular cyclization of unsaturated dialdehydes⁵³, primary amine catalyzed vinylogous Michael addition of cyclic enones to nitroalkenes⁵⁴ (Scheme 1.12). The common character for the substrate is β -aliphatic-substituted γ -enolizable unsaturated carbonyl compounds.

Scheme 1.12 Other Examples of Dienamine Catalytic Transformations




1.2.1.5 Oxidative Enamine Catalysis


Enamine and SOMO catalysis represent powerful approaches to α -functionalization of aldehydes. In 2011, Wang group reported the seminal work about the direct β -functionalization of simple aldehydes rather than enals in a catalytic enantioselective manner⁵⁵. Although the strategy involving transformation of an iminium ion to an enamine in iminium catalysis has enjoyed tremendous successes (Figure 1.4, eq

(1)), the reverse process in which an enamine is converted to an iminium species has not been explored. Dr. Wang and co-worker reported that the use of *o*-iodoxybenzoic acid (IBX) as an oxidant in the presence of a secondary amine catalyst serves as an effective system for promoting rapid conversion of enamines to iminium ions (Figure 1.4, eq (2))⁵⁵. The preparative power of this process has been demonstrated in the context of direct asymmetric β -functionalization of simple aldehydes. Furthermore, a variety of enantioselective cascade transformations, including triple and quadruple cascades, have been developed for the ‘one-pot’ synthesis of versatile chiral building blocks and structural frameworks (Scheme 1.13)⁵⁵.

Figure 1.4 Aminocatalysis: (1) Iminium Catalysis. (2) Oxidative Enamine Catalysis.



Scheme 1.13 Oxidative Enamine Catalysis in Multiple Cascade Reactions




1.2.2 Non-Covalent Organocatalysts

Understanding of the mechanism of enzymatic catalysis identified the key-role for non-covalent interactions⁵⁶. Now, non-covalent interactions-particularly H-bonding has emerged as a widely applicable approach in organocatalysis. Unlike covalent catalysts which rely on strong covalent interactions with substrates, non-covalent catalysts active substrates and stabilize the transition state through well-defined non-covalent interactions.


The simultaneous donation of two hydrogen bonds has proven to be a highly successful strategy for electrophile activation because such interactions benefit from increased strength and directionality relative to a single hydrogen bond⁵⁷. In 1998,

Sigman and Jacobsen reported that urea and thiourea derivatives can catalyze enantioselective Strecker reaction (Scheme 1.14)⁵⁸. In 2002, a mechanistic analysis revealed that the thiourea functionality is responsible for catalytic activity and that the imine substrate can interact with the catalyst via a dual H-bond interaction to the urea protons⁵⁹. Shortly later, Jacobsen's group identified these chiral (thio)urea as highly versatile, effective catalysts for Mannich reaction (Scheme 1.15), launching the generic use of enantioselective hydrogen-bonding catalysis⁶⁰. Along this line, several examples^{57, 61} have been presented such as Aza-Baylis-Hillman reaction⁶², hydrophosfonylation of imines⁶³ and Acyl-Pictet Spengler reaction⁶⁴ (Scheme 1.16). It should be noted that the Jacobsen (thio)ureas are monofunctional catalysts that work through the sole activation of electrophiles.


Scheme 1.14 Thiourea-Catalyzed Asymmetric Strecker Reaction



Scheme 1.15 Thiourea-Catalyzed Asymmetric Mannich Reaction




Scheme 1.16 Other Examples of Thiourea Catalyst as Double H-Bonding Catalyst




The amino acid arginine possessing a guanidinium functional group, frequently acts as a double H-bond donor in biological systems. Similarly, in synthetic chemistry, guanidinium ion is also capable of participating in double H-bonding interactions⁶⁵. Pioneering work in this area was reported by Corey and Grogan, who demonstrated that chiral bicyclic guanidine mediates asymmetric Strecker reactions (Scheme 1.17)⁶⁶. The utilizations of chiral guanidines as catalysts in many different kinds of reactions such as Henry reaction⁶⁷, Michael reaction⁶⁸, Diels–Alder reaction⁶⁹, and Mannich reaction⁷⁰ have been investigated thoroughly in the past few years⁶⁵.

Scheme 1.17 Bicyclic Guanidine Catalyst for Asymmetric Strecker Reaction




Recently, inspired by enzyme-mediated catalysis, which relies on the synergistic cooperation of a number of functional groups, organic chemists have developed bifunctional organocatalysts which commonly combine the H-bond donors and Brønsted or Lewis base functionalities in a chiral scaffold. Starting with Jacobsen's monofunctional (thio)ureas catalyst, in 2003, Takemoto and co-workers developed novel bifunctional catalyst with the introduction of an additional basic, nucleophile-activating group in the thiourea⁷¹. As reported, the thiourea catalyst **I** promoted the Michael reaction of malonates to various nitroolefins with high enantioselectivities (Scheme 1.18). In the bifunctional catalyst, thiourea moiety activates the Michael acceptor through H-bonding interaction with nitro group and the tertiary amine group serves as Lewis base to activate the nucleophilic enol species simultaneously (Scheme 1.18)⁷². The scope of Takemoto catalyst has also been expanded to transformations using substantially different electrophiles and nucleophiles, such as aza-Henry reaction⁷³, Mannich reaction⁷⁴, cyanosilylation⁷⁵, Baylis–Hillman reaction⁷⁶ and various Michael additions⁷⁷.

Scheme 1.18 Michael Additions of Malonates to Nitroolefins Catalyzed by Takemoto Thiourea Catalyst **I**



The natural products cinchona alkaloids have also been utilized for the creation of a new class of bifunctional organocatalysts by incorporation of thiourea moieties⁷⁸. Pioneering work was reported by Soós and co-workers⁷⁹ about Michael addition of nitromethane to chalcones catalyzed by cinchona alkaloid derived bifunctional thioureas catalyst **II** (Scheme 1.19).

Scheme 1.19 Michael Addition of Nitromethane to Chalcones Catalyzed by Cinchona Alkaloid Derived Bifunctional Thioureas Catalyst **II**




Since then, cinchona alkaloid derived bifunctional thioureas catalysts have been

intensively explored for asymmetric organic transformations. Notable examples include Mannich reaction⁸⁰, Diels-Alder reaction⁸¹ and a wide range of Michael addition⁸².

Wang and co-workers identified chiral naphthyl-derived bifunctional thiourea catalyst **III**. It has been demonstrated to promote the asymmetric Morita-Baylis-Hillman reactions of cyclohexenone with a variety of aldehydes to afford highly functionalized, synthetically useful chiral allylic alcohols (Scheme 1.20)⁸³.

Scheme 1.20 Catalyst **III** Promoted Asymmetric Morita-Baylis-Hillman Reaction



1.3 Summary and Outlook

Over the past ten years, the field of enantioselective organocatalysis has had a significant impact on chemical synthesis and has developed into a practical synthetic paradigm crucial for many applications including drug discovery^{3c,9, 11b}. In principle, the most pivotal to the success of organocatalysis in the past decade has been the invention or identification of generic modes of catalyst activation^{3b}. Based on them, a myriad of asymmetric reactions have been developed and the number of publications on the topic of organocatalysis has recently increased markedly. Given the significance of new

activation modes, it is desirable for chemists to focus their research efforts on uncovering new modes.

Very recently, another valuable achievement is the combination of different activation modes or catalytic strategies into one single transformation to generate previously unattainable molecules⁸⁴. A number of multicatalysis mechanisms have been identified including cooperative metal-organocatalyst system⁸⁵ and organocatalytic photoredox strategy⁸⁶. Although multicatalysis concept is still in its infancy and it has inherent problems such as catalyst compatibility, we believe that it will continue to grow and offer solutions to the challenges; ultimately, it will be recognized as a powerful catalysis strategy.

1.4 Reference

- [1] Anon, "Recognizing the Best in Innovation: Breakthrough Catalyst". *R&D Magazine*, September **2005**, pg 20.
- [2] A. M. Thayer, *Chem. Eng. News*, **2007**, 85, 11-19.
- [3] (a) E. N. Jacobsen, D. W. C. MacMillan, *Proc. Natl. Acad. Sci.* **2010**, 107, 20618-20619; (b) D. W. C. MacMillan, *Nature* **2008**, 455, 304-308; (c) M. J. Gaunt, C. C.C. Johansson, A. McNally, N. T. Vo, *Drug Discovery Today*. **2007**, 12, 8-27.
- [4] G. Bredig, W. S. Fiske, *Biochem.Z.* **1912**, 7.

- [5] (a) H. Pracejus, *Justus Liebigs Ann. Chem.* **1960**, 634, 9-22; (b) H. Pracejus, H. Matje, *J. Prakt. Chem.* **1964**, 24, 195-205.
- [6] (a) Z. G. Hajos, D. R. Parrish, *German Patent* **1971**, DE2102623; (b) Z. G. Hajos, D. R. Parrish, *J. Org. Chem.* **1974**, 39, 1615-1621; (c) U. Eder, G. Sauer, R. Wiechert, *German Patent* **1971**, DE2014757; (d) U. Eder, G. Sauer, R. Wiechert, *Angew. Chem. Int. Ed.* **1971**, 10, 496-497.
- [7] B. List, R. A. Lerner, C. F. III Barbas, *J. Am. Chem. Soc.* **2000**, 122, 2395-2396.
- [8] K. A. Ahrendt, C. J. Borths, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2000**, 122, 4243-4244.
- [9] P. I. Dalko, L. Moisan, *Angew. Chem. Int. Ed.* **2004**, 43, 5138-5175.
- [10] H. Pellissier, *Tetrahedron* **2007**, 63, 9267-9331.
- [11] (a) A. Berkessel, H. Gröger, *Asymmetric Organocatalysis: From Biomimetic Concepts to Applications in Asymmetric Synthesis*, Wiley-VCH, Weinheim, **2005**; (b) P. I. Dalko, *Enantioselective Organocatalysis*, Wiley-VCH, Weinheim, **2007**.
- [12] B. List, *Chem. Commun.* **2006**, 819-824.
- [13] S. Mukherjee, J. W. Yang, S. Hoffmann, B. List, *Chem. Rev.* **2007**, 107, 5471-5569.
- [14] W. Notz, F. Tanaka, C. F. Barbas III, *Acc. Chem. Res.* **2004**, 37, 580-591.
- [15] B. List, P. Pojarliev, W. T. Biller, H. J. Martin, *J. Am. Chem. Soc.* **2002**, 124, 827-833.

- [16] A. Córdova, W. Notz, G. Zhong, J. M. Betancort, C. F. Barbas III, *J. Am. Chem. Soc.* **2002**, *124*, 1842-1843.
- [17] A. Córdova, S. Watanabe, F. Tanaka, W. Notz, C. F. Barbas, III, *J. Am. Chem. Soc.* **2002**, *124*, 1866-1867.
- [18] W. Wang, J. Wang, H. Li, *Angew. Chem. Int. Ed.* **2005**, *44*, 1369-1371.
- [19] B. List, *J. Am. Chem. Soc.* **2002**, *124*, 5656-5657.
- [20] N. Kumaragurubaran, K. Juhl, W. Zhuang, A. Bøgevig, K. A. Jørgensen, *J. Am. Chem. Soc.* **2002**, *124*, 6254-6255.
- [21] P. Merino, T. Tejero, *Angew. Chem. Int. Ed.* **2004**, *43*, 2995-2997.
- [22] M. Marigo, D. Fielenbach, A. Branton, A. Kjærsgaard, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2005**, *44*, 3703-3706.
- [23] S. Bertelsen, N. Halland, S. Bachmann, M. Marigo, A. Branton, K. A. Jørgensen, *Chem. Commun.* **2005**, 4821-4823.
- [24] M. P. Brochu, S. P. Brown, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2004**, *126*, 4108-4109.
- [25] M. Marigo, T. C. Wabnitz, D. Fielenbach, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2005**, *44*, 794-797.
- [26] M. Tiecco, A. Carlone, S. Sternativo, F. Marini, G. Bartoli, P. Melchiorre, *Angew. Chem. Int. Ed.* **2007**, *46*, 6882-6885.

- [27] H. Sundén, R. Rios, A. Córdova, *Tetrahedron Lett.* **2007**, 48, 7865-7869.
- [28] N. Vignola, B. List, *J. Am. Chem. Soc.* **2004**, 126, 450-451.
- [29] (a) L. F. Tietze, U. Beifuss, In *Comprehensive Organic Synthesis*; B. M. Trost, I. Fleming, Eds.; Pergamon: Oxford, U.K., **1991**; Vol. 2, p 341; (b) E. Knoevenagel, *Chem. Ber.* **1894**, 27, 2345.
- [30] K. C. Blanchard, D. L. Klein, J. MacDonald, *J. Am. Chem. Soc.* **1931**, 53, 2809.
- [31] T. I. Crowell, D. W. Peck, *J. Am. Chem. Soc.* **1953**, 75, 1075-1077.
- [32] K. A. Ahrendt, C. J. Borths, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2000**, 122, 4243-4244.
- [33] W. S. Jen, J. J. M. Wiener, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2000**, 122, 9874-9875.
- [34] M. Harmata, S.K. Ghosh, X. Hong, S. Wacharasindhu, P. Kirchhoefer, *J. Am. Chem. Soc.* **2003**, 125, 2058-2059.
- [35] N. A. Paras, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2001**, 123, 4370-4371.
- [36] S. P. Brown, N. C. Goodwin, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2003**, 125, 1192-1194.
- [37] S. Brandau, A. Landa, J. Franzen, M. Marigo, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2006**, 45, 4305-4309.

- [38] S. G. Ouellet, J. B. Tuttle, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2005**, *127*, 32-33.
- [39] R. K. Kunz, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2005**, *127*, 3240-3241.
- [40] (a) M. Marigo, J. Franzén, T.B. Poulsen, W. Zhuang, K.A. Jørgensen, *J. Am. Chem. Soc.* **2005**, *127*, 6964-6965; (b) W. Zhuang, M. Marigo, K.A. Jørgensen, *Org. Biomol. Chem.* **2005**, 3883-3885; (c) H. Sundén, I. Ibrahim, A. Córdova, *Tetrahedron Lett.* **2006**, *47*, 99-103.
- [41] A. Erkkil, I. Majander, P. M. Pihko, *Chem. Rev.* **2007**, *107*, 5416-5470.
- [42] C. Grondal, M. Jeanty, D. Enders, *Nature Chemistry*, **2010**, *2*, 167-178.
- [43] C. Palomo, A. Mielgo, *Angew. Chem. Int. Ed.* **2006**, *45*, 7876-7880.
- [44] (a) G. Bartoli, P. Melchiorre, *Synlett* **2008**, *12*, 1759-1772; (b) Y.-C. Chen, *Synlett* **2008**, *13*, 1919-1930.
- [45] T.-Y. Liu, H.-L. Cui, Y. Zhang, K. Jiang, W. Du, Z.-Q. He, Y.-C. Chen, *Org. Lett.* **2007**, *9*, 3671-3674.
- [46] P. Galzerano, F. Pesciaioli, A. Mazzanti, G. Bartoli, P. Melchiorre, *Angew. Chem. Int. Ed.* **2009**, *48*, 7892-7894.
- [47] K. Narasaka, T. Okauchi, K. Tanaka, M. Murakami, *Chem. Lett.* **1992**, *21*, 2099-2102.
- [48] T. D. Beeson, A. Mastracchio, J. B. Hong, K. Ashton, D. W. C. MacMillan, *Science*, **2007**, *316*, 582-585.

- [49] H.-Y. Jang, J.-B. Hong, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2007**, *129*, 7004-7005.
- [50] H. Kim, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2008**, *130*, 398-399.
- [51] A. Mastracchio, A. A. Warkentin, A. M. Walji, D. W. C. MacMillan, *PNAS*, **2010**, *107*, 20648-20651.
- [52] S. Bertelsen, M. Marigo, S. Brandes, P. Dinér, K. A. Jørgensen, *J. Am. Chem. Soc.* **2006**, *128*, 12973-12980.
- [53] R. M. de Figueiredo, R. Fröhlich, M. Christmann, *Angew. Chem. Int. Ed.* **2008**, *47*, 1450-1453.
- [54] G. Bencivenni, P. Galzerano, A. Mazzanti, G. Bartoli, P. Melchiorre, *PNAS*, **2010**, *107*, 20642-20647.
- [55] S.-L. Zhang, H.-X.Xie, J. Zhu, H. Li, X.-S. Zhang, J. Li, W. Wang, *Nature Communications*, **2011**, DOI: 10.1038/ncomms1214.
- [56] R. B. Silverman, *The Organic Chemistry of Enzyme-Catalyzed Reactions*; Academic Press: San Diego, CA, **2002**.
- [57] M. S. Taylor, E. N. Jacobsen, *Angew. Chem. Int. Ed.* **2006**, *45*, 1520-1543.
- [58] M. S. Sigman, E. N. Jacobsen, *J. Am. Chem. Soc.* **1998**, *120*, 4901-4902.
- [59] P. Vachal, E. N. Jacobsen, *J. Am. Chem. Soc.* **2002**, *124*, 10012-10014.

- [60] A. G. Wenzel, E. N. Jacobsen, *J. Am. Chem. Soc.* **2002**, *124*, 12964-12965.
- [601] A. G. Doyle, E. N. Jacobsen, *Chem. Rev.* **2007**, *107*, 5713-5743.
- [62] I. T. Raheem, E. N. Jacobsen, *Adv. Synth. Catal.* **2005**, *347*, 1701-1708.
- [63] G. D. Joly, E. N. Jacobsen, *J. Am. Chem. Soc.* **2004**, *126*, 4102-4103.
- [64] M. S. Taylor, E. N. Jacobsen, *J. Am. Chem. Soc.* **2004**, *126*, 10558-10559.
- [65] D. Leow, C.-H. Tan, *Chem. Asian J.* **2009**, *4*, 488-507.
- [66] E. J. Corey, M. J. Grogan, *Org. Lett.* **1999**, *1*, 157-160.
- [67] (a) R. Chinchilla, C. Nájera, P. Sánchez-Agulló, *Tetrahedron: Asymmetry* **1994**, *5*, 1393-1402; (b) A. Howard-Jones, P. J. Murphy, D. A. Thomas, *J. Org. Chem.* **1999**, *64*, 1039-1041; (c) M. T. Allingham; A. Howard-Jones, P. J. Murphy, D. A. Thomas, P. W. R. Caulkett, *Tetrahedron Lett.* **2003**, *44*, 8677-8680.
- [68] (a) W. Ye, D. Leow, S. L. M. Goh, C.-T. Tan, C.-H. Chian, C.-H. Tan, *Tetrahedron Lett.* **2006**, *47*, 1007-1010; (b) T. Ishikawa, Y. Araki, T. Kumamoto, H. Seki, K. Fukuda, T. Isobe, *Chem. Commun.* **2001**, 245-246; (c) M. Terada, T. Ikehara, H. Ube, *J. Am. Chem. Soc.* **2007**, *129*, 14112-14113.
- [69] (a) J. Shen, T. T. Nguyen, Y.-P. Goh, W. Ye, X. Fu, J. Xu, C.-H. Tan, *J. Am. Chem. Soc.* **2006**, *128*, 13692-13693; (b) J. Shen, C.-H. Tan, *Org. Biomol. Chem.* **2008**, *6*, 4096-4098.
- [70] S. Kobayashi, R. Yazaki, K. Seki, Y. Yamashita, *Angew. Chem. Int. Ed.* **2008**, *47*,

5613-5615.

[71] T. Okino, Y. Hoashi, Y. Takemoto, *J. Am. Chem. Soc.* **2003**, *125*, 12672-12673.

[72] X.-H. Yu, W. Wang, *Chem. Asian J.* **2008**, *3*, 516-532.

[73] T. Okino, S. Nakamura, T. Furukawa, Y. Takemoto, *Org. Lett.* **2004**, *6*, 625-627.

[74] Y. Yamaoka, H. Miyabe, Y. Yasui, Y. Takemoto, *Synthesis* **2007**, 2571-2575.

[75] D. E. Fuerst, E. N. Jacobsen, *J. Am. Chem. Soc.* **2005**, *127*, 8964-8965.

[76] Y. Sohtome, A. Tanatani, Y. Hashimoto, K. Nagasawa, *Tetrahedron Lett.* **2004**, *45*, 5589-5592.

[77] (a) H. Li, L.-S. Zu, J. Wang, W. Wang, *Tetrahedron Lett.* **2006**, *47*, 3145-3148; (b) Y. Hoashi, T. Okino, Y. Takemoto, *Angew. Chem. Int. Ed.* **2005**, *44*, 4032-4035; (c) H. Li, J. Wang, L.-S. Zu, W. Wang, *Tetrahedron Lett.* **2006**, *47*, 2585-2589.

[78] S. J. Connon, *Chem. Commun.* **2008**, 2499-2510.

[79] B. Vakulya, S. Varga, A. Csámpai, T. Soós, *Org. Lett.* **2005**, *7*, 1967-1969.

[80] J. Song, Y. Wang, L. Deng, *J. Am. Chem. Soc.* **2006**, *128*, 6048-6049.

[81] Y. Wang, H. Li, Y.-Q. Wang, Y. Liu, B. M. Foxman, L. Deng, *J. Am. Chem. Soc.* **2007**, *129*, 6364-6365.

[82] (a) J. Wang, H. Li, L.-S. Zu, W. Jiang, H.-X. Xie, W.-H. Duan, W. Wang, *J. Am. Chem. Soc.* **2006**, *128*, 12652-12653; (b) J.-X. Ye, D. J. Dixon, P. S. Hynes, *Chem.*

Commun. **2005**, 4481-4483; (c) S. H. McCooey, S. J. Connon, *Angew. Chem. Int. Ed.* **2005**, *44*, 6367-6370.

[83] J. Wang, H. Li, X.-H. Yu, L.-S. Zu, W. Wang, *Org. Lett.* **2005**, *7*, 4293-4296.

[84] A. E. Allen, D. W. C. MacMillan, *Chem. Sci.* **2012**, *3*, 633-658.

[85] (a) M. Ikeda, Y. Miyake, Y. Nishibayashi, *Angew. Chem. Int. Ed.* **2010**, *49*, 7289-7293; (b) A. S. K. Hashmi, C. Hubbert, *Angew. Chem. Int. Ed.* **2010**, *49*, 1010-1012; (c) P. de Armas, D. Tejedor, F. García-Tellado, *Angew. Chem. Int. Ed.* **2010**, *49*, 1013-1016; (d) I. Ibrahim, A. Córdova, *Angew. Chem. Int. Ed.* **2006**, *45*, 1952-1956; (e) A. Duschek, S. F. Kirsch, *Angew. Chem. Int. Ed.* **2008**, *47*, 5703-5705.

[86] (a) D. A. Nicewicz, D. W. C. MacMillan, *Science*, **2008**, *322*, 77-80; (b) D. A. Nagib, M. E. Scott, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2009**, *131*, 10875-10877; (c) H.-W. Shih, M. N. Vander Wal, R. L. Grange, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2010**, *132*, 13600-13603.

Chapter 2


Organocatalytic Regio- and Stereoselective Michael Addition of 1*H*-1, 2, 3-Triazole to Alkynals

2.1 Background and Significance


The catalytic Michael addition reactions which efficiently construct C-C and C-heteroatom bonds held persistent interest in the area of asymmetric organocatalysis. Notably, high effective asymmetric organocatalytic additions of various nucleophiles to α , β -unsaturated cyclic and acyclic enals, enones, nitro olefins, vinyl sulfones and vinyl phosphonates have been developed¹. Among those, Michael additions to enals represent an enormous and fruitful research branch². In contrast, the protocols for conjugate additions to the corresponding alkynals are relatively rare. This is due to the fact that (i) alkynals possess no prochiral center at the β -carbon atom; (ii) alkynals are known to be a challenging class of electrophiles in asymmetric synthesis, and it is expected that they would behave differently to the well-established enals in terms of reactivity and stereocontrol, because they adopt different structures and geometry (linear sp hybridized alkynes versus $trans\ sp^2$ hybridized olefins). There are only a very small number of reports dealing with Michael additions to alkynones or acetylenic esters which introduce chiral centers by the Michael donors (Scheme 2.1)^{3a-3c}. And the examples of substituted alkynals or alkynones act as acetylenic dienophiles in Diels-Alder reactions are sporadic

(Scheme 2.2)^{3d-3h}.

Scheme 2.1 Examples of Catalytic Asymmetric Michael Additions to Alkynones or Acetylenic Esters



Scheme 2.2 Examples of Acetylenic Dienophiles in Asymmetric Diels-Alder Reaction



2.2 Research Design

Motivated by the dearth of applications of alkynals in organocatalysis (versus the widely explored enals), we are pursuing to develop alkynals involved organocatalytic single-step reaction. To the best of our knowledge, in the case of these Michael additions to alkynals or alkynones, the iminium-ion activation mechanism which facilitates the addition of the nucleophile to the β -carbon atom has not been reported although the similar mechanism for activation of enals is already well-established.

In order to successfully develop the conjugate addition to alkynal, we surmise that the key issue is to choose the Michael donor. After a brief screening, we chose to test the Michael addition of 1*H*-1, 2, 3-triazole to alkynals. This is based on the fact that *N*-(1-Alkenyl)azoles are involved in the synthesis of polymers⁴ and heteropentalene mesomeric betaines of type B⁵. Moreover, *N*-(1-Alkenyl)azoles have found to display intriguing biological activities⁶.

2.3 Introduction

The efficient regio- and stereoselective construction of trisubstituted alkenes is a long standing goal in organic synthesis. Carbonyl olefination such as Wittig⁷ and Horner-Wadsworth-Emmons reaction⁸ is the most classical strategy. Transition-metal mediated carbometallation of alkynes⁹ and cross-coupling reactions¹⁰ has emerged as most widely used methods for formation of trisubstituted alkenes. In all above strategies, regio- and stereocontrol are the two key issues and remain a formidable challenge. Accordingly, we

envisioned to develop a regio- and stereoselective synthetic strategy for trisubstituted alkenes starting from alkynals by employing organocatalyst via an iminium-ion activation mechanism.

2.4 Results and Discussion

At the outset of our investigations, the model reaction between phenyl-propynal **1a** and 1*H*-1, 2, 3-triazole was examined in dichloromethane at room temperature in the presence of 20 mol% organocatalyst (Figure 2.1 and Table 2.1). Several types of secondary amine catalysts **I-VII** (Figure 2.1) were screened because they can activate alkynal via iminium-ion mechanism. It was found that the catalysts tested exhibited significantly different catalytic activity in the event. The diarylprolinol silyl ethers catalyst **I** and **II** promoted the process to achieve 100% conversion in 2 h with good yield (~93%) and good *E/Z* ratio (1:0.19 and 1:0.18 respectively) (Table 2.1, entry 1, 2). It is worthy of point out that the reaction also proceeded with pretty good regioselectivity, affording the *N*-2-alkenyl substituted triazole derivative **2a** as the major product. However, the either moieties had little impact on the *E/Z* ratio due to the small Michael donor 1*H*-1, 2, 3-triazole¹¹. The reaction proceeded to afford **2a** in 83% yield and a little poor *E/Z* ratio (1:0.23) after 60 h using diphenylprolinol **III** as catalyst (Table 2.1, entry 3). Pyrrolidine-based diamine catalyst **IV** was also good promoter for the reaction, resulted in high yield (94%), good *E/Z* ratio (1:0.17) and excellent regioselectivity (*N*-2-addition). The stereoselectivity between catalyst **II** and **IV** was nearly the same for the above model reaction (Table 2.1, entry 2, 4). So it was necessary to change the Michael

acceptor phenyl-propynal **1a** to (4-methoxy-phenyl)-propynal **1b** to evaluate which catalyst was more tolerate to versatile substrates. Correspondingly, the *E/Z* ratio was 1:0.29 and 1:0.20 when catalyst **II** and **IV** were employed respectively. Poor regioselectivity for Wang's catalyst **VI** was observed (Table 2.1, entry 6). The catalytic activity of L-proline **V** and MacMillan's catalyst **VII** were very poor; the conversion is only 58% and 29% after 60 h respectively. Moreover, catalyst **V** and **VII** afforded *N*-1-alkenyl substituted triazole derivative **3a**. Thereby, the diamine catalyst **IV** was chosen for further optimization reaction condition. Then, the effects of additives on the *E/Z* ratio were probed. The *E/Z* ratio can be improved slightly by adding 1eq. of NaOAc (Table 2.1, entry 8). It is considered as the result of facilitating the deprotonation of 1*H*-1, 2, 3-triazole. Generally, the relatively strong base such as K₂CO₃ was considered as a detriment to the *E/Z* ratio (Table 2.1, entry 9). The steric organic base quinine (Table 2.1, entry 10) and DABCO (Table 2.1, entry 11) cannot achieve better results. Et₃N (Table 2.1, entry 12) is also a poor additive although without sacrificing the yield. In the iminium-ion activation model, PhCO₂H additive usually favors the formation of iminium-ion, as a consequence, improve the reactivity of Michael acceptor. However, in this instance, PhCO₂H (20 mol%) additive can react with phenyl-propynal **1a** completely in 1 h which lead to more complicated reaction system and low yield (Table 2.1, entry 13). The screening of the reaction solvent indicated the use of dichloromethane gave the best results (Table 2.1, entry 14, 15 and 16). At last, we found unexpectedly lowering the reaction temperature to -40°C, the *E/Z* ratio became worse along with the prolonged reaction time (Table 2.1, entry 17). The configuration of the dominant component of

compound **2a** prepared under the optimal condition was determined by X-ray crystallography to be *E* configuration; in other words, the organocatalytic Michael addition process was kinetic controlled process.

Figure 2.1 Secondary Amine Catalysts Screened





Table 2.1 Optimization Reaction Conditions for the Organocatalytic Regio- and Stereocontrolled Michael Addition of **1a** and 1*H*-1, 2, 3-Triazole to Afford **2a**^a.



Entry	Catalyst	Additive	Solvent	<i>t</i> (h)	Yield (%) ^b	<i>E/Z</i> ratio ^c
1	I	None	CH ₂ Cl ₂	2	93	1:0.19
2	II	None	CH ₂ Cl ₂	2	93	1:0.18
3	III	None	CH ₂ Cl ₂	60	83	1:0.23
4	IV	None	CH ₂ Cl ₂	2	94	1:0.17
5	V	None	CH ₂ Cl ₂	60	<5	ND ^d
6	VI	None	CH ₂ Cl ₂	24	64 ^e	1:0.46
7	VII	None	CH ₂ Cl ₂	60	<5	ND ^d

8	IV	NaOAc	CH ₂ Cl ₂	2	94	1:0.16
9	IV	K ₂ CO ₃	CH ₂ Cl ₂	2	92	1:0.24
10	IV	quinine	CH ₂ Cl ₂	3	90	1:0.16
11	IV	DABCO	CH ₂ Cl ₂	2	85	1:0.26
12	IV	Et ₃ N ^f	CH ₂ Cl ₂	2	92	1:0.26
13	IV	PhCOOH ^g	CH ₂ Cl ₂	2	58 ^e	1:0.28 ^e
14	IV	None	DMSO	9	47 ^e	1:0.56
15	IV	None	toluene	5	93	1:0.23
16	IV	None	THF	48	<5	ND ^d
17 ^h	IV	NaOAc	CH ₂ Cl ₂	48	85	1:0.26

^a Reaction conditions: unless specified, a mixture of phenyl-propynal **1a** (0.05mmol), 1*H*-1, 2, 3-triazole (0.06mmol), catalyst (0.01mmol) and additive (0.05mmol) in solvent was stirred at r.t. for a specified time.

^b Isolated yields. ^c Determined by ¹HNMR of crude reaction solution based on the integral of hydrogen of aldehydes or the integral of hydrogen of alkenes. ^d Not determined. ^e Calculated from ¹HNMR of crude reaction solution based on the integral of hydrogen of aldehyde of two regioisomers. ^f 40 mol% Et₃N was employed. ^g 20 mol% PhCOOH was employed. ^h Performed between (4-Methoxy-phenyl)-propynal **1b** and 1*H*-1, 2, 3-triazole at -40°C.

As mentioned above, L-proline **V** and MacMillan's catalyst **VII** afforded *N*-1-alkenyl substituted triazole derivative **3a** although the conversion was low. The results encouraged us to further explore optimal protocol for efficient synthesis of the regioisomer **3a**. An extensive screening of catalysts (Figure 2.1, 2.2), additives and solvents were carried out and some representative results are shown in Table 2.2. Generally speaking, acid additives usually favored the formation of *N*-1-alkenyl substituted triazole derivatives for most secondary amine catalysts. It was found that TFA

additive lead to remarkable reaction rate acceleration together with improvement in the regio- and stereoselectivity. As seen from the results, employing imidazolidine-2-carboxylic acid **IX** as catalyst, TFA as additive and dichloroethane as solvent afforded the best outcome (Table 2.2, entry 8).

There exist tautomerism for 1*H*-1, 2, 3-triazole (Figure 2.3 and Figure 2.4). In a neutral solution, **1B**-configuration was more stable than **1A**-configuration due to the lone pair/lone pair repulsion of adjacent nitrogen atoms. So the predominant more electron rich species **1B** can be deprotonated, followed by Michael addition to alkynals at *N*-2 position. By contrast, under an acidic condition, **2B** exists as a dominant protonated form, which renders less nucleophilic. Therefore, **2A** serves as a nucleophile for *N*-1 addition. It should be noted that the bifunctional catalyst **X** also afforded the *N*-1 addition product exclusively (Table 2.2, entry 11). This might be because the thiourea part of catalyst **X** form H-bonding with two adjacent nitrogen atoms of 1*H*-1, 2, 3-triazole, which rendered them less nucleophilicity. The stereoconfiguration of the major component of the compound **3a** prepared under the optimal condition was determined by X-ray crystallography to be *E* configuration based on enol **4** derived from **3a**. The observed configuration is consistent with the proposed mechanism of the process.

Figure 2.2 Other Secondary Amine Catalysts Screened





Table 2.2 Optimization Reaction Conditions for the Organocatalytic Regio- and Stereocontrolled Michael Addition of **1a** and 1*H*-1, 2, 3-Triazole to Afford **3a**^a.



Entry	Catalyst	Additive	Solvent	<i>t</i> (h)	Yield (%) ^b	E/Z ratio ^c
1	VIII	HCl	DCM	60	83	1:0.89
2	II	TFA	DCM	30	85	1:0.69
3	VII	TFA	DCM	7	89	1:0.60
4	IX	<i>p</i> TsOH•H ₂ O	DCM	10	84	1:0.49
5	IX	TFA	DCM	8	87	1:0.45
6	IX	CH ₃ SO ₃ H	DCM	10	85	1:0.63
7	IX	CF ₃ SO ₃ H	DCM	6	91	1:0.53
8	IX	TFA	DCE	9	88	1:0.41
9	IX	TFA	toluene	10	83	1:0.45
10	IX	TFA	MeOH	60	47 ^d	1:0.45 ^d
11	X	None	DCM	48	90	1:0.83

^a Reaction conditions: unless specified, a mixture of phenyl-propynal **1a** (0.05mmol), 1*H*-1, 2, 3-triazole (0.058mmol), catalyst (0.01mmol) and additive (0.01mmol) in solvent was stirred at r.t. for a specified time.

^b Isolated yields. ^c Determined by ¹HNMR of isolated product based on the integral of hydrogen of aldehydes or the integral of hydrogen of alkenes. ^d Determined by ¹HNMR of crude reaction solution.

Figure 2.3 Tautomerism of Neutral 1*H*-1, 2, 3-Triazole





Figure 2.4 Tautomerism of Triazolium Ion




Having established the optimal protocols for the reactions, we proceeded to examine the substrate scope with the results summarized in Table 2.3. In general, the additions of 1*H*-1, 2, 3-triazole to alkynals **1** which afforded compounds **2** proceeded with high yields (75%-96%) and good stereoselectivity (*E/Z* ratio ranged from 82:18 to 95:5) within 2 hours. It is notable that alkynals were more active toward the nucleophile 1*H*-1, 2, 3-triazole under the present condition compared with the corresponding enals; for example, cinnamaldehyde gave only low conversions (<20%)¹². Phenyl-propynal derivatives bearing electron-withdrawing group on the *meta*- and *para*-position of the benzene ring gave slightly poor stereoselectivity (Table 2.3, entry 9-12). The similar stereoselectivity were observed for *meta*- and *para*- position halogen substituted phenyl-propynal (Table 2.3, entry 4-7). Only strong electron-donating group resulted in a little poor stereoselectivity (Table 2.3, entry 2 vs. 3). To our surprise, among the substrates we surveyed, *ortho*-fluoro and *meta*-trifluoromethane substituted phenyl-propynal afforded

distinctive stereoselectivity (*E/Z* ratio were 95:5 and 91:9 respectively). The aliphatic alkynal also works well with good stereoselectivity although the reaction yield is relatively low (Table 2.3, entry 14).

As for the reactions which synthesized compound **3**, longer reaction time was required. For aromatic alkynals **1**, regardless of electron-donating and electron-withdrawing substituents on the phenyl ring, participated in the process in moderate yield (79%-90%). However, the *E/Z* ratio ranged from 66:34 to 77:23. It is worthy of point out that the reaction proceeded with much better stereoselectivity (*E/Z* ratio was 89:11) for aliphatic alkynal (Table 2.3, entry 14).

Table 2.3 Scope of the Organocatalytic Regio- and Stereoselective Michael Addition of 1*H*-1, 2, 3-Triazole to Substituted Phenyl-propynal **1a-m** to Afford **2^a** and **3^b** Respectively.



Entry	R	2			3		
		<i>t</i> (h)	Yield (%) ^c	<i>E/Z</i> ratio ^d	<i>t</i> (h)	Yield (%) ^c	<i>E/Z</i> ratio ^d
1	Ph	2	94	86:14	9	88	71:29
2	<i>p</i> -CH ₃ OC ₆ H ₄	2	92	84:16	10	89	69:31
3	<i>p</i> -CH ₃ C ₆ H ₄	2	89	85:15	10	87	70:30
4	<i>p</i> -BrC ₆ H ₄	2	90	85:15	10	88	71:29
5	<i>p</i> -ClC ₆ H ₄	2	93	84:16	10	90	70:30

6	<i>p</i> -FC ₆ H ₄	2	90	86:14	10	83	68:32
7	<i>m</i> -FC ₆ H ₄	2	96	86:14	10	91	77:23
8	<i>o</i> -FC ₆ H ₄	2	96	95:5	10	86	73:27
9	<i>m</i> -NO ₂ C ₆ H ₄	2	95	82:18	11	89	69:31
10	<i>p</i> -NO ₂ C ₆ H ₄	2	88	84:16	10	83	68:32
11	<i>m</i> -CNC ₆ H ₄	2	90	83:17	10	79	69:31
12	<i>p</i> -CNC ₆ H ₄	2	92	82:18	10	80	66:34
13	<i>m</i> -CF ₃ C ₆ H ₄	2	93	91:9	10	82	66:34
14	<i>n</i> -C ₅ H ₁₁	2	75	92:8	11	67	89:11

^a Reaction conditions: unless specified, a mixture of substituted phenyl-propynal **1** (0.08mmol), 1*H*-1, 2, 3-triazole (0.096mmol), catalyst **IV** (0.016mmol) and NaOAc (0.08mmol) in dichloromethane (0.15ml) was stirred at r.t. for 2 h. ^b Reaction conditions: unless specified, a mixture of substituted phenyl-propynal **1** (0.08mmol), 1*H*-1, 2, 3-triazole (0.092mmol), catalyst **IX** (0.016mmol) and trifluoroacetic acid (0.016mmol) in DCE (0.15ml) was stirred at r.t. for a specified time. ^c Isolated yields. ^d Determined by ¹HNMR of isolated product based on the integral of hydrogen of aldehydes or the integral of hydrogen of alkenes.

2.5 Conclusions

In summary, we have developed a new strategy for regio- and stereocotrolled synthesis of trisubstituted alkenes by employing appropriate organocatalysts and additives via iminium-ion mechanism starting from alkynals with high efficiency and good stereoselectivity. Under the catalysis of (*S*)-(+)-1-(2-pyrrolidinylmethyl) pyrrolidine and sodium acetate as additive, the *N*-2 addition product can be achieved in very good yield and the *Z/E* ratio is up to 5:95. Under the catalysis of imidazolidine-2-carboxylic

acid and TFA as additive, the *N*-1 addition product can be obtained in good yield and the *Z/E* ratio is nearly 1:2.

2.6 Supporting Information

General information: (This information is valid for all the experimental parts)


All commercial reagents and solvents were used without further purification unless otherwise noted. Analytical thin-layer chromatography (TLC) was performed on Sorbent Technologies silica gel plates with fluorescence F_{254} indicator. And column chromatography was performed using the indicated solvent on Merck 60 silica gel (230-400 mesh). ^1H NMR and ^{13}C NMR spectra were recorded on Bruker Avance 500. ^1H NMR data were reported as follows: chemical shifts in ppm using 7.26 signal of CDCl_3 and 2.50 signal of $\text{DMSO}-d_6$ as reference respectively, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, and br = broad), coupling constant (*J*), and integration. And ^{13}C NMR data were reported in ppm using 77.0 signal of CDCl_3 and 39.50 signal of $\text{DMSO}-d_6$ as reference respectively.

General procedure for Michael addition of 1*H*-1, 2, 3-triazole to substituted phenylpropynal to afford compound 2

To a solution of substituted phenylpropynal (0.08mmol), (*s*)-(+)-1-(2-pyrrolidinylmethyl) pyrrolidine (2.61ul, 0.016mmol) and sodium acetate (6.56mg, 0.08mmol) in dichloromethane (0.15ml) was added 1*H*-1, 2, 3-triazole (5.56ul, 0.096mmol). The resulting solution was stirred at room temperature for a specified time. Then the reaction


mixture was directly purified by column chromatography, eluted with hexane/EtOAc to afford the desired product.

The following spectra information is related with the major isomer.




3-Phenyl-3-[1, 2, 3] triazol-2-yl-propenal (2a) (Table 2.3, Entry 1)

The title compound was prepared according to the general procedure, as described above in 94% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.51 (d, $J = 8.0$ Hz, 1H), 7.87 (s, 2H), 7.59~7.56 (m, 1H), 7.53~7.50 (m, 2H), 7.47 (d, $J = 7.5$ Hz, 2H), 7.06 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.1, 155.5, 137.8, 131.1, 129.8, 128.5, 118.7.




3-(4-Methoxy-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2b) (Table 2.3, Entry 2)

The title compound was prepared according to the general procedure, as described above in 92% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.54 (d, $J = 7.5$ Hz, 1H), 7.87 (s, 2H), 7.40 (d, $J = 8.0$ Hz, 2H), 7.01 (d, $J = 8.5$ Hz, 2H), 6.97 (d, $J = 8.0$ Hz, 1H), 3.87 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.2, 161.9, 155.5, 137.7, 132.9, 121.9, 118.5, 114.0, 55.4.




3-*p*-Tolyl-3-[1, 2, 3] triazol-2-yl-propenal (2c) (Table 2.3, Entry 3)

The title compound was prepared according to the general procedure, as described above in 89% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.53 (d, *J* = 8.0 Hz, 1H), 7.86 (s, 2H), 7.37~7.31 (m, 4H), 7.02 (d, *J* = 8.0 Hz, 1H), 2.44 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 192.2, 155.7, 141.6, 137.7, 131.1, 129.2, 126.9, 118.6, 21.5.




3-(4-Bromo-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2d) (Table 2.3, Entry 4)

The title compound was prepared according to the general procedure, as described above in 90% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.51 (d, *J* = 8.0 Hz, 1H), 7.87 (s, 2H), 7.66 (d, *J* = 8.5 Hz, 2H), 7.34 (d, *J* = 8.5 Hz, 2H), 7.06 (d, *J* = 8.0 Hz, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 191.4, 154.2, 137.9, 132.5, 131.9, 128.7, 125.9, 118.8.




3-(4-Chloro-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2e) (Table 2.3, Entry 5)

The title compound was prepared according to the general procedure, as described above in 93% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.51 (d, $J = 8.0$ Hz, 1H), 7.87 (s, 2H), 7.50 (d, $J = 8.0$ Hz, 2H), 7.41(d, $J = 8.5$ Hz, 2H), 7.06 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.4, 154.2, 137.9, 137.2, 132.4, 129.8, 128.9, 118.9.



3-(4-Fluoro-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2f) (Table 2.3, Entry 6)


The title compound was prepared according to the general procedure, as described above in 90% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.46 (d, $J = 8.0$ Hz, 1H), 7.82 (s, 2H), 7.43~7.41 (m, 2H), 7.19~7.15 (m, 2H), 7.01 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.6, 164.3 ($J = 251$ Hz), 154.3, 137.9, 133.2 ($J = 8.6$ Hz), 118.9, 115.9 ($J = 22$ Hz).



3-(3-Fluoro-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2g) (Table 2.3, Entry 7)


The title compound was prepared according to the general procedure, as described above in 96% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.51 (d, $J = 8.0$ Hz, 1H), 7.87 (s, 2H), 7.52~7.48 (m, 1H), 7.30~7.27 (m, 2H), 7.20~7.18 (m, 1H), 7.06 (d, $J = 8.0$ Hz, 1H); ^{13}C

NMR (CDCl₃, 125 MHz): δ 191.4, 162.3 ($J = 247$ Hz), 153.8, 137.9, 131.7 ($J = 7.8$ Hz), 130.2 ($J = 8.1$ Hz), 127.0, 118.9, 118.2 ($J = 21$ Hz).




3-(2-Fluoro-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2h) (Table 2.3, Entry 8)

The title compound was prepared according to the general procedure, as described above in 96% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.50 (d, $J = 8.0$ Hz, 1H), 7.89 (s, 2H), 7.60~7.56 (m, 1H), 7.47~7.44 (m, 1H), 7.32~7.29 (m, 1H), 7.25~7.21 (m, 1H), 7.15 (d, $J = 8.0$ Hz, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 191.2, 160.4 ($J = 250$ Hz), 150.0, 137.8, 133.1 ($J = 8.0$ Hz), 132.7, 124.2, 119.2, 118.1 ($J = 15$ Hz), 116.2 ($J = 21$ Hz).




3-(3-Nitro-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2i) (Table 2.3, Entry 9)

The title compound was prepared according to the general procedure, as described above in 95 % yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.50 (d, $J = 8.0$ Hz, 1H), 8.44 (d, $J = 8.5$ Hz, 1H), 8.36 (s, 1H), 7.89 (s, 2H), 7.82 (d, $J = 7.5$ Hz, 1H), 7.74 (t, $J = 8.0$ Hz, 1H), 7.17 (d, $J = 8.0$ Hz, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 190.4, 152.4, 148.2, 138.2, 136.7, 131.4, 129.7, 126.0, 125.6, 119.2.




3-(4-Nitro-phenyl)-3-[1, 2, 3] triazol-2-yl-propenal (2j) (Table 2.3, Entry 10)

The title compound was prepared according to the general procedure, as described above in 88 % yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.49 (d, $J = 8.0$ Hz, 1H), 8.39 (d, $J = 9.0$ Hz, 2H), 7.89 (s, 2H), 7.68 ((d, $J = 8.5$ Hz, 2H), 7.16 (d, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.4, 152.6, 149.2, 138.2, 136.0, 132.2, 123.7, 119.2.




3-(3-Oxo-1-[1, 2, 3] triazol-2-yl-propenyl)-benzonitrile (2k) (Table 2.3, Entry 11)

The title compound was prepared according to the general procedure, as described above in 90 % yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.48 (d, $J = 8.0$ Hz, 1H), 7.89 (s, 2H), 7.86 (d, $J = 7.5$ Hz, 1H), 7.78 (s, 1H), 7.72 (d, $J = 8.0$ Hz, 1H), 7.66 (t, $J = 7.5$ Hz, 1H), 7.14 (d, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.5, 152.6, 138.2, 135.1, 134.4, 134.2, 131.2, 129.5, 119.2, 117.5, 113.3.




4-(3-Oxo-1-[1, 2, 3] triazol-2-yl-propenyl)-benzonitrile (2l) (Table 2.3, Entry 12)

The title compound was prepared according to the general procedure, as described above in 92 % yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.47 (d, $J = 8.0$ Hz, 1H), 7.88 (s, 2H), 7.82 (d, $J = 8.0$ Hz, 2H), 7.61 (d, $J = 8.0$ Hz, 2H), 7.13 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.6, 153.0, 138.2, 132.2, 131.7, 129.2, 119.1, 117.7, 114.9.



3-[1, 2, 3] Triazol-2-yl-3-(3-trifluoromethyl-phenyl)-propenal (2m) (Table 2.3, Entry 13)

The title compound was prepared according to the general procedure, as described above in 93% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.48 (d, $J = 8.0$ Hz, 1H), 7.89 (s, 2H), 7.84~7.83 (m, 1H), 7.74 (s, 1H), 7.68 (d, $J = 4.5$ Hz, 2H), 7.13 (d, $J = 8.0$ Hz, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.1, 153.6, 138.1, 134.3, 130.7, 129.2, 127.8 ($J = 8.8$ Hz), 122.2 ($J = 50$ Hz), 119.1.




3-(2H-1,2,3-triazol-2-yl)oct-2-enal(2n) (Table 2.3, Entry 14)

The title compound was prepared according to the general procedure, as described above in 75% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.09 (d, $J = 7.5$ Hz, 1H), 7.85 (d, $J = 6.0$ Hz, 2H), 6.94 (d, $J = 7.5$ Hz, 1H), 3.30 (t, $J = 7.5$ Hz, 2H), 1.75~1.68 (m, 2H), 1.43~1.32 (m, 4H), 0.89 (t, $J = 7.0$ Hz, 3H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.1, 156.8, 137.3, 116.3, 31.4, 29.2, 27.5, 22.3, 13.8.

General procedure for Michael addition of 1*H*-1, 2, 3-triazole to substituted phenylpropynal to afford compound 3


To a solution of substituted phenylpropynal (0.08mmol), imidazolidine-2-carboxylic acid organocatalyst (3.52mg, 0.016mmol) and trifluoroacetic acid (1.23ul, 0.016mmol) in 1, 2-dichloroethane (0.15ml) was added 1*H*-1, 2, 3-triazole (5.33ul, 0.092mmol). The resulting solution was stirred at room temperature for 2 hours. Then the reaction mixture was directly purified by column chromatography, eluted with hexane/EtOAc to give the desired product.



3-Phenyl-3-[1, 2, 3] triazol-1-yl-propenal (3a) (Table 2.3, Entry 1)


The title compound was prepared according to the general procedure, as described above in 88% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.54 (d, $J = 8.0$ Hz, 1H), 7.76 (s, 1H), 7.57~7.53 (m, 2H), 7.47~7.44 (m, 4H), 7.01 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125

MHz): δ 191.4, 152.3, 134.5, 132.3, 130.6, 129.3, 127.7, 124.1, 120.2.




3-(4-Methoxy-phenyl)-3-[1, 2, 3] triazol-1-yl-propenal (3b) (Table 2.3, Entry 2)

The title compound was prepared according to the general procedure, as described above in 89% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.56 (d, $J = 8.0$ Hz, 1H), 7.76 (s, 1H), 7.53 (s, 1H), 7.36 (d, $J = 8.5$ Hz, 2H), 7.03 (d, $J = 8.5$ Hz, 2H), 6.87 (d, $J = 7.5$ Hz, 1H), 3.88 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.5, 162.5, 152.3, 134.4, 132.4, 126.3, 124.2, 119.8, 114.7, 55.6.




3-p-Tolyl-3-[1, 2, 3] triazol-1-yl-propenal (3c) (Table 2.3, Entry 3)

The title compound was prepared according to the general procedure, as described above in 87% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.55 (d, $J = 8.0$ Hz, 1H), 7.75 (d, $J = 0.5$ Hz, 1H), 7.49 (d, $J = 1$ Hz, 1H), 7.34 (d, $J = 8.5$ Hz, 2H), 7.32 (d, $J = 8.5$ Hz, 2H), 6.96 (d, $J = 7.5$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.5, 152.5, 142.6, 134.4, 130.6, 129.9, 127.6, 124.1, 120.0, 21.5.




3-(4-Bromo-phenyl)-3-[1, 2, 3] triazol-1-yl-propenal (3d) (Table 2.3, Entry 4)

The title compound was prepared according to the general procedure, as described above in 88% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.55 (d, $J = 8.0$ Hz, 1H), 7.78 (s, 1H), 7.70 (d, $J = 8.5$ Hz, 2H), 7.52 (s, 1H), 7.32 (d, $J = 8.0$ Hz, 2H), 6.95 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.6, 151.0, 134.6, 132.6, 132.0, 128.9, 126.7, 123.7, 120.4.




3-(4-Chloro-phenyl)-3-[1, 2, 3] triazol-1-yl-propenal (3e) (Table 2.3, Entry 5)

The title compound was prepared according to the general procedure, as described above in 90% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.55 (d, $J = 8.0$ Hz, 1H), 7.78 (s, 1H), 7.54 (d, $J = 8.0$ Hz, 2H), 7.52 (s, 1H), 7.39 (d, $J = 8.0$ Hz, 2H), 6.95 (d, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.7, 151.0, 138.5, 134.7, 131.9, 129.7, 128.8, 123.8, 120.5.




3-(4-Fluoro-phenyl)-3-[1,2,3]triazol-1-yl-propenal (3f) (Table 2.3, Entry 6)

The title compound was prepared according to the general procedure, as described above in 83% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.53 (d, $J = 7.5$ Hz, 1H), 7.77 (s, 1H), 7.51 (s, 1H), 7.46~7.43 (m, 2H), 7.26~7.22 (m, 2H), 6.93 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.8, 164.7 ($J = 253$ Hz), 151.2, 134.6, 132.8 ($J = 8.8$ Hz), 126.1, 123.8, 120.4, 116.7 ($J = 22$ Hz).




3-(3-Fluoro-phenyl)-3-[1, 2, 3] triazol-1-yl-propenal (3g) (Table 2.3, Entry 7)

The title compound was prepared according to the general procedure, as described above in 91% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.57(d, $J = 8.0\text{Hz}$, 1H), 7.80 (s, 1H), 7.60~7.55 (m, 1H), 7.54 (s, 1H), 7.37~7.34 (m, 1H), 7.30~7.27 (m, 1H), 7.19 (d, $J = 8.5\text{Hz}$, 1H), 7.01 (d, $J = 8.0\text{Hz}$, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.7, 162.7 ($J = 249$ Hz), 150.6, 134.6 ($J = 25\text{Hz}$), 131.2 ($J = 8\text{Hz}$), 126.5, 123.8, 120.5, 119.0 ($J = 21\text{Hz}$), 117.6 ($J = 23\text{Hz}$).




3-(2-Fluoro-phenyl)-3-[1, 2, 3] triazol-1-yl-propenal (3h) (Table 2.3, Entry 8)

The title compound was prepared according to the general procedure, as described above in 86% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.55 (d, $J = 7.5$ Hz, 1H), 7.80 (s, 1H), 7.68~7.64 (m, 1H), 7.58 (s, 1H), 7.49~7.47 (m, 1H), 7.38 (t, $J = 7.5$ Hz, 1H), 7.32~7.28 (m, 1H), 7.08 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.5, 160.2 ($J = 251$ Hz), 146.6, 134.6, 133.9 ($J = 8.1$ Hz), 132.4, 125.0 ($J = 15$ Hz), 123.1, 120.7, 116.8 ($J = 21$ Hz).




3-(3-Nitro-phenyl)-3-[1, 2, 3] triazol-1-yl-propenal (3i) (Table 2.3, Entry 9)

The title compound was prepared according to the general procedure, as described above in 89% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.54 (d, $J = 7.5$ Hz, 1H), 8.48 (d, $J = 7.5$ Hz, 1H), 8.34 (s, 1H), 7.84~7.77 (m, 3H), 7.65 (s, 1H), 6.95 (d, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.6, 149.6, 148.5, 136.2, 133.0, 131.6, 130.5, 126.4, 125.4, 123.4, 120.8.




3-(4-Nitro-phenyl)-3-[1,2,3]triazol-1-yl-propenal (3j) (Table 2.3, Entry 10)

The title compound was prepared according to the general procedure, as described above in 83% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.53 (d, $J = 7.5$ Hz, 1H), 8.40 (d, $J = 8.5$ Hz, 2H), 7.81 (s, 1H), 7.69 (d, $J = 8.5$ Hz, 2H), 7.62 (s, 1H), 6.96 (d, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.7, 149.7, 135.9, 135.0, 131.8, 128.6, 124.5, 124.3, 120.8.




3-(3-Oxo-1-[1,2,3] triazol-1-yl-propenyl)-benzonitrile (3k) (Table 2.3, Entry 11)

The title compound was prepared according to the general procedure, as described above in 79% yield. ^1H NMR ($\text{DMSO}-d^6$, 500 MHz): δ 9.42 (d, $J = 8.0$ Hz, 1H), 8.65 (d, $J = 1.0$ Hz, 1H), 8.21 (s, 1H), 8.11 (d, $J = 8.0$ Hz, 1H), 8.02 (d, $J = 1.0$ Hz, 1H), 7.92 (d, $J = 8.0$ Hz, 1H), 7.78 (t, $J = 8.0$ Hz, 1H), 6.95 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR ($\text{DMSO}-d^6$, 125 MHz): δ 191.3, 150.2, 135.5, 134.7, 134.6, 134.5, 131.9, 129.9, 125.1, 119.7, 118.0, 112.0.




4-(3-Oxo-1-[1, 2, 3] triazol-1-yl-propenyl)-benzonitrile (3l) (Table 2.3, Entry 12)

The title compound was prepared according to the general procedure, as described above in 80% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.52 (d, $J = 7.5$ Hz, 1H), 7.86 (d, $J = 8.0$ Hz, 2H), 7.81 (s, 1H), 7.60 (d, $J = 8.5$ Hz, 2H), 7.58 (s, 1H), 6.95 (d, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.8, 150.0, 134.9, 134.2, 132.8, 131.3, 123.4, 120.7, 117.3, 115.7.



3-[1, 2, 3] Triazol-1-yl-3-(3-trifluoromethyl-phenyl)-propenal (3m) (Table 2.3, Entry 13)



The title compound was prepared according to the general procedure, as described above in 82% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.52 (d, $J = 7.5$ Hz, 1H), 7.89 (d, $J = 7.5$ Hz, 1H), 7.80 (s, 1H), 7.72 (s, 2H), 7.67 (d, $J = 7.5$ Hz, 1H), 7.54 (s, 1H), 7.00 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.3, 150.5, 134.9, 133.8, 130.8 ($J = 9.3$ Hz), 130.0, 128.5, 127.2, 124.4, 123.6, 120.7.



3-(1H-1,2,3-triazol-1-yl)oct-2-enal(3n) (Table 2.3, Entry 14)

The title compound was prepared according to the general procedure, as described above in 67% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.11 (d, $J = 7.5$ Hz, 1H), 7.94 (s, 1H), 7.82 (s, 1H), 6.44 (d, $J = 7.5$ Hz, 1H), 3.32 (t, $J = 8.0$ Hz, 2H), 1.71~1.65 (m, 2H), 1.42~1.27 (m, 4H), 0.87 (t, $J = 7.5$ Hz, 3H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.5, 154.9, 134.8, 121.5, 117.7, 31.3, 28.8, 28.4, 22.2, 13.8.


Determination of Configuration:



The compound **4** was prepared according to the general procedure:

The obtained compound **3a** (39.8mg, 0.2mmol) in 2ml anhydrous MeOH at 0°C was treated with cerium (III) chloride heptahydrate (89.4mg, 0.24mmol) and sodium borohydride (9.1mg, 0.24mmol). After 1 hour, the reaction was quenched by the addition of saturated aqueous ammonium chloride solution (2ml), and extracted with dichloromethane (3 × 20ml). The combined organic layer was then washed once with brine (50ml), dried over anhydrous Na₂SO₄, filtered and concentrated on vacuo. Purification of the residue by column chromatography affords the two isomeric enols in 72% yield totally.

¹H NMR (CDCl₃, 500 MHz): δ 7.79 (s, 1H), 7.56 (s, 1H), 7.35 (t, *J* = 8.0 Hz, 3H), 7.14 (d, *J* = 7.0 Hz, 2H), 6.40 (t, *J* = 7.0 Hz, 1H), 4.11 (d, *J* = 7.0 Hz, 2H), 2.96 (br, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 137.5, 135.3, 133.7, 129.7, 128.9, 126.6, 126.5, 125.1, 57.9.



The title compound was prepared according to the general procedure, as described above.

¹H NMR (CDCl₃, 500 MHz): δ 7.67 (s, 1H), 7.43~7.41 (m, 3H), 7.39 (s, 1H), 7.25 (d, *J* = 6.0 Hz, 2H), 6.68 (d, *J* = 7.0 Hz, 1H), 4.30 (d, *J* = 7.5 Hz, 2H). ¹³C NMR (CDCl₃, 125 MHz): δ 137.2, 133.5, 132.4, 129.9, 129.3, 128.8, 123.7, 123.6, 58.8.

2.7 Reference

[1] For recent reviews on organocatalytic asymmetric Michael additions, see: (a) D. Almasi, D. A. Alonso, C. Najera, *Tetrahedron: Asymmetry*, **2007**, *18*, 299-365; (b) S. B. Tsogoeva, *Eur. J. Org. Chem.*, **2007**, *11*, 1701-1716; (c) D. Enders, C. Wang, J. X. Liebich, *Chem. Eur. J.* **2009**, *15*, 11058-11076; (d) J. L. Vicario, D. Badía, L. Carrillo, *Synthesis* **2007**, 2065-2092; (e) S. Sulzer-Mossé, A. Alexakis, *Chem. Commun.* **2007**, 3123-3135.

[2] For selected examples of organocatalytic asymmetric Michael additions to enals, see: (a) S. Brandau, A. Landa, J. Franzén, M. Marigo, K. A. Jørgensen, *Angew. Chem. Int. Ed.*, **2006**, *45*, 4305-4309; (b) H. Gotoh, H. Ishikawa, Y. Hayashi, *Org. Lett.*, **2007**, *9*, 5307-5309; (c) S. Bertelsen, P. Dinr, R. L. Johansen, K. A. Jørgensen, *J. Am. Chem. Soc.*, **2007**, *129*, 1536-1537; (d) P. Dinér, M. Nielsen, M. Marigo, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2007**, *46*, 1983-1987; (e) U. Uria, J. L. Vicario, D. Badía, L. Carrillo, *Chem. Commun.* **2007**, 2509-2511; (f) X.-W. Wang, B. List, *Angew. Chem. Int. Ed.* **2008**, *47*, 1119-1122; (g) S.-L. Zhang, Y.-N. Zhang, Y.-F. Ji, H. Li, W. Wang, *Chem. Commun.* **2009**, 4886-4888.

[3] There are few examples of substituted acetylenic carbonyls as Michael acceptors in asymmetric catalysis. For alkynones, see: (a) M. Bella, K. A. Jørgensen, *J. Am. Chem. Soc.* **2004**, *126*, 5672-5673; (b) Z.-H. Chen, M. Furutachi, Y. Kato, S. Matsunaga, M. Shibasaki, *Angew. Chem. Int. Ed.* **2009**, *48*, 2218-2220; For acetylenic esters, see: (c) X.

Wang, M. Kitamura, K. Maruoka, *J. Am. Chem. Soc.* **2007**, *129*, 1038-1039; For substituted acetylenic aldehydes and ketones in Diels–Alder reactions, see: (d) K. Ishihara, S. Kondo, H. Kurihara, H. Yamamoto, S. Ohashi, S. Inggaki, *J. Org. Chem.* **1997**, *62*, 3026-3027; (e) E. J. Corey, T.W. Lee, *Tetrahedron Lett.* **1997**, *38*, 5755-5758; (f) K. Ishihara, M. Fushimi, *J. Am. Chem. Soc.* **2008**, *130*, 7532-7533; (g) J. N. Payette, H. Yamamoto, *Angew. Chem. Int. Ed.* **2009**, *48*, 8060-8062; (h) S. B. Jones, B. Simmons, D.W. C. MacMillan, *J. Am. Chem. Soc.* **2009**, *131*, 13606-13607.

[4] N. L. Mazyar, V. V. Annenkov, V. A. Klugova, S. M. Anańev, E. N. Danilovtseva, A. V. Rokhin, S. V. Zinchenko, *Russ. Chem. Bull.*, **2000**, *109*, 2013-2017.

[5] Y.-A. Choi, K. Kim, Y.-J. Park, *Tetrahedron Lett.*, **2003**, *44*, 7507-7511.

[6] M. Ogata, H. Matsumhimizu, S. Kida, M. Shiro, K. Tawara, *J. Med. Chem.*, **1987**, *30*, 1348-1354.

[7] (a) G. Wittig, G. Geissler, *Justus Liebigs Ann. Chem.* **1953**, *580*, 44-57; (b) B. E. Maryanoff, A. B. Reitz, *Chem. Rev.* **1989**, *89*, 863-927.

[8] (a) W. S. Wadsworth, W. D. Emmons, *J. Am. Chem. Soc.* **1961**, *83*, 1733-1738; (b) J. Boutagy, R. Thomas, *Chem. Rev.* **1974**, *74*, 87-99; (c) W. S. Wadsworth, Jr. *Org. React.* **1977**, *25*, 73-253; (d) S. E. Kelly, In *Comprehensive Organic Synthesis*; B. M. Trost, I. Fleming, Ed.; Pergamon: Oxford, **1991**, *1*, 729-817. For applications in natural product synthesis, see: (e) K. C. Nicolaou, M. W. Härter, J. L. Gunzner, A. Nadin, *Liebigs Ann./Recueil* **1997**, *97*, 1283-1301.

[9] For selected examples of transition-metal mediated carbometallation of alkynes, see: (a) D. G. Hall, D. Chapdelaine, P. Prévile, P. Deslongchamps, *Synlett*, **1994**, 660-662; (b) E. Shirakawa, K. Yamasaki, H. Yoshida, T. Hiyama, *J. Am. Chem.Soc.* **1999**, *121*, 10221-10222 ; (c) Z. Lu, S.-M. Ma, *J. Org. Chem.* **2006**, *71*, 2655-2660; (d) D.-H. Zhang, J. M. Ready, *J. Am. Chem. Soc.* **2006**, *128*, 15050-15051.

[10] (a) E.-i Negishi, *In Metal-catalyzed Cross-coupling Reactions*, F. Diederich, P. J. Stang, Eds. Wiley-VCH: New York, **1998**, pp. 167-202; (b) A. Arefolov, N. F. Langille, J. S. Panek, *Org. Lett.* **2001**, *3*, 3281-3284. For other alkene stereocontrol related work, see also (c) C. McFarland, J. Hutchison, M. C. McIntosh, *Org. Lett.* **2005**, *7*, 3641-3644.

[11] In fact, in the cases with larger Michael donor, the ether moieties had great influence on the *E/Z* ratio. In order to prove this point, we investigated the Michael addition between phenyl-propynal **1a** and 5-phenyl-1*H*-tetrazole in DCM at r.t. in the presence of 20 mol% catalysts **I** and **II**. When changing from less bulkyTMS catalyst **I** to bulky TBS catalyst **II**, the*E/Z* ratio can be enhanced significantly from 1:1.08 to 1:0.32.

[12] P. Dinér, M. Nielsen, M. Marigo, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2007**, *46*, 1983-1987.

Chapter 3

Alkynyl Involved Organocatalytic Cascade Reactions


3.1 Background and Significance

Highly efficient constructions of complex organic molecules continue to receive paramount attention in modern synthetic realm. So far, organocatalytic cascade or tandem reactions have grown into one of the most robust and appealing synthetic methodologies which enable rapid assemble versatile biologically relevant compounds containing higher levels of molecular complexity with high enantiomeric purity in an economically and environmentally friendly manner¹. Of crucial importance is that the remarkable achievements definitely accelerate the development of pharmaceutical industry².


Chiral secondary amines are probably the most successful organocatalyst in organocatalytic asymmetric cascade reactions to date, as secondary amines are capable of both enamine and iminium catalysis³. The versatile possibilities of combination of these two catalytic activation modes in one protocol allows for rapid conversion of simple achiral starting materials into complex, highly enantioenriched products although their scope is mainly limited to carbonyl systems¹. Among them, the iminium-enamine combination has turned out to be very powerful. The enal substrate is first activated through iminium-ion formation, and then facilitates the nucleophilic conjugate addition. The resulting enamine can then undergo a second reaction with an electrophile to afford

the products, which in general contain two new stereocenters (Scheme 3.1). In 2004, Jørgensen and co-workers developed an organocatalytic Michael-aldol cascade for the assembly of optically active cyclohexanones⁴. Today, numerous examples of this combination can be found in the literature^{1a}.

Scheme 3.1 the Iminium-Enamine Combination



Scheme 3.2 Organocatalytic Michael-aldol Cascade for the Assembly of Optically Active Cyclohexanones




Besides the widely explored iminium-enamine cascade sequence, many other cascade reactions were achieved by simply changing the order of activation modes such as enamine-enamine activation, enamine-iminium activation^{1a,1f}.

The concept of organocatalytic cascade reaction is also extended to a more complex scenario later. The leading example is reported by Enders and co-workers about the three component triple cascade reaction for the stereocontrolled formation of tetrasubstituted cyclohexene carbaldehyde bearing up to four stereogenic centres⁵. The

reaction proceeds through a Michael-Michael-aldol condensation cascade which can be explained as enamine-iminium-enamine sequence (Scheme 3.3).

Scheme 3.3 Orgacatalytic Michael-Michael-aldol Triple Cascade Reaction



This strategy was substantially extended to quadruple cascade reactions such as Michael-Michael-Michael-aldol condensation reaction which go through an iminium-

enamine-iminium-enamine catalytic cycle. They enable the consecutive formation of four new bonds and provide efficient methods for construction of highly functionalized trisubstituted cyclohexene carbaldehydes⁶, spirooxindole motif⁷ and marine meroterpene (+)-conicol⁸.


3.2 Research Design

Recently, tremendous efforts have been directed to expand the scope of organocatalyzed cascade reactions triggered by Michael additions to enals which efficiently furnish valuable chiral building blocks via combined iminium-enamine activation strategy^{1a} (Figure 3.1, Eq.(1)) to control the enantioselectivity at the β -carbon atom of α , β -unsaturated aldehydes. Importantly, the geometry of the *trans*-enals controls the Michael-initiated cascade reactions to afford high enantio and/or diastereoselectivity. To the best of our knowledge, in contrast to these widely reported iminium-enamine catalyzed systems, the paradigms for organocatalytic cascade/tandem reactions involving conjugate additions to the corresponding alkynals in an asymmetric fashion have not yet been reported so far presumably owing to the fact that the alkynal substrates possess no prochiral center at the β -carbon atom. Recently, MacMillan and co-workers reported the use of propynal as the dienophile in an efficient iminium activated [4+2] cycloaddition reaction⁹.

To perform an enantioselective cascade reaction on an alkynal substrate, we envision that their use instead of an enal in the initial Michael addition to an iminium ion

formed *in situ* would afford an unprecedented chiral allenamine intermediate (Figure 3.1, Eq. (2))¹⁰⁻¹³. The allenamine formed would then act as the nucleophile in a subsequent enantioselective reaction with an electrophile to produce a new stereogenic center. In this context, the stereochemistry is governed by a poorly understood chiral allenamine¹¹⁻¹³ that is formed *in situ* in the second step of the cascade reaction, whilst in the classic iminium-enamine cascade processes, the enantioselectivity is generally controlled by the initial conjugate addition step.

Figure 3.1 Conventional Iminium-Enamine (1) and Novel Iminium-Allenamine (2) Activation Strategies in Organocatalytic Enantioselective Cascade Reactions.




3.3 Iminium–Allenamine Cascade Catalysis: One-Pot Access to Chiral 4*H*-Chromenes by a Highly Enantioselective Michael–Michael Sequence¹⁴

3.3.1 Introduction

4*H*-Chromenes are a core structural feature of an array of fascinating natural products that have intriguing biological activities (Figure 3.2). For example, rhodomyrtone and rhodomyrtosone B show potent antibiotic activities¹⁵, whereas the dimeric 4*H*-chromene acylphloroglucinol, myrtucommulone E, serves as a promising α -glucosidase inhibitor with antibacterial activity¹⁶. These frameworks have become attractive targets in organic synthesis^{17, 18}, and despite significant advances having been made, asymmetric methods for their construction are still extremely rare¹⁹.

Figure 3.2 Selected Examples of 4*H*-Chromenes as Substructures in Biologically Interesting Natural Products



3.3.2 Results and Discussion

Motivated by the above idea and in continuation of our previous research interest in organocatalytic asymmetric cascade reactions for the construction of chromanes or chromenes²⁰, we ventured into the development of the new double Michael additions between 2-(*E*)-(2-Nitrovinyl)-phenols and alkynals considering the nitroalkenes are among the most reactive Michael acceptors to explore the feasibility of the iminium–allenamine cascade catalysis.

We started our investigations by conducting the model reaction between phenylpropynal **1a** and 2-(*E*)-(2-Nitrovinyl)-phenol **2a** in dichloromethane at room temperature in the presence of 20 mol% organocatalyst **I** in view of the fact that chiral secondary amine, especially, diarylprolinol silyl ether organocatalysts are capable of both iminium and enamine catalysis in promoting enals cascade reactions²¹ (Figure 3.3 and Table 3.1). Encouragingly, the reaction proceeded to completion within 1 h to yield the desired product **3a** in an excellent yield (95%) and in an excellent ee (97%) (Table 3.1, entry 1).

This reaction shows that alkynals are more active than enals. Despite all that good results, we still made a careful optimization. A screen of other diarylprolinol silyl ether analogues **II–IV** revealed that the more bulky TBDMS catalyst **III** gave slightly higher enantioselectivities (Table 3.1, entries 2-4), although diamine **V**, reported by Barbas and Betancort, showed poor enantiocontrol (Table 3.1, entry 5)²². Accordingly, catalyst **III** was chosen for further optimization of the reaction conditions. A brief survey of different reaction solvents revealed toluene to be the most suitable for this procedure (Table 3.1, entries 6 and 7). The process displayed low dependence on the base additive (Table 3.1, entry 8). Lowering the reaction temperature to 0 °C and the catalyst loading to 15 mol% resulted in a higher ee (>99%) and higher yield (97%) without considerably prolonging the reaction time (Table 3.1, entry 9 and 10).

Figure 3.3 Secondary Amine Catalysts Screened





Table 3.1 Optimization Reaction Conditions for the Organocatalytic Asymmetric Cascade Oxa-Michael-Michael Reaction between Phenyl-propynal **1a** and 2-(*E*)-(2-Nitrovinyl)-phenol **2a**^a.




Entry	Catalyst	Additive	Solvent	<i>t</i> (h)	Yield (%) ^b	ee (%) ^c
1	I	None	CH ₂ Cl ₂	1	95	97
2	II	None	CH ₂ Cl ₂	1	93	98
3	III	None	CH ₂ Cl ₂	1	92	98
4	IV	None	CH ₂ Cl ₂	1	90	96
5	V	None	CH ₂ Cl ₂	1	91	66
6	III	None	Toluene	1	94	99
7	III	None	DCE	1	92	98
8	III	NaOAc	Toluene	1	92	99
9	III ^d	None	Toluene	5	88	99
10	III ^e	None	Toluene	4	97	>99

^a Reaction conditions: unless specified, a mixture of phenyl-propynal **1a** (0.11mmol), 2-(*E*)-(2-Nitrovinyl)-phenol **2a** (0.10mmol), catalyst (0.02mmol) and additive (0.02mmol) in solvent (0.8ml) was stirred at r.t. for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralpak AS-H). ^d 10 mol% of catalyst **III** was applied. ^e Performed at 0°C and 15 mol% of catalyst **III** was applied.

Having established the optimal conditions for this cascade oxa-Michael–Michael reaction, we then investigated the scope of this process. As shown in Table 3.2, the one-pot reaction promoted by organocatalyst **III** serves as a general and atom economical approach to “privileged” chiral 4*H*-chromenes, with formation of two C-C bonds, one new stereogenic center, and the incorporation of two versatile nitro and aldehyde functionalities that are available for further elaboration. Notably, in all cases, the reactions were completed in 4-5 hours, with excellent levels of enantioselectivity (98->99% ee) and in high yields (92-98%). Moreover, a broad substrate scope was observed.

Aromatic alkynals **1** that have electron-donating (Table 3.2, entries 2, 12, and 13) or electron withdrawing substituents (Table 3.2, entries 3-6 and 11) were investigated, and the effects of the substituent on the reaction was found to be very limited. These reactions proceeded very smoothly, affording excellent yields (92-98%) and excellent enantioselectivities (99->99%). A similar trend was observed with the structural variations of substrate **2** (Table 3.2, entries 7-9 and 11-13). Heteroaromatic alkyne thiophen-2-yl-propynal also effectively engaged in the cascade process (Table 3.2, entry 10). Finally, the reaction of the highly sterically hindered aliphatic alkyne also proceeded successfully, in 93% yield and 98% ee (Table 3.2, entry 14). The absolute configuration of product **3g** prepared under the optimal condition was determined to be *R* configuration by using single crystal X-ray diffraction (Figure 3.4).

Table 3.2 Catalyst **III** Promoted Asymmetric Cascade Oxa-Michael-Michael Reactions of Propynal **1** with 2-(*E*)-(2-Nitrovinyl)-phenol **2^a**.




Entry	R	X	<i>t</i> (h)	Yield (%) ^b	ee (%) ^c
1	Ph	H	4	97	>99
2	4-MeOC ₆ H ₄	H	5	93	99
3	4-ClC ₆ H ₄	H	4	97	99
4	4-BrC ₆ H ₄	H	4	95	99
5	4-NO ₂ C ₆ H ₄	H	4	95	99

6	3-NO ₂ C ₆ H ₄	H	4	93	99
7	Ph	5-Cl	4	92	99
8	Ph	3-MeO	4	95	98
9	Ph	5-MeO	4	97	99
10	Thienyl	H	4	92	99
11	4-BrC ₆ H ₄	5-MeO	4	98	99
12	4-MeC ₆ H ₄	5-Cl	4	94	99
13	4-MeC ₆ H ₄	5-MeO	4	98	99
14	<i>t</i> -Bu	H	4	93	98

^a Reaction conditions: unless specified, a mixture of propynal **1** (0.11mmol), 2-(*E*)-(2-Nitrovinyl)-phenol **2** (0.10mmol), catalyst **III** (0.015mmol) in Toluene (0.8ml) was stirred at 0°C for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralpak AS-H or IC).

Figure 3.4 the Single-Crystal X-ray Structure of Compound **3g**, with Ellipsoids Set at 20% Probability




3.3.3 Conclusion

In conclusion, we have developed a novel asymmetric oxa-Michael–Michael reaction involving an unprecedented chiral iminium–allenamine cascade. Ynals are used for the first time in organocatalyzed asymmetric cascade reactions. This process is a viable one-pot approach to synthetically and biologically significant chiral 4*H*-chromenes in high yields and with excellent enantioselectivities. A broad substrate scope has been successfully employed in this reaction, including aromatic and aliphatic alkynals as Michael acceptors, and 2-(*E*)-(2-nitro-vinyl)-phenols as Michael donors/acceptors with significant structural variation. The proposed iminium-allenamine activation mode has potential applications in the development of new organocatalytic enantioselective cascade reactions.

3.3.4 Supporting Information


General procedure for Cascade Michael-Michael addition reaction:

To a solution of alkynal (0.11mmol), organocatalyst **III** (5.51mg, 0.015mmol) in toluene (0.8ml) was added 2-(*E*)-(2-Nitrovinyl)-phenol (0.10mmol). The resulting solution was stirred at 0°C for a specified time. Then the reaction mixture was directly purified by column chromatography, eluted with hexane/EtOAc to afford the desired product.




(*R*)-4-Nitromethyl-2-phenyl-4*H*-chromene-3-carbaldehyde (3a) (Table 3.2, Entry 1):

The title compound was prepared according to the general procedure, as described above in 97% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.59 (s, 1H), 7.61 (d, $J = 7.5$ Hz, 2H), 7.58~7.56 (m, 1H), 7.51 (t, $J = 7.5$ Hz, 2H), 7.34~7.29 (m, 2H), 7.24~7.17 (m, 2H), 4.73~4.69 (m, 2H), 4.66~4.62 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.2, 169.8, 150.6, 131.7, 130.6, 130.3, 129.3, 128.6, 128.5, 125.9, 119.7, 117.1, 111.5, 79.6, 32.2; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 30/70, flow rate = 0.7 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 31.00$ min, $t_{\text{major}} = 21.57$ min, ee > 99%; $[\alpha]_{\text{D}}^{27} = +15.7$ (c = 1.0 in CHCl_3).



(R)-2-(4-Methoxy-phenyl)-4-nitromethyl-4H-chromene-3-carbaldehyde (3b) (Table 3.2, Entry 2):


The title compound was prepared according to the general procedure, as described above in 93% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.62 (s, 1H), 7.58 (d, $J = 8.5$ Hz, 2H), 7.36~7.29 (m, 2H), 7.23~7.19 (m, 2H), 7.02 (d, $J = 8.5$ Hz, 2H), 4.73~4.62 (m, 3H), 3.90 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.1, 169.7, 162.6, 150.7, 132.1, 129.2, 128.5, 125.8, 122.8, 120.1, 117.1, 114.1, 110.7, 79.7, 55.5, 32.3; HPLC (Chiralpak IC, *i*-PrOH/hexane = 30/70, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 104.07$ min, $t_{\text{major}} = 39.79$ min, ee = 99%; $[\alpha]_{\text{D}}^{27} = -16.0$ (c = 1.0 in CHCl_3).



(R)-2-(4-Chloro-phenyl)-4-nitromethyl-4H-chromene-3-carbaldehyde (3c) (Table 3.2,

Entry 3):

The title compound was prepared according to the general procedure, as described above in 97% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.59 (s, 1H), 7.57 (d, $J = 8.5$ Hz, 2H), 7.51 (d, $J = 8.5$ Hz, 2H), 7.37~7.31 (m, 2H), 7.26~7.18 (m, 2H), 4.76~4.64 (m, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.7, 168.5, 150.6, 138.2, 131.6, 129.4, 129.1, 128.5, 126.1, 119.6, 117.1, 111.8, 79.6, 32.3; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 30/70, flow rate = 0.7 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 34.53$ min, $t_{\text{major}} = 25.80$ min, ee = 99%; $[\alpha]_{\text{D}}^{27} = +14.0$ (c = 1.0 in CHCl_3).




(R)-2-(4-Bromo-phenyl)-4-nitromethyl-4H-chromene-3-carbaldehyde (3d) (Table 3.2,

Entry 4):


The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.59 (s, 1H), 7.68 (d, $J = 8.5$ Hz, 2H), 7.50 (d, $J = 8.0$ Hz, 2H), 7.36~7.31 (m, 2H), 7.26~7.23 (m, 1H), 7.18 (d, $J = 8.0$ Hz, 1H),

4.76~4.64 (m, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.7, 168.5, 150.5, 132.0, 131.7, 129.5, 129.4, 128.5, 126.6, 126.1, 119.5, 117.1, 111.9, 79.6, 32.3; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 30/70, flow rate = 0.7 mL/min, λ = 254nm): t_{minor} = 41.68 min, t_{major} = 26.58 min, ee = 99%; $[\alpha]_{\text{D}}^{27}$ = -6.9 (c = 0.86 in CHCl_3).




(R)-4-Nitromethyl-2-(4-nitro-phenyl)-4H-chromene-3-carbaldehyde (3e) (Table 3.2, Entry 5):

The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.59 (s, 1H), 8.39 (d, J = 8.5 Hz, 2H), 7.83 (d, J = 8.5 Hz, 2H), 7.39~7.34 (m, 2H), 7.28 (t, J = 7.5 Hz, 1H), 7.19 (d, J = 8.5 Hz, 1H), 4.83~4.79 (m, 1H), 4.72~4.67 (m, 2H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.0, 166.8, 150.4, 149.6, 136.6, 131.3, 129.6, 128.5, 126.4, 123.8, 119.1, 117.2, 113.1, 79.5, 32.3; HPLC (Chiralpak IC, *i*-PrOH/hexane = 40/60, flow rate = 0.7 mL/min, λ = 254nm): t_{minor} = 112.58 min, t_{major} = 93.90 min, ee = 99%; $[\alpha]_{\text{D}}^{27}$ = -5.6 (c = 1.0 in CHCl_3).



(R)-4-Nitromethyl-2-(3-nitro-phenyl)-4H-chromene-3-carbaldehyde (3f) (Table 3.2, Entry 6):


The title compound was prepared according to the general procedure, as described above in 93% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.60 (s, 1H), 8.51 (s, 1H), 8.46 (d, *J* = 8.5 Hz, 1H), 7.96 (d, *J* = 7.5 Hz, 1H), 7.76 (t, *J* = 8.0 Hz, 1H), 7.39~7.33 (m, 2H), 7.27 (t, *J* = 7.5 Hz, 1H), 7.21 (d, *J* = 8.0 Hz, 1H), 4.83~4.78 (m, 1H), 4.72~4.68 (m, 2H); ¹³C NMR (CDCl₃, 125 MHz): δ 189.0, 166.7, 150.4, 148.4, 136.2, 132.4, 129.9, 129.6, 128.5, 126.4, 126.2, 124.8, 119.1, 117.2, 112.9, 79.5, 32.3; HPLC (Chiralpak IC, *i*-PrOH/hexane = 40/60, flow rate = 0.7 mL/min, λ = 254nm): *t*_{minor} = 110.28 min, *t*_{major} = 87.38 min, ee = 99%; [α]_D²⁷ = +42.4 (c = 1.0 in CHCl₃).



(R)-6-Chloro-4-nitromethyl-2-phenyl-4H-chromene-3-carbaldehyde (3g) (Table 3.2, Entry 7):


The title compound was prepared according to the general procedure, as described above in 92% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.59 (s, 1H), 7.62~7.51 (m, 5H), 7.31~7.30 (m, 2H), 7.15~7.13 (m, 1H), 4.79~4.76 (m, 1H), 4.67~4.63 (m, 2H); ¹³C NMR (CDCl₃, 125 MHz): δ 190.1, 169.6, 149.3, 131.9, 130.9, 130.3, 129.5, 128.7, 128.2, 121.5, 118.6, 111.0, 79.3, 32.2; HPLC (Chiralpak IC, *i*-PrOH/hexane = 30/70, flow rate = 0.7 mL/min,

$\lambda = 254\text{nm}$): $t_{\text{minor}} = 27.23 \text{ min}$, $t_{\text{major}} = 22.94 \text{ min}$, $ee = 99\%$; $[\alpha]_{\text{D}}^{27} = +68.3$ ($c = 1.0$ in CHCl_3).




(R)-8-Methoxy-4-nitromethyl-2-phenyl-4H-chromene-3-carbaldehyde (3h) (Table 3.2, Entry 8):

The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.62 (s, 1H), 7.66 (d, $J = 8.0$ Hz, 2H), 7.60~7.57 (m, 1H), 7.52 (t, $J = 7.5$ Hz, 2H), 7.16 (t, $J = 8.0$ Hz, 1H), 6.93 (d, $J = 8.5$ Hz, 1H), 6.87 (d, $J = 7.5$ Hz, 1H), 4.74~4.62 (m, 3H), 3.91 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.2, 169.7, 148.3, 140.4, 131.8, 130.6, 128.7, 125.8, 121.0, 119.6, 111.6, 111.3, 79.6, 56.1, 32.3; HPLC (Chiralpak IC, *i*-PrOH/hexane = 40/60, flow rate = 0.7 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 49.63 \text{ min}$, $t_{\text{major}} = 19.57 \text{ min}$, $ee = 98\%$; $[\alpha]_{\text{D}}^{27} = +90.2$ ($c = 1.0$ in CHCl_3).



(R)-6-Methoxy-4-nitromethyl-2-phenyl-4H-chromene-3-carbaldehyde (3i) (Table 3.2, Entry 9):


The title compound was prepared according to the general procedure, as described above in 97% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.59 (s, 1H), 7.62~7.57 (m, 3H), 7.52 (t, *J* = 7.5 Hz, 2H), 7.13 (d, *J* = 9.0 Hz, 1H), 6.89~6.86 (m, 1H), 6.79 (s, 1H), 4.74~4.65 (m, 3H), 3.81 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 190.3, 170.0, 157.3, 144.7, 131.7, 130.8, 130.3, 128.6, 120.6, 118.1, 115.2, 112.4, 110.7, 79.5, 55.7, 32.6; HPLC (Chiralpak IC, *i*-PrOH/hexane = 40/60, flow rate = 0.7 mL/min, λ = 254nm): *t*_{minor} = 27.12 min, *t*_{major} = 21.48 min, ee = 99%; [α]_D²⁷ = +50.6 (c = 1.0 in CHCl₃).



(R)-4-Nitromethyl-2-thiophen-2-yl-4H-chromene-3-carbaldehyde (3j) (Table 3.2, Entry 10):

The title compound was prepared according to the general procedure, as described above in 92% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.90 (s, 1H), 7.70 (d, *J* = 5.0 Hz, 1H), 7.55 (d, *J* = 3.5 Hz, 1H), 7.37~7.34 (m, 1H), 7.30~7.20 (m, 4H), 4.73 (t, *J* = 5.5 Hz, 1H), 4.64 (d, *J* = 5.5 Hz, 2H); ¹³C NMR (CDCl₃, 125 MHz): δ 189.2, 163.0, 150.4, 133.6, 132.2, 131.7, 129.4, 128.5, 127.8, 126.0, 120.0, 117.1, 111.5, 79.5, 32.4; HPLC (Chiralpak IC, *i*-


PrOH/hexane = 30/70, flow rate = 0.7 mL/min, λ = 254nm): t_{minor} = 70.58 min, t_{major} = 31.78 min, ee = 99%; $[\alpha]_{\text{D}}^{27}$ = -8.8 (c = 1.0 in CHCl₃).



(R)-2-(4-Bromo-phenyl)-6-methoxy-4-nitromethyl-4H-chromene-3-carbaldehyde (3k)


(Table 3.2, Entry 11):

The title compound was prepared according to the general procedure, as described above in 98% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.57 (s, 1H), 7.66 (d, J = 8.5 Hz, 2H), 7.49 (d, J = 8.0 Hz, 2H), 7.12 (d, J = 9.0 Hz, 1H), 6.89~6.86 (m, 1H), 6.79 (d, J = 2.5 Hz, 1H), 4.76~4.71 (m, 1H), 4.68~4.64 (m, 2H), 3.81 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 189.7, 168.7, 157.4, 144.6, 132.0, 131.7, 129.7, 126.5, 120.4, 118.1, 115.3, 112.4, 111.0, 79.4, 55.8, 32.6; HPLC (Chiralpak IC, *i*-PrOH/hexane = 40/60, flow rate = 0.7 mL/min, λ = 254nm): t_{minor} = 31.79 min, t_{major} = 25.25 min, ee = 99%; $[\alpha]_{\text{D}}^{27}$ = +20.7 (c = 1.0 in CHCl₃).



(R)-6-Chloro-4-nitromethyl-2-p-tolyl-4H-chromene-3-carbaldehyde (3l) (Table 3.2, Entry 12):


The title compound was prepared according to the general procedure, as described above in 94% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.60 (s, 1H), 7.50 (d, *J* = 8.0 Hz, 2H), 7.34~7.29 (m, 4H), 7.15~7.13 (m, 1H), 4.77~4.72 (m, 1H), 4.66~4.62 (m, 2H), 2.46 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 190.1, 169.8, 149.3, 142.7, 130.8, 130.3, 129.4, 128.2, 127.4, 121.7, 118.5, 110.6, 79.3, 32.2, 21.6; HPLC (Chiralpak IC, *i*-PrOH/hexane = 30/70, flow rate = 0.7 mL/min, λ = 254nm): *t*_{minor} = 30.28 min, *t*_{major} = 25.69 min, ee = 99%; [α]_D²⁷ = +36.5 (c = 1.0 in CHCl₃).



(R)-6-Methoxy-4-nitromethyl-2-p-tolyl-4H-chromene-3-carbaldehyde (3m) (Table 3.2, Entry 13):

The title compound was prepared according to the general procedure, as described above in 98% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.59 (s, 1H), 7.51 (d, *J* = 8.0 Hz, 2H), 7.32 (d, *J* = 8.0 Hz, 2H), 7.13 (d, *J* = 8.5 Hz, 1H), 6.88~6.86 (m, 1H), 6.78 (d, *J* = 2.5 Hz, 1H), 4.71~4.63 (m, 3H), 3.81 (s, 3H), 2.45 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 190.3, 170.2, 157.2, 144.8, 142.4, 130.3, 129.3, 127.9, 120.8, 118.1, 115.2, 112.4, 110.4, 79.5, 55.7, 32.6, 21.6; HPLC (Chiralpak IC, *i*-PrOH/hexane = 30/70, flow rate = 0.7 mL/min, λ

= 254nm): $t_{\text{minor}} = 38.12$ min, $t_{\text{major}} = 29.81$ min, ee = 99%; $[\alpha]_{\text{D}}^{27} = +30.8$ (c = 1.0 in CHCl_3).



(R)-2-tert-Butyl-4-nitromethyl-4H-chromene-3-carbaldehyde (3n) (Table 3.2, Entry 14):

The title compound was prepared according to the general procedure, as described above in 93% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.38 (s, 1H), 7.31~7.28 (m, 1H), 7.24~7.22 (m, 1H), 7.19~7.16 (m, 1H), 7.10 (d, $J = 8.5$ Hz, 1H), 4.72~4.70 (m, 1H), 4.49~4.43 (m, 2H), 1.51 (s, 9H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 188.6, 178.5, 150.6, 129.0, 128.1, 125.7, 120.1, 116.5, 110.4, 79.9, 38.9, 32.5, 31.0; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 30/70, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 14.16$ min, $t_{\text{major}} = 12.08$ min, ee = 98%; $[\alpha]_{\text{D}}^{27} = -44.9$ (c = 1.0 in CHCl_3).


3.4 Development of Organocatalytic Asymmetric Cascade Double Michael Addition Reactions of 2-(2, 2-bis-benzenesulfonyl-vinyl)-phenol to Alkynals

3.4.1 Introduction

Sulfones are widely useful intermediates in synthetic and medicinal chemistry²³ due to the fact that the sulfonyl moiety can be removed by reduction²⁴ or transformed into

other useful functionalities²⁵. In spite of the great chemical versatility of the sulfonyl group, only recently have the sulfones begun to be incorporated systematically into the field of organocatalysis. In particular, the vinyl monosulfones or vinyl bis(sulfones)²⁶ have been used as very appealing Michael acceptors in organocatalytic enantioselective conjugate additions²⁷ due to the strong electron-withdrawing nature of the sulfonyl groups (Scheme 3.4).

Scheme 3.4 Selected Examples of Michael Additions to Vinyl Sulfones



Despite all of the aforementioned excellent advances, the employment of vinyl sulfones as one of the Michael acceptors in organocatalytic enantioselective cascade reactions remains an unexplored field and a challenging task. In continuation of our efforts on developing more general and variable strategies for construction of biologically relevant compounds, we envisioned that if the versatile sulfonyl group were incorporated into the compounds formed via enantioselective cascade reactions, a variety of analogues

can be readily available by taking advantage of the chameleonic ability of sulfones. Undoubtedly, the strategy has paramount importance on the rapid development of pharmaceutical industry.

Although the wonderful reaction scenario on the oxa-Michael-Michael sequence of 2-(*E*)-(2-nitrovinyl)-phenols with alkynals using iminium–allenamine cascade catalysis has been recognized by our group recently¹⁴, it is more fascinating to incorporate sulfonyl groups instead of nitro group into the “privileged” chiral 4*H*-chromene motifs in view of the fact that prior to reductive desulfonylation, miscellaneous electrophiles can be readily introduced to efficiently generate a great many of analogues.

3.4.2 Results and Discussion

To fulfill our expectation, we proposed to develop organocatalytic asymmetric cascade reactions between 2-(2, 2-bis-benzenesulfonyl-vinyl)-phenol and alkynalsto synthesize versatile biologically interesting 4*H*-chromene scaffolds.

We started our investigations by conducting the model reaction between phenylpropynal **1a** and 2-(2, 2-bis-benzenesulfonyl-vinyl)-phenol **4a** in dichloromethane at room temperature in the presence of 20 mol% organocatalyst **IV** (Figure 3.5 and Table 3.3). To our gratification, the initial trial gave encouraging result: the reaction between **1a** and **4a** finished within 2 h to yield the desired product **5a** with a moderate isolated yield (80%) and a moderate ee value (78%) (Table 3.3, entry 1). A brief investigation of reaction medium revealed that dichloromethane is the most suitable solvent for the

process (Table 3.3, entry 1-6). The screening of several other organocatalysts **I-VI** revealed that the bulky TBDMS catalyst **III** gave a slightly higher enantioselectivity and a slightly lower yield (Table 3.3, entry 9). However, the Barbas's catalyst **V** does not show enantioselectivity (Table 3.3, entry 10). The MacMillan catalyst **VI** shows poor enantioselectivity (Table 3.3, entry 11). Thereby, the catalyst **III** and dichloromethane solvent were chosen for further optimization reaction condition. Basic additives screen including both inorganic and organic base indicated that NaOAc additive lead to a slightly increased enantioselectivity (Table 3.3, entry 12-14). Further lowering the reaction temperature to -20°C and increasing the catalyst loading to 30 mol% resulted in better ee (92%) and yield (78%) (Table 3.3, entry 15).

Figure 3.5 Secondary Amine Catalysts Screened





Table 3.3 Optimization Reaction Conditions for the Organocatalytic Asymmetric Cascade Michael-Michael Reaction between Phenyl-propynal **1a** and 2-(2, 2-Bis-benzenesulfonyl-vinyl)-phenol **4a**^a.




Entry	Catalyst	Additive	Solvent	<i>t</i> (h)	Yield (%) ^b	ee (%) ^c
1	IV	None	DCM	2	80	78
2	IV	None	DCE	2	78	74
3	IV	None	Toluene	2	83	60
4	IV	None	CH ₃ CN	23	69	73
5	IV	None	Et ₂ O	6	71	69
6	IV	None	MeOH	17	62	78
7	I	None	CH ₂ Cl ₂	2	78	82
8	II	None	CH ₂ Cl ₂	2	76	83
9	III	None	CH ₂ Cl ₂	2	76	85
10	V	None	CH ₂ Cl ₂	2	77	0
11	VI	None	CH ₂ Cl ₂	20	68	50
12	III	NaOAc	DCM	1	77	86
13	III	Na ₂ CO ₃	DCM	1	75	79
14	III	Et ₃ N	DCM	1	76	84
15	III ^{d,e}	NaOAc	DCM	48	78	92

^a Reaction conditions: unless specified, a mixture of phenyl-propynal **1a** (0.055mmol), 2-(2, 2-bis-benzenesulfonyl-vinyl)-phenol **4a** (0.050mmol), catalyst (0.01mmol) and additive (0.025mmol) in solvent (0.4mL) was stirred at r.t. for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralpak OD-H). ^d Performed at -20°C. ^e 30 mol% of catalyst **III** was applied.

With the optimal reaction condition in hands, we examined the scope of the cascade oxa-Michael-Michael process with the results summarized in Table 3.4. As shown, the reaction has rather broad applicability. Aromatic alkynal **1**, regardless of neutral substituent (entries 1, 10), electron-donating substituents (entries 2, 3) and electron-withdrawing substituents (entries 4-9), participated in the process in

moderate yield (60%~83%) and with excellent enantioselectivities (87%~94%). On the other hand, 2-(2, 2-bis-benzenesulfonyl-vinyl)-phenol bearing electron-withdrawing group is tolerated although relatively low yield is observed (entry 10). Furthermore, low yield and enantioselectivity were observed for heteroaromatic alkynal such as thiophen-2-yl-propynal (entry 11). It is worthy of point out that the highly branched aliphatic alkynal was compatible with this methodology (entry 12). The absolute configuration of **5a** prepared under the optimal condition was determined by X-ray crystallography to be *S* configuration (Figure 3.6).

Table 3.4 Catalyst **III** Promoted Asymmetric Cascade Michael-Michael Reactions of Propynal **1** with 2-(2, 2-Bis-benzenesulfonyl-vinyl)-phenol **4**^a.




Entry	R	X	<i>t</i> (h)	5	Yield (%) ^b	ee (%) ^c
1	Ph	H	48	5a	78	92
2 ^d	4-MeOC ₆ H ₄	H	70	5b	63	89
3	4-MeC ₆ H ₄	H	70	5c	68	92
4	4-BrC ₆ H ₄	H	46	5d	82	94
5	4-ClC ₆ H ₄	H	46	5e	81	92
6	2-FC ₆ H ₄	H	46	5f	83	91
7	3-FC ₆ H ₄	H	46	5g	78	92
8	3-CF ₃ C ₆ H ₄	H	46	5h	80	91
9	3-CNC ₆ H ₄	H	46	5i	77	90

10 ^d	Ph	3-Cl	50	5j	60	87
11 ^d	Thienyl	H	48	5k	65	77
12	<i>t</i> -Bu	H	47	5l	83	88

^a Reaction conditions: unless specified, a mixture of propynal **1** (0.055mmol), 2-(2, 2-bis-benzenesulfonyl-vinyl)-phenol **4** (0.050mmol), NaOAc (0.025mmol) and catalyst **III** (0.015mmol) in DCM (1.0ml) was stirred at -20 °C for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralcel OD-H or Chiralpak AS-H). ^d Reaction was performed at -13°C.


Figure 3.6 the Single-Crystal X-ray Structure of Compound **5a**



To demonstrate the synthetic utility of the obtained enantioenriched *4H*-chromene motifs **5**, related post-modifications were carried out (Scheme 3.5). In general, the reduction of the aldehyde functionality may be necessary for minimizing the side reactions of the reductive desulfonylation step. Treatment of adduct **5a-1** with magnesium/methanol/dichloromethane gave the desired desulfonated product **5a-2** bearing one chiral methyl group in moderate yield, unfortunately, the enantioselectivity was seriously dropped from 92% to 17% (Scheme 3.5, eq (1)). Since fluorine-containing

compounds usually widely exist in pharmaceutical and agrochemical products²⁸, the efficient and selective incorporation of fluorine-containing moieties into organic compounds to modulate their biological properties has become a powerful strategy in drug design. Considering the significance of fluorine in drug discovery, we incorporated the fluorine atom in compound **5a**, in combination with subsequent reduction and desulfonylation, giving access to optically active monofluoromethylation analogue **5a-5** without erosion of enantioselectivity (Scheme 3.5, eq (2)).

Scheme 3.5 Synthetic Utility of Adduct **5**




3.4.3 Conclusion

In summary, we have disclosed the first organocatalytic enantioselective oxa-Michael-Michael cascade reaction applying vinyl sulfones as one of the Michael acceptors. This reaction proceeded with moderate yield and excellent enantioselectivity, as well as broad compatibility. The described approach together with facile post-modifications represents an efficient strategy to access versatile enantioenriched 4*H*-chromene motifs although further investigations on the utilities of the adduct are needed.

3.4.4 Supporting Information


General procedure for the synthesis of 2-(2,2-bis(phenylsulfonyl)vinyl)phenol:



To a solution of 2-acetoxybenzaldehyde (1.11g, 6.75 mmol), Bis(phenylsulfonyl)methane (1g, 3.37 mmol) and sodium fluoride (21.3mg, 0.51 mmol) in toluene (20ml) was added diethylamine hydrochloride (0.74g, 6.75 mmol). The resulting solution was refluxed with Dean-Stark apparatus for 3days. Then evaporate the toluene and the resulting crude solution was purified directly by running column chromatography to give the product with 45% isolated yield.

General procedure for Cascade Oxa-Michael-Michael addition reaction (Table 3.4):


To a solution of alkynal **1** (0.055 mmol), organocatalyst **III** (5.51mg, 0.015 mmol) and sodium acetate (2.05mg, 0.025 mmol) in dichloromethane (0.8ml) was added 2-(2,2-bis(phenylsulfonyl)vinyl)phenol **4** (0.05 mmol). The resulting solution was stirred at -20°C for a specific time. Then the reaction mixture was directly purified by column chromatography, eluted with hexane/EtOAc to afford the desired product.



(S)-4-(bis(phenylsulfonyl)methyl)-2-phenyl-4H-chromene-3-carbaldehyde (5a)

(Table 3.4, Entry 1):


The title compound was prepared according to the general procedure, as described above in 78% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.56 (s, 1H), 8.10~8.09 (m, 2H), 7.85 (d, J = 8.0 Hz, 1H), 7.71~7.55 (m, 10H), 7.48 (t, J = 7.5 Hz, 1H), 7.31~7.24 (m, 3H), 7.16~7.10 (m, 2H), 5.82 (s, 1H), 5.46 (d, J = 1.5 Hz, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.0, 170.7, 151.0, 140.5, 138.6, 134.0, 133.8, 131.7, 130.9, 130.8, 130.6, 129.7, 129.5, 129.1, 128.55, 128.50, 128.3, 125.6, 117.1, 116.8, 112.5, 85.5, 31.7. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, λ = 254nm): t_{minor} = 29.19 min, t_{major} = 23.12 min, ee = 92%; $[\alpha]_{\text{D}}^{25}$ = -80.6 (c = 1.0 in CHCl_3).



(S)-4-(bis(phenylsulfonyl)methyl)-2-(4-methoxyphenyl)-4H-chromene-3-carbaldehyde (5b) (Table 3.4, Entry 2):


The title compound was prepared according to the general procedure, as described above in 63% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.51 (s, 1H), 8.06 (d, J = 8.0 Hz, 2H), 7.78 (d, J = 7.5 Hz, 1H), 7.65 (t, J = 7.5 Hz, 1H), 7.59 (d, J = 8.5 Hz, 4H), 7.55 (t, J = 8.0 Hz,

2H), 7.42 (t, $J = 7.5$ Hz, 1H), 7.24~7.18 (m, 3H), 7.09 (d, $J = 8.5$ Hz, 1H), 7.04 (t, $J = 7.5$ Hz, 1H), 7.00 (d, $J = 8.5$ Hz, 2H), 5.74 (s, 1H), 5.41 (s, 1H), 3.89 (s, 3H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.8, 170.6, 162.5, 150.9, 140.6, 138.6, 134.0, 133.8, 132.4, 130.7, 129.7, 129.4, 129.1, 128.6, 128.3, 125.5, 123.0, 117.1, 117.0, 114.0, 111.6, 85.7, 55.5, 31.8. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.7 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 60.09$ min, $t_{\text{major}} = 28.98$ min, ee = 89%; $[\alpha]_{\text{D}}^{25} = -28.1$ ($c = 1.0$ in CHCl_3).




(S)-4-(bis(phenylsulfonyl)methyl)-2-p-tolyl-4H-chromene-3-carbaldehyde (5c) (Table 3.4, Entry 3):

The title compound was prepared according to the general procedure, as described above in 68% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.51 (s, 1H), 8.06~8.05 (m, 2H), 7.79 (d, $J = 7.5$ Hz, 1H), 7.65 (t, $J = 7.5\text{Hz}$, 1H), 7.61~7.59 (m, 2H), 7.57~7.53 (m, 4H), 7.42 (t, $J = 7.5\text{Hz}$, 1H), 7.31 (d, $J = 7.5$ Hz, 2H), 7.24~7.19 (m, 3H), 7.09~7.05 (m, 2H), 5.75 (s, 1H), 5.41 (d, $J = 1.5$ Hz, 1H), 2.45 (s, 3H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.0, 170.9, 151.0, 142.4, 140.5, 138.6, 134.0, 133.8, 130.8, 130.6, 129.7, 129.5, 129.3, 129.1, 128.6, 128.3, 128.0, 125.5, 117.1, 116.8, 112.1, 85.6, 31.7, 21.6. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 37.56$ min, $t_{\text{major}} = 20.30$ min, ee = 92%; $[\alpha]_{\text{D}}^{25} = -22.9$ ($c = 1.0$ in CHCl_3).



**(S)-4-(bis(phenylsulfonyl)methyl)-2-(4-bromophenyl)-4H-chromene-3-carbaldehyde
(5d) (Table 3.4, Entry 4):**


The title compound was prepared according to the general procedure, as described above in 82% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.50 (s, 1H), 8.02 (d, $J = 8.0$ Hz, 2H), 7.79 (d, $J = 7.5$ Hz, 1H), 7.67~7.63 (m, 3H), 7.59 (d, $J = 8.5$ Hz, 2H), 7.56~7.52 (m, 4H), 7.43 (t, $J = 7.5$ Hz, 1H), 7.23~7.19 (m, 3H), 7.09~7.05 (m, 2H), 5.75 (s, 1H), 5.37 (s, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.4, 169.5, 150.9, 140.5, 138.6, 134.0, 133.9, 132.0, 131.9, 130.8, 129.7, 129.6, 129.1, 128.5, 128.4, 126.5, 125.8, 117.1, 116.7, 112.9, 85.5, 31.7. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 72.60$ min, $t_{\text{major}} = 25.63$ min, ee = 94%; $[\alpha]_{\text{D}}^{26} = -45.0$ (c = 1.0 in CHCl_3).



**(S)-4-(bis(phenylsulfonyl)methyl)-2-(4-chlorophenyl)-4H-chromene-3-carbaldehyde
(5e) (Table 3.4, Entry 5):**


The title compound was prepared according to the general procedure, as described above in 81% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.50 (s, 1H), 8.01 (d, $J = 7.5$ Hz, 2H), 7.79

(d, $J = 8.0$ Hz, 1H), 7.65 (t, $J = 7.5$ Hz, 1H), 7.61~7.58 (m, 4H), 7.54 (t, $J = 8.0$ Hz, 2H), 7.50 (d, $J = 8.0$ Hz, 2H), 7.43 (t, $J = 7.5$ Hz, 1H), 7.27~7.24 (m, 1H), 7.21 (t, $J = 8.0$ Hz, 2H), 7.10~7.05 (m, 2H), 5.76 (s, 1H), 5.37 (d, $J = 1.5$ Hz, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.4, 169.4, 150.9, 140.4, 138.5, 138.1, 134.0, 133.9, 131.8, 130.8, 129.7, 129.6, 129.3, 129.1, 128.9, 128.43, 128.35, 125.8, 117.1, 116.7, 112.8, 85.4, 31.7. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 63.73$ min, $t_{\text{major}} = 24.12$ min, ee = 92%; $[\alpha]_{\text{D}}^{26} = -55.5$ (c = 1.0 in CHCl_3).



**(S)-4-(bis(phenylsulfonyl)methyl)-2-(2-fluorophenyl)-4H-chromene-3-carbaldehyde
(5f) (Table 3.4, Entry 6):**


The title compound was prepared according to the general procedure, as described above in 83% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.42 (s, 1H), 7.98 (d, $J = 8.0$ Hz, 2H), 7.83 (d, $J = 7.5$ Hz, 1H), 7.66~7.61 (m, 4H), 7.57~7.50 (m, 3H), 7.45 (t, $J = 7.0$ Hz, 1H), 7.30 (t, $J = 7.5$ Hz, 2H), 7.27~7.21 (m, 3H), 7.10~7.07 (m, 2H), 5.73 (s, 1H), 5.31 (s, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.4, 161.5, 159.4, 151.2, 140.2, 138.8, 134.1, 133.9, 133.4, 133.3, 132.7, 130.8, 129.7, 129.1, 128.5, 128.4, 125.7, 124.4, 119.2, 119.1, 117.2, 116.5, 116.3, 114.2, 85.4, 31.7. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 29.83$ min, $t_{\text{major}} = 21.14$ min, ee = 91%; $[\alpha]_{\text{D}}^{26} = -70.0$ (c = 1.0 in CHCl_3).



(S)-4-(bis(phenylsulfonyl)methyl)-2-(3-fluorophenyl)-4H-chromene-3-carbaldehyde


(5g) (Table 3.4, Entry 7):

The title compound was prepared according to the general procedure, as described above in 78% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.53 (s, 1H), 8.01 (d, $J = 8.0$ Hz, 2H), 7.79 (d, $J = 7.5$ Hz, 1H), 7.65 (t, $J = 7.0$ Hz, 1H), 7.59 (d, $J = 8.0$ Hz, 2H), 7.54 (t, $J = 8.0$ Hz, 2H), 7.50~7.47 (m, 1H), 7.45~7.42 (m, 2H), 7.37 (d, $J = 7.5$ Hz, 1H), 7.30~7.25 (m, 2H), 7.21 (t, $J = 8.0$ Hz, 2H), 7.11~7.06 (m, 2H), 5.76 (s, 1H), 5.37 (s, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.5, 169.0, 163.4, 161.4, 150.9, 140.4, 138.5, 134.0, 133.9, 132.82, 132.76, 130.7, 130.3, 130.2, 129.7, 129.1, 128.41, 128.38, 126.6, 125.8, 118.8, 118.7, 117.4, 117.2, 117.1, 116.6, 113.0, 85.4, 31.6. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 37.31$ min, $t_{\text{major}} = 22.88$ min, ee = 92%; $[\alpha]_{\text{D}}^{26} = -47.5$ (c = 1.0 in CHCl_3).




(S)-4-(bis(phenylsulfonyl)methyl)-2-(3-(trifluoromethyl)phenyl)-4H-chromene-3-carbaldehyde (5h) (Table 3.4, Entry 8):

The title compound was prepared according to the general procedure, as described above in 80% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.48 (s, 1H), 8.02 (d, $J = 8.0$ Hz, 2H), 7.87~7.84 (m, 3H), 7.80 (d, $J = 7.5$ Hz, 1H), 7.68~7.64 (m, 2H), 7.59 (d, $J = 8.0$ Hz, 2H), 7.54 (t, $J = 8.0$ Hz, 2H), 7.44 (t, $J = 7.5$ Hz, 1H), 7.29 (d, $J = 7.5$ Hz, 1H), 7.22 (t, $J = 8.0$ Hz, 2H), 7.13 (d, $J = 8.5$ Hz, 1H), 7.09 (t, $J = 7.5$ Hz, 1H), 5.79 (s, 1H), 5.36 (s, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.2, 168.8, 150.9, 140.4, 138.5, 134.1, 134.0, 131.7, 131.5, 131.2, 130.8, 129.7, 129.3, 129.1, 128.4, 128.3, 126.9, 125.9, 117.2, 116.6, 113.4, 85.4, 31.6. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 25.84$ min, $t_{\text{major}} = 19.40$ min, ee = 91%; $[\alpha]_{\text{D}}^{27} = -79.1$ ($c = 1.0$ in CHCl_3).



(S)-3-(4-(bis(phenylsulfonyl)methyl)-3-formyl-4H-chromen-2-yl)benzonitrile (5i) (Table 3.4, Entry 9):

The title compound was prepared according to the general procedure, as described above in 77% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.47 (s, 1H), 7.97~7.94 (m, 3H), 7.91 (d, *J* = 8.0 Hz, 1H), 7.87 (d, *J* = 7.5 Hz, 1H), 7.80 (d, *J* = 8.0 Hz, 1H), 7.68~7.62 (m, 2H), 7.58 (d, *J* = 8.0 Hz, 2H), 7.52 (t, *J* = 8.0 Hz, 2H), 7.45 (t, *J* = 7.5 Hz, 1H), 7.29 (t, *J* = 7.5 Hz, 1H), 7.24~7.21 (m, 2H), 7.13~7.09 (m, 2H), 5.78 (s, 1H), 5.33 (s, 1H). ¹³C NMR (CDCl₃, 125 MHz): δ 189.7, 168.0, 150.8, 140.3, 138.4, 134.8, 134.7, 134.1, 134.0, 133.7, 132.3, 130.8, 129.8, 129.7, 129.6, 129.1, 128.4, 128.3, 126.1, 117.6, 117.1, 116.5, 113.7, 113.3, 85.2, 31.6. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.7 mL/min, λ = 254nm): *t*_{minor} = 99.49 min, *t*_{major} = 58.73 min, ee = 90%; [α]_D²⁷ = -82.6 (c = 1.0 in CHCl₃).




(S)-4-(bis(phenylsulfonyl)methyl)-6-chloro-2-phenyl-4H-chromene-3-carbaldehyde

(5j) (Table 3.4, Entry 10):

The title compound was prepared according to the general procedure, as described above in 60% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.50 (s, 1H), 8.02 (d, *J* = 8.0 Hz, 2H), 7.78 (s, 1H), 7.65~7.63 (m, 5H), 7.60~7.47 (m, 6H), 7.28~7.25 (m, 2H), 7.20 (d, *J* = 8.5 Hz, 1H), 7.03 (d, *J* = 9.0 Hz, 1H), 5.69 (s, 1H), 5.38 (s, 1H). ¹³C NMR (CDCl₃, 125 MHz): δ 190.8, 170.4, 149.6, 140.2, 138.5, 134.1, 131.9, 130.7, 130.6, 130.5, 129.7, 129.4, 129.2, 128.9, 128.6, 128.5, 128.4, 118.4, 112.1, 85.3, 31.6. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, λ = 254nm): *t*_{minor} = 51.19 min, *t*_{major} =


21.46 min, ee = 87%; $[\alpha]_D^{27} = -110.4$ (c = 1.0 in CHCl_3).



(S)-4-(bis(phenylsulfonyl)methyl)-2-(thiophen-2-yl)-4H-chromene-3-carbaldehyde

(5k) (Table 3.4, Entry 11):


The title compound was prepared according to the general procedure, as described above in 65% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.78 (s, 1H), 8.04 (d, $J = 7.5$ Hz, 2H), 7.80 (d, $J = 7.5$ Hz, 1H), 7.69~7.68 (m, 1H), 7.66 (t, $J = 7.5$ Hz, 1H), 7.59~7.54 (m, 4H), 7.47~7.46 (m, 1H), 7.42 (t, $J = 7.5$ Hz, 1H), 7.28 (d, $J = 7.5$ Hz, 1H), 7.21 (t, $J = 7.5$ Hz, 2H), 7.18~7.16 (m, 1H), 7.13 (d, $J = 8.0$ Hz, 1H), 7.08 (t, $J = 7.5$ Hz, 1H), 5.73 (s, 1H), 5.37 (d, $J = 2.0$ Hz, 1H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 190.0, 164.0, 150.7, 140.4, 138.8, 134.2, 134.1, 133.9, 131.8, 130.7, 129.5, 129.4, 129.1, 128.9, 128.6, 128.4, 127.6, 125.7, 117.1, 116.8, 112.3, 85.5, 31.9. HPLC (Chiralcel OD-H, *i*-PrOH/hexane = 20/80, flow rate = 0.7 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 49.19$ min, $t_{\text{major}} = 30.01$ min, ee = 77%; $[\alpha]_D^{27} = -42.3$ (c = 1.0 in CHCl_3).



(S)-4-(bis(phenylsulfonyl)methyl)-2-tert-butyl-4H-chromene-3-carbaldehyde (5l)

(Table 3.4, Entry 12):

The title compound was prepared according to the general procedure, as described above in 83% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.3 (s, 1H), 7.90 (d, $J = 7.5$ Hz, 2H), 7.77 (d, $J = 8.0$ Hz, 1H), 7.60 (t, $J = 7.5$ Hz, 1H), 7.54 (d, $J = 7.5$ Hz, 2H), 7.49~7.43 (m, 3H), 7.27~7.21 (m, 3H), 7.09~7.05 (m, 2H), 5.68 (s, 1H), 5.05 (d, $J = 1.0$ Hz, 1H), 1.54 (s, 9H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.2, 180.1, 150.9, 140.4, 139.0, 133.83, 133.75, 130.3, 129.6, 129.3, 129.0, 128.3, 125.5, 116.9, 116.7, 110.9, 86.1, 38.9, 32.0, 31.1. HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 14.66$ min, $t_{\text{major}} = 18.88$ min, ee = 88%; $[\alpha]_{\text{D}}^{27} = -83.0$ ($c = 1.0$ in CHCl_3).




(S)-(4-methyl-2-phenyl-4H-chromen-3-yl)methanol (5a-2)

A round-bottom flask is charged with 100 mg **5a** (0.188 mmol, 92 % e.e.) in methanol (5 ml) and dichloromethane (1 ml). To the well-stirred solution at 0°C is added $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (140.4 mg, 0.377 mmol), then sodium borohydride (14.3 mg, 0.377 mmol). The reaction mixture is stirred at 0°C for 1h and monitored by TLC. After the reaction finish, 20 ml of saturated ammonium chloride aqueous solution is added at 0°C to quench the reaction. Evaporating solvent and the resulting residue is dissolved in dichloromethane. Then it is transferred to a separatory funnel. The aqueous phase is separated and extracted with dichloromethane (3x10ml). The organic layers are combined and washed with brine,

dried with MgSO_4 , filtered, and concentrated. The crude product is purified by column chromatography to give the alcohol **5a-1** in quantitative yield.

Compound **5a-1** (100 mg, 0.188 mmol) was dissolved in 5 mL MeOH. Activated Mg (prepared from Mg and $\text{BrCH}_2\text{CH}_2\text{Br}$ in Et_2O) (90 mg, 3.76 mmol) was added to this solution and the mixture was stirred at room temperature for 2 h. HCl (25 mL, 1 mol/L) was added and extracted with EtOAc for three times. The combined organic layers were then washed with water and brine, dried (Na_2SO_4), and concentrated. The residue was then purified by column chromatography to give 19.4 mg the desired product **5a-2** in 41% yield and 17% e.e. ^1H NMR (CDCl_3 , 500 MHz): δ 7.50 (d, $J = 7.5$ Hz, 2H), 7.42~7.41 (m, 3H), 7.21~7.16 (m, 2H), 7.08 (t, $J = 7.5$ Hz, 1H), 7.00 (d, $J = 7.5$ Hz, 1H), 4.32~4.25 (m, 2H), 3.80~3.76 (m, 1H), 1.46 (d, $J = 6.5$ Hz, 3H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 151.4, 148.8, 134.2, 128.9, 128.2, 128.1, 127.2, 127.0, 123.6, 116.1, 113.9, 61.6, 31.4, 24.4.



(S)-4-(fluoromethyl)-2-phenyl-4H-chromen-3-ylmethanol (**5a-5**)

A round-bottom flask is charged with 100 mg **5a** (0.188 mmol, 92 % e.e.) in anhydrous DMF (3 ml). To the well-stirred solution is added $t\text{-BuOK}$ (63.4 mg, 0.564 mmol) in one portion, then stir at r.t. for 20 min. Selectflour (200.3 mg, 0.564 mmol) in 2 ml anhydrous DMF is added dropwise. The reaction mixture is stirred at r.t. for 8h and monitored by TLC. After the reaction finish, water and EtOAc are added to extract the reaction solution.

The aqueous layer is separated and extracted with EtOAc for three times. The combined organic layers were then washed with water and brine, dried (Na₂SO₄), and concentrated. The residue was then purified by column chromatography to give the desired product **5a-3** in 92% yield. The compound **5a-4** and **5a-5** are prepared according to the general procedure described for **5a-1** and **5a-2**.

For compound **5a-5**, ¹H NMR (CDCl₃, 500 MHz): δ 7.60~7.59 (m, 2H), 7.44~7.43 (m, 3H), 7.29~7.25 (m, 2H), 7.14~7.09 (m, 2H), 4.66~4.48 (m, 2H), 4.34 (d, *J* = 12 Hz, 1H), 4.23 (d, *J* = 12 Hz, 1H), 4.01~3.95 (m, 1H), 1.91 (s, 1H); ¹³C NMR (CDCl₃, 125 MHz): δ 152.2, 151.6, 133.7, 129.4, 129.0, 128.8, 128.3, 128.2, 123.8, 120.8, 120.7, 116.3, 108.8, 87.5, 86.1, 62.6, 39.8, 39.6. HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 20/80, flow rate = 0.5 mL/min, λ = 254nm): t_{minor} = 15.28 min, t_{major} = 11.38 min, ee = 93%; [α]_D²⁸ = -13.7 (c = 1.0 in CHCl₃).

3.5 “One-Pot” Access to 4*H*-Chromenes with Formation of a Chiral Quaternary Stereogenic Center by a Highly Enantioselective Iminium-allenamine Involved Oxa-Michael Aldol Cascade²⁹

3.5.1 Introduction

Quaternary stereogenic centers are essential structural unit in many biologically active natural products and pharmaceuticals³⁰. The catalytic enantioselective construction of complex molecules containing quaternary stereocenters still pose a demanding

challenge for synthetic organic chemists although significant improvements have been achieved in this area³¹. Notably, organocatalytic formation of quaternary stereocenters^{30e,31g} with impressive levels of stereocontrol has received special attention over the past decade by virtue of relative simple and mild catalytic system. Among the developed organocatalytic methods suitable for this purpose, the chiral functionalized tertiary alcohols³² which are known to be important building blocks of bioactive natural products and drugs are commonly obtained by the direct aldol reactions of prochiral ketones³³.


4H-chromenes are an important class of core structures featured in a number of naturally occurring and biologically active molecules^{17, 34}. The ability to access the complex members of the chiral *4H*-chromene family in an enantioselective and cascade manner allows chemists to explore their potential biological properties. Therefore, the development of catalytic asymmetric strategies for the efficient construction of structurally diverse chiral *4H*-chromenes is highly valuable to the fields of organic synthesis and chemical biology/medicinal chemistry.

Toward this end, our group has recently reported the enantioselective cascade oxa-Michael Michael reactions of ynals involving an unprecedented iminium-allenamine activation mode (Scheme 3.6, eq 1)¹⁴. Despite the extensively investigated iminium-enamine strategy with cascade sequences^{1a}, the respective iminium-allenamine chemistry is much less appreciated. Expanding the scope of the chemistry will not only add a new domain in organocatalysis but also, more importantly, afford new potentially useful platforms for the facile assembly of important molecular scaffolds. We questioned


whether the conceptually novel tactic could be extended to an oxa-Michael aldol cascade sequence, which would generate the structurally diverse 4*H*-chromenes with installation of new functionalities. Herein, we wish to report the results which have led to an asymmetric iminium-allenamine approach to chiral 4*H*-chromenes bearing chiral-hydroxy carboxylate functionality in high yields (84~99%) and with excellent enantioselectivities (93~>99% ee) for a wide range of substrates (Scheme 3.6, eq 2). A powerful “one-pot” oxa-Michael aldol cascade reaction between (2-hydroxy-phenyl)-2-oxoacetates and ynals is realized for the first time. Furthermore, a quaternary stereogenic center involving formation of versatile α -hydroxy carboxylate motif, a still synthetically unmet issue in organic synthesis, is created highly enantioselectively.

Scheme 3.6 Alkynals Participating in Enantioselective Cascade Oxa-Michael-Michael and Aldol Reactions Promoted by a Chiral Secondary Amine Involving a Novel Iminium-allenamine Activation Mode

Oxa-Michael-Michael Cascade (Eq. 1)



Oxa-Michael-Aldol Cascade (Eq. 2)




3.5.2 Results and Discussion

Initial efforts on the proposed organocatalytic enantioselective oxa-Michael-aldol cascade reaction of simple salicylaldehyde or -ketone with phenyl ynal in the presence of

20 mol % (*S*)-diphenylpyrrolinol TMS ether **I** in toluene were frustrated by the failure of delivering the desired products as a result of very poor reaction yield. These studies underscored the completely different reaction behavior between ynals and enals since we have shown that enals can efficiently engage in the Michael aldol cascade process with salicylaldehydes^{20b}. We turned our attention to the more reactive ketone ester **6a** (Scheme 3.6 and Table 3.5)³³. To our delight, the oxa-Michael-aldol cascade sequence proceeded smoothly to give the desired product **7a** in an excellent yield (97%) and with excellent enantiomeric excess (ee) (97%) under the same reaction conditions in toluene at r.t. in the presence of 20 mol % catalyst **I** within 3 h (Table 3.5, entry 1), albeit the highly steric ketoester substrate employed. Notably, the process provides an efficient entry to useful chiral α -hydroxycarboxylates with formation of a quaternary stereogenic center in a cascade fashion. The screening of other diarylpyrrolinol silyl ether analogues **II-III** revealed that the bulky TBDMS catalyst **III** gave a slightly higher enantioselectivity (Table 3.5, entries 2~3). Quinine alone was not effective for this transformation (Table 3.5, entry 4). Therefore, the catalyst **III** was defined as the optimal promoter for this process. A survey of solvents led to choose toluene as the reaction medium for the process (Table 3.5, entries 5~7). Better e.e. (99%) was observed when decreasing the reaction temperature to -15~-10 °C and lowering the catalyst loading to 15 mol% (Table 3.5, entry 8).

Table 3.5 Development and Optimization of Reaction Conditions of an Organocatalytic Enantioselective Cascade Oxa-Michael-aldol Cascade Reaction^a




Entry	Catalyst	Solvent	<i>t</i> (h)	Yield (%) ^b	ee (%) ^c
1	I	Toluene	3	97	97
2	II	Toluene	3	98	98
3	III	Toluene	3	98	98
4	(-)-quinine	THF	2	<5	ND ^d
5	III	Xylene	3	96	98
6	III	CH ₂ Cl ₂	3	99	97
7	III	Cl(CH ₂) ₂ Cl	3	94	96
8 ^e	III	Toluene	6	98	99

^a Reaction conditions: unless specified, a mixture of phenyl-propynal **1a** (0.080 mmol), ethyl 2-(2-hydroxy-4-methoxyphenyl)-2-oxoacetate **6a** (0.080 mmol), and catalyst (0.016 mmol) in solvent (0.5 mL) was stirred at r.t. for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralcel AD). ^d Not determined. ^e Performed at -15~-10 °C, and 15 mol % of catalyst **III** was applied.

With the optimal conditions in hand, the scope of the cascade oxa-Michael-aldol reaction was investigated. Remarkably, as shown in Table 3.6, the cascade process exhibits a broad substrate tolerance, and a quaternary stereogenic center is generated highly enantioselectively in all cases. A full range of aromatic ynals **1**, including electron-neutral (entries 1~5), -donating (entries 6 and 7), and -withdrawing substituents (entries 8~15), as well as heterocyclic examples (entries 16 and 17), can participate in the cascade reactions in excellent yields (84%~99%) and with excellent enantioselectivities (93~>99%). It is noteworthy that the substrate scope can also be expanded to highly steric demanding aliphatic alkynal, providing the desired product in 91% yield and 98% ee (entry 18). Moreover, alkyl alkynals (R=CH₂CH₂Ph and *n*-C₅H₁₁, entries 19 and 20, respectively) were also probed for the cascade process. It was found that they efficiently participated in the reaction with excellent enantioselectivities (97 and 99% e.e., respectively), although the reaction occurred slower (60 h) than aryl ones and higher catalyst loading (25 mol %) was needed as a result of their poorer reactivity. The variation in ethyl 2-(2-hydroxyphenyl)-2-oxoacetate components **6** was also possible, as demonstrated by the aromatic structures carrying electron-donating (entries 1, 3, 4, 6, 10, and 12~18), -neutral (entries 2 and 11), and -withdrawing (entry 5) groups. It is observed that with respect to substrate **6e** having electron-withdrawing Cl (entry 5) poor e.e. was observed. However, we found that quinine as co-catalyst was essential to ensure high enantioselectivity presumably due to its interaction with the ketone moiety in the ketoester. The absolute configuration of **7h** was determined to be *R* configuration by X-ray crystallographic analysis (Figure 3.7).

Table 3.6 Catalyst **III** Promoted Enantioselective Cascade Oxa-Michael-aldol Reactions of Alkynals **1** with Ethyl 2-(2-Hydroxyphenyl)-2-oxoacetates **6**^a




Entry	R	X	<i>t</i> (h)	Yield (%) ^b	ee (%) ^c
1	Ph	5-MeO	6	98	99
2	Ph	H	8	97	99
3	Ph	4-MeO	8	84	96
4	Ph	4-Me	9	93	98
5 ^d	Ph	4-Cl	6	99	93
6	4-MeOC ₆ H ₄	5-MeO	6	99	99
7	4-MeC ₆ H ₄	5-MeO	6	99	99
8	4-ClC ₆ H ₄	5-MeO	6	97	99
9	4-BrC ₆ H ₄	5-MeO	6	99	99
10	4-BrC ₆ H ₄	4-Me	10	92	98
11	4-BrC ₆ H ₄	H	9	98	98
12	4-FC ₆ H ₄	5-MeO	6	99	>99
13	3-FC ₆ H ₄	4-Me	9	93	98
14	4-NO ₂ C ₆ H ₄	5-MeO	6	98	99
15	3-NO ₂ C ₆ H ₄	5-MeO	6	98	99
16	2-thienyl	5-MeO	10	92	99
17	2-furanyl	5-MeO	10	97	96
18	<i>t</i> -Bu	5-MeO	10	91	98

19 ^e	PhCH ₂ CH ₂	4-Me	60	69	97
20 ^e	<i>n</i> -C ₅ H ₁₁	4-Me	60	75	99

^a Reaction conditions: unless specified, see footnote *a* in Table 3.5 and Supporting Information. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralcel AD or Chiralpak IC). ^d 15 mol % of **I** and 10 mol % of (-)-Quinine were used. ^e 25 mol % of **III** was used.

Figure 3.7 X-ray Structure of **7h**



3.5.3 Conclusion


In conclusion, we have described a new protocol for asymmetric organocatalysis through the first iminium-allenamine engaged oxa-Michael-aldol cascade reaction. In addition to the potential of this powerful “one-pot” approach to the catalytic generation of chiral 4*H*-chromenes bearing a quaternary stereogenic center, it is a rare example of a highly enantioselective cascade reaction with ynals. The operationally friendly reaction conditions and high efficiency (cascade, high yields and e.e., and readily available achiral

starting materials) render the process particularly attractive in the practice of organic synthesis and medicinal chemistry. Chiral 4*H*-chromene products with a versatile α -hydroxycarboxylate moiety are valuable structures for the synthesis of biologically active molecules.

3.5.4 Supporting Information

General procedure for Cascade Michael-Aldol addition reaction:

To a solution of alkynal **1** (0.08 mmol), organocatalyst **III** (4.41 mg, 0.012 mmol) in toluene (1.0 mL) was added ethyl 2-(2-hydroxyphenyl)-2-oxoacetate **6** (0.08 mmol). The resulting solution was stirred at -15~-10 °C for a specified time. Then the reaction mixture was directly purified by column chromatography, eluted with hexane/EtOAc to afford the desired product.




(R)-Ethyl 3-formyl-4-hydroxy-7-methoxy-2-phenyl-4*H*-chromene-4-carboxylate (7a)

(Table 3.5, entry 1).

The title compound was prepared according to the general procedure, as described above in 98% yield. ¹H NMR (CDCl₃, 300 MHz): δ 9.56 (s, 1H), 7.51~7.65 (m, 5H), 7.45 (d, *J* =


9.0 Hz, 1H), 6.82 (dd, $J = 9.0, 2.4$ Hz, 1H), 6.72 (d, $J = 2.4$ Hz, 1H), 4.76 (s, 1H), 4.13~4.29 (m, $J = 7.2$ Hz, 2H), 3.81 (s, 3H), 1.18 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR(CDCl₃, 75 MHz): δ 191.6, 173.3, 166.8, 160.8, 150.0, 131.4, 130.7, 130.1, 128.6, 128.1, 116.2, 113.6, 113.2, 101.3, 68.0, 62.6, 55.5, 13.9; HPLC (Chiralcel AD, *i*-PrOH/hexane = 20/80, flow rate = 1.0 mL/min, $\lambda = 254$ nm): $t_{\text{minor}} = 48.86$ min, $t_{\text{major}} = 51.31$ min, ee = 99%; $[\alpha]_{\text{D}}^{25} = -2$ ($c = 1.0$ in CHCl₃); HRMS (ESI): C₂₀H₁₈O₆+Na, calcd: 377.0996, found: 377.0989; IR (CHCl₃, cm⁻¹): 3478, 2978, 2935, 2860, 1738, 1663, 1635, 1506, 1334, 1306, 1236, 1197, 1173, 1114, 1067, 1032, 887, 775, 701.



(R)-Ethyl 3-formyl-4-hydroxy-2-phenyl-4H-chromene-4-carboxylate (7b) (Table 3.5, entry 2).


The title compound was prepared according to the general procedure, as described above in 99% yield. ^1H NMR (CDCl₃, 300 MHz): δ 9.57 (s, 1H), 7.22~7.67 (m, 9H), 4.85 (s, 1H), 4.14~4.31 (m, $J = 7.2$ Hz, 2H), 1.18 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (CDCl₃, 75 MHz): δ 191.6, 173.2, 167.0, 149.1, 131.5, 130.7, 130.2, 130.1, 128.6, 127.2, 125.8, 121.4, 117.2, 116.0, 68.2, 62.8, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 10/90, flow rate = 1.0 mL/min, $\lambda = 254$ nm): $t_{\text{minor}} = 44.92$ min, $t_{\text{major}} = 46.69$ min, ee = 99%; $[\alpha]_{\text{D}}^{25} = -5$ ($c = 1.0$ in CHCl₃); HRMS (ESI): C₁₉H₁₆O₅+Na, calcd: 347.0890, found: 347.0895; IR (CHCl₃,

cm⁻¹): 3483, 2981, 2932, 2860, 1740, 1664, 1578, 1396, 1334, 1239, 1219, 1192, 1118, 1071, 1039, 890, S3 838, 765, 702.



(R)-Ethyl 3-formyl-4-hydroxy-6-methoxy-2-phenyl-4H-chromene-4-carboxylate (7c)
(Table 3.5, entry 3).


The title compound was prepared according to the general procedure, as described above in 96% yield. ¹HNMR (CDCl₃, 300 MHz): δ 9.56 (s,1H), 7.51~7.66 (m,5H), 7.16 (d, *J* = 9.0 Hz,1H), 7.01 (d, *J* = 3.0 Hz, 1H), 6.95 (dd, *J* = 9.0, 3.0 Hz, 1H), 4.87 (s,1H), 4.14~4.31 (m, *J* = 7.2 Hz, 2H), 3.81 (s,3H), 1.19 (t, *J* = 7.2Hz, 3H); ¹³CNMR(CDCl₃, 75 MHz): δ 191.8, 173.2, 167.1, 157.2, 143.3, 131.4, 130.8, 130.2, 128.6, 121.8, 118.4, 113.6, 117.6, 115.1, 109.6, 68.7, 62.8, 55.8, 14.0; HPLC (Chiralcel AD, EtOH/hexane = 20/80, flow rate = 1.0 mL/min, λ = 254 nm): t_{minor} = 25.17 min, t_{major} = 31.75 min, ee = 96%; [α]_D²⁵ = -18 (*c* = 1.0 in CHCl₃); HRMS (ESI): C₂₀H₁₈O₆+Na, calcd: 377.0996, found: 377.0991; IR (CHCl₃, cm⁻¹): 3473, 2980, 2923, 2855, 1738, 1654, 1619, 1331, 1303, 1235, 1198, 1153, 1098, 1060, 1038, 817, 773, 753, 700, 664.



(R)-Ethyl 3-formyl-4-hydroxy-6-methyl-2-phenyl-4H-chromene-4-carboxylate (7d)

(Table 3.5, entry 4).

The title compound was prepared according to the general procedure, as described above in 93% yield. ¹HNMR (CDCl₃, 500 MHz): δ 9.55 (s, 1H), 7.65~7.63 (m, 2H), 7.60~7.57 (m, 1H), 7.53 (t, *J* = 7.5 Hz, 2H), 7.34 (s, 1H), 7.18 (d, *J* = 8.0 Hz, 1H), 7.11 (d, *J* = 8.5 Hz, 1H), 4.82 (s, 1H), 4.32~4.26 (m, 1H), 4.19~4.12 (m, 1H), 2.35 (s, 3H), 1.18 (t, *J* = 7.0 Hz, 3H); ¹³CNMR (CDCl₃, 125 MHz): δ 191.7, 173.3, 167.0, 147.1, 135.6, 131.3, 131.0, 130.8, 130.1, 128.6, 127.1, 120.9, 116.9, 115.8, 68.3, 62.7, 20.9, 13.9; HPLC (Chiralpak IC, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, λ = 254nm): *t*_{minor} = 57.56 min, *t*_{major} = 44.41 min, ee = 98%; [α]_D²⁰ = -45.6 (*c* = 1.0 in CHCl₃).




(R)-Ethyl 6-chloro-3-formyl-4-hydroxy-2-phenyl-4H-chromene-4-carboxylate (7e)

(Table 3.5, entry 5).


The title compound was prepared according to the general procedure, as described above in 99% yield. ¹HNMR (CDCl₃, 300 MHz): δ 9.56 (s, 1H), 7.51~7.66 (m, 6H), 7.34 (dd, *J* = 8.7, 2.4 Hz, 1H), 7.18 (d, *J* = 8.7 Hz, 1H), 4.87 (s, 1H), 4.13~4.36 (m, *J* = 7.2 Hz, 2H), 1.20 (t, *J* = 7.2 Hz, 3H); ¹³CNMR (CDCl₃, 75 MHz): δ 191.4, 172.7, 166.6, 147.6, 131.6, 130.8, 130.4, 130.3, 130.2, 128.7, 127.1, 122.9, 118.7, 115.6, 68.1, 63.0, 13.9; HPLC

(Chiralcel AD, EtOH/hexane = 15/85, flow rate = 1.0 mL/min, λ = 254 nm): t_{minor} = 20.54 min, t_{major} = 22.93 min, ee = 93%; $[\alpha]_{\text{D}}^{25}$ = -49 (c = 1.0 in CHCl_3); HRMS (ESI): $\text{C}_{19}\text{H}_{15}\text{O}_5\text{Cl}+\text{Na}$, calcd: 381.0500, found: 381.0499; IR (CHCl_3 , cm^{-1}): 3475, 2980, 2925, 2858, 1741, 1662, 1632, 1478, 1331, 1238, 1184, 1094, 1069, 1037, 890, 771, 701.



(R)-Ethyl 3-formyl-4-hydroxy-7-methoxy-2-(4-methoxy-phenyl)-4H-chromene-4-carboxylate (7f) (Table 3.5, entry 6).


The title compound was prepared according to the general procedure, as described above in 99% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.57 (s, 1H), 7.59 (dd, J = 6.6, 1.8 Hz, 2H), 7.43 (d, J = 8.7 Hz, 1H), 7.04 (dd, J = 6.6, 1.8 Hz, 2H), 6.81 (dd, J = 8.7, 2.7 Hz, 1H), 6.72 (d, J = 2.7 Hz, 1H), 4.76 (s, 1H), 4.12~4.28 (m, J = 7.2 Hz, 2H), 3.89 (s, 3H), 3.81 (s, 3H), 1.17 (t, J = 7.2 Hz, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 191.6, 173.4, 166.7, 162.1, 160.8, 150.0, 131.8, 128.0, 122.8, 115.7, 114.6, 113.7, 113.0, 101.3, 68.2, 62.5, 55.5, 55.4, 21.5, 13.9; HPLC (Chiralcel AD, *i*-PrOH/hexane = 20/80, flow rate = 1.0 mL/min, λ = 254 nm): t_{minor} = 116.37 min, t_{major} = 97.17 min, ee = 99%; $[\alpha]_{\text{D}}^{25}$ = +1 (c = 1.0 in CHCl_3); HRMS (ESI): $\text{C}_{21}\text{H}_{20}\text{O}_7+\text{Na}$, calcd: 407.1101, found: 407.1110; IR (CHCl_3 , cm^{-1}): 3469, 2958, 2926, 2850, 1738, 1632, 1607, 1508, 1333, 1292, 1254, 1197, 1174, 1116, 1066, 1032, 889, 834.



(R)-Ethyl 3-formyl-4-hydroxy-7-methoxy-2-p-tolyl-4H-chromene-4-carboxylate (7g)


(Table 3.5, entry 7).

The title compound was prepared according to the general procedure, as described above in 99% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.55(s, 1H), 7.53(d, $J = 8.1\text{ Hz}$, 2H), 7.44 (d, $J = 8.7\text{ Hz}$, 1H), 7.33 (d, $J = 8.1\text{ Hz}$, 2H), 6.81 (dd, $J = 8.7, 2.4\text{ Hz}$, 1H), 6.71(d, $J = 2.4\text{ Hz}$, 1H), 4.76(s, 1H), 4.12~4.28 (m, $J = 6.9\text{ Hz}$, 2H), 3.82 (s, 3H), 2.45 (s, 3H), 1.17 (t, $J = 6.9\text{ Hz}$, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 191.8, 173.4, 167.0, 160.8, 150.0, 142.0, 130.1, 129.3, 128.1, 127.8, 116.0, 113.7, 113.1, 101.3, 68.2, 62.6, 55.6, 21.5, 14.0.; HPLC (Chiralcel AD, EtOH/hexane = 15/85, flow rate = 1.0 mL/min, $\lambda = 254\text{ nm}$): $t_{\text{minor}} = 62.05\text{ min}$, $t_{\text{major}} = 54.31\text{ min}$, ee = 99%; $[\alpha]_{\text{D}}^{25} = -4$ ($c = 1.0$ in CHCl_3); HRMS (ESI): $\text{C}_{21}\text{H}_{20}\text{O}_6 + \text{Na}$, calcd: 391.1152, found: 391.1149; IR (CHCl_3 , cm^{-1}): 3483, 2957, 2921, 2851, 1739, 1663, 1634, 1506, 1332, 1236, 1197, 1172, 1115, 1067, 1033, 889, 827.



(R)-Ethyl 2-(4-chloro-phenyl)-3-formyl-4-hydroxy-7-methoxy-4H-chromene-4-carboxylate (7h) (Table 3.5, entry 8).


The title compound was prepared according to the general procedure, as described above in 97% yield. ¹HNMR (CDCl₃, 300 MHz): δ 9.54 (s, 1H), 7.69 (m, 2H), 7.52 (m, 2H), 7.44 (d, *J* = 8.7 Hz, 1H), 6.83 (dd, *J* = 8.7, 2.7 Hz, 1H), 6.72 (d, *J* = 2.7 Hz, 1H), 4.75 (s, 1H), 4.11~4.29(m, *J* = 6.9 Hz, 2H), 3.82(s, 3H), 1.18 (t, *J* = 6.9 Hz, 3H); ¹³CNMR (CDCl₃, 300 MHz): δ 191.0, 173.2, 165.6, 160.9, 149.9, 137.8, 131.5, 129.1, 129.0, 128.1, 116.5, 113.5, 113.3, 101.3, 68.0, 62.7, 55.6, 14.0; HPLC (Chiralcel AD, EtOH/hexane = 15/85, flow rate = 1.0 mL/min, λ = 254nm): t_{minor} = 73.10 min, t_{major} = 61.60 min, ee = 99%; [α]_D²⁵ = -2 (*c* = 1.0 in CHCl₃); HRMS(ESI): C₂₀H₁₇O₆Cl+Na, calcd: 411.0606, found: 411.0617; IR (CHCl₃, cm⁻¹): 3481, 2976, 2932, 2855, 1739, 1665, 1635, 1507, 1333, 1237, 1197, 1173, 1115, 1067, 1032, 888, 838, 762.



(R)-Ethyl 2-(4-Bromo-phenyl)-3-formyl-4-hydroxy-7-methoxy-4H-chromene-4-carboxylate (7i) (Table 3.5, entry 9).

The title compound was prepared according to the general procedure, as described above in 99% yield. ¹HNMR (CDCl₃, 300 MHz): δ 9.54 (s, 1H), 7.68 (d, *J* = 8.3 Hz, 2H), 7.52 (d,


$J = 8.3$ Hz, 2H), 7.44 (d, $J = 8.7$ Hz, 1H), 6.83 (dd, $J = 8.7, 2.4$ Hz, 1H), 6.71 (d, $J = 2.4$ Hz, 1H), 4.73 (s, 1H), 4.14~4.29 (m, $J = 6.9$ Hz, 2H), 3.82 (s, 3H), 1.18 (t, $J = 6.9$ Hz, 3H); ^{13}C NMR(CDCl_3 , 75 MHz): δ 191.0, 173.2, 165.6, 160.9, 149.9, 132.0, 131.6, 129.6, 128.1, 126.2, 116.5, 113.4, 113.3, 101.3, 68.0, 62.7, 55.6, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 20/80, flow rate = 1.0 mL/min, $\lambda = 254$ nm): $t_{\text{minor}} = 65.97$ min, $t_{\text{major}} = 56.51$ min, ee = 99%; $[\alpha]_{\text{D}}^{25} = -5$ ($c = 1.0$ in CHCl_3); HRMS (ESI): $\text{C}_{20}\text{H}_{17}\text{BrO}_6 + \text{Na}$, calcd: 455.0101, found: 455.0110; IR (CHCl_3 , cm^{-1}): 3480, 2977, 2933, 2861, 1738, 1664, 1634, 1507, 1331, 1237, 1197, 1173, 1115, 1067, 1032, 887, 834, 757.



(R)-Ethyl 2-(4-bromo-phenyl)-3-formyl-4-hydroxy-6-methyl-4H-chromene-4-carboxylate (7j) (Table 3.5, entry 10).


The title compound was prepared according to the general procedure, as described above in 92% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.54 (s, 1H), 7.67 (d, $J = 8.5$ Hz, 2H), 7.51 (d, $J = 8.0$ Hz, 2H), 7.32 (s, 1H), 7.18 (d, $J = 8.5$ Hz, 1H), 7.10 (d, $J = 8.5$ Hz, 1H), 4.80 (s, 1H), 4.30~4.14 (m, 2H), 2.35 (s, 3H), 1.18 (t, $J = 7.0$ Hz, 3H); ^{13}C NMR(CDCl_3 , 125 MHz): δ 191.0, 173.1, 165.7, 147.0, 135.8, 131.9, 131.6, 131.0, 129.7, 127.1, 126.1, 120.8, 116.8, 116.1, 68.2, 62.7, 20.9, 13.9; HPLC (Chiralpak IC, *i*-PrOH/hexane = 20/80,

flowrate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 58.48\text{ min}$, $t_{\text{major}} = 46.10\text{ min}$, ee = 98%;
[α]_D²⁰ = -24.3 ($c = 1.0$ in CHCl₃).




(R)-Ethyl 2-(4-bromo-phenyl)-3-formyl-4-hydroxy-4H-chromene-4-carboxylate (7k)
(Table 3.5, entry 11).

The title compound was prepared according to the general procedure, as described above in 98% yield. ¹HNMR (CDCl₃, 500 MHz): δ 9.55 (s, 1H), 7.68 (d, $J = 8.0\text{ Hz}$, 2H), 7.55~7.52 (m, 3H), 7.41~7.38 (m, 1H), 7.28~7.21 (m, 2H), 4.83 (s, 1H), 4.30~4.16 (m, 2H), 1.18 (t, $J = 7.5\text{ Hz}$, 3H); ¹³CNMR (CDCl₃, 125 MHz): δ 190.9, 173.0, 165.6, 148.9, 131.9, 131.6, 130.2, 129.6, 127.2, 126.2, 125.9, 121.3, 117.1, 116.2, 68.1, 62.8, 13.9; HPLC (Chiralpak IC, *i*-PrOH/hexane = 10/90, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 125.31\text{ min}$, $t_{\text{major}} = 87.56\text{ min}$, ee = 98%; [α]_D²⁰ = -11.3 ($c = 1.0$ in CHCl₃).



(R)-Ethyl 2-(4-Fluoro-phenyl)-3-formyl-4-hydroxy-7-methoxy-4H-chromene-4-carboxylate (7l) (Table 3.5, entry 12).


The title compound was prepared according to the general procedure, as described above in 99% yield. ¹HNMR (CDCl₃, 300 MHz): δ 9.54 (s,1H) ,7.66 (m, 2H), 7.44 (d, *J* = 8.7 Hz, 1H), 7.22 (m,2H),6.83(dd, *J* = 8.7, 2.4 Hz, 1H), 6.72 (d, *J* = 2.4 Hz, 1H), 4.77 (s,1H), 4.14~4.29 (m, *J* = 7.2 Hz, 2H), 3.82(s, 3H), 1.18 (t, *J* = 7.2 Hz, 3H); ¹³CNMR (CDCl₃, 75 MHz): δ 191.2, 173.3, 165.8, 160.9, 149.9, 132.4, 132.3, 128.1, 126.8, 116.2, 115.8, 113.5, 113.3, 101.3, 68.0, 62.7, 55.6, 14.0; HPLC (Chiralcel AD, EtOH/hexane = 20/80, flow rate = 1.0 mL/min, λ = 254nm): t_{minor} = 48.76 min, t_{major} = 43.86 min, ee = >99%; [α]_D²⁵ = 0 (*c* = 1.0 in CHCl₃); HRMS (ESI): C₂₀H₁₇O₆F+Na, calcd: 395.0901, found: 395.0905; IR (CHCl₃, cm⁻¹): 3483, 2962, 2933,2856, 1740, 1666, 1636, 1507, 1333, 1235, 1198, 1172, 1115, 1068, 1033, 891, 846, 814, 627,517.



(R)-Ethyl 2-(3-Fluoro-phenyl)-3-formyl-4-hydroxy-6-methyl-4H-chromene-4-carboxylate (7m) (Table 3.5, entry 13).

The title compound was prepared according to the general procedure, as described above in 93% yield. ¹HNMR (CDCl₃, 500 MHz): δ 9.56 (s, 1H), 7.53~7.48 (m, 1H), 7.42~7.38 (m, 2H), 7.33 (s, 1H), 7.31~7.26 (m, 1H), 7.19 (d, *J* = 8.0 Hz, 1H), 7.11 (d, *J* = 8.0


Hz, 1H), 4.80 (s, 1H), 4.32~4.26 (m, 1H), 4.19~4.13 (m, 1H), 2.35 (s, 3H), 1.18 (t, $J = 7.0$ Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.1, 173.1, 165.3, 163.4, 161.5, 147.0, 135.8, 132.7 ($J = 8.3\text{Hz}$), 131.0, 130.3 ($J = 8.1$ Hz), 127.1, 126.2, 120.8, 118.4 ($J = 21$ Hz), 117.1, 116.9 ($J = 7.9$ Hz), 116.2, 68.2, 62.7, 20.9, 13.9; HPLC (Chiralpak IC, *i*-PrOH/hexane = 20/80, flow rate = 0.6mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 53.52$ min, $t_{\text{major}} = 44.73$ min, ee = 98%; $[\alpha]_{\text{D}}^{20} = -42.4$ ($c = 1.0$ in CHCl_3).



(R)-Ethyl 3-formyl-4-hydroxy-7-methoxy-2-(4-nitro-phenyl)-4H-chromene-4-carboxylate (7n) (Table 3.5, entry 14).


The title compound was prepared according to the general procedure, as described above in 98% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.53 (s, 1H), 8.40 (d, $J = 8.7$ Hz, 2H), 7.85 (d, $J = 8.7$ Hz, 2H), 7.44 (d, $J = 8.7$ Hz, 1H), 6.86 (dd, $J = 8.7, 2.4$ Hz, 1H), 6.72 (d, $J = 2.4$ Hz, 1H), 4.74 (s, 1H), 4.16~4.30 (m, $J = 7.2$ Hz, 2H), 3.83 (s, 3H), 1.18 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 191.1, 172.9, 164.0, 161.0, 149.7, 149.4, 136.7, 131.2, 128.2, 123.8, 117.4, 113.6, 113.4, 101.3, 67.8, 62.9, 55.6, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 30/70, flow rate = 1.0 mL/min, $\lambda = 254$ nm): $t_{\text{minor}} = 78.30$ min, $t_{\text{major}} = 102.15$ min, ee = 99%; $[\alpha]_{\text{D}}^{25} = -10$ ($c = 1.0$ in CHCl_3); HRMS (ESI): $\text{C}_{20}\text{H}_{17}\text{NO}_8 + \text{Na}$,

calcd: 422.0846, found: 422.0838; IR (CHCl₃, cm⁻¹): 3472, 2976, 2931, 2861, 1738, 1666, 1637, 1524, 1347, 1238, 1197, 1174, 1114, 1067, 1031, 932, 851, 753, 702.




(R)-Ethyl 3-Formyl-4-hydroxy-7-methoxy-2-(3-nitro-phenyl)-4H-chromene-4-carboxylate (7o) (Table 3.5, entry 15).

The title compound was prepared according to the general procedure, as described above in 98% yield. ¹HNMR (CDCl₃, 300 MHz): δ 9.54(s, 1H), 8.56 (s, 1H), 8.45 (d, *J* = 8.1 Hz, 1H), 7.96 (d, *J* = 7.5 Hz, 1H), 7.75 (dd, *J* = 8.1, 7.5 Hz, 1H), 7.44 (d, *J* = 8.7 Hz, 1H), 6.85 (dd, *J* = 8.7, 2.4 Hz, 1H), 6.74 (d, *J* = 2.4 Hz, 1H), 4.74 (s, 1H), 4.16~4.30 (m, *J* = 6.9 Hz, 2H), 3.83 (s, 3H), 1.17 (t, *J* = 6.9 Hz, 3H); ¹³CNMR (CDCl₃, 75 MHz): δ 190.0, 172.9, 163.8, 161.0, 148.3, 136.0, 132.4, 129.8, 128.1, 126.0, 124.9, 117.3, 113.7, 113.4, 101.3, 67.8, 62.9, 55.7, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 20/80, flow rate = 1.0 mL/min, λ = 254 nm): *t*_{minor} = 65.94 min, *t*_{major} = 51.11 min, ee = 99%; [α]_D²⁵ = + 2 (*c* = 1.0 in CHCl₃); HRMS (ESI): C₂₀H₁₇NO₈+Na, calcd: 422.0846, found: 422.0840; IR (CHCl₃, cm⁻¹): 3423, 3088, 2966, 2931, 2867, 1738, 1666, 1637, 1533, 1348, 1237, 1198, 1174, 1114, 1067, 1033, 935, 854, 814, 763, 741, 699.




(R)-Ethyl 3-formyl-4-hydroxy-7-methoxy-2-thiophen-2-yl-4H-chromene-4-carboxylate (7p) (Table 3.5, entry 16).

The title compound was prepared according to the general procedure, as described above in 92% yield. ¹HNMR (CDCl₃, 300 MHz): δ 10.02 (s, 1H), 9.84 (s, 1H), 7.81 (d, *J* = 3.9 Hz, 1H), 7.54 (d, *J* = 3.9 Hz, 1H), 7.41 (d, *J* = 8.7 Hz, 1H), 6.83 (dd, *J* = 8.7, 2.7 Hz, 1H), 6.73 (d, *J* = 2.7 Hz, 1H), 4.73 (s, 1H), 4.13~4.28 (m, 2H), 3.83 (s, 3H), 1.17 (t, *J* = 7.2 Hz, 3H); ¹³CNMR (CDCl₃, 75 MHz): δ 189.8, 182.8, 172.8, 161.0, 149.5, 147.1, 140.1, 134.9, 133.1, 128.0, 117.7, 113.7, 113.2, 101.2, 68.0, 62.9, 55.6, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 20/80, flow rate = 1.0 mL/min, λ = 254 nm): *t*_{minor} = 74.02 min, *t*_{major} = 66.40 min, ee = 99%; [α]_D²⁵ = -50 (*c* = 1.0 in CHCl₃); HRMS (ESI): C₁₈H₁₆O₆S+Na, calcd: 383.0565, found 383.0561; IR (CHCl₃, cm⁻¹): 3470, 3094, 2954, 2925, 2853, 1740, 1665, 1631, 1507, 1341, 1286, 1236, 1211, 1169, 1111, 1065, 1029, 854, 818.



(R)-Ethyl 3-formyl-2-furan-2-yl-4-hydroxy-7-methoxy-4H-chromene-4-carboxylate (7q) (Table 3.5, entry 17).


The title compound was prepared according to the general procedure, as described above in 97% yield. ¹HNMR (CDCl₃, 300 MHz): δ 10.30 (s, 1H), 7.69 (d, *J* = 1.8 Hz, 1H), 7.42 (d, *J* = 8.7 Hz, 1H), 7.15 (d, *J* = 3.0 Hz, 1H), 6.80 (dd, *J* = 8.7, 2.4 Hz, 1H), 6.73 (d, *J* = 2.4 Hz, 1H), 6.64 (dd, *J* = 3.0, 1.8 Hz, 1H), 4.11~4.26 (m, *J* = 7.2 Hz, 2H), 3.83 (s, 3H), 1.15 (t, *J* = 7.2 Hz, 3H); ¹³CNMR (CDCl₃, 75 MHz): δ 191.5, 173.3, 160.9, 154.1, 149.4, 146.5, 146.0, 128.0, 116.8, 115.2, 113.3, 113.1, 112.1, 101.1, 68.0, 62.6, 55.6, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 20/80, flow rate = 1.0 mL/min, λ = 254 nm): *t*_{minor} = 46.07 min, *t*_{major} = 38.55 min, ee = 96%; [α]_D²⁵ = -27 (*c* = 1.0 in CHCl₃); HRMS (ESI): C₁₈H₁₆O₇+Na, calcd: 367.0788, found: 367.0782; IR (CHCl₃, cm⁻¹): 3455, 3148, 2973, 2927, 1735, 1631, 1400, 1234, 1208, 1111, 1064, 1027, 832, 764.



(R)-Ethyl 2-tert-butyl-3-formyl-4-hydroxy-7-methoxy-4H-chromene-4-carboxylate (7r) (Table 3.5, entry 18).

The title compound was prepared according to the general procedure, as described above in 91% yield. ¹HNMR (CDCl₃, 300 MHz): δ 10.35 (s, 1H), 7.39 (d, *J* = 8.7 Hz, 1H), 6.76 (dd, *J* = 8.7, 2.4 Hz, 1H), 6.63 (d, *J* = 2.4 Hz, 1H), 4.67 (s, 1H), 4.05~4.24 (m, *J* = 7.2 Hz, 2H), 3.81 (s, 3H), 1.54 (s, 9H), 1.14 (t, *J* = 7.2 Hz, 3H); ¹³CNMR (CDCl₃, 75 MHz): δ 190.5, 174.7, 173.6, 160.7, 149.3, 127.8, 115.8, 113.1, 113.0, 100.8, 68.8, 62.2, 55.5,


38.5, 31.3, 13.9; HPLC (Chiralcel AD, EtOH/hexane = 15/85, flow rate = 1.0 mL/min, λ = 254 nm): t_{minor} = 17.29 min, t_{major} = 21.26 min, ee = 98%; $[\alpha]_{\text{D}}^{25}$ = -34 (c = 1.0 in CHCl_3); HRMS (ESI): $\text{C}_{18}\text{H}_{22}\text{O}_6 + \text{Na}$, calcd: 357.1309, found: 357.1318; IR (CHCl_3 , cm^{-1}): 3511, 2959, 2923, 2853, 1739, 1653, 1632, 1571, 1508, 1291, 1231, 1205, 1178, 1117, 1036, 934, 757.



(R)-Ethyl 3-formyl-4-hydroxy-6-methyl-2-phenethyl-4H-chromene-4-carboxylate (7s)

(Table 3.5, Entry 19)

The title compound was prepared according to the general procedure, as described above in 69% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.66 (s, 1H), 7.29~7.26 (m, 3H), 7.22~7.18 (m, 3H), 7.14 (d, J = 8.0 Hz, 1H), 7.03 (d, J = 8.5 Hz, 1H), 4.64 (s, 1H), 4.24~4.18 (m, 1H), 4.13~4.07 (m, 1H), 3.14~3.08 (m, 4H), 2.32 (s, 3H), 1.13 (t, J = 7.0 Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 188.8, 173.3, 168.0, 146.7, 139.4, 135.4, 130.8, 128.7, 128.5, 127.0, 126.7, 121.0, 116.5, 116.0, 67.8, 62.5, 33.9, 32.0, 20.8, 13.9; HPLC (Chiralpak IC, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, λ = 254nm): t_{minor} = 126.8 min, t_{major} = 78.9 min, ee = 97%; $[\alpha]_{\text{D}}^{17}$ = -32.2 (c = 1.0 in CHCl_3).



(R)-Ethyl 3-formyl-4-hydroxy-6-methyl-2-pentyl-4H-chromene-4-carboxylate (7t)
(Table 3.5, Entry 20)

The title compound was prepared according to the general procedure, as described above in 75% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.93 (s, 1H), 7.28 (s, 1H), 7.14 (d, $J = 8.5$ Hz, 1H), 7.03 (d, $J = 8.0$ Hz, 1H), 4.71 (s, 1H), 4.28~4.21 (m, 1H), 4.14~4.07 (m, 1H), 2.89~2.78 (m, 2H), 2.32 (s, 3H), 1.81~1.76 (m, 2H), 1.41~1.35 (m, 4H), 1.15 (t, $J = 7.0$ Hz, 3H), 0.91 (t, $J = 7.0$ Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 189.2, 173.4, 169.7, 146.7, 135.3, 130.8, 127.0, 121.0, 116.5, 115.6, 67.9, 62.5, 31.1, 29.8, 27.7, 22.3, 20.8, 13.9; HPLC (Chiralpak IC, *i*-PrOH/hexane = 20/80, flow rate = 0.6 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 75.1$ min, $t_{\text{major}} = 50.4$ min, ee = 99%; $[\alpha]_{\text{D}}^{17} = -48.0$ ($c = 1.0$ in CHCl_3).


3.6 Efficient Synthesis of Polysubstituted Quinolines and Chiral 1,4-Dihydroquinolines by a Divergent Organocatalytic Cascade Approach. An Unexpected Effect of “N” Protecting Groups on Formed Products

3.6.1 Introduction

The development of new synthetic methodologies that enable to facilitate the construction of synthetically and biologically important complex molecular architectures in an efficient way from readily available starting materials is of considerable significance in chemical

synthesis. Quinolines **A** and related chiral hydroquinolines **B-D** are ‘privileged’ structures, widely distributed in natural products and synthetic substances with a broad spectrum of intriguing biological properties (Figure 3.8)³⁵. Therefore, they have long been a mainstay of organic synthesis³⁶. The general strategies for the construction of quinoline scaffolds include Friedländer³⁷, Skraup³⁸, Doebner-VonMiller³⁹, Pfitzinger⁴⁰, and Combes⁴¹ annulation reactions. Moreover, recently, new improving catalytic approaches also have been developed to meet their synthetic demand⁴². On the other hand, while significant efforts have been made toward the chiral hydroquinolines **B**^{43,44} and 1,2-dihydroquinolines **C**^{45,46}, surprisingly asymmetric synthesis of 1,4-dihydroquinoline architectures **D** remains elusive⁴⁷ and only a single example, reported by Mangeney, Alexakis and co-workers, was found by using chiral auxiliary as stereo-control⁴⁸. To our knowledge, a catalytic version has not been reported.

Figure 3.8 Skeletons of Quinolines **A** and Hydroquinolines **B-D**



In this part, we would like to report a powerful divergent organocatalytic cascade approach involving an unprecedented aza-Michael-aldol and/or aromatization sequence to both chiral 1,4-dihydroquinolines and quinolines, respectively. Notably, in the study we have made an unexpected discovery that the nature of products formed is governed by the form of *N*-protective groups⁴⁹. When aryl sulfonyls bearing electron-donating groups

are used as protecting groups for the nitrogen, a Michael-aldol-aromatization cascade proceeds to give polysubstituted quinolines; whereas chiral 1,4-dihydroquinolines are produced instead via a highly enantioselective Michael-aldol cascade with electron-withdrawing moieties such as trifilic group.


3.6.2 Results and Discussion

3.6.2.1 An unexpected organocatalytic aza-Michael-aldol-aromization cascade

1) Exploration of aza-Michael-aldol cascade: an unanticipated effect of “N”protecting groups on products formed.

As shown above, recently we have uncovered organocatalytic enantioselective allenamine involved oxa-Michael-aldol (Scheme 3.7, Eq. 1) and -Michael cascade reactions⁵⁰. In our continuing effort on expanding the efficient cascade strategy, we envisioned that the use of nitrogen-centered nucleophiles instead of “O” could lead to a new aza-Michael-aldol cascade approach to chiral 1,4-dihydroquinolines, a synthetically much less studied class of compounds (Scheme 3.7, Eq. 2).

Scheme 3.7 Organocatalytic Michael-aldol Cascade Reactions Involving Chiral Allenamines



Our initial investigation focused on the model reaction of *N*-tosyl (Ts) 2-aminobenzaldehyde **8a** with phenylpropargyl aldehyde **1a** in the presence of 30 mol% of diphenylpyrrolinol TMS ether **I** in CHCl₃ at r.t. (Table 3.7, entry 1). It is noteworthy that the Ts was selected as protecting group for “*N*” since its strong electron-withdrawing capacity enhances the NH acidity for easy ionization to produce more nucleophilic “*N*” anion for the initial Michael addition^{46a}. TLC and crude ¹H NMR analysis showed that the seemingly aza-Michael-aldol product **9aa** was produced. However, when the reaction mixture was subjected to purification on silica gel, unexpected compound **10a** was obtained instead in 54% yield (Table 3.7, entry 1). It appeared that the product **9aa** was transformed into **10a** in the presence of silica gel. The unanticipated discovery prompted us to investigate the interesting process in depth since the successful realization of the reaction could generate a new approach to the synthetically valuable polysubstituted quinolines.

14 ^f	8c	none	CHCl ₃	30	rt	4	0	32
15 ^g	8c	none	CHCl ₃	30	rt	4	0	28
16	8c	Cs ₂ CO ₃ (1.0 equiv.)	CHCl ₃	30	rt	5	0	31
17	8c	Na ₂ CO ₃ (1.0 equiv.)	CHCl ₃	30	rt	42	0	45
18	8c	K ₂ CO ₃ (1.0 equiv.)	CHCl ₃	30	rt	7	0	56
19	8c	K ₂ CO ₃ (0.1 equiv.)	CHCl ₃	10	50	16	0	99 ^h


^a Reaction conditions: unless specified, a mixture of ynal **1a** (0.16 mmol) and compound **8** (0.15 mmol) and cat. **I** (0.045 mmol) and an additive in a solvent (1.0 mL) was stirred at specified temperature for a defined time and see Experimental Section and Supporting Information. ^b Isolated yields. ^c Determined by chiral HPLC (Chiralpak AS-H column). ^d In parenthesis, ee value. ^e Calculated based on crude ¹H NMR. ^f **1a** in chloroform (1.0 mL, 0.013 mol/L) was injected into the reaction solution at 0.05 mL/min. ^g **8c** in chloroform (1.0 mL, 0.006 mol/L) was injected into the reaction solution at 0.03 mL/min. ^h After reaction completion, then silica gel (80 mg) was added and stirred at r.t. for 24 h. Et₃N (0.18 mmol) was added and stirred at r.t. for 3 h to get free deprotonated quinoline **10a**.

2) Optimization of reaction conditions.

(1) *Effect of “N” protecting groups.* It is believed that acidic silica gel promotes an aromatization process through a dehydration-deprotection of the sulfonyl group cascade sequence (Scheme 3.8). The driving force for the observed product may come from the dehydration of the aromatization tending 1,4-dihydroquinolines under an acidic condition. An aromatic sulfonamide bearing an electron-donating moiety assists the carbocation character development for a favorable dehydration-aromatization process; while a strong electron-withdrawing one will retard it. Indeed, as shown, when Tf was

used, no aromatization product **10a** was formed (Table 3.7, entry 2). Instead, a stable chiral 1,4-dihydroquinoline **9ab** (R = CF₃SO₂) was obtained in 64% yield but with only 7% ee after silica gel chromatography. We also found that the Michael-aldol cascade proceeded rapidly (5 h) presumably due to the more acidic TfNH moiety to easier produce TfN⁻. The electronic effect was further demonstrated in cases of aromatic sulfonyls bearing electron-donating (**8c**, Table 3.7, entry 3), -neutral (**8d**, Table 3.7, entry 4), and -withdrawing (**8e**, Table 3.7, entry 5) substituents. With the electron-donating (**8c**, Table 3.7, entry 3) and -neutral (**8d**, Table 3.7, entry 4) groups, quinoline **10a** was obtained exclusively when treated with silica gel. However, the electron-withdrawing *p*-NO₂C₆H₄SO₂- moiety dictated product 1,4-dihydroquinoline **9ae**. Moreover, it should be noted that there is an essential electronic balance of the “N” protecting form for nucleophilic reactivity and stabilizing the carbocation feature. It is anticipated that although a sulfonyl functionality having an electron-donating group facilitates the aromatization process, the initial conjugate addition becomes more difficult as a result of reducing NH acidity. As shown, only 19% yield of product **10a** (Table 3.7, entry 6) and no reaction was observed when Ms- (Table 3.7, entry 6,) and Cbz- (Table 3.7, entry 7) were used, respectively. We decided to choose *p*-MeOC₆H₄SO₂ as masking group to further optimize the reaction conditions.

Scheme 3.8 Organocatalytic aza-Michael-aldol-aromatization Reaction



2) *Effects of reaction temperature, solvents, reactant addition sequence and additives.* In consideration of higher temperature facilitating a reaction, accordingly we raised the reaction temperature to 60 °C (Table 3.7, entry 8). However, it became worse and a significant amount of **1a** was decomposed with only 50% conversion observed. The desired product was calculated based on 1H NMR in <10% yield. Solvents play an important role in governing the reaction rate and products formed. Therefore, we probed the effect of reaction medium on the I-promoted cascade process. Solvents including commonly used EtOAc, MeCN, toluene, MeOH and THF were screened (Table 3.7, entries 9-13). It was found that $CHCl_3$ was a preferred medium for the process. Since substrate ynal **1a** is relatively labile, its slow addition in $CHCl_3$ was employed (Table 3.7, entry 14). Disappointingly, a lower reaction yield (37%) was obtained. A similar outcome was observed in a reverse order addition (Table 3.7, entry 15). We envisioned that due to the relatively weak acidic property of NH group induced by the *para*-methoxysulfonyl group, addition of a base may assist the deprotonation to produce more nucleophilic negative nitrogen species, which may speed up the process. Accordingly, base additives Cs_2CO_3 , K_2CO_3 , and Na_2CO_3 were examined in $CHCl_3$. Indeed, it appeared that they were beneficial to the process (Table 3.7, entries 16~18). The best


condition was identified that the use of K_2CO_3 (0.1 equiv.) at 50 °C with 10 mol% catalyst loading (Table 3.7, entry 19). It should be noted that pretreatment of the reaction mixture with TEA was necessary in order to get a consistent yield of the free deprotonated quinoline product **10a**.

3) Scope of the cascade Michael-aldol-aromatization reactions.

The optimal reaction conditions, uncovered in the exploratory effort, are exploited to probe the scope of the organocatalyst **I** catalyzed Michael-aldol-aromatization cascade reactions. As revealed in Table 3.8, the tandem process serves as a general approach to the preparation of polysubstituted quinolines. In the cascade process, notably the reactions proceeded in high yield (76-99%) with a broad substrate scope. It seems that the electronic effect of aromatic alkynals **1** has limited influence on the process. The neutral (Table 3.8, entries 1, 15, 17-20, and 23), electron-donating (entries 3-4, 9, and 11) and -withdrawing (entries 2, 5-8, 12, 16, 21-22, and 24) substituents could be tolerated with a significant structural variation. A similar trend was observed for heteroaromatic alkynals, as shown with thiophen-2-yl-propynal (Table 3.8, entry 10). Furthermore, the reaction also worked well with less reactive aliphatic alkynals **1** (Table 3.8, entries 13-14) although a higher catalyst loading (20 mol%) was needed and relatively low yields were observed. On the other hand, the reaction could be applied to substrates **8** with a broad structural scope. Again the survey of the electronic effect reveals that such impact is limited. Both electron-donating (Table 3.8, entries 15-16) and -withdrawing (Table 3.8, entries 17-18) groups can be well tolerated. Moreover, significantly more hindered and

less reactive ketones instead of aldehydes in substrates **8** (e.g., $R^2 \neq H$) are compatible with this methodology (Table 3.8, entries 19-24). Structurally diverse ketones can be engaged in the process to give trisubstituted quinolines **10** in high efficiency. A trend of increase of the steric hindrance of R^1 group from H- to $-CH_3$, $(E)\text{-CH=CHC}_6\text{H}_5$ and $-\text{Ph}$ requiring more drastic conditions for aromatization is seen, but still relatively mild conditions with achieving high yields (76-98%).

Table 3.8 Substrate Scope of **I**-Catalyzed “one-pot” Synthesis of Quinolines **10**^a



Entry	R^1	R^2	X	10	$t(\text{h})$	% yield ^b
1	Ph	H	H	10a	16	99
2	4-Br- C_6H_4	H	H	10b	16	97
3	4- CH_3 - C_6H_4	H	H	10c	24	92
4	4- CH_3O - C_6H_4	H	H	10d	21	95
5	2-Cl- C_6H_4	H	H	10e	16	96
6	4-F- C_6H_4	H	H	10f	16	98
7 ^c	4-CN- C_6H_4	H	H	10g	20	91
8	4-Cl- C_6H_4	H	H	10h	16	92
9 ^d	2- CH_3O - C_6H_4	H	H	10i	16	95
10	2-thienyl	H	H	10j	16	92
11 ^d	3- CH_3O - C_6H_4	H	H	10k	16	97

12 ^c	3-CN-C ₆ H ₄	H	H	10l	13	95
13 ^c	<i>n</i> -C ₅ H ₁₁	H	H	10m	16	80
14 ^c	PhCH ₂ CH ₂	H	H	10n	20	83
15 ^d	Ph	H	6-CH ₃	10o	24	98
16 ^d	4-Cl-C ₆ H ₄	H	6-CH ₃	10p	24	90
17	Ph	H	4-Cl	10q	16	94
18	Ph	H	5-Cl	10r	16	91
19 ^c	Ph	CH ₃	H	10s	22	84
20 ^{c,e}	Ph	(<i>E</i>)-CH=CHPh	H	10t	12	80
21 ^{c,e}	4-Cl-C ₆ H ₄	(<i>E</i>)-CH=CHPh	H	10u	12	78
22 ^e	4-F-C ₆ H ₄	(<i>E</i>)-CH=CHPh	H	10v	16	76
23 ^f	Ph	Ph	H	10w	16	96
24 ^f	4-Br-C ₆ H ₄	Ph	H	10x	16	98

^a Reaction conditions: unless specified, a mixture of propynal **1** (0.16mmol), compound **8** (0.15mmol), specific amount of organocatalyst **I** and K₂CO₃ (0.015mmol) in chloroform (1.0 mL) was stirred at 50°C for a specified time. After reaction finish, then 80mg silcal gel was added and stirred at r.t. for 24h, then Et₃N (0.18mmol) was added and stirred at r.t. for 3h. ^b Isolated yields. ^c 20 mol% **I** use. ^d 15 mol% **I** used. ^e Aromatization step required heating at 50 °C for 3 h in the presence of silica gel. ^f 1.0 eq of NaHSO₄·H₂O was added and stirred at 50 °C for 24 h.


3.6.2.2 Organocatalytic enantioselective Michael-aldol cascade reactions

1) Optimization of reaction conditions for one-pot synthesis of chiral 1,4-dihydroquinolines.

Having established an efficient protocol for the preparation of quinolines using an organocatalytic aza-Michael-aldol-aromatization cascade process, we turned our attention on the optimization of aza-Michael-aldol cascade for the “one-pot” preparation of structurally different chiral 1,4-dihydroquinolines. The above study unveiled that the use of Tf as masking group led to the product without subsequent aromatization (Table 3.7, entry 2). However, only 7% ee was obtained with catalyst **I**. We found that when more bulky ketone **11a** was used, the enantioselectivity induced by (*S*)-diphenylpyrrolinol TMS ether **I** under similar reaction conditions was improved dramatically (Table 3.9, entry 1, 76% ee, 99% yield). A range of chiral α , α -diaryl prolinol silyl ether catalysts (**I-IV**) were probed accordingly. Disappointedly, the results were not encouraging (Table 3.9, entries 2-4). It should be pointed out that for the model reaction of phenylpropargyl aldehyde **1a** with 2'-(trifluoromethanesulfonyl)aminochalcone **11a**, the formed product **12a** and **11a** have the same polarity which rendered the optimization process tedious because of difficult purification. To minimize the work load, we chose more polar 3-nitrophenyl propargyl aldehyde as model substrate for the subsequent optimization. A similar level of enantioselectivity was attended with catalyst **I** (Table 3.9, entry 5, 72% ee). Gratifyingly, when C_2 -symmetry catalyst (*2R*, *5R*)-diphenylpyrrolidine **IX**⁵¹ was employed, the enantioselectivity was significantly enhanced to 87% and full conversion was obtained in only 1 h (Table 3.9, entry 6). Furthermore, solvent screening revealed that the enantioselectivity was highly solvent dependent. Remarkably, compared with dichloromethane, significantly higher enantioselectivity was observed for toluene and TBME (*t*-BuOMe) (Table 3.9, entries 8 and 10). Polar protic solvent, such as MeOH, had

a deleterious effect on both the reaction yield and enantioselectivity (Table 3.9, entry 9). Given the fast reaction rate and the practical advantage of carrying out the reaction at 0 °C, the catalyst loading was dramatically reduced to 1 mol% at 0 °C in toluene and the reaction afforded the desired product in almost quantitative yield and with 99% ee within only 3 h (Table 3.9, entry 11).

Table 3.9 Optimization of Reaction Conditions for Organocatalytic Enantioselective Aza-Michael-aldol Cascade Reactions^a



Entry	R	Cat.	Solvent	<i>t</i> (h)	Yield (%) ^b	ee (%) ^c
1	Ph	I	CH ₂ Cl ₂	12	99	76
2	Ph	II	CH ₂ Cl ₂	12	93	75
3	Ph	III	CH ₂ Cl ₂	12	92	71
4	Ph	IV	CH ₂ Cl ₂	12	84	40
5	3-NO ₂ C ₆ H ₄	I	CH ₂ Cl ₂	12	93	72
6	3-NO ₂ C ₆ H ₄	IX	CH ₂ Cl ₂	1	97	87
7	3-NO ₂ C ₆ H ₄	IX	Cl(CH ₂) ₂ Cl	1	99	94
8	3-NO ₂ C ₆ H ₄	IX	Toluene	1	99	98
9	3-NO ₂ C ₆ H ₄	IX	MeOH	6	61	75
10	3-NO ₂ C ₆ H ₄	IX	TBME	4	96	98

11 ^d	3-NO ₂ C ₆ H ₄	IX	Toluene	3	99	99
-----------------	---	-----------	---------	---	----	----


^a Reaction conditions: unless specified, a solution of alkynal **1** (0.08 mmol), organocatalyst (0.016 mmol) in a solvent (0.8 mL) was added *N*-protected 2'-aminochalcone **11a** (0.08 mmol). The resulting solution was stirred at r.t. for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralpak AS-H or IB column). ^d Reaction performed at 0 °C with 1 mol% of catalyst loading.

2) Scope of IX-catalyzed aza-Michael-aldol cascade reactions.

As revealed in Table 3.10, the optimized protocol has been demonstrated to be general to a variety of ynals **1** and 2'-(trifluoromethanesulfonyl)aminoketones **11**. Notably, the reactions served as synthetically efficient “one-pot” access to enantioenriched diverse 1,4-dihydroquinolines with creation of a quaternary stereogenic center. They proceeded in high yields (70-99%) and with excellent levels of enantioselectivities (94-99%). Both substrates ynals **1** and aminoketones **11** with significant structural variations can be tolerated. The electronic and steric factors associated with the α,β -unsaturated ketone moieties of 2'-(trifluoromethanesulfonyl)aminochalcone **11** appeared to have minimal impact on the reaction efficiencies in terms of enantioselectivity and yields (Table 3.10, entries 1-14). It is observed that electronic effect of ynals **1** play a role in the reaction rate. The electron-withdrawing groups on aromatic ynal substrates tended to accelerate the reaction illustrated by the higher turnover (1 mol% of catalyst loading and short reaction time (Table 3.10, entries 3-9, 3-9 h). While for the ynals **1** bearing neutral (Table 3.10, entries 1 and 2) or electron-donating substituents (Table 3.10, entry 2), increase of catalyst loading is necessary to assure full conversion. Less reactive aliphatic ynals can also efficiently participate in the process to

give desired products **12l** and **12m** in excellent yields (99 and 98%, respectively) and with excellent ee values (96 and 97%, respectively) except relatively slow reaction rates (Table 3.10, entries 13 and 14). We have also investigated more steric aliphatic alkynals, such as *tert*-butyl propynal and cyclopentyl propynal, however, no reaction was observed. Structural alternation of ketones besides chalcone moieties in **11**, as shown with acetophenone and benzophenone is also amenable to the protocol with high efficiency (Table 3.10, entries 15 and 16). The absolute configuration of **12g** prepared under the optimal condition was determined by X-ray crystallography (Figure 3.9).

Table 3.10 Enantioselective Organocatalytic Cascade Aza-Michael-aldol Reactions between Ynals **1** and 2'-(Trifluoromethanesulfonyl)aminoketones **11**^a




Entry	R ¹	R ²	12	<i>t</i> (h)	% yield ^b	% ee ^c
1 ^d	Ph	<i>trans</i> -PhCH=CH	12a	18	95	97
2 ^d	Ph	<i>trans</i> -4-BrC ₆ H ₄ CH=CH	12b	18	93	98
3	3-NO ₂ C ₆ H ₄	<i>trans</i> -PhCH=CH	12c	3	99	99
4	3-NO ₂ C ₆ H ₄	<i>trans</i> -4-BrC ₆ H ₄ CH=CH	12d	3	99	98
5	4-ClC ₆ H ₄	<i>trans</i> -PhCH=CH	12e	4	99	98
6	4-CNC ₆ H ₄	<i>trans</i> -4-BrC ₆ H ₄ CH=CH	12f	3	91	99
7	4-CNC ₆ H ₄	<i>trans</i> -3-MeOC ₆ H ₄ CH=CH	12g	8	98	98

8	4-BrC ₆ H ₄	<i>trans</i> -4-NO ₂ C ₆ H ₄ CH=CH	12h	5	96	98
9	4-BrC ₆ H ₄	<i>trans</i> -2-MeC ₆ H ₄ CH=CH	12i	9	99	97
10 ^d	4-MeOC ₆ H ₄	<i>trans</i> -PhCH=CH	12j	18	92	96
11	2-thienyl	<i>trans</i> -PhCH=CH	12k	9	90	96
12 ^d	PhCH ₂ CH ₂	<i>trans</i> -PhCH=CH	12l	12	99	96
13 ^e	<i>n</i> -C ₅ H ₁₁	<i>trans</i> -PhCH=CH	12m	20	98	97
14 ^f	<i>trans</i> -PhCH=CH	<i>trans</i> -PhCH=CH	12n	18	70	94
15 ^f	Ph	CH ₃	12o	18	96	95
16 ^{e,g}	3-NO ₂ C ₆ H ₄	Ph	12p	3	84	99

^a Reaction conditions: unless specified, a solution of alkynal **1** (0.08mmol), organocatalyst **IX** (1~20 mol%) in toluene (0.8mL) was added 2'-(trifluoromethanesulfonyl)aminoketones **11** (0.08mmol). The resulting solution was stirred at 0°C for a specified time. ^b Isolated yields. ^c Determined by chiral HPLC analysis (Chiralpak AS-H or IB column). ^d 5 mol% of catalyst **IX** used. ^e 20 mol% of catalyst **IX** used. ^f 8 mol% of catalyst **IX** used. ^g Reaction carried out at r.t..

Figure 3.9 X-ray Crystal Structure of Compound **12g**




3.6.3 Conclusion

While organocatalytic cascade reactions have been established as a viable route in organic synthesis¹, the examples of combining powerful divergent synthesis and organocatalytic cascade strategy are rare⁵². In this investigation, we have illustrated a divergent organocatalytic cascade approach to synthetic valuable polysubstituted quinolines and highly enantioenriched 1,4-dihydroquinolines. The key to the fate of products formed depends on the protecting sulfonyls used for the “*N*” initiated an aza-Michael-aldol cascade. The electron-donating aryl sulfonamides facilitate the dehydration-aromatization of the aza-Michael-aldol adducts to give polysubstituted quinolines in good to high yields (76-99%) with a broad substrate scope. However, when strong electron-withdrawing Tf (CF₃SO₂-) is used, chiral 1,4-dihydroquinolines products are produced. Notably, the process, efficiently catalyzed by (2*R*, 5*R*)-diphenylpyrrolidine


using as low as 1 mol% loading, leads to the first catalytic enantioselective access to the chiral scaffold with excellent level of enantioselectivity (94-99% ee). The studies show a scarce example of the electronic effect on the nature of reaction course and products⁴⁹.

3.6.4 Supporting Information


General Procedure for aza-Michael-aldol-aromatization Cascade Reactions (Table 3.8): To a solution of ynal **1** (0.16 mmol), organocatalyst **I** (10~20 mol%) in chloroform (1.0 mL) was added compound **8** (0.15 mmol) and K₂CO₃ (0.015 mmol). The resulting solution was stirred at 50 °C for a specified time. After reaction finish monitored by TLC, then silica gel (80 mg) was added and stirred at r.t. for 24h, then Et₃N (0.18 mmol) was added and stirred at r.t. for 3h. The reaction mixture was directly purified by column chromatography, eluted with hexane/EtOAc to afford the desired product.




2-phenylquinoline-3-carbaldehyde (10a) (Table 3.8, Entry 1): The title compound was prepared according to the general procedure, as described above in 99% yield. ¹H NMR (CDCl₃, 500 MHz): δ 10.19 (s, 1H), 8.85 (s, 1H), 8.22 (d, *J* = 8.5 Hz, 1H), 8.02 (d, *J* = 8.5 Hz, 1H), 7.90~7.86 (m, 1H), 7.70~7.69 (m, 2H), 7.66~7.63 (m, 1H), 7.59~7.54 (m, 3H); ¹³C NMR (CDCl₃, 75MHz): δ 191.5, 160.3, 149.6, 138.1, 137.8, 132.6, 130.2, 129.6, 129.43, 129.38, 128.7, 127.6, 127.5, 126.4.




2-(4-bromophenyl)quinoline-3-carbaldehyde (10b) (Table 3.8, Entry 2): The title compound was prepared according to the general procedure, as described above in 97% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.17 (s, 1H), 8.84 (s, 1H), 8.19 (d, $J = 8.5$ Hz, 1H), 8.02 (d, $J = 8.0$ Hz, 1H), 7.90~7.87 (m, 1H), 7.70 (d, $J = 8.5$ Hz, 1H), 7.67~7.64 (m, 1H), 7.57 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 190.9, 158.9, 149.6, 138.6, 136.7, 132.8, 131.9, 131.8, 129.6, 129.4, 127.7, 127.5, 126.4, 124.1.




2-p-tolylquinoline-3-carbaldehyde (10c) (Table 3.8, Entry 3): The title compound was prepared according to the general procedure, as described above in 92% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.19 (s, 1H), 8.82 (s, 1H), 8.20 (d, $J = 8.5$ Hz, 1H), 8.00 (d, $J = 8.0$ Hz, 1H), 7.87~7.84 (m, 1H), 7.63~7.58 (m, 3H), 7.37 (d, $J = 8.0$ Hz, 2H), 2.46 (s, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.7, 160.3, 149.6, 139.5, 138.0, 134.9, 132.5, 130.2, 129.5, 129.4, 127.7, 127.3, 126.3, 21.3.




2-(4-methoxyphenyl)quinoline-3-carbaldehyde (10d) (Table 3.8, Entry 4): The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.19 (s, 1H), 8.80 (s, 1H), 8.18 (d, $J = 8.5$ Hz, 1H), 7.99 (d, $J = 8.5$ Hz, 1H), 7.86~7.83 (m, 1H), 7.64 (d, $J = 8.5$ Hz, 2H), 7.62~7.59 (m, 1H), 7.08 (d, $J = 8.5$ Hz, 2H), 3.90 (s, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.7, 160.7, 159.8, 149.7, 138.2, 132.5, 131.7, 130.2, 129.5, 129.4, 127.6, 127.2, 126.2, 114.2, 55.4.




2-(2-chlorophenyl)quinoline-3-carbaldehyde (10e) (Table 3.8, Entry 5): The title compound was prepared according to the general procedure, as described above in 96% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.95 (s, 1H), 8.86 (s, 1H), 8.21 (d, $J = 8.5$ Hz, 1H), 8.04 (d, $J = 8.0$ Hz, 1H), 7.90~7.87 (m, 1H), 7.69~7.65 (m, 1H), 7.61~7.59 (m, 1H), 7.54~7.51 (m, 1H), 7.49~7.47 (m, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 190.4, 158.0, 149.7, 137.5, 136.9, 133.1, 132.7, 131.4, 130.6, 129.64, 129.61, 129.5, 127.9, 127.7, 127.4, 126.8.




2-(4-fluorophenyl)quinoline-3-carbaldehyde (10f) (Table 3.8, Entry 6): The title compound was prepared according to the general procedure, as described above in 98% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.17 (s, 1H), 8.83 (s, 1H), 8.19 (d, $J = 8.5$ Hz, 1H), 8.01 (d, $J = 8.0$ Hz, 1H), 7.88 (t, $J = 8.0$ Hz, 1H), 7.69~7.62 (m, 3H), 7.26 (t, $J = 8.5$ Hz, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.1, 165.3, 161.9, 159.0, 149.5, 138.5, 133.9, 132.8, 132.2, 132.1, 129.5, 129.4, 127.6, 126.4, 116.0, 115.7.




4-(3-formylquinolin-2-yl)benzonitrile (10g) (Table 3.8, Entry 7): The title compound was prepared according to the general procedure, as described above in 91% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.16 (s, 1H), 8.88 (s, 1H), 8.21 (d, $J = 8.5$ Hz, 1H), 8.05 (d, $J = 8.5$ Hz, 1H), 7.94~7.91 (m, 1H), 7.86 (d, $J = 8.0$ Hz, 2H), 7.81 (d, $J = 8.0$ Hz, 2H), 7.70 (t, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 190.2, 157.8, 149.5, 142.4, 139.4, 133.2, 132.4, 130.8, 129.7, 129.5, 128.2, 127.4, 126.6, 118.4, 113.2.



2-(4-chlorophenyl)quinoline-3-carbaldehyde (10h) (Table 3.8, Entry 8): The title compound was prepared according to the general procedure, as described above in 92% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.17 (s, 1H), 8.84 (s, 1H), 8.19 (d, $J = 8.5$ Hz, 1H), 8.01 (d, $J = 8.5$ Hz, 1H), 7.90~7.87 (m, 1H), 7.66~7.63 (m, 3H), 7.54 (d, $J = 8.5$ Hz, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 190.9, 158.9, 149.6, 138.6, 136.2, 135.8, 132.8, 131.5, 129.6, 129.4, 128.9, 127.7, 127.5, 126.4.




2-(2-methoxyphenyl)quinoline-3-carbaldehyde (10i) (Table 3.8, Entry 9): The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.99 (s, 1H), 8.77 (s, 1H), 8.20 (d, $J = 8.5$ Hz, 1H), 8.00 (d, $J = 8.0$ Hz, 1H), 7.85~7.82 (m, 1H), 7.69 (dd, $J_1 = 1.5$ Hz, $J_2 = 7.5$ Hz, 1H), 7.63~7.60 (m, 1H), 7.52~7.49 (m, 1H), 7.20 (t, $J = 8$ Hz, 1H), 7.01 (d, $J = 8.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.5, 157.6, 156.6, 150.1, 136.2, 132.0, 131.4, 131.2, 129.6, 129.4, 128.2, 127.3, 126.9, 126.6, 121.7, 110.7, 55.3.




2-(thiophen-2-yl)quinoline-3-carbaldehyde (10j) (Table 3.8, Entry 10): The title compound was prepared according to the general procedure, as described above in 92%

yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.45 (s, 1H), 8.77 (s, 1H), 8.15 (d, $J = 9.0$ Hz, 1H), 7.96 (d, $J = 8.5$ Hz, 1H), 7.86~7.83 (m, 1H), 7.61~7.58 (m, 2H), 7.353~7.346 (m, 1H), 7.23~7.21 (m, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.2, 153.0, 149.6, 141.2, 138.5, 132.7, 130.8, 129.6, 129.3, 128.0, 127.5, 127.4, 126.1.




2-(3-methoxyphenyl)quinoline-3-carbaldehyde (10k) (Table 3.8, Entry 11): The title compound was prepared according to the general procedure, as described above in 97% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.18 (s, 1H), 8.84 (s, 1H), 8.21 (d, $J = 8.5$ Hz, 1H), 8.02 (d, $J = 8.0$ Hz, 1H), 7.89~7.86 (m, 1H), 7.65~7.62 (m, 1H), 7.46 (t, $J = 8.0$ Hz, 1H), 7.26 (s, 1H), 7.19 (d, $J = 7.5$ Hz, 1H), 7.09~7.07 (m, 1H), 3.90 (s, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.5, 160.2, 159.9, 149.5, 139.1, 138.0, 132.6, 129.7, 129.6, 129.4, 127.7, 127.5, 126.4, 122.8, 115.4, 115.3, 55.4.




3-(3-formylquinolin-2-yl)benzonitrile (10l) (Table 3.8, Entry 12): The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.17 (s, 1H), 8.88 (s, 1H), 8.21 (d, $J = 8.5$ Hz, 1H), 8.05 (d,

$J = 7.5$ Hz, 2H), 7.94~7.91 (m, 1H), 7.89~7.87 (m, 1H), 7.84~7.82 (m, 1H), 7.71~7.66 (m, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 190.2, 157.4, 149.5, 139.5, 139.3, 134.4, 133.5, 133.2, 132.7, 129.6, 129.4, 129.4, 128.2, 127.4, 126.6, 118.2, 113.2.




2-pentylquinoline-3-carbaldehyde (10m) (Table 3.8, Entry 13): The title compound was prepared according to the general procedure, as described above in 80% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.39 (s, 1H), 8.62 (s, 1H), 8.08 (d, $J = 8.5$ Hz, 1H), 7.93 (d, $J = 8.5$ Hz, 1H), 7.83 (t, $J = 8.0$ Hz, 1H), 7.57 (t, $J = 7.5$ Hz, 1H), 3.35 (t, $J = 8.0$ Hz, 2H), 1.81~1.75 (m, 2H), 1.49~1.43 (m, 2H), 1.42~1.36 (m, 2H), 0.91 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.1, 162.5, 149.4, 142.0, 132.5, 129.1, 128.9, 127.6, 126.8, 126.1, 36.4, 31.9, 30.1, 22.5, 14.0.




2-phenethylquinoline-3-carbaldehyde (10n) (Table 3.8, Entry 14): The title compound was prepared according to the general procedure, as described above in 83% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.27 (s, 1H), 8.60 (s, 1H), 8.13 (d, $J = 8.5$ Hz, 1H), 7.95 (d, $J = 8.0$ Hz, 1H), 7.88~7.85 (m, 1H), 7.60 (t, $J = 7.5$ Hz, 1H), 7.33~7.28 (m, 4H), 7.21 (t, $J = 7.0$ Hz, 1H), 3.69~3.66 (m, 2H), 3.13 (t, $J = 8.0$ Hz, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.0, 161.0, 149.3, 142.7, 141.4, 132.6, 129.03, 128.97, 128.6, 128.4, 127.8, 127.0,


126.1, 38.2, 35.7.




5-methyl-2-phenylquinoline-3-carbaldehyde (10o) (Table 3.8, Entry 15): The title compound was prepared according to the general procedure, as described above in 98% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.20 (s, 1H), 9.02 (s, 1H), 8.05 (d, $J = 8.5$ Hz, 1H), 7.76~7.73 (m, 1H), 7.70~7.68 (m, 2H), 7.58~7.53 (m, 3H), 7.44 (d, $J = 7.0$ Hz, 1H), 2.80 (s, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.7, 159.9, 150.1, 137.7, 136.9, 134.6, 132.5, 130.3, 129.3, 128.7, 127.8, 127.0, 125.9, 18.7.




2-(4-chlorophenyl)-5-methylquinoline-3-carbaldehyde (10p) (Table 3.8, Entry 16): The title compound was prepared according to the general procedure, as described above in 90% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.18 (s, 1H), 9.01 (s, 1H), 8.03 (d, $J = 8.5$ Hz, 1H), 7.76 (t, $J = 8.0$ Hz, 1H), 7.64 (d, $J = 8.5$ Hz, 2H), 7.54 (d, $J = 8.0$ Hz, 2H), 7.46 (d, $J = 7.0$ Hz, 1H), 2.79 (s, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.2, 158.5, 150.1, 136.9, 136.2, 135.8, 135.1, 132.7, 131.6, 128.9, 128.0, 127.8, 126.9, 126.0, 18.7.



7-chloro-2-phenylquinoline-3-carbaldehyde (10q) (Table 3.8, Entry 17): The title compound was prepared according to the general procedure, as described above in 94% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.16 (s, 1H), 8.80 (s, 1H), 8.20 (s, 1H), 7.94 (d, $J = 9.0$ Hz, 1H), 7.68~7.67 (m, 2H), 7.58~7.56 (m, 4H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.1, 161.3, 149.8, 138.8, 137.9, 137.3, 130.5, 130.2, 129.7, 128.8, 128.7, 127.7, 124.7.




6-chloro-2-phenylquinoline-3-carbaldehyde (10r) (Table 3.8, Entry 18): The title compound was prepared according to the general procedure, as described above in 91% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.17 (s, 1H), 8.74 (s, 1H), 8.14 (d, $J = 9.0$ Hz, 1H), 7.98 (s, 1H), 7.79 (d, $J = 9.0$ Hz, 1H), 7.69~7.67 (m, 2H), 7.57~7.56 (m, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 191.1, 160.4, 147.9, 137.4, 137.1, 133.4, 133.3, 131.2, 130.2, 129.6, 128.8, 128.2, 127.7, 126.9.




4-methyl-2-phenylquinoline-3-carbaldehyde (10s) (Table 3.8, Entry 19): The title

compound was prepared according to the general procedure, as described above in 84% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.12 (s, 1H), 8.25 (d, $J = 8.0$ Hz, 1H), 8.19 (d, $J = 8.5$ Hz, 1H), 7.85~7.82 (m, 1H), 7.66~7.63 (m, 3H), 7.53~7.52 (m, 3H), 3.02 (s, 3H); ^{13}C NMR (CDCl_3 , 75MHz): δ 194.2, 160.4, 148.04, 147.98, 139.0, 131.6, 130.3, 130.1, 129.2, 128.7, 127.1, 127.0, 125.2, 14.8.




(E)-2-phenyl-4-styrylquinoline-3-carbaldehyde (10t) (Table 3.8, Entry 20): The title compound was prepared according to the general procedure, as described above in 80% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.21 (s, 1H), 8.41 (d, $J = 8.5$ Hz, 1H), 8.23 (d, $J = 8.5$ Hz, 1H), 7.93 (d, $J = 16.5$ Hz, 1H), 7.89~7.86 (m, 1H), 7.67 (d, $J = 7.5$ Hz, 4H), 7.64~7.61 (m, 1H), 7.56~7.52 (m, 3H), 7.47~7.44 (m, 2H), 7.40~7.37 (m, 1H), 6.89 (d, $J = 17$ Hz, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 193.1, 160.4, 149.2, 148.2, 139.5, 139.0, 136.3, 132.0, 130.1, 129.2, 128.9, 128.6, 127.2, 127.1, 125.4, 125.1, 123.1.




(E)-2-(4-chlorophenyl)-4-styrylquinoline-3-carbaldehyde (10u) (Table 3.8, Entry 21):

The title compound was prepared according to the general procedure, as described above in 78% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.23 (s, 1H), 8.38 (d, $J = 8.5$ Hz, 1H), 8.20 (d, $J = 8.5$ Hz, 1H), 7.92~7.86 (m, 2H), 7.66~7.59 (m, 5H), 7.51 (d, $J = 8.5$ Hz, 2H), 7.45 (t, $J = 7.5$ Hz, 2H), 7.39 (t, $J = 7.5$ Hz, 1H), 6.88 (d, $J = 16.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 192.6, 158.9, 149.0, 148.6, 140.1, 137.6, 136.1, 135.5, 132.2, 131.3, 130.1, 129.0, 128.9, 128.8, 127.4, 127.2, 127.0, 125.5, 125.1, 122.5.




(E)-2-(4-fluorophenyl)-4-styrylquinoline-3-carbaldehyde (10v) (Table 3.8, Entry 22):

The title compound was prepared according to the general procedure, as described above in 76% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.22 (s, 1H), 8.39 (d, $J = 8.5$ Hz, 1H), 8.21 (d, $J = 8.5$ Hz, 1H), 7.91 (d, $J = 16.5$ Hz, 1H), 7.88~7.86 (m, 1H), 7.67~7.61 (m, 5H), 7.45 (t, $J = 8.0$ Hz, 2H), 7.39 (t, $J = 7.5$ Hz, 1H), 7.23 (t, $J = 8.5$ Hz, 2H), 6.89 (d, $J = 16.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 75MHz): δ 192.8, 165.1, 159.1, 149.0, 148.5, 139.9, 136.2, 135.2, 132.1, 132.0, 131.9, 130.0, 129.0, 128.9, 127.3, 127.1, 127.0, 125.4, 125.2, 122.7, 115.8, 115.5.



2,4-diphenylquinoline-3-carbaldehyde (10w) (Table 3.8, Entry 23): The title compound was prepared according to the general procedure, as described above in 96% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.04 (s, 1H), 8.25 (d, $J = 8.5$ Hz, 1H), 7.86~7.82 (m, 1H), 7.66~7.65 (m, 2H), 7.61 (d, $J = 8.5$ Hz, 1H), 7.57~7.49 (m, 7H), 7.39~7.37 (m, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 192.5, 159.0, 151.9, 148.5, 139.5, 134.6, 131.9, 129.8, 129.7, 128.9, 128.6, 128.4, 128.3, 127.3, 126.6, 126.3.



2-(4-bromophenyl)-4-phenylquinoline-3-carbaldehyde (10x) (Table 3.8, Entry 24): The title compound was prepared according to the general procedure, as described above in 98% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 10.01 (s, 1H), 8.22 (d, $J = 8.0$ Hz, 1H), 7.85 (t, $J = 7.0$ Hz, 1H), 7.64~7.61 (m, 3H), 7.56~7.50 (m, 6H), 7.39~7.38 (m, 2H); ^{13}C NMR (CDCl_3 , 75MHz): δ 192.2, 157.5, 152.9, 148.5, 138.7, 134.1, 132.1, 131.5, 131.2, 129.8, 128.9, 128.5, 127.5, 127.2, 126.3, 123.4.

General procedure for the preparation of substrates 11:


Substrates were synthesized via the following two steps.

Step 1: Claisen-Schmidt condensation of benzaldehyde with 2'-amino acetophenone to trans-2'-amino-chalcone⁵³

A mixture of 2'-amino acetophenone (3.0 mmol, 0.36 mL) and benzaldehyde (3.0 mmol) in ethanol was added sodium hydroxide (15%, 0.9 mL) under stirring with ice-bath. Then the solution was stirred at r.t. for 24h. The reaction mixture was neutralized with 1M HCl. The solid was filtrated and crystallized from ethanol.

Step 2: Synthesis of corresponding trifluoromethanesulfonamides⁵⁴


A solution of trans-2'-amino-chalcone (1.5 mmol) and triethylamine (0.23 mL, 1.65 mmol) in dry CH₂Cl₂ at -78 °C was treated dropwise with trifluoromethanesulfonic anhydride (0.28 mL, 1.65 mmol). The mixture was stirred at -78 °C for 1.5h and then at room temperature for 2.5h. The reaction was partitioned between dichloromethane and water. The aqueous phase was extracted with dichloromethane, and the combined organic phases was washed with brine, dried and concentrated. The residue was purified by column chromatography, eluted with hexane/EtOAc to afford the desired product.



(E)-N-(2-cinnamoylphenyl)-1,1,1-trifluoromethanesulfonamide


The title compound was prepared according to the general procedure, as described above in 57% yield for two steps. ¹H NMR (CDCl₃, 500 MHz): δ 8.07 (d, *J* = 8.0 Hz, 1H), 7.92

(d, $J = 15.5$ Hz, 1H), 7.83 (d, $J = 8.5$ Hz, 1H), 7.68~7.67 (m, 2H), 7.64~7.58 (m, 2H), 7.48~7.45 (m, 3H), 7.33 (t, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.9, 147.3, 138.3, 134.9, 134.1, 131.3, 130.9, 129.0, 128.8, 124.6, 124.0, 123.6, 121.0, 120.8, 119.9, 118.4, 115.9.



(E)-N-(2-(3-(4-bromophenyl)acryloyl)phenyl)-1,1,1-trifluoromethanesulfonamide


The title compound was prepared according to the general procedure, as described above in 52% yield for two steps. ^1H NMR (CDCl_3 , 500 MHz): δ 8.05 (d, $J = 8.0$ Hz, 1H), 7.84~7.81 (m, 2H), 7.63~7.56 (m, 4H), 7.52 (d, $J = 8.5$ Hz, 2H), 7.33 (t, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.7, 145.8, 138.4, 135.1, 133.1, 132.4, 130.8, 130.1, 125.8, 124.6, 124.1, 123.6, 121.6, 121.0, 120.2, 118.5, 115.9.



(E)-1,1,1-trifluoro-N-(2-(3-o-tolylacryloyl)phenyl)methanesulfonamide


The title compound was prepared according to the general procedure, as described above in 47% yield for two steps. ^1H NMR (CDCl_3 , 500 MHz): δ 8.23 (d, $J = 15.5$ Hz, 1H), 8.08 (d, $J = 7.5$ Hz, 1H), 7.83 (d, $J = 8.5$ Hz, 1H), 7.72 (d, $J = 7.5$ Hz, 1H), 7.62 (t, $J =$

7.5 Hz, 1H), 7.53 (d, $J = 15.0$ Hz, 1H), 7.37~7.32 (m, 2H), 7.30~7.26 (m, 2H), 2.51 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.0, 144.9, 138.9, 138.5, 135.0, 133.2, 131.1, 130.9, 126.6, 126.5, 124.6, 124.2, 123.6, 122.0, 121.1, 120.1, 118.5, 115.9, 19.7.



(E)-1,1,1-trifluoro-N-(2-(3-(4-nitrophenyl)acryloyl)phenyl)methanesulfonamide


The title compound was prepared according to the general procedure, as described above in 41% yield for two steps. ^1H NMR (CDCl_3 , 500 MHz): δ 8.30 (d, $J = 8.0$ Hz, 2H), 8.08 (d, $J = 8.0$ Hz, 1H), 7.91 (d, $J = 15.5$ Hz, 1H), 7.85~7.82 (m, 3H), 7.72~7.64 (m, 2H), 7.36 (t, $J = 7.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.3, 149.0, 143.8, 140.2, 138.6, 135.6, 131.0, 129.3, 125.0, 124.8, 124.3, 123.9, 121.0, 120.4, 118.4.



(E)-1,1,1-trifluoro-N-(2-(3-(3-methoxyphenyl)acryloyl)phenyl)methanesulfonamide

The title compound was prepared according to the general procedure, as described above in 69% yield for two steps. ^1H NMR (CDCl_3 , 500 MHz): δ 8.06 (d, $J = 8.0$ Hz, 1H), 7.87~7.81 (m, 2H), 7.62~7.55 (m, 2H), 7.38~7.31 (m, 2H), 7.26 (d, $J = 7.5$ Hz, 1H), 7.16 (s, 1H), 7.02~7.00 (m, 1H), 3.86 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.9, 160.0,


147.3, 138.4, 135.5, 135.0, 130.9, 130.1, 124.6, 124.2, 123.6, 121.4, 121.3, 121.0, 120.1, 118.5, 117.1, 115.9, 113.8, 55.3.



***N*-(2-acetylphenyl)-1,1,1-trifluoromethanesulfonamide**


The title compound was prepared according to the general procedure, as described above for step 2 in 75% yield. ¹H NMR (CDCl₃, 500 MHz): δ 7.95 (d, *J* = 7.5 Hz, 1H), 7.77 (d, *J* = 8.5 Hz, 1H), 7.58 (t, *J* = 8.0 Hz, 1H), 7.26 (t, *J* = 7.5 Hz, 1H), 2.69 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 203.2, 138.1, 135.5, 132.0, 124.5, 123.6, 122.6, 121.0, 119.4, 118.4, 115.9, 28.0.

General Procedure for Cascade aza-Michael-aldol Reaction (Table 3.10): To a solution of ynal **1** (0.08 mmol), organocatalyst **IX** (1 mol%) in toluene (0.8 mL) was added 2'-NHTf ketone **11** (0.08 mmol). The resulting solution was stirred at 0 °C for a specified time. Then the reaction mixture was directly purified by column chromatography, eluted with hexane/EtOAc to afford the desired product and ees are determined by chiral HPLC analysis.



(R)-(E)-4-hydroxy-2-phenyl-4-styryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12a) (Table 3.10, Entry 1):


The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.35 (s, 1H), 7.94 (d, $J = 7.5$ Hz, 1H), 7.58~7.51 (m, 7H), 7.47~7.44 (m, 1H), 7.37 (d, $J = 7.0$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 2H), 7.26~7.22 (m, 1H), 6.58 (d, $J = 16.0$ Hz, 1H), 6.45 (d, $J = 16.0$ Hz, 1H), 5.50 (s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.1, 153.3, 137.2, 135.9, 133.9, 133.7, 131.3, 131.1, 130.8, 130.3, 128.7, 128.5, 128.4, 128.3, 128.1, 127.0, 125.9, 122.6, 120.2, 117.6, 72.3; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 10/90, flow rate = 0.45 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 19.16$ min, $t_{\text{major}} = 24.81$ min, ee = 97%; $[\alpha]_{\text{D}}^{30} = +117.1$ ($c = 1.0$ in CHCl_3).



(R)-(E)-4-(4-bromostyryl)-4-hydroxy-2-phenyl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12b) (Table 3.10, Entry 2):


The title compound was prepared according to the general procedure as described above in 93% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.34 (s, 1H), 7.92 (d, $J = 7.5$ Hz, 1H), 7.58~7.55 (m, 1H), 7.52~7.44 (m, 7H), 7.41 (d, $J = 8.5$ Hz, 2H), 7.22 (d, $J = 8.5$ Hz, 2H), 6.48 (d, $J = 16.0$ Hz, 1H), 6.42 (d, $J = 15.5$ Hz, 1H), 5.53 (s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.2, 153.5, 137.1, 134.9, 133.8, 131.8, 131.7, 131.4, 130.7, 129.2, 128.7,

128.5, 128.4, 125.9, 122.6, 122.1, 120.1, 117.6, 72.3; HPLC (Chiralpak IB, *i*-PrOH/hexane = 15/85, flow rate = 0.45 mL/min, λ = 254nm): t_{minor} = 11.90 min, t_{major} = 19.76 min, ee = 98%; $[\alpha]_{\text{D}}^{30}$ = +97.1 (c = 1.0 in CHCl₃).




(R)-(E)-4-hydroxy-2-(3-nitrophenyl)-4-styryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12c) (Table 3.10, Entry 3):

The title compound was prepared according to the general procedure as described above in 99% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.35 (s, 1H), 8.43 (d, J = 8.0 Hz, 1H), 8.38 (s, 1H), 7.94 (d, J = 7.0 Hz, 1H), 7.85 (d, J = 6.5 Hz, 1H), 7.73 (t, J = 8.0 Hz, 1H), 7.55~7.50 (m, 3H), 7.37~7.35 (m, 2H), 7.30 (t, J = 7.5 Hz, 2H), 7.26~7.25 (m, 1H), 6.54 (d, J = 16.0 Hz, 1H), 6.45 (d, J = 15.5 Hz, 1H), 5.23 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz): δ 191.6, 149.7, 147.9, 136.6, 135.5, 133.2, 132.5, 130.8, 130.5, 129.6, 129.0, 128.6, 128.5, 128.3, 126.9, 126.2, 125.8, 122.5, 120.9, 116.6, 72.2. HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 15/85, flow rate = 0.55 mL/min, λ = 254nm): t_{minor} = 24.82 min, t_{major} = 47.05 min, ee = 99%; $[\alpha]_{\text{D}}^{30}$ = +104.9 (c = 1.0 in CHCl₃).



(R)-(E)-4-(4-bromostyryl)-4-hydroxy-2-(3-nitrophenyl)-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12d) (Table 3.10, Entry 4):


The title compound was prepared according to the general procedure, as described above in 99% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.35 (s, 1H), 8.43 (d, $J = 7.0$ Hz, 1H), 8.38 (s, 1H), 7.93 (d, $J = 7.5$ Hz, 1H), 7.85 (d, $J = 7.0$ Hz, 1H), 7.75~7.72 (m, 1H), 7.56~7.50 (m, 3H), 7.43~7.41 (m, 2H), 7.22 (d, $J = 8.5$ Hz, 2H), 6.47~6.41 (m, 2H), 5.26 (s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.6, 150.0, 148.0, 136.5, 135.5, 134.5, 133.3, 132.5, 131.8, 131.3, 129.8, 129.7, 129.1, 128.8, 128.5, 126.3, 126.0, 122.6, 122.3, 120.1, 117.5, 72.2; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 15/85, flow rate = 0.55 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 25.76$ min, $t_{\text{major}} = 44.58$ min, ee = 98%; $[\alpha]_{\text{D}}^{30} = +59.5$ (c = 1.0 in CHCl_3).



(R)-(E)-2-(4-chlorophenyl)-4-hydroxy-4-styryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12e) (Table 3.10, Entry 5):


The title compound was prepared according to the general procedure as described above in 99% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.35 (s, 1H), 7.93 (d, $J = 7.5$ Hz, 1H), 7.53~7.44 (m, 7H), 7.35 (d, $J = 7.5$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 2H), 7.25~7.22 (m, 1H), 6.54 (d, $J = 15.5$ Hz, 1H), 6.43 (d, $J = 16.0$ Hz, 1H), 5.39 (s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.5, 151.8, 137.9, 137.1, 135.8, 134.5, 133.6, 130.9, 130.5, 129.3, 128.8, 128.6, 128.4, 128.2, 127.0, 126.1, 122.6, 120.2, 117.6, 72.3; HPLC (Chiralpak IB, *i*-

PrOH/hexane = 5/95, flow rate = 0.40 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 19.14\text{ min}$, $t_{\text{major}} = 37.65\text{ min}$, ee = 98%; $[\alpha]_{\text{D}}^{30} = +123.4$ (c = 1.0 in CHCl_3).




(R)-4-(4-(4-bromostyryl)-3-formyl-4-hydroxy-1-(trifluoromethylsulfonyl)-1,4-dihydroquinolin-2-yl)benzonitrile (12f) (Table 3.10, Entry 6):

The title compound was prepared according to the general procedure as described above in 91% yield. $^1\text{H NMR}$ (CDCl_3 , 500 MHz): δ 9.32 (s, 1H), 7.92 (d, $J = 7.5\text{ Hz}$, 1H), 7.81 (d, $J = 8.0\text{ Hz}$, 2H), 7.64 (d, $J = 8.0\text{ Hz}$, 2H), 7.55~7.48 (m, 3H), 7.41 (d, $J = 8.0\text{ Hz}$, 2H), 7.20 (d, $J = 8.0\text{ Hz}$, 2H), 6.44 (d, $J = 16.0\text{ Hz}$, 1H), 6.40 (d, $J = 16.0\text{ Hz}$, 1H), 5.26 (s, 1H); $^{13}\text{C NMR}$ (CDCl_3 , 125 MHz): δ 191.8, 150.5, 136.5, 135.3, 135.2, 134.5, 133.4, 132.2, 132.1, 131.8, 131.3, 130.1, 129.7, 129.1, 128.8, 128.5, 126.3, 125.4, 123.7, 122.5, 122.3, 120.0, 117.5, 115.3, 72.2; HPLC (Chiralpak IB, *i*-PrOH/hexane = 15/85, flow rate = 0.45 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 27.43\text{ min}$, $t_{\text{major}} = 93.65\text{ min}$, ee = 99%; $[\alpha]_{\text{D}}^{30} = +142.2$ (c = 1.0 in CHCl_3).



(R)-(E)-4-(3-formyl-4-hydroxy-4-(3-methoxystyryl)-1-(trifluoromethylsulfonyl)-1,4-dihydroquinolin-2-yl)benzonitrile (12g) (Table 3.10, Entry 7):


The title compound was prepared according to the general procedure as described above in 98% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.32 (s, 1H), 7.93 (d, $J = 7.5$ Hz, 1H), 7.81 (d, $J = 8.0$ Hz, 2H), 7.65 (d, $J = 7.5$ Hz, 2H), 7.55~7.47 (m, 3H), 7.21 (t, $J = 7.5$ Hz, 1H), 6.94 (d, $J = 8.0$ Hz, 1H), 6.87 (s, 1H), 6.81~6.79 (m, 1H), 6.50 (d, $J = 16.0$ Hz, 1H), 6.42 (d, $J = 16.0$ Hz, 1H), 5.24 (s, 1H), 3.79 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.7, 159.8, 150.4, 137.0, 136.7, 135.3, 133.3, 132.1, 130.9, 130.8, 129.6, 129.0, 128.8, 128.6, 126.3, 122.5, 120.1, 119.6, 117.5, 115.2, 114.2, 113.6, 112.1, 72.2, 55.2; HPLC (Chiralpak IB, *i*-PrOH/hexane = 15/85, flow rate = 0.50 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 79.73$ min, $t_{\text{major}} = 65.06$ min, ee = 98%; $[\alpha]_{\text{D}}^{30} = +156.1$ (c = 1.0 in CHCl_3).



(R)-(E)-2-(4-bromophenyl)-4-hydroxy-4-(4-nitrostyryl)-1-(trifluoromethylsulfonyl)-1,4-dihydroquinolin-2-yl

1,4-dihydroquinoline-3-carbaldehyde (12h) (Table 3.10, Entry 8):

The title compound was prepared according to the general procedure, as described above in 96% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.35 (s, 1H), 8.17~8.14 (m, 2H), 7.93~7.90 (m, 1H), 7.67 (d, $J = 8.7$ Hz, 2H), 7.57~7.54 (m, 1H), 7.53~7.46 (m, 4H), 7.39 (d, $J = 8.4$ Hz, 2H), 6.60~6.54 (m, 2H), 5.56 (s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.6, 152.5, 147.4, 142.2, 136.5, 135.2, 133.9, 133.6, 131.9, 129.4, 129.0, 128.8, 128.3, 127.6, 126.5, 126.0, 124.0, 122.7, 120.1, 117.5, 72.2; HPLC (Chiralpak IB, *i*-PrOH/hexane = 15/85, flow rate = 0.50 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 19.13$ min, $t_{\text{major}} = 50.19$ min, ee = 98%; $[\alpha]_{\text{D}}^{30} = +122.7$ (c = 1.0 in CHCl_3).




(R)-(E)-2-(4-bromophenyl)-4-hydroxy-4-(2-methylstyryl)-1-

(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12i) (Table 3.10, Entry 9):


The title compound was prepared according to the general procedure as described above in 99% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.36 (s, 1H), 7.94 (d, $J = 7.0$ Hz, 1H), 7.67 (d, $J = 8.5$ Hz, 2H), 7.53~7.44 (m, 3H), 7.41~7.38 (m, 3H), 7.15~7.12 (m, 3H), 6.79 (d, $J = 16.0$ Hz, 1H), 6.35 (d, $J = 15.5$ Hz, 1H), 5.33 (s, 1H), 2.30 (s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.5, 151.6, 137.0, 136.1, 134.7, 134.4, 133.4, 132.2, 131.8, 130.3, 129.8,

128.8, 128.4, 128.3, 128.1, 126.2, 126.1, 126.0, 122.5, 120.1, 117.6, 72.3, 19.7; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 5/95, flow rate = 0.40 mL/min, λ = 254nm): t_{minor} = 21.89 min, t_{major} = 39.93 min, ee = 97%; $[\alpha]_{\text{D}}^{30}$ = +127.0 (c = 1.0 in CHCl₃).



(R)-(E)-4-hydroxy-2-(4-methoxyphenyl)-4-styryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12j) (Table 3.10, Entry 10):


The title compound was prepared according to the general procedure as described above in 92% yield. ¹H NMR (CDCl₃, 500 MHz): δ 9.34 (s, 1H), 7.92 (d, J = 7.5 Hz, 1H), 7.51~7.49 (m, 2H), 7.46~7.43 (m, 3H), 7.36 (d, J = 7.5 Hz, 2H), 7.30~7.27 (m, 2H), 7.22 (t, J = 7.0 Hz, 1H), 7.01 (d, J = 8.5 Hz, 2H), 6.56 (d, J = 16.0 Hz, 1H), 6.43 (d, J = 15.5 Hz, 1H), 5.53 (s, 1H), 3.89 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 193.2, 162.2, 153.4, 137.5, 136.0, 133.8, 133.3, 131.2, 130.2, 128.6, 128.5, 128.2, 128.1, 127.0, 125.9, 122.9, 122.7, 120.2, 117.7, 113.9, 72.3, 55.5; HPLC (Chiralpak AS-H, *i*-PrOH/hexane = 10/90, flow rate = 0.45 mL/min, λ = 254nm): t_{minor} = 25.85 min, t_{major} = 71.24 min, ee = 96%; $[\alpha]_{\text{D}}^{30}$ = +126.3 (c = 1.0 in CHCl₃).



(R)-(E)-4-hydroxy-4-styryl-2-(thiophen-2-yl)-1-(trifluoromethylsulfonyl)-1,4-

dihydroquinoline-3-carbaldehyde (12k) (Table 3.10, Entry 11):


The title compound was prepared according to the general procedure, as described above in 90% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.50 (s, 1H), 7.92 (d, $J = 7.0$ Hz, 1H), 7.67 (d, $J = 5.0$ Hz, 1H), 7.52~7.49 (m, 2H), 7.46~7.44 (m, 2H), 7.36 (d, $J = 7.5$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 2H), 7.25~7.23 (m, 1H), 7.18~7.16 (m, 1H), 6.55 (d, $J = 15.5$ Hz, 1H), 6.42 (d, $J = 16.0$ Hz, 1H), 5.36 (s, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.5, 146.0, 137.2, 135.8, 135.5, 134.5, 133.5, 131.5, 131.3, 130.8, 130.7, 128.8, 128.5, 128.4, 128.2, 127.4, 127.0, 126.0, 123.0, 120.3, 117.8, 72.6; HPLC (Chiralpak IB, *i*-PrOH/hexane = 5/95, flow rate = 0.40 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 18.93$ min, $t_{\text{major}} = 21.13$ min, ee = 96%; $[\alpha]_{\text{D}}^{30} = +76.3$ ($c = 1.0$ in CHCl_3).



(R)-(E)-4-hydroxy-2-phenethyl-4-styryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12l) (Table 3.10, Entry 12):


The title compound was prepared according to the general procedure as described above in 99% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.67 (s, 1H), 7.87 (d, $J = 7.5$ Hz, 1H), 7.49 (t, $J = 7.5$ Hz, 1H), 7.42 (d, $J = 8.0$ Hz, 1H), 7.36 (d, $J = 8.0$ Hz, 1H), 7.30~7.24 (m, 4H), 7.21~7.19 (m, 4H), 6.94~6.93 (m, 2H), 6.40 (d, $J = 15.5$ Hz, 1H), 6.25 (d, $J = 16.0$ Hz, 1H), 5.46 (s, 1H), 3.45~3.39 (m, 1H), 3.24~3.18 (m, 1H), 3.11~3.06 (m, 1H), 2.83~2.77 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 191.0, 153.8, 138.3, 137.7, 135.8, 135.1, 133.3, 130.5, 130.0, 128.7, 128.6, 128.52, 128.47, 128.3, 128.0, 127.0, 126.9, 125.7, 122.4,

120.4, 117.8, 115.2, 72.4, 35.5, 32.7; HPLC (Chiralpak IB, *i*-PrOH/hexane = 15/85, flow rate = 0.45 mL/min, λ = 254nm): t_{minor} = 14.17 min, t_{major} = 23.86 min, ee = 96%; $[\alpha]_{\text{D}}^{30}$ = +10.8 (c = 1.0 in CHCl₃).




(R)-(E)-4-hydroxy-2-pentyl-4-styryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12m) (Table 3.10, Entry 13):

The title compound was prepared according to the general procedure, as described above in 98% yield. ¹H NMR (CDCl₃, 300 MHz): δ 10.1 (s, 1H), 7.88~7.85 (m, 1H), 7.49~7.42 (m, 2H), 7.40~7.36 (m, 1H), 7.34~7.29 (m, 4H), 7.23~7.20 (m, 1H), 6.42 (d, J = 16.0 Hz, 1H), 6.27 (d, J = 16.0 Hz, 1H), 5.60 (s, 1H), 3.28~3.20 (m, 1H), 2.87~2.77 (m, 1H), 1.75~1.72 (m, 1H), 1.55~1.52 (m, 1H), 1.334~1.327 (m, 4H), 0.88 (t, J = 7.0 Hz, 3H); ¹³C NMR (CDCl₃, 125 MHz): δ 191.2, 156.5, 137.7, 135.9, 134.0, 133.5, 130.6, 130.0, 128.6, 128.5, 128.3, 128.0, 126.9, 125.7, 122.4, 120.4, 117.8, 72.4, 31.3, 30.7, 29.8, 22.2, 13.8; HPLC (Chiralpak IB, *i*-PrOH/hexane = 15/85, flow rate = 0.45 mL/min, λ = 254nm): t_{minor} = 11.67 min, t_{major} = 12.83 min, ee = 97%; $[\alpha]_{\text{D}}^{30}$ = +35.8 (c = 1.0 in CHCl₃).




(R)-4-hydroxy-2,4-distyryl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12n) (Table 3.10, Entry 14):

The title compound was prepared according to the general procedure, as described above in 70% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.80 (s, 1H), 7.92~7.89 (m, 1H), 7.57~7.54 (m, 2H), 7.52~7.49 (m, 2H), 7.46~7.42 (m, 4H), 7.36~7.33 (m, 2H), 7.31~7.22 (m, 3H), 7.03 (d, $J = 15.6$ Hz, 1H), 6.87 (d, $J = 15.6$ Hz, 1H), 6.50 (d, $J = 15.9$ Hz, 1H), 6.39 (d, $J = 15.9$ Hz, 1H), 5.50 (s, 1H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 191.3, 151.4, 145.1, 136.8, 135.9, 134.4, 134.2, 133.2, 131.1, 130.5, 130.2, 129.1, 128.5, 128.2, 128.1, 127.8, 126.9, 125.8, 122.8, 121.3, 117.9, 117.1, 72.3; HPLC (Chiralpak IB, *i*-PrOH/hexane = 10/90, flow rate = 0.45 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 16.66$ min, $t_{\text{major}} = 27.54$ min, ee = 94%; $[\alpha]_{\text{D}}^{29} = +360.0$ (c = 1.0 in CHCl_3).



(R)-4-hydroxy-4-methyl-2-phenyl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12o) (Table 3.10, Entry 15):

The title compound was prepared according to the general procedure, as described above in 96% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.28 (s, 1H), 7.89~7.85 (m, 1H), 7.56~7.38 (m, 8H), 5.18 (s, 1H), 1.83 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.3, 152.0, 139.4, 136.0, 133.1, 131.2, 130.8, 128.6, 128.3, 128.0, 125.2, 122.5, 121.0, 116.8, 69.6, 31.6; HPLC (Chiralpak IB, *i*-PrOH/hexane = 10/90, flow rate = 0.45 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 10.48$ min, $t_{\text{major}} = 15.88$ min, ee = 95%; $[\alpha]_{\text{D}}^{29} = +187.2$ (c = 1.0 in CHCl_3).



(R)-4-hydroxy-2-(3-nitrophenyl)-4-phenyl-1-(trifluoromethylsulfonyl)-1,4-dihydroquinoline-3-carbaldehyde (12p) (Table 3.10, Entry 16):

The title compound was prepared according to the general procedure, as described above in 84% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.39 (s, 1H), 8.45~8.41 (m, 1H), 8.38 (s, 1H), 8.10~8.07 (m, 1H), 7.86 (d, $J = 7.5$ Hz, 1H), 7.73 (t, $J = 7.8$ Hz, 1H), 7.59~7.48 (m, 3H), 7.45~7.29 (m, 5H), 5.38 (s, 1H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 191.9, 150.4, 148.0, 143.3, 137.6, 136.5, 133.5, 132.7, 129.7, 129.0, 128.7, 128.5, 126.6, 126.0, 122.4, 120.7, 116.5, 73.9; HPLC (Chiralpak IB, *i*-PrOH/hexane = 20/80, flow rate = 0.45 mL/min, $\lambda = 254\text{nm}$): $t_{\text{minor}} = 17.52$ min, $t_{\text{major}} = 51.50$ min, ee = 99%; $[\alpha]_{\text{D}}^{30} = +136.7$ (c = 1.0 in CHCl_3).

3.7 Reference

[1] For recent reviews of organocatalytic cascade/tandem/domino reactions, see: (a) D. Enders, C. Grondal, M. R. M. Hüttl, *Angew. Chem., Int. Ed.* **2007**, *46*, 1570-1581; (b) H.-C. Guo, J.-A. Ma, *Angew. Chem., Int. Ed.* **2006**, *45*, 354-366; (c) X. Yu, W. Wang, *Org. Biomol. Chem.* **2008**, *6*, 2037-2046; (d) G. Guillena, D. J. Ramon, M. Yus, *Tetrahedron: Asymmetry* **2007**, *18*, 693-700; (e) A. Dondoni, A. Massi, *Angew. Chem. Int. Ed.* **2008**,

- 47, 4638-4660; (f) C. Grondal, M. Jeanty, D. Enders, *Nature Chemistry*, **2010**, *2*, 167-178.
- [2] M. J. Gaunt, C. C. C. Johansson, A. McNally, N. T. Vo, *Drug Discovery Today*. **2007**, *12*, 8-27.
- [3] B. List, *Chem. Commun.* **2006**, 819-824, and references therein.
- [4] N. Halland, P. S. Aburel, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2004**, *43*, 1272-1272.
- [5] D. Enders, M. R. M. Hüttl, C. Grondal, G. Raabe, *Nature* **2006**, *441*, 861-863.
- [6] F.-L. Zhang, A.-W. Xu, Y.-F. Gong, M.-H. Wei, X.-L. Yang, *Chem. Eur. J.* **2009**, *15*, 6815-6818.
- [7] K. Jiang, Z.-J. Jia, X. Yin, L. Wu, Y.-C. Chen, *Org. Lett.* **2010**, *12*, 2766-2769.
- [8] B.-C. Hong, P. Kotame, C.-W. Tsai, J.-H. Liao, *Org. Lett.* **2010**, *12*, 776-779.
- [9] S. B. Jones, B. Simmons, D.W. C. MacMillan, *J. Am. Chem. Soc.* **2009**, *131*, 13606-13607.
- [10] For reviews of allenes in organic synthesis, see: (a) *Modern Allene Chemistry* (Eds.: N. Krause, A. S. K. Hashmi), Wiley-VCH, Weinheim, **2004**; (b) S. Ma, *Chem. Rev.* **2005**, *105*, 2829-2871; (c) H. Kim, L. J. Willams, *Curr. Opin. Drug Discovery Dev.* **2008**, *11*, 870.
- [11] Examples of catalytic asymmetric synthesis of chiral allenes are very rare; for a recent review, see: (a) A. Hoffmann-Röder, N. Krause, *Angew. Chem. Int. Ed.* **2002**, *41*,

2933-2935; for examples, see: (b) W. de Graaf, J. Boersma, G. van Koten, C. J. Elsevier, *J. Organomet. Chem.* **1989**, 378, 115-124; (c) M. Ogasawara, H. Ikeda, T. Nagano, T. Hayashi, *J. Am. Chem. Soc.* **2001**, 123, 2089-2090; (d) J. W. Han, N. Tokunaga, T. Hayashi, *J. Am. Chem. Soc.* **2001**, 123, 12915-12916; (e) T. Hayashi, N. Tokunaga, K. Inoue, *Org. Lett.* **2004**, 6, 305-307; (f) C.-Y. Li, X.-B. Wang, X.-L. Sun, Y. Tang, J.-C. Zheng, Z.-H. Xu, Y.-G. Zhou, L.-X. Dai, *J. Am. Chem. Soc.* **2007**, 129, 1494-1495; (g) B. M. Trost, J. Xie, *J. Am. Chem. Soc.* **2008**, 130, 6231-6242; (h) H. Liu, D. Leow, G.-W. Huang, C.-H. Tan, *J. Am. Chem. Soc.* **2009**, 131, 7212-7213.

[12] For the use of alkynyl imines in non-asymmetric conjugate addition reactions and the proposed involvement of allenamines in these processes, see: (a) I. Hachiya, K. Ogura, M. Shimizu, *Org. Lett.* **2002**, 4, 2755-2757; (b) M. Shimizu, T. Nishi, A. Yamamoto, *Synlett* **2003**, 1469-1473.

[13] For the use of silyloxyallenes in enantioselective synthesis, see: T. E. Reynolds, K. A. Scheidt, *Angew. Chem. Int. Ed.* **2007**, 46, 7806-7809.

[14] X.-S. Zhang, S.-L. Zhang, W. Wang, *Angew. Chem. Int. Ed.* **2010**, 49, 1481-1484.

[15] D. Salni, M. V. Sargent, B. W. Skelton, I. Soediro, M. Sutisna, A. H. White, E. Yulinah, *Aust. J. Chem.* **2002**, 55, 229-232.

[16] F. Shaheen, M. Ahmad, S. N. Khan, S. S. Hussain, S. Anjum, B. Tashkhodjaev, K. Turgunov, M. N. Sultankhodzhaev, M. I. Choudhary, Atta-ur-Rahman, *Eur. J. Org. Chem.* **2006**, 2371-2377.

[17] For reviews of chemistry and biology of chromenes, see: (a) E. E. Schweizer, O. Meeder-Nycz in *Chromenes, Chromanes, Chromones* (Eds.: G. P. Ellis), Wiley-Interscience, New York, **1977**; (b) B. A. Keay in *Comprehensive Heterocyclic Chemistry II*, Vol. 2 (Eds.: A. R. Katritzky, C.W. Rees, E. F. V. Scriven), Pergamon, Oxford, **1996**, p. 395; (c) K. C. Nicolaou, J. A. Pfefferkorn, A. J. Roecker, G.-Q. Cao, S. Barluenga, H. J. Mitchell, *J. Am. Chem. Soc.* **2000**, *122*, 9939-9953, and references therein.

[18] For recent examples of the racemic synthesis of 4*H*-chromenes, see: (a) Y. Nishibayashi, Y. Inada, M. Hidai, S. Uemura, *J. Am. Chem. Soc.* **2002**, *124*, 7900-7901; (b) W. A. L. van Otterlo, E. L. Ngidi, S. Kuzvidza, G. L. Morgans, S. S. Moleele, C. B. de Koning, *Tetrahedron* **2005**, *61*, 9996-10006; (c) M. Kidwai, S. Saxena, M. K. R. Khan, S. S. Thukral, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4295-4298; (d) Y.-L. Shi, M. Shi, *Org. Lett.* **2005**, *7*, 3057-3060; (e) Y.-L. Shi, M. Shi, *Chem. Eur. J.* **2006**, *12*, 3374-3378; (f) Y. W. Fang, C. Z. Li, *J. Org. Chem.* **2006**, *71*, 6427-6431; (g) L. Ye, X. Sun, C. Zhu, Y. Tang, *Org. Lett.* **2006**, *8*, 3853-3856; (h) M. N. Elinson, A. S. Dorofeev, S. K. Feducovich, S. V. Gorbunov, R. F. Nasybullin, N. O. Stepanov, G. I. Nikishin, *Tetrahedron Lett.* **2006**, *47*, 7629-7633; (i) J. Fan, Z. Wang, *Chem. Commun.* **2008**, 5381-5383; (j) S. Makarem, A. A. Mohammadi, A. R. Fakhari, *Tetrahedron Lett.* **2008**, *49*, 7194-7196; (k) L. Lu, J. Wei, J. Chen, J. Zhang, H. Deng, M. Shao, H. Zhang, W. Cao, *Tetrahedron* **2009**, *65*, 9152-9156; (l) K. Funabiki, T. Komeda, Y. Kubota, M. Matsui, *Tetrahedron* **2009**, *65*, 7457-7463.

[19] To the best of our knowledge, only one study of the catalytic asymmetric synthesis of 4*H*-chromenes using a Pd^{II}-(*S,S*)-chiraphos complex for conjugate addition-

dehydration (two steps) has been reported: T. Nishikata, Y. Yamamoto, N. Miyaoura, *Adv. Synth. Catal.* **2007**, *349*, 1759-1764.

[20] Our laboratory has developed several organocatalytic enantioselective cascade reactions for the construction of chromanes or chromenes: (a) W. Wang, H. Li, J. Wang, L. Zu, *J. Am. Chem. Soc.* **2006**, *128*, 10354-10355; (b) H. Li, J. Wang, T. E-Nunu, L. Zu, W. Jiang, S. Wei, W. Wang, *Chem. Commun.* **2007**, 507-509; (c) L.-S. Zu, S.-L. Zhang, H.-X. Xie, W. Wang, *Org. Lett.* **2009**, *11*, 1627-1630; for work by other groups on organocatalytic oxa-Michael cascades, see the review: (d) C. F. Nising, S. Bräse, *Chem. Soc. Rev.* **2008**, *37*, 1218-1228; (e) T. Govender, L. Hojabri, F. M. Moghaddam, P. I. Arvidsson, *Tetrahedron: Asymmetry* **2006**, *17*, 1763-1767; (f) M. M. Biddle, M. Lin, K. A. Scheidt, *J. Am. Chem. Soc.* **2007**, *129*, 3830-3831; (g) H. Sundén, I. Ibrahim, G. L. Zhao, L. Eriksson, A. Córdova, *Chem. Eur. J.* **2007**, *13*, 574-581; (h) M. Rueping, E. Sugiono, E. Merino, *Angew. Chem. Int. Ed.* **2008**, *47*, 3046-3049; (i) K. Liu, A. Chougnet, W.-D. Woggon, *Angew. Chem. Int. Ed.* **2008**, *47*, 5827-5829; (j) D.-Q. Xu, Y.-F. Wang, S.-P. Luo, S. Zhang, A.-G. Zhong, H. Chen, Z.-Y. Xu, *Adv. Synth. Catal.* **2008**, *350*, 2610-2616; (k) P. Kotame, B.-C. Hong, J. Liao, *Tetrahedron Lett.* **2009**, *50*, 704-707; (l) E. Reyes, G. Talavera, J. L. Vicario, D. Badia, L. Carrillo, *Angew. Chem. Int. Ed.* **2009**, *48*, 5701-5704; (m) F.-L. Zhang, A.-W. Xu, Y.-F. Gong, M.-H. Wei, X.-L. Yang, *Chem. Eur. J.* **2009**, *15*, 6815-6818.

[21] For a recent review of prolinol ether catalysis, see: (a) C. Palomo, A. Mielgo, *Angew. Chem. Int. Ed.* **2006**, *45*, 7876-7880; (b) A. Mielgo, C. Palomo, *Chem. Asian J.* **2008**, *3*, 922-948.

- [22] J. M. Betancort, C. F. Barbas III, *Org. Lett.* **2001**, *3*, 3737-3740.
- [23] (a) N. S. Simpkins, *Sulfones in Organic Synthesis*, Pergamon Press, Oxford, **1993**; (b) *The Chemistry of Sulfones and Sulfoxides*; S. Patai, Z. Rapoport, C. Stirling, Eds.; J. Wiley & Sons: Chichester, U.K., **1988**; (c) T. Toru, C. Bolm, *Organosulfur Chemistry in Asymmetric Synthesis*, Wiley-VCH, Weinheim, **2008**; (d) J.-E. Bäckvall, R. Chinchilla, C. Najera, M. Yus, *Chem. Rev.* **1998**, *98*, 2291-2312.
- [24] (a) A. C. Brown, L. A. Carpino, *J. Org. Chem.* **1985**, *50*, 1749-1750; (b) C. Nájera, M. Yus, *Tetrahedron*, **1999**, *55*, 10547-10658.
- [25] (a) B. Trost, *Bull. Chem. Soc. Jpn.* **1998**, *61*, 107; (b) J. C. Carretero, R. Arrayás, J. Adrio, in *Sulfones in Asymmetric Catalysis*, (Eds.: T. Toru, C. Bolm), Wiley, **2008**, ch. 9, pp. 291-320.
- [26] For a review on the chemistry of vinyl sulfones, see: N. S. Simpkins, *Tetrahedron*, **1990**, *46*, 6951-6984.
- [27] For selected examples of organocatalytic enantioselective Michael additions to vinyl sulfones, see: (a) S. Pinheiro, A. Guingant, D. Desmaële, J. d'Angelo, *Tetrahedron: Asymmetry* **1992**, *3*, 1003-1006; (b) D. Desmaële, S. Delarue-Cochin, C. Cave, J. d'Angelo, G. Morgant, *Org. Lett.* **2004**, *6*, 2421-2424; (c) H. Li, J. Song, X. Liu, L. Deng, *J. Am. Chem. Soc.* **2005**, *127*, 8948-8949; (d) S. Mosse, A. Alexakis, *Org. Lett.* **2005**, *7*, 4361-4364; (e) Q. Zhu, Y. Lu, *Org. Lett.* **2008**, *10*, 4803-4806; (f) Q. Zhu, L. Cheng, Y. Lu, *Chem. Commun.* **2008**, 6315-6317; (g) Q. Zhu, Y. Lu, *Org. Lett.* **2009**, *11*, 1721-1724; (g) S. Sulzer-Mossé, A. Alexakis, J. Mareda, G. Bollot, G. Bernardinelli, Y. Filinchuk,

Chem. Eur. J. **2009**, *15*, 3204-3220; (h) A. Landa, M. Maestro, C. Masdeu, Á. Puente, S. Vera, M. Oiarbide, C. Palomo, *Chem. Eur. J.* **2009**, *15*, 1562-1565; (i) Q. Zhu, Y. Lu, *Angew. Chem. Int. Ed.* **2010**, *49*, 7753-7756.

[28] J. Hu, W. Zhang, F. Wang, *Chem. Commun.* **2009**, 7465-7478 and references therein.

[29] C.-L. Liu, X.-S. Zhang, R. Wang, W. Wang, *Org. Lett.* **2010**, *12*, 4948-4951.

[30] For selected examples of synthesis natural products bearing quaternary stereocenters, see: (a) A. W. G. Burgett, Q. Li, Q. Wei, P. G. Harran, *Angew. Chem. Int. Ed.* **2003**, *42*, 4961-4966; (b) B. Jiang, M. Xu, *Angew. Chem. Int. Ed.* **2004**, *43*, 2543-2546; (c) A. Steven, L. E. Overman, *Angew. Chem. Int. Ed.* **2007**, *46*, 5488-5508; (d) G. E. Veitch, E. Beckmann, B. J. Burke, A. Boyer, C. Ayats, S. V. Ley, *Angew. Chem. Int. Ed.* **2007**, *46*, 7633-7635; (e) R. M. Marcia de Figueiredo, M. Christmann, *Eur. J. Org. Chem.* **2007**, 2575-2600.

[31] For recent reviews of catalytic asymmetric construction of quaternary stereocenters, see: (a) *Quaternary Stereocenters-Challenges and Solutions for Organic Synthesis*; J. Christoffers, A. Baro, Eds.; Wiley-VCH: Weinheim, **2005**; (b) K. Fuji, *Chem. Rev.* **1993**, *93*, 2037-2066; (c) E. J. Corey, A. Guzman-Perez, *Angew. Chem. Int. Ed.* **1998**, *37*, 388-401; (d) J. Christoffers, A. Mann, *Angew. Chem. Int. Ed.* **2001**, *40*, 4591-4597; (e) C. J. Douglas, L. E. Overman, *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 5363-5367; (f) J. Christoffers, A. Baro, *Adv. Synth. Catal.* **2005**, *347*, 1473-1482; (g) M. Bella, T. Gasperi, *Synthesis*, **2009**, *10*, 1583-1614; (h) S. F. Martin, *Tetrahedron*, **1980**, *36*, 419-460.

[32] For recent reviews, see: (a) O. Riant, J. Hannedouche, *Org. Biomol. Chem.* **2007**, *5*, 873-888; (b) M. Shibasaki, M. Kanai, *Chem. Rev.* **2008**, *108*, 2853-2873.

[33] For selected examples of direct aldol reactions which aimed at stereoselective construction of quaternary stereocenters, see: (a) A. Bøgevig, N. Kumaragurubaran, K. A. Jørgensen, *Chem. Commun.*, **2002**, 620-621; (b) O. Tokuda, T. Kano, W.-G. Gao, T. Ikemoto, K. Maruoka, *Org. Lett.* **2005**, *7*, 5103-5105; (c) P. Dambruoso, A. Massi, A. Dondoni, *Org. Lett.* **2005**, *7*, 4657-4660; (d) Z. Tang, L.-F. Cun, X. Cui, A.-Q. Mi, Y.-Z. Jiang, L.-Z. Gong, *Org. Lett.* **2006**, *8*, 1263-1266; (e) S. Samanta, C.-G. Zhao, *J. Am. Chem. Soc.* **2006**, *128*, 7442-7443.

[34] For selected examples of bioactive natural products containing 4*H*-chromene core structures, see: (a) J. Demyttenaere, K. V. Syngel, A. P. Markusse, S. Vervisch, S. Debenedetti, N. D. Kimpe, *Tetrahedron* **2002**, *58*, 2163-2166; (b) Z.-W. Huang, US006660871B2, **2003**; (c) S. G. Das, J. M. Doshi, D. F. Tian, S. N. Addo, B. Srinivasan, D. L. Hermanson, C.-G. Xing, *J. Med. Chem.* **2009**, *52*, 5937-5949; (d) E. J. Corey, L.-I. Wu, *J. Am. Chem. Soc.* **1993**, *115*, 9327-9328; (e) F. Shaheen, M. Ahmad, S. N. Khan, S. S. Hussain, S. Anjum, B. Tashkhodjaev, K. Turgunov, M. N. Sultankhodzhaev, M. I. Choudhary, A. Rahman, *Eur. J. Org. Chem.* **2006**, *10*, 2371-2377.

[35] For reviews, see: (a) G. Jones, *Comprehensive Heterocyclic Chemistry*, Pergamon, New York, **1984**; (b) M. Johannsen, K. A. Jørgensen, *Chem. Rev.* **1998**, *98*, 1689-1708; (c) T. E. Müller, M. Beller, *Chem. Rev.* **1998**, *98*, 675-704; (d) A. R. Katritzky, A. F. Pozharskii, *Handbook of Heterocyclic Chemistry*, 2nd ed.; Pergamon: Oxford, **2000**; p

616; (e) J. P. Michael, *Nat. Prod. Rep.* **2005**, *25*, 627-646; (d) J. P. Michael, *Nat. Prod. Rep.* **2007**, *24*, 223-246; (f) J. P. Michael, *Nat. Prod. Rep.* **2008**, *25*, 166-187; (g) R. P. Verma, C. Hansch, *Chem. Rev.* **2009**, *109*, 213-235; (h) V. R. Solomon, H. Lee, *Curr. Med. Chem.* **2011**, *18*, 1488-1508. (i) M. M. Ghorab, F. A. Ragab, H. I. Heiba, W. M. Ghorab, *J. Heterocycl. Chem.* **2011**, *48*, 1269-1279; (j) N. T. Patil, V. Singh, *J. Organomet. Chem.* **2011**, *696*, 419-432.

[36] For selected reviews, see: (a) A. R. Katritzky, S. Rachwal, B. Rachwal, *Tetrahedron* **1996**, *52*, 15031-15070; (b) V. V. Kouznetsov, L. Y. Vargas Méndez, C. M. Meléndez Gómez, *Curr. Org. Chem.* **2005**, *9*, 141-161; (c) M. Arisawa, Y. Terada, C. Theeraladanon, K. Takahashi, M. Nakagawa, A. J. Nishida, *Organomet. Chem.* **2005**, *690*, 5398-5406; (d) H. Ila, O. Baron, A. J. Wagner, P. Knochel, *Chem. Commun.* **2006**, 583-593; (e) S. A. Yamashkin, E. A. Oreshkina, *J. Heterocycl. Chem.* **2006**, *42*, 701-718; (f) V. Sridharan, P. A. Suryavanshi, J. C. Menéndez, *Chem. Rev.* **2011**, *111*, 7157-7259.

[37] For selected examples, see: (a) P. Friedlander, *Chem. Ber.* **1882**, *15*, 2572; (b) a review: J. Marco-Contelles, E. Pérez-Mayoral, A. Samadi, M. C. Carreiras, E. Soriano, *Chem. Rev.* **2009**, *109*, 2652-2671; (c) B. L. Miller, B. R. Mcnaughton, *Org. Lett.* **2003**, *5*, 4257-4259; (d) Y.-Z. Hu, G. Zhang, R. P. Thummel, *Org. Lett.* **2003**, *5*, 2251-2253; (e) R. Martínez, D. J. Ramón, M. Yus, *J. Org. Chem.* **2008**, *73*, 9778-9780; (f) D. S. Bose, M. Idrees, N. M. Jakka, J. V. Rao, *J. Comb. Chem.* **2010**, *12*, 100-110.

[38] For reviews, see: (a) R. H. F. Manske, M. Kukla, *Org. React.* **1953**, *7*, 59-98; (b) B. C. Ranu, A. Hajra, S. S. Dey, U. Jana, *Tetrahedron* **2003**, *59*, 813-819.

[39] For selected examples, see: (a) S. E. Denmark, S. Venkatraman, *J. Org. Chem.* **2006**, *71*, 1668-1676; (b) A. R. Mackenzie, C. J. Moody, C. W. Rees, *Tetrahedron* **1986**, *42*, 3259-3268; (c) S. Y. Tanaka, M. Yasuda, A. Baba, *J. Org. Chem.* **2006**, *71*, 800-803.

[40] H. R. Henze, D. W. Carroll, *J. Am. Chem. Soc.* **1954**, *76*, 4580-4584.

[41] (a) A. Combes, *Bull. Soc. Chim. Fr.* **1888**, *49*, 89-91; (b) L. Jerry, J. Born, *J. Org. Chem.* **1972**, *37*, 3952-3953.

[42] For selected examples, see: (a) M. Beller, O. R. Thiel, H. Trauthwein, C. G. Hartung, *Chem. Eur. J.* **2000**, *6*, 2513-2522; (b) B. Jiang, Y.-G. Si, *J. Org. Chem.* **2002**, *67*, 9449-9451; (c) R. P. Korivi, C.-H. Cheng, *J. Org. Chem.* **2006**, *71*, 7079-7082; (d) L. Li, W. D. Jones, *J. Am. Chem. Soc.* **2007**, *129*, 10707-10713; (e) C. S. Yi, S. Y. Yun, *Org. Lett.* **2005**, *7*, 2181-2183; (f) Z. Zhang, J. Tan, Z. Wang, *Org. Lett.* **2008**, *10*, 173-175; (g) B. Gabriele, R. Mancuso, G. Salerno, E. Lupinacci, G. Ruffolo, M. Costa, *J. Org. Chem.* **2008**, *73*, 4971-4977; (h) B. Gabriele, R. Mancuso, G. Salerno, G. Ruffolo, P. Plastina, *J. Org. Chem.* **2007**, *72*, 6873-6877; (i) Y. Matsubara, S. Hirakawa, Y. Yamaguchi, Z. Yoshida, *Angew. Chem. Int. Ed.* **2011**, *50*, 7670-7673.

[43] For examples of asymmetric organometallic catalysis, see: (a) W.-B. Wang, S.-M. Lu, P.-Y. Yang, X.-W. Han, Y.-G. Zhou, *J. Am. Chem. Soc.* **2003**, *125*, 10536-10537; (b) S. M. Lu, Y. Q. Wang, X. W. Han, Y. G. Zhou, *Angew. Chem., Int. Ed.* **2006**, *45*, 2260-2263; (c) Z.-J. Wang, G.-J. Deng, Y. Li, Y.-M. He, W.-J. Tang, Q.-H. Fan, *Org. Lett.* **2007**, *9*, 1243-1246; (d) Q. S. Guo, D. M. Du, J. Xu, *Angew. Chem., Int. Ed.* **2008**, *47*, 759-762; (e) H. F. Zhou, Z. W. Li, Z. J. Wang, T. L. Wang,; L. J. Xu, Y. He, Q. H. Fan, J.

Pan, L. Q. Gu, A. S. C. Chan, *Angew. Chem., Int. Ed.* **2008**, *47*, 8464-8467; (f) C. Wang, C. Q. Li, X. F. Wu, A. Pettman, J. L. Xiao, *Angew. Chem., Int. Ed.* **2009**, *48*, 6524-6528; (g) Z.-Y. Han, Z.-H. Chen, L.-Z. Gong, *J. Am. Chem. Soc.* **2009**, *131*, 9182-9183; (h) D. S. Giera, C. Schneider, *Org. Lett.* **2010**, *12*, 4884-4887.

[44] For selected examples of organocatalysis, see: (a) N. Halland, P. S. Aburel, K. A. Jørgensen, *Angew. Chem., Int. Ed.* **2003**, *42*, 661-665; (b) M. Rueping, A. R. Antonchick, T. Theissmann, *Angew. Chem., Int. Ed.* **2006**, *45*, 3683-3686; (c) S. Fustero, J. Moscardó, D. Jiménez, M. D. Pérez-Carrión, M. del Pozo, C. Sánche-Roselló, *Chem. Eur. J.* **2008**, *14*, 9868-9872; (d) H. Liu, G. Dagousset, G. Masson, P. Retailleau, J. Zhu, *J. Am. Chem. Soc.* **2009**, *131*, 4598-4599; (e) Y. K. Kang, S. M. Kim, D. Y. Kim, *J. Am. Chem. Soc.* **2010**, *132*, 11847-11849; (f) M. Rueping, T. Theissmann, *Chem. Sci.* **2010**, *1*, 473-476; (g) H. Xu, S. J. Zuend, M. G. Woll, Y. Tao, E. N. Jacobsen, *Science* **2010**, *327*, 986-990; (h) X. Liu, Y. Lu, *Org. Lett.* **2010**, *12*, 5592-5595; (i) Z.-X. Jia, Y.-C. Luo, P.-F. Xu, *Org. Lett.* **2011**, *13*, 832-835; (j) K. Mori, K. Ehara, K. Kurihara, T. Akiyama, *J. Am. Chem. Soc.* **2011**, *133*, 6166-6169; (k) L. Ren, T. Lei, J.-X. Ye, L.-Z. Gong, *Angew. Chem., Int. Ed.* **2012**, *51*, 771-774; (l) H.-R. Tan, H.-F. Ng, J. Chang, J. Wang, *Eur. J. Chem.* **2012**, DOI: 10.1002/chem.201103136.

[45] For recent selected examples of organometallic catalysis, see: (a) M. Takamura, K. Funabashi, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.* **2000**, *122*, 6327-6328; (b) P. A. Evans, J. E. Robinson, K. K. Moffett, *Org. Lett.* **2001**, *3*, 3269-3271; (c) S. J. Pastine, S.-W. Youn, D. Sames, *Tetrahedron* **2003**, *59*, 8859-8868; (d) P. D. Pohlhaus, R. K. Bowman, J. S. Johnson, *J. Am. Chem. Soc.* **2004**, *126*, 2294-2295; (e) G.-L. Lu, H. C.

Malinakova, *J. Org. Chem.* **2004**, *69*, 4701-4715; (f) L. Cointeaus, A. Alexakia, *Tetrahedron: Asymmetry* **2005**, *16*, 925-929.

[46] For examples of organocatalysis, see: (a) H. Li, J. Wang, H.-X. Xie, L.-S. Zu, W. Jiang, E. N. Duesler, W. Wang, *Org. Lett.* **2007**, *9*, 965-968; (b) H. Sundén, R. Rios, I. Ibrahim, G.-L. Zhao, L. Eriksson, A. Cordova, *Adv. Synth. Catal.* **2007**, *349*, 827-832.

[47] For selected non-asymmetric methods of 1,4-dihydroquinoles, see: (a) Y. Hitotsuyanagi, M. Kobayashi, M. Fukuyo, K. Takeya, H. A. Itokawa, *Tetrahedron Lett.* **1997**, *38*, 8295-8296; (b) J. N. Kim, H. S. Kim, J. H. Gong, Y. M. Chung, *Tetrahedron Lett.* **2001**, *42*, 8341-8344; (c) C. Tratat, S. Giorgi-Renault, H.-P. Husson, *Org. Lett.* **2002**, *4*, 3187-3189; (d) M.-L. Bennasar, T. Roca, M. Monerri, D. García-Díaz, *Tetrahedron Lett.* **2005**, *46*, 4035-4038; (e) X.-J. Wu, R. Jiang, B. Wu, X.-M. Su, X.-P. Xu, S.-J. Ji, *Adv. Synth. Catal.* **2009**, *351*, 3150-3156; (f) A. Kumar, A. E. Alegria, *J. Heterocyclic Chem.* **2010**, *47*, 1275-1282; (g) X.-J. Wu, X.-P. Xu, X.-M. Su, G. Chen, Y. Zhang, S.-J. Ji, *Eur. J. Org. Chem.* **2009**, 4963-4970; (h) G. Viault, D. Grée, T. Roisnel, S. Chandrasekhar, R. Grée, *Tetrahedron* **2009**, *65*, 10149-10154; (i) T. Watanabe, T. Oishi, N. Fujii, H. Ohno, *Org. Lett.* **2007**, *9*, 4821-4824; (j) C. Shi, J. Wang, H. Chen, D.-Q. Shi, *J. Comb. Chem.* **2010**, *12*, 430-434; (k) S. S. Koval'skaya, N. G. Kozlov, E. A. Dikumar, *Russ. J. Org. Chem.* **2010**, *46*, 1121-1134; (l) N. G. Kozlov, A. B. Tereshko, *Russ. J. Org. Chem.* **2010**, *46*, 1223-1225; (m) N. G. Kozlov, S. L. Bondarevb, V. N. Knyukshtoc, B. A. Odnoburtseva, L. I. Basalaeva, *Russ. J. Org. Chem.* **2010**, *46*, 1645-1651; (n) K. Namba, M. Kanaki, H. Suto, M. Nishizawa, K. Tanino, *Org. Lett.* **2012**, *14*, DOI: 10.1021/ol2034492.

[48] F. Rezgui, P. Mangeney, A. Alexakis, *Tetrahedron Lett.* **1999**, *40*, 6241-6244.

[49] Fu and co-workers have reported an elegant study of the effect of “*N*” protecting form on stereoselectivity: E. C. Lee, B. L. Hodous, E. Bergin, C. Shih, G. C. Fu, *J. Am. Chem. Soc.* **2005**, *125*, 11586-11587.

[50] (a) Michael-aldol: see reference [29]; (b) Michael-Michael: see reference [14]; (c) Michael-Mannich: J. Aleman, A. Nunez, L. Marzo, V. Marcos, C. Alvarado, J. L. Garcia Ruano, *Chem. Eur. J.* **2010**, *16*, 9453-9456.

[51] For examples of (*2R*, *5R*)-diphenylpyrrolidine organocatalyst promoted reactions, see (a) M. P. Brochu, S. P. Brown, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2004**, *126*, 4108-4109; (b) O. O. Fadeyi, M. L. Schulte, C. W. Lindsley, *Org. Lett.* **2010**, *12*, 3276-3278; (c) N. Halland, M. A. Lie, A. Kjærsgaard, M. Marigo, B. Schiøtt, K. A. Jørgensen, *Chem. Eur. J.* **2005**, *11*, 7083-7090; (d) S. Bertelsen, N. Halland, S. Bachmann, M. Marigo, A. Braunton, K. A. Jørgensen, *Chem. Commun.* **2005**, 4821-4823; (e) N. Halland, A. Braunton, S. Bachmann, M. Marigo, K. A. Jørgensen, *J. Am. Chem. Soc.* **2004**, *126*, 4790-4791.

[52] (a) We have reported a divergent organocatalytic Michael-alkylation cascade: H.-X. Xie, L.-S. Zu, H. Li, J. Wang, W. Wang, *J. Am. Chem. Soc.* **2007**, *129*, 10886-10894; an other example from Ma and co-workers: (b) W. Yao, L. Pan, Y. Zhang, G. Wang, X. Wang, C. Ma, *Angew. Chem., Int. Ed.* **2010**, *49*, 9210-9214.

[53] G. Litkei, A. L. Tökés, *Synthetic Communications*, **1991**, *15&16*, 1597-1609.

[54] D. G. Batt, R. Goodman, D. G. Jones, J. S. Kerr, L. R. Mantegna, C. McAllister, R. C. Newton, S. Nurnberg, P. K. Welch, M. B. Covington, *J. Med. Chem.* **1993**,*36*,1434-1442.

Chapter 4

Organocatalytic 1,6-Addition Reactions


4.1 Background and Significance

The catalytic 1,4-addition¹ of various nucleophiles to electron-deficient olefin has developed into one of the most reliable methods for efficiently construction of C-C and C-heteroatom bonds. In this field, attention has been focused on both metal-catalyzed² and organocatalyst-catalyzed³ processes (Scheme 4.2, eq (1)). Furthermore, major advances such as high regioselectivity (1,4- vs. 1,2-addition) and excellent control of stereoselectivity have been achieved in catalytic 1,4- addition.


On the other hand, catalytic 1,6-addition to extended conjugate systems has developed less rapidly due to the fact that there is considerable difficulties in controlling both the regioselectivity (1,2-, 1,4- vs. 1,6-addition) and stereoselectivity. Except metal-catalyzed nonasymmetric 1,6-selective conjugate addition⁴, successful reports of asymmetric 1,6-addition to various $\alpha,\beta,\gamma,\delta$ -unsaturated Michael acceptors such as 2,4-dienones, 2,4-dienoates and 2,4-dienamides catalyzed by copper⁵, rhodium⁶, iridium⁷ and iron⁸ complex have recently appeared. In contrast, the organocatalytic 1,6-addition to extended conjugated systems remains underdeveloped. The 1,4-additions to active dienes or enynes have been reported⁹. In these cases, the β -carbon is highly active in comparison with δ -carbon which accounts for the regioselectivity. Only two reports of asymmetric

1,6-addition promoted by organocatalyst are developed. In 2007, Jørgensen and co-workers¹⁰ disclosed the first organocatalytic enantioselective 1,6-addition of β -ketoesters and benzophenone imine to electron-deficient δ -unsubstituted dienes using chiral phase-transfer catalyst (Scheme 4.1). It is noteworthy that the strategy for the control of regioselectivity here is to reduce the steric hindrance at the δ -carbon to bias the system towards 1,6-addition. Very recently, asymmetric enamine 1,6-conjugate addition of aldehydes to 1,3-bis-(sulfonyl) butadienes in remarkable stereoselectivity (99% e.e. and 1:99 d.r.) was reported by A. Alexakis and J. C. Stephens¹¹ (Scheme 4.1). The exclusive regioselectivity is probably associated with the specific structural feature of the substrate designed based on the fact that a single activating sulfone group was not strong enough to activate the alkene to generate an enamine 1,4-addition product.


Scheme 4.1 Examples of Organocatalytic 1,6-Addition



2.



3.




4.2 Research Design

Due to the poor propagation of electronic effect through the extended conjugated system, the development of catalytic 1,6-addition remain a challenging task and the preliminary successful studies were strongly dependent on the nature and structure of the substrates. Regioselective 1,6-conjugate addition of $\alpha,\beta,\gamma,\delta$ -unsaturated aldehydes face even more significant obstacles. Unlike ketones, 1,2-addition of aldehydes is favorable. Moreover, in iminium catalysis, 1,4-addition occurs exclusively with dienic aldehydes without 1,6-additions^{3a}. Considering alkyne is more active than enal in the presence of secondary amine (eq (2)), we hypothesized that insertion of another double bond between carbonyl group and alkyne group¹² where the 1,6-acceptor carbon is sp^2 -hybridized while the 1,4-carbon acceptor is sp^2 -hybridized would be able to promote the 1,6-addition over


1,4-addition (eq (3)). The plausible reasons are: (1) the sp hybridized carbon of a $C\equiv C$ bond is more electron deficient than sp^2 $C=C$ bond; (2) the steric effect may also favor the addition to the $C\equiv C$ bond.

Scheme 4.2 Proposal for Organocatalytic 1,6-Addition


Extensively studied 1,4-addition by iminium catalysis (Eq. 1)



Rapid 1,4-addition of ynals (Eq. 2)



Unprecedented organocatalyzed 1,6-addition of ynals (Eq. 3)




4.3 Results and Discussion

1) Investigation of sulfur-based nucleophiles

With the uncertainty and possibility of the chemistry faced, we started to investigate the simple conjugate addition of various nucleophiles to δ -substituted-2-en-4-ynal substrates. Our initial studies focused on the sulfur nucleophile. At first, the model reaction between (*E*)-5-phenylpent-2-en-4-ynal **1a** and thiophenol in dichloromethane at room temperature in the presence of 20 mol% of organocatalyst **I** was investigated. To my surprise, the reaction indeed happened with the exclusive formation of 1,6-addition product **2a** in high efficiency (87% yield) although the *Z* and *E* mixture were obtained (Scheme 4.3). To improve the *Z/E* selectivity, catalysts with different steric recognition were briefly screened. And it should make sense that the *Z/E* selectivity was not affected at all even simple pyrrolidine was applied because the catalyst is too far away from the δ -carbon position. One of the limitations related with the process is 1,4-addition product was generated exclusively using aliphatic thiol, such as ethanethiol and benzyl mercaptan.


Scheme 4.3 Organocatalytic 1,6-Addition between (*E*)-5-Phenylpent-2-en-4-ynal **1a** and Thiophenol



In effort to gain an insight into the reaction mechanism, the reaction was conducted at lower temperature (-78 °C) or in the presence of lower catalyst loading (3 mol%) and monitored carefully by running crude ¹HNMR. Unexpected, I found the


reaction went through 1,4-addition product **1a-I** which then underwent 1,4-migration of thio group resulted in an allene intermediate **1a-II**. The active allene intermediate **1a-II** would transformed into highly conjugated 1,6-addition product **2a** spontaneously (Scheme 4.4, eq (1)). Another possibility is the 1,3-migration of thio group to form compound **2aa** (eq (2))¹³.

Scheme 4.4 Proposed Mechanism for 1,6-Addition of Thiophenol to (*E*)-5-Phenylpent-2-en-4-ynal **1a**



At this moment, it is necessary to verify the structure of 1,6-addition product **2a**. First, the compound **2a-ester-I** was prepared through Michael addition of thiophenol to 3-phenyl-2-propynal followed by Wittig chain extension (Scheme 4.5, eq (1)). And the ¹HNMR spectra of compound **2a-ester-I** is the same as the compound **2a-ester-II** transformed from the 1,6-addition product **2a**.


Scheme 4.5 Verification of the Structure of 1,6-Addition Product **2a**



2) Investigation of oxygen-based nucleophiles

Then the oxygen nucleophile was investigated. The model reaction between (*E*)-5-phenylpent-2-en-4-ynal **1a** and phenol in dichloromethane at room temperature in the presence of 20 mol% of organocatalyst **I** was tested. Different from thiophenol which is more nucleophilic towards the yenal substrate, the conversion of the reaction is unsatisfactory. Ensuring the complete conversion, the amount of phenol was increased to 2 eq.. Various base additives such as K_2CO_3 , Cs_2CO_3 , LiOH, NaH and *t*-BuOK were evaluated. Without base additive, the reaction proceeded slowly with low yield (28%) after 5 d (entry 1). By adding 0.1eq of K_2CO_3 additive, the reaction yield can be improved a little (entry 2). Accordingly, 1.0 eq of various base additives were tested with results listed in Table 4.1, entries 3~7. Higher yield was observed with 1.0 eq of K_2CO_3 additive (entry 3). The additional raising of the reaction temperature to 45°C leads to similar result as room temperature (entry 8).

Table 4.1 Optimization of Reaction Condition of 1,6-Addition between (*E*)-5-Phenylpent-2-en-4-ynal **1a** and Phenol^a



Entry	Temp. (°C)	Time (h)	Additive(eq)	Yield (%)
1	r.t.	116 (5d)	None	28 ^c
2	r.t.	116 (5d)	K ₂ CO ₃ (0.1eq)	36 ^c
3	r.t.	45.5	K ₂ CO ₃ (1eq)	54 ^b
4	r.t.	69.5	Cs ₂ CO ₃ (1eq)	37 ^c
5	r.t.	65	LiOH (1eq)	23 ^c
6	r.t.	65	NaH (1eq)	32 ^c
7	r.t.	65	<i>t</i> -BuOK (1eq)	29 ^c
8	45	22.5	K ₂ CO ₃ (1eq)	50


^a Reaction conditions: unless specified, a mixture of (*E*)-5-phenylpent-2-en-4-ynal **1a** (0.05mmol), phenol (0.10mmol), catalyst (0.01mmol) and additive in CHCl₃ was stirred at rt for a specified time. ^b Isolated yields. ^c Calculated yield based on crude ¹HNMR.

Crude ¹H NMR was conducted during the reaction process in the presence of 3 mol% of organocatalyst **I**, no 1,4-addition intermediate was observed at all. A controlled study is also performed without catalyst **I**. No reaction occurs even heating for 60 h at 66 °C, indicating that the reaction indeed involves an iminium catalysis.

3) Investigation of nitrogen-based nucleophiles

As for the nitrogen nucleophile, I tried aniline with various electron-withdrawing protection groups such as *para*-methoxysulfonyl, 2,4-dinitrobenzenesulfonyl and trifluoromethanesulfonyl protected aniline. Among those, only trifluoromethanesulfonyl protected aniline **4a** that has the strongest electron-withdrawing electronic property and smallest size act as excellent Michael donor. Gratifyingly, the reaction between (*E*)-5-phenylpent-2-en-4-ynal **1a** and trifluoromethanesulfonyl protected aniline **4a** in chloroform at room temperature in the presence of 20 mol% of organocatalyst **I** give the 1,6-addition product **5a** in 90% yield with 6:1 *Z/E* selectivity (Scheme 4.6, eq (1)). However, in sharp contrast, K₂CO₃ base additive would retard the reaction process which is totally opposite with phenol nucleophile (only 50% conversion was observed in the presence of additional 0.5eq of K₂CO₃ base additive). Besides that, triazole and tetrazole heterocycles were studied. Generally speaking, the nucleophilicity of tetrazole is better than that of triazole. For the 5-phenyl-1*H*-tetrazole nucleophile, in the presence of 20 mol% of organocatalyst **I** in chloroform at room temperature, the reaction proceeded slowly. As expected, conducting the reaction at elevated temperature (60 °C) gave the desired 1,6-addition product **6a** in 62% yield (Scheme 4.6, eq (2)).


Scheme 4.6 1,6-Addition of Nitrogen Nucleophiles to (*E*)-5-Phenylpent-2-en-4-ynal **1a**



4) Substrate scope of 1,6-addition to ynals

The generality of the 1,6-addition was subsequently investigated. A diverse range of aryl-substituted S, O, N nucleophiles and (*E*)-5-phenylpent-2-en-4-ynals **1** were selected to evaluate the scope of the process. As revealed in Table 4.2, the 1,6-addition processes were tolerant of Michael acceptors **1** with significant structural variations. Remarkably, the aromatic moieties bearing electron-withdrawing (entries 6, 10, 12), -neutral (entries 1-3, 5, 9, 11) and -donating (entry 7) substituents could efficiently participate in the process with high efficiency. Furthermore, heteroaromatic ynals also effectively engaged in the 1,6-addition process (entry 4, 8).

Table 4.2 Scope of 1,6-Addition Process ^a



Entry	R	Nu	Solvent	T(°C)	t(h)	#	Z:E	% yield ^b
1	Ph	PhSH	DCM	25	5	2a	5:3	87
2	Ph	4-Methylbenzenethiol	DCM	25	7	2b	5:4	92
3	Ph	2-Chlorobenzenethiol	DCM	25	7	2c	5:3	84
4	2-Thienyl	PhSH	DCM	25	7	2d	3:1	84
5	Ph	PhNHTf	CHCl ₃	25	24	5a	6:1	90
6	4-BrC ₆ H ₄	PhNHTf	CHCl ₃	25	24	5b	7:1	85
7	3-CH ₃ C ₆ H ₄	PhNHTf	CHCl ₃	25	24	5c	7:1	91
8	2-Thienyl	PhNHTf	CHCl ₃	25	24	5d	5:4	95
9	Ph	5-Phenyl-1H-tetrazole	CHCl ₃	60	7	6a	1:1	62
10	4-BrC ₆ H ₄	5-Phenyl-1H-tetrazole	CHCl ₃	60	4.5	6b	1:1	65
11 ^c	Ph	Phenol	CHCl ₃	25	45	3a	6:1	54
12 ^c	4-ClC ₆ H ₄	Phenol	CHCl ₃	25	24	3b	10:1	60

^a Unless specified, see the Supporting Information section for reaction conditions. ^b Isolated yield ^c 2eq of phenol and 1eq of K₂CO₃ were applied.


4.4 Conclusions

In conclusion, we have discovered an unprecedented organocatalytic 1,6-addition to yenals with exclusive δ -regioselectivity to generate synthetically useful dienic aldehydes. This chemistry can be extended to “O”, “S” and “N” based nucleophiles. The cascade reactions initiated by the organocatalytic 1,6-addition to yenals will be further developed in our lab.

4.5 Supporting Information

General procedure for the 1,6-addition reactions:


A solution of yenal **1** (0.10 mmol), organocatalyst **Rac-I** (6.5 mg, 0.02 mmol) in a solvent (0.1 mL) was added nucleophile (0.12 mmol). The resulting solution was stirred at specific temperature for a specified time as described in Table 4.2. Then the reaction solution was directly purified by running column chromatography to give the desired product.



(2E)-5-phenyl-5-(phenylthio)penta-2,4-dienal (**2a**) (Table 4.2, Entry 1):


The title compound was prepared according to the general procedure, as described above in 87% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.68 (d, $J = 8.1$ Hz, 1H), 9.34 (d, $J = 8.1$ Hz, 0.6H), 8.00 (dd, $J_1 = 11.1$ Hz, $J_2 = 15.3$ Hz, 1H), 7.62~7.59 (m, 2H), 7.54~7.51 (m, 1.6H),

7.44~7.39 (m, 5.6H), 7.28~7.25 (m, 3H), 7.20~7.08 (m, 6H), 7.04~6.98 (m, 1.6H), 6.41~6.33 (m, 1H), 6.13~6.09 (m, 0.6H), 5.97~5.89 (m, 0.6H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.8, 193.6, 154.4, 148.1, 147.7, 138.3, 136.0, 134.7, 133.7, 133.3, 131.3, 130.4, 130.0, 129.6, 129.53, 129.46, 128.9, 128.6, 128.42, 128.38, 128.1, 126.8, 122.0.




(2E)-5-phenyl-5-(p-tolylthio)penta-2,4-dienal (2b) (Table 4.2, Entry 2)

The title compound was prepared according to the general procedure, as described above in 92% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.67 (d, $J = 8.0$ Hz, 1H), 9.32 (d, $J = 8.5$ Hz, 0.8H), 8.00 (dd, $J_1 = 11.0$ Hz, $J_2 = 15.5$ Hz, 1H), 7.583~7.576 (m, 2H), 7.43~7.42 (m, 7H), 7.26~7.22 (m, 6H), 7.08 (d, $J = 8.0$ Hz, 2H), 7.02 (dd, $J_1 = 11.5$ Hz, $J_2 = 15.0$ Hz, 1H), 6.94 (d, $J = 9.0$ Hz, 2H), 6.35 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.5$ Hz, 1H), 6.03 (d, $J = 11.5$ Hz, 1H), 5.90 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.0$ Hz, 0.8H), 2.39 (s, 3H), 2.20 (s, 3H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.8, 193.6, 155.2, 148.4, 148.2, 147.8, 139.9, 138.4, 136.9, 136.1, 134.8, 133.0, 130.7, 130.5, 130.3, 129.9, 129.7, 129.4, 129.3, 129.2, 128.5, 128.4, 128.3, 128.0, 126.6, 121.2, 21.3, 20.9.




(2E)-5-(2-chlorophenylthio)-5-phenylpenta-2,4-dienal (2c) (Table 4.2, Entry 3):

The title compound was prepared according to the general procedure, as described above in 84% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.69 (d, $J = 8.0$ Hz, 1H), 9.36 (d, $J = 8.0$ Hz, 0.6H), 7.99 (dd, $J_1 = 11\text{Hz}$, $J_2 = 15.0$ Hz, 1H), 7.61~7.60 (m, 2H), 7.56 (d, $J = 7.5$ Hz, 0.6H), 7.49~7.46 (m, 2H), 7.41~7.40 (m, 2H), 7.32 (t, $J = 7.5$ Hz, 1H), 7.30~7.27 (m, 5H), 7.08~7.02 (m, 4H), 6.99~6.96 (m, 1H), 6.38 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.0$ Hz, 1H), 6.12 (d, $J = 11.5$ Hz, 0.6H), 5.98 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.0$ Hz, 0.6H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.8, 193.6, 151.6, 147.9, 147.5, 146.4, 138.5, 138.0, 136.7, 135.6, 134.1, 133.7, 133.0, 132.2, 131.6, 130.9, 130.5, 130.2, 129.8, 129.6, 128.5, 128.1, 127.9, 127.6, 127.0, 122.9.




(2E)-5-(phenylthio)-5-(thiophen-2-yl)penta-2,4-dienal (2d) (Table 4.2, Entry 4):

The title compound was prepared according to the general procedure, as described above in 84% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.63 (d, $J = 8.0$ Hz, 1H), 9.47 (d, $J = 8.0$ Hz, 0.3H), 7.93 (dd, $J_1 = 11.0$ Hz, $J_2 = 15.0$ Hz, 1H), 7.50~7.46 (m, 1H), 7.41~7.39 (m, 2H), 7.29 (d, $J = 5.0$ Hz, 1H), 7.26~7.14 (m, 6H), 7.09 (t, $J = 4.5$ Hz, 0.3H), 6.94 (t, $J = 4.5$ Hz, 1H), 6.36 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.0$ Hz, 1H), 6.19 (d, $J = 11.5$ Hz, 0.3H), 6.01 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.0$ Hz, 0.3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.6, 147.7, 145.2, 143.7, 138.3, 137.9, 134.7, 134.0, 133.3, 131.1, 130.54, 130.49, 129.9, 129.6, 129.3, 129.2, 128.8, 128.7, 128.3, 128.0, 127.6, 126.7, 123.6.



1,1,1-trifluoro-*N*-((1*Z*)-5-oxo-1-phenylpenta-1,3-dienyl)-*N*-phenylmethanesulfonamide (5a) (Table 4.2, Entry 5):


The title compound was prepared according to the general procedure, as described above in 90% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.65 (d, $J = 8.0$ Hz, 1H), 9.47 (d, $J = 8.0$ Hz, 0.15H), 7.60~7.59 (m, 2H), 7.54~7.37 (m, 9H), 7.31~7.29 (m, 2H), 7.14~7.03 (m, 1.2H), 6.91 (d, $J = 11.0$ Hz, 0.15H), 6.42 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.5$ Hz, 1H), 6.38~6.36 (m, 0.12H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.1, 192.9, 146.8, 146.6, 144.0, 143.7, 139.4, 136.9, 135.43, 135.35, 134.6, 133.2, 130.4, 130.2, 129.8, 129.4, 129.3, 129.1, 129.0, 128.7, 128.5, 127.7, 127.0, 126.5, 123.9, 122.0, 117.7.



***N*-((1*Z*)-1-(4-bromophenyl)-5-oxopenta-1,3-dienyl)-1,1,1-trifluoro-*N*-phenylmethanesulfonamide (5b) (Table 4.2, Entry 6):**


The title compound was prepared according to the general procedure, as described above in 85% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.65 (d, $J = 7.5$ Hz, 1H), 9.48 (d, $J = 7.5$ Hz, 0.1H), 7.54~7.51 (m, 3H), 7.48~7.38 (m, 6H), 7.32~7.28 (m, 1.8H), 7.18 (d, $J = 8.5$ Hz, 0.4H), 7.11 (d, $J = 11.0$ Hz, 1H), 7.02~6.96 (m, 0.15H), 6.90 (d, $J = 11.5$ Hz, 0.15H),

6.43 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.5$ Hz, 1H), 6.38~6.35 (m, 0.1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 192.9, 192.7, 145.7, 145.4, 143.5, 142.5, 139.1, 135.7, 135.1, 134.4, 132.2, 132.1, 131.3, 129.9, 129.6, 129.5, 128.93, 128.87, 128.4, 127.8, 127.0, 124.9, 124.8, 123.8, 121.1, 118.5.




1,1,1-trifluoro-*N*-((1*Z*)-5-oxo-1-*m*-tolylpenta-1,3-dienyl)-*N*-phenylmethanesulfonamide (5c) (Table 4.2, Entry 7):

The title compound was prepared according to the general procedure, as described above in 91% yield. ^1H NMR (CDCl_3 , 500 MHz): δ 9.64 (d, $J = 7.5$ Hz, 1H), 9.47 (d, $J = 8.0$ Hz, 0.14H), 7.53~7.50 (m, 1H), 7.47~7.45 (m, 2H), 7.42~7.37 (m, 4H), 7.31~7.28 (m, 3H), 7.23~7.18 (m, 1.5H), 7.13~7.07 (m, 1.5H), 6.88 (d, $J = 11.0$ Hz, 0.14H), 6.41 (dd, $J_1 = 8.0$ Hz, $J_2 = 15.0$ Hz, 1H), 6.37~6.34 (m, 0.1H), 2.37 (s, 3H), 2.35 (s, 0.4H); ^{13}C NMR (CDCl_3 , 75 MHz): δ 193.1, 193.0, 147.0, 146.9, 144.1, 143.9, 139.5, 138.8, 138.5, 136.9, 135.4, 135.2, 134.4, 133.1, 131.3, 131.0, 130.2, 129.8, 129.6, 129.4, 129.3, 129.1, 128.8, 128.5, 128.2, 127.6, 127.2, 126.3, 124.3, 123.9, 123.4, 122.0, 117.7, 113.4, 21.4, 21.3.



1,1,1-trifluoro-*N*-((1*Z*)-5-oxo-1-(thiophen-2-yl)penta-1,3-dienyl)-*N*-phenylmethanesulfonamide (5d) (Table 4.2, Entry 8):


The title compound was prepared according to the general procedure, as described above in 95% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.69 (d, $J = 7.8$ Hz, 1H), 9.58 (d, $J = 7.8$ Hz, 0.8H), 7.58~7.46 (m, 4H), 7.43~7.31 (m, 9.8H), 7.18~7.17 (m, 0.8H), 7.08 (d, $J = 11.4$ Hz, 1H), 7.05~7.01 (m, 1.8H), 6.95 (d, $J = 11.1$ Hz, 0.8H), 6.47~6.39 (m, 1.8H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.0, 192.9, 146.0, 143.9, 139.9, 139.5, 138.8, 137.6, 137.2, 135.8, 135.2, 134.8, 132.0, 130.0, 129.9, 129.5, 129.3, 128.9, 128.8, 128.5, 128.34, 128.26, 128.1, 127.7, 127.2, 124.4, 123.9, 121.3, 121.0, 118.7, 118.5.



(2*E*)-5-phenyl-5-(5-phenyl-2*H*-tetrazol-2-yl)penta-2,4-dienal (6a) (Table 4.2, Entry 9):


The title compound was prepared according to the general procedure, as described above in 62% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.58 (d, $J = 7.8$ Hz, 1H), 9.56 (d, $J = 7.8$ Hz, 1H), 8.26~8.19 (m, 4H), 7.60~7.43 (m, 16H), 7.31~7.27 (m, 2H), 7.23~7.15 (m, 2H), 7.07 (d, $J = 11.4$ Hz, 1H), 6.51 (dd, $J_1 = 7.8$ Hz, $J_2 = 12.9$ Hz, 1H), 6.46 (dd, $J_1 = 7.8$ Hz, $J_2 = 12.6$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.1, 192.8, 165.6, 165.1, 145.4,

143.5, 142.9, 141.5, 135.9, 135.7, 133.7, 131.0, 130.9, 130.9, 130.6, 130.1, 129.03, 128.95, 128.8, 127.19, 127.15, 127.1, 126.55, 126.47, 123.8, 120.7.




(2E)-5-(4-bromophenyl)-5-(5-phenyl-2H-tetrazol-2-yl)penta-2,4-dienal (6b) (Table 4.2, Entry 10):

The title compound was prepared according to the general procedure, as described above in 65% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.58 (d, $J = 7.8$ Hz, 1H), 9.58 (d, $J = 7.8$ Hz, 1H), 8.24~8.17 (m, 4H), 7.75~7.66 (m, 3H), 7.61~7.50 (m, 9H), 7.39~7.33 (m, 2H), 7.22~7.13 (m, 3H), 7.04 (d, $J = 11.4$ Hz, 1H), 6.52 (dd, $J_1 = 7.8$ Hz, $J_2 = 13.8$ Hz, 1H), 6.47 (dd, $J_1 = 7.8$ Hz, $J_2 = 13.5$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.0, 192.7, 165.7, 165.2, 144.6, 143.2, 141.6, 140.3, 136.3, 136.1, 132.7, 132.3, 132.2, 132.1, 131.1, 131.0, 129.1, 129.0, 128.6, 127.2, 126.4, 126.3, 125.6, 125.5, 124.0, 121.0.



(2E)-5-phenoxy-5-phenylpenta-2,4-dienal (3a) (Table 4.2, Entry 11):

The title compound was prepared according to the general procedure, as described above in 54% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.55 (d, $J = 7.8$ Hz, 1H), 9.41 (d, $J = 8.1$ Hz, 0.15H), 7.62~7.57 (m, 3H), 7.54~7.49 (m, 0.8H), 7.44~7.33 (m, 3.5H), 7.32~7.23 (m, 2.5H), 7.14~7.12 (m, 0.3H), 7.03~6.95 (m, 3H), 6.68 (d, $J = 11.4$ Hz, 1H), 6.29 (dd, $J_1 = 8.1$ Hz, $J_2 = 15.6$ Hz, 1H), 6.02 (dd, $J_1 = 8.1$ Hz, $J_2 = 15.3$ Hz, 0.15H), 5.81 (d, $J = 11.4$ Hz, 0.15H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.8, 193.2, 165.6, 157.2, 149.9, 145.7, 133.5, 131.5, 130.4, 130.1, 130.0, 129.8, 129.6, 129.5, 128.8, 128.6, 126.7, 125.2, 122.7, 120.8, 116.3, 114.6, 107.4.



(2E)-5-(4-chlorophenyl)-5-phenoxy-penta-2,4-dienal (3b) (Table 4.2, Entry 12):

The title compound was prepared according to the general procedure, as described above in 60% yield. ^1H NMR (CDCl_3 , 300 MHz): δ 9.54 (d, $J = 8.1$ Hz, 1H), 9.42 (d, $J = 7.8$ Hz, 0.1H), 7.59~7.46 (m, 3.4H), 7.42~7.39 (m, 0.4H), 7.32~7.20 (m, 4H), 7.12~7.09 (m, 0.2H), 7.05~7.00 (m, 1H), 7.00~6.93 (m, 2H), 6.65 (d, $J = 11.1$ Hz, 1H), 6.29 (dd, $J_1 = 7.8$ Hz, $J_2 = 15.3$ Hz, 1H), 6.03 (dd, $J_1 = 8.1$ Hz, $J_2 = 15.3$ Hz, 0.1H), 5.81 (d, $J = 11.7$ Hz, 0.1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 193.7, 193.0, 157.0, 156.0, 154.3, 149.0, 145.3, 136.5, 136.1, 132.1, 131.9, 130.7, 130.1, 130.0, 129.9, 129.1, 129.0, 128.0, 125.3, 122.9, 120.7, 116.2, 114.9, 107.7.

4.6 Reference

[1] For recent reviews on catalytic 1, 4-addition reactions, see: (a) N. Krause, A. Hoffmann-Röder, *Synthesis* **2001**, 171-196; (b) M. P. Sibi, S. Manyem, *Tetrahedron* **2000**, *56*, 8033-8061; (c) K. Tomioka, Y. Nagaoka in *Comprehensive Asymmetric Catalysis*, Vol. 3 (Eds. : E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer, Berlin, **1999**, chap. 31.1, pp. 1105-1120; (d) M. Kanai, M. Shibasaki in *Catalytic Asymmetric Synthesis*, 2nd ed. (Ed. : I. Ojima), Wiley, New York, **2000**, p. 569; (e) P. Perlmutter, *Conjugate Addition Reactions in Organic Synthesis*. (Tetrahedron Organic Chemistry Series, Vol. 9, series editor: J. E. Baldwin), **2003**; (f) B. H. Lipshutz in *Organometallics in Organic Synthesis. A Manual*, 2nd ed. (Ed. : M. Schlosser), Wiley, Chichester, **2002**, p. 665; (g) J. Christoffers, G. Koripelly, A. Rosiak, M. RJsle, *Synthesis* **2007**, 1279-1300.

[2] For selected reviews on enantioselective copper-catalyzed conjugate addition, see: (a) A. Alexakis, C. Benhaim, *Eur. J. Org. Chem.* **2002**, 3221-3236; (b) B. L. Feringa, R. Naasz, R. Imbos, L. A. Arnold, *Modern Organocopper Chemistry* (Ed. : N. Krause), Wiley-VCH, Weinheim, **2002**, p. 224-258; (c) B. L. Feringa, *Acc. Chem. Res.* **2000**, *33*, 346-353. For selected reviews on enantioselective rhodium-catalyzed conjugate addition, see: (d) T. Hayashi, *Synlett* **2001**, 879-887; (e) K. Fagnou, M. Lautens, *Chem. Rev.* **2003**, *103*, 169-196; (f) T. Hayashi, K. Yamasaki, *Chem. Rev.* **2003**, *103*, 2829-2844; (g) T. Hayashi, *Bull. Chem. Soc. Jpn.* **2004**, *77*, 13-21. For a recent example of palladium, see: (h) T. Nishikata, Y. Yamamoto, N. Miyaura, *Chem. Commun.* **2004**, 1822-1823. For a recent example of nickel, see: (i) Y.-S. Kwak, E. J. Corey, *Org. Lett.* **2004**, *6*, 3385-3388.

[3] For recent reviews on organocatalytic asymmetric Michael additions, see: (a) D. Almasi, D. A. Alonso, C. Najera, *Tetrahedron: Asymmetry*, **2007**, *18*, 299-365; (b) S. B. Tsogoeva, *Eur. J. Org. Chem.*, **2007**, *11*, 1701-1716; (c) D. Enders, C. Wang, J. X. Liebich, *Chem. Eur. J.* **2009**, *15*, 11058-11076; (d) J. L. Vicario, D. Badía, L. Carrillo, *Synthesis* **2007**, 2065-2092; (e) S. Sulzer-Mossé, A. Alexakis, *Chem. Commun.* **2007**, 3123-3135.

[4] For recent examples of metal-catalyzed nonasymmetric 1,6-selective conjugate addition, see: (a) K. Fukuhara, H. Urabe, *Tetrahedron Lett.* **2005**, *46*, 603-606; (b) G. de laHerrán, C. Murcia, A. G. Csáký, *Org. Lett.* **2005**, *7*, 5629-5632; (c) G. de laHerrán, A. G. Csáký, *Synlett* **2009**, 585-588; (d) T. Nishimura, Y. Yasuhara, T. Hayashi, *Angew. Chem. Int. Ed.* **2006**, *45*, 5164-5166.

[5] (a) T. den Hartog, S. R. Harutyunyan, D. Font, A. J. Minnaard, B. L. Feringa, *Angew. Chem. Int. Ed.* **2008**, *47*, 398-401; (b) H. Hénon, M. Mauduit, A. Alexakis, *Angew. Chem. Int. Ed.* **2008**, *47*, 9122-9124; (c) K.-S. Lee, A. H. Hoveyda, *J. Am. Chem. Soc.* **2010**, *132*, 2898-2900; (d) E. Fillion, A. Wilsily, E-T. Liao, *Tetrahedron: Asymmetry* **2006**, *17*, 2957-2959.

[6] (a) T. Hayashi, S. Yamamoto, N. Tokunaga, *Angew. Chem. Int. Ed.* **2005**, *44*, 4224-4227; (b) T. Hayashi, N. Tokunaga, K. Inoue, *Org. Lett.* **2004**, *6*, 305-307.

[7] T. Nishimura, Y. Yasuhara, T. Sawano, T. Hayashi, *J. Am. Chem. Soc.* **2010**, *132*, 7872-7873.

[8] S. Okada, K. Arayama, R. Murayama, T. Ishizuka, K. Hara, N. Hirone, T. Hata, H.

Urabe, *Angew. Chem. Int. Ed.* **2008**, *47*, 6860-6864.

[9] (a) A. G. Csákÿ, G. de la Herrán, M. C. Murcia, *Chem. Soc. Rev.* **2010**, *39*, 4080-4102; (b) S. Belot, A. Massaro, A. Tenti, A. Mordini, A. Alexakis, *Org. Lett.* **2008**, *10*, 4557-4560; (c) S. Belot, K. A. Vogt, C. Besnard, N. Krause, A. Alexakis, *Angew. Chem. Int. Ed.* **2009**, *48*, 8923-8926; (d) S. Belot, A. Quintard, N. Krause, A. Alexakis, *Adv. Synth. Catal.* **2010**, *352*, 667-695; (e) T. He, J.-Y. Qian, H.-L. Song, X.-Y. Wu, *Synlett*, **2009**, 3195-3197; (f) C. G. Oliva, A. M. S. Silva, F. A. A. Paz, J. A. S. Cavaleiro, *Synlett*, **2010**, 1123-1127.

[10] L. Bernardi, J. López-Cantarero, B. Niess, K. A. Jørgensen, *J. Am. Chem. Soc.* **2007**, *129*, 5772-5778.

[11] J. J. Murphy, A. Quintard, P. McArdle, A. Alexakis, J. C. Stephens, *Angew. Chem. Int. Ed.* **2011**, *50*, 5095-5098.

[12] Enantioselective reductive coupling of 1,3-enynes to heterocyclic aromatic aldehydes and ketones via rhodium-catalyzed asymmetric hydrogenation, see: V. Komanduri, M. J. Krische, *J. Am. Chem. Soc.* **2006**, *128*, 16448-16449.

[13] N. Yoshikai, T. Yamashita, E. Nakamura, *Chem. Asian J.* **2006**, *1*, 322-330.