


11-4-2017

Virulence Phenotypes of *Rotylenchulus reniformis*: Evaluation of Host Status of Cotton and Utility of Single Nucleotide Polymorphisms (SNPs) for Identification

Churamani Khanal

Louisiana State University and Agricultural and Mechanical College, ckhana1@lsu.edu

Follow this and additional works at: https://digitalcommons.lsu.edu/gradschool_dissertations

 Part of the [Agricultural Science Commons](#), [Molecular Genetics Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Khanal, Churamani, "Virulence Phenotypes of *Rotylenchulus reniformis*: Evaluation of Host Status of Cotton and Utility of Single Nucleotide Polymorphisms (SNPs) for Identification" (2017). *LSU Doctoral Dissertations*. 4129.
https://digitalcommons.lsu.edu/gradschool_dissertations/4129

This Dissertation is brought to you for free and open access by the Graduate School at LSU Digital Commons. It has been accepted for inclusion in LSU Doctoral Dissertations by an authorized graduate school editor of LSU Digital Commons. For more information, please contact gradetd@lsu.edu.

**VIRULENCE PHENOTYPES OF *ROTYLENCHULUS RENIFORMIS*:
EVALUATION OF HOST STATUS OF COTTON AND UTILITY OF
SINGLE NUCLEOTIDE POLYMORPHISMS (SNPS) FOR
IDENTIFICATION**

A Dissertation

Submitted to the Graduate Faculty of the
Louisiana State University and
Agricultural and Mechanical College
in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

in

The Department of Plant Pathology and Crop Physiology

by

Churamani Khanal

B.S., Tribhuvan University, Nepal, 2010

M.S., University of Arkansas, 2014

December 2017

I would like to dedicate this dissertation to my parents Pitambar Khanal and Jamuna Khanal and loving wife Durga Bhattarai

ACKNOWLEDGMENTS

I would like to thank my parents (Pitambar Khanal and Jamuna Khanal), my living god. The academic milestone I have achieved today was the result of foundation of education they have built in me from childhood. I am really thankful to my dear wife, Durga Bhattarai, who has been a great support throughout my Ph.D. studies. It would not be possible to accomplish this degree without her love, care, and patience. I am blessed to have Dr. Edward C. McGawley as my major advisor. He always encouraged me to do a better science. He has become more of my friend rather than an adviser. He was always there to help me whether it were my personal or academic problems. It would not be possible to achieve this success without his guidance, help, and endless motivation. I am equally thankful to my co-advisor Dr. Charles Overstreet for being a source of motivation and constant support. I have been a big fan of his pleasant personality and social behavior. He exposed me to field experiments and made me more capable of conducting any experiments independently in future. Without his support, this degree would not have been possible.

I would like to express my gratitude to the members of my graduate committee, Dr. Salliana R. Stetina, Dr. Gerald O. Myers, and Dr. James M. Fannin for their time and guidance over the last several years. Dr. Stetina has been a great help from raising cotton seeds to guidance about the molecular part of my research. Similarly, Dr. Myers provided regular guidance and support throughout the research. I am also thankful to Dr. Fannin for sparing time during important events of my Ph.D. studies.

I am indebted to help I received from Dr. Jeffery D. Ray. He has been a very important part of my research involving genetic aspects of reniform nematode. Also, I am obliged to Ms. Angie Davis for her help during our visit to USDA ARS, Mississippi. My sincere thank you goes

to Dr. David Blouin of the Department of Experimental Statistics at LSU for review and approval of experimental designs and data analysis employed in this research.

Many thanks go out to Ms. Claudette Oster, Stephen Salzer, Josh Granger, and Bill Salzer for their help during greenhouse and microplot research. I am thankful for the help that my colleagues Manjula Kularathna, Benjamin McInnes, Felipe Godoy provided during experiment setup and processing of samples. Also, thanks goes to Deborah Xavier-Mis, research associate of Dr. Overstreet, who helped order lab and greenhouse supplies. I and my wife enjoyed hospitality of Ms. Helen and Ms. Karen. I will never forget the Christmas gift from Ms. Helen and delicious food from Ms. Karen in Thanksgiving. I really enjoyed interacting with other professors and students in the department of plant pathology and crop physiology at LSU. I sincerely thank Ms. Dolores Dyess, Robert M. Carver, and Charletta Warr for their help in administrative process. All of these people will always be in my memory.

TABLE OF CONTENTS

ACKNOWLEDGMENTS	iii
ABSTRACT.....	vi
CHAPTER 1. INTRODUCTION.....	9
1.1 Cotton.....	9
1.2 Cotton growth and development	10
1.3 Nematodes of cotton.....	11
1.4 The reniform nematode	11
1.5 Crop losses by reniform nematode.....	13
1.6 Reniform nematode management.....	14
CHAPTER 2. REPRODUCTION AND PATHOGENICITY OF ENDEMIC POPULATIONS OF <i>ROTYLENCHULUS RENIFORMIS</i> ON COTTON	16
2.1 Introduction	16
2.2 Materials and methods	18
2.3 Results	21
2.4 Discussion	32
CHAPTER 3. SINGLE NUCLEOTIDE POLYMORPHISM ANALYSIS PROVIDES AN INSIGHT ON GENETIC VARIABILITY IN <i>ROTYLENCHULUS RENIFORMIS</i>	36
3.1 Introduction	36
3.2 Methodology	39
3.3 Results	46
3.4 Discussion	51
CHAPTER 4. SUMMARY AND CONCLUSIONS	56
REFERENCES	59
APPENDIX 1. FLANKING SEQUENCE INFORMATION FOR SNP ASSAYS TESTED IN THIS STUDY. THE "SNP ID" PROVIDES A COMMON NAME INCLUDING ORGANISM ABBREVIATION "RREN" <i>ROTYLENCHULUS RENIFORMIS</i> (NEMATODES), THE REFERENCE ASSEMBLY CONTIG (FIRST NUMBER) AND THE SEQUENCE LOCATION (SECOND NUMBER) OF THE SNP POSITION.	68
APPENDIX 2. FLANKING SEQUENCE INFORMATION FOR UNTESTED SNP. THE "SNP ID" PROVIDES A COMMON NAME INCLUDING ORGANISM ABBREVIATION "RREN" <i>ROTYLENCHULUS RENIFORMIS</i> , THE REFERENCE ASSEMBLY CONTIG (FIRST	

NUMBER), AND THE SEQUENCE LOCATION (SECOND NUMBER) OF THE SNP POSITION.	76
APPENDIX 3: LETTER REQUESTING PERMISSION TO USE COMMON METHODOLOGY IN CHAPTER 3	109
APPENDIX 4: LETTER OF PERMISSION TO USE COMMON METHODOLOGY IN CHAPTER 3	110
VITA.....	111

ABSTRACT

Comparative reproduction and pathogenicity of reniform nematode (*Rotylenchulus reniformis*) populations derived from single-egg masses and collected from West Carroll (WC), Rapides (RAP), Morehouse (MOR), and Tensas (TEN) parishes in Louisiana were evaluated in microplot and greenhouse trials. Data from microplot trials showed significant differences among isolates of reniform nematode in both reproduction and pathogenicity on upland cotton (*Gossypium hirsutum*) cultivars PhytoGen 499 WRF, Deltapine 1133 B2RF, and PhytoGen 333 WRF. Across all cotton cultivars, MOR and RAP isolates had the greatest and the least reproduction values of 331.8 and 230.2, respectively. Reduction in plant dry weight, number of bolls, seed cotton weight, and lint weight was the greatest and the least for MOR and RAP isolate, respectively. The reproduction and pathogenicity of WC and TEN isolates were intermediate. In the greenhouse experiment, reproduction of MOR and RAP isolates across all cotton genotypes (three cultivars used in microplot experiment, cultivar Stoneville 4946 GLB2, and two resistant germplasm lines MT2468 Ren3 and M713 Ren5) was the greatest (reproduction value 10.7) and the least (reproduction value 7.9), respectively. Although reproduction values of reniform nematode were lower in the germplasm lines than the cultivars, the germplasm lines sustained greater plant weight loss. The variability in reproduction and pathogenicity among endemic populations of reniform nematode in both the microplot and greenhouse experiments adds further support to the hypothesis that virulence phenotypes of *R. reniformis* exist. In order to determine genetic variability in reniform nematode populations, 31 KASP SNP primers sets were evaluated against 13 reniform nematode isolates that include MOR and RAP isolates from Louisiana as well as other 11 isolates from Mississippi, Arkansas, Hawaii, and Alabama. Twenty-six SNP assays amplified genomic DNA of reniform nematode

isolates while five failed to successfully amplify. Five SNP assays identified genetic differences within and among populations of reniform nematode from Louisiana, Mississippi, and Arkansas. Similarly, eight SNP assays identified genetic differences among samples from Hawaii, and Alabama. The SNP markers developed in this study will be useful in resistance breeding programs as well as in the assessment of the genetic diversity, origin, and subsequent distribution of this nematode.

CHAPTER 1. INTRODUCTION

1.1 Cotton

Cotton (*Gossypium* spp.) is a very important textile fiber crop with an estimated worldwide production of 125 million bales (Dighe *et al.*, 2009; Dabbert and Gore, 2014). The genus *Gossypium* comprises almost 50 species, four of which are commercial cotton crops (Wallace *et al.*, 2009). Cotton is principally cultivated for the production of natural fiber for the textile industry. In addition to cotton fiber, cottonseed is also regarded as an economically important product of cotton. Approximately 6 million tons of cottonseed is produced annually in the United States (Koenning *et al.*, 2004; Anonymous, 2017). Cottonseed is utilized in the animal feed industry as an excellent source of protein (Koenning *et al.*, 2004). Similarly, a high quality vegetable oil, equivalent to or better than soybean oil suitable for human consumption, is produced from cotton (Mauney and Stewart, 1986; Thorp *et al.*, 2014). Cottonseed oil is also a possible source of biofuel production (Thorp *et al.*, 2014).

Cotton production in the United States is intensive and highly mechanized (Koenning *et al.*, 2004; Starr *et al.*, 2007). The United States is one of the leading cotton producers that accounts for almost a quarter of world lint supply (Koenning *et al.*, 2004). In terms of cotton production, the United States ranks third after India and China (Meyer and MacDonald, 2016). Most of the United States cotton is upland cotton (*Gossypium hirsutum* L.) grown as an annual crop from seed planted each year (Anonymous, 2010). Commercial production of upland cotton is mainly carried out in the southern United States with concentrations in the Texas High Plains, the southeast, and the Mississippi Delta (Anonymous, 2010). According to United States Department of Agriculture, National Agricultural Statistics Service (USDA-NASS) report of 2017, Texas is the top cotton producer in the United States followed by Georgia, Mississippi,

Arkansas, Alabama, Oklahoma, Tennessee, Missouri, Arizona, North Carolina, and Louisiana (Anonymous, 2017). In 2016, upland cotton was planted in approximately 4 million hectares with production of approximately 16 million bales (Anonymous, 2017).

Cotton is one of the most important row crops in Louisiana (Lofton *et al.*, 2014). In 2016, upland cotton was planted on 57 thousand hectares in Louisiana with an estimated yield of 260 thousand bales (Anonymous, 2017). A bale of cotton contains 480 pounds of lint.

1.2 Cotton growth and development

Cotton is a woody perennial crop which passes through distinct developmental stages. An excellent overview of growth and development of cotton has been provided by Oosterhuis and Bourland (2001), and Mauney (2015). Briefly, under favorable conditions for germination, cotton seedlings emerge from soil in four to nine days after planting. Elongation of the apical meristem gives rise to a main stem which has monopodial and indeterminate growth habit meaning the terminal bud continues to grow as a central leader shoot. The main stem contains several nodes and internodes and branches develop from a bud located at a node in a leaf axil. Branches and leaves are spirally arranged on the stem in a three to eight phyllotaxy. Cotton produces vegetative and fruiting branches. Vegetative branches have monopodial growth while fruiting branches have sympodial growth meaning the fruiting branches terminate in a square. Fruiting branches are produced by both the main stem and vegetative branches. First square occurs 30-40 days after emergence. Fertilization followed by anthesis takes place near dawn in 20-25 days after the development of square. Flower consists of three bracts, calyx whorl (five sepals), corolla (five petals), and 4- or 5-locule ovary which contains six to nine ovules. Cotton is a self-pollinated crop, but cross pollination, predominantly by insects, can also occur. Boll maturity is reached in about 50 days after fertilization. Fibers are produced by extension of

epidermal cells of seed coat. Fibers are first elongated which is followed by secondary thickening. Approximately 500 bolls are required to produce a kilogram of cotton fiber. Depending on growing conditions, cotton is harvested in 130 to 160 days after planting.

1.3 Nematodes of cotton

Cotton is attacked by several plant parasitic nematodes. The predominant plant parasitic nematodes of cotton are the southern root-knot nematode (*Meloidogyne incognita*), and the reniform nematode (*Rotylenchulus reniformis*) (Koenning *et al.*, 2004; Starr *et al.*, 2011). Amount of crop losses caused by southern root-knot and reniform nematodes is more or less equally divided between the two (Weaver, 2015). Other economically important nematodes of cotton are sting nematode (*Belonolaimus longicaudatus*), and Columbia lance nematode (*Hoplolaimus columbus* Sher) (Weaver, 2015). Economic damage threshold of sting and lance nematode on cotton is approximately one nematode per cubic centimeter of soil (Noe, 1993; Crow *et al.*, 2000). Sting and lance nematode are distributed in very sandy soils (>80% sand) and thus have little impact on cotton as very little cotton is grown in heavily sandy soil (Weaver, 2015).

1.4 The reniform nematode

Reniform nematode (*Rotylenchulus reniformis* Linford and Oliveira) was first observed in Oahu island of Hawaii in 1940 attacking cowpea, pineapple, and several weeds (Linford and Oliveira, 1940). The adult female of this nematode is kidney shaped and hence got the vernacular name. The genus *Rotylenchulus* was proposed because of its morphological similarity to *Rotylenchus* and life history similar to *Tylenchulus*. Reniform nematode belongs to the superfamily Heteroderoidea and the family Nacobbidae. The genus *Rotylenchulus* has 10 previously described species (Robinson *et al.*, 1997) and a newly described species (Berg *et al.*,

2016). Of the 11 described species, only *R. reniformis* and *R. parvus* have been reported from the United States (Lehman and Inserra, 1990).

Reniform nematodes are categorized into five groups based on lip morphology and hyaline length of tail, with *R. reniformis* being the only species in group III (Robinson *et al.*, 1997). Some diagnostic features of *R. reniformis* are the presence of males, adult female is 324-383µm long by 161-168 µm wide, 16-21µm long stylet, head sharply set off from the expanded neck region, posterior vulva ($v>63\%$), later stage becomes immobile and develops feeding site, and posterior part of the body enlarges assuming reniform shape (Linford and Oliveira, 1940). Juveniles, young females, and males are vermiform in shape and dorsal gland opening is usually one stylet length behind stylet (Linford and Oliveira, 1940).

Following first report of occurrence from Hawaii, *Rotylenchulus reniformis* was reported from cotton and cowpea roots in Baton Rouge, Louisiana in 1941 (Smith and Taylor, 1941). Subsequently, scientists from other parts of the United States reported occurrence of this nematode. Currently, reniform nematode is the predominant parasitic nematode of upland cotton in the mid-south area of the United States (Stetina and Young, 2006; Robinson, 2007; Starr *et al.*, 2011).

An excellent overview of life cycle of reniform nematode has been provided by Robinson *et al.* (1997). Briefly, a mature adult female lays eggs covered in a gelatinous mass. Upon embryogenesis, first stage juvenile (J1) is formed. J1 molts and becomes second stage juvenile (J2) which emerges from the egg by breaking the egg shell with the help of stylet. Subsequent molting gives rise to third and fourth stage juveniles (J3 and J4). The J4 becomes adult and differentiates into either male or female. Only the J4 female is infective while male remains non infective. The adult female infects the host root cortex, develops a feeding site, and becomes

sedentary. Only the anterior part of the body is embedded in the root and posterior part swells as the nematode feeds and gains reproductive maturity. The mature adult female lays approximately 60 eggs covered in a gelatinous mass. This process completes the life cycle of reniform nematode. Multiple life cycles can occur in a single crop growing season.

Under optimum condition, life cycle of reniform nematode can be completed in as few as 25 days. In other words, the minimum life cycle can be roughly divided as 8 days for hatching, 8 days for completion of molting, and 9 days for laying eggs (Linford and Oliveira, 1940). The life cycle can be extended to several years if nematodes enter the state of anhydrobiosis (Birchfield and Martin, 1967; Gaur and Perry, 1991).

1.5 Crop losses by reniform nematode

Reniform nematodes are distributed in tropical and subtropical regions of the world attacking at least 314 plant species belonging to 77 families (Robinson *et al.*, 1997). The reason for its wide distribution may be attributed to the wide host range and the ability to survive in a state of anhydrobiosis for several years without host plants (Birchfield and Jones, 1961; Birchfield and Martin, 1967; Gaur and Perry, 1991). Using a stylet, the adult female nematode punctures host cells and delivers proteolytic and pectolytic enzymes to degrade cell wall (Dieterich and Sommer, 2009). Once inside host, nematode differentiates host cells to form multinucleated syncytial cells as a source of food. Upon infection, reniform nematode adversely affects plant growth, delays flowering and fruiting times, reduces number and size of the bolls, and decreases lint quality (Robinson, 2007). According to Dighe *et al.* (2009), cotton production in the United States has been greatly compromised by reniform nematodes. In upland cotton, loss to reniform nematode in 2016 was approximately 205 thousand bales including 8 thousand bales from Louisiana (Lawrence *et al.*, 2017).

In Louisiana reniform nematode has become the major nematode pathogen of cotton and soybean over the last three decades (Overstreet and McGawley, 1994; Overstreet and McGawley, 1998; Overstreet, 2006, 2015). According to Robinson *et al.* (2007), Louisiana, Mississippi, and Alabama sustain almost three quarters of the United States cotton crop losses to the reniform nematode. Each year approximately 3 to 5% of the Louisiana cotton crop is lost as a result of reniform nematode pathology (Bell *et al.*, 2014; Lawrence *et al.*, 2017).

1.6 Reniform nematode management

Some common management methods for reniform nematode include, but are not limited to, chemical, biological, crop rotation, tolerance, and resistance. Commonly used chemicals for reniform nematode management are fumigants (Vapam, Telone), non-fumigants (aldicarb, Vydate, Velum), and seed treatments (Avicta, Aeris). For economic and environmental reasons, the use of nematicides is not a preferred management option (Agudelo *et al.*, 2005; Blessitt *et al.*, 2012). Some biological control agents such as *Pasteuria penetrans*, *Bacillus firmus*, and *Paecilomyces lilacinus* have been effective to manage reniform nematode in lab conditions, however, their efficacy is greatly reduced under field conditions. Crop rotation with non-host crops such as corn and resistant soybean reduces the nematode populations greatly; however, the populations usually resurge quickly in a subsequent single year of cotton production (Robinson *et al.*, 2007, Stetina *et al.*, 2007). Crop rotation with milo, peanuts, and sugarcane can also be used to reduce reniform nematodes (Overstreet *et al.*, 2014). Use of reniform nematode resistant cotton has been a most widely desired method of reniform nematode. More than three decades of research in developing reniform nematode resistant upland cotton cultivars has not been successful. Because no commercial cotton cultivars that are resistant to reniform nematode are

available, the best management options currently available are the use of tolerant cultivars and crop rotation.

CHAPTER 2. REPRODUCTION AND PATHOGENICITY OF ENDEMIC POPULATIONS OF *ROTYLENCHULUS RENIFORMIS* ON COTTON

2.1 Introduction

The reniform nematode (*Rotylenchulus reniformis* Linford and Oliveira) was first observed in Hawaii in 1940 attacking cowpea, pineapple and several weeds (Linford and Oliveira, 1940). Subsequently, this nematode was reported from cotton and cowpea roots in Baton Rouge, Louisiana in 1941 (Smith and Taylor, 1941). Currently, reniform nematode is the predominant parasitic nematode of upland cotton in the mid-south area of the United States (Stetina and Young, 2006; Robinson, 2007; Starr *et al.*, 2011). The female nematode infects cotton roots producing approximately 60 eggs per egg mass in as less as 25 days (Linford and Oliveira, 1940). Upon infection, reniform nematode adversely affects plant growth, delays flowering and fruiting times, reduces number and size of the bolls, and decreases lint quality (Robinson, 2007). Because of crop damage caused by reniform nematode, cotton production in the United States has been greatly compromised (Dighe *et al.* 2009). Each year approximately 205 thousand bales of United States upland cotton is lost to reniform nematode (Lawrence *et al.*, 2017).

With an aim of durable management of reniform nematode in upland cotton production, studies on identification of resistant cultivars and germplasm lines have been conducted since early 1960s. Birchfield and Brister (1963) evaluated 24 upland cotton cultivars in the greenhouse and found that none were resistant to reniform nematode. Yik and Birchfield (1984) reported that *Gossypium longicalyx* was a non-host while *G. stocksii*, *G. somalense*, and *G. barbadense* 'Texas 110' showed high levels of resistance. Robinson *et al.* (1999, 2007) emphasized the absence of a source of resistance to the reniform nematode as a major constraint of upland cotton production

in the United States. In 2007, two cotton germplasm lines, LONREN-1 and LONREN-2, having a resistance gene introgressed from *G. longicalyx* Hutch. & Lee were released by the USDA-ARS, Texas Agricultural Experiment Station and Cotton Incorporated (Bell *et al.*, 2014a). Reniform nematode resistance genes *Ren*₁ (previously *Ren*^{lon}), *Ren*₂ (previously *Ren*^{ari}), and *Ren*₂^{GB713} have been identified from *G. longicalyx*, *G. aridum* (Rose & Standl.) Skov, and *G. barbadense* L. GB713 (PI 608139), respectively (Dighe *et al.*, 2009; Romano *et al.*, 2009; Fang and Stetina, 2011; Bell *et al.*, 2014b). Subsequently, reniform nematode resistant germplasm lines TAM RKRNR-9, TAM RKRNR-12, and BARBREN-713 were released (Starr *et al.*, 2011; Bell *et al.*, 2014b). All of these germplasm lines suppressed reniform nematode reproduction by 40-90% (Starr *et al.*, 2011; Bell *et al.*, 2014a, 2014b). Similarly, a joint effort by USDA, ARS, Mississippi State University, and College Station, Texas released several sources of reniform nematode resistant germplasm lines that include TX 110, M713 Ren1, M713 Ren2, M713 Ren5, MT2468 Ren1, MT2468 Ren2, and MT2468 Ren3 (Wallace *et al.*, 2009; McCarty *et al.*, 2012, 2013).

Because no commercial cotton cultivars that are resistant to reniform nematode are available, the best management options currently available are the use of tolerant cultivars and crop rotation. A field study conducted by Blessitt *et al.* (2012) to evaluate cotton cultivars in Mississippi identified six cultivars viz. Cropland Genetics 3520 B2RF, DynaGrow 2520 B2RF, Stoneville 5242 BR, Stoneville 5599 BR, Deltapine 488 BG/RR, and Fibermax 960 B2R as tolerant to reniform nematodes. Tolerant cultivars are those that have the capacity to support reproduction while sustaining satisfactory yields (Schafer, 1971; Blessitt *et al.*, 2012). Crop rotation with non-host crops such as corn and resistant soybean reduces the nematode populations greatly; however, the populations usually resurge quickly in a subsequent single year

of cotton production (Robinson *et al.*, 2007). Use of chemicals can possibly lead to better management, however, the use of nematicides is no longer a viable management option because of economic and environmental reasons (Agudelo *et al.*, 2005; Blessitt *et al.*, 2012).

The success of management of a disease is based on understanding of variability in a pathogen (Werlemark *et al.*, 2006; Silva *et al.*, 2008). A few studies have reported the existence of variability in geographic populations of reniform nematode. A report from the 1960s indicated that Louisiana populations of *R. reniformis* were physiologically different from other reniform nematode populations suggesting the existence of races (Birchfield, 1962). Subsequent reports demonstrated physiological variation in reproduction and pathogenicity of geographic populations of reniform nematode (McGawley and Overstreet, 1995; McGawley *et al.*, 2010, 2011). Studies have also been conducted to determine the amount of genetic variability in reniform nematode populations. While Agudelo *et al.* (2005) did not find any obvious genetic variability in reniform nematode populations collected from ten states in the United States, other genetic studies reported variability in geographic populations of reniform nematode (Tilahun *et al.*, 2008, Arias *et al.*, 2009; Leach *et al.*, 2012). A better understanding of variability in populations of reniform nematode can help scientists develop a better management strategy. The main objective of this research was to determine whether or not there was reproductive and pathological variation in populations of *R. reniformis* endemic in Louisiana.

2.2 Materials and methods

2.2.1 General procedure

Isolates of reniform nematode were collected from West Carroll (WC), Rapides (RAP), Morehouse (MOR), and Tensas (TEN) parishes in Louisiana, confirmed morphometrically as *R. reniformis*, and used to establish single-egg mass cultures. Axenic cultures were maintained

under greenhouse conditions on tomato (*Solanum lycopersicum* L. cultivar Rutgers PS, Seedway; Hall, New York 14463). Reniform nematode isolates from four parishes were employed in greenhouse and microplot studies with the most widely planted upland cotton cultivars Phytogen 499 WRF, Deltapine 1133 B2RF, and Phytogen 333 WRF (Anonymous, 2015). The cultivars Phytogen 499 WRF, Deltapine 1133 B2RF, and Phytogen 333 WRF hereafter will be abbreviated as PHY499, DP1133, and PHY333, respectively. Exact details of greenhouse and microplot studies are presented below under the appropriate subheadings.

A soil mixture consisting of three parts Commerce silt loam soil (fine-silty, mixed, superactive, nonacid, thermic Fluvaquentic Endoaquepts) and one part sand, and pots, unless stated otherwise, used in all experiments was steam sterilized for 5 hours at 135°C prior to use. In each test, two cotton seeds were planted to a depth of 2.5 cm and thinned to one per pot after germination. Soils were infested by pipetting aqueous suspensions of vermiform individuals of *R. reniformis* into three depressions arranged into a triangular pattern, 0.5-cm diam. × 5- to 7.5-cm deep, surrounding a 7-d-old seedling. The infestation level for the microplot experiments was 50,000 vermiform life stages per microplot. Similarly, the infestation level for the greenhouse experiments was 4,000 vermiform life stages per pot, which is equivalent to the number of vermiform life stages per gram of soil in microplot trials. Half of the inoculum was added to soil in microplots at 10 days after planting and the remainder at 21 days. Standard fertilization, weeding and insect management practices were employed in all trials.

In all cases, nematodes were extracted from a 250 cm³ subsample of soil from each pot and processed using a semi-automatic elutriator (Byrd *et al.*, 1976) and the centrifugal/sugar flotation technique (Jenkins, 1964). Vermiform life-stages were enumerated using a dissecting microscope at 40x magnification. Eggs were extracted from whole root systems in greenhouse

experiments by agitating root in 0.6% NaOCl for 10 min to dislodge eggs from egg masses (Hussey and Barker, 1973) and counting at 40x magnification. All plant materials were dried at 30-35 °C for two weeks and weighed.

2.2.2 Microplot studies

Terra cotta pots having top diameters of 35.6 cm were used as microplots. Microplots were placed in depressions in soil so that only the rim was exposed. Each microplot was filled with 20 kg of soil mixture. The entire microplot area was bounded by aluminum Quonset hut skeletal frame which was open at both ends. The skeletal frame was covered with one layer of 6 mm polyethylene to protect plants in microplots from excessive summer rainfall and one layer of 20% reflective foilcloth for optimal sunlight. This cover was equipped with overhead fans and an irrigation system. The entire system allowed for maintenance of near field conditions (McGawley *et al.*, 2010). A total of 75 microplots were established to evaluate 3 widely planted upland cotton cultivars (PHY499, DP1133, and PHY333), 4 isolates of reniform nematode, a non-inoculated control for each cultivar and 5 replications. Establishment of plants, inoculation with nematodes, and processing of plant and nematode materials were the same as that described above. Additional plant data collected in microplot studies included number of harvestable bolls per plant, seed cotton weight, and lint weight.

2.2.3 Greenhouse studies

This study involved six genotypes of cotton: three cultivars (PHY499, DP1133, and PHY333), one susceptible cultivar (Stoneville 4946 GLB2), and two germplasm lines showing moderate to high levels of resistance (MT2468 Ren3 and M713 Ren5). The cotton genotypes Stoneville 4946 GLB2, MT2468 Ren3, and M713 Ren5 hereafter will be abbreviated as ST4946, MT2468, and M713, respectively. Terra cotta pots with a top diameter of 15 cm and containing

1.6 kg of soil mixture were used. A total of 150 pots were established to evaluate the 6 genotypes, 4 isolates of the nematode, a non-inoculated control for each genotype and 5 replications. The experiment was terminated after 60 days and nematode life stages in soil and roots were quantified as described above.

2.2.3 Data analysis

Each experiment employed a factorial treatment structure and was established as a randomized block design with five replications. Each experiment was repeated once. Analysis of variance was conducted using test as a fixed effect and there were no significant trial by treatment interactions in any of the tests described herein. Therefore data from all like trials was combined for analysis. Data were examined by analysis of variance (ANOVA) for a factorial design using the “Fit Y by X” module of SAS JMP Pro, version 13.0 (SAS Institute; Cary, NC). Means of data were separated by Fisher’s LSD at $P \leq 0.05$.

2.3 Results

2.3.1 Microplot studies

Data from two microplot trials were combined for analysis because of the absence of year by treatment interactions. Reniform nematode isolates, across all cotton cultivars, produced significant differences in reproduction and effect on plant dry weight, seed cotton weight, and lint weight (Table 1). Significant interactions were not observed between cotton cultivars and reniform nematode isolates for any of the growth parameter measured. Statistical main effects of the 4 isolates of the nematode across the 3 cultivars are presented in Table 2. Population density at harvest ranged approximately from 288 to 415 thousand individuals per 500 cm³ of soil, representing reproductive values (number of vermiform stages per microplot divided by the infestation level) of 230.2 to 331.8.

Table 1. Main and interaction effects (*P* values) of four isolates of *Rotylenchulus reniformis* endemic in Louisiana on three cotton cultivars in a microplot environment^x.

Source	DF	Vermiform life stages	Plant weight	Number of bolls	Seed cotton weight	Lint weight
Cultivar			0.0025*	0.0398*	0.0005**	0.0004**
(C) ^y	2	0.8956				
Isolate			<0.0001**	<0.0001**	<0.0001**	<0.0001*
(I) ^z	3	<0.0001**				
C × I	6	0.7831	0.5983	0.2375	0.7521	0.7732

^x Data were combined over two full season trials and are means of ten replications. Plant material was dried at 30-35 °C. Data were analyzed as a 2 × 3 factorial with ANOVA ($P \leq 0.05$); * and ** indicate *P* values significant at the 0.05% and 0.01% levels, respectively.

^y Cultivars were Phytogen 499 WRF, Deltapine 1133 B2RF, and Phytogen 333 WRF that were recommended for use in Louisiana in 2015.

^z Isolates were each derived from a single-egg mass from roots of cotton from West Carroll, Rapides, Morehouse, and Tensas parishes in Louisiana.

Across all cotton cultivars, reproduction of the MOR isolate was significantly greater than that by the other 3 isolates. Reproduction by the RAP was least while TEN and WC had intermediate reproduction. The isolate from MOR parish significantly reduced plant dry weight, number of bolls, seed cotton weight, and lint weight compared to those of the non-inoculated control and other three nematode isolates. Cotton plant dry weight of inoculated plants ranged from 138g to 278g compared to 305.6g for the non-inoculated control. MOR, WC, TEN, and RAP isolates lowered plant dry weight of cotton by 55%, 25%, 21%, and 9%, respectively. The RAP isolate produced the least reduction in plant dry weight, number of bolls, seed cotton weight, and lint weight, although not always significantly different from the control.

Statistical main effects of the 3 varieties of cotton across the 4 isolates of the nematode are presented in Table 3. Across all 4 isolates of the nematode, dry weight for DP1133 was reduced significantly more than were those for the 2 Phytogen cultivars. Results for numbers of

Table 2. Main effect of isolate of *Rotylenchulus reniformis* on vermiform life stages, plant weight, number of bolls, seed cotton, and lint weights across three cultivars of cotton in a full season microplot environment^w.

Isolate ^x	Vermiform life stages (1000's) per 500 cm ³ of soil ^y	Reproduction value ^z	Plant weight (g)	Number of bolls	Seed cotton weight (g)	Lint weight (g)
WC	337.3 b	269.8	228.4 b	16.5 b	82.6 b	35.6 b
RAP	287.8 c	230.2	278 ab	23.9 a	99.7 b	45.2 ab
MOR	414.8 a	331.8	138 c	9.5 c	48.5 c	19.7 c
TEN	333.3 bc	266.6	241.9 b	17.3 b	89.5 b	39.6 ab
Control	0	0	305.6 a	27.7 a	117.4 a	55.5 a

^wData were combined over two full season trials and are means of ten replications. Plant material was dried at 30-35 °C. Cultivars of cotton were PhytoGen 499 WRF, Deltapine 1133 B2RF, and PhytoGen 333 WRF.

^xReniform nematode isolates were each derived from single egg masses isolated from roots of soybean from West Carroll (WC), Rapides (RAP), Morehouse (MOR), and Tensas (TEN) parishes in Louisiana.

^yData were analyzed with ANOVA and Fisher's LSD test ($P \leq 0.05$). Within columns, means followed by a common letter are not significantly different.

^zReproduction values were calculated by dividing the estimated number of vermiform stages per microplot (20 kg of soil) by the infestation level of 50,000 vermiform life stages.

bolls, and seed and lint weights were similar, except for PHY499 where numbers of bolls were not significantly different, averaging 15.7.

2.3.2 Greenhouse studies

Because there was no significant trial by treatment interactions, data from the two greenhouse trials were also combined. For almost all parameters, there were highly significant genotype and isolate main effects as well as genotype by isolate interactions (Table 4). Genotype

Table 3. Main effect of three cultivars of cotton on plant weight, number of bolls, seed cotton, and lint weights across four isolates of *Rotylenchulus reniformis* in a full season microplot environment^x.

Cultivar ^y	Plant weight (g) ^z	Number of bolls	Seed cotton weight (g)	Lint Weight (g)
Phytogen 499 WRF	268.7 a	20.2 ab	100.9 a	45.3 a
Deltapine 1133 B2RF	196.7 b	15.7 b	64.4 b	28.2 b
Phytogen 333 WRF	251.4 a	21.1 a	98.2 a	44.3 a

^xData were combined over two full season trials and are means of ten replications. Plant material was dried at 30-35 °C.

^yCultivars were recommended for use in Louisiana in 2015.

^zData were analyzed with ANOVA and Fisher's LSD test ($P \leq 0.05$). Within columns, means followed by a common letter are not significantly different.

Table 4. Main and interaction effects (P values) of four isolates of *Rotylenchulus reniformis* endemic in Louisiana on six genotypes of cotton in a greenhouse environment^x.

Source	DF	Vermiform life stages	Eggs/root system	Root weight	Shoot weight	Plant weight
Genotype (G) ^y	5	<0.0001**	<0.0001**	0.0027*	<0.0001**	<0.0001**
Isolate (I) ^z	4	<0.0001**	<0.0001**	<.0001**	<0.0001**	<0.0001**
G × I	20	<0.0001**	<0.0001**	0.5167	0.0001**	0.0002**

^xData were combined over two 60-day trials and are means of ten replications. Plant material was dried at 30-35 °C. Data were analyzed as a 5 × 4 factorial with ANOVA ($P \leq 0.05$); * and ** indicate P values significant at the 0.05% and 0.01% level, respectively.

^yGenotypes were the cultivars Phytogen 499 WRF, Deltapine 1133 B2RF, Phytogen 333 WRF, and Stoneville 4946 GLB2 and the germplasm germplasm lines MT2468 Ren3, and M713 Ren5.

^zIsolates were each derived from a single-egg mass from roots of cotton from West Carroll, Rapides, Morehouse, and Tensas parishes in Louisiana.

main effects on nematode vermiform stages and eggs as well as shoot and plant dry weights were significant at the 1% level and root dry weight was significant at the 5% level. Isolate main

Table 5. Main effect of isolates of *Rotylenchulus reniformis* on vermiform life stages, eggs per root system, root, shoot, and plant weights across six genotypes of cotton in a greenhouse environment^w.

Isolate ^x	Vermiform life stages (1000's) per 500 cm ³ of soil ^y	Reproduction value ^z	Eggs/root system	Root weight (g)	Shoot weight (g)	Plant weight (g)
WC	11.4 b	9.1	4484 a	2.0 b	7.5 b	9.5 c
RAP	9.8 b	7.9	4093 a	2.2 b	8.2 b	10.3 b
MOR	14.4 a	10.7	4713 a	1.9 b	6.6 c	8.6 d
TEN	11.1 b	8.8	4228 a	2.1 b	7.8 b	9.9 bc
Control	0 c	0	0 b	2.7 a	9.1 a	11.7 a

^wData were combined over two 60-day trials and are means of ten replications. Plant material was dried at 30-35 °C. Genotypes were the cultivars Phytogen 499 WRF, Deltapine 1133 B2RF, Phytogen 333 WRF, and Stoneville 4946 GLB2 and the germplasm lines MT2468 Ren3, and M713 Ren5.

^xReniform nematode isolates were each derived from single egg masses isolated from roots of soybean from West Carroll (WC), Rapides (RAP), Morehouse (MOR) and Tensas (TEN) parishes in Louisiana.

^yData were analyzed with ANOVA and Fisher's LSD test ($P \leq 0.05$). Within columns, means followed by a common letter are not significantly different.

^zReproduction values were calculated by dividing the estimated number of vermiform stages per pot (1.6 kg of soil) by the infestation level of 4,000 vermiform life stages.

effects were significant at the 1% across both nematode and plant parameters. Genotype by isolate interactions were significant at the 1% levels for both nematode and plant values except root dry weight where they were not significant.

Statistical main effects of the six genotypes of cotton across the 4 isolates of the nematode in the greenhouse environment are presented in Table 5. Reniform nematode

population density in soil ranged from approximately 10 thousand to 14 thousand individuals per 500 cm³ of soil with corresponding reproductive values of 7.9 to 10.7. The numbers of eggs per root system was similar among the isolates and averaged 4 to 5 thousand per root system. Across all cultivars, all isolates of the nematode caused significant reductions in root weight compared with controls, but there were no differences among the isolates. Results for weights of shoots was similar except that the MOR isolate caused greater reductions than the other 3 isolates. The isolate of *R. reniformis* from MOR parish also caused a reduction in final plant weight which was greater than that caused by the other 3 isolates, 8.6 g compared with 9.5 g for the WC isolate, 10.3 g for the RAP isolate, and 9.9 g for the TEN isolate.

The two PhytoGen cultivars, along with DP1133 and ST4946 supported the highest, but not significantly different, numbers of nematodes which ranged from 9,688 to 13,022 individuals per 500 cm³ of soil (Table 6). Significantly fewer numbers, 7,315, were recovered for MT2468 Ren3. Vermiform stages per 500 cm³ of soil averaged 4,573 for M713 Ren5 and were significantly less than the averages across the 4 isolates for the other genotypes.

Egg production by the nematode, across isolates, was similar and not significantly different for PHY499, DP1133, PHY333, and ST4946 (Table 6). Respectively, eggs per root system averaged 4,667, 4,534, 4,637 and 5,181. Significantly fewer, 1,360 and 642, were collected from roots of MT2468 Ren3 and M713 Ren5.

There were highly significant genotype by isolate interactions which influenced both soil and root stages of the nematode (Table 4). Individual treatment means illustrating soil population levels of the nematode across the 6 genotype / 4 isolate combinations are presented in Figure 1. Nematode numbers associated with PHY499 ranged from 12,795 for the TEN isolate to 15,520 per 500 cm³ of soil for the MOR isolate with intermediate values of 13,232 for WC and 14,454

Table 6. Main effect of genotypes of cotton on vermiform life stages, eggs per root system, root, shoot, and plant weights across four isolates of *Rotylenchulus reniformis* in a greenhouse environment^x.

Genotype ^y	Vermiform life stages ^z	Eggs/root system	Root weight (g)	Shoot weight (g)	Plant weight (g)
Phytogen 499 WRF	11200 a	4667 a	2.2 abc	8.0 b	10.3 b
Deltapine 1133 B2RF	9688 ab	4534 a	2.4 a	9.2 a	11.6 a
Phytogen 333 WRF	10174 a	4637 a	2.3 ab	8.2 b	10.4 b
Stoneville 4946 GLB2	13022 a	5181 a	2.1 bc	7.9 b	9.9 b
MT2468 Ren3	7315 b	1360 b	2.0 c	6.6 c	8.6 c
M713 Ren5	4573 c	642 b	2.1 bc	7.1 c	9.1 c

^xData were combined over two full season trials and are means of ten replications. Plant material was dried at 30-35 °C.

^yGenotypes were the cultivars Phytogen 499 WRF, Deltapine 1133 B2RF, Phytogen 333 WRF, and Stoneville 4946 GLB2 and the germplasm lines MT2468 Ren3, and M713 Ren5.

^zData were analyzed with ANOVA and Fisher's LSD test ($P \leq 0.05$). Within columns, means followed by a common letter are not significantly different.

for RAP and no significant differences in numbers among the 4 isolates. An average soil population level of 15,592 individuals per 500 cm³ of soil was estimated for the WC isolate on DP1133 and this was significantly greater than the 8,907 and 11,829 for the RAP and MOR isolates but not the 12,112 per 500 cm³ with the TEN isolate. With the genotype PHY333, nematode populations in soil averaged 13,728, 10,520, 14,768 and 11,856 nematodes per 500 cm³'s for the WC, RAP, MOR and TEN isolates, respectively. These population levels did not differ significantly from one another. For ST4946 there were significantly greater numbers of the MOR and TEN isolates (20,856 and 17,728, respectively) in soil than for the other 2 isolates with averages of 8,672 for WC and 12,056 for RAP. There were no significant differences in




Figure 1. Vermiform life stages of *Rotylenchulus reniformis* per 500 cc of soil, after 60 days in a greenhouse environment from cotton genotypes PhytoGen 499 WRF (PHY499), Deltapine 1133 B2RF (DP1133), PhytoGen 333 WRF (PHY333), Stoneville GLB2 (ST4946), MT2468 Ren3, and M713 Ren5. Data are means of 10 replications averaged over two trials. Bars with common letters are not significantly different based on Fisher's LSD test ($P < 0.05$)

population levels of the nematode in soil for any of the isolates with the genotype MT2468. Numbers per 500 cm³ of soil ranged from 6,776 for the TEN isolate to 10,192 for the MOR isolate. Overall, the lowest populations of the nematode in soil were found with the genotype M713. Nematode numbers for the RAP and TEN isolates were similar and not significantly different averaging 3,488 and 4,992, respectively. Greater population levels, 7,016 for the WC isolate and 7,369 for MOR were associated with M713.

Individual treatment means showing numbers of eggs of the nematode recovered per root system across the 6 genotype / 4 isolate combinations are presented in Figure 2. The histograms illustrate clearly that significantly greater numbers of eggs were produced by all isolates of the nematode on the 4 named cultivars, PHY499, DP1133, PHY333 and ST4946, than on the




Figure 2. Egg stages of *Rotylenchulus reniformis* from whole root systems of cotton genotypes PhytoGen 499 WRF (PHY499), Deltapine 1133 B2RF (DP1133), PhytoGen 333 WRF (PHY333), Stoneville 4946 GLB2 (ST4946), MT2468 Ren3, and M713 Ren5 after 60 days in a greenhouse environment. Data are means of 10 replications averaged over two trials. Bars with common letters are not significantly different based on Fisher's LSD test ($P < 0.05$).

germplasm lines MT2468REN3 and M713REN5. For PHY499, there were significant differences in egg numbers per root system across the 4 isolates of the nematode where RAP had significantly higher reproduction than MOR and TEN, but not WC with means averaging 5,816, 4,664, 6,432 and 6,424 for WC, RAP, MOR and TEN, respectively. For DP1133, there were also no significant differences in numbers of eggs with very similar numbers estimated for each isolate; 5,896 for WC, 5,224 for RAP, 5,936 for MOR and 5,616 for TEN. Numbers of eggs per root system of PHY333 were not significantly different for the WC, RAP and MOR isolates, respectively averaging 6,152, 5,984 and 6,192. Numbers of eggs per root system from PHY333 averaged 4,856 for the TEN isolate of the nematode and this was significantly lower than that numbers from the MOR and WC isolates but not the RAP isolate. Numbers of eggs for each of

the 4 isolates of the nematode were very similar for ST4946 and not significantly different with means averaging 6,576 for WC, 6,608 for RAP, 6,600 for MOR and 6,120 for TEN. Mean numbers of eggs from roots of MT2468 averaged 1,760 for the WC isolate of *R. reniformis*, 1,472 for RAP, 1,888 for MOR and 1,680 for TEN and there were no significant differences in these averages. Similarly, there were no significant differences in numbers of eggs recovered from M713 for each of the isolates of the nematode. Averages for eggs per root system were 704, 608, 1,228 and 672 for WC, RAP, MOR and TEN, respectively.

There were also highly significant genotype by isolate interactions indicated in Table 4 that influenced the cotton shoot and plant dry weights. Inspection of individual treatment means for both plant parameters reveal a similar pattern and, therefore, only those for plant weight are presented as Figure 3. Overall, the figure illustrates clearly that the isolate of the nematode from MOR parish was the most pathogenic and it was significantly more so on the germplasm lines MT2468 and M713. Relative to the non-inoculated control at 60 days after inoculation, weights of PHY499 plants were reduced significantly by the reniform nematode isolates from WC, MOR and TEN parishes but not by the one from RAP. Control plant weight averaged 11.9 g, those for WC, MOR and TEN were 10.1, 8.8 and 9.6 g, respectively and that for RAP was 10.9 g. Three of the 4 isolates, WC, RAP and TEN, did not cause significant reductions in plant weight for DP1133 when compared with the average for the control. Respectively, these plant dry weights averaged 11.2, 12.2, 11.6 and 12.6 g. The WC and TEN isolates caused significant damage to PHY333 and the RAP and MOR isolates did not. The mean plant weight for non-inoculated PHY333 was 11.7 g. Weights for PHY333 inoculated with isolates from WC and TEN were 9.2 and 9.9 g, respectively and those for RAP and MOR were 10.4 and 10.9. With ST4946 the isolates from RAP and MOR reduced weights of plants significantly below the 11.6 g value of




Figure 3. Effect of four isolates of *Rotylenchulus reniformis* on plant dry weight of cotton genotypes PhytoGen 499 WRF (PHY499), Deltapine 1133 B2RF (DP1133), PhytoGen 333 WRF (PHY333), Stoneville 4946 GLB2 (ST4946), MT2468 Ren3, and M713Ren5. Nematode isolates each were derived from a single-egg mass isolated from West Carroll, Rapides, Morehouse, and Tensas Parishes of Louisiana. Plant material was dried at 30-35 °C. Data were combined over two 60-day trials and are means of ten replications. Data were analyzed with ANOVA and Fisher's LSD test ($P \leq 0.05$). Means followed by a common letter are not significantly different.

the non-inoculated control. Isolates from WC and TEN did not reduce weights of ST4946 significantly. Weights were 8.6 g for RAP, 9.2 g for MOR, and 10.3 g for WC and TEN. For both MT2468 and M713, three of the four isolates of the nematode reduced weights of plants significantly relative to the control. Plant weights for non-inoculated MT2468 and M713 each averaged 10.8 g. For MT2468 significant reductions in plant weight were caused by WC, MOR, and TEN isolates with mean plant weights of 7.7, 5.8, and 8.0 g, respectively. Plant weight for the RAP isolate was 10.2 g and not significantly different from the control. For M713 the isolates which caused significant plant damage were from WC, RAP, and MOR parishes. Respectively,

plant weights were 8.2, 9.6, and 6.3 g and the isolate which did not elicit significant plant damage was from TEN parish and the final plant weight averaged 10.0 g.

2.4 Discussion

Nematologists have documented the existence of reproductive, pathogenic, and/or genetic variability in a range of plant parasitic nematodes including burrowing nematode (Ducharme and Birchfield, 1956; Huettel and Yaegashi, 1988), stem and bulb nematode (Seinhorst, 1957), soybean cyst nematode (Riggs *et al.*, 1981; Niblack *et al.*, 2002), and root-knot nematode (Barker *et al.*, 1985; Noe, 1992; Van der Beek *et al.*, 1999; Khanal *et al.*, 2016). The existence of morphological, physiological, and/or genetic variability among geographic isolates of *R. reniformis* have been proposed by some studies as well. Dasgupta and Seshadri (1971) designated two races (Race A, Race B) of *R. reniformis* based on their differential reproduction in castor, cotton, and cowpea. A study published in 1983 in Japanese and translated in English by Nakasono in 2004 designated 3 distinct biological types (male-numerous, male-rare, and male-absent) of *R. reniformis* that originated from Japan and the United States. McGawley and Overstreet (1995) studied 17 populations of reniform nematode collected from Louisiana, Arkansas, Hawaii, Mississippi, and Texas in greenhouse and laboratory tests and found variation among populations with respect to reproduction on and/or damage to cotton and soybean. A study conducted by Agudelo *et al.* (2005) on selected cotton and soybean cultivars involving 13 amphimictic populations of reniform nematode collected from major cotton growing area in the United States showed that considerable variation in reproduction and morphology exists within and among the geographic populations. Agudelo *et al.* (2005) further evaluated the ribosomal internal transcribed spacer region-1 (ITS1) of populations of reniform nematodes from the United States as well as from Brazil, Colombia, Honduras, and Japan. They found that all the

populations, except one from Japan, did not differ genetically for the studied nuclear region. The population from Japan, that was parthenogenetic, showed a considerable amount of nucleotide variation (41/348 bp) suggesting that a difference in genotypic make up can introduce considerable variation into a population. They suggested further that other molecular markers such as amplified fragment length polymorphism and microsatellites would be useful in assessing variation in nematode populations. In contrast to the results from Agudelo *et al.* (2005), Tilahun *et al.* (2008) found significant amount of variation in ITS1 and 18S ribosomal DNA of seven reniform nematode populations in Alabama. Such contrasting results have created confusion about the suitability of ribosomal DNA for assessment of genetic variability in this nematode.

As the genetic variability studies specifically focused on ITS and 18s rDNA has been elusive, Arias *et al.* (2009) readily distinguished reniform nematode populations from Texas, Louisiana, Mississippi, and Georgia using microsatellite markers. Furthermore, a greenhouse experiment by Arias *et al.* (2009) supported the notion of variability in geographic isolates. Studies of the variability of geographic isolates described heretofore were short duration greenhouse or lab studies that did not always employ populations derived from single-egg mass cultures, and the experiment may or may not have been repeated. In contrast to the methodology employed in previous research, McGawley *et al.* (2010, 2011) conducted microplot tests involving cotton and soybean to assess reproduction and pathogenicity of reniform nematode populations derived from single-egg mass cultures collected from Alabama, Arkansas, Hawaii, Louisiana, Mississippi, and Texas. They collected data from two full-season microplot trials and found significant differences among isolates of reniform nematode in both reproduction and pathogenicity. Microplot trials conducted by McGawley *et al.* (2010, 2011) reported that

reniform nematode inoculated cotton and soybean cultivars, respectively, sustained 38.6% and 27.9% plant dry weight reduction compared to those of the non-inoculated controls. Further research including several isolates of reniform nematode endemic in Louisiana are necessary to support that reniform nematode is more damaging to cotton than soybean. Greenhouse study conducted by Bhandari *et al.* (2015) reported significant variability in reproduction and pathogenicity of Louisiana populations, although not derived from single egg mass cultures, of reniform nematode on susceptible cotton genotypes and resistant germplasm lines.

Research detailed in this report provides an indication of the amount of variation in endemic populations of reniform nematode from cotton growing regions in Louisiana. Over the course of this research, two microplot and two greenhouse experiments were conducted to determine the pathogenicity and reproduction of four endemic populations of reniform nematode on cotton genotypes. Data obtained from both greenhouse and microplot experiments demonstrated that the MOR and RAP isolates caused the greatest and the least damage, respectively. Furthermore, least reproduction and lower pathogenicity of reniform nematode isolates, especially MOR and RAP, was evident on the germplasm lines rather than on commercial cultivars (Figure 1-4).

This research is the first report that employs a series of microplot and greenhouse experiments to demonstrate reproductive and pathogenic variation among populations of *R. reniformis* endemic to Louisiana. Evidence strongly suggesting the existence of virulence phenotypes in endemic populations makes it essential for breeding programs aimed at developing reniform nematode resistant cultivars employ as many isolates as possible in order to produce durable sources of resistance.

Parallel research conducted by a fellow nematology student here at LSU, Mr. Manjula Kularathna, employs the same populations of reniform nematode, but uses soybean as the host plant. Data from his research also shows differences in reproduction and pathology of the nematode on soybean. A major difference in results from these two parallel lines of research involve the level of reproduction of MOR isolate on two different hosts. Across soybean genotypes, the MOR isolate exhibited the lowest level of reproduction, but caused the greatest amount of damage. Conversely, with cotton, the MOR isolate exhibited the greatest level of reproduction and caused the greatest level of damage. Across all cotton and soybean genotypes, respectively, MOR isolate reduced plant dry weight by 54.8% and 29.8% relative to those of the non-inoculated controls. This difference in pathogenicity of MOR isolate on cotton and soybean is possibly a function of host.

CHAPTER 3. SINGLE NUCLEOTIDE POLYMORPHISM ANALYSIS PROVIDES AN INSIGHT ON GENETIC VARIABILITY IN *ROTYLENCHULUS RENIFORMIS*

3.1 Introduction

The reniform nematode (*Rotylenchulus reniformis* Linford and Oliveira) has established itself as the major plant parasitic nematode of cotton and soybean in the United States (Stetina and Young, 2006; Robinson, 2007; Arias *et al.*, 2009; Leach *et al.*, 2012; Allen *et al.*, 2017). Approximately 12% of cotton and 5% of soybean yields are lost to reniform nematode in the United States cotton belt with Louisiana, Alabama, and Mississippi sustaining the greatest damage (Lawrence *et al.*, 2017; Allen *et al.*, 2017).

Current practice of reniform nematode management is based primarily on use of chemicals or crop rotation. Chemical control is not always the preferred method of reniform nematode management because of health and environmental issues (Agudelo *et al.*, 2005). Selection of a suitable crop for rotation can help reduce losses caused by the nematode, however, reniform nematode populations resurge quickly to a damaging level in a single crop growing season when a susceptible crop is planted (Robinson *et al.*, 2007). Use of host plant resistance has been the most desirable, but least available method of reniform nematode management in cotton. Unfortunately, reniform nematode resistant cotton cultivars are not available. Some cotton breeding lines and soybean cultivars showing moderate to high level of resistance to reniform nematode are available, however, their yield performance is not always similar when tested across different geographic locations (Yik and Birchfield, 1984; Robinson *et al.*, 1997; Robinson *et al.*, 2004; Weaver *et al.*, 2007). It is well known that durable host plant resistance is mainly dependent on amount of variability present in a pathogen (Riggs *et al.*, 1981; Noe, 1992; Van der Beek *et al.*, 1999; Niblack *et al.*, 2002). The inconsistency in suppression of

reproduction of reniform nematode in different geographic locations by some available resistant cultivars/breeding lines have been reported (Robinson *et al.*, 1997; Robinson *et al.*, 2004; Yik and Birchfield, 1984; Weaver *et al.*, 2007; Wallace *et al.*, 2013; Bell *et al.*, 2015). This inconsistency could have been caused by the existence of physiological and genetic variability in geographic isolates of reniform nematode. Utilization of molecular techniques to better understand the genetic variability in reniform nematode populations would be helpful in developing durable reniform nematode resistant cultivars.

Molecular techniques such as restriction fragment length polymorphism (RFLP), random amplified polymorphic DNA (RAPD), amplified fragment length polymorphism (AFLP), inter-simple sequence repeats (ISSR), simple sequence repeat (SSR), and single nucleotide polymorphisms (SNP) have been used in the last three decades to study genetic variations and characterization of population of many organisms (Grover and Sharma, 2016). The first generation of molecular markers such as RFLPs are not currently much used for genetic variability assays because they are complex, costly, and identify lower rates of polymorphism (Gao *et al.*, 2016; Yang *et al.*, 2017). SSRs are often considered the second generation of genomic markers and are useful in determining a higher polymorphism rate (Gao *et al.*, 2016). However, SSR markers are often considered cumbersome to use in high throughput genotyping protocols and they may not be widely and evenly distributed in the genome (Salem *et al.*, 2012). The third generation of markers (Gao *et al.*, 2016), SNPs, are superior in studying genetic variability as they are abundant and widely distributed in the genome (Salem *et al.*, 2012). Advent of next generation sequencing (NGS) technology has enabled more efficient study of genetic variation in an organism using SNP analysis (Gao *et al.*, 2016). The utility of SNPs allows genomic sequences of an organism to be compared in a rapid, reliable, and highly

efficient way (Graves *et al.*, 2016; Yang *et al.*, 2017). SNP analyses have been useful in discriminating species of many organisms including fungi, bacteria, virus, nematode, plants, and animals (Faga *et al.*, 2001; Morais *et al.*, 2006; Rattei, *et al.*, 2007; Samson-Himmelstjerna *et al.*, 2007; Figueiredo *et al.*, 2013; Lu *et al.*, 2013; Ojeda *et al.*, 2014; Gao *et al.*, 2016; Linlokken *et al.*, 2017; Yang *et al.*, 2017).

The existence of morphometric, physiological, and genetic variabilities within *R. reniformis* populations have been documented from different parts of the world including Japan (Nakasono, 2004), India (Dasgupta and Seshadri, 1971), Africa (Germani, 1978), Brazil (Rosa *et al.*, 2003; Soares *et al.*, 2003, 2004), and the United States (McGawley and Overstreet, 1995; Agudelo *et al.*, 2005; Tilahun *et al.* 2008, Arias *et al.*, 2009, McGawley *et al.*, 2010; McGawley *et al.*, 2011). The variability among geographic populations of *R. reniformis*, particularly genetic variability, has not always been obvious. A study conducted by Agudelo *et al.* (2005) showed no variation in the first internal transcribed spacer (ITS1) region of reniform nematode collected from ten states in the United States. In contrast to the report by Agudelo *et al.* (2005), Tilahun *et al.* (2008) found fairly substantial variation in ITS1 as well as in the 18S region of reniform nematode populations from Alabama. Regarding the lack of correlation between phenotypic and genotypic variation, Agudelo *et al.* (2005) suggested that use of microsatellite markers could provide a more reliable way to evaluate populations. Studies conducted by Arias *et al.* (2009), and Leach *et al.* (2012) using microsatellite markers reported genetic variability in geographic isolates of reniform nematode. With the advent of NGS technology, it is possible to analyze whole genomic DNA of reniform nematode in much more detail using a robust molecular assay such as SNPs. Analysis of SNP data would be useful in determining genetic variability within isolates and among geographic populations of reniform nematode. To date, there have not been

any published reports of SNP molecular markers specific to reniform nematodes. The major objective of this research was to identify SNP molecular markers and evaluate their use in determining genetic variability among reniform nematode isolates.

3.2 Methodology

3.2.1 Isolation and extraction of nematode

A total of 13 geographic populations of reniform nematode, two from Louisiana, six from Mississippi, three from Arkansas, and one each from Hawaii, and Alabama, were used to evaluate putative SNPs and measure genetic variability (Table 7). Each population was derived from single egg mass and maintained on tomato (*Solanum lycopersicum* L. cultivar Rutgers PS, Seedway; Hall, New York 14463) in a greenhouse (Table 7). From each population, 300 to 400 gravid females were excised from tomato roots with the help of stereoscopic microscope, sterilized needles, and tweezers. The excised gravid females from each population were placed in distilled water in petri plates and subsequently transferred to labelled 2 ml Eppendorf tubes.

3.2.2 Genomic DNA extraction from nematode

From nematode samples DNA was extracted using a Maxwell 16 (Promega, Madison, WI, USA) automated DNA isolation machine. To Eppendorf tubes containing the nematodes in a water solution, 500 μ l of CTAB buffer, 30 μ l of Proteinase K (20 mg/ μ l), 2 μ l of RNase A (10 mg/ml, catalog No. EN0531), and 2 μ l of lysozyme (500 ng/ μ l) was added. The tubes were briefly vortexed vigorously. The tubes were then incubated for 2 hours at 60 °C with gentle shaking at 350 rpm. The tubes were then vortexed for 5 seconds to mix the solution. The solution obtained was processed using Maxwell 16 FFS Nucleic Acid Extraction System, Custom (Catalog No. X9431). The supernatant, which is essentially genomic DNA, obtained at the end of process was transferred to labelled 1.5 ml Eppendorf tubes. The genomic DNA was quantified on

Table 7. Sample ID, origin, population type, and source of reniform nematode populations used for SNP analysis.

Sample ID	Origin of sample ^x	Isolate ^y	Source ^z
LA1	Rapides, Louisiana	Single egg mass	Nematode advisory service, LSU
LA2	Morehouse, Louisiana	Single egg mass	Nematode advisory service, LSU
MS1	Humphreys, Mississippi	Single egg mass	Salliana Stetina
MS2	Holmes, Mississippi	Single egg mass	Salliana Stetina
MS3	Holmes, Mississippi	Single egg mass	Salliana Stetina
MS4	Stark, Mississippi	Single egg mass	Gary Lawrence
MS5	Elizabeth, Mississippi	Single egg mass	Salliana Stetina
MS6	Washington, Mississippi	Single egg mass	Salliana Stetina
AR1	Kibler, Arkansas	Single egg mass	Robert Robbins
AR2	Hawkins, Arkansas	Single egg mass	Robert Robbins
AR3	Fayetteville, Arkansas	Single egg mass	Robert Robbins
HI1	Oahu, Hawaii	Single egg mass	Brent Sipes
AL1	Auburn, Alabama	Single egg mass	Kathy Lawrence

^xLocation from where reniform nematode populations were originally collected.

^yReniform nematode populations were maintained on tomato in greenhouse either as single-egg mass or mixed populations.

^zPersons or lab that originally collected and supplied the nematode samples.

Synergy H1 (BioTek®, Winooski, VT, USA) microplate spectrophotometer at 260 nm UV absorption. DNA from the single egg reniform nematode samples were isolated and whole genome amplified.

3.2.3 Whole genome amplification of genomic DNA

In order to conduct multiple SNP analyses, a larger amount of genomic DNA was required. Sufficient amounts of genomic DNA for each reniform nematode population was obtained by the process of whole genome amplification (WGA) using GenomePlex® Complete Whole Genome Amplification kits (Sigma-Aldrich, St. Louis, MO; Cat. No. WGA2) following manufacturer's instructions. The genomic DNA isolations served as the DNA template for whole genome amplification. The procedure consisted of three broad steps: 1) Fragmentation, 2) Library Preparation, and 3) Amplification. Briefly, for Step 1 (Fragmentation): 1 µl of 10X fragmentation buffer and 10 µl of DNA (1 ng/µl) were pipetted in a 200 µl PCR tube. The tube was placed in a PTC-200 thermal cycler (MJ Research, Waltham, MA) at 95 °C for 4 minutes. Immediately the sample was cooled by placing the tube on ice for 2 minutes followed by a brief centrifugation to consolidate the contents. For Step 2 (Library Preparation): to the tube from Step 1, 2 µl of 1X library preparation buffer, 1 µl of library stabilization solution were added and thoroughly vortexed. The tube was consolidated by centrifugation and placed in a PTC-200 thermal cycler (MJ Research, Waltham, MA) at 95 °C for 2 minutes. The sample was cooled again by placing the tube on ice for 2 minutes, then consolidated by centrifugation and returned to ice. A 1 µl of library preparation enzyme was added to the tube, thoroughly vortexed and briefly centrifuged. The tube was placed in a PTC-200 thermal cycler (MJ Research, Waltham, MA) and incubated with following conditions: 16 °C for 20 minutes, 24 °C for 20 minutes, 37 °C for 20 minutes, and 75 °C for 5 minutes. Tubes were removed from the thermal cycler and briefly centrifuged. For Step 3 (Amplification): the 15 µl library sample obtained in Step 2 was used for the subsequent amplification process by adding 7.5 µl of 10X Amplification Master Mix, 47.5 µl of water (molecular biology grade), and 5 µl of WGA DNA polymerase for a total volume of 75

µl. The tube was thoroughly vortexed, briefly centrifuged, and placed in a PTC-200 thermal cycler for amplification. The thermal cycler was set with following conditions: an initial denaturation at 95°C for 3 minutes; followed by 28 cycles of denaturation at 94°C for 15 seconds, and annealing/extension at 65°C for 5 minutes. A 5 µl aliquot of the final product, WGA amplified DNA, was run out on a 1.5% Agarose gel to confirm the procedure was successful. The remaining volume of WGA DNA was purified using a GenElute™ PCR Clean-Up Kit from Sigma-Aldrich (Catalog Number NA1020).

The WGA amplified DNA from each reniform nematode population was quantified on Synergy H1 microplate spectrophotometer and stored at -20°C. In order to obtain enough DNA for the subsequent SNP analyses, all WGA DNA samples were reamplified using the GenomePlex WGA Reamplification Kit (Sigma-Aldrich, Catalog Number WGA3) following manufacturer's instructions. Reamplified DNA samples were purified using the GenElute kits and concentrations determined on the microplate spectrophotometer.

3.2.4 Identification of SNPs

Putative SNPs were derived from reniform genomic DNA analysis in a previous study using nextRAD (Nextera-tagmented Reductivity-Amplified DNA; SNPsaurus, Eugene, OR USA) technology (Dr. Jeffery D. Ray, USDA-ARS, unpublished data). Flanking sequences of 162 putative sequences are shown in Appendix 1 and 2. The 31 putative SNPs shown in Appendix 1 were selected for testing in the current study. SNPs were specifically selected to be at different genomic locations (i.e. on different contigs) as reported for the reniform genome (RREN 1.0) at NCBI (https://www.ncbi.nlm.nih.gov/assembly/GCA_001026735.1/). The flanking sequences of the 31 selected SNPs (Table 8) were sent to LGC Genomics (Teddington, UK) where KASP (kompetitive allele-specific PCR) genotyping assays were designed for each SNP.

Table 8. Summary of SNP ID, contig, sequence position, fluorescence label for each SNP, GC content, and LGC Genomics reference number for the SNP-specific primers used in this research. The SNP and flanking sequences are shown in Appendix 1.

SNP ID	Contig	Sequence Position	Allele FAM ^x	Allele HEX ^y	GC% FAM	GC% HEX	GC% common	LGC Genomics ^z
RREN_4410_3972	4,410	3,972	T	C	54.5	57.1	61.9	1140749440
RREN_1572_36933	1,572	36,933	T	C	45.8	52	59.1	1140749445
RREN_4410_3979	4,410	3,979	A	G	56	59.1	73.7	1140749464
RREN_43396_315	43,396	315	T	G	44	50	59.1	1140749374
RREN_523_19992	523	19,992	A	C	42.3	45.8	48	1140749391
RREN_367_3958	367	3,958	T	G	52.2	54.5	39.3	1140749398
RREN_4834_4618	4,834	4,618	A	G	42.3	45.8	48	1140749415
RREN_5033_5267	5,033	5,267	T	C	42.3	44	59.1	1140749422
RREN_845_36717	845	36,717	A	G	35.7	38.5	48	1140749439
RREN_3215_15723	3,215	15,723	T	C	42.3	44	37.9	1140749446
RREN_1660_513	1,660	513	T	C	30	35.7	33.3	0216484048
RREN_4410_3946	4,410	3,946	A	G	65	68.4	59.1	0216484047
RREN_7711_4758	7,711	4,758	T	C	42.3	45.8	48	0216484024
RREN_514_63176	514	63,176	A	G	54.5	57.1	59.1	0216484023
RREN_925_39379	925	39,379	C	G	33.3	31	48	0216484000
RREN_514_63173	514	63,173	T	G	59.1	65	54.2	0216483999
RREN_91287_201	91,287	201	A	C	44	45.8	33.3	0216483976
RREN_91287_193	91,287	193	T	C	44	45.8	37.9	0216484070
RREN_43396_339	43,396	339	A	C	29	30	59.1	0216484049

(Table continued)

SNP ID	Contig	Sequence	Allele	Allele	GC%	GC%	GC%	LGC
		Position	FAM ^x	HEX ^y	FAM	HEX	common	
RREN_1990_6847	1,990	6,847	A	G	37	44	46.2	0216484046
RREN_20709_1089	20,709	1,089	A	C	33.3	37	37.9	0216484025
RREN_258_12977	258	12,977	A	G	52	52.2	48	0216484022
RREN_269_9935	269	9,935	A	G	24.2	30	46.2	0216484001
RREN_456_104249	456	104,249	T	G	34.5	37	48	0216483998
RREN_43396_325	43,396	325	T	G	37	40.7	59.1	0216483977
RREN_901_49990	901	49,990	A	G	30	31	37.9	0216484069
RREN_1886_12077	1,886	12,077	C	G	37	37	48	0216484050
RREN_16875_158	16,875	158	T	C	24.2	30	73.7	0216484045
RREN_251_23034	251	23,034	T	C	52.2	54.5	50	0216484026
RREN_1895_31360	1,895	31,360	A	T	44	44	33.3	0216484021
RREN_9137_320	9,137	320	T	C	35.7	42.3	37.9	0216484002

^wSNPs were assigned and chosen across the reniform nematode genome so that each SNP is not clustered together with the others.

^xTail of primers were labelled with FAM and HEX which generate specific fluorescence signals that is detected by LightCycler 480 software.

^yPercentage of Guanine and Cytosine in a SNP sequence.

^z LGC Genomics reference number for each SNP-specific assay tested in this research.

3.2.5 KASP genotyping assay

SNPs indicate a single base change in the genome which can be detected by designing primers that amplify the respective base change. For example in the middle of the sequence below the letters in brackets “[T/C]” represent a SNP, where one allele is a “T” and another allele is a “C” and heterozygotes represent a mixture of both alleles:

AAGACGCCCCGGGGGAAGGA[T/C]AGAGGGAATTCCCCTCTCC.

The primers are designed in such a way as to specifically amplify one base or the other, in this case “T” or “C”. KASP assays are designed as dual emission fluorescent reactions where different wavelengths represent one or the other allele (i.e. “T” or “C” in the above case). After PCR amplification of the assay, the specific fluorescent emissions are read on a fluorimeter and analyzed to determine which allele (or both alleles) are present in the sample. In this study, a LightCycler 480 (Rouche Diagnostics Corporation, Indianapolis, IN, USA) real-time PCR machine was used to determine the fluorescent emissions and proprietary Rouche software (LightCycler® 480 Software ver. 1.5.1.62SP3) was used to call the alleles. In the software, one allele was denoted as X and the other as Y based on emission wavelengths. Mixtures of both alleles were denoted as “H” for heterozygotes.

A total of 17 samples, that included 4 no-template controls and the 13 reniform nematode samples described above, were tested. The Amplification Reaction Mix was prepared so that each reaction had 10 µl of 1X KASP Master Mix, and 0.4 µl of 1X KASP Assay Mix (containing the allele specific primers unique to each SNP), and 9.6 µl of WGA DNA (at a concentration of 12.5 ng µl⁻¹). The PCR reactions were assembled in 96-well semi-skirted PCR plates with white wells and clear frames (4ti-0951, 4titude Ltd., Wotton, Surrey UK) using a Janus robot (Perkin Elmer, Shelton, CT). The plate was sealed with QPCR adhesive seals (9095-10055, KBio, Beverly, MA) and placed in a PTC-200 thermocycler. PCR was conducted as follows: an initial denaturation at 94°C for 15 minutes; followed by 10 cycles of denaturation at 94°C for 20 seconds, and annealing/extension at 65°C for 1 minute with a temperature reduction of -0.8 °C per cycle; and subsequent 26 cycles of denaturation at 94°C for 20 seconds, and annealing/extension at 57°C for 1 minute.

3.3 Results

The procedures of WGA and WGA reamplification successfully produced sufficient quantities and quality of DNA for SNP analysis. This technology allows the molecular analysis of minute quantities of DNA derived from individual nematodes, thereby facilitating the analysis of genetic diversity among and within nematode populations. One approach is to examine genetic diversity among samples using molecular markers such as SNPs. However, to date, no SNPs have been reported specifically for reniform nematodes.

As part of an on-going project, 162 putative reniform nematode specific SNPs were identified (Jeffery D. Ray, USDA-ARS, Stoneville, MS; personal communication) but have not been reported elsewhere. However, none of these putative SNPs had been previously evaluated to determine if they actually perform as intended. Therefore a subset of 31 putative SNPs were designed and manufactured to function as dual emission fluorescent KASPTM (kompetitive allele-specific PCR) primers (Table 8, LGC Genomics reference number). These KASP primers enable the bi-allelic scoring of SNPs at specific loci including those in complex genomes. The 31 KASP SNP primers sets were evaluated against 13 reniform nematode isolates.

The 13 isolates of reniform nematode were collected from Louisiana, Mississippi, Arkansas, Hawaii, and Alabama (Table 7). Of the 31 SNPs tested, 13 amplified genomic DNA of reniform nematode isolates from different geographic locations while five SNPs, for the most part, failed to successfully amplify. A summary of results from KASP genotyping assays is presented in Table 9.

In the case of reniform nematode samples from Louisiana, 25 SNPs amplified while six SNPs failed to amplify. Of the 25 positive SNPs, twelve appeared to identify only one allele

while six identified both alleles (i.e. heterozygous DNA). The remaining seven SNPs identified allelic variants (i.e. genetic differences) among the reniform nematode isolates from Louisiana.

For nematode samples from Mississippi, two of the SNPs failed to amplify half of the samples. Of the remaining 29 SNPs, eight were monomorphic (except for MS11 which the LightCycler could not differentiate) while two detected heterozygous loci (except for MS10 which was not amplified). The other 15 SNPs identified genetic differences among the reniform nematode isolates from Mississippi. The remaining three SNPs were not very good because of inability to detect amplified samples by the LightCycler.

Table 9. Likely alleles as called by LightCycler 480 software after running 31 single nucleotide polymorphism (SNP) analysis on isolates from, Mississippi (MS), Arkansas (AR), Hawaii (HI), Alabama (AL), and Louisiana (LA)^{xyz}.

SNP ID	MS1	MS2	MS3	MS4	MS5	MS6	AR1	AR2	AR3	HI1	AL1	LA1	LA2
RREN_4410_3972	X	X	X	X	UNK	X	H	X	H	Y	H	X	X
RREN_1572_36933	X	X	UNK	UNK	X	X	Y	X	Y	H	H	Y	Y
RREN_4410_3979	Y	H	X	Y	X	X	X	X	X	X	X	X	X
RREN_43396_315	X	X	UNK	X	-	-	X	X	-	X	-	-	-
RREN_523_19992	H	H	H	H	H	H	H	H	H	H	H	H	H
RREN_367_3958	Y	Y	X	Y	X	X	X	UNK	X	Y	Y	Y	Y
RREN_4834_4618	Y	Y	X	X	X	H	H	Y	X	Y	X	H	H
RREN_5033_5267	X	X	H	UNK	UNK	X	Y	H	UNK	Y	UNK	X	H
RREN_845_36717	H	H	H	-	H	H	H	H	H	H	H	H	H
RREN_3215_15723	X	Y	UNK	H	H	Y	H	UNK	H	Y	UNK	Y	H
RREN_1660_513	X	X	H	UNK	H	UNK	H	UNK	H	H	H	H	H
RREN_4410_3946	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
RREN_7711_4758	Y	Y	Y	Y	Y	Y	Y	X	Y	Y	Y	Y	Y

(Table continued)

SNP ID	MS1	MS2	MS3	MS4	MS5	MS6	AR1	AR2	AR3	HI1	AL1	LA1	LA2
RREN_514_63176	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	X	Y
RREN_925_39379	H	Y	H	H	H	H	H	Y	H	H	H	H	H
RREN_514_63173	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	X	Y
RREN_91287_201	Y	Y	-	Y	-	Y	-	-	-	-	-	-	-
RREN_91287_193	H	H	-	H	-	Y	H	-	-	-	-	-	-
RREN_43396_339	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	-	-
RREN_1990_6847	X	X	Y	X	X	X	X	X	X	X	X	X	X
RREN_20709_1089	X	UNK	X	X	UNK	UNK	Y	X	Y	Y	UNK	X	X
RREN_258_12977	X	X	X	X	X	X	X	X	X	X	X	X	X
RREN_269_9935	Y	H	H	UNK	H	UNK	Y	UNK	Y	H	H	H	Y
RREN_456_104249	X	X	X	X	X	X	X	X	X	X	X	X	X
RREN_43396_325	Y	H	-	Y	-	-	-	H	-	-	-	-	-
RREN_901_49990	UNK	X	H	Y	H	UNK	H	X	X	X	UNK	Y	X
RREN_1886_12077	X	X	X	X	X	X	X	X	H	X	X	X	X

(Table continued)

SNP ID	MS1	MS2	MS3	MS4	MS5	MS6	AR1	AR2	AR3	HI1	AL1	LA1	LA2
RREN_16875_158	Y	Y	-	Y	-	-	-	Y	-	-	-	-	-
RREN_251_23034	UNK	UNK	H	UNK	Y	UNK	X	UNK	UNK	UNK	H	X	H
RREN_1895_31360	H	Y	H	Y	H	H	H	H	H	H	H	Y	Y
RREN_9137_320	H	UNK	H	Y	UNK	Y	UNK	UNK	Y	UNK	H	H	H

^x SNPs were assigned and chosen across the reniform nematode genome so that each SNP is not clustered together with the others.

^y Samples were collected from different locations in Mississippi (MS), Arkansas (AR), Hawaii (HI), Alabama (AL), and Louisiana and maintained in tomato.

^z X, Y, and H represent LightCycler 480 calls for FAM, HEX, and both fluorescent labels, respectively. “UNK” indicates LightCycler 480 could not distinguish the fluorescence while “-” indicates the failure of SNP on that particular isolate.

Only 28 SNPs amplified nematode samples from Arkansas while three mostly failed to amplify. Eight SNPs were monomorphic while other three detected heterozygous loci in samples from Arkansas. Of the remaining 17 SNPs, nine identified allelic differences in reniform nematode isolates from Arkansas.

Five SNPs failed to successfully amplify genomic DNA on samples from Hawaii, and Alabama. Of the remaining 26 SNPs, two identified allelic differences among samples from Hawaii and Alabama.

Evaluation of multiple isolates of reniform nematode from Louisiana, Mississippi, and Arkansas resulted in detection of genetic differences between and among the geographic isolates. Of the 26 SNPs that, for the most part, amplified genomic DNA of reniform nematode isolates from different geographic locations, five (RREN_5033_5267, RREN_3215_15723, RREN_269_9935, RREN_901_49990, and RREN_251_23034) were able to identify genetic differences between and among isolates of reniform nematode from Louisiana, Mississippi, and Arkansas. Because only a single isolate of reniform nematode from Hawaii and Alabama were available, genetic difference among the isolates from these locations could not be elucidated. However, among the isolates from Hawaii, and Alabama, eight SNPs were able to detect genetic variability (Table 9).

3.4 Discussion

To date no SNPs have been reported in reniform nematodes. Herein 162 putative SNPs (Appendices 1 and 2) are reported of which a select group were tested to confirm their functionality. These SNPs were tested on a group of 13 geographic isolates of reniform nematode from Louisiana, Mississippi, Arkansas, Hawaii, and Alabama. This study employed SNPs, a third generation marker (Gao *et al.*, 2016), to elucidate the genetic differences in

geographic isolates of reniform nematode from Louisiana, Mississippi, Arkansas, Hawaii, and Alabama. Existence of genetic variability in *Caenorhabditis elegans* and various plant parasitic nematodes, including but not limited to cyst nematodes (Caswell-Chen *et al.*, 1992; Folkertsma *et al.*, 1994; Kalinski and Huettel, 1988; Silva *et al.*, 2000), root-knot nematode (Guirao *et al.*, 1995; Semblat *et al.*, 1999; Tigano *et al.*, 2010; Khanal *et al.*, 2016), rice white tip nematode (Figueiredo *et al.*, 2013), and reniform nematode (Agudelo *et al.*, 2005; Tilahun *et al.*, 2008; Arias *et al.*, 2009) have been documented. Previous identifications, however, were based on use of first and second generation markers. This study is the first to report genetic variability among geographic isolates of reniform nematode using SNP analysis.

A subset of 31 SNPs (Table 8) were used to design bi-allelic KASP genotyping assays and tested on genomic DNA of 13 isolates of reniform nematode collected from various geographic locations in the United States in order to detect genetic differences among the isolates (Table 7). Most SNPs identified the two SNP alleles as well as heterozygotes across all samples (Table 9). However, some SNPs only identified one allele or the other and many only identified heterozygous alleles (Table 9). Monomorphic SNPs likely indicate a lack of genetic diversity at that genomic location where as heterozygotes likely indicate the samples were not necessarily pure. Increasing the number of different genotypes (i.e. isolates) tested would most likely increase the frequency of all alleles detected.

Five of the 31 SNPs evaluated did not work because of poor amplification. Better amplification could possibly be achieved by optimizing PCR conditions specifically for each SNP. Although some SNPs failed to amplify, 26 of 31 SNPs amplified reniform nematode DNA giving a success rate of 83.9%. This rate of success is comparable to similar studies conducted in plants with success rates of 78.5% to 88.4% (Cockram *et al.*, 2012; Saxena *et al.*, 2012; Semagn

et al., 2014; Graves *et al.*, 2016). Assuming a similar success rate for the putative SNPs not tested, then most of those SNPs shown in Appendix 2 should function.

Prior to this research no reniform nematode specific SNPs have been available. The sequence information shown for each SNP in Appendices 1 and 2 can be used to prepare reniform nematode specific SNP assays. For the 31 SNPs tested in this study using KASP assays (Appendix 1), the sequence information can be used to design other types of SNP assays. Alternatively the LGC Genomics Reference number shown in Table 8 can be used to order the KASP assays used in this study directly from LGC Genomics.

Seven SNPs did appear to identify non-heterozygous differences among the reniform nematode isolates from Louisiana indicating there are identifiable molecular differences between the isolates. As described in Chapter 2 of this dissertation, significant differences in reproduction and pathogenicity among the reniform nematode isolates was evident from greenhouse and microplot experiments that employed cotton as host. Morehouse isolate (MOR) had the greatest reproduction and pathogenicity on cotton whereas Rapides isolate (RAP) had the least. A parallel research conducted by Manjula and colleagues (personal communication) involving the same isolates of reniform nematode and soybean as host found that MOR isolate had the least reproduction and the greatest pathogenicity. This study involving SNP analysis also indicated that significant variability does occur in isolates of reniform nematode from Louisiana. Six SNPs were found to be effective in distinguishing between the MOR and RAP isolates. Salem *et al.* (2012) reported that some SNPs are associated with biological functions in an organism. Because the isolates having different levels of reproduction and pathogenicity were also found to be genetically different in SNP analysis, some of the SNPs are possibly associated with reproduction and pathogenicity functions. The association of SNPs with biological functions in

reniform nematode should be explored in future experiments. Although a reference sequence for reniform nematode is available (RREN 1.0 assembly at NCBI) the sequence has not been fully annotated. As the sequence annotation becomes more complete, genes in the areas around the SNPs that were found to identify genetic differences in this study (as well as future studies) can be investigated for potential relationships to reproduction and pathogenicity functions.

Several SNPs were polymorphic not only for reniform nematode isolates within a single geographic location but also across several locations. This suggests that a significant amount of genetic variability exists in reniform nematode isolates from within and among geographic locations. Variability in reproduction and pathogenicity in geographic isolate of reniform nematode have previously been reported in a microplot study involving cotton and soybean as hosts (McGawley *et al.*, 2010; 2011). Results from this study further supported the existence of variability in geographic isolates of reniform nematode.

Mostly the SNPs functioned and identified molecular differences among a broad range of nematode samples. Moreover, five SNPs detected genetic differences between and among the geographic isolates of reniform nematode (Table 9) indicating these SNPs could be valuable markers to distinguish reniform nematode isolates in larger geographic areas. A greater number of SNPs and utilization of a greater number of reniform nematode isolates would provide more robust data that can distinguish the pathogen from larger geographic areas. It was also observed that some SNPs polymorphic in one geographic isolate of *R. reniformis* were not always polymorphic in other geographic isolates. For example, RREN_514_63173 was polymorphic for reniform nematode isolates from Louisiana while it was monomorphic on reniform nematode from other geographic locations (Table 9). Finding of additional such SNPs would be useful in determining which geographic isolates of reniform nematode is more genetically diverse.

Evaluation of a greater number of SNPs and isolates could lead to the development of a SNP panel that could be used to categorize reniform isolates from any location.

Several studies were conducted in the past to understand the amount of genetic variability in geographic isolates of reniform nematode, although with contrasting results. Agudelo *et al.* (2005) did not find any differences in ITS1 marker of reniform nematode from the United States, Brazil, Colombia, and Honduras except for a population from Japan. In contrast to the findings of Agudelo *et al.* (2005), Tilahun *et al.* (2008) found fairly substantial nucleotide difference in ITS1 marker of the reniform nematode collected from Alabama. Microsatellite analyses have also been employed to determine genetic variability in populations of reniform nematode (Arias *et al.*, 2009; Leach *et al.*, 2012). Although microsatellite analysis is useful in determining polymorphism in a pathogen, SNPs are much more efficient and powerful markers (Salem *et al.*, 2012). Results from this research, by utilizing SNP analysis, provided a strong evidence for existence of genetic variability within and among geographic isolates of *R. reniformis*. An extensive characterization of genetic variability comprising larger number of reniform nematode isolates representing greater geography are necessary to further describe the genetic variability in this pathogen. The SNP markers developed in this study will be extremely useful in the assessment of the genetic diversity, origin, and subsequent distribution of this nematode. Additionally, once SNPs are found to be associated with genes for pathogenicity in reniform nematode, these markers will be of importance for the breeders involved in development of reniform nematode resistant crops.

CHAPTER 4. SUMMARY AND CONCLUSIONS

The reniform nematode (*Rotylenchulus reniformis* Linford and Oliveira) has become one of the major constraints of cotton production in the United States cotton belt. Approximately 205 thousand bales of the United States cotton was lost to this nematode in 2016 (Lawrence *et al.* 2017). Among the 16 states in cotton belt, cotton yield in Mississippi, Alabama, and Louisiana sustain greatest damage to this pathogen (Lawrence *et al.*, 2017).

Some common management options for reniform nematode in cotton include chemical, crop rotation, and biological. Chemical control is the most widely used method of reniform nematode management in cotton, however, it is least desirable because of health and environmental issues. Crop rotation with a non-host crop can help reduce the nematode population, however population usually resurges to a damaging level in a single season of susceptible crop plantation (Robinson *et al.*, 2007). Use of host plant resistance is the most widely desired method of reniform nematode management. Unfortunately, reniform nematode resistant upland cotton cultivars are not available to date. Some upland cotton breeding lines showing moderate to high levels of reniform nematode resistance are available, but their performance is not consistent across wide geographic area (Yik and Birchfield, 1984; Robinson *et al.*, 1997; Robinson *et al.*, 2004; Weaver *et al.*, 2007). This inconsistency, to the most part, is caused by existence of physiological and genetic differences in geographic populations of reniform nematode. A few studies conducted in the past using geographic isolates of reniform nematode have suggested the existence of virulence of phenotypes (McGawley *et al.*, 2010, 2011). Additionally, several studies involving internal transcribed spacer-1 (ITS1), 18S, and microsatellite marker analysis reported a significant amount of genetic variation in geographic isolates of reniform nematode while one study was unable to detect any differences making

genetic variability more elusive (Agudelo *et al.*, 2005; Tilahun *et al.*, 2008; Arias *et al.*, 2009, Leach and Agudelo, 2012). Reniform nematode populations derived from single-egg mass and collected from West Carroll, Rapides, Morehouse, and Tensas parishes in Louisiana were used in this study to determine variability in this pathogen. Variability in reniform nematode was studied from two perspectives: (1) physiological variability as determined from reproduction and pathogenicity data obtained from microplot and greenhouse experiments, and (2) genetic variability determined from single nucleotide polymorphism (SNP) analysis. Two full season (150 days) microplot experiments were conducted using three most widely planted upland cotton cultivars (Phytogen 499 WRF, Deltapine 1133 B2RF, and Phytogen 333 WRF) that were recommended for use in Louisiana in 2015. Similarly, two 60-day greenhouse experiments were conducted to determine reproduction of four reniform nematode isolates endemic in Louisiana using three upland cotton cultivars that were used in microplot experiment, two cotton germplasm lines showing moderate to high level of resistance (MT2468 Ren3, and M713 Ren5), and one susceptible control. The reasons behind employing SNP analysis to determine genetic variability in geographic populations of reniform nematode are that SNPs are robust, more efficient, and most suitable to analyze genomic data obtained from next generation sequencing.

Results from microplot experiments suggest a significant difference in reproduction and pathogenicity of reniform nematode populations. In the microplot experiment, Morehouse and Rapides isolates, respectively, had the greatest and the least reproduction while West Carroll and Tensas isolates had intermediate levels of reproduction. Effect of reproduction was reflected in plant yield. The Morehouse isolate caused the greatest reduction in plant dry weight, number of bolls, seed cotton weight, and lint weight while the Rapides isolate caused the least.

Data from two greenhouse experiments showed similar reproduction results to that of microplot experiment. In other words, Morehouse isolates had significantly higher reproduction while Rapides isolate had significantly lower reproduction. Reproduction of West Carroll and Tensas isolates on cotton was intermediate. Differences in reproduction and pathogenicity of reniform nematode isolates was more pronounced on the germplasm lines in greenhouse experiments. Greenhouse and microplot studies conducted in the past have reported significant differences in reproduction and pathogenicity of geographic populations of reniform nematode suggesting existence of virulence phenotypes (McGawley *et al.*, 2010, 2011). Results from this experiment support the existence of virulence phenotypes in reniform nematode.

Kompetitive allele-specific PCR (KASP) genotyping assay was conducted to test 31 SNPs on 13 reniform nematode isolates collected from Louisiana, Mississippi, Arkansas, Hawaii, and Alabama. Of 25 SNPs that amplified reniform nematode isolates, five SNPs were of the most interest as they identified genetic differences between and among geographic isolates of reniform nematode from Louisiana, Mississippi, and Arkansas. This study is the first to report genetic variability in geographic isolates of reniform nematode employing SNP assay. Further studies comprising larger number of SNPs and greater number of reniform nematode isolates are necessary to understand SNP polymorphism and its association with biological function in this pathogen. The SNP markers developed in this study will be extremely useful in resistance breeding programs as well as in the assessment of the genetic diversity, origin, and subsequent distribution of this nematode.

REFERENCES

- Agudelo, P., R. T. Robbins, J. M. Stewart, and A. L. Szalanski. 2005. Intraspecific variability of *Rotylenchulus reniformis* from cotton-growing regions in the United States. *Journal of Nematology* 37:105-114.
- Allen, T. W., C. A. Bradley, J.P Damicone, N. S. Dufault, T. R. Faske, C. A. Hollier, T. Isakeit, R. C. , Kemerait, N. M. Kleczewski, R. J. Kratochvil, H. L. Mehl, J. D. Mueller, C. Overstreet, P. P. Price, E. J. Sikora, T. N. Spurlock, L. Thiessen, W. J. Wiebold, and H. Young. 2017. Southern United States soybean disease loss estimates for 2017. Proceeding of the 44th Southern Soybean Disease Workers meeting, Pensacola, FL p. 3-8.
- Anonymous. 2010. Field crops usual planting and harvesting dates. United States Department of Agriculture, National Agricultural Statistics Service. Online. <http://usda.mannlib.cornell.edu/usda/current/planting/planting-10-29-2010.pdf>.
- Anonymous. 2015. Cotton varieties planted 2015 crop. U. S. Department of Agriculture, Agricultural Marketing Service. Online.<http://www.ams.usda.gov/mnreports/cnavar.pdf>.
- Anonymous. 2016. An introduction to next generation sequencing. Online. http://www.illumina.com/content/dam/illumina-marketing/documents/products/illumina_sequencing_introduction.pdf.
- Anonymous. 2017. Crop production 2016 summary. United States Department of Agriculture, National Agricultural Statistics Service. Online. <http://usda.mannlib.cornell.edu/usda/current/CropProdSu/CropProdSu-01-12-2017.pdf>.
- Arias, R. S., S. R. Stetina, J. L. Tonos, L. A. Scheffler, and B. E. Scheffler. 2009. Microsatellites reveal genetic diversity in *Rotylenchulus reniformis* populations. *Journal of Nematology* 41:146-156.
- Barker, K. R., C. C. Carter, and J. N. Sasser. 1985. An advanced treatise on *Meloidogyne*, volume II: methodology. North Carolina State University Graphics.
- Bell, A. A., A. F. Robinson, J. Quintana, N. D. Dighe, M. A. Menz, D. M. Stelly, X. Zheng, J. E. Jones, C. Overstreet, E. Burris, R. G. Cantrell, and R. L. Nichols. 2014a. Registration of LONREN-1 and LONREN-2 germplasm lines of upland cotton resistant to reniform nematode. *Journal of Plant Registrations* 8:187-190.
- Bell, A. A., A. F. Robinson, J. Quintana, S. E. Duke, J. L. Starr, D. M. Stelly, X. Zheng, S. Prom, V. Saladino, O. A. Gutiérrez, S. R. Stetina, and R. L. Nichols. 2014b. Registration of BARBREN-713 germplasm line of upland cotton resistant to reniform and root-knot nematodes. *Journal of Plant Registrations* 8:89-93.
- Berg, E. V., J. E. Palomares-Rius, N. Vovlas, L. R. Tiedt, P. Castillo, and S. A. Subbotin. 2016. Morphological and molecular characterisation of one new and several known species of

- the reniform nematode, *Rotylenchulus* Linford & Oliveira, 1940 (Hoplolaimidae: Rotylenchulinae), and a phylogeny of the genus. *Nematology* 18:67-107.
- Bhandari, B., G. O. Myers, M. O. Indest, and C. Overstreet. 2015. Response of five resistant cotton genotypes to isolates of *Rotylenchulus reniformis* collected from reniform nematode infested fields of Louisiana. *Nematropica* 45:252-262.
- Birchfield, W. 1962. Host-parasite relations of *Rotylenchulus reniformis* on *Gossypium hirsutum*. *Phytopathology* 52:862-865.
- Birchfield, W. and J. E. Jones. 1961. Distribution of the reniform nematode in relation to crop failure of cotton in Louisiana. *Plant Disease Reporter* 45:671-673.
- Birchfield, W., and L. R. Brister. 1963. Susceptibility of cotton and relatives to reniform nematode in Louisiana. *Plant Disease Reporter* 47:990-992.
- Birchfield, W., and W. J. Martin. 1967. Reniform nematode survival in air-dried soil. *Phytopathology* 57:804 (abstract).
- Blessitt, J. A., S. R. Stetina, T. P. Wallace, P. T. Smith, and G. L. Sciumbato. 2012. Cotton (*Gossypium hirsutum*) cultivars exhibiting tolerance to the reniform nematode (*Rotylenchulus reniformis*). *International Journal of Agronomy* doi:10.1155/2012/893178.
- Byrd, D. W., K. R. Barker, H. Ferris, C. J. Nusbaum, W. E. Griffin, R. H. Small, and C. A. Stone. 1976. Two semi-automatic elutriators for extracting nematodes and certain fungi from soil. *Journal of Nematology* 8:206-212.
- Caswell-Chen, E. P., V. M. Williamson, and F. F. Wu. 1992. Random amplified polymorphic DNA analysis of *Heterodera cruciferae* and *H. schachtii* populations. *Journal of Nematology* 24:343-351.
- Cockram, J., H. Jones, C. Norris, and D.M. O'Sullivan. 2012. Evaluation of diagnostic molecular markers for DUS phenotypic assessment in the cereal crop, barley (*Hordeum vulgare* ssp. *vulgare* L.). *Theoretical and Applied Genetics* 125:1735-1749.
- Crow W. T., D. W. Dickson, D. P. Weingartner, R. McSorley, and G. L. Miller. 2000. Yield reduction and root damage to cotton induced by *Belonolaimus longicaudatus*. *Journal of Nematology* 32:205-209.
- Dabbert, T. A., and M. A. Gore. 2014. Challenges and perspectives on improving heat and drought stress resilience in cotton. *The Journal of Cotton Science* 18:393-409.
- Dasgupta, D. R., and A. R. Seshadri. 1971. Races of the reniform nematode, *Rotylenchulus reniformis* Lindford and Oliveira, 1940. *Indian Journal of Nematology* 1:21-24.

- Dieterich, C., and R. J. Sommer. 2009. How to become a parasite – lessons from the genomes of nematodes. *Trends Genet* 25:203-209.
- Dighe, N. D., A. F. Robinson, A. A. Bell, M. A. Menz, R. G. Cantrell, and D. M. Stelly. 2009. Linkage mapping of resistance to reniform nematode in cotton (*Gossypium hirsutum* L.) following introgression from *G. longicalyx* (Hutch. & Lee). *Crop Science* 49:1151-1164.
- Ducharme, E. P., and W. Birchfield. 1956. Physiologic races of the burrowing nematode. *Phytopathology* 46:615-616.
- Faga, B. W. Maury, D. A. Bruckner, and C. Grose. 2001. Identification and mapping of single nucleotide polymorphisms in the varicella-zoster virus genome. *Virology* 280:1-6.
- Fang, D. D., and S. R. Stetina. 2011. Improving cotton (*Gossypium hirsutum* L.) plant resistance to reniform nematodes by pyramiding *Ren1* and *Ren2*. *Plant Breeding* 130:673-678.
- Figueiredo, J., M. J. Simoes, P. Gomes, C. Barroso, D. Pinho, L. Conceicao, L. Fonseca, I. Abrantes, M. Pinheiro, and C. Egas. 2013. Assessment of the geographic origins of pinewood nematode isolates via single nucleotide polymorphism in effector genes. *Plos One* 8:e83542.
- Folkertsma, R. T., J. N. A. M. R. Van Der Voort, M. P. E. Van Gent-Pelzer, K. E. Groot, W. J. Van Den Bos, A. Schots, J. Bakker, and F. J. Gommers. 1994. Inter- and intraspecific variation between populations of *Globodera rostochiensis* and *G. pallida* revealed by random amplified polymorphic DNA. *Phytopathology* 84:807-811.
- Gao, L., J. Jia, and X. Kong. 2016. A SNP-based molecular barcode for characterization of common wheat. *PLoS ONE* 11: e0150947.
- Gaur, H. S., and R. N. Perry. 1991. The role of the moulted cuticles in the desiccation survival of adults of *Rotylenchulus reniformis*. *Revue de Nématology* 14:491-496.
- Germani, G. 1978. Caractères morpho-biométriques de trois espèces ouest-africaines de *Rotylenchulus* Linford & Oliveirs 1940 (Nematoda: Tylenchida). *Revue de Nematologie* 1:241-250.
- Grada, A., and K. Weinbrech. 2013. Next-generation sequencing: methodology and application. *Journal of Investigative Dermatology* 133:1-4.
- Graves, H., A. L. Rayburn, J. L. Gonzalez-Hernandez, G. Nah, Do-Soon Kim, and D. K. Lee. 2016. Validating DNA polymorphisms using KASP assay in prairie cordgrass (*Spartina pectinata* Link) populations in the U.S. *Frontiers in Plant Science* doi: 10.3389/fpls.2015.01271
- Grover, A., and P. C. Sharma. 2014. Development and use of molecular markers: past and present. *Critical Reviews in Biotechnology* 36: 290-302.

- Guirao, P., A. Moya, and J. L. Cenis. 1995. Optimal use of random amplified polymorphic DNA in estimating the genetic relationship of four major *Meloidogyne* spp. *Phytopathology* 85:547-551.
- Huettel, R. N., and T. Yaegashi. 1988. Morphological differences between *Radopholus citrophilus* and *R. similis*. *Journal of Nematology* 20:150-157.
- Hussey, R. S., and K. R. Barker. 1973. A comparison of methods for collecting inocula for *Meloidogyne* spp., including a new technique. *Plant Disease Reporter* 57:1025-1028.
- Jenkins, W. R. 1964. A rapid centrifugal-flotation technique for separating nematodes. *Plant Disease Reporter* 48:692.
- Kalinski, A. and R. N. Huettel. 1988. DNA restriction fragment length polymorphism in races of the soybean cyst nematode, *Heterodera glycines*. *Journal of Nematology* 20:532-538.
- Khanal, C., R. T. Robbins, T. R. Faske, A. L. Szalanski, E. C. McGawley, and C. Overstreet. 2016. Identification and haplotype designation of *Meloidogyne* spp. of Arkansas using molecular diagnostics. *Nematropica* 46:261-270.
- Koenning, S. R., T. L. Kirkpatrick, J. L. Starr, J. A. Wrather, N. R. Walker, and J. D. Mueller. 2004. Plant-parasitic nematodes attacking cotton in the United States: old and emerging challenges. *Plant Disease* 88:100-113.
- Lawrence, K., A. Hagan, R. Norton, T. Faske, R. Hutmacher, J. Mueller, D. Wright, I. Small, B. Kemerait, C. Overstreet, P. Price, G. Lawrence, T. Allen, S. Atwell, A. Jones, S. Thomas, N. Goldberg, R. Boman, J. Goodson, H. Kelly, J. Woodward, and H. L. Mehl. 2017. Cotton disease loss estimate committee report, 2016. Pp. 150-152 *in*: Proceedings of the 2017 Beltwide Cotton Conferences, Dallas, TX. Cordova: National Cotton Council.
- Lawrence, K., A. Hagan, R. Norton, T. Faske, R. Hutmacher, J. Mueller, D. Wright, I. Small, B. Kemerait, C. Overstreet, P. Price, G. Lawrence, T. Allen, S. Atwell, A. Jones, S. Thomas, N. Goldberg, R. Boman, J. Goodson, H. Kelly, J. Woodward, and H. L. Mehl. 2017. Cotton disease loss estimate committee report, 2016. Pages 150-152 *in*: Proceedings of the Beltwide Cotton Conferences, Dallas, TX. Cordova: National Cotton Council.
- Leach, M., P. Agudelo, and A. Lawton-Rauh. 2012. Genetic variability of *Rotylenchulus reniformis*. *Plant Disease* 96:30-36.
- Lehman, P. S., and R. N. Inserra. 1990. Morphometric variation of *Rotylenchulus parvus* and *Rotylenchulus reniformis* populations in the southern United States. *Soil and Crop Science Society of Florida Proceedings* 49:220-226.
- Linford, M. B., and J. M. Oliveira. 1940. *Rotylenchulus reniformis*, nov. gen., n sp., a nematode parasite of roots. *Proceedings of the Helminthological Society* 7:35-42.

- Linlokken, A. N., T. O. Haugen, M. P. Kent, and S. Lien. 2017. Genetic differences between wild and hatchery-bred brown trout (*Salmo trutta* L.) in single nucleotide polymorphisms linked to selective traits. *Ecology and Evolution* 7:4963-4972.
- Lofton, J., B. Haggard, D. Fromme, and B. Tubana. 2014. Utilization of poultry litter, tillage, and cover crops for cotton production on highly degraded soils in northeast Louisiana. *The Journal of Cotton Science* 18:376-384.
- Lu, S., M. C. Edwards, and T. L. Friesen. 2013. Genetic variation of single nucleotide polymorphisms identified at the mating type locus correlates with form-specific disease phenotype in the barley net blotch fungus *Pyrenophora teres*. *European Journal of Plant Pathology* 135:49-65.
- Mauney, J. R. 2015. Anatomy and morphology of cultivated cottons. Pp. 77-96 in D. D. Fang, and R. G. Percy, 2 ed. *Cotton*. Madison, WI: American Society of Agronomy.
- Mauney, J. R., and J. M. Stewart. 1986. *Cotton physiology: the cotton foundation reference book series*. Memphis, Tennessee: the cotton foundation.
- McCarty, J. C., J. N. Jenkins, M. J. Wubben, O. A. Gutierrez, R. W. Hayes, F. E. Callahan, and D. Deng. 2013. Registration of three germplasm lines of cotton derived from *Gossypium barbadense* L. accession GB713 with resistance to the reniform nematode. *Journal of Plant Registrations* 7:220-223.
- McCarty, J. C., J. N. Jenkins, M. J. Wubben, R. W. Hayes, and J. M. LaFoe. 2012. Registration of three germplasm lines of cotton derived from *Gossypium hirsutum* L. accession T2468 with moderate resistance to the reniform nematode. *Journal of Plant Registrations* 6:85-87.
- McGawley, E. C., and C. Overstreet. 1995. Reproduction and pathological variation in populations of *Rotylenchulus reniformis*. *Journal of Nematology* 27:508 (abstract).
- McGawley, E. C., C. Overstreet, and M. J. Pontif. 2011. Variation in reproduction and pathogenicity of geographic isolates of *Rotylenchulus reniformis* on soybean. *Nematropica* 41:12-22.
- McGawley, E. C., M. J. Pontif, and C. Overstreet. 2010. Variation in reproduction and pathogenicity of geographic Isolates of *Rotylenchulus reniformis* on cotton. *Nematropica* 40:275-288.
- Meyer, L., and S. MacDonald. 2016. *Cotton and Wool Outlook*. Economic Research Service, United States Department of Agriculture. Online.
<http://usda.mannlib.cornell.edu/usda/ers/CWS//2010s/2017/CWS-04-13-2017.pdf>
- Morais, S., R. Marco-Moles, R. Puchades, and A. Maquieira. 2006. DNA microarraying on compact disc surfaces. Application to the analysis of single nucleotide polymorphisms in Plum pox virus. *Chemical Communications* 14:2368-2370.

- Nakasono, K. 2004. Studies on morphological and physio-ecological variations of the reniform nematode, *Rotylenchulus reniformis* Lindford and Oliveira, 1940 with an emphasis on differential geographical distribution of amphimictic and parthenogenetic populations in Japan. *Journal of Nematology* 36:356-420.
- Niblack, T. L., P. R. Arelli, G. R. Noel, C. H. Opperman, J. H. Orf, D. P. Schmitt, J. G. Shannon, and G. L. Tylka. 2002. A revised classification scheme for genetically diverse populations of *Heterodera glycines*. *Journal of Nematology* 34:279-282.
- Noe, J. P. 1992. Variability among populations of *Meloidogyne arenaria*. *Journal of Nematology* 24:404-414.
- Noe, J. P. 1993. Damage functions and population changes of *Hoplolaimus columbus* on cotton and soybean. *Journal of Nematology* 25:440-445.
- Ojeda, D. I., B. Dhillon, C. K. M. Tsui, and R. C. Hamelin. 2014. Single-nucleotide polymorphism discovery in *Leptographium longiclavatum*, a mountain pine beetle-associated symbiotic fungus, using whole-genome resequencing. *Molecular Ecology Resources* 14:401-410.
- Oosterhuis, D. M., and F. M. Bourland. 2001. Development of the cotton plant. Pp 3-7 in T. L. Kirkpatrick, and C. S. Rothrock, 2 ed. *Compendium of Cotton Diseases*. St. Paul, MN: The American Phytopathological Society Press.
- Overstreet, C. 2006. The impact of reniform nematode on cotton production in the USA. *Nematropica* 26:216 (abstract.).
- Overstreet, C. 2015. Using resistance to manage nematodes in soybean. Louisiana crop Newsletter. Online. <http://www.lsuagcenter.com/MCMS/RelatedFiles/%7BF12F62D7-128B-47FD-8D34-BFE1841CBA1B%7D/Louisiana-Crops-Newsletter-March-2015.pdf>
- Overstreet, C., and E. C. McGawley. 1994. Cotton production and *Rotylenchulus reniformis* in Louisiana. *Journal of Nematology* 26:562-563 (abstract).
- Overstreet, C., and E. C. McGawley. 1998. The emergence of *Rotylenchulus reniformis* as the major nematode pest in Louisiana, U.S.A. *Nematropica* 28:142 (abstract).
- Overstreet, C., E. C. McGawley, C. Hollier, D. D. Fromme, and T. Price. 2014. Nematode management. Louisiana cotton production. LSU AgCenter. Online. <http://www.lsuagcenter.com/profiles/lblack/articles/page1473702737099>.
- Rattei, T., S. Ott, M. Gutacker, J. Rupp, M. Maass, S. Schreiber, W. Solbach, T. Wirth, and J. Gieffers. 2007. Genetic diversity of the obligate intracellular bacterium *Chlamydomphila pneumoniae* by genome-wide analysis of single nucleotide polymorphisms: evidence for highly clonal population structure. *BMC Genomics* 8:355. Doi: 10.1186/1471-2164-8-355.

- Riggs, R. D., M. L., Hamblen, and L. Rakes. 1981. Infra-species variation in reactions to hosts in *Heterodera glycines* populations. *Journal of Nematology* 13:171-179.
- Robinson, A. F. 2007. Reniform in U.S. cotton: when, where, why, and some remedies. *Annual Review of Phytopathology* 45:263-88.
- Robinson, A. F., A. A. Bell, N. Dighe, M. A. Menz, R. L. Nichols, and D. M. Stelly. 2007. Introgression of resistance to nematode *Rotylenchulus reniformis* into upland cotton (*Gossypium hirsutum*) from *G. longicalyx*. *Crop Science* 47:1865-1877.
- Robinson, A. F., A. C. Bridges, and A. E. Percival. 2004. New sources of resistance to the reniform (*Rotylenchulus reniformis*) and root-knot (*Meloidogyne incognita*) nematode in upland (*Gossypium hirsutum* L.) and sea island (*G. barbadense* L.) cotton. *Journal of Cotton Science* 8:191-197.
- Robinson, A. F., C. G. Cook, and A. E. Percival. 1999. Resistance to *Rotylenchulus reniformis* and *Meloidogyne incognita* race 3 in the major cotton cultivars planted since 1950. *Crop Science* 39:850-858.
- Robinson, A. F., R. N. Inserra, E. P. Caswell-Chen, N. Vovlas, and A. Troccoli. 1997. *Rotylenchulus* species: Identification, distribution, host ranges, and crop plant resistance. *Nematropica* 27:127-180.
- Romano, G. B., E. J. Sacks, S. R. Stetina, A. F. Robinson, D. D. Fang, O. A. Gutierrez, and J. A. Scheffler. 2009. Identification and genomic location of a reniform nematode (*Rotylenchulus reniformis*) resistance locus (*Ren^{ari}*) introgressed from *Gossypium aridum* into upland cotton (*G. hirsutum*). *Theoretical and Applied Genetics* 120:139-150.
- Rosa, R. C. T., R. M. Moura, E. M. R. Pedrosa, and A. Chaves. 2003. Ocorrência de *Rotylenchulus reniformis* em Cana de açúcar no Brasil. *Nematologia Brasileira* 27:9395.
- Salem, M., R. L. Vallejo, T. D. Leeds, Y. Palti, S. Liu, A. Sabbagh, C. E. Rexroad III, and J. Yao. 2012. RNA-Seq identifies SNP markers for growth traits in rainbow trout. *PLoS ONE* 7:e36264.
- Samson-Himmelstjerna, G. V., W. J. Blackhall, J. S. McCarthy, and P. J. Skuce. Single nucleotide polymorphism (SNP) markers for benzimidazole resistance in veterinary nematodes. *Parasitology* 134:1077-1086.
- Saxena, R. K., R. V. Penmetsa, H. D. Upadhyaya, A. Kumar, N. Carrasquilla-Garcia, J. A. Schlueter, A. Farmer, A. M. Whaley, B. K. Sarma, G. D. May, D. R. Cook, and R. K. Varshney. 2012. Large-scale development of cost-effective single-nucleotide polymorphism marker assays for genetic mapping in pigeon pea and comparative mapping in legumes. *DNA Research* 19:449-461.
- Schafer, J. F. 1971. Tolerance to plant disease. *Annual Review of Phytopathology* 9:235-252.

- Seinhorst, J. W. 1957. Some aspects of the biology and ecology of stem eelworms. *Nematologica*: 2:355-361.
- Semagn, K., R. Babu, S. Hearne, and M. Olsen. 2014. Single nucleotide polymorphism genotyping using Kompetitive Allele Specific PCR (KASP): overview of the technology and its application in crop improvement. *Molecular Breeding* 33:1-14.
- Semblat, J. P., M. Bongiovanni, E. Wajnberg, A. Dalmasso, P. Abad, and P. Castagnone-Sereno. 1999. Virulence and molecular diversity of parthenogenetic root-knot nematodes, *Meloidogyne* spp. *Heredity* 84:81-89.
- Silva A. T. D., J. C. V. Penna, Goulart L. R., M. A. D. Santos, and N. E. Arantes. 2000. Genetic variability among and within races of *Heterodera glycines* Ichinohe assessed by RAPD markers. *Genetics and Molecular Biology* 23:323-329.
- Silva, K. J. D., E. A. Souza, A. Sartorato, and C. N. Souza-Freire. 2008. Pathogenic variability of isolates of *Pseudocercospora griseola*, the cause of common bean angular leaf spot, and its implications for resistance breeding. *Journal of Phytopathology* 156:602-606.
- Smith A. L., and A. L. Taylor. 1941. Nematode distribution in the 1940 regional cotton-wilt plots. *Phytopathology* 31:771.
- Soares, P. L. M., J. M. dos Santos, and A. S. Ferraudo. 2004. Estudo morfofométrico comparativo de 58 populacões Brasileiras de *Rotylenchulus reniformis* (Nemata: Rotylenchulinae). *Fitopatologia Brasileira* 29:419-424.
- Starr, J. L., C. W. Smith, K. Ripple, E. Zhou, R. L. Nichols, and T. R. Faske. 2011. Registration of TAM RKRNR-9 and TAM RKRNR-12 germplasm lines of upland cotton resistant to reniform and root-knot nematodes. *Journal of Plant Registrations* 5:393-396.
- Starr, J. L., S. R. Koenning, T. L. Kirkpatrick, A. F. Robinson, P. A. Roberts, and R. L. Nichols. 2007. The future of nematode management in cotton. *Journal of Nematology* 39:283-294.
- Stetina, S. R., and L. D. Young. 2006. Comparisons of female and egg assays to identify *Rotylenchulus reniformis* resistance in cotton. *Journal of Nematology* 38:326-332.
- Stetina, S. R., L. D. Young, W. T. Pettigrew, and H. A. Bruns. 2007. Effect of corn-cotton rotations on reniform nematode populations and crop yield. *Nematropica* 37:237-248.
- Thorp, K. R., S. Ale, M. P. Bange, E. M. Barnes, G. Hoogenboom, R. J. Lascano, A. C. McCarthy, S. Nair, J. O. Paz, N. Rajan, K. R. Reddy, G. W. Wall, and J. W. White. 2014. Development and application of process-based simulation models for cotton production: a review of past, present, and future directions. *The Journal of Cotton Science* 18:10-47.
- Tigano, M., K. de Siqueira, P. Castagnone-Sereno, K. Mulet, P. Queiroz, M. dos Santos, C. Teixeira, M. Almeida, J. Silva, and R. Carneiro. 2010. Genetic diversity of the root-knot

- nematode *Meloidogyne enterolobii* and development of a SCAR marker for this guava-damaging species. *Plant Pathology* 59:1054-1061.
- Tilahun, M., K. Soliman, K. S. Lawrence, L. J. Cseke, and J. W. Ochieng. 2008. Nuclear ribosomal DNA diversity of a cotton pest (*Rotylenchulus reniformis*) in the United States. *African Journal of Biotechnology* 7:3217-3224.
- Van der Beek, J. G., P. W. Th. Maas, G. J. W. Janssen, C. Zijlstra, and C. H. Van Silfhout. 1999. A pathotype system to describe intraspecific variation in pathogenicity of *Meloidogyne chitwoodi*. *Journal of Nematology* 31:386-392.
- Wallace T. P., D. Bowman, B. T. Campbell, P. Chee, O. A. Gutierrez, R. J. Kohel, J. McCarty, G. Myers, R. Percy, F. Robinson, W. Smith, D. M. Stelly, J. M. Stewart, P. Thaxton, M. Ulloa, and D. B. Weaver. 2009. Status of the USA cotton germplasm collection and crop vulnerability. *Genetic Resources and Crop Evolution* 56:507-532.
- Wallace, T. P., B. Golden, P. M. Thaxton, J. Scheffler, K. S. Lawrence, and D. Weaver. 2013. Agronomic performance of *barbadense* and *longicalyx* derived breeding lines. Pp. 1005 in *Proceedings of the Beltwide Cotton Conferences*, San Antonio, TX.
- Weaver, D. B. 2015. Cotton nematodes. Pp. 547-570 in D. D. Fang, and R. G. Percy, Cotton. Madison, WI: American Society of Agronomy.
- Weaver, D. B., K. S. Lawrence, and E. Van Santen. 2007. Reniform nematode resistance in upland cotton germplasm. *Crop Science* 47:19-24.
- Werlemark, G., B. U. Carlson-Nilsson, and C. G. Davidson. 2006. Genetic variation in the rose pathogen *Marssonina rosae* estimated by RAPD. *International Journal of Horticultural Science* 12:63-67.
- Yang, C. H., K. C. Wu, H. U. Dahms, L. Y. Chuang, and H. W. Chang. 2017. Single nucleotide polymorphism barcoding of cytochrome c oxidase I sequences for discriminating 17 species of Columbidae by decision tree algorithm. *Ecology and Evolution* 7:4717-4725.
- Yik, C. P., and W. Birchfield. 1984. Resistant germplasm in *Gossypium* species and related plants to *Rotylenchulus reniformis*. *Journal of Nematology* 16:146-153.

APPENDIX 1. FLANKING SEQUENCE INFORMATION FOR SNP ASSAYS TESTED IN THIS STUDY. THE "SNP ID" PROVIDES A COMMON NAME INCLUDING ORGANISM ABBREVIATION "RREN" ROTYLENCHULUS RENIFORMIS (NEMATODES), THE REFERENCE ASSEMBLY CONTIG (FIRST NUMBER) AND THE SEQUENCE LOCATION (SECOND NUMBER) OF THE SNP POSITION.

S.N.	SNP ID	SNP	Surrounding Sequence 5'-3' (approximately 100 bp each side)
1	RREN_4410_3972	[T/C]	AAGCAGACAGCGAAAAAGCCCCACTCGTGACCGCGTAGAGGGATAGCGAGGAA GGGATGGGGAGGCGACGAGCGAAAAAGACGCCCCGGGGGAAGGA[T/C]AGAGG GAATTCCCCTCTCCCCAGGGAAGCAAGTACGGGGAAACCACTCAGATGCGATG AGAACGAAGGGTTTTTCGCTTAGGAAAAGGCAATGCGAGAGGAT
2	RREN_1572_36933	[T/C]	TTTTGGACATCTTTCGCTTCTCCTGGACAATTTTCTATCTTTCGAATATTTTGGAC TTTTTTGGACACCTTATGATTGACCATTTACAGCCCCATCC[T/C]GCTGGCCAAGC GGTCTCCTACTCGGCCAGCAGAAGAACCTGTTGATGTGGGCGGTGGCCGTCGG CTCCATGCTCGGCACTTTCCCCTTCGCCTGGCTCTAC
3	RREN_4410_3979	[A/G]	CAGCGAAAAAGCCCCACTCGTGACCGCGTAGAGGGATAGCGAGGAAGGGATGG GGAGGCGACGAGCGAAAAAGACGCCCCGGGGGAAGGATAGAGGG[A/G]ATTCC CACTCTCCCCAGGGAAGCAAGTACGGGGAAACCACTCAGATGCGATGAGAACG AAGGGTTTTTCGCTTAGGAAAAGGCAATGCGAGAGGATTCGCTGT

8 RREN_5033_5267 [T/C] ACTTCCATCTCCAAGTTGTTTATAGAGATGTTTGCCGAGTTCAGTGGGATTCGTG
ATATTCGAAGGGGGCACTGTTACACGCACCTCCTGCCCTGTT[T/C]GCCATACAA
CTCGCATATATTGCTGTTCCATAGTTCCCAAGGATGGCCATGAATGGGGATAAA
CCATCACTGCAAGCCCCACATACCAATCTGAATTGAATT

9 RREN_845_36717 [A/G] GACTTTCTGCATGGCTTTGAGGAGTAAAATTCTTGCCTAAAATTACAATCTTGTT
TTATTAGTTTTTTTATTCAAAAAAATAGCTTACAGCAGAGGTC[A/G]TGAACAATG
AGATGATGATGTTGGAGCACGCGTGCTCCTCGACCTGAATTATGAAAAAGTTTA
TTTTTCTCGAATAAAAAATATCTAATTTATAAAAAACATAC

10 RREN_3215_15723 [T/C] TTTTTCATGAGCACTCTTTTCGTTTTCTTCAACACTTTTTCTGAGCAATCTCTCGT
TTCAACGAACACTTTTTTTATGGGGGTAAACCGTACAATA[T/C]AAGAGCCGACT
TATTTTGTACAGCGTACACTCTACACTGTATTGCAAAATAGAATTAATAAAATAA
AAATAGTCTGGTACTTAGGTATATAGTTTACGTGACC

11 RREN_1660_513 [T/C] GGGTGCGGTAGTCGGTTCGGCTTATGGCGTAGAGCTGATCAGTGAGAACCAGAT
CCCGTCCGCACTCCAAGACATATTCCAAATTGGTGATGCTTAA[T/C]GAAAATGT
TTGATCTTCTGACATGAACCAATCATCAAACATGGTTCCTTTCATTTCTGCATC
AACACGTACCGGCATTGTGTGTCCGCTTTGGGGCGGTTC

12 RREN_4410_3946 [A/G] GGAAAGTGTGCTCTCGACGAAAAGGAAGCAGACAGCGAAAAAGCCCCACTCG
TGACCGCGTAGAGGGATAGCGAGGAAGGGATGGGGAGGCGACGA[A/G]CGAAA
AAGACGCCCCGGGGGAAGGATAGAGGGAATTCCCACTCTCCCCAGGGAAGCAA
GTACGGGGAAACCACTCAGATGCGATGAGAACGAAGGGTTTTTCG

13 RREN_7711_4758 [T/C] GGAGGTGAGAGAGTGTAGAGTGGTGAAGTGGAGGTGAGAGAGTACGGTATGTG
AGAGAGTACGGTATGTGACAGTACCGTGTTGTCCACGACCACGA[T/C]ACACTCT
TTGTTGGTTGCCTTCACAACTCGGCAAACGGCCTCAATATCGATGACTTTCAGCA
ATGGGTGGACGGTGTTCGAACCAGACCATCTATGGATG

14 RREN_514_63176 [A/G] TGCCTCAAATCCTCGGATTTTTTCAGAAATTCGTCAAATTTTATTGGCATTTTTT
CTGTGTAGAGAGTTTATTGGAAGTCGGGAGGTGTGGCTGAA[A/G]AGCATCTCCA
ACCTGTTGCCGCGGCACATCCTCAAGGCCTCATTGGCACTGCAGTCGGTGGTGC
ACCAGTACGAGCCGGACGCCATGATGCCAATCCCGTCA

15 RREN_925_39379 [C/G] GAGGCAAGAGGCATCGAACAAATGGATCAATCTGTCCCTACTTCCGGAAGCCAG
CAAATGAAGCTGATCTGGTGCTAAATTTAACCTTATGTATTCA[C/G]TTGGAAAT
AGCAAAAATTGATAAAATGAAAAAATGGACTAACCAGCCGAAGGATAGTTGTA
TTCCAGACACAACACCTCGCTGTCATGTGCCTCTAATTCGA

16 RREN_514_63173 [T/G] TTTTGCCTCCAAATCCTCGGATTTTTCAGAAATTCGTCAAATTTTATTGGCATT
TTTCTGTGTAGAGAGTTTATTGGAAGTCGGGAGGTGTGGCT[T/G]AAGAGCATCT
CCAACCTGTTGCCGCGGCACATCCTCAAGGCCTCATTGGCACTGCAGTCGGTGGT
GCACCAGTACGAGCCGGACGCCATGATGCCAATCCCG

17 RREN_91287_201 [A/C] CTCTGAATTCCTCGTATTATGAAAATGAGTACAGCTATTCGCAAGTCTTACCATA
CATATATTCTAATTAATAGTTTTCTTCTACCGATGTTCCCTC[A/C]CTCTCTGAATT
CCTCGTATTATGAAAATGAGTACAGCTATTCGCAAGTCTTACCATACATATATTC
TAATTAATAGTTTTCTTCTACCGATGTTCCCTCCGC

18 RREN_91287_193 [T/C] CTCTGAATTCCTCGTATTATGAAAATGAGTACAGCTATTCGCAAGTCTTACCATA
CATATATTCTAATTAATAGTTTTCTTCTACCGATGTTCCCTC[T/C]GTATTCACCTC
TCTGAATTCCTCGTATTATGAAAATGAGTACAGCTATTCGCAAGTCTTACCATAC
ATATATTCTAATTAATAGTTTTCTTCTACCGATGT

19 RREN_43396_339 [A/C] CTGAGACCACCATCGTGTAGAGCATGGTCGATAATAATAAGGAAGTGACATCCT
TTTTTGGCACAAACCCTGGTTAAATTTTGAGTGAATTTTAA[A/C]ATATTTTTT
CCACGTGCTTCAAGCACGGGTCATCGGTGCTAAAAATGTCTTTTGGTCAACAAA
GCTCAATAAGTTAAAATTAATAAAGAAAAAAAATGCAG

20 RREN_1990_6847 [A/G] TAGGTGCTCGATTTCCCGACCATCCATTATGTCCGCCGTTCCTTTTCCGCTCGAGT
GCTAGCCGGATGCTATATATTGTCCGGACTGTGTAGAGTAT[A/G]GCCAAGAAGA
TTGTGAGCAGAATGGCCAGATAGCAGAAAAGATGAGTCCAGATGCTGTTCCCCA
AGTTTTTGCAAAGATAGGCAAGCGGGTTGTGCGGCTCA

21 RREN_20709_1089 [A/C] AATAGGCCAATGCCTTTTTTCTGCTCATATGAAATTCGACATTTTTGCCTTTTTG
GTGGAGTTGGGGTGTATTCAGAAGAGCTTGATTTTTGATCG[A/C]CTTAAATAAA
GGATATTTACAAATTTAGAACATATTTTCTTACCATTTCCCTGTTCCGGATTCATCG
GAACTCTCGGATTCGCCTTCTCCATCTGACGACACT

22 RREN_258_12977 [A/G] AGTGTTCTGTTAGACAGTATAGGCAATTAGTTAGTATTTTCACCATTTGCTCTGC
ATCACCGTTCGGCTAATGGCTAGATGAAGGGATATGCTCCCC[A/G]CGGGCTTGA
ATATATGTCTGCACGGCGGTGGGATTCGAACCCACGTCCC GGGATTTAGCGGTC
CCGTGTGATAGACCACTACACCACGCCGCCGACTCTACA

23 RREN_269_9935 [A/G] GAGCCTTGCAATAGTGA ACTATGTATCAAGGGAATCAAAGA ACTAAAAAATTGG
TTGAAAAAATTTTAGCAATGGAAAAAACTTGAATAAATTGCA[A/G]AGAGAAT
CAGCTAAGATCTGGTCGGGATAAGAGTTGACAACATCTTAAATAGTAACGATTT
TTTGTCATTAGAAAGAAAATAATCACTTGTTATAAAAAGTAA

24 RREN_456_104249 [T/G] TTTAACCGTCCCATTAATTTTTAGCCGTCCCATCACAGTTTAACCGTCCCATCA
AATTTTTAACAGTCCAACCAGCTTCAATTTCCGACAAAATTA[T/G]TTTGTCAACA
GAATAGAAATATATAATCGCGGAACATGTTGAACCGGGAAGTACGATTGTGTCG
GATGGATGGCGCTCTTATGGCGGTATTAGAGCTCTACA

25 RREN_43396_325 [T/G] GGAATTCGCATATTCTGAGACCACCATCGTGTAGAGCATGGTCGATAATAATAA
GGAAGTGACATCCTTTTTTGGCACAAACCCCTGGTTAAATTTT[T/G]AGTGAATTT
TTAAAATATTTTTTCCACGTGCTTCAAGCACGGGTCATCGGTGCTAAAAATGTCT
TTTGGTCAACAAAGCTCAATAAGTTAAAATTAATAAAA

26 RREN_901_49990 [A/G] AATTCGCATATTCTGACAACACCATCGTGTAGAGAGCAATAAGTAGTAAGGAAG
TGATATCCTTTTTTGGCATAAACCCCTGCTGGTTAAAATATTA[A/G]TGAATTTTC
AAACCAAATTTTCCACATGCCTTAAGCGCGGGTCATCGGTGGGTCACAGGCCAT
TATGGTCAGCCAAATTTCAAAAAACAACTAGAGTAAAC

27 RREN_1886_12077 [C/G] GTCCAATGTGTGGAAAGGGTACATGGATGGTGGAGTCATTTCCCGGGAACACGA
ACACGTCATCAGCTCCACCAGCATCACCACCATCATTGTACAG[C/G]ATGGGATC
GCTAACAAAGAAAATTTCCCTAGATTTAACTAAAGGTAAAAAGACTCACGCTTCT
CCATCAATATCATAATGTTCCAAGACGAGGGCCTTCACCG

28 RREN_16875_158 [T/C] CAGTAACATCAACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCGCCTTAACACCGCC
GGGTCCAACAACACTGCGCCAGCCACCGGGGACATTTCGAGCCA[T/C]ATGTTCAATT
TGTTTCATTCATAACCATCTATTTCAACTGCTCAAAGCAGTAACATCAACTCTCTTCT
CTCCGTTTCGCCTCCTCCTGCCGCCTTAACACCGCCG

29 RREN_251_23034 [T/C] CGGCGGTTCCGCCAGCTTTGCCTGCCAAAAAATCGGCCAAAATGGTCGATGGACA
CAGATGGATGCAGGCATTTCGATGGGTTTGGTGTAGAGCGCCGG[T/C]CTAACACA
TTTCAGGGCGCCAGGGCAAGAGCAGTTCACCTGCTCAATCCATTGCAGAAAGGT
AGGGGAGGGGGCCATTTTTTCAGAATTGGAAGTGTAATGG

30 RREN_1895_31360 [A/T] GTAGAGAGCAAAAAGAGATTAATTA AACCTAAATTTGTCCATGCCCGACTGAG
TTGAAAAGAAAATTTATAGACACGAATAGTTGTAGATGAGGG[A/T]TAGAAGA
AATGGTGTAGTATTTTGAGGAAAAGATCGAAAGAAAACGTGAGACAAAGGGAA
ATTTTAGTTTCGAATACTTTTCTAACATCAATCAAAGGCTCT

31 RREN_9137_320 [T/C] ACGGATAGACCCATATCTATCCAAGGTCCATATTTGGATTTCAACAGACATTCCC
ACCCATATACGGATAGACCCATATCTATCCAAGGTCCATATT[T/C]GGATTTCAA
CAGGTATTCACATCCATATACGGATAGACCAATTTTCCCGCCTCTACCCCCATCC
CAAGCCTCATGCACACCCATCAAGTTCGAGCAGTACAA

APPENDIX 2. FLANKING SEQUENCE INFORMATION FOR UNTESTED SNP. THE "SNP ID" PROVIDES A COMMON NAME INCLUDING ORGANISM ABBREVIATION "RREN" *ROTYLENCHULUS RENIFORMIS*, THE REFERENCE ASSEMBLY CONTIG (FIRST NUMBER), AND THE SEQUENCE LOCATION (SECOND NUMBER) OF THE SNP POSITION.

S.N.	SNP ID	SNP	Surrounding Sequence 5'-3' (approximately 100 bp each side)
1	RREN_8_16622	[T/C]	GGCGCTCTACGACCTGTACACCCAGCCGGCCACCAAGTGCGGCCCTTCCTCGT CGGCCTCCTGCTCGGC GTGTTACCCCTCCGTCCTCCTCCTTCCGCT[T/C]CCTCCTC CCCGTCTTCCGCTTCCCTCCGCCTCCTCCCTACTCTTCTGGATCGGCTTCCTCCTTG CGCTGGGCACCATCTACGGCATTCTGCCGGAGTATTG
2	RREN_8_16659	[T/G]	TGCGGCCCTTCCTCGTCGGCCTCCTGCTCGGC GTGTTACCCCTCCGTCCTCCTCC TTCCGCTCCCTCCTCCCCGTCTTCCGCTTCCCTCCGCCTCCTCCC[T/G]ACTCTTCTG GATCGGCTTCCTCCTTGCGCTGGGCACCATCTACGGCATTCTGCCGGAGTATTGG CACCCGGACCAGGGGGTCAACCCTCTACAACACCCTC
3	RREN_12_209115	[A/C]	TGCGCTCCATTGCACATTCTAAAATAGCGAAAATGGGATGTTGTTGATGCCCTAT AAGATGGAAATTGTGTTAAATTGACCCACAACCCATGCTTTTAAG[A/C]TCTCAG TTCTAGTTACCGGTTTTAAATGGAAAATATGTAAATTATTACATTACCATCGCTA TTTATGGCAACACAAGTGCCAATCTTATTGCGAAGATAC

4 RREN_24_69296 [A/G] GGACGCGTAGAGGTCATCGACGCACGAGGGTGAGTCATTTGCATATTGTATGAC
GGATTA AAAATGAAAATTGGGCACTGATTAACCAGCTTGGAAACCATC[A/G]TGAA
TATGAAGCGAATTCCTTATTATAGAATGTTACGGGAAGAGTGAATGAACAGAGA
AAAAGAAGTCAAACAACAATATATTTTTTAACCCAGGGTTCTC

5 RREN_24_69332 [C/G] ATTTGCATATTGTATGACGGATTA AAAATGAAAATTGGGCACTGATTAACCAGCTT
GGAACCATCGTGAATATGAAGCGAATTCCTTATTATAGAATGTTA[C/G]GGGAAG
AGTGAATGAACAGAGAAAAAGAAGTCAAACAACAATATATTTTTTAACCCAGGG
TTCTCATCGAAAAATTTAAAAAATCAGGACATCATGACCAA

6 RREN_32_65082 [A/C] AGCATTATTTTCTGTATATTTTTGCTTCTTACAGGTCTACCCTGACAAGGTTTCTA
AATTTGGCTGATCAAAGTCGCGGTTGACCCACCGATGACACGTG[A/C]TTAAAGC
ACGTGGAAAAAATATTTTTAAAAATTCACTAATATTTTAGCAAGGGGTTTGTGCC
AAAAACGGATGTCACTTCTTTATTATTATCGACCATGCT

7 RREN_61_242003 [A/T] TGTAGAGCACCCCTCACAAAGCATCGTATATTTTTGGCTTTTTTTGGCCCGCTAACA
TAGTGAGGGGTTGAAAAATCGGTTTCAATTTTAAATGAACGGTTCG[A/T]TCATCC
ACGCGAATGACCTGTCCATTGGATGACATCCACCTTCAATGTAAAATAATATTT
AAACATGATTATTTTTTGCTTTATATTCATCAAATTATC

8 RREN_61_242024 [T/C] ATCGTATATTTTTGGCTTTTTTGGCCCGCTAACATAGTGAGGGGTGAAAAATCG
GTTTCAATTTTAAATGAACGGTCGTTTCATCCACGCGAATGACCTG[T/C]CCATTG
GATGACATCCACCTTTCAATGTAAAATAATATTTAAACATGATTATTTTTTGCTTT
ATATTCATCAAATTATCATATAAAAATAATTCCCGGACA

9 RREN_125_72000 [T/C] GCCGATGCCGAGAGCACTGGGCGCAAAGCCAATCAACGAAGGGAACAACCAAC
CAGGAGGCCAAATACAAAACGTACTACACCAGCACCCGCAACAACGA[T/C]CAC
TACCACAGTGGCAACCCCCACAGCAACCACTACCAAAGCTCCGGAAACCCCGAG
CACTGTCACAACCTCGCCCTCAAACCTCTCACCACAGTCACAACCT

10 RREN_125_72023 [A/G] CAAAGCCAATCAACGAAGGGAACAACCAACCAGGAGGCCAAATACAAAACGTA
CTACACCAGCACCCGCAACAACGACCACTACCACAGTGGCAACCCCC[A/G]CAG
CAACCACTACCAAAGCTCCGGAAACCCCGAGCACTGTCACAACCTCGCCCTCAA
CTCTCACCACAGTCACAACCTACAAAACCAACCGATAACCCAGG

11 RREN_125_72052 [A/C] CCAGGAGGCCAAATACAAAACGTACTACACCAGCACCCGCAACAACGACCACT
ACCACAGTGGCAACCCCCACAGCAACCACTACCAAAGCTCCGGAAAC[A/C]CCG
AGCACTGTCACAACCTCGCCCTCAAACCTCTCACCACAGTCACAACCTACAAAACCA
CCGATAACCCAGGCATCTCTACAGTCAGTCCACCAGTTGTGA

12 RREN_128_40796 [A/G] TCAATTTTTCAAGCAACAATTTACGAAATTTATTTCTTATTTGGAATTTTTTGATT
GATTTTCGCCATTTTCGTCACCTCTGCAGAATTCTTTGAGTTCA[A/G]CGCTAAAT
TTCAGTTCCTCTACACGAGACAAAGGGTTGCGAAATGGTTCTTGGACCAGCCGG
AACCAGCCGGAAACCGGAACCAGCCAGACACATTCCAT

13 RREN_159_85688 [A/T] CCAATTCACTCCTCTCTCCCTATCCTCTCTGTTCTTTCTGGTCACTCTAAATTC
TCTTTCTCCCCCTCCGCTACATATCTATCCTTCTCTCCATCCT[A/T]TACCACACTC
TCTCGTCCTCTTTCCGGTCATTCTCAATGTGTTTCTCCCCATTTTCCCCACTCTCTC
AACCTCCTCTGCATCTCTACCCGATCTCTCCA

14 RREN_190_41184 [T/C] ACGGATGATCCGCGTGGGTGAACGACCGTTCGTTTGTGTTTTGAAACCGATTTTC
TACCCTTCACTTCCTGTGCGGGCCAAAAAGCCAAAAATATACGA[T/C]GCAATG
TGAGGGTGCTCTACAAGATAGGATGGGTAAATTATGATCTGAATACTCACACC
TCGGTATGTTCCCTTGTTAGGTAATTTATAGAGGGTATAG

15 RREN_190_43474 [A/G] GAAGAGCGACGACTATCCCCTTTGTAAAGAAGATTCGTTTTCGAAATATTCTACT
GACTAACTTGTAAGAAAGTGGCGAAAGCATAAATTATATTCCCAAG[A/G]CCGA
GTAAGTTCACGTATGCTCATTTTACTTATTAGTACATATGATTCATGATTGGGAT
TACAGTATGCCATTATCTATATCCTGGACAAGCTCTAAGGT

16 RREN_190_43475 [T/C] AAGAGCGACGACTATCCCCTTTGTAAAGAAGATTTCGTTTCGAAATATTCTACTG
ACTAACTTGTAAGAAAGTGGCGAAAGCATAAATTATATTCCCAAGA[T/C]CGAGT
AAGTTCACGTATGCTCATTTTACTTATTAGTACATATGATTCATGATTGGGATTA
CAGTATGCCATTATCTATATCCTGGACAAGCTCTAAGGTG

17 RREN_202_14786 [A/G] AACTTAATCAAGAGCATCAACGGTGCAGAGCCACCAAAGAACAACAGTCCGCT
GATCTTCAATGGTGTACCGTGCCTAGACAACAGCAAGTGTGCCAACA[A/G]GCTG
AATGCCTTGTTCCACCAACGACCAACCGGCAAACCTGTCAATACTGGCCGGGCC
GCCAAACGACTAATCACCGCACAGGCCAAAGCCGCGGAGCAC

18 RREN_208_6302 [A/C] GCACCCTCATATTCCATCGTATATTTTTGGCTTTTTCTCCCCGCTCAGGAAGTGA
AGGGTAGAAAAATCGATTTTAAAAATGATCGGTCGTTCCATCCACA[A/C]GGACC
ATCCACCTATTGGATGAAATCCACCTTACTATGTATATAATAATGATTAATCATG
ACTGTTTTCTGGCTTATATTCATCCTTTTAGCTTAGCAAT

19 RREN_228_25614 [A/G] GCATGGTACGACCGACCCATTCCAAAGGGTGTGTGGATTTGCGTGGATGAACG
ACCGTTTATTTTTAAAATCAAATTTCTACCCTTCACTATTTGAGC[A/G]GGCCAA
AAAAGCCAAAAATATACGATGCAATGTGAGGGTGTCTACAAAATAGGATGGG
TTAAATTTTCGATCTGAGTGGTGACAGCTCGATATGTTCCCT

20 RREN_228_25645 [T/C] GTGTGGATTTGCGTGGATGAACGACCGTTTATTTTTAAAATCAAATTTCTACCC
TTC ACTATTTGAGCAGGCCAAAAAGCCAAAAATATACGATGCAA[T/C]GTGAG
GGTGTTC TACAAAATAGGATGGGTAAATTTTCGATCTGAGTGGTGACAGCTCGA
TATGTTCCCTTGTGAATTTTTTAAAGGAGGTGCTTAGTTGC

21 RREN_242_77265 [T/C] CGGAACTCACCAGCGAAGAGTACGAAGAAAAGTCAATATATAAAGGACGAAGA
AAGCCCCAAAAACCATCACGTGTAGAGCGGAGCAGACTCAGTTTTAT[T/C]ACAT
TTGTGAACAACATAACTGCGCCAACAACCCGCAGTGAACAAGGGTATTCCTTGA
CACAGTTTTTTTTATGTTTTTCATTTGATTTGTGGAAATTTGGA

22 RREN_242_77303 [A/G] TATAAAGGACGAAGAAAGCCCCAAAAACCATCACGTGTAGAGCGGAGCAGACT
CAGTTTTATTACATTTGTGAACAACATAACTGCGCCAACAACCCGCA[A/G]TGAA
CAAGGGTATTCCTTGACACAGTTTTTTTTATGTTTTTCATTTGATTTGTGGAAATTTG
GATAATTATAATAAGAAAATATTCACCGATTATTTCTAAT

23 RREN_269_61835 [A/T] TGTTCCTGCAGCCCATGGATTCCGATGAACAGTTTGTAGAGCCAGGCGCCGATC
AGAGCCCCGAGGAAGGGAATAGCCATTGGGATCCAGAAGTAGAAGT[A/T]GTTG
TTGCTGGGGAAGGATAAGTACAGTATAGCAAGAGAGTATGTACAGTATGCAAG
GTCAATAGTCCCAATGGAACCTAAACACTTCCCAGCCAAGTCC

24 RREN_295_80337 [A/G] CGTGGGTAAACGACCGTTCGTTTGTGTTTTGAAACCAATTTTTCTACCCTTCACTAT
CTGTGCGGGCCAAAAAGCCAAAAATATACGATGAAATGTGAGG[A/G]TGCTCT
ACAAGATAGGGTGCGTTAAATTTAGATCTGAGTGGTCACACCTCGGTATGTTCC
CTTGTCAGTACTAAAAACACTGAAAACTACTGTACTGAT

25 RREN_297_7718 [T/G] CCCGGTCCCAGGAGCTTGCCTCGTTGGGCATCCCCGACAAGACCCGCAGTCCA
TGGTGGTCTCTGCCGAGCGGATCATGTACCAGCACGCGATTGATCT[T/G]TGCCA
GTCGGCCGCTTTGGATGAGCTCTTTGGCAACCCGCAGTTGTGCCCCAAACGCTAC
CAGACCGCACACATGATGCTGCACACGCTGCTCTACACGG

26 RREN_301_61722 [T/C] GGTGATGTGGTCGTGGCATTGGCACCGTCAGTTCCAGATGGGGAAGGCGGCGTA
GAGGTGGAAGTGGTGGCAGAGGTGGTAGGGGGCACCGAAGTGGTGG[T/C]ATTG
AGGCATTCGCCAACGGAATCCCTTCGTTGTCTGTCTGGTTGACCAAAACACGG
ATCCCATTCTGAAAAAGCAAAAATCCATTGAGTTAGGGTCAC

27 RREN_308_72119 [A/T] AATATTTTCATCGATGATGTAGAGCATTGGATGTCCGATGGGGTATTGCGTTATAA
AATTGATGAAAGGGTAGAGGCTGGTGACGTCGTAGTAGGAGATTT[A/T]CTCTCC
TAACCGCAAAGCATGGAAAAGTTTAAGCGGTCCCGTTCGGCCTGAAAAATTCAA
AAGGATTTTTTAAATATTACTTTAAAAATCTCACCTCCCA

28 RREN_336_82853 [T/C] CCCCGGTAGAGAAATAGAGAAAGATTTGAGGGATTTATTTGTTTCGTGTCGAAAA
ACCGCCGATGAACGCGCGGACGACGAGTGCCACCGGCGTGGAAAAG[T/C]TGAA
AATTTGCGTGTCATGGGTTTTGTTGAAAAACAAATGTTTTGTATGGGAATTTGT
GCTTTATTCATCTTATTATTAGTATGGATTATTTGTATTCT

29 RREN_349_16421 [T/C] CACTCCGTAGAGCTGTGTCCAGTCAGCCTCCCATCAGCCAGCCAGCACATACCA
ACGGGGAGTATTAGCCTCGTCAGTCCTTCTCCCCCAGCAGCCTGTC[T/C]TGCCTC
TGCATCCGCCTTTTGTGCGCCTCGGATCGACCGAAATGAATTGAACGCGTGTC
GTGTACTTTTCCAAAGCAGCATGAACAGAGAAAGAGAAA

30 RREN_371_58464 [T/C] GACTCATAAATGGCTTGCGAATTTTGC GTTTGTACCCCTAGCTGTGGCTGCTGAA
TGTATGTTTGGCTTCGGGGTTGCTGCTGAATAATTCGCTGCGCTC[T/C]CGCCTGC
CTCACTACATGTTGCTGGGCTATTGGTTGCTGCACAACACCAGCCCTTGCTTGCC
GGCTCCTACCCATGTTATGCCCAACATATCATTGCT

31 RREN_431_1809 [A/G] AGAGTGCCAACCTTAGAGTCCCAAGTTGAAAACATGTTTTTAGTGATTTTTGA
GCTTAGGTTTCTGTTACAAAAATGTAGAGCGTATGGAAAAACATG[A/G]TGTGAT
AACAAAAAATTTTAGGCTTAGGTTAGGCCTAAGAAATTTTTTTGGGAAATTTCTA
AAATTTCCGGGACACTAAGTGGGCCTAAGGCATTTGTGC

32 RREN_432_101500 [A/G] TACATATTCATTGGCTAAAGCTGGTGCGTTTATTCAAGATTATTATTATTTTTCTG
TTTTATAAGATAATTTGATAAAATACTCACTTTGGTGTAGAGCA[A/G]AGGGAAC
ACAACACCAAAAAAGCGTTCGATGGTCAACGAAACAATGGCCATGCAACTGAT
GTAGACGGGGGTGTTGAAGAGGTACTCGGTGAGGAGGCAG

33 RREN_514_63263 [A/G] AGGTGTGGCTGAAGAGCATCTCCAACCTGTTGCCGCGGCACATCCTCAAGGCCT
CATTGGCACTGCAGTCGGTGGTGCACCAGTACGAGCCGGACGCCAT[A/G]ATGC
CAATCCCGTCATGGCAATGGGTGGACAGGTAGGGGGGCCAAGAAAATTCGCCA
AATTCGGAAGAAAATTAGACCAATTTTCCCATAAAAATCGGGA

34 RREN_521_25222 [A/C] GCTATTCTGTCCCTGCCAACTCTGGTTTGGCACCCATCGGCTCCCTTGGGGGCGC
ATTCCTCCTTGCTCACCCCTCTCTTTCTCTCTGTTCCCATCTCTT[A/C]TCTTCTCTT
CACCATCTTCTTTTTTCATTTCTCGCGAAATTGCGTTTATTTCTGCTCTCATTTCCCT
CCATCTCCACACGAGCGATCACCTCCGTCTCCAA

35 RREN_523_12130 [T/C] TTCCATCTCCAAGTTGTTTATAGAGATGTTTGCCGAGTTCAGTGGGATTCGTGAT
ATTCGAAGGGGGCACTGTTACACGCACCTCCTGCCCTGTTGCGCA[T/C]ACAAC
CGCATATATTGCTGTTCCATTGTTCCCAGGGAAGGCCACGAATGGGGATAAACC
ATCACTGCAAGCCCCACATAACCAATCTGAATTGAATTGCA

36 RREN_523_13494 [T/C] TGACTTCTTTCTGATGTTTTTTGCCTCCGTAAACAACGGGCTGACACTGGTAAAT
GCTACTGGTGAATGTATATCCTTGTAGAGACGCTCCAATAGACGC[T/C]TTATGG
ATGCCTTCATTGCACGTTACCGCCACAATGATGGTTTAAAAGAAAACTGTTACT
CACTACTCGACTTTTGGTTACGTTTTTGTTCACCTTTTT

37 RREN_526_81331 [A/C] TCTGCTGTAAGCTATATTATCATTTTAACTCATAAAATTAATTTTCAACCGTTTTT
ATGGACCATTAGATTTTAGTGTTTTTGCATGGTACGACCGACGC[A/C]ATTCCAA
AGGGTGTGTGGATTTGCGTGGATGAACGACCGTTTATTTTTAAAATCAAATTTTC
TACCCTTCATTATTTGAGCGGGCCAAAAAGGCCAAAAA

38 RREN_526_81371 [T/C] ATTTTCAACCGTTTTTATGGACCATTAGATTTTAGTGTTTTTGCATGGTACGACCG
ACGCAATTCCAAAGGGTGTGTGGATTTGCGTGGATGAACGACCG[T/C]TTATTTT
TAAAATCAAATTTTCTACCCTTCATTATTTGAGCGGGCCAAAAAGGCCAAAAAC
ATACGATGCAATGTGGGGGTGCTCTACAAGATAGGATGG

39 RREN_526_81423 [A/G] ACCGACGCAATTCCAAAGGGTGTGTGGATTTGCGTGGATGAACGACCGTTTATT
TTTAAAATCAAATTTTCTACCCTTCATTATTTGAGCGGGCCAAAAA[A/G]GCCAA
AAACATACGATGCAATGTGGGGGTGCTCTACAAGATAGGATGGGTTTAAATTCG
ATCTGAGTGGTGACAGCTCGATATGTTCCCTTGTAAGAACT

40 RREN_526_81432 [T/C] ATTCCAAAGGGTGTGTGGATTTGCGTGGATGAACGACCGTTTATTTTTAAAATCA
AATTTTCTACCCTTCATTATTTGAGCGGGCCAAAAAGGCCAAAAA[T/C]ATACGA
TGCAATGTGGGGGTGCTCTACAAGATAGGATGGGTTTAATTTTCGATCTGAGTGG
TGACAGCTCGATATGTTCCCTTGTAAGAACTTTATTTTCT

41 RREN_526_81448 [A/G] GGATTTGCGTGGATGAACGACCGTTTATTTTTAAAATCAAATTTTCTACCCTTCA
TTATTTGAGCGGGCCAAAAAGGCCAAAAACATACGATGCAATGTG[A/G]GGGTG
CTCTACAAGATAGGATGGGTTTAATTTTCGATCTGAGTGGTGACAGCTCGATATGT
TCCCTTGTAAGAACTTTATTTTCTCTAATACTCAAAAATC

42 RREN_558_68223 [T/C] ATTCTGTCATTTCTTATGCAATTCCCTCCTCATTGTCAACTATATAAGCAATGCA
TTTTCAATCATTTGTCACTTCCATTCCCATTCCAGCTCCCATTT[T/C]CCTTATTCC
AATTCATCTATTGTCCTATATTGTCTTCAATAAATTCTTCACGAGGACACAACAA
TTTGGCGCAGTCACGAAAACGACTCTACGCAATGCC

43 RREN_590_31647 [T/C] TCTTACAATCAAATCTGCCCTTGGGAGCTTTTTATGGAAATTTTTTCAAGGAAA
ATCATGAAACTGGACAATAAGATTAAAACTCTGCCAGAACCACCA[T/C]CTGGCC
AAGTTTGCGCTGCGGCGTCGTGACGGCGACTTTTTGTCCATCGAAGCAGCATTGT
TCTTGGCCGAACAGGGCGCCGCTGTGGTGGTGGAAACATC

44 RREN_625_19284 [T/C] ACATCTGCATTTGGGGAAGGGGGCGAATTTTGTAATAATAGGCAAAAATCGAAG
GTGAACAAGGGGGGTAGGAGATGGATGTTTTTCATCATCCATTACA[T/C]CCAAC
ACATCCAATTCTGAAAAATGGCCCCCTCCCCAACTTTTCTGCAGTGGATCGAACA
GCTGAATTGCTCATGCCCTGATGGTCTGAAGTGTGTTCGA

45 RREN_718_5249 [A/G] ACAAGGCATGTAGAGGACGACGCAGAAGGGACCAATTGTACAGAATATTGATG
ATCAAAGCCCCTAGCACGCCACCCTCCGAACCAGCAGGACCCACCTC[A/G]GGC
TGAACAAAAGGATTAGAGTAAATAAAGAACTGGAATGAGTGTAAATAATACCAT
GTAGGGAACAAAAACGGCACTGGCCAAATATCCGCCAATTCCGG

46 RREN_721_22152 [A/G] AAAC TTTCGTATCGGATTCTCCTGCTGCTGTCCGTCAATTCCTGAACGACCCAAA
AATTGAAGTGGACTTTATAGAACA ACTAAACGAGGAATGCGTCCT[A/G]ATCCG
ATACACACCACTAAAAGAATGGATCGAGGAGCACA ACTGTTCAAACATTGTGCT
CTCTCTATGGACAACGGCGGCGGCACGACTACATCTCCTCA

47 RREN_721_22224 [A/G] TAGAACAACTAAACGAGGAATGCGTCCTAATCCGATACACACCACTAAAAGAAT
GGATCGAGGAGCACA ACTGTTCAAACATTGTGCTCTCTCTATGGAC[A/G]ACGGC
GGCGGCACGACTACATCTCCTCAAGCTAATGCAAAAAGTGGCGTCCACTCCCAA
CTGTGTACTCCTCTACACGGACACGGACAGTTTGATTTTCG

48 RREN_780_29960 [A/G] TCCACACAATATAAGCACTTGGCCAAGGTCAAAATTTTCATTTTTACTCTAAATT
TTTTTCTGACAATTTTAAACCCTTCAGCTTCTCGCCCCTCCTTCC[A/G]CCCACAG
CGAACCATTTGATGTGCCACGTCCCTTTCCATGTCCATTTGTGCGATTCATGCCGCG
CTTTGTGCTCTTTTGGAGTTGTCCATGTCCGAGTCCGC

49 RREN_800_50912 [T/C] ACAAGGGAAGGACAAAACATGTTAATCGTGGATCAATTCGAGACCCGTATAATA
TTAGAGCATCCGCAACTTCCGTCTTAAATTTTAATATTAGCTGCCC[T/C]CGTCCA
CAAGAAAAAAAAAGTTATGAATATTTATTTTCATAATCCGCGATCCGCGCGGGTCA
AGGGAGTGACATACCTATTCAGATTTCTAAAAATTACGTC

50 RREN_929_9795 [A/C] CGGAGGCGCGAGTTTGGGCTAGTACCGTAAAAATGGGAAAAAAAAATAGAGTTTA
CGTTCCGAAAATAAATTGGGCCAAATTATATACCATAGTGTAGAGCT[A/C]GACG
AGCTGAGTACGAATATGTAATTATTTTTTTGGCGCAAACCACTTTAAACCGGTTTT
TGAACCCTTTCAAGTTCTTATCCAAAAATGCAAAACACTTG

51 RREN_981_25656 [T/C] CATCCAGCGGTTTGTTCGGAGCGTAAATCCGTATTCCTTCGCGACTTGTGCACCT
GGACACGGATGTGTAGAGTTGTCCATGAGCGAAAGGTTCTTCGCT[T/C]AGGTCG
ATCCCAGCCTTTCGATGGTCTGACCTTGGGCCTTCGCGAATGTCATCGCGAAGG
CTACCCGTACCGGGAACCTGGAATCGTTCAAAGGGCACAT

52 RREN_1012_21981 [T/C] AACTTTTCTCTCAAATCAGCAAAAATCAAAAAACATGGCAAAAAGCATGGCAAA
TTTTAAAAACATGGCATTGTCATGTATACATGGTAATTTGGCCT[T/C]CCTGTT
CAAAAATCCGATATAAAAGTGACCCCTTCTTCCGAAAATAATTCATTTCTTCGCG
GGCATCTTCGTTGTTTCTACTTCTACATCTCTACACAAC

53 RREN_1120_48434 [C/G] TTTTACTACTTTTTCTTTTGATTTTATGCATTTTGGCAAAGTGTGCAAGTGAAAT
TGCACAGAATTTAAAAGGGTTCAAAAACCGGTTTAAAGTGGTTG[C/G]CGCCAA
AAAATATTTATATATTCGTGCTCAGCTCGACGAGCTCTACACGATGGTATATAAT
TTTGGTCAGTTTGACTTCCGGAACATAAACTGCATTTTT

54 RREN_1123_57666 [A/C] NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNTGGG
AAGGTAAGCGGAAGGGTTTGGGAAGGTAAGCGGAAGGGTCTGGGAAGGA[A/C]
GCGGAAGGGTTTGGGAAGGTAAGCGGAAGGGTCTGGGAAGGCTGTGTGGACGA
ATTTTGGTAGCGCCGAAAGTCTGGATCTCCCTAAAAAAGAACGGACC

55 RREN_1175_25864 [T/G] TACCAATTTTTCGATAATTTAACAAAAATGCTGTCATTTTGGATATGCATGCATG
TGTATGCGCGTAGAGTGCGCGCGTGTGTGACAAGGGTATGCGCGT[T/G]ACAAA
GAAGTACAAATTCACCATGTTAGTTGGTCAGTATCTTGGCTAAGTTGGTCTCGTC
CTTGGTTAAATGCATCATCGTCGTCGTCTGAATTTCCCT

56 RREN_1187_12082 [A/C] TATTATCTCTTATAAATTAATTGCCAAACATTTTTATGACATTTACATTAAGTG
 TTTATGCATGGTACAATCGACGTCATCCCAAAGGGCGAATGATC[A/C]GCGTGGA
 TGAACGACCGTTCATTTTTAAAACCGATTTTTTCGACCCCTCACTATCTGAGCGGG
 CAGAAAAGCCAAAAATATACGATGTAATGTGAGGATG

57 RREN_1187_25663 [A/G] GTGCCAAAGACGGCCGTACCAGCAAACGAGACCATTGTGCCTCCGCCCGCATCC
 GATCCCTCTTCTTCTTCCATTGCGACAGTGCAATCCGCTGGAACAC[A/G]CTGGC
 ACCCATCACGACCTTCACCAACCTCGGCCCTTCCGCTCTGCACTCGGCCTTCCC
 ATCCGATACACAGTGACTGGCATCGCTTCCACGCCCAATT

58 RREN_1215_1686 [A/C] TCGAATTCAATTTAAGATTCAATAAATTAGAATACCAAAAAACCTCTAGAACT
 TTCATAAAGCTAATTATAAGCAGATAAACAATATATTCAAAAATAC[A/C]GACTG
 GAATACTCTCTTTTCAAATAGTATAGAAATTGATCATCTATATCAAACATTTAGC
 CACAATATCCATAAAACCATCGAAGATCATATTCCTATCA

59 RREN_1572_36973 [A/C] TCTTTCGAATATTTTGGACTTTTTTGGACACCTTATGATTGACCATTTACAGCCCC
 ATCCCGCTGGCCAAGCGGTCTCCTACTCGGCCAGCAGAAGAAC[A/C]TGTTGAT
 GTGGGCGGTGGCCGTCGGCTCCATGCTCGGCACTTTCCCCTTCGCTGGCTCTAC
 ACCCGGCACGGTGCCCGCTGGGTTCTGTTCGGTGCCGG

60 RREN_1660_519 [A/T] TGCGGTAGTCGGTTCGGCTTATGGCGTAGAGCTGATCAGTGAGAACCAGATCCC
GTCCGCACTCCAAGACATATTCCAAATTGGTGATGCTTAATGAAAA[A/T]GTTTG
ATCTTCTGACATGAACCAATCATCAAACATGGTTCCTTTCATTTCCCTGCATCAAC
ACGTACCGGCATTGTGTGTCCGCTTTGGGGCGGTTCCATA

61 RREN_1695_18038 [A/C] AAAAAGTAATATTGTGCTGAATTTTATGCTCTATCTTCTGGGATTTATAATTCGG
CCAAAAAATTGGAAATATCCCCTAAAACCTTATTTTTTCAGAGTAA[A/C]TTTTTG
GTAGAATTTAAATAAATAATAGAAGATTCTGCACAATGGCTTTTTTATAGTTTTTG
GCCCTAGCACCGATCAACCCTCTACACGGTATACCATT

62 RREN_1695_18063 [A/C] ATGCTCTATCTTCTGGGATTTATAATTCGGCCAAAAAATTGGAAATATCCCCTAA
AACCTTATTTTTTCAGAGTAACTTTTTGGTAGAATTTAAATAAATA[A/C]TAGAAG
ATTCTGCACAATGGCTTTTTTATAGTTTTTGGCCCTAGCACCGATCAACCCTCTAC
ACGGTATACCATTTTAGTGGCGGACCAATTAATTGTGGT

63 RREN_1721_18343 [T/C] GGCCCCAAATGTTCCCTGAATATGTACACGCCTCTCTCAATTTATAGTTTAGGTG
TACCCTCGCGATTAAATTGTAAATACGCCCATGGCCACCGTCCTC[T/C]CTTTTCC
TACACACATCTTGATAGAATATTTTCAATAGCTCTATTTTCAAAGTTGTGATAAA
TAATCAACATTATTTCCCTATGATCAATATTCTCACTC

64 RREN_1741_30669 [T/C] TCCCATTCCTGCACTTTTTTTTATATTTTCTGTTCCCTCAATTACTTTTCTTCGCAAT
 CCAAATCCGTATTTAAGGGTGTTC AATTTCTCTTTTTGCTCA[T/C]TTAAACTTG
 GACTTATCTTCACTGTCCCTCACCCAATTTACTAATTCGCCGTCCCAGCTCTACA
 CTATTCACTTTTGGTGGGTACACATCCCATTTTCGAT

65 RREN_1874_61680 [A/G] CTATTGATTTTAAGCATTTTTGGTTGTGTGCCAGTGAAATTTGCGTGGA ACTAAA
 GGGGTTCAAAAACCGGTTTAAAGTGGTTTGCGCCAAAATATAATT[A/G]CATATT
 CGTACTCAGCTCGACGAGCTCTACACGATGGTATATAATTTGGGCCCGATTTATT
 TTCGGAACGTAAACTGCATTTTTTACCCAGAATTTTATGC

66 RREN_1886_8608 [T/C] TAATTTGTAAGGCTCCCGCAGGATTTTCGCATCAAAACAAGATAATGCGAATTAT
 TTCGTGCGACCTTCAATTCTTTCGTGAACAGATGCTGAGTGACTAA[T/C]ACCAC
 ACTCATACCCCAATTGTGGGAACCCCGAGTGAAA ACTGTGTCTAAAAATTGCTG
 ACGCATCCCGACCATCAAATCATCCAATACAACCAGCAGAT

67 RREN_1886_12029 [T/C] GTCTATAATTTTTATGGGCCTCATAGGTGTAGAGATCGGCCAGCGGATTGCGTCC
 AATGTGTGGAAAGGGTACATGGATGGTGGAGTCATTTCCCGGGAA[T/C]ACGAA
 CACGTCATCAGCTCCACCAGCATCACCACCATCATTGTACAGCATGGGATCGCT
 AACAAAGAAAATTTCTAGATTTAACTAAAGGTAAAAAGAC

68 RREN_1915_307 [T/C] CGTCTAGAGACTCAATCGTGGAGACTCAATCGTGGAGACTCAATCGTAGAGACT
CAATCGACCCCCACCGGATAAACGACGCTCTTCATCTTTTGCCAAC[T/C]CAAAT
GTGAAAGATAAGCAGAAACTGAGAGAAACAAAACGGTAGAGAAAAGAGTATG
AGAGAGAACATAAACGATAAAGAAAGAGTTCGTTTCGGAGGCT

69 RREN_1928_25815 [T/G] CCATAGGTTATTTTTCGTTCTCTATTTTAAATTTTATTAATCACCCATTGCCTTATT
ATTCAAATTTTACATTTTCTAAATACAAAAAATTAAGATGG[T/G]TTTCCGCTC
TAAATTGTGCAATTTTTTATTATGTGAAATTTATCCGCGCTTCAATATTCATCATC
TAAGGCATTTCTCTACACATCCTCTCCAACCACA

70 RREN_1930_18838 [A/T] ATTTTGTGCATTTTGGAAAAGTGTACCAGTGAAATTGTACAGAACTTAAAAGG
GTTCAAACCCAGGTTTAAAGTGGTTGCCGCCAAAAAATATTTACAT[A/T]TTCGT
GCTCAGCTCGACGATCTCTACACGTTGGCAGATAATTTTGCCCAGTTTGGCTTCC
GGAACATAAACTGCATTTTTTCCAAAAACAAAACATCCTT

71 RREN_2082_17935 [A/C] CATTCTTTACAACACTATCACAAAATGGTGTCTATTGAACATTATTAAGAATTCGA
CTAAGGGTATCGACACAGGATAAGCACCGCTTAAACCGGTGAATG[A/C]CCATTT
TTTCGTTATACGGTATCGATTTAATGCATGGTACAGTAACCAATTCAATACCACA
ATTTTTGTAACGTAACCTTCTCTACACTATCCCCAATGAC

72 RREN_2082_17995 [T/C] GGTATCGACACAGGATAAGCACCGCTTAAACCGGTGAATGCCCATTTTTTCGTT
 ATACGGTATCGATTTAATGCATGGTACAGTAACCAATTCAATACCA[T/C]AATTT
 TTGTAACGTAACCTTCTCTACACTATCCCCAATGACCTTCTACCGTATGCTCCTTTT
 ATCGTGTCGTTTAGGATGACTGCATACCATATTCCAGCT

73 RREN_2193_23293 [C/G] ATTGGCGACAGAGAAGAAGGCAGCGACTGTGGGGAGTTCGGCGACAGAGAAAA
 AACCAGCGACAATGACAGAGAAGCTTCTGGAGAAGAAGAATGGATTC[C/G]AGA
 CACAGCTTTAAGTAGTAAGAATGGTGCCACATAAATCATGCATAAAAAACCGGT
 ATCGGGGCAACCGAATAGTGGTCATGGACACCAAGAAATATGA

74 RREN_2229_14581 [A/G] ATGAGGTGAGTCATAAAAGTTTTAACGATGGGTTGATTGGGTGTTATTGGACAC
 CATCAGAGGTAAAGGAGGACACGTGTACAGCCAACGCTATGTCGCA[A/G]AGAA
 ACCGGATGTCCCACCCAAGGATGTGTATCAGCTATCGGTATGGAATAAGAGGTG
 GAGATGAATGGATCAAAGATCTATTTGGACCGGAAGAACAGC

75 RREN_2245_19256 [A/G] ATTCCGGTGCCCCACGGAAGAAGTTATTCAAAGACAGCCCAAACCCGCACTGG
 TCATCCTTGATGATCTCCTTTATTCCATCGATCTCAAGTTTCTGGC[A/G]GATCTC
 TACACCAAGAAGAGTCACCATGGTAATTTTGGAAATCGTTATGCTCACTCAGGAT
 TTGTTTGATCGTAAGATGAAGGTGGTTAGGCAAAATTCAA

76 RREN_2295_24784 [A/G] TGCCTATCCACGTGTCCAAGCAGCTACTGCGGCTTTGGAAGCAATCCTTGAATG
GCTGACCAACAACCCACAGTCTTCTGCGGTTGAAAAGATCACACTT[A/G]TGGTC
TCTAATCCAAATGACCAAGGCTCTACAAAGATCTACTTCAACGGGCTAAGCGT
CAAATTGTAGGCTCCAGAACAGCAAGTCGCGCATCTTCCAG

77 RREN_2301_5490 [A/G] TTGAATGCAAGAAACAATAGACAATGGCGAGCTGGGATGATAGCCAAGTCCA
AGAATGAGGGAAAATCACTAGCTTATATACATGCAGGCGTGAAGGAA[A/G]GGG
AAGGAAGAAAGAGAGAAAACAATACAATAGCGACAGACAAATGTGAAAGGGA
ACATAATGCCAAACAATACGAATGAACAATACATTTGAATCACA

78 RREN_2380_20166 [T/C] ACGGCAAGGTCTCCATTCCCGGTCTTATCCGCCCTTTCTCATAGTGTGGTCTATA
ATTTTTGTGGGCCTGATAGGTGTAGAGATCAGACACTGGAGTTCG[T/C]CCGATC
TGTGGAAACGGAACATTGATGGAAGAGTCGTTGCCCGCAAATACAAACACATCA
TCATCCGCGTTAAAATTGTCCAGAGCAGGATCGCTAAATT

79 RREN_2380_20294 [A/C] GAAGAGTCGTTGCCCGCAAATACAAACACATCATCATCCGCGTTAAAATTGTCC
AGAGCAGGATCGCTAAATTGGTCACGAAATTTTATACCCAGAATAT[A/C]TACAC
CCACATTTGACCATCGATGTCATAATGTTCCAAGACAATCGCTTTCCTGTTTCG
AAATTGAGCAATTGATGGTTGGCATAGGAGAGGCGAAAGC

80 RREN_2496_36896 [T/G] ATCACATTGCATCGTATATTTTTGGCTTTTTTGTCTCGCACAGATAGTGAAGAGT
AGAAAAATCGGTTTTAAAAATAAACGAATGGTCGTTTACCCACGC[T/G]GATCAT
CCGTCCGTTGAATGATAGCTATCTTACAATGAATAAAATTTTTTAACAATGACTA
TTTTCTGGTTTTATATTCATCATTATATCATATAGAATA

81 RREN_2611_11791 [A/T] AAAATTGTAGAGCGTGTGTCAAACATAAAGTATGATCAAAAAATTTTGAGGTT
AGGTTAGGCCTAAGAAAAATTTTCGGGAAATTTTCGAAAAATTTTC[A/T]GGTTT
ATTGGAGGCCTAATAGAAATGTGGTTTGTAAATTGATATTTTGAGCTGATTTTTTG
TACTCAGGGGTTTTTCGAGGGTGCTGAATCCGAATATGACA

82 RREN_2644_249 [T/C] CATTCTCCGACTTGAAGGGAACGGGCGCGTTTGTCCGGTATGTGGCCATCGAACC
TAAAAATAATTTATACTAAATAAGTATAGTATTTTAATCCAAAAGA[T/C]GAACC
TAAAAATAATTTATACTAAATAAGTATAGTATTTTAATCCAAAAGAAAGCTAAC
CGAATGTTGACATACCTATCCTGAAGCCGCCAAAAACCCTG

83 RREN_3007_13959 [A/G] TTTCGGTTTTAGGGCCAAAACACTGATTTTCGGTTTTTAAAATTTGACGGCAGC
GTGGTGTAGTGGTCTATCACACAGGACACAAATTCTGGGACGTGG[A/G]TTCGAA
TCCCACCGCTGTGTTGGTCATATACTCAAGCCCGTGGGGAGCATATCCCTTCACG
TAGCCATTAGCCGAGCGGTGATGCAAAGCAAATGGTGAA

84 RREN_3315_26470 [A/C] GAGACTGCGGATGAGGAAGAGGGTGTGGAGGATGGAGCTGGGGGTGATGTACT
GCTTGGAGATGATGGATTTGAAGAATTTAATGGTGCAGAGGAAGATG[A/C]GGA
GGTGGAAATTGGAAGGGATGGAAGATGAGGGGGAGGAAGATGAAGAGGAGGAT
GAAGACACGTTTGACGCTGGCAGCGAGCTGGAGTCCAGGGCCCAG

85 RREN_3415_20865 [T/C] GCATGGGTCTCTGGAGGTGTAGAGCGTCAAAGCGGCCGCGTTTTTTATGGAAAT
TGTTGAAAAAAGGTTTTTCCGCGCGAATTTAAATGTGAATCTTTT[T/C]TGATTA
GGGACAATGCGACACGTATGGCATTGATTAGGAAGTATATCAATCCCGGTACCA
CTATTCATTCTGATTGTTAGAAAGCGTATGCAAACATGGA

86 RREN_3722_17382 [A/C] CTCGCGTAACTCGTTCCGGCATCCGAGCATTATAAATAGTGAAAGGTGGCATAT
CCAACGGACGACATGATGAAGCATGGGTTTGCCTTACCCATCTGAT[A/C]ACCTA
GAGAAGGTTAAAAGGGAAAGGATATATTAGAGGAAAAGTTGAAAACCTTACATT
TTCTAAACCCTCTTCCGAGTCGGGAATTGTCATTTGAAAGC

87 RREN_3853_3705 [A/G] AACACTTTGCGAGCCTCTATCCACAAACGCTGTACTCCAGCAAAGCTGATGGG
GATTTTGGGTCATTGTAGAGTCGATCCAAGGTAGCCTTCACTGTGG[A/G]CCTAG
AATTATAAAGTAATTCAAGGTTATTTTTCTCGAATGAAAAAATACCTAAATTTAT
TTCAGTAAATTAAGGTTAGCTTTTTCTAAAGAAAAATCTT

88 RREN_4099_6882 [A/T] TTGGGTGTACGGGGGGATGTAGAGGTGCAGGCTGACGGCCGGATCGGCCGTGGG
AAGGGTTTTCCATCCGGTGCAGGCCGATCTTGTCTGTGAGATCAAAG[A/T]TCAG
TGATCAATATCAGCCGGCCTTTTTTCTCATTACCTATTTATATTCTGTGGGGATAT
GACTTTTATTTAAATATTATTTAAAGCGGATACCAAAGTA

89 RREN_4280_20028 [T/C] CACGTAGAGCTGTGTGTGAATTCGGGATTGAGCTAGTCGGGATTCGGGATTTC
GAGATTCGGAATTTTGCCATTGTTGATCTCTAACCTCGACTATTT[T/C]GATTTG
GAAACTGCGGAATTTCCCAATTA AAAATATGACAATTTTGGAGGCATATAGGT
AGTAGTGTGGAGTGAGTGAAGTAGGCTTCCTCACGATGCT

90 RREN_4396_2332 [T/C] AACTAGATTAAGAAAATAGTGGGGTAGAGAAAAAGAGAGACGAAATAATAGG
TAGAGAACTAGAGTAAGAAAATAGTGGGGTAGAGAAAACAGAGGGA[T/C]GA
AATAATAGGTAGAGAATAGAGAAAACAGAAGACAGCGAAAGAAACAAATATA
GATTAACGAGAGAAAGCAGAGAAAAAATATAGCCTTAGGCATGTAC

91 RREN_4396_2346 [A/G] AATAGTGGGGTAGAGAAAAAGAGAGACGAAATAATAGGTAGAGAACTAGA
GTAAGAAAATAGTGGGGTAGAGAAAACAGAGGGACGAAATAATAGGTA[A/G]A
GAATAGAGAAAACAGAAGACAGCGAAAGAAACAAATATAGATTAACGAGAGA
AAGCAGAGAAAAAATATAGCCTTAGGCATGTACAATGACCACCAACC

92 RREN_4491_8860 [A/T] GACGAAACACCTAATTTGAAATGTTAAACCTTTTATTGTATTAAAGTTGCTGGGC
AACCTTGCATAACTTGATGCAATTTTATTAATTACGTGTTTTTTT[A/T]ACTTCATT
TTACTGCGTTTTCTATTGATTTTATGCATTTTTGGTTATTTTTTCAGTGAAATTTGC
GTAAAGTAAATGGGTTCAAAAACCGGTTTAAAGTG

93 RREN_4741_9023 [C/G] AGCTCTTCGGGAATAGACAAAACCCGAAAACGTAAATATGACACCGAAGAAGT
CAATGGGATGCCAAATGACGAGGGTCCATGGCTCAGCAAAGCGGGTG[C/G]TGG
AGTGAACAAAAAGCTGCTTTGGATGCTTTGCAAAAACAAGTATCAACTCGGATT
CGATTTGTGCACCGACTGCAATCAGGCCCTCTACAAGAAGAAG

94 RREN_4834_4812 [A/G] ATTTTCGTCTCTTTTTTCTCTACCCACTTATTTCTTACTCTAGTTTCTCTACCTC
TTATTTTCGTCTCTTTTTTCGCTAACCCAGTATATTCTTACTCT[A/G]GTTCTCTACC
TATTATTTTCATCTCTCCGTTTTCTCTGTCCCACTATTTTCTTACTCTAGCTTCTCTA
CCCCACTATTTTCTTTCTCTAGTTTCTCTACCT

95 RREN_4834_7584 [T/C] GTGCACTTTGGCACCTTAACTCTGCGTTTGGTTCATCCACATCGCCAGTTCTGC
TTACCAAAAATGGCCCACTTGGAGCTTCAGCATTCAATGCCTGGG[T/C]TCACAG
AGAGTCAAGCAACCCTGGCTTCATACCCATTTAGAGTTTGAGAATAGGTTAAGG
ACATTTTCGTCCCAAGTCCTCTAATCATTTCGCTTTACCGA

96 RREN_4985_4041 [A/T] TCCAAAAACATGAAAATAATAAATGTAATAATTAAGAAAAAATATGTA
CACATAGGCTGACGGGGACAATTACACAATCGGATTTTTGTACAAGA[A/T]AATA
CCCAAGCATTGGACAAGTATAATTTGAATATCGGTTTGGACAATAACCAAGAA
TTTGGCCTGAAGTTGCAAGGCGATGAGTGGTGAATAAAACT

97 RREN_5033_11813 [A/G] GTGGAAAGGGCACATGGATGCTCGAGTCATTTCCCGGAAACACGAATACATCAT
CAGCTCCACCAGCATCACCACCATCATTGTACAGCATGGGATCGCT[A/G]ACAAA
GAAAATTCCTAGATTTAACTAAAGGTAACAAACTCACGCTTCTCCATCAATATC
ATAATGTGCCAAGACGAGGGCCTTCACCGTTTCAAATTC

98 RREN_5033_11816 [A/G] GAAAGGGCACATGGATGCTCGAGTCATTTCCCGGAAACACGAATACATCATCAG
CTCCACCAGCATCACCACCATCATTGTACAGCATGGGATCGCTGAC[A/G]AAGAA
AATTCCTAGATTTAACTAAAGGTAACAAACTCACGCTTCTCCATCAATATCATA
ATGTGCCAAGACGAGGGCCTTCACCGTTTCAAATTC AAC

99 RREN_5385_4749 [A/G] CTGTAACATGTTACCTTGCGTGCTGACAAGGGAACATACCGAGGTGTGACCACT
CAGATCTTAATTTAACCATTCTATCTTGTAGAGCACCTCACATT[A/G]CATCGT
ATATTTTTGGCTTTTTTGGCCCGCACAGATAGTGAAGGGTAGAAAAATCGGTTTC
AAAAATAAACGGTCGTTTCATCCACGCGGATCATTCGCCC

100 RREN_5385_4848 [A/C] TGCATCGTATATTTTTGGCTTTTTTGGCCCGCACAGATAGTGAAGGGTAGAAAAA
TCGGTTTCAAAAATAAACGGTTCGTTTCATCCACGCGGATCATTTCGC[A/C]CTTTGG
AATGGCGTCGATCGTACCATGCATAAACACAAAAATGTAATTCTCCATAATTTG
GGTTGGAAATTATTCTATATGATATAACGATGAATATAAA

101 RREN_5497_2602 [A/G] TGTCGGCTTGTCCCCTGCTCAATCGCTGCTACCGGCGAATAATTCCGTGTAGAGG
GGCGGAAAGGCACTGAGCACATTTTGC GGATGAAAATTCTGTAAG[A/G]AAGTG
TAAAAACAAATTGAATTTTGAAGCTTTGGATATATTCAAAAAAATTTAAACTAA
TAACTTATCAGAACAAGACGAGGAAAATGAAGAATGAAAT

102 RREN_5940_3890 [T/G] AGACTGTAGAGACCGAAGAGGAGGCAGAGGAGAGAGTGGCAGAGCAGGTAGG
CGAACTTGTGGCTGATCATTATCTTTTTTGTGTCTGAAGGGGACAATCA[T/G]CGA
AGATATTCTGGACAAGGACAAGACAGAACAGACTGGAAGGGCAATGCAACTGC
AAGAAGCCGATCAGCGGCGGCACTGAACAACGCCATCATGTTGG

103 RREN_5940_3923 [A/T] GAGTGGCAGAGCAGGTAGGCGAACTTGTGGCTGATCATTATCTTTTTTGTGTCTGA
AGGGGACAATCAGCGAAGATATTCTGGACAAGGACAAGACAGAAC[A/T]GACTG
GAAGGGCAATGCAACTGCAAGAAGCCGATCAGCGGCGGCACTGAACAACGCCA
TCATGTTGGCATTGGCACTGAACAGCATAGGCTGAAGGAGGG

104 RREN_6983_4756 [A/G] GACAAGGGAACATACCGAGGTGTGACTATTCAGATCATAATTTAACCCATCCTA
TCTTGTAGAGCACCTCACATTGCATCGTATATTTTTGGCTTTTTT[A/G]GCCCCG
ACAAATAGTGAAGGGTAGAAAATTCGTTTTTAAAAATAAACGGTCGTTTCATCCA
CGCGGATCATTGCCCCTTTGGAATGGCGTCGATCGTACCA

105 RREN_7324_5103 [T/C] GCAGTGTGGCAACCAGGTGTTCTCTCAAAGAGACAGAGAAGCAGATAATTATC
GATTTTTTCCGAGGGTCAGAGACGCAGAAAGGACGCCAACTTTTGG[T/C]GGGA
AAAGCGGCGTCACCTTTCCTGCCAGCCACCTCTCTACACTCTCTGTAAGACAGAG
AGCGAAATTTATCGATTGATTTTTGGGTTAGAGACGCAGAGG

106 RREN_7711_4765 [T/C] GTGAGAGAGTGTAGAGTGGTGAAGTGGAGGTGAGAGAGTACGGTATGTGAGAG
AGTACGGTATGTGACAGTACCGTGTTGTCCACGACCACGACACACTC[T/C]TTGT
TGGTTGCCTTCACAACTCGGCAAACGGCCTCAATATCGATGACTTTCAGCAATG
GGTTGGACGGTGTTCGAACCAGACCATCTATGGATGATTAT

107 RREN_7711_4768 [A/G] AGAGAGTGTAGAGTGGTGAAGTGGAGGTGAGAGAGTACGGTATGTGAGAGAGT
ACGGTATGTGACAGTACCGTGTTGTCCACGACCACGACACACTCTTT[A/G]TTGG
TTGCCTTCACAACTCGGCAAACGGCCTCAATATCGATGACTTTCAGCAATGGGTT
GGACGGTGTTCGAACCAGACCATCTATGGATGATTATGTA

108 RREN_8907_4239 [A/G] GTGAACAGAACTTTAAATATGGCGGAGATCGAGTTGACAAAGAAACGCTTAAA
AAACTTGACAAAATGCTCAGGAAACACCATCCTTTGGCAAAGAATT[A/G]ATG
AATTTCCACACACAATACCAGCGGGAATTAGCTCTAAACGGACCTGATGCCGTT
GCAAACCTACCGTTTCACGATTCTCGAGGCACGTGATGCACCGA

109 RREN_9458_2923 [T/C] GGAATGGACAGCAGAAAGTTGGATGGCGGAAATGGACGGGAGAGAAAGTGAA
GAGCAGAGGTGGACGGCGGACATGAGCGGCGGATTTAAACGGCGGAAG[T/C]G
AACGGCAGGTGTCGGTGGCTGACGGCGGATAAGTGAACAGCGGAAATAGACGG
TGGAGGGTTGACGGTGGAGCTTGACGGCGGACGTGGACAGGACCGA

110 RREN_10201_2603 [A/T] AATGTTTCGAAACAATAGGGAATGGATCTCACGATCCCTCGACTTTGAAGGGTAC
TTTGGTCAGGGGACCAGGGATTAGCCCCTATTTAAAGCCGCTCCAA[A/T]AATGA
GAGAGGCATGTTCTGTTCAATTCATTCATTCCAAGTCTCAAAGCAGCAACATCA
ACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCGCCTTAACA

111 RREN_10854_3216 [T/C] ACCACCTACACAACGAGAGCCTTATGCTTCCAGTCAAGGAGCACAACCTACATGC
TCAGCAAGCAGTTCCTGGCCAAATGCCGCCATCCACTTCATCCAAA[T/C]TTCCT
CGCCACAAACAACGTCCCAGCCAGGCTCATGAAGCAGACCCTTCCGTCCAAGTT
TTGGAAGGAAGTCGACCAAGCTCTACAGGCCGCGGACAACG

112 RREN_10854_3286 [A/G] GGCAAATGCCGCCATCCACTTCATCCAAACTTCCTCGCCACAAACAACGTCCC
 AGCCAGGCTCATGAAGCAGACCCTTCCGTCCAAGTTTTGGAAGGAA[A/G]TCGA
 CCAAGCTCTACAGGCCGCGGACAACGACCACAACAAATGCAGCGCCAACATCC
 ACACAACCACAGTCCAAACGGCCAAGCAGCGACAAAGGAGCAC

113 RREN_16875_122 [T/C] AAGCAGTAACATCAACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCGCCTTAACAC
 CGCCGGGTCCAACAACACTGCGCCAGCCACCGGGGACATTCGAGCCA[T/C]AGCCC
 CTATTTAAGCCGCTCCAACAATGAGAGAGGCATGTTCAATTTGTTTCATTCATACC
 ATCTATTTCAACTGCTCAAAGCAGTAACATCAACTCTCTT

114 RREN_16875_146 [T/C] AAGCAGTAACATCAACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCGCCTTAACAC
 CGCCGGGTCCAACAACACTGCGCCAGCCACCGGGGACATTCGAGCCA[T/C]AATGA
 GAGAGGCATGTTCAATTTGTTTCATTCATACCATCTATTTCAACTGCTCAAAGCAG
 TAACATCAACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCG

115 RREN_16875_190 [T/C] AAGCAGTAACATCAACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCGCCTTAACAC
 CGCCGGGTCCAACAACACTGCGCCAGCCACCGGGGACATTCGAGCCA[T/C]CAACT
 GCTCAAAGCAGTAACATCAACTCTCTTCTCTCCGTTTCGCCTCCTCCTGCCGCCTT
 AACACCGCCGGGTCCAACAACACTGCGCCAGCCACCGGGGAC

116 RREN_23053_791 [T/C] GCAAAGGTCGTTTTGGCATTCTTCACAATAAAAATTTACCGTTAAATGCGTCTAAG
 CAATAATTGCAAATGGTTAAATGATGACGAAAATCATAGTTCCAT[T/C]TTAAAT
 TATTGCAGAAGCCGCGCTCTCTACAATCCCCTGAATATCAATAAACCATGATC
 CCTTTGTCACTGAAAATAAAGACTGTAGGAAATTTATGAT

117 RREN_23053_809 [T/C] TTCTTCACAATAAAAATTTACCGTTAAATGCGTCTAAGCAATAATTGCAAATGGTT
 AAATGATGACGAAAATCATAGTTCCATTTTAAATTATTGCAGAAG[T/C]CGCGCT
 CTCTACAATCCCCTGAATATCAATAAACCATGATCCCTTTGTCACTGAAAATAA
 AGACTGTAGGAAATTTATGATGCTGAGATTGTGGCTCCC

118 RREN_28983_551 [T/C] ACATGTTTTTTCTTCGTTTTACTGCGTTTTCTATTGATTTTATGCATTTTTGTAAAG
 TGTACCATCGAAATTTGCGTGATAACTAAAGGGGTTCAAAAAC[T/C]GGTTTAAA
 GTGGTTTCCCCCAAAAATAATTACATATTCGTA CT CAGCTCGACGAGCTCTACA
 CGATGGTATAAATTCTACTCGATTTGTCTTCCGGTAC

119 RREN_36168_664 [A/C] GAATTTAATTTGTTTCAGCGCAACTCTCAAGCAAAAATGCGGTGAATGTTCTGCGC
 AGGCGGA ACTATAAAAAGGGTGTTGATTTTCGCACCGTAACCACCACC[A/C]GCCAT
 CACTCGCTCTACACGCAACAAACACGCACTCAACTCTCCGCGTTCCCAGCAGCC
 GCTTCAACAACACAACACTGTTTCGTCGACCCTCTGGACAAG

120 RREN_37161_217 [A/G] CCGAAATACTAAAAACACGTAAAAACTTCGAAGGATCATAACTCTGCTACAGCA
TATCCATGCAAGACGAGCAATATACCAATCAATAGAGTAACATGTC[A/G]CGTA
AAAACTTCGAAGGATCATAACTCTGCTACAGCATATCCATGCAAGACGAGCAAT
ATACCAATCAATAGAGTAACATGTCCTCCACTAATCCCAGAA

121 RREN_37161_226 [C/G] CCGAAATACTAAAAACACGTAAAAACTTCGAAGGATCATAACTCTGCTACAGCA
TATCCATGCAAGACGAGCAATATACCAATCAATAGAGTAACATGTC[C/G]TTCGA
AGGATCATAACTCTGCTACAGCATATCCATGCAAGACGAGCAATATACCAATCA
ATAGAGTAACATGTCCTCCACTAATCCCAGAAAAGAACATG

122 RREN_43396_312 [T/C] TTTTCTACTGTAAACAGGAATTCGCATATTCTGAGACCACCATCGTGTAGAGCAT
GGTCGATAATAATAAGGAAGTGACATCCTTTTTTGGCACAAACCC[T/C]TGGTTA
AATTTTGAGTGAATTTTTAAAATATTTTTTCCACGTGCTTCAAGCACGGGTCATC
GGTGCTAAAAATGTCTTTTGGTCAACAAAGCTCAATAAG

123 RREN_47097_473 [C/G] CATTGTTTTGTATGCTTGTATTTGTATTGTTTTCCCTCTTCTTCCCTTCCCTTCCCT
CACGCCTGCATGTATATAAGCTAGTGATTTTCCCTCATTCTT[C/G]GCAGTTGGTT
GTCACCCCAGCTCGCCATTGTTTATTGTCTATTGTATCTTCAATAAACACTTCTTC
TCGGGTTCGGACTTCACTTTGGGGTTGTCCTTAC

124 RREN_47097_533 [T/C] CGCCTGCATGTATATAAGCTAGTGATTTTCCCTCATTCTTCGCAGTTGGTTGTCA
 CCCAGCTCGCCATTGTTTATTGTCTATTGTATCTTCAATAAACA[T/C]TTCTTCTC
 GGGTTCGGACTTCACTTTGGGGTTGTCCTTACTCCTAAACTATCAATCTTCATTG
 GGCAGTGGCTGTCCTACATTTTTCTCCGATTATTACT

125 RREN_47097_539 [A/C] CATGTATATAAGCTAGTGATTTTCCCTCATTCTTCGCAGTTGGTTGTCACCCCAG
 CTCGCCATTGTTTATTGTCTATTGTATCTTCAATAAACACTTCTT[A/C]TCGGGTTC
 GGACTTCACTTTGGGGTTGTCCTTACTCCTAAACTATCAATCTTCATTGGGCAGT
 GGCTGTCCTACATTTTTCTCCGATTATTACTTGGTAG

126 RREN_53141_284 [A/T] GACACCACCATCGGAATCAGCATCTACGAAAACCTCGAGTACTAAAAATTGGC
 TCAAATATCAATAATAACAACATTTCCATTAGGCCCCCAAAAA[A/T]TTAGG
 CCCCCAAAAACCTGAAATTTTTTCGAAAATTTCCCAAAAATTTTCTTAGGCCT
 AACCTAAGGTCAAATTTTTTGACTAATTTTATGTTTTCG

127 RREN_53141_365 [T/G] AATAATAACAACATTTCCATTAGGCCCCCAAAAAACCTGAAATTTTTTCGAAA
 ATTTCCCAAAAATTTTCTTAGGCCTAACCTAAGGTCAAATTTTT[T/G]GACTAA
 TTTTATGTTTTCGCACATGCTCTACAACCTTAAAAAATAAACCATGCCTCTAAAC
 CCCTTATAAACACTTCAAATAGTCCTGTCACGTATGATT

128 RREN_53845_394 [T/G] ACGCTTATGGTAGAGGGGATACGATAACGCCTTACTGTAGAGGGAATGCCGCA
ACACATTTAGGGTAGAAGGGATACGGTAACACACTTAAGGGAGGGG[T/G]AGGA
TACGGTACCACAATACGTTAACACAATTGATGTGATACGGTACCACACCTTAAG
TAGAGGGATACGGTAACACACTTATGGTAGAGGAAATACGAG

129 RREN_57146_374 [T/C] TGCTGATTTGGGGATCCTTTGTCCGTCGGCCTTTTGTCCACAATCCGATGTCATA
TTCGGATTCAGCAGCCTCGATAACCCCCGAGTACCAAAAATCAGC[T/C]CAAAT
ATCAATTACAAATAGCATTTCATTAGGCCTCCAAAAACCTGAAATTTTTTTGA
AAATTTCCCGAAAATTTTTCTTAGGCCCAACCTAAGGTC

130 RREN_86325_225 [A/G] CTCTCTAACTCTCCCTCGTCGTCTATACTCTCTCCCAAACCACACTATCTTATAT
TTTTTATGCAATATTCCATCCCCTCTTCACGCTTTTCCAATCAC[A/G]TACTCTCTC
CCCAAACCACACTATCTTATATTTTTTATGCAATATTCCATCCCCTCTTCACGCTT
TTCCAATCACTCCCCATCGCTTACGACCATACCG

131 RREN_90419_227 [A/G] TGAATAAGTGTGCCAGTGAAATTTGCAGTAACTAAAAGGGTTCAAAAACCGGTT
TAAAGTGGTTTCCCCCAAAAATAATTACATATTCGTAICTCAGCTC[A/G]GTAAC
TAAAAGGGTTCAAAAACCGGTTTAAAGTGGTTTCCCCCAAAAATAATTACATA
TTCGTAICTCAGCTCGACGAGCTCTACACGATGGTATATAAT

**APPENDIX 3: LETTER REQUESTING PERMISSION TO USE COMMON
METHODOLOGY IN CHAPTER 3**

Churamani Khanal
Department of Plant Pathology and Crop Physiology
302 Life Sciences Building
Louisiana State University
Baton Rouge, La 70808


Herath Kularathna
Department of Plant Pathology and Crop Physiology
302 Life Sciences Building
Louisiana State University
Baton Rouge, La 70808

Dear Mr. Kularathna,

Would you please grant me permission to use common methodology from our collaborative research on chapter three of my dissertation? I look forward to hearing from you about your decision.

Thank you.

Sincerely,


.....
Churamani Khanal

**APPENDIX 4: LETTER OF PERMISSION TO USE COMMON METHODOLOGY IN
CHAPTER 3**

Herath Kularathna
Department of Plant Pathology and Crop Physiology
302 Life Sciences Building
Louisiana State University
Baton Rouge, La 70808

Churamani Khanal
Department of Plant Pathology and Crop Physiology
302 Life Sciences Building
Louisiana State University
Baton Rouge, La 70808

Dear Mr. Khanal,

I hereby grant you permission to use common methodology from our collaborative research on chapter three of your dissertation.

Thank you.

Sincerely,



.....
Herath Kularathna

VITA

Churamani Khanal hails from Nepal. He received a B.S. in agriculture with plant pathology major from Tribhuvan University, Nepal in 2010. From 2011 to 2012, he served as an agriculture instructor as well as coordinator of Junior Technician Program in a governmental institution, Rapti Technical School, Dang, Nepal. He enrolled in the Department of Plant Pathology at the University of Arkansas, Fayetteville, Arkansas in 2012 and earned an M.S. degree in 2014 under supervision of Dr. Robert T. Robbins.