

2011

Biomechanical evaluation of medial and lateral approaches for experimentally created medial condylar fractures of the equine third metacarpal bone

Saybl Beauton Sprinkle

Louisiana State University and Agricultural and Mechanical College, ssprin3@tigers.lsu.edu

Follow this and additional works at: https://digitalcommons.lsu.edu/gradschool_theses



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Sprinkle, Saybl Beauton, "Biomechanical evaluation of medial and lateral approaches for experimentally created medial condylar fractures of the equine third metacarpal bone" (2011). *LSU Master's Theses*. 729.

https://digitalcommons.lsu.edu/gradschool_theses/729

This Thesis is brought to you for free and open access by the Graduate School at LSU Digital Commons. It has been accepted for inclusion in LSU Master's Theses by an authorized graduate school editor of LSU Digital Commons. For more information, please contact gradetd@lsu.edu.

BIOMECHANICAL EVALUATION OF MEDIAL AND LATERAL APPROACHES FOR
EXPERIMENTALLY CREATED MEDIAL CONDYLAR FRACTURES OF THE EQUINE
THIRD METACARPAL BONE

A Thesis
Submitted to the Graduate Faculty of the
Louisiana State University and
Agricultural and Mechanical College
In partial fulfillment of the
Requirements for the degree of
Master of Science

In

The Interdepartmental Program in
Veterinary Medical Sciences
Through the
Department of Veterinary Clinical Sciences

by
Saybl Beauton Sprinkle
B.S., Louisiana State University, 2007
August 2011

DEDICATION

I would like to dedicate this work to:

Dr. Gary A. Sod, and Dr. Laura M. Riggs for their encouragement, support, and belief in me throughout the years and this program.

ACKNOWLEDGEMENTS

The author would like to thank her major professor, mentor, and friend, Dr. Gary A. Sod for his continued patience and guidance throughout her graduate program. Without his constant encouragement, confidence, hours spent in necropsy helping prepare samples and support throughout this program the author could not have made it through this program.

The author would also like to thank the other members of her committee, Drs. Laura M. Riggs, Daniel J. Burba, and Colin F. Mitchell, for their encouragement, support during the research, and for reviewing and giving advice for the improvements of the manuscript. The author would like to thank Dr. Mitchell for asking about the research and always offering to help anytime needed. The author would also like to thank Dr. Riggs for helping her with her project throughout this program. The author would also like to thank Dr. Burba for agreeing to join her committee on a short notice.

The author would also like to take this opportunity to thank Niki Marie Hansen, for her friendship, encouragement, support, advice throughout this program and life, and her confidence she bestowed within the author. The author would also like to thank her colleagues Dr. Nan Huff and Dr. Josh Cartmill and Jennifer Sloan who also helped encouraged and reassured that everything would be okay.

Finally, the author would like to take this time to thank her parents, Douglas and Kathleen Sprinkle, for their support throughout the years. The author would like to thank Kathleen Sprinkle for always expressing how proud she is of the author, always encouraging her to go for her dreams, and always being there when needed.

TABLE OF CONTENTS

DEDICATION.....	ii
ACKNOWLEDGEMENT.....	iii
LIST OF TABLES.....	vi
LIST OF FIGURES.....	viii
ABSTRACT.....	x
CHAPTER 1. GENERAL INTRODUCTION.....	1
1.1 The Metacarpo(-tarso)phalangeal Joint.....	2
1.2 The Equine Third Metacarpal Bone	3
1.3 Pathophysiology of Condylar Fractures	3
1.4 The AO/ASIF.....	9
1.4.1 The 4.5-mm AO Cortical Screw	10
1.4.2 The Lag Screw Principle.....	11
1.5 Repair of Medial Condylar Fractures.....	12
<p>CHAPTER 2. COMPARISON OF COMPRESSION OF VARYING FRAGMENT THICKNESSES OF EXPERIMENTALLY CREATED MEDIAL CONDYLAR FRACTURES OF THE EQUINE THIRD METACARPAL BONE WITH A 4.5-MM AO CORTICAL BONE SCREW USING A MEDIAL VS. LATERAL</p>	
APPROACH.....	14
2.1 Introduction.....	15
2.2 Hypothesis and Objectives.....	16
2.3 Methods and Materials.....	16
2.4 Results.....	18
2.5 Discussion.....	21
<p>CHAPTER 3. COMPARISON OF TORQUE TO FAILURE OF VARYING FRAGMENT THICKNESSES OF EXPERIMENTALLY CREATED MEDIAL CONDYLAR FRACTURES OF THE EQUINE THIRD METACARPAL BONE WITH A 4.5-MM AO CORTICAL BONE SCREW USING A MEDIAL VS LATERAL</p>	
APPROACH.....	23
3.1 Introduction.....	24
3.2 Hypothesis and Objectives.....	25
3.3 Methods and Materials.....	25
3.4 Results.....	27
3.5 Discussion.....	32

CHAPTER 4. FINAL DISCUSSION AND CONCLUSIONS.....	34
4.1 Summary.....	35
BIBLIOGRAPHY.....	36
VITA.....	43

LIST OF TABLES

Table 2.1 Compression values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of an 8 mm thick experimentally created medial condylar fracture in equine MC3 bones.....	18
Table 2.2 Compression values during tightening of a 4.5 mm AO screw Placed in lag fashion for reduction of an 12 mm thick experimentally created medial condylar fracture in equine MC3 bones.....	19
Table 2.3 Compression values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of an 20 mm thick experimentally created medial condylar fracture in equine MC3 bones.....	19
Table 3.1 Torque to failure values and failure modes of 4.5 mm AO screws used for repair of experimentally created 8 mm thick medial condylar fractures in equine MC3 bones.....	29
Table 3.2 Torque to failure values and failure modes of 4.5 mm AO screws used for repair of experimentally created 12 mm thick medial condylar fractures in equine MC3 bones.....	30
Table 3.3 Torque to failure values and failure modes of 4.5 mm AO screws used for repair of experimentally created 20 mm thick medial condylar fractures in equine MC3 bones.....	31

LIST OF FIGURES

Figure 1.1 Photographic view of the palmar distal region of the distal joint surfaces of the MC3 bones from a 3-year-old male Thoroughbred	5
Figure 1.2 Scanning electron microscopic views of the failure surface from a 5-year-old racing Thoroughbred with a catastrophic MC3 lateral condylar fracture	6
Figure 1.3 Standard dorso-palmar radiographic images of a Type 1 (A), Type 2 (B), Type 3 (C), and Type 4 (D) condylar fractures.....	8
Figure 2.1 Example of a graph generated by the I-scan system recording software during the tightening of an AO screw to repair an experimentally created medial condylar fracture in an equine MC3 bone. Data was collected at a rate of 100 frames per second.....	17
Figure 2.2 Mean fracture plane force values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of experimentally created medial fracture of equine third metacarpal.....	20
Figure 2.3 Mean fracture plane pressure values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of experimentally created medial condylar fracture of equine third metacarpal.....	21
Figure 3.1 Equine cadaveric third metacarpal bones that were tested for torque to failure of experimentally created medial condylar fractures of various thicknesses repaired with a 4.5 mm AO screw placed in lag fashion. Types of failure that can occur with repair of condylar fractures include bone failure with the threads stripping or screw failure with the screw breaking at the screw head junction.....	26

Figure 3.2 Image of the condylar region of an equine third metacarpal bone showing where a cortical screw engaged the bone with lag screw fixation of a fracture.....

27

ABSTRACT

Objective -To compare the compression produced in reduction of experimentally created medial condylar fractures using lag screw fixation with medial vs. lateral approach, and to determine the maximum torque at screw failure.

Materials and Methods- Twelve (12) pairs (left and right) 3rd metacarpal bones (MC3) were collected from adult (2-7 years) Thoroughbreds euthanized for reasons unrelated to orthopedic disease. Complete parasagittal medial condylar osteotomies were created at a measurement of 9, 13, and 21 mm axial to the epicondylar fossa on four pairs each of cadaveric MC3 bones resulting in fracture fragments measuring 8, 12, and 20 mm in thickness. For each pair of cadaveric MC3, a lateral or medial approach was randomly selected to repair the condylar fracture using a single 4.5 mm AO cortical screw. Each repair was tested for fracture plane compression and screw torque to failure.

Results-There was no significant difference in compression between the medial and lateral approaches for the 8 or 12-mm fragment groups. There was significantly more compression generated in the lateral approach when compared to the medial approach for the 20-mm fragment group. Failure occurred at significantly lower torque in the 8-mm group. There was no significant difference between medial and lateral approach in torque to failure for the 12 and 20-mm groups.

Conclusion-Based on this data we have concluded that there was no significant difference in torque to failure between a medial vs. lateral approach for the 12 mm fragments but there was a

significant difference for the 8 mm fragments and that a lateral approach may be acceptable for the repair of medial condylar fractures in 12-mm or thicker fragments. The compression achieved by a medial approach was not significantly greater for the 8, 12 or 20-mm groups.

Clinical Relevance- Based on our results the 20 mm size fragments reaches a higher compression at a faster rate when compared to the 8 and 12 mm size fragments. We recommend using caution when repairing medial condylar fractures with a lateral approach for fragment sizes measuring 8-mm thick. The smaller fragment torque to failure was low and not much higher than the insertional torque. Failure resulted from the screws stripping in the bone fragment. The screws in the thicker fragments (12 and 20 mm) engage more bone and have a higher torque to failure as a result.

CHAPTER 1
GENERAL INTRODUCTION

1.1 The Metacarpo(-tarso)phalangeal Joint

The equine metacarpophalangeal and metatarsophalangeal joints (fetlock joints) are made up of the third metacarpal bone, both proximal sesamoid bones, and the first phalanx. It is classified as a ginglymus (hinged) joint, thus it has only one axis of movement in the sagittal plane through flexion and extension. This limitation is due to the presence of the prominent medial and lateral collateral ligaments of the fetlock. It is the one joint in the horse with the highest range of motion ranging from 120° of extension to 120° of flexion. This is best appreciated during athletic activity such as racing or jumping (Bertone, 2004). In the standing horse, the fetlock is supported by the suspensory ligament, intersesamoidean ligament, distal sesamoidean ligaments, deep and superficial flexor tendons, the common digital and lateral digital extensor tendons, and the medial and lateral collateral ligaments.

The locomotor functions of the digit and fetlock include the flexion essential to movement, extension when the foot is off the ground, the diminution of concussion when the hoof contacts the ground, and the recovery from extension (Kainer, 2002). The normal movements of the digit were elaborated by Rooney (1974). When the body weight of the horse is applied to the leg, the fetlock joint extends and translates downward as a result of distal interphalangeal joint motion. With extension, the cannon bone and the first phalanx (P1) tend to rotate about their long axes in a medial to lateral direction. As the limb is unloaded the fetlock joint begins to flex, allowing the dorsal angle of the fetlock to open, and the pastern elevates because of the distal interphalangeal joint motion in the opposite direction. During this opening and elevating movement, the cannon bone and the first phalanx again rotate, however this time about their long axes in

a lateral to medial direction. This movement necessitates that the two bones must move in synchrony.

1.2 The Equine Third Metacarpal Bone

The MC3 is a substantial bone, measuring approximately 26 cm in length from the midsagittal ridge to the carpometacarpal joint. At the distal metaphysis, it is approximately 50 wide from the lateral to medial supracondylar fossae. Cortical and trabecular densities for equine bone have been previously measured and determined to be 1035.25 mg hydroxyapatite/ml and 1048.55 mg hydroxyapatite/ml, respectively (Furst et al, 2008). Mean tensile strengths and modulus elasticity have been reported to be 2137.9 to 2295.7 MPa (El Shorafa et al, 1979) and 16.3 GPa (Batson et al, 2000), respectively.

1.3 Pathophysiology of Condylar Fractures

It has been theorized that condylar fractures result due to asynchronous movement between the cannon bone and the proximal phalanx. The asynchronous movement leading to lateral condylar fractures was described (Alexander and Rooney 1972, Rooney 1974) as the proximal phalanx remains stationary with the fetlock in a fully dorsiflexed position while the cannon bone rotates in a lateral to medial direction, owing to fracture of the lateral condyle as it strikes the stationary lateral aspect of proximal P1. This asynchrony would be expected near the end of the support phase of the stride when the fetlock joint dorsal angle is opening (Rooney

1974). No work has been performed to support this theory, however more recent efforts have been made to determine predisposing factors associated with condylar fracture formation.

Recent work evaluating the pathophysiology of the development of condylar fractures suggests that they originate from microfractures in the subchondral bone that develop secondary to repetitive loading from exercise related stress remodeling (Fig. 1.1A and 1.1B), and is associated with significant cyclic shear loading of the condyle in dorsopalmar (-plantar) bending which then leads to propagation of a dominant crack proximally (Radtke et al. 2002). Ultrastructural work further revealed that these microfractures develop into clusters and most notably in the palmar aspect of the condyle; these macrofractures propagate along cement lines and interfaces between bone lamellae (Fig. 1.2A and 1.2B) (Stepnik et al. 2004).

Condylar fractures involving the third metacarpal and metatarsal bones have long been recognized as a significant problem among racehorses, and in severe cases can require euthanasia (Rooney 1974, Pool 1990, Wilson et al 1993, Stover 1994, Kane et al 1996, Kane et al 1998). Severe condylar fractures account for 20% (Johnson et al, 1994) to 25% (McKee, 1995) of catastrophic injuries in Thoroughbred racehorses in California and the United Kingdom, respectively. Condylar fractures of the left third metacarpal bone are the most commonly noted site in Thoroughbred racehorses, accounting for 39% to 57.1% of all condylar fractures (Rick et al. 1983, Ellis et al. 1994); whereas there appears to be a more even distribution with regards to left vs. right in Standardbred racehorses. This increased prevalence for occurrence in the left forelimb among Thoroughbred racehorses has been theorized to be due to the direction in which




Figure 1.1. Photographic view of the palmar distal region of the distal joint surfaces of the MC3 bones from a 3-year-old male Thoroughbred. (A) In the right MC3 bone, a parasagittal defect can be seen in the articular cartilage of the lateral condylar groove (black arrows). Adjacent to this lesion is circular area of cartilage degeneration in the lateral condylar (white arrow heads). A similar lesion is also present to a lesser extent in the medial condyle. Parasagittal linear wear lines in the articular cartilage are also visible. (B) In the left MC3 bone, a parasagittal condylar fracture is present in the lateral condylar groove (white arrowheads). In the medial condylar groove, a branching array of subchondral cracks can be seen (black arrows). In the lateral condylar groove, comminution of this subchondral bone developed during propagation of the fracture. The articular cartilage was removed by treatment with 0.1 M NaOH to permit the articular surface of the subchondral bone to be examined. (Pictures taken from Radtke et al. 2002)




Figure 1.2. Scanning electron microscopic views of the failure surface from a 5-year-old racing Thoroughbred with a catastrophic MC3 lateral condylar fracture. (A) In adapted palmar subchondral bone from the distal end of the MC3 bone, microcracks were often seen propagating along cement lines and the interfaces between bone lamellae. (B) In adapted dorsal subchondral bone from the distal end of the MC3 bone, arrays of branching microcracks were also seen, which were similar to the palmar/plantar region of the condyle. (Pictures taken from Stepnik et al. 2004)

racehorses are run, which places the left limb on the inside, thus increasing the load placed on this limb, when compared to the right limb. This same observation has been made in England, however one study reported that the majority of condylar fractures were incurred during training, which involves galloping mainly straight distances (Ellis et al. 1994).

Condylar fractures can occur in either the lateral or medial condyle; however, fractures of the lateral condyle are the most commonly occurring type (Alexander et al 1972, Meagher 1976, Rick et al 1983, Ellis 1994, Johnson et al 1994, Bassage et al 1998, Zekas 1999, LeJeune 2003), reported as being approximately 85% in various racehorse populations (Zekas et al 1999; Bassage et al 1998). Condylar fractures are typically described as being one of four types: 1) incomplete – no evidence of joint malalignment or complete extension of the fracture through the proximal cortex; 2) complete-nondisplaced fractures – neither a step at the joint surface nor evidence of separation of the fragment proximally – although the fracture line penetrates through the cortex; 3) complete-displaced fractures – malalignment at the joint surface and abaxial displacement at the proximal cortical surface; 4) special longitudinal diaphyseal fractures – either complete or incomplete – involving the medial condyle and extending various distances up the diaphysis (Figure 1.3) (Rick et al. 1983).

Historically, an onset of severe lameness following intense exercise occurs typically either immediately following injury or within a few hours of exercise. Typical physical examination findings reveal a variable lameness which does not correlate well with severity of the fracture, as severe non-weightbearing lameness is commonly seen with incomplete, nondisplaced fractures, yet a milder weightbearing lameness is



Figure 1.3. Standard dorso-palmar radiographic images of a Type 1 (A), Type 2 (B), Type 3 (C), and Type 4 (D) condylar fractures.

typically appreciated with displaced condylar fractures (Richardson, 2006). Additional findings include variable effusion of the metacarpo- or metatarso-phalangeal joint of the affected limb,

elicitation of pain on flexion of the joint, and either the presence or absence including the lateromedial and dorsopalmar (-plantar) are diagnostic, however it is strongly recommended that a horizontal dorsopalmar projection with the fetlock placed in slight flexion be performed highlighting the palmar aspect of MC3/MT3 to rule out the presence of palmar comminution, a common finding associated with condylar fractures (Richardson, 2006).

Treatment goals for type 1, 2, and 3 condylar fractures are aimed at either conservative management consisting of prolonged periods of stall confinement, immobilization via external coaptation by heavy bandaging or distal limb casting (Meagher, 1976, Rick, et al. 1983), or surgical reduction; however, surgical reduction is typically recommended for the best possible prognosis. Surgical repair is most commonly performed through fracture fragment fixation via placement of either 4.5-mm or 5.5-mm AO cortical bone screws (Synthes, Paoli, PA) using the lag screw principle to allow for adequate compression and allows for fracture healing as well as to align the articular surfaces of the fracture to minimize the chances for development of degenerative joint disease (Richardson. 2006).

1.4 The AO/ASIF

The Arbeitsgemeinschaft für Osteosynthesefragen (Association for the Study of Internal Fixation) was formed in 1958 to further research into the concepts of immediate functional rehabilitation after rigid internal fixation. They have done this through research into osteosynthesis, and the development of instruments and implants that of crepitation on palpation of the limb. Radiographic evaluation of the fetlock joint typically confirms the presence of condylar fracture. Traditional radiographic projections promote rigid internal fixation. This has

evolved into a worldwide organization evaluating not only the development of implants in human orthopedics, but as well in veterinary orthopedics (Colton. 1981).

1.4.1 The 4.5-mm AO Cortical Screw

The 4.5-mm AO cortical screw is made of implant quality 316L stainless steel, which contains roughly 62.5% iron, 17.6% chromium, 14.5% nickel, 2.8% molybdenum, and minor alloy additions. It is meant to be resistant to corrosion due to its low carbon content (Texhammar. 1981). This screw was designed for use in the lag principle or for plate fixation. It is a fully threaded non-self-tapping screw. In cortical bone, the screw has a holding strength of approximately 2500 N.

The dimensions of the screw are as follows:

Head diameter: 8.0-mm

Hexagonal socket width: 3.5-mm

Core diameter: 3.0-mm

Thread diameter: 4.5-mm

Pitch: 1.75-mm

Glide hole diameter: 4.5-mm

Thread hole diameter: 3.2-mm

Tap diameter: 4.5-mm

The 4.5-mm cortical screw also has characteristics similar to that of the other large AO/ASIF screws including first a spherical screw head which ensures optimal screw to plate contact even if a screw is placed at an angle. Second, the AO/ASIF screw thread which is characterized as a buttress thread profile which allows excellent holding in cortical bone due to the shallow thread and fine pitch leading to a large screw-to-bone contact area. The third common characteristic is the core diameter which is the solid stem of the screw from which the threads protrude.

1.4.2 The Lag Screw Principle

The lag screw was defined by Perren and Buchanan (1981) as *the production of interfragmental compression by compressing the bone under the screw head against the fragment in which the screw threads are anchored*. The steps for proper lag screw principle for a 4.5-mm cortical screw are as follows:

1. The fracture is reduced and held with reduction forceps.
2. The glide hole is drilled through the near cortex or fragment with a 4.5-mm drill bit protected by the drill sleeve.
3. The 3.2-mm drill sleeve is then inserted into the glide hole until it comes into contact with the far cortex or the parent bone.
4. The thread hole is then drilled in a coaxial direction in the trans-cortex or parent bone with a 3.2-mm drill bit.

5. The cis-cortex is countersunk with the large countersink.
6. Screw length is then measured with the large depth gauge
7. The thread hole is then tapped with a 4.5-mm bone tap. By turning two turns clockwise and one-half turn counterclockwise, the cut bone is directed into the channels of the cutting flutes to be removed.
8. A 4.5-mm cortical screw is inserted with the large hexagonal screwdriver, ensuring engagement of the trans cortex.

1.5 Repair of Medial Condylar Fractures

Medial condylar fractures typically differ from lateral condylar fractures by having a longitudinally originating configuration that typically does not toward the ipsilateral cortex. Medial condylar fractures tend to be more complex as they classically spiral as it propagates from the articular surface. These fractures typically have a configuration where the dorsal cortical fracture line propagates medially and the palmar/plantar cortical fracture line propagates laterally. Thus the fracture plane propagates counterclockwise when views from the distal aspect of the left limb (Turner 1977; Barr 1989; Ellis 1994). Traditionally these fractures have been repaired with the screws being placed medially. The two fragments in a spiraling medial condylar fracture are approximately the same size, thus a medial approach offers no biomechanical advantage over a lateral approach. It is technically easier to perform surgery with the affected limb uppermost (Auer 2006) but this requires a lateral approach. Repair of a dorsomedial-palmaro/plantarolateral spiral configuration fracture from the medial aspect with

multiple screws in lag fashion risks screw interference with the soft tissue structures on the palmar/plantar aspect of MC3/MT3 and the second metacarpal/metatarsal bone may prevent insertion perpendicular to the fracture plane (Richardson 1990). By inserting screws from the lateral aspect, the contour of the fracture will necessitate that the holes are drilled from a more dorsal direction, naturally taking the screw heads away from the palmar/plantar structures of the leg. The advantage of direct fracture observation is the ability to insert screws perpendicularly to the fracture plane, in a biomechanically optimal position, along the entire length of the fracture (Rick, O'Brien et al. 1983; Smith, Greet et al. 2009; Wright and Smith 2009). In two case series involving Thoroughbred racehorses with medial condylar fractures, it was concluded the repair of propagating sagittal and spiral fractures of the medial condyle of MC3/MT3 with diaphyseal involvement through a lateral approach with periosteal reflection permitted stable fixation with minimal complications (Smith 2009, Wright 2009).

CHAPTER 2

COMPARISON OF COMPRESSION OF VARYING FRAGMENT THICKNESSES OF EXPERIMENTALLY CREATED MEDIAL CONDYLAR FRACTURES OF THE EQUINE THIRD METACARPAL BONE WITH A 4.5-MM AO CORTICAL BONE SCREW USING A MEDIAL VS. LATERAL APPROACH

2.1 Introduction

Fractures of the condyles of the third metacarpal (MC3) or metatarsal (MT3) bones are relatively common injuries in Thoroughbred racehorses and are the most common long bone fracture of horses in training (Ellis 1994; Wright and Smith 2009). These fractures occur more frequently in the lateral condyle while medial condylar fractures typically originate closer to the sagittal ridge and differ from lateral condylar fractures by having longitudinally originated configuration that typically does not extend towards the ipsilateral cortex and classically spirals as it propagates from the articular surface (Rick, O'Brien et al. 1983; Bassage and Richardson 1998; Smith, Greet et al. 2009). Medial condylar fractures are traditionally repaired through a medial approach; however recent publications suggest a lateral approach is acceptable clinically and is technically easier due to patient positioning (Wright and Smith 2009).

Recently, two case series have been published which describe lateral approach for repair of medial condylar fractures. These investigations had positive results and stated that fractures of medial MC3 or MT3 can be repaired by a lateral approach without greater complication rates or negative effect on outcome associated with the side of approach. However to our knowledge no controlled studies have been published comparing the biomechanical differences between lateral and medial approaches for repair of medial condylar fractures. Our study will compare the compression values of a single AO cortical screw placed in lag fashion using a medial vs. lateral approach.

2.2 Hypothesis and Objectives

We hypothesized that there will not be any significant difference in compression values when repairing an experimentally created medial condylar fracture of equine third metacarpal with lag screw fixation from a lateral approach versus a medial approach.

2.3 Materials and Methods

Twelve (12) pairs (left and right) of MC3 bones were collected from adult (2-7 years) Thoroughbreds euthanized for reasons unrelated to orthopedic disease. Each MC3 was wrapped in saline (0.9% NaCl) soaked towels with each pair identified and stored in a freezer at -20 C. The MC3 were thawed at room temperature for 24 hours prior to testing. For the preparation of the fracture, the cadaveric metacarpal bone was secured in a positioning jig and a lapidary table saw (Covington Slab Saw) with a blade thickness of 0.8 mm was used to create a medial fracture on the condyle at the level of the epicondylar fossa. The limbs were randomly divided into three groups (four pairs each). Complete parasagittal condylar osteotomies were created at 9, 13, or 21 mm axial to the medial epicondylar fossa resulting in fracture fragments measuring 8 (group 1), 12 (group 2), and 20 mm (group 3). A transverse cut was made proximal to the condyle to make a complete fracture model. In each group, a lateral or medial approach was randomly selected for each limb in the pairs, to repair the medial condylar fracture.

A Tekscan k-scan 6220 sensor with a 2 mm stay pin hole and a center hole to accommodate screws was placed in the osteotomy plane between the fracture fragments and connected to a computer. The MC3 was secured in the vice press and a 4.5 mm AO (Synthes, Paoli, PA) screw was placed in routine lag fashion through the epicondylar fossa of each bone. The screw measured 60 mm in length was positioned and tightened with a digital torque wrench

to an insertional torque of 5.4 N m. The compression pressure and force was acquired at a rate of 100 frames per second via the I-Scan System during tightening and was recorded for data analysis (Figure 2.1). Results were compared with a student t-test for paired samples and analyzed using Prism software. P-value was set at <0.05.




Figure 2.1 Example of a graph generated by the I-scan system recording software during the tightening of an AO screw to repair an experimentally created medial condylar fracture in an equine MC3 bone. Data was collected at a rate of 100 frames per second.

2.4 Results

There were no significant differences in compression (kPa) between medial and lateral approach for the 8 mm ($p=0.42$) or 12 mm thick fragments ($p=0.65$) (Tables 2.1 & 2.2; Figures 2.3 & 2.4). For the 20 mm fragment group, the lateral approach resulted in significantly greater compression when compared to the medial approach ($p=0.023$) (Table 2.3; Figures 2.3 & 2.4).

Table 2.1 Compression values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of an 8 mm thick experimentally created medial condylar fracture in equine MC3 bones.

Approach	Leg	Area (mm)	Force (N)	Pressure (kPa)
Medial				
	R	452	734.8	1625.6
	L	387	1242.7	3211.0
	R	495	1402.9	2834.1
	R	531	1979.2	3727.3
Standard Deviation		466.3 ± 61.9	1339.2 ± 512.6	2849.5 ± 894.3
Lateral				
	L	444	855.2	1926.2
	R	364	471.1	1294.1
	L	321	343.7	1070.7
	L	419	572.4	1366.0
Standard Deviation		387 ± 55.3	560.6 ± 217.6	1414.3 ± 363.7

R=right, L=left

Table 2.2 Compression values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of a 12 mm thick experimentally created medial condylar fracture in equine MC3 bones.

Approach	Leg	Area (mm)	Force (N)	Pressure (kPa)
Medial				
	R	485	1373.1	2831.1
	L	511	1896.9	3712.1
	L	393	783.3	1993.0
	R	447	1174.7	2627.9
Standard Deviation		459.0 ± 51.3	1307.0 ± 463.4	2791.1 ± 710.3
Lateral				
	L	526	1398.1	2658.1
	R	469	1534.9	3272.7
	R	378	675.2	1786.2
	L	412	1047.7	2543.0
Standard Deviation		446.3 ± 65.1	1164.0 ± 385.1	2564.5 ± 610.7

R=right, L=left

Table 2.3 Compression values during tightening of a 4.5 mm AO screw placed in lag fashion for reduction of a 20 mm thick experimentally created medial condylar fracture in equine MC3 bones.

Approach	Leg	Area (mm)	Force (N)	Pressure (kPa)
Medial				
	R	509	617.9	1213.9
	L	473	404.1	1854.4
	R	406	255.4	629.1
	R	452	477.9	1057.4
Standard Deviation		460.0 ± 43.0	438.8 ± 151.0	938.7 ± 253.5
Lateral				
	L	556	1091.8	1963.7
	R	504	781.1	1727.1
	L	472	547.5	1297.3
	L	517	974.4	2086.4

Standard Deviation		512.3 ± 34.8	848.7 ± 238.2	1639.5 ± 366.7
--------------------	--	--------------	---------------	----------------

R=right, L=left




Figure 2.2 Mean fracture plane force values obtained during tightening to 4.5 N-m of a 4.5 mm AO screw placed in lag fashion for reduction of experimentally created medial condylar fractures of equine third metacarpi.




Figure 2.3 Mean fracture plane pressure values obtained during tightening to 5.4 N-m of a 4.5 mm AO screw placed in lag fashion for reduction of experimentally created medial condylar fractures of equine third metacarpi.

2.5 Discussion

Adequate fracture compression using lag screw technique for the repair of condylar fractures requires substantial engagement of the bone by the screw in the thread hole. The compression achieved is related to the number of threads engaging the bone although a minimum number of threads and thus minimum value for “adequate” compression is not defined for this fracture configuration. The 4.5 mm cortical screw used here has a thread pitch, the distance from one thread to the next, of 1.75 mm. Given this value, a thread hole created in 8, 12 and 20 mm fragments will engage 4.6, 6.9 and 11.4 threads, respectively, a step-wise increase of 50% and 67%.

The results of this study indicate that there is no significant difference in compression values between a medial and lateral approach for a medial condylar fracture with an 8 or 12 mm fragment. This result is in agreement with our hypothesis and appears to indicate that even small fracture fragments (8-12 mm) will engage enough of the screw to produce compression at an insertional torque of 5.4 N-m. This may allow for adequate compression and thus fracture healing without a gap at the articular surface.

The 20 mm fracture fragment also demonstrated adequate compression from both the medial and lateral approach. Interestingly the medial force and pressure were actually much lower, by approximately half, than the lateral approach and these results were statistically significant. Greater compression was expected with the medial approach due to the number of screw threads the bone would engage. However, that was not the case and cannot be easily explained. One possibility is that there is a threshold amount of screw engagement, to the point that compression can no longer increase. These findings in the larger fracture fragment bring up new questions to be answered. While it doesn't appear to be clinically relevant because of the substantial values obtained, it is an interesting phenomenon. Potentially, there is a point in the ratio of fracture fragments at which the bone and screw interface have a physical property change and different biomechanical principles are in effect. This theory remains to be proven.

It appears that the use of a lateral approach for the repair of medial condylar fractures is acceptable with regards to the compressive forces applied at the distal screw. This is an important point of repair for this fracture configuration and is vital for the congruency at the articular surface. Lateral repair of medial condylar fractures are technically easier to perform and allow for more direct approach to the fracture without interference by other structures. With increased use in clinical cases, we may find that this approach improves prognosis.

CHAPTER 3

COMPARISON OF TORQUE TO FAILURE FOR VARYING FRAGMENT THICKNESSES OF EXPERIMENTALLY CREATED MEDIAL CONDYLAR FRACTURES OF THE EQUINE THIRD METACARPAL BONE WITH A 4.5-MM AO CORTICAL BONE SCREW USING A MEDIAL VS LATERAL APPROACH

3.1 Introduction

Fractures of condyles of the third metacarpal (MC3) or Metatarsal (MT3) bone are relatively common injuries in Thoroughbred racehorses. They are amongst the most common long bone fractures of horses in training (Ellis 1994; Wright and Smith 2009). These fractures occur more frequently in the lateral condyle. Medial condylar fractures originate closer to the sagittal ridge and differ from lateral condylar fractures by having longitudinally originated configuration that typically does not extend towards the ipsilateral cortex and classically spirals as it propagates from the articular surface (Smith, Greet et al. 2009) (Rick, O'Brien et al. 1983; Bassage and Richardson 1998) Typically when screws are inserted in lag fashion the smaller bone fragment contains the glide hole and is compressed to the larger bone fragment. Stability is achieved by compression with a screw that is oriented perpendicular to the fracture (Smith, Greet et al. 2009). The compression that can be achieved is related to the strength of the screw-bone construct.

Recently, case series have been published which describe lateral approach for repair of medial condylar fractures. These reports presented positive results and stated that fractures of medial MC3/MT3 can be repaired by a lateral approach without greater complication rates or negative effect on outcome associated with the side of approach. However to our knowledge no controlled studies have been published comparing lateral vs. medial approaches for repair of medial condylar fractures. Thus this study will compare the torque to failure of 4.5 mm AO cortical screws placed in lag fashion for repair of experimentally created medial condylar fractures of various thicknesses in equine MC3 bones using a medial vs. a lateral approach.

3.2 Hypothesis and Objectives

We hypothesized that there will be significant difference in measurements of torque to failure of 4.5 mm cortical screws with a medial versus lateral approach.

3.3 Materials and Methods

Twelve pairs (left and right) of MC3 bones were collected from adult (2-7 years) Thoroughbred race horses euthanized for reasons unrelated to orthopedic disease. Each MC3 was wrapped in saline (0.9% NaCl) soaked towels with each pair identified and stored in a freezer at -20 C. The MC3 were thawed for 24 hours at room temperature prior to testing. For the preparation of the fracture the cadaveric metacarpal bone was secured in a positioning jig, and a lapidary table saw (Covington Slab Saw) with a blade thickness of 0.8 mm was used to create a medial fracture of the condyle at the level of the epicondylar fossa. The limbs were randomly divided into 3 groups (four pairs each). Complete parasagittal condylar osteotomies were created at 9, 13, or 21 mm axial to the medial epicondylar fossa resulting in fracture fragments measuring 8 (group 1), 12 (group 2), or 20 mm (group 3). A transverse cut was made proximal to the condyle to make a complete fracture model (Figure 2.5). In each group, a lateral or medial approach was randomly selected for each limb in the pairs to repair the medial condylar fracture. A Tekscan k-scan 6220 sensor with a 2 mm stay pin hole and a center hole to accommodate screws was placed in the osteotomy plane between the fracture fragments and connected to a computer (Figure 2.1). The MC3 was secured in the vice press and a 4.5 mm x 60 mm length AO screw was placed in routine lag fashion through the epicondylar fossa of each bone. The screws were tightened with a torque wrench with a digital display.

After the initial torque measurements were obtained the screws were tightened to the point of failure with the digital torque wrench and that measurement was recorded and compared between fragment thicknesses (Figure 3.1). In the figure 3.1 fragment thicknesses of 12 mm and 20 mm were used with the lateral and medial approach both showing screw failure with lag screw fixation. Medial and lateral approaches were compared with a student t-test for paired samples using Prism software. The p-value was set at <0.05 . The mode failure was recorded for each limb.



Figure 3.1 Equine cadaveric third metacarpal bones that were tested for torque to failure of experimentally created medial condylar fractures of various thicknesses repaired with a 4.5 mm AO screw placed in lag fashion. Types of failure that can occur with repair of condylar fractures include bone failure with the threads stripping or screw failure with the screw breaking at the screw head junction.



Figure 3.2 Image of the condylar region of an equine third metacarpal bone showing where a cortical screw engaged the bone with lag screw fixation of a fracture.

3.4 Results

The mean value for the torque to failure of 8 mm, 12 mm, and 20 mm fragments for the medial approach were 9.76 ± 0.33 N-m, 9.45 ± 0.73 N-m, and 10.15 ± 0.38 N-m, respectively. For the lateral approach with fragment sizes of 8 mm, 12 mm, and 20 mm the torque to failure measurements were 5.88 ± 0.19 N-m, 10.0 ± 0.30 N-m, and 10.07 ± 0.41 N-m, respectively. For the 8 mm fragment group there was a significant increase in maximum torque at failure for the medial approach when compared to the lateral approach ($p < 0.000008$) while there was no significant difference between medial and lateral approach for the 12 and 20 mm fragments, $p = 0.22$ and $p = 0.78$, respectively.

The mode of failure was dependent on fragment thickness and approach. For the 8 mm fragment, all screws introduced through a medial approach failed by screw breakage while all

screws introduced through a lateral approach failed by stripping of the screw threads (Table 3.1). For the 12 mm fragment the results were mixed (Table 3.2) and for the 20 mm fragments all screws failed by screw breakage (Table 3.3).

Table 3.1 Torque to failure values and failure modes of 4.5 mm AO screws placed in lag fashion used for repair of experimentally created 8 mm thick medial condylar fractures in equine MC3 bones.

Approach	Leg	Failure Torque (N-m)	Failure Mode
Medial			
	R	9.47	Broken Screw
	L	9.83	Broken Screw
	R	9.55	Broken Screw
	R	10.21	Broken Screw
Standard Deviation		9.76 ± 0.33	
Lateral			
	L	6.11	Threads Stripped
	R	5.96	Threads Stripped
	L	5.67	Threads Stripped
	L	5.79	Threads Stripped
Standard Deviation		5.88 ± 0.19	

R=right, L=left

Table 3.2 Torque to failure values and failure modes of 4.5 mm AO screws placed in lag fashion used for repair of experimentally created 12 mm thick medial condylar fractures in equine MC3 bones.

Approach	Leg	Failure Torque (N-m)	Failure Mode
Medial			
	R	10.33	Broken Screw
	L	9.61	Broken Screw
	L	9.29	Broken Screw
	R	8.56	Broken Screw
Standard Deviation		9.45 ± 0.73	
Lateral			
	L	9.68	Broken Screw
	R	10.37	Threads Stripped
	R	9.89	Broken Screw
	L	10.14	Threads Stripped
Standard Deviation		10.0 ± 0.30	

R=right, L=left

Table 3.3 Torque to failure values and failure modes of 4.5 mm AO screws placed in lag fashion used for repair of experimentally created 20 mm thick medial condylar fractures in equine MC3 bones.

Approach	Leg	Failure Torque (N-m)	Failure Mode
Medial			
	R	10.54	Broken Screw
	L	10.39	Broken Screw
	R	9.72	Broken Screw
	R	9.96	Broken Screw
Standard Deviation		10.15 ± 0.38	
Lateral			
	L	9.93	Broken Screw
	R	9.87	Broken Screw
	L	10.67	Broken Screw
	L	9.80	Broken Screw
Standard Deviation		10.07 ± 0.41	

R=right, L=left

3.5 Discussion

The results of torque to failure testing indicate a distinct difference in failure for both the three fragment thicknesses and approach used. The differences are likely due to the amount of bone engaged by the screw with varying fragment thicknesses and surgical approach. These differences could impact the success of a clinical case.

We have concluded that there is a significant difference in the maximum torque to failure between a medial and lateral approach in the 8 mm fragment group. Failure torque values for the 8 mm fragment (Table 3.1) using the medial approach were significantly greater, almost twice that of the lateral approach. While the mean torque to failure was 9.76 N-m for the medial approach, it was only 5.88 N-m for the lateral approach.

It is concerning that the mean torque to failure for the lateral group was only slightly higher than the insertional torque of 5.4N-m.. Because insertional torque routinely exceed 5.4 N-m this minimal difference highlights a substantial risk of screw stripping when a screw is tightened under common clinical conditions.. This is supported by the failure mode for the 8 mm fragments. While each 8 mm fragment compressed from a traditional medial approach failed by screw breakage, the screw stripped in each of the lateral approach specimens indicating a lack of adequate purchase in the bone. The 4.5 mm cortical screw used here has a thread pitch, the distance from one thread to the next, of 1.75 mm. Given this value, a thread hole created in 8, 12 and 20 mm fragments will engage 4.6, 6.9 and 11.4 threads, respectively, a step-wise increase of 50% and 67%. These values indicate a minimum acceptable number of threads engaged to decrease the risk of screw breakage.

For the 12 mm fragments there was no significant difference in the failure torque between the medial and lateral approach. Also the failure mode revealed both screw breakage and stripping for both medial and lateral approaches.

For the 20 mm fragments there was no significant difference in the failure torque between the medial and lateral approach. The failure mode (Table 3.3) for all specimens in this group was screw breakage. Therefore the screws were engaged in enough bone that the screw failed instead of stripping when maximally tightened.

The results of this study do not support a definitive acceptable minimum for fragment size in the case of a lateral approach for a medial condylar fracture. They do, however, support the conclusion that the risk of stripping a screw at insertion is great when the fragment containing the thread hole engages few screw threads as is the case in these 8 mm fragments. Clinically, the medial condylar fracture is more likely to propagate from an axial location on the articular surface and therefore, be a thicker fragment in the 12-20mm range. This lessens but does not completely mitigate the risk of failure by the screw at insertion under normal conditions and torque.

The use of the lateral approach for medial condylar fractures has been used successfully and in the majority of clinical situations use of the approach is supported by the data presented here. It remains to be determined if the risks associated with the approach outweigh the benefits.

CHAPTER 4
FINAL DISCUSSION AND CONCLUSION

4.1 Summary

In summary, we have compared a medial vs. lateral approach of the repair of medial condylar fractures by measuring the interfragmentary compression and torque to failure for 3 different simulated fracture fragment thicknesses in adult equine third metacarpal bones. We have concluded that there is no significant difference in compression between a medial and lateral approach for a medial condylar fracture with an 8 or 12 mm thick fragment. A possible explanation is that the screws were not tightened to a maximal torque. This maximal torque would have most likely been different for medial and lateral approaches due to the differences in the number of threads engaged in the bone. This in turn would have most likely resulted in differences in interfragmentary compression between the medial and lateral approaches.

There is a significant difference in the torque to failure between a medial and lateral approach for a medial condylar fracture. A possible explanation for this is due to the amount of screw that is engaged into the bone with the varying fragment thicknesses. The larger the fragment the more surface area of bone that is in contact with the screw causes more friction thus causing the screw to fail due to threads stripping before reaching the maximal torque because the amount of the screw that is engaged with the bone. The failure torque for the 8 mm fragments is not much greater than the insertional torque of 5.4 Nm. The insertional torque of 5.4 N m is not an adequate amount of torque used to tighten screws in a repair of a condylar fracture. Screws are typically tightened farther than 5.4 N m of torque and tightened closer to the maximal torque around 9 N m. Thus in conclusion, the lateral approach may be an acceptable alternative for repair of the medial condylar fracture of at least 12mm in thickness.

BIBLIOGRAPHY

- Stashack TS (ed.): *Adam's Lameness in Horses* (4th ed.). Philadelphia: Lea&Febiger, 1987.
- Adams T, Blevins, and Shamis. Surgical Repair of Metacarpal Condylar Fractures with Palmar Osteochondral Communion in Two Thoroughbred Horses. *Vet Surg* 1985;14(1): 4.
- Alexander J and Rooney JR. Biomechanics, surgery, and prognosis of equine fractures 1967-1971. *Proc Am Assoc Equine Pract* 1972;19: 18.
- Auer JA, Stick J (eds). *Equine Surgery* (3rd ed.), St. Louis: Saunders, 2006.
- Barr ARS, Sridhar B, Denny HR. Long incomplete longitudinal fractures of the third metacarpal and metatarsal bone in horses. *Vet Rec* 1989;124: 3.
- Bassage LH and Richardson DW. (1998). Longitudinal fractures of the condyles of the third metacarpal and metatarsal bones in racehorses: 224 cases (1986-1995). *J Am Vet Med Assoc* 1998; 212(11): 1757-64.
- Bathe AP. (1994). Fractures in thoroughbred racehorses: results of a 2-year prospective study in Newmarket. *Proc Am Assoc Equine Pract* 1994; 40: 2.
- Batson EL, Reilly GC, Currey JD, et al. Postexercise and positional variation in mechanical properties of the radius in young horses. *Equine Vet J* 2000; 32: 95-100.
- Bertone AL: Distal limb: fetlock and pastern. In Hinchcliff KW, Kaneps AJ, Geor RJ (eds): *Equine Sports Medicine and Surgery: Basic and Clinical Sciences of the Equine Athlete*. Edinburgh: Saunders, 2004:289-318.
- Boyde A, Haroon Y, et al. Three dimensional structure of the distal condyles of the third metacarpal bone of the horse. *Equine Vet J* 1999;31(2): 122-129.

Brinker P (ed.). *Handbook of Small Animal Orthopedics and Fracture Treatment*, Philadelphia: W. B. Saunders, 1990.

Colton C: History of Osteosynthesis. In Texhammar R, Colton C (eds): *AO/ASIF Instruments and Implants: A Technical Manual*. New York: Springer-Verlag, 1981: 3-4.

DM, M. Lateral condylar fractures of the metacarpus and metatarsus in horses. *Proc. Am Assoc Equine Pract* 1976:22: 8.

DW, R. Medial condylar fractures of the third metatarsal bone in horses. *Am Vet Med Assoc* 1984:185: 5.

Dyce, S., Wensing (2002). *Veterinary Anatomy*, Saunders.

Ellis DR. Some observations on condylar fractures of the third metacarpus and third metatarsus in young thoroughbreds. *Equine Vet J* 1994:26(3): 178-83.

El Shorafa WM, Feaster JP, Ott EA: Horse metacarpal bone: age, ash content, cortical area and failure stress interrelationships. *J Anim Sci* 1979; 49: 979-982.

Fischer KJ, Jacobs CR and Carter DR. Computational methods for determination of bone and joint loads using bone density distributions. *Biomech* 1995: 28: 9.

Fürst A, Meier D, Michel S, et al: Effect of age on bone mineral density and micro architecture in the radius and tibia of horses: An Xtreme computed tomographic study. *BMC Vet Res* 2008; 4:3.

Galuppo LD, Simpson EL, et al. A clinical evaluation of a headless, titanium, variable-pitched, tapered, compression screw for repair of nondisplaced lateral condylar fractures in thoroughbred racehorses. *Vet Surg* 2006: 35(5): 423-30.

GE Fackelman, D. N. *Manual of Internal fixation of the Horse*.

Gilman, M. A. (1983). "Injuries on the racetrack." *Am Assoc Equine Pract*(1): 3.

Goodrich, L. R. (2004). Spiraling medial condylar fractures. Proceedings of the 14th Annual American College of Veterinary Surgeons Symposium, Columbia, MO.

Gregory L. Ferraro, D. (1990). "Lameness Diagnosis and Treatment in the Thoroughbred Racehorse." *Veterinary Clinics of North America:Equine Practice* **6**(1): 63-84.

Hans K. Uthoff (August 1973). "Mechanical Factors Influencing the Holding Power of Screws in Compact Bone." *The Journal of Bone and Joint Surgery* **55 B**(No. 3): 633-639.

Hilary M. Clayton, P. F. F., Diana S. Rosenstein (2005). *Clinical Anatomy of the Horse*, Mosby Elsevier.

Ian M. Wright, M. R. W. S. (2009). "A Lateral Approach to the Repair of Propagating Fractures of the Medial Condyle of the Third Metacarpal and Metatarsal Bone in 18 Racehorses." *Veterinary Surgery* **38**: 7.

John, C. J. (2009). "An In Vitro Biomechanical Comparison of Dynamic Condylar Screw Plate Combined with a Dorsal Plate Fixation of Distal Diaphyseal Radial Osteotomies in Adult Horses." *Veterinary Surgery* **38**: 13.

Johnson AL, Houlton JEF, Vannini R (eds). *AO Principles of Fracture Management in the Dog and Cat*. Stuttgart:Thieme, 2005.

Johnson BJ, Stover SM, Daft BM, et al: Causes of death in racehorses over a 2 year period. *Equine Vet J* 1994;26:327-330.

Jorg A. Auer, J. A. S. (1999). *Equine Surgery*, W. B. Saunders.

Joseph Schatzker, M., B.Sc. (MED.), F.R.C.S. (C), J.G. Horne, M.B. CH.B., and G. Sumner-Smith, B.V.Sc., F.R.C.V.S (1975). "The Reaction of Cortical Bone to Compression by Screw Threads." *Clinical Orthopaedics and Related Research* **111**: 263-265.

Kane, A. J., S. M. Stover, et al. (1996). "Horseshoe characteristics as possible risk factors for fatal musculoskeletal injury of thoroughbred racehorses." *Am J Vet Res* **57**(8): 1147-52.

Kane AJ, Stover SM, Gardner IA, et al: Hoof size, shape, and balance as possible risk factors for catastrophic musculoskeletal injury of Thoroughbred racehorses. *Am J Vet Res* 1998;59:1545-1552.

Kainer RA: Functional Anatomy of Equine Locomotor Organs. In Stashak TS (ed): *Adams Lameness in Horses*. Philadelphia: Lippincott, Williams, and Wilkins, 2002:1-72.

LeJeune SS, MacDonald MH, Stover SM, et al: Biomechanical Investigation of the Association Between Suspensory Ligament Injury and Lateral Condylar Fracture in Thoroughbred Racehorses. *Vet Surg* 2003;32:585-597.

Lewis C. R. Smith, T. R. C. G., and Andrew P. Bathe (2009). "A Lateral Approach for Screw Repair in Lag Fashion of Spiral Third Metacarpal Medial Condylar Fractures in Horses." *Veterinary Surgery* **38**: 8.

Martin, G. S. (2000). "Factors associated with racing performance of Thoroughbreds undergoing lag screw repair of condylar fractures of the third metacarpal or metatarsal bone." *JAVMA* **217**(12): 8.

McKee, S. L. (1995). "An update on racing fatalities in the UK. ." *Equine Veterinary Education* **7**: 3.

Meagher, D. (1976). "Lateral condylar fractures of the metacarpus and metatarsus in horses" *Proc Am Assoc Equine Pract* **22**: 8.

Mueller (1999). *Fracture Healing*.

Nixon, A. J. (1996). *Equine Fracture Repair*. Philadelphia, W. B. Saunders Company.

Nunamaker, D. M. (2000). *General techniques and biomechanics*, AO Publishing.

Parkin, T. D., P. D. Clegg, et al. (2006). "Catastrophic fracture of the lateral condyle of the third metacarpus/metatarsus in UK racehorses - fracture descriptions and pre-existing pathology." *Vet J* **171**(1): 157-65.

Parkin, T. D. H., Clegg, P. D., French N. P., Proudman, C.J., Riggs, C.M. Singer, E.R., Webbon, P.M., Morgan, K.L. (2004). "Risk of fatal distal limb fractures among thoroughbreds involved in the five types of racing in the United Kingdom." *Veterinary Records* **154**: 5.

Perren SM, Buchanan J: Basic AO/ASIF Technique: Aims and Principles. In Texhammar R, Colton C (eds): *AO/ASIF Instruments and Implants: A Technical Manual*. New York: Springer-Verlag, 1981: 26.

Pool, R. R. and D. M. Meagher (1990). "Pathologic findings and pathogenesis of racetrack injuries." *Vet Clin North Am Equine Pract* **6**(1): 1-30.

Pool, R. R., McIlwraith, C.W. (1995). "Necropsy of Prairie Bayou: A look at the facts." *Proc Am Assoc Equine Pract* **41**: 3.

Radtke CL, Danova NA, Scollay MC, et al: Macroscopic changes in the distal ends of the third metacarpal and metatarsal bones of Thoroughbred racehorses with condylar fractures. *Am J Vet Res* 2003;64:1110-1116. R.D. Frandson, W. L. W., Anna Dee Fails (2003).

Anatomy and Physiology of Farm Animals, Lippincott Williams and Wilkins.
Rice, J. L., Cowin, S.C., and Bowman, J.A. (1988). "On the dependence of the elasticity and strength of cancellous bone on apparent density." *Biomech* **21**: 14.

Richardson DW: The Metacarpal and Metatarsal Bones. In Auer JA, Stick JA (eds): *Equine Surgery*. Philadelphia: Elsevier, 2006:1238–1253.

Rick, M. C., T. R. O'Brien, et al. (1983). "Condylar fractures of the third metacarpal bone and third metatarsal bone in 75 horses: radiographic features, treatments, and outcome." *J Am Vet Med Assoc* **183**(3): 287-96.

Riggs, C. M. (1999). "Aetiopathogenesis of parasagittal fractures of the distal condyles of the third metacarpal and third metatarsal bones--review of the literature." *Equine Vet J* **31**(2): 116-20.

Riggs, C. M., G. H. Whitehouse, et al. (1999). "Pathology of the distal condyles of the third metacarpal and third metatarsal bones of the horse." *Equine Vet J* **31**(2): 140-8.

- Riggs, C. M., G. H. Whitehouse, et al. (1999). "Structural variation of the distal condyles of the third metacarpal and third metatarsal bones in the horse." *Equine Vet J* **31**(2): 130-9.
- Rooney, J. R. (1969). *Biomechanics of Lameness in Horses*. Baltimore, Williams and Wilkins.
- Rooney, J. R. (1974). "Distal condylar fractures of the cannon bone in the horse." *Mod Vet Pract*: 113-114.
- Russell TM, M. A. (2006). "Standing surgical repair of propagating metacarpal and metatarsal condylar fractures in racehorses." *Equine Vet J* **38**: 5.
- Smith, L. C., T. R. Greet, et al. (2009). "A lateral approach for screw repair in lag fashion of spiral third metacarpal and metatarsal medial condylar fractures in horses." *Vet Surg* **38**(6): 681-8.
- Stepnik MW, Radtke CL, Scollay MC, et al: Scanning electron microscopic examination of third metacarpal/third metatarsal bone failure surfaces in thoroughbred racehorses with condylar fracture. *Vet Surg* 2004;33:2-10.
- Stover, S., Read DH, Johnson BJ (1994). "Lateral condylar fracture histomorphology in racehorses." *Proc Am Assoc Equine Pract* **40**: 2.
- Texhammar R: AO/ASIF Instrumentation. In Texhammar R, Colton C (eds): *AO/ASIF Instruments and Implants: A Technical Manual*. New York: Springer-Verlag, 1981: 73-78. Uhl RL: The biomechanics of screws. *Orthop Rev* 1989; 18(12):1302-1307.
- T.M. Russell, A. A. M. (2006). "Standing surgical repair of propagating metacarpal and metatarsal condylar fractures in racehorses." *Equine Vet J* **38**: 5.
- Thomas P. Ruedi, W. M. M. (2000). *AO Principles of Fracture Management*, Thieme.
- Turner, A. (1977). "Surgical repair of fractures of the third metatarsal bones in a Standardbred gelding." *J Am Vet Med Assoc* **171**: 4.

Wilson JH, H. S., Jensen RC (1993). "Injuries sustained during racing at racetracks in the U.S." *Proc Am Assoc Equine Pract* **39**: 2.

Wright, I. M. and M. R. Smith (2009). "A lateral approach to the repair of propagating fractures of the medial condyle of the third metacarpal and metatarsal bone in 18 racehorses." *Vet Surg* **38**(6): 689-95.

Zekas, L. J., L. R. Bramlage, et al. (1999). "Characterisation of the type and location of fractures of the third metacarpal/metatarsal condyles in 135 horses in central Kentucky (1986-1994)." *Equine Vet J* **31**(4): 304-8.

Zekas, L. J., L. R. Bramlage, et al. (1999). "Results of treatment of 145 fractures of the third metacarpal/metatarsal condyles in 135 horses (1986-1994)." *Equine Vet J* **31**(4): 309-13.

VITA

Saybl Sprinkle was born in Austin, Texas, in 1986. At the age of three her dad gave her a mean Tom cat that she named Wilbur, her fascination with animals grew from that first experience of owning and caring for her first pet. This was followed with an abundance and variety of species that only further expanded her passion for all shapes and sizes of animals. She matured and her fondness for animals grew during her High school years when she joined an organization known as FFA where she learned leadership, showmanship, and animal husbandry skills. She participated in leadership competitions, meetings, conventions, and livestock shows at the local, regional and state level. Saybl attended Louisiana State University for both her undergraduate and graduate studies. Her goal throughout school was to attend veterinary school and has plans to apply to veterinary schools around the country. Through her graduate program she developed a wonderful mentorship with her mentors and a wonderful friendship with a fellow grad student Niki Marie Hansen.