

2012

Identification of algal strains by PCR amplification and evaluation of their fatty acid profiles for biodiesel production

Reynaldo Moreno

Louisiana State University and Agricultural and Mechanical College, rmoren1@tigers.lsu.edu

Follow this and additional works at: https://digitalcommons.lsu.edu/gradschool_theses



Part of the [Life Sciences Commons](#)

Recommended Citation

Moreno, Reynaldo, "Identification of algal strains by PCR amplification and evaluation of their fatty acid profiles for biodiesel production" (2012). *LSU Master's Theses*. 2479.

https://digitalcommons.lsu.edu/gradschool_theses/2479

This Thesis is brought to you for free and open access by the Graduate School at LSU Digital Commons. It has been accepted for inclusion in LSU Master's Theses by an authorized graduate school editor of LSU Digital Commons. For more information, please contact gradetd@lsu.edu.

**IDENTIFICATION OF ALGAL STRAINS BY PCR AMPLIFICATION AND
EVALUATION OF THEIR FATTY ACID PROFILES FOR BIODIESEL PRODUCTION**

A Thesis
Submitted to the Graduate Faculty of the
Louisiana State University and
Agricultural and Mechanical College
In partial fulfillment of the
Requirements for the degree of
Master of Science

In

The Department of Food Science

by
Reynaldo Moreno Uclés
B.S., Agricultural Engineering and Business Administration
Universidad de San Pedro Sula, Honduras, 2002
B.M., Music
Louisiana State University, 2008
May, 2012

DEDICATION

This thesis is dedicated to my parents Reynaldo Moreno and Carmen Suyapa Uclés de Moreno who provided their never-ending support, love, inspiration, encouragement to achieve my goals in life, and their efforts to provide me with the best preparation and education possible. Likewise, I extend this dedication to my sisters Karen Suyapa, and Gina Maria, and to my brother Carlos Armando for caring as well and for being such outstanding role models, and to my nieces Karen Daniela, Grace Marie and my nephews Mario Alfredo, Diego Ernesto and Luc Wyatt.

ACKNOWLEDGEMENTS

First and foremost, I would like to thank God the creator, as I believe He is the provider of the blessings, the one who has allowed for all of this to happen, for me to be here and be able to accomplish this goal.

I would like to express my deepest gratitude to my advisor Dr. Giovanna M. Aita, for her financial support, patience, guidance, motivation and inspiration to excel. I am extremely thankful for all the advice I have received from her through these two years and for this opportunity, which allows me to culminate such an important milestone in my professional life.

I extend my appreciation to my committee members Dr. Donal Day and Dr. Zhimin Xu for their help and suggestions.

I would specially like to thank Dr. Dina L. Gutierrez and the Comparative Biomedical Science department at the School of Veterinary Medicine at Louisiana State University for the invaluable help, knowledge and expertise in molecular biology. Without her guidance, support and patience, I would have not been able to complete this project. Special thanks are extended to Dr. Shaomian Yao for allowing me to perform a significant portion of my research in their facility.

Special thanks go to Dr. Barry Hurlburt and Suzanne Brashear at the United States Department of Agriculture- Agricultural Research Service in New Orleans for their collaboration which greatly assisted me on this project.

My gratitude also goes to everyone at the Audubon Sugar Institute, in particular to Dr. Lee Madsen for his patience, support, and advice, Dr. Misook Kim, Dr. Michelle Walker, Dr. Deepti Salvi, Dr. Derek Dorman and Diana Roberts, Melati Tessier, Irina Tischeckina, Julie King, Stuart L. Goudeau, Adam Hoogewind, Christian Lohrey as well as my laboratory mates Akanksha, Shuo, Zenghui, Cong and Saeed for their help through these years.

I would also like to thank Dr's. Jack Losso, Joan King, Subramaniam Sathivel and other faculty members in Food Science Department who had perhaps, inadvertently helped me too. Special thanks go to Terri Gilmer for her advice and support.

My gratitude is also extended to Dr. Matthew Brown from the Socolofsky Microscopy Center in the Department of Biological Sciences at Louisiana State University for his help with confocal microscopy, Phillip Riekenberg and Alvaro Armas at the Department of Oceanography and Coastal Sciences for the spectroradiometer, and Javed Iqbal at the Callegari Institute, Dr. Damien Narcisse and Alfredo Prudente in the Food Science Department for their help with the fatty acids analysis.

My sincere thanks to all my friends from the Food Science Department, Christ the King Church, Zachary United Methodist Church, Group "Fortaleza de Dios" in Honduras, friends like Marjan, Jose Daniel, Jose Pablo and Kenneth who have helped me out with this project in one way or another, and with whom I shared good times, and to all the wonderful people I have been able to meet during the last few years whose support has helped make this project a reality, and finally, to my relatives in the United States for their encouragement and for being there for me at all times.

TABLE OF CONTENTS

DEDICATION	ii
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	viii
LIST OF ABBREVIATIONS	ix
ABSTRACT	x
1. LITERATURE REVIEW	1
1.1. Introduction	1
1.2. Microalgae as a Source for Biodiesel	3
1.3. Photosynthesis in Microalgae	5
1.4. Microalgae Classification	6
1.4.1. Diatoms	6
1.4.2. Green Algae	7
1.4.3. Golden-Brown Algae	7
1.4.4. Prymnesiophytes	7
1.4.5. Eustigmatophytes	8
1.4.6. Cyanobacteria	8
1.5. Growth Requirements of Microalgae	8
1.5.1. Light	9
1.5.2. Temperature	10
1.5.3. Carbon Dioxide and Organic Carbon Sources	10
1.5.4. Nutrients	12
1.5.4.1. Nitrogen	12
1.5.4.2. Phosphorus	13
1.5.4.3. Other Macro, Micronutrients and Chelates	13
1.5.5. Salinity	14
1.6. Microalgae Identification	15
1.6.1. Polymerase Chain Reaction	15
1.6.2. Enhancing Agents	16
1.6.3. Magnesium Concentration	17
1.6.4. Amplification of 16S rRNA, 18S rRNA and Other Genes	17
1.7. Microalgae Lipid Content	20
1.8. Goal of This Study	21
2. MATERIALS AND METHODS	23
2.1. Sampling and Collection of Organisms	23
2.2. Growth Conditions	23
2.3. Colony Selection	24
2.4. Colony Homogenization	25
2.5. Microorganism Selection and Harvesting	25
2.6. Genomic DNA Extraction	26
2.7. DNA Purification	27

2.8. Standard Polymerase Chain Reaction (PCR) and Optimization	27
2.9. Gel Electrophoresis	28
2.10. Gel Extraction and Purification	30
2.11. Cloning of PCR Products	30
2.11.1. LB Agar Plates and Broth Media Preparation	30
2.11.2. Colony Screening	30
2.11.3. Cloning Reaction and Transformation with Chemically Competent <i>E. coli</i>	31
2.12. Purification of Plasmid DNA	31
2.13. Restriction Enzyme Digestion	32
2.14. Sequencing	32
2.15. Lipid Characterization and Quantification	33
2.15.1. Algae Inoculum Density	33
2.15.2. Growth Conditions	33
2.15.3. Lipid Extraction and Chlorophyll Determination	34
2.15.4. Fatty Acids Analysis	35
3. RESULTS	37
3.1. Algal Isolation	37
3.2. DNA Extraction	37
3.3. PCR Products	38
3.4. PCR Optimization	38
3.5. Purification of Plasmid DNA	38
3.6. Restriction Enzyme Digestion	38
3.7. Sequencing	39
3.8. BLAST	40
3.9. Algae Inoculum	40
3.10. Biomass for Lipid Extraction	41
3.11. Total Extracted Lipid and Chlorophyll Content	41
4. DISCUSSION	45
5. CONCLUSIONS	56
REFERENCES	58
APPENDIX 1. F/2 MEDIA COMPOSITION	72
APPENDIX 2. BLUE-GREEN 11 MEDIA COMPOSITION	73
APPENDIX 3. BOLD'S BASAL MEDIA COMPOSITION	74
APPENDIX 4. SAMPLES DILUTIONS FOR HEMACYTOMETER	75
APPENDIX 5. COUNTING PROCEDURES FOR HEMACYTOMETER READINGS	76
APPENDIX 6. FREEZE-DRYER PROGRAM	77
VITA	78

LIST OF TABLES

Table 1.1. Comparison of some sources of biodiesel	4
Table 1.2. Fatty acid composition of laboratory batch cultures of <i>Chaetoceros</i> sp. (CS256) grown at different temperatures	11
Table 1.3. Lipid content of some microalgae	21
Table 2.1 Salinities of sampling locations through total conductivity.	24
Table 2.2. 16S rRNA and 18S rRNA primers used for PCR reactions.	28
Table 2.3. Reaction volumes and concentrations for PCR master mixes.	29
Table 2.4. PCR cycle conditions for 16S rRNA.	29
Table 2.5. PCR cycle conditions for 18S rRNA.	29
Table 3.1. Codes for isolated microorganisms with tentative identification based on morphology, similarity between amplified sequences and nearest recorded sequence in NCBI using BLAST and accession numbers for NCBI sequences.	41
Table 3.2. Algae inoculums for Batch A and Batch B in cells ml ⁻¹	41
Table 3.3. Total dry biomass for lipid extraction.....	42
Table 3.4. Total lipid extracted from biomass (g/100 g dry biomass).....	43
Table 3.5. Fatty acid profiles for evaluated strains grown in air (0.037 % CO ₂) or 5 % CO ₂ in mg g ⁻¹ dry biomass.	44
Table 4.1. Flue gas composition of Wyodak PC power plant.	53
Table 4.2. Typical untreated flue gas composition from a power plant burning low sulfur eastern bituminous coal.	53

LIST OF FIGURES

Figure 1.1 Transesterification Reaction	3
Figure 1.2. Algal Biomass Product Streams	5
Figure 2.1 Map of Sampling Locations	23
Figure 2.2. Tentatively Identified Microorganisms. FC2: diatom, N3: cyanobacteria, FC1 and D2: green algae.	26
Figure 3.1. Microscopic Images for A) N3: <i>Synechococcus sp.</i> , B) FC2: <i>Sellaphora pupula</i> , C) FC1: <i>Chlorella sorokiniana</i> , D) D2: <i>Scenedesmus abundans</i> , and E) <i>Chlorella vulgaris</i> UTEX 259 (control) as identified by BLAST.	37
Figure 3.2. PCR Amplification Results Gel. PCR product quality of nine conditions for N3 (A), FC2 (B), FC1 (C), and D2 (D) mixed with MgCl ₂ at 1.5 mM (low concentration, L), 3 mM (medium concentration, M), and 4.5 mM (high concentration, H).	39
Figure 3.3. Restriction Enzyme Digestion Gel.	39

LIST OF ABBREVIATIONS

CLS-Y: Cell Lysis Solution for yeast, algae or fungi containing 1-5 % SDS and 1-5 % DL-Dithiothreitol.

SEWS: Washing solution containing 10-20 % TRIS hydrochloride, less than 1 % TRIS base and 95 % ethanol.

DES: Ultra-pure water DNA elution solution.

TE: 10 mM Tris-HCl and 1 mM EDTA elution buffer (pH 8.0).

QG: pH indicator buffer for optimal DNA binding containing guanidine thiocyanate at 50-100%.

PE: Wash buffer containing ethanol at 70-80 %.

EB: Elution buffer containing 10 mM Tris-Cl (pH 8.5).

P1: Resuspension buffer containing 50 mM Tris-Cl and 10 mM EDTA.

P2: Lysis buffer containing 200 mM NaOH and 1 % SDS.

N3: Purification buffer containing 25-50 % guanidium chloride and 10-25 % acetic acid.

PB: Binding buffer for DNA clean-up containing guanidinium chloride at 25-50 % and propan-2-ol at 25-50 %.

ECO RI buffer: Restriction endonuclease buffer containing 15.8 % Tris buffered with HCl, 3 % NaCl, and 1 % MgCl.

SOC Medium: Medium containing 2 % tryptone, 0.5 % yeast extract, 10 mM sodium chloride, 2.5 mM potassium chloride, 10 mM magnesium chloride, 10 mM magnesium sulfate and 20 mM glucose per liter.

PNNL: Pacific Northwest National Laboratory.

EN 14214: European Standard that describes the requirements and test methods for FAME.

ASTM D6751: American Society for Testing and Materials (ASTM) standards and specifications for biodiesels blended with middle distillate fuels.

ABSTRACT

Four microalgae, obtained from different brackish and fresh water sources within the state of Louisiana's Southeast region, were evaluated for their potential use in the production of biodiesel. The microorganisms were isolated and identified using genomic DNA, and 16S rRNA or 18S rRNA gene amplification followed by sequencing. The resultant sequences were compared with those available on the NCBI website database through the BLAST bioinformatic tool. The results showed high correlation with known nucleotide sequence identities at 99 % with *Synechococcus* sp., 98 % with *Sellaphora pupula*, 99 % with *Chlorella sorokiniana*, 99 % with *Scenedesmus abundans*, and 99 % with *Chlorella vulgaris* (control).

The fatty acid profiles of the identified organisms grown using 5 % CO₂ aeration into the growth media were evaluated and were found to be different to the control group (0.037 % CO₂). In *Synechococcus* sp., total fatty acids (TFA) decreased from 20.63 g kg⁻¹ to 17.62 g kg⁻¹ dry biomass with the appearance of C18:2 and C18:3, which were absent in the control. TFA from *Sellaphora pupula* decreased from 54.8 g kg⁻¹ to 24.4 g kg⁻¹ dry biomass and contained the greatest C16:1. The extracted TFA from *Scenedesmus abundans* increased from 14.14 g kg⁻¹ to 31.63 g kg⁻¹ and displayed the highest content of C18:1. For *Chlorella vulgaris* UTEX 259, TFA content increased from 15.14 g kg⁻¹ to 47.83 g kg⁻¹, and 50 % of that total was C18:3. The TFA content from *Chlorella sorokiniana* decreased from 29.82 g kg⁻¹ to 23.99 g kg⁻¹; however, it had a lower C18:3 which allowed for a balanced fatty acid profile in terms of cetane number, oxidative stability, viscosity and low temperature conditions. The aforementioned conditions, plus owning the best biomass productivity when using 5 % CO₂, deemed *Chlorella sorokiniana* as the best candidate of the strains evaluated for the production of biodiesel.

Keywords: microalgae• biodiesel• fatty acid methyl esters• carbon dioxide• *Chlorella sorokiniana*

1. LITERATURE REVIEW

1.1. Introduction

The rising need for energy in developing nations is giving place to vehement competition for the world's decreasing energy resources (Pienkos and Darzins, 2009). The increased use of fossil fuels results in larger greenhouse gases (GHG) emissions, and this is usually considered the main reason for global climate change (Wuebbles and Atul, 2001). Fossil fuels are the largest contributor of GHGs to the atmosphere (EIA, 2006). With the increase in anthropogenic GHG emissions due to the extensive use of fossil fuels for transport, new techniques for the development of electricity and thermal energy generation are needed (Brennan and Owende, 2010). The goal of a 5.2 % reduction in GHG emissions worldwide from 1990 values was proposed during The Kyoto Protocol back in 1997 (Wang et al., 2008). One hundred and ninety-three countries are currently part of this protocol, with the United States being the only remaining signatory nation that has not ratified it (Status of ratification of Kyoto Protocol, 2011). Greater use of biofuels, which compete and have partially displaced petroleum based fuels for use in transportation, could help meet that reduction in emissions objective (Macedo et al., 2008).

In terms of CO₂ reduction, first generation biofuels are said to have a limited performance and demand the use of large amounts of land (Schubert, 2006) and have now reached economic levels of production. Examples are food and oil crops (e.g. biodiesel from rapeseed oil, and ethanol from sugarcane, sugar beet, and corn) (FAO, 2008) as well as animal fats (FAO, 2007). Strong controversy surrounds the use of first generation biofuels, usually due to their negative impacts like contributing to the increase in food prices, deforestation and biodiversity losses,

hence, the extent of their ultimate contribution to the reduction of GHG is frequently questioned (Gomez et al., 2011).

Second generation biofuels are produced from lignocellulosic biomass (e.g. agricultural waste products, forest residues, municipal waste) (Aita and Kim, 2010). Technologies for the conversion of lignocellulosic biomass into biofuels have yet to reach the scales for commercialization (FAO, 2008). In order for a biofuel resource to be technically and economically viable, it needs to compete with petroleum fuels' prices, improve air quality (e.g. through CO₂ sequestration) and require low usage of water and land (Khosla, 2009).

Microalgae are projected to be the source for the third generation of biofuels (Vieira Costa and Greque de Morais, 2011). High biomass productivity, accumulation of up to 20-50 % (w/w dry weight) triacylglycerols, the lack of requirement for high quality agricultural land and water renewal when compared to terrestrial crops are among the advantages to be offered by microalgae (Singh and Gu, 2010; Scott et al., 2010). Despite those advantages, several intrinsic and applied research and development obstacles need to be addressed for the commercialization of algal-based fuels at a cost that can compete with that of petroleum-based fuels (Pienkos and Darzins, 2009).

Like plants and some photosynthetic bacteria, algae are photosynthetic organisms, which efficiently utilize energy from the sun to convert water, CO₂ and O₂ into biomass (Sheehan et al., 1998). The most comprehensive research effort regarding the production of fuels from algae has been the "The Aquatic Species Program (ASP)" led by the Department of Energy (DOE) from 1978 to 1996. DOE invested approximately \$25 million to study a variety of aquatic species such as macroalgae, cattails and microalgae for use in renewable energy production, and several important advances in the technology arose from this effort (Sheehan et al., 1998).

Microalgae are photosynthetic organisms which can be prokaryotic or eukaryotic and can grow in a wide range of environmental conditions (Mata et al., 2010). Microalgae produce lipids in the form of triacylglycerols or triglycerides (Sheehan et al., 1998). These triglycerides can be reacted with an alcohol in a reaction called transesterification or alcoholysis (Figure 1.1) to obtain fatty acid methyl or ethyl esters for the production of biodiesel (Gouveia and Oliveira, 2009).




Figure 1.1 Transesterification Reaction.

1.2. Microalgae as a Source for Biodiesel

The use of microalgae as fuel feedstock was first proposed over 50 years ago for the production of methane gas (Meier, 1955). Yields for biodiesel production from microalgae can be 10 to 20 times higher than those obtained from oleaginous seeds and/or vegetable oils (Table 1.1). The oil content in some microalgae can be fairly high and can be induced to produce even higher concentrations of lipids through the implementation of low nitrogen media, varying Fe^{3+} concentration and increased light intensity (Illman et al., 2000; Liu et al., 2007; Rodolfi et al., 2007; Solovchenko et al., 2008; Tornabene et al., 1983). CO_2 removal from power plants through biofixation represents an interesting method for reducing GHG emissions and assisting in the increase of microalgal biomass, lipid and biodiesel yields (Wang et al., 2008).

Several different types of renewable fuels can be obtained from microalgae. Among those, we can include methane produced by anaerobic digestion of the algal biomass (Spolaore et al., 2006); biodiesel derived from microalgal oil (Roessler et al., 1994; Sawayama et al., 1995; Dunahay et al., 1996; Sheehan et al., 1998; Banerjee et al., 2002; Gavrilescu and Chisti, 2005); and biohydrogen which can be produced photobiologically (Ghirardi et al., 2000; Akkerman et al., 2002; Melis, 2002; Fedorov et al., 2005; Kapdan and Kargi, 2006).

Some microalgae have also a convenient fatty acids profile for transesterification and an unsaponifiable fraction which allows for the production of biodiesel with high oxidation stability (Dote et al., 1994; Ginzburg, 1993; Milne et al., 1990; Minowa et al., 1995) and physical and fuel properties (e.g. density, viscosity, acid value, and heating value) which are comparable to those found in fossil diesel (Miao and Wu, 2006; Rana and Spada, 2007).

Table 1.1. Comparison of some sources of biodiesel (Chisti, 2007).

Crop	Oil yield (L ha ⁻¹)
Corn	172
Soybean	446
Canola	1,190
Jatropha	1,892
Coconut	2,689
Palm	5,950
*Microalgae ^a	136,900
*Microalgae ^b	58,700

* Based on experimentally demonstrated biomass productivity in photobioreactors.

a 70% oil (by wt) in biomass

b 30% oil (by wt) in biomass

Another advantage consists on the potential use of the residual biomass (mainly carbohydrate and protein), collected after the removal of the lipid component for the generation of energy, liquid or gaseous fuels, or higher value by-products (Figure 1.2) (Pienkos and Darzins, 2009). When compared to fossil diesel, biodiesel has advantages in that it is renewable, biodegradable, and produces less SOx and particulate emissions when burned besides not

requiring engine modification for its usage (Sheehan et al., 1998). Microalgae can be an alternative option to produce fuels because of their versatility as biomass source. Microalgae have higher photosynthetic efficiency, higher biomass productivities, and faster growth rates than higher plants, as well as the highest CO₂ fixation and O₂ production rates, positioning microalgae as one of the Earth's most important renewable fuel crops (Campbell, 1997; Chisti, 2007; Chisti, 2008).




Figure 1.2. Algal Biomass Product Streams (Pienkos and Darzins, 2009).

1.3. Photosynthesis in Microalgae

Photosynthesis is a process where sunlight energy is used to convert CO₂ into glucose and oxygen. In this process CO₂ and light are converted into organic matter by photoautotrophs. Photosynthesis serves as a source of energy for metabolism and growth for basically all forms of life on Earth either in a direct or indirect way (Masojidek et al., 2004). Algae are photosynthetic,

non-vascular plants containing chlorophyll *a* as pigment (Vonshak and Maske, 1982). The chlorophyll *a*, functioning as the principal photochemically active compound, receives light in order for photosynthesis to occur. Hence, the content of this pigment in microalgae influences photosynthetic activity (MacIntyre et al., 2002). The concentration of this light harvesting pigment has an impact on biomass production in microalgae as well as on the accumulation of target products (Su et al., 2007).

Nearly half of the total photosynthesis taking place on Earth is associated with marine phytoplankton (Camacho et al., 2003). Light availability is related to the growth and production performances of a photosynthetic organism. When grown in a photobioreactor, the amount of light absorbed by an algal cell depends on the incident flux on the particular cell, culture density, and cell pigmentation as well as on the photosynthesis-irradiance relationship (the P-I curve). The P-I curve helps predict the culture performance and understand the relationship between the growth observed and the amount of light received (Camacho et al., 2003).

1.4. Microalgae Classification

The main groups of microalgae differ primarily in terms of pigment composition, biochemical constituents, ultrastructure, and life cycle. The groups include diatoms (Class Bacillariophyceae), green (Class Chlorophyceae), golden brown (Class Chrysophyceae), prymnesiophytes (Class Prymnesiophyceae), eustigmatophytes (Class Eustigmatophyceae), and blue-green or cyanobacteria (Class Cyanophyceae) (Sheehan et al., 1998).

1.4.1. Diatoms. Diatoms, with close to 100,000 known species, are among the most common group of algae in existence and tend to dominate the phytoplankton in oceans, but are also commonly found in fresh and brackish water habitats (Sheehan et al., 1998). The golden-brown color is due to the presence of fucoxanthin and β -carotene which mask the green color chlorophyll *a* and chlorophyll *c* (Tomaselli, 2004). Two major groups of diatoms can be

distinguished: the pennates having bilateral symmetry and the centrals with radial symmetry (Tomaselli, 2004). The main storage compounds of diatoms are lipids and a β -1,3-linked carbohydrate known as chrysolaminarin. Substantial amounts of polymerized silicate (Si) can be found in their cell walls (Sheehan et al., 1998).

1.4.2. Green Algae. Members of this group are among the most common microalgae, especially in freshwater (Neenan, 1986). Approximately 8,000 species are estimated to be in existence (Sheehan et al., 1998). This group contains chlorophyll *a* and chlorophyll *b* and several carotenoids which can be synthesized under stress conditions and make the alga change in color (Tomaselli, 2004). Starch is their primary storage component. In certain species, however, N-deficiency has been found to boost lipid accumulation (Sheehan et al., 1998). Higher plants are considered descendants of the green algae, and for that reason, have received more attention than other algal groups with *Chlorella* and *Chlamydomonas* being two commonly studied genera (Sheehan et al., 1998).

1.4.3. Golden-Brown Algae. This group is also known as chrysophytes and shares similarities in biochemical and pigment composition with diatoms. Approximately 1,000 species are known, and are primarily present in freshwater habitats (Sheehan et al., 1998), especially in oligotrophic waters low in calcium (Tomaselli, 2004). Major carbon storage forms in this group are usually lipids and chrysolaminarin (Sheehan et al., 1998).

1.4.4. Prymnesiophytes. They are also known as haptophytes. Prymnesiophytes are constituted by approximately 500 species (Sheehan et al., 1998). They are mostly found in the oceans, and form a major part of marine phytoplankton (Tomaselli, 2004). Just like in diatoms and chrysophytes, fucoxanthin imparts a brown color to the cells, with both lipids and chrysolaminarin constituting the major storage products (Sheehan et al., 1998). Scales embedded

in a mucilage can be usually found covering the cells and these can sometimes be calcified (Tomaselli, 2004).

1.4.5. Eustigmatophytes. This group encompasses a significant portion of the “picoplankton” (Sheehan et al., 1998). Eustigmatophytes include unicellular and coccoid organisms that produce a small number of zoospores and are mostly found in soil and fresh waters (Tomaselli, 2004). *Nannochloropsis*, a marine species within this class, is a source of polyunsaturated fatty acids with high amounts of eicosapentaenoic acid (Tomaselli, 2004).

1.4.6. Cyanobacteria. Cyanobacteria are prokaryotic organisms that contain chlorophyll *a* (Tomaselli, 2004). The name of blue–green has been given because of the presence of phycocyanin and phycoerythrin which usually masks the chlorophyll pigmentation (Tomaselli, 2004). They are similar to bacteria in that their cells lack a nucleus, chloroplasts, and also, have a different gene structure (Neenan et al., 1986; Sheehan et al., 1998). An estimated 2,000 species belong to this group. Some cyanobacteria are known to assimilate atmospheric nitrogen (N) thus eliminating the need to provide fixed N to the cells. None of them have been found to produce significant quantities of lipid as storage (Sheehan et al., 1998).

1.5. Growth Requirements of Microalgae

The biochemical composition of microalgae can be affected by such factors as growth rate, environmental conditions, and life cycle (Richmond, 1986). Microalgal growth and chemical composition are mainly controlled by light, temperature, available carbon dioxide, pH, and nutrients (Tzovenis et al., 1997, Zhu et al., 1997). Other factors, such as salinity, can be of vital importance to some species (Chu et al., 1996). As for synergistic or symbiotic effects, association with bacteria can be beneficial to algae for growth. *Azospirillum* sp. is a known plant-growth- promoting bacterium that aids in the growth and yields of many plants and can also promote the growth of several freshwater species belonging to the *Chlorella* genus (Bashan and

Levanony, 1990; Okon and La-bandera Gonzalez, 1994; Bashan and Holguin, 1997; Bashan et al., 2004). Increased nitrogen content, mass, length and yields have been shown in crops grown using strains of *Azospirillum lipoferum* (Govedarica et al., 1993; Govedarica et al., 1994; Favilli et al., 1993). Studies have found an accumulation of lipids with a greater variety of fatty acids when *Azospirillum brasilense* was co-immobilized in alginate beads in conjunction with certain *Chlorella* species (de-Bashan et al., 2002).

1.5.1. Light. The use of monocultures is required in various microalgal applications and controlled cultivation systems. This requirement has favored the development of closed photo bioreactors (Barbosa et al., 2001). Efficient utilization of light is one of the major challenges in microalgal biotechnology, especially when an increase in the biomass yield is desired (Barbosa et al., 2001). Photoacclimation or photoadaptation is a process that controls the effect of light on the biochemical composition of photosynthetic algae. During this process, algal cells undergo changes in cell composition and alterations in their ultrastructural, biophysical and physiological properties can also be observed (Dubinsky et al., 1995). Lee and Lee (2001) showed that cell concentration on a *Chlorella kessleri* inoculum with a density of 10^5 cells ml^{-1} increased to 1.6×10^7 cells ml^{-1} after three days of continuous light. However, the total cell concentration in flasks with L/D (light/dark) lighting scheme was increased to 9.1×10^6 cells ml^{-1} during the same period, only 57 % of that under continuous illumination. This difference in cell concentration between the two lighting schemes was proportional to the duration of the light exposure. The reduction in cell biomass under L/D lighting scheme seemed to be caused by night biomass loss by respiration. In one study using *Chaetoceros* sp. under a continuous photon flux of $125 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, provided by either a Cool White (CW) or by Gro-Lux and Gro-Lux wide spectrum agricultural lamps (GRO and GRO/WS), the protein concentration was found to be

higher whereas lipids were lower at the end of the exponential growth compared to the stationary phase (Sanchez and Voltolina, 2006). The concentration of proteins, however, remained consistently higher than other treatments with GRO lamps in both stages, possibly due to the “high emission of blue light”. As for lipids, the highest yields were observed in algae grown using CW; whereas, algae grown using GRO/WS yielded the best carbohydrate production during both growth phases.

1.5.2. Temperature. The effect temperature exerts on biochemical reactions and how it affects the biochemical composition of algae, makes temperature one of the most important environmental factors (Hu, 2004). Renaud et al. (2002) evaluated the effect of temperature on fatty acid composition of four tropical Australian microalgal species (the diatom *Chaetoceros* sp. (CS256); two cryptomonads, *Rhodomonas* sp. (NT15) and *Cryptomonas* sp. (CRFI01) and an unidentified prymnesiophyte (NT19) and in a commercially available *Isochrysis* sp. used as control. Results showed that the diatom *Chaetoceros* had the highest percentage of lipid (16.8 % dry weight) when cells were cultured at 25°C. However, *Rhodomonas* sp., *Cryptomonas* sp., NT19, and *Isochrysis* sp. had significantly higher amounts of lipids at 15.5, 12.7, 21.4, and 21.7 % dry weight, respectively, when grown at temperatures within the range of 27-30°C. The effect of temperature on fatty acid composition of *Chaetoceros* is summarized in Table 1.2.

1.5.3. Carbon Dioxide and Organic Carbon Sources. Algae use carbon dioxide (CO₂) as the source of carbon to synthesize organic compounds such as lipids. Since carbon demand and productivity increase proportionally, intensive mass culture of microalgae requires a vast source of CO₂ (Neenan et al., 1986). Carbon dioxide is present in air at a very small concentration (0.037 % by volume in dry air). On the marine diatom *Chaetoceros wighamii*, addition of CO₂ increased the protein content, lowered the concentration of carbohydrates, but had no significant

Table 1.2. Fatty acid composition of laboratory batch cultures of *Chaetoceros* sp. (CS256) grown at different temperatures (Renaud et al., 2002).

	Temperature (°C)				
	25	27	30	33	35
Saturated fatty acids					
12:0	– ¹	–	–	–	–
14:0	23.6 ^{m,2}	22.3 ^m	22.3 ^m	23.8 ^m	28.3 ⁿ
16:0	9.2	8.6	8.9	8.7	8.4
17:0	–	–	–	–	–
18:0	0.7	0.7	0.9	1.3	0.9
Sum%	33.5	31.6	32.1	33.8	38.2
Monounsaturated					
16:1n-7	36.5 ^m	36.4 ^m	39.1 ^m	36.9 ^m	33.5 ⁿ
18:1n-9	1.7	1.4	1.4	1.9	1.8
18:1n-7	1.2	1.2	1.3	1.1	1.2
Sum%	39.4	39.3	42.5	40.0	36.5
Polyunsaturated					
16:2n-7	0.9	1.2	1.4	1.4	1.8
16:3n-4	2.6	4.3	3.9	4.2	4.8
16:4n-1	0.5	0.7	0.7	0.7	0.7
18:2n-6	0.4	1.2	1.1	1.3	1.2
18:3n-6	0.9	0.6	0.8	0.8	1.0
18:3n-3	0.5	0.6	0.4	0.5	1.0
18:4n-3	0.6	0.4	0.2	0.5	0.3
18:5n-3	–	–	–	–	–
20:4n-6	4.1 ^m	4.5 ^m	4.0 ⁿ	3.6 ⁿ	2.7 ⁿ
20:5n-3	8.0 ^m	6.7 ⁿ	6.5 ⁿ	6.9 ⁿ	6.6 ⁿ
22:5n-3	–	–	–	–	–
22:6n-3	1.0	0.6	0.4	0.4	0.3
Sum%	19.5	20.8	19.0	19.8	20.4

Data as mean percentage of total fatty acids; n=3 for flask cultures.

1 (–) Indicates that fatty acid was less than 0.1% of total fatty acids.

2 Different superscripts across a row (m to n) indicate significant difference between means (ANOVA, Tukey's test; P<0.05).

effect on the lipid content (Castro and Tavano, 2005). As for microalgae grown on the heterotrophic mode, experiments using diverse sources such as molasses (containing 25 %

glucose, 25 % fructose and 30 % sucrose on average (Becker, 1994a), acetic acid, and hydrocarbons (such as *n*-heptadecane) have been performed and shown to work at different concentrations (Becker, 1994a).

1.5.4. Nutrients. Nutrients are inorganic or organic compounds other than carbon dioxide and water, used for growth whose presence in the cell is necessary for cellular function (Neenan et al., 1986). Some algae require specific organic compounds synthesized by other organisms. However, many algae require only inorganic nutrients, and it is likely that these algae could be used as feedstock for biomass fuel production. Limiting nutrients to algae are nitrogen, phosphorus, silica (for diatoms) and iron (Neenan et al., 1986).

1.5.4.1. Nitrogen. After carbon, nitrogen (N) is the most important nutrient contributing to the production of biomass. The nitrogen content in the microalgal biomass can range from 1 % to more than 10 % and depending on its supply and availability, it can vary between different groups (e.g. low in diatoms) and within particular species (Grobbelaar, 2004). Discoloration of the cells is a frequent response to nitrogen limitation due to a decrease in chlorophyll content and an increase in carotenoids, as well as the accumulation of organic carbon compounds such as polysaccharides and certain oils like polyunsaturated fatty acids (PUFAs) (Becker, 1994b). Nitrogen is mostly supplied as nitrate, but often ammonia and urea are used, both displaying similar growth rates (Kaplan et al., 1986). Many microorganisms tend to prefer ammonia as a nitrogen source, and the assimilation of either nitrate or ammonia is said to be related to the pH of the growth media (Hu, 2004). A drop on pH can be observed when ammonia is used as the only source of N, especially during active growth owing to the release of H⁺ ions (Grobbelaar, 2004). An increase in pH is observed when nitrate is used as the only N source (Grobbelaar, 2004). Ammonia loss due to volatilization is an important factor to be considered when deciding

whether to supply either one. Urea can be considered another attractive source of nitrogen. About 42 % of the weight of urea is nitrogen; therefore, more material than that of ammonia has to be handled. Lin and Lin (2011) reported that the microalgae fed with a combination of urea and sodium nitrate had the highest ash-free dry biomass content with a yield of $4.15 \pm 0.38 \text{ g L}^{-1}$. The urea molecule contains a carbon atom as well as two nitrogen atoms. This carbon atom is released as CO_2 when urea is utilized and presumably, it is available for photosynthetic assimilation (Neenan et al., 1986), therefore, urea has the potential of providing both the nitrogen and 1.5 to 10 % of the carbon requirement (Neenan et al., 1986).

1.5.4.2. Phosphorus. Phosphorus is essential for growth in many cellular processes such as energy transfer and during the biosynthesis of nucleic acids. Orthophosphate is the preferred form in which it is provided to algae and its uptake is said to be energy dependant (Grobbelaar, 2004). In spite of the fact that algal biomass contains less than 1 % phosphorous (P), it usually becomes one of the most important growth limiting factors in algal culture (Neenan, et al., 1986). This happens because P binds easily to other ions (e.g. carbonate and iron) resulting in its precipitation. The insolubility of the resulting phosphate makes this essential nutrient unavailable for algal uptake (Grobbelaar, 2004). The supply of P also influences the composition of the produced biomass. Similar effects to the ones obtained in algae grown under nitrogen starvation, such as the tendency to accumulate large amount of lipids, with a decreased amount of proteins, chlorophyll and nucleic acids content have been reported on phosphate deficient cultures (Becker, 1994a).

1.5.4.3. Other Macro, Micronutrients and Chelates. Sulfur, potassium, sodium, iron, magnesium, calcium and trace elements like boron, copper, manganese, zinc, molybdenum, cobalt, vanadium, and selenium are also important in algal nutrition (Grobbelaar, 2004). Silicon, present in the cell

walls of many algal groups, is an important component in diatoms where it constitutes an essential nutrient for their growth and production (Healy, 1973). Silicon limitation, which is prone to happen, can also lead to the accumulation of secondary metabolites, such as lipids (Grobbelaar, 2004). Experiments conducted with silicon-deficient cells of *Cyclotella cryptic* (a diatom species that accumulates lipids under non-growing conditions) indicated that lipid accumulation occurs as a function of both increased partitioning of newly photo-assimilated carbon into lipids and slow conversion of non-lipid compounds (Roessler, 1987).

1.5.5. Salinity. Protein, lipids and carbohydrates seem slightly affected by a wide range of salinity for most microalgae species. However, in some species, increases in ash and lipid content were observed at higher salinity. Studies by Raghavan et al. (2008) indicated that for *Chaetoceros calcitrans*, in terms of growth and chemical composition, a salinity of 25 mM NaCl was optimum. This study also showed that at a salinity of 35 mM NaCl, carbohydrates were increased while lipids and protein decreased (Raghavan et al., 2008). Vasquez-Duhalt and Arredondo-Vega (1991) tested two strains of *Botryococcus Braunii* (Austin and Gottingen) under different NaCl concentrations. The protein content in the cell biomass of the Austin strain showed a decrease with increasing salinity. The effect was small in the case of the Austin and more pronounced in the Gottingen strain. Carbohydrate content of Austin strain cells was not affected by media NaCl concentrations up to 0.2 M. Higher concentrations of NaCl induced an increase in the carbohydrate content of cells. However, large standard deviations were reported (Vasquez-Duhalt and Arredondo, 1991). The lipid content in both strains was not affected by the salt concentration. This suggests that saline stress does not induce lipid accumulation in these organisms (Vasquez-Duhalt and Arredondo-Vega, 1991). Mishra and Jha (2009) found similar results on *Dunaliella* cultures harvested 20 days after inoculation and grown at different salt

concentrations. Salt induces osmotic pressure in the broth which causes the microorganisms to protect themselves from it by producing exopolysaccharides (EPSs) (Abbasi and Amiri, 2008). EPSs of 944 mg l^{-1} were obtained from *Dunaliella* grown in 5M NaCl relative to the 56 mg l^{-1} observed in media containing 0.5 M NaCl (Mishra and Jha, 2009).

Salinity can have an effect on growth rate along with other factors like temperature. Castro and Tavano (2005) observed that temperature had a significant effect on the growth of *Chaetoceros wighamii* under salinity of 25 mM, but not at 35 mM. At 30°C, the growth rate was lower at 25 mM when no CO₂ had been added. Rao et al. (2006) identified a marginal increase in carbohydrate content at 17 mM and 34 mM salinity cultures with *Botryococcus braunii*. The total fat content of alga grown at different salinity varied from 24-28 % (w/w) compared to only 20 % (w/w) for control without salts. The fatty acid profile indicated the presence of C16:0, C16:1, C18:0, C18:1, C18:2, C22:0, C22:1 and C24:0 fatty acids. Stearic and linoleic acids were higher in proportion in the control culture while palmitoleic and oleic acids were the major fatty acids for cultures grown in 34mM and 85mM salinity (Rao et al., 2006).

1.6. Microalgae Identification

1.6.1. Polymerase Chain Reaction. The polymerase chain reaction (PCR) is a powerful and sensitive technique which amplifies specific DNA sequences exponentially through a three-step process done in multiple cycles (Saiki et al., 1985). First, the double-stranded DNA template is denatured at a high temperature. Then, sequence-specific primers are annealed to the target sequence followed by the addition of a thermostable DNA polymerase, such as *Taq* DNA polymerase (Chien et al., 1976; Kaledin et al., 1980; Lawyer et al., 1993; Longley et al., 1990; Lyamichev et al., 1993). This enzyme is responsible for extending the annealed primers, and doubling the amount of the original DNA sequence. The new product then becomes an additional template for subsequent cycles of amplification. These three steps are usually repeated in cycles

for 20 to 30 times, resulting in an increase of target DNA concentration of 10^5 to 10^9 times the original amount.

The use of PCR to obtain large amounts of a desired product can have both positive and negative aspects. If amplification is not successful, this can lead to the generation of many undesired products leading even to the exclusion of the target product (Roux, 1995). The opposite would be that no product may be amplified. In regards to optimization, several variables have been recognized to contribute to this effect (Saiki et al., 1988). Main optimization variables include magnesium (Mg^{2+}) concentrations, buffer pH, and cycling conditions. Within cycling conditions, the annealing temperature is of utmost importance. The interdependence between variables adds difficulty to the situation. For instance, increasing the amount of deoxynucleotide triphosphates (dNTPs) lowers the concentration of free Mg^{2+} available to exert an effect on polymerase function because dNTPs directly chelate a proportional number of Mg^{2+} ions (Roux, 1995).

1.6.2. Enhancing Agents. Several additives and enhancing agents such as dimethyl sulfoxide (DMSO), N,N,N,-trimethylglycine (betaine), formamide, glycerol, non-ionic detergents, bovine serum albumin, polyethylene glycol and tetramethyl ammonium chloride can be included in PCR reactions with the aim of increasing yield, specificity and consistency (Frackman et al., 1998).

Application of the Taguchi method (Taguchi, 1986), which focuses only on the main effects and two factor interactions, can eliminate the need of a full multivariate matrix analysis for each of the variables tested, which can become a burdensome and costly task. With this method, the size of the matrix can be trimmed down significantly and several key variables can be altered simultaneously (Cobb and Clarkson, 1994).

1.6.3. Magnesium Concentration. Magnesium chloride serves as an essential co-factor for the DNA polymerase and optimization for every primer/template pair should always be attained (Kolmodin and Felton, 1997). Magnesium ion is closely bound to the phosphate-sugar backbone in nucleotides and nucleic acids; therefore, different amounts can have strong and complex effects on experiments on which nucleic acids take part (Blanchard et al., 1993). Because of the needed role of free magnesium as enzyme co-factor in PCR, its total ion concentration has to exceed the total dNTP concentration. Generally, magnesium ion's concentration varies in series of 1.5 - 4 mM with 0.5-mM step increments (Kolmodin and Felton, 1997).

1.6.4. Amplification of 16S rRNA, 18S rRNA and Other Genes. 16s rRNA is the most commonly used genetic marker for the study of bacterial phylogeny and taxonomy (Janda and Abbott, 2007). Reasons for this include: (1) its presence in almost all bacteria as operons, (2) no change in 16s rRNA gene's function has been observed over time and, (3), the 16s rRNA gene's length (1,500 bp) is suitable for informatics purposes (Janda and Abbott, 2007). Erwin and Thacker (2008) characterized partial 16S rRNA and the entire 16S- 23S internal transcribed spacer (ITS) sequences from *Synechococcus spongiarum* in an attempt to assess the phylogenetic utility of rRNA sequence data in resolving the phylogeny of sponge-associated bacteria. Their results showed that 16S rRNA sequences were highly conserved, exhibiting less than 1 % sequence divergence among symbiont clades; whereas, ITS gene sequences displayed a much higher variability than 16S rRNA sequences.

In addition to molecular techniques based on PCR amplification targeting conserved regions inside the 16S rRNA gene (Komarek , 2006), other genes such as *hetR* (heterocyst differentiation control protein) (Han et al., 2009; Schleifer, 2009) and *nifH*, which encodes for nitrogenase reductase (Zehr and McReynolds, 1989; Janson et al., 1998), have also demonstrated

to have potential for phylogenetic discrimination. However, considering that several disagreements between traditional morphological classification and phylogenetic analysis still remain, the utilization of chemotaxonomic markers, such as lipids and their fatty acids, have been considered as complementary approaches since they provide data for taxonomic position assignment as well as some correlations with morphological properties of cyanobacteria (Galhano et al., 2011). In their study, Galhano et al. (2011) obtained molecular information regarding the 16S rRNA gene as well as the *hetR* and *nifH* genes from filamentous cyanobacterial strains isolated from two Portuguese freshwater ecosystems, a eutrophic shallow lake (genus *Aphanizomenon*) and rice fields (genera *Anabaena* and *Nostoc*). Features such as genotypic, morphological and biochemical attributes (fatty acid methyl ester profiles) were used for strain characterization. Their results showed that the unknown *Aphanizomenon* had 99 % similarity with 16S rDNA sequence with *Aphanizomenon gracile* 219 (isolated from a Danish lake), the unknown *Anabaena* had 99 % with *Anabaena flosaquae* PCC 7905 (type strain) and 99 % similarity with *Anabaena cylindrica* NIES19. The unknown *Nostoc* showed 98 % 16S rDNA sequence similarity with *Nostoc* sp. 8938 and *Nostoc muscorum* I. The *hetR* gene fragments with approximately 450 bp showed 97 % similarity with *Aphanizomenon* sp. TR183 and 98 % similarity with *Anabaena flos-aquae* SAG 30.87; whereas, a 100 % similarity was seen with *Nostoc* PCC 7906. The sequence of the *nifH* gene fragment (319 bp) of the unknown *Aphanizomenon* had high similarity (98 %) with a nitrogenase reductase sequence of an uncultured cyanobacterium clone Gt1463 and a different strain's *nifH* partial sequence showed high similarity (99 %) with *Nostoc muscorum* CC1090A1; whereas, the *Anabaena*'s strain's *nifH* partial sequence had similarities of 99 and 100 % with the partial nitrogenase reductase gene

sequences of a freshwater strain of *Anabaena oscillarioides* and *Anabaena cylindrica* UTEX B629, respectively (Galhano et al., 2001).

Considered one of the most important molecular markers, 18S rRNA is a gene used in several applications among which molecular phylogenetic analyses and biodiversity screening can be included (Meyer et al., 2010). Buchheim et al., 1990, 1996, 1997, have conducted phylogenetic studies using 18S rRNA gene which have demonstrated significant diversity within the green algal genus *Chlamydomonas*. Armbrust et al. (2001) used 18S primers to amplify the 18S rRNA genes from three closely related centric diatoms. Degenerate PCR primers were designed and used to amplify a portion of *Sig1* (sexually induced gene 1), a gene strongly upregulated during sexual reproduction in the centric diatom *Thalassiosira weissflogii*. This gene has been hypothesized to encode a protein involved in the gamete recognition in three closely related species of the cosmopolitan genus *Thalassiosira* sp., *oceanica*, *guillardii*, and *pseudonana*. In their study, identification of *Sig1* facilitated the development of this gene as a molecular marker for diatom sexual events and how these events affect diatom speciation and population dynamics.

Kaczmarek et al. (2005) amplified the small subunit rDNA (SSU rDNA) gene from the domoic acid-producing diatom *Pseudo-nitzschia multiseriata* by PCR using the primer pair 27F and 1492R. Isolation and identification of bacteria associated with the *P. multiseriata* clones and sub-clones to the closest characterized bacterium was done using the Ribosomal Database Project II. Their results showed that five of the eight bacterial strains belonged to the Alphaproteobacteria; whereas, the remaining bacteria were related to the Gammaproteobacteria and the bacteroides. Additionally, BLAST analysis of some of the above mentioned bacterial strains showed that they shared their highest SSU rRNA gene sequence identities (99.1, 99.0,

93.9 and 99.0 %, respectively) with the bacteria affiliated to *Stappia* (AY258082), *Sulfitobacter* (AY258079 and AJ534214), *Flexibacter* (AY258133) and *Sphingomonas* (AY258097). BLAST (Basic Local Alignment Search Tool) is a bioinformatic tool that helps identify members of gene families by finding regions of local similarity between sequences and by comparing nucleotide or protein sequences to sequence databases. Statistical significances are calculated for those matches ([http:// www.ncbi.nlm.nih.gov/blast/](http://www.ncbi.nlm.nih.gov/blast/)). A method described by Rasoul-Amini et al. (2009) used DNA extraction and PCR amplification of the 18S rRNA gene to obtain sequences on 12 microalgae species belonging to the class Chlorophyceae. Bioinformatic tools (BLAST) were also used. The results of PCR products BLASTed with other sequenced microalgae in NCBI showed similarity to the 18S small subunit rRNA of other microalgae at 99-100 %.

1.7. Microalgae Lipid Content

Lipids are extractable to non-polar solvents (e.g. ether, chloroform, alkanes). This definition includes not only the triglycerides (triesters of fatty acids and glycerol), but compound lipids (e.g. phospholipids, glycolipids), steroids, chlorophylls, carotenoids, and hydrocarbons as well (Benemann and Weissman, 1984). The main phospholipids in algae are phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, phosphatidylserine, phosphatidylglycerol, phosphatidic acid, and diphosphatidyl glycerol (Pohl, 1982). The major algal glycolipids are monogalactosyldiglyceride, digalactosyldiglyceride, and sulphoquinovosyldiglyceride (Pohl, 1982). A novel class of algal lipids are chlorosulpholipids which are derivatives of N-docosane-1,14-diol and of N-tetracosane-1,15-diol disulphates found in Chrysophyceae, Xanthophyceae, Chlorophyceae, and Cyanophyceae (Pohl, 1982).

Several reports exist on lipid production by microalgae, but the majority of this data refers only to fatty acids, most of which are C14 to C20 (Benemann and Weissman, 1984). Most of the fatty acids of algae are bound to the above lipids. In some algae, however, minor quantities of free

fatty acids can be found. Larger amounts of even-numbered saturated fatty acids (12:0, 14:0, 16:0,18:0) are primarily produced by algae, with only minor quantities of odd numbered fatty acids (13:0 to 19:0) [(Pohl, 1982)]. Small amounts of branched chain fatty acids have also been reported (Pohl, 1982). Fatty acids with C12 to C22 carbon chains and 1 to 6 double bonds are predominant. These double bonds are usually in the cis- configuration being an exception trans-3-(16:1) which is found in the phosphatidylglycerol of photosynthesizing algae. Different from bacteria and fungi, it seems apparent that algae do not synthesize fatty acids with unusual structures, such as acetylenic, hydroxyl, epoxy, oxo, cyclopropanoic, and cyclopropenoic acids (Pohl, 1982). The average lipid content in algae can range between 1 and 40 % dry weight (Becker, 1994b). However, numbers as high as 85 % dry weight can be achieved under certain conditions (Becker, 1994b). The lipid content of various species of microalgae is presented in Table 1.3.

1.8. Goal of This Study

The goal of this research was to compare different media and growth conditions for the isolation and identification of microalgal strains that are native to both Louisiana’s fresh and brackish water bodies for their potential use in the alternative fuel industry. Selected strains were identified through genomic DNA in sequencing of the 16S rRNA and 18SrRNA genes. Lipid and fatty acid content were characterized and quantified post genomic identification.

Table 1.3. Lipid content of some microalgae (Gouveia and Oliveira, 2009). Originally adapted from Becker (1994b); Illman et al. (2000); Liu et al. (2007); Miao and Wu. (2006); Natrah et al. (2008); Spolaore et al. (2006); Tornabene et al. (1983); and Xiong et al. (2008).

Species	Lipids (% dry matter)
<i>Scenedesmus obliquus</i>	11–22/35–55

(Table 1.3, continued)

<i>Scenedesmus dimorphus</i>	6-7/16-40
<i>Chlorella vulgaris</i>	14-40/56
<i>Chlorella emersonii</i>	63
<i>Chlorella protothecoides</i>	23/55
<i>Chlorella sorokiana</i>	22
<i>Chlorella minutissima</i>	57
<i>Dunaliella bioculata</i>	8
<i>Dunaliella salina</i>	14-20
<i>Neochloris oleoabundans</i>	35-65
<i>Spirulina maxima</i>	4-9

2. MATERIALS AND METHODS

2.1. Sampling and Collection of Organisms

Water samples were collected from sources at different locations (gulf, bays, lakes, bayous, ditches) within Louisiana's Southeast region with the aim of isolating and identifying algae as potential feedstock for the production of biodiesel. On April 17, 2010, water samples were taken from Dulotte Canal (Empire, LA), Bay Adams (Empire, LA), Myrtle Grove Marina (Port Sulphur, LA) and on May 18, 2010 from Magnolia ditch and End of Magnolia (Raceland, LA), Cocodrie Marina (Chauvin, LA), and Chacahoula ditch, Lake Palourde, Brownell Memorial's lake, and Adam's landing (Morgan City, LA) (Figure 2.1). Samples were stored at 4°C. Salinities of water samples were analyzed by total conductivity and are depicted in Table 2.1.



Figure 2.1. Map of Sampling Locations. (Adapted from www.maps.com).

2.2. Growth Conditions

Media for the isolation of algal strains were prepared by mixing filtered (55 mm, Whatman filters, Kent, United Kingdom) and sterilized natural brackish or freshwater (same source from where samples were taken) with either 50X Guillard's (F/2) (APPENDIX 1) marine

water enrichment solution (Sigma-Aldrich, St. Louis, MO) for brackish water or Chu's Media for freshwater samples. The Chu's Media contained 40 mg Ca (NO₃), 0.8 mg FeCl₃, 100 mg

Table 2.1. Salinities of sampling locations through total conductivity.

Location	Salinity ^{**} (mg/L)
Dulotte Canal	3,727
Bay Adams	4,304
Myrtle Grove	1,064
Magnolia Ditch	58
End of Magnolia	51
Cocodrie Marina	3,317
Chacahoula	53
Lake Palourde	108
Brownell Memorial	65
Adam's Landing	93

^{**} The sum of chromatographic sodium, potassium, magnesium and calcium.

K₂HPO₄, 250 mg MgSO₄.7H₂O, 20 mg Na₂CO₂, 25 mg NaSiO₃. 9 H₂O per liter. Petri plates of each medium were prepared containing 1.5 % w/w agar (Difco Bacterius Limited, Houston, Texas). Supplemental Na₂ SiO₃. 9 H₂O w/v at 1 % using a stock solution containing 30 g L⁻¹ was added to F/2 enriched media at pH 7 to enhance the isolation of diatoms (Perez, 2006).

2.3. Colony Selection

One ml of each water sample (brackish or freshwater) was inoculated on both plates and into flasks containing 25 ml broth of each medium. Inoculated plates and flasks were placed in a GyromaxTM 777 incubator (Amerex Instruments Inc., Lafayette, CA) under continuous fluorescent lighting at 23 μmol photons m⁻² s⁻¹ measured using an LI-193 spherical quantum sensor along with a LI-1400 data logger (LI-COR, Lincoln, NE) at 30°C ± 2 and slow swirling (flasks only) at 80 RPM for 2 to 4 weeks.

2.4. Colony Homogenization

Colonies were selected based on color differences and transferred to fresh agar plates. In the case of samples grown in liquid media, a 1 ml aliquot was transferred to agar plates and incubated under similar conditions for an additional 2 to 3 weeks. Homogenization and purification of isolated colonies were carried out through the inoculation of colonies on both Blue Green Media (BG-11) (APPENDIX 2) which favors blue greens and diatoms, and Bold's Basal Media (BBM) (APPENDIX 3) which favors green algae. Microscopic observations at 100X magnification with oil immersion (DC3-163 Microscope, National Optical and Scientific Instruments Inc., San Antonio, Texas) and serial dilution were made until a unialgal culture was obtained.

2.5. Microorganism Selection and Harvesting

Out of 17 isolates, four algae were selected for identification. Their selection was based on rapid growth, morphological and class diversity, and culture homogeneity. Based on morphological differences observed through microscopic analysis, they were tentatively identified as a cyanobacteria (N3) isolated from Cocodrie Marina, a diatom (FC2) isolated from a Chacahoula ditch and two green algae isolated from a Chacahoula ditch (FC1) and Dulotte Canal (D2), respectively (Figure 2.2). *Chlorella vulgaris* UTEX 259 (University of Texas Culture Collection of Algae, Austin, TX), provided by Dr. Misook Kim from the Audubon Sugar Institute (St. Gabriel, LA), was used as control during the identification process. N3 and FC2 isolates were grown in 1 L flasks containing 800 ml of BG-11; whereas, FC1, D2, and *Chlorella vulgaris* were grown in BBM. Flasks were placed under continuous fluorescent lighting $20 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ at 30°C with slow shaking at 80 RPM in a shaker incubator Gyromax 777 and harvested after 15 days by centrifugation (Sorvall RC-5 Superspeed Refrigerated Centrifuge, Dupont, England) at either 1,200 x g for N3 or 6,500 x g for FC2, FC1, D2 and *Chlorella*

vulgaris UTEX 259 at 4°C. Post harvest, 100 mg of wet cell biomass were washed with 100 mM Tris buffer (pH 7.5) and stored at -20°C.




Figure 2.2. Tentatively Identified Microorganisms. FC2: diatom, N3: cyanobacteria, FC1 and D2: green algae.

2.6. Genomic DNA Extraction

Frozen cells were thawed and genomic DNA extraction was performed using a Fast DNA Kit (MP Biomedicals, Solon, OH). Briefly, 1 ml of CLS-Y was added to each tube containing 100 mg of wet cells and then transferred to a new set of tubes each containing a ball and garnet. Cells were processed using a FastPrep®-24 Instrument (MP Biomedicals) for 10 s at a speed setting of 5 m s^{-1} , placed on ice for 1 h and then centrifuged for 15 min at 4°C at $15,700 \times g$ (Hermle Z233 MK-2 High Speed Refrigerated Centrifuge, Hermle Labortechnik, Wehingen, Germany). After centrifuging, 600 μl of supernatant fluid was placed in a fresh tube and to that, 600 μl of binding resin (guanidine thiocyanate at 60-70%, water at 10-20 % and silica at 10-20 %) was added. The mixture was thoroughly mixed and incubated for 5 min at 25°C. Tubes were centrifuged for 1 min at $15,700 \times g$ (Centrifuge 5415 D- Eppendorf, Hamburg, Germany) and after supernatant removal, 500 μl of SEWS solution was added to each tube, mixed, and centrifuged for 1 min, and then supernatants were discarded. To re-wash, another 500 μl of SEWS was added, mixed, centrifuged, supernatants removed again, and tubes were centrifuged for 10 s to remove any residual supernatants. To elute DNA from the resin, 100 μl of DES

solution was added and incubated for 3 min at 25°C. After incubation, DNA was collected by pipetting, centrifuged for 1 min at 15,700 x g, and supernatants were collected; an additional centrifugation was carried out to make certain no resin was present. Studies were conducted in duplicate.

2.7. DNA Purification

Ten μ l of 3M sodium acetate solution (pH 5.2) and 275 μ l of 95 % v/v ethanol were added to each sample's DNA, mixed by flipping and placed in dry ice for 15 min. The samples were then centrifuged for 15 min at 15,700 x g at 4°C (Hermle Z233 MK-2 High Speed Refrigerated Centrifuge) to remove the supernatant. The pellets were each washed once with 1 ml of 95% v/v cold ethanol and centrifuged for 5-6 mins. After supernatants were discarded, DNA pellets were placed in an oven (Isotemp Vacuum Oven Model 280A, Fisher Scientific, Pittsburgh, PA) at 37°C for 10 min to dry under a vacuum of -25 inches of Hg. After drying, 80 μ l of TE buffer was added and the DNA concentration was determined with a ND-1000 Spectrophotometer (Nanodrop, Wilmington, DE). Two readings were taken per sample (2 μ l).

2.8. Standard Polymerase Chain Reaction (PCR) and Optimization

Nine different conditions were tested using a set of primers (Integrated DNA Technologies, Coralville, IA) for either a 16S rRNA gene (cyanobacteria), since this gene is better suited for prokaryotes, or an 18S rRNA gene (diatom and greens) which is a universal gene commonly used for the identification of eukaryotes (Table 2.2).

Three PCR master mixes were prepared using a Taq PCR Kit (New England Biolabs, Ipswich, MA). The components of each PCR master mix are indicated in Table 2.3. Dimethyl sulfoxide (DMSO) (Sigma-Aldrich, St. Louis, MO) and glycerol (Sigma-Aldrich, St. Louis, MO) were added to the second and third master mixes, respectively. Each mix was divided into three tubes of 80 μ l and combined with 20 μ l of three MgCl₂ concentrations (15 mM, 30 mM and 45

mM) , resulting in a PCR reaction mixture with a final volume of 100 μ l. The PCR reactions were performed on a thermocycler (PTC-100, MJ Research, Inc., Waltham, MA) using a program designed for the amplification of either 16S rRNA or 18S rRNA genes (Tables 2.4 and 2.5).

Table 2.2. 16S rRNA and 18S rRNA primers used for PCR reactions.

Primer	Sequence
16S rRNA	
16S bottom ^a	5'- AGAGTTTGATCMTGGCTC-3'
16S top ^a	5'-ACGGGCGGTGTG-3'
16S bottom 2 ^b	5'-TTGGGCGTAAAGCGT-3'
18S rRNA	
18S-C ^a	5'-TGATCCTTCYGCAGGTTCAC-3'
18S-D ^a	5'-ACCTGGTTGATCCTGCCAG-3'
18S C-2 ^b	5'- ATTGGAGGGCAAGTCTGGT-3'
18S D-2 ^b	5'- ACTAAGAACGGCCATGCAC-3'

a = Used for PCR amplification; b = Used for sequencing

2.9. Gel Electrophoresis

DNA quality and PCR products were determined using 0.8 % agarose cast gels (Genepure LE agarose, ISE Bioexpress, Kaysville, UT) on Tris-acetate-EDTA (TAE) buffer. Ethidium bromide was added to the gel mix to a final concentration of 0.2 μ g/ml, and samples were mixed with 6X loading buffer (0.25 % w/v bromophenol blue and 40 % w/v sucrose). Ten μ l of 2 log DNA ladder (0.1-10 kb, New England Biolabs) were loaded onto the agarose gel and the electrophoresis was run at 105 watts for 45 min (BRL-Model 250, Life Technologies, Carlsbad, CA). Bands were visualized with a UV lamp at a wavelength of 312 nm (UV Fotoprep I- Fotodyne Kodak, Rochester, NY).

Table 2.3. Reaction volumes and concentrations for PCR master mixes.

Component	Master Mixes*			Final Concentration
	Volume			
	1	2	3	
	μl	μl	μl	
10X Mg-free buffer	40	40	40	1X
Primer 1 (50 μM)	4	4	4	0.5 μM
Primer 2 (50 μM)	4	4	4	0.5 μM
Template (sample)	X	X	X	0.25 ng/μl
dNTPs (10 mM)	8	8	8	0.2 mM
Taq DNA (5U/μl)	2	2	2	0.025 U/μl
DMSO (neat)	0	20	0	5%
Glycerol (50%)	0	0	80	10%
Nuclease Free Water (to a final volume of 320 μl)	Y	Y	Y	

X= Amount of template required to reach final concentration.

Y= Amount of Nuclease free water up to 320 μl.

*Each master mix was divided into 3 tubes of 80 μl. Next, 20μl of a MgCl₂ stock solution (Low (L)=15 mM, Medium (M)=30 mM, and High (H)=45 mM) was added to get the following combinations: 1L, 1M 1H, 2L, 2M, 2H, 3L, 3M, and 3H . Final concentrations of MgCl₂ were 1.5, 3, and 4.5 mM for L, M, and H, respectively.

Table 2.4. PCR cycle conditions for 16S rRNA.

Initial Denaturation	95°C	30 s
30 cycles	95°C	30 s
	61°C	1 min
	68°C	1 min
Final Extension	68°C	5 min
Hold	4°C	∞

Table 2.5. PCR cycle conditions for 18S rRNA.

Initial Denaturation	95°C	30 s
30 cycles	95°C	30 s
	67°C	1 min
	72°C	1 min
Final Extension	72°C	5 min
Hold	4°C	∞

2.10. Gel Extraction and Purification

Electrophoresis was conducted as previously described. DNA bands were visualized using a UV lamp and the gel bands excised using a clean glass cover slip and then placed in a 1.5 ml microfuge tube. DNA was extracted using a QIAquick Gel Extraction Kit (Qiagen, Valencia, CA). Briefly, 1 ml of buffer QG was added to each microfuge tube and mixed by vortexing (VortexGenie 2-VWR, Radnor, PA) for 30 s. The tubes were then incubated on a heating block (ISOTEMP 125D, Fisher Scientific, Pittsburg, PA) at 50°C for 10 min with vortexing for 2-3 s every 2 min. The contents of each tube were transferred to a QIAquick spin column (QIAquick Gel Extraction Kit) containing a silica membrane with a 10 kb cut-off and centrifuged at 15,700 x g for 1 min. The DNA was bound to the membrane in the spin column. Buffer PE (750 µl) was used to wash each column followed by 1 min centrifugation. After the removal of supernatant columns were centrifuged once more to remove residual buffer PE, then they were placed into a new set of tubes and 25 µl of buffer EB was added to elute the DNA. After 1 min, tubes were centrifuged for 1 min and the DNA concentration was measured with a ND-1000 spectrophotometer and stored at 4°C.

2.11. Cloning of PCR Products

2.11.1. LB Agar Plates and Broth Media Preparation. Preceding the cloning reaction and transformation with chemically competent *Escherichia coli* (*E. Coli*) cells, plates containing Luria- Bertani (LB) (10g Tryptone, 5g Yeast extract, and 10g sodium chloride per liter) [(Fisher Scientific, Pittsburgh, PA)], with 1.5 % agar, and broth were prepared. Kanamycin to a final concentration of 50 µg/ml was added for the selection of transformant colonies.

2.11.2. Colony Screening. Transformant colonies which were grown at 37°C for 18 h were determined by blue/white screening by evenly distributing 50 µl of 2 % X-gal to each LB agar

plate containing kanamycin. Mostly white cells were observed which was indicative of positive cell transformation.

2.11.3. Cloning Reaction and Transformation with Chemically Competent *E. coli*. The TOPO TA Cloning Kit for sequencing with the plasmid vector PCR[®] 4-TOPO[®] and one shot[®]-TOP 10 chemically competent cells (Invitrogen, Carlsbad, CA) was used. Four μl of gel purified DNA from each sample was mixed with 1 μl of salt solution (containing 1.2 M NaCl and 0.06 M MgCl_2) and 1 μl of TOPO vector (6 μl total reaction volume). After gentle mixing by tapping, the mixes were left to stand for 30 min at 25°C. Vials containing the chemically competent *E. coli* kept at -80°C were thawed on ice (~ 0°C) for 5 mins. After the 30 min incubation, 2 μl of the cloning reaction mix was transferred to the vial containing the *E. coli* competent cells, gently mixed by tapping 6-8 times and incubated on ice (~ 0°C) for 30 min. The vials were placed in a water bath (C76 Water Bath Shaker, New Brunswick Scientific, Edison, NJ) at 42°C for 30 s to produce a heat shock, and then immediately placed on ice followed by the addition of 250 μl of SOC medium. Vials were placed in a shaker incubator at 37°C for 1 h at 200 RPM. The contents of each vial were split and spread onto two LB agar plates containing kanamycin and x-gal, adding 80 % of the content to one plate and 20 % to the other one, and incubated overnight at 37°C. After 18 h of incubation, three white colonies were selected from each sample. Each selected colony was grown overnight in 3 ml of LB broth containing 50 $\mu\text{g/ml}$ kanamycin at 37°C at 250 RPM.

2.12. Purification of Plasmid DNA

The QIAprep Spin Miniprep Kit (Qiagen, Valencia, CA) was used for the purification step. *E. coli* cells grown overnight were centrifuged for 3 min at 9,300 x g and the supernatant decanted. The pelleted bacterial cells were resuspended in 250 μl of buffer P1 (containing lyse-

blue reagent which is used as a visual identification system to prevent handling errors), thoroughly mixed followed by the addition of 250 µl of buffer P2. Samples were mixed gently with lysis time not exceeding 5 min. Buffer N3 (350 µl) was added last and samples were centrifuged for 10 min at 15,700 x g. Centrifugates containing the plasmid DNA were then transferred to a column containing a 50 kb cut-off silica membrane and centrifuged for 1 min. Buffer PB (0.5 ml) was used to wash the columns and then centrifuged for 1 min. All flow-through material was discarded. This step was repeated twice to remove all supernatant fluid. The DNA was eluted by transferring the columns to new 1.5 ml tubes and adding 50 µl of EB buffer. Columns were let stand for 1 min followed by centrifugation at 15,700 x g for 1 min. DNA concentration was determined as previously described.

2.13. Restriction Enzyme Digestion

EcoRI restriction enzyme and 10X EcoRI Buffer (New England Biolabs, Ipswich, MA) were used. Sample DNA up to a maximum amount of 800 ng was mixed with nuclease-free water up to a volume of 17.5 µl, followed by the addition of 2 µl of 10X Eco RI buffer and 0.5 µl of restriction enzyme in a 1.5 ml centrifuge tube. The contents were gently mixed by tapping. Sample tubes were incubated in a water bath at 37°C for 2 h. The desired enzyme cleavage was confirmed by electrophoresis following the conditions previously described.

2.14. Sequencing

Plasmid DNA from each sample was sent out for sequencing to the GeneLab in the School of Veterinary Medicine at Louisiana State University, Baton Rouge, LA. Nucleotide sequences were determined by automated sequence analysis using a Perkin Elmer/ABI Prism 3130 four capillary based DNA sequencer (Perkin Elmer, Foster City, CA) and Applied BioSystems BigDye Terminator version 3.1 (Perkin Elmer, Foster City, CA) and analyzed using ABI Sequencing Analysis 5.3.1 software. The sequences obtained were compared against

sequences in the GenBank nucleotide collection through the Basic Local Alignment Search Tool (BLAST) available on the National Center for Biotechnology Information (NCBI) website (www.ncbi.nlm.nih.gov/). Sequencing was done twice from independent PCR runs to compensate for any Taq polymerase errors (Cline et al., 1996) that could have occurred during the PCR process.

2.15. Lipid Characterization and Quantification

2.15.1. Algae Inoculum Density. Cell counts for all seed cultures were determined by the use of an Improved Neubauer 0.1 mm Deep Bright-Line hemacytometer (American Optical Corporation, Buffalo, NY). In brief, dilutions of the four strains and control were made (APPENDIX 4), thoroughly mixed, placed on the hemacytometer, and finally observed under a microscope (Leitz SM-LUX Microscope, Wetzlar, Germany). The cells present on either four or five specific squares (APPENDIX 5) were counted and the cell concentration per milliliter determined using the following calculations:

- Total cell count in 4 squares x 2,500 x dilution factor [For Diatom (FC2) only]
- Total cell count in 5 squares x 50,000 x dilution factor

2.15.2. Growth Conditions. Two batches (A and B) were grown using slightly different conditions to evaluate the fatty acid content of the identified strains.

Batch A: Five liters of each of the identified strains were grown in 3 gallon glass carboys using BG-11 media (Cyanobacteria-N3), 50X F/2 enriched with 1 % Na₂SiO₃·9 H₂O % w/v (Diatom-FC2) and BBM (Green algae FC1, D2 and *Chlorella vulgaris* UTEX 259). The inoculum used was 50 ml L⁻¹ on all strains except for the Diatom-FC2, for which 100 ml L⁻¹ were used. Air (0.037 % CO₂) was used for mixing at a flow rate of 190 SCCM (standard cubic centimeters per minute) under continuous illumination (59 μmol photons m⁻² s⁻¹, measured using an LI-193 spherical quantum sensor along with a LI-1400 data logger) at 25°C for 15 days with the

exception of FC2 (diatom), which was grown for 21 days. Cultures were harvested by centrifugation at 13,300 x g using a Sorvall RC-5 Superspeed refrigerated centrifuge. Harvested biomass was washed with DI water 2-3 times to reduce the salt content. Microalgae paste was subsequently freeze-dried using a Millrock Technology LD53 freeze-dryer (Millrock Technology Inc., Kingston, NY) with a preset program (APPENDIX 6) for 15 h. The total dry weight was determined for each of the strains. Analyses were run in duplicate.

Batch B: Strains were grown using the same conditions as in Batch A, but providing mixing through pumping of 5 % CO₂ enriched air (American Air Liquide, Houston, TX) and the addition of a 0.05 M Sodium Phosphate buffer to adjust the pH in the growth media. Cultures were harvested after 10 days and after 21 days for FC2 (diatom). Analyses were run in duplicate.

2.15.3. Lipid Extraction and Chlorophyll Determination. Lipids were extracted from freeze-dried biomass using a variation of the Folch Method (Folch et al., 1957; Iverson et al., 2001). In brief, 20 parts of 2:1 chloroform/methanol were added to 1 part biomass. Next, 8-10 glass beads were placed in each tube, closed tightly and vortexed for 2 min. This mixture was then filtered using Whatman Grade 42 ashless filters with vacuum and then washed several times with a solution of 2:1 chloroform/methanol. A solution containing 0.88 % NaCl in water was added to obtain a combined filtrate with a final ratio of 8:4:3 chloroform/methanol/water. The final biphasic system was centrifuged, and the lower phase was collected into a preweighed glass tube. Chlorophyll *a* and *b* were determined at 664 and 646 nm, respectively, using a Beckman Coulter DU 800 UV/VIS spectrophotometer (Beckman Coulter Inc., Brea, CA). Absorbance units obtained were divided by molar absorptivity coefficients proposed by Jeffrey and Welschmeyer, (1997) where:

- Chlorophyll *a*: 87.67 L g⁻¹ cm⁻¹

- Chlorophyll b: $51.36 \text{ L g}^{-1} \text{ cm}^{-1}$

in the equation to calculate pigment concentration:

- Concentration of Pigment (mg/L) :
$$\left(\frac{\text{Absorbance units (Au)}}{\text{Specific Absorption Coefficient (L g}^{-1} \text{ cm}^{-1})} \right)$$

The results for both types of chlorophyll were subtracted from the total lipid. Following this, the extract was evaporated to dryness under N_2 using a water bath at 30°C and a handmade evaporator. The tubes were wiped dry using kimwipes, dried in an oven at 55°C for an hour, let cool to 20°C in a dessicator for 30 min and the weights were recorded. Extractions for each sample were done in duplicate.

2.15.4. Fatty Acids Analysis. Fatty acid methyl esters (FAME) were prepared by transesterification of lipid extracts as described by Xu et al. (2010). Briefly, $40 \mu\text{l}$ of tricosanoic acid ($1000 \mu\text{g/ml}$) were added to each of the tubes containing a minimum of 2 mg of lipid. Two ml of MeOH:Benzenes at a ratio of 4:1 ($\rho = 0.8045 \pm 0.012 \text{ g/cm}^3$) were added to each tube and vortexed for 30 s. The samples were chilled at -74°C for 15 min followed by the addition of $200 \mu\text{l}$ of acetyl chloride. All tubes were flushed with N_2 for five seconds. Samples were transesterified by keeping them in the dark at 20°C for 24 h. Five ml of a saturated solution of NaHCO_3 was added to each tube to neutralize the mixture and stop the reaction. Samples were vortexed for 30 s and $40 \mu\text{l}$ of nervonic acid methyl ester ($1000 \mu\text{g/ml}$) was added as internal standard. Samples were centrifuged for 10 min at $900 \times g$. The top layer ($150 \mu\text{l}$) was collected and $3 \mu\text{l}$ injected into a GC (5890 Series II gas chromatograph, Hewlett-Packard, Avondale, PA) equipped with a flame ionization detector. FAME analyses were carried out with a $60 \text{ m} \times 0.32 \text{ mm} \times 0.5 \mu\text{m}$ ZB-Wax + (Phenomenex, Torrance, CA) column. The GC temperature program

was as follows: 150°C for 3 min, 200°C at increments of 8°C per minute for 15.5 min, 250°C at increments of 8°C per minute for 20.5 min and at 280°C at increments of 10°C per minute for 11 min. Helium was used as the carrier at a flow rate of 1.2 ml/min. Fatty acid identification was done by comparison of retention times with known standards (Supelco F.A.M.E. mix C4-C24, product number 18919-1AMP, Sigma-Aldrich, St. Louis, MO). Percent recovery for tricosanoic acid and nervonic acid methyl ester was calculated using the formula: $C_{\text{exp}} / C_{\text{Theo}} \times 100$, where C_{exp} = concentration of the standard based on the GC peak area (chromatogram) and C_{Theo} = concentration of the standard added to each sample.

3. RESULTS

3.1. Algal Isolation

After growth for 2-3 weeks, cultures were heterogeneous at this stage. Colony homogenization was achieved by a serial dilution process. Samples were plated onto BG-11 and BBM for 2-4 weeks and observed under a light microscope at 250X. Four out of the 17 initially isolated colonies were selected for this study based on rapid growth, morphological and color diversity, and homogeneity. They were tentatively identified as cyanobacteria (N3), diatom (FC2), and green algae (FC1 and D2) (Figure 3.1).




Figure 3.1. Microscopic Images for A) N3: *Synechococcus* sp., B) FC2: *Sellaphora pupula*, C) FC1: *Chlorella sorokiniana*, D) D2: *Scenedesmus abundans*, and E) *Chlorella vulgaris* UTEX 259 (control) as identified by BLAST. Images were taken at 250X using a Leica TCS SP2 spectral confocal microscope.

3.2. DNA Extraction

Cells (N3, FC2, FC1, D2 and *Chlorella vulgaris* UTEX-259) were harvested through centrifugation and their genomic DNA extracted (in duplicate) and quantified. DNA concentration ranged from 13- 440 ng/µl. Samples with DNA bands that showed high molecular weights and brightness were selected for PCR analysis.

3.3. PCR Products

Following PCR reactions, the quality of the PCR products was determined through agarose gel electrophoresis. Bands were observed in all nine conditions with isolate N3; whereas, five conditions showed bands with isolate D2 and only three conditions showed bands with isolates FC2 and FC1 (Figure 3.2). The molecular weights obtained for PCR products were 1.6 kb for N3 using the 16S primers, 1.8 kb for FC2 with the 18S primers, 1.8 kb for FC1 using the 18S primers, and 2.2 kb for D2 with the 18S primers. A PCR product of 1.8 kb was obtained for *Chlorella vulgaris* UTEX 259 using the 18S primers.

3.4. PCR Optimization

The best PCR conditions, based on PCR product quality results, were determined for each organism and were as follow: N3 (1L) no DMSO or glycerol at 1.5 mM MgCl₂, FC2 (1H) no DMSO or glycerol at 4.5 mM MgCl₂, FC1 (2M) with DMSO at 3 mM MgCl₂, D2 (3M) with glycerol at 3 mM MgCl₂ and *Chlorella vulgaris* UTEX-1 (3M) (gel not shown) with glycerol and 3 mM MgCl₂ (Figure 3.2).

3.5. Purification of Plasmid DNA

Plasmid DNA concentration for each microorganism was determined after purification using 2 µl of sample DNA and reading it on a nanodrop spectrophotometer. Plasmid DNA concentration for the samples averaged 390 ng/µl for N3, 109 ng/µl for FC2, 273 ng/µl for FC1, 300 ng/µl for D2, and 405 ng/µl for *Chlorella vulgaris* UTEX.

3.6. Restriction Enzyme Digestion

After EcoRI enzymatic digestion all selected transformant colonies had inserts (Figure 3.3). For N3, two bands that were close together could be seen on each of the samples which corresponded to the PCR product insert bands. With regards to the samples amplified using 18S

rRNA gene primers, it was noticeable that the inserts for D2 (Lanes 7- 9) were larger than the rest.




Figure 3.2. PCR Amplification Results Gel. PCR product quality of nine conditions for N3 (A), FC2 (B), FC1 (C), and D2 (D) mixed with $MgCl_2$ at 1.5 mM (low concentration, L), 3 mM (medium concentration, M), and 4.5 mM (high concentration, H).




Figure 3.3. Restriction Enzyme Digestion Gel. Lane M, molecular weight marker. Lane 1, N3-1. Lane 2, N3-2. Lane 3, N3-3. Lane 4, FC1-1. Lane 5, FC1-2. Lane 6, FC1-3. Lane 7, D2-1. Lane 8, D2-2. Lane 9, D2-3. Lane 10, *C. vulgaris* UTEX-1. Lane 11, *C. vulgaris* UTEX-2. Lane 12, *C. vulgaris* UTEX-3. Lane 13, FC2-1. Lane 14, FC2-2. Lane 15, FC2-3.

3.7. Sequencing

Plasmid DNA for each strain was sent out for sequencing and the sequences obtained were compared against sequences on the GenBank at the NCBI website. It was noticed that the

sequences amplified using M13 forward and M13 reverse primers (GeneLab) did not possess the length required to cover the amplified sequence of either 16S rRNA or 18S rRNA genes on the samples. Therefore, new primers were designed to fill that gap (Table 2.1). After the new sequences were obtained from all microorganisms from the LSU GeneLab, they were analyzed by pair-wise comparison and a final sequence was assembled. The obtained sequences were once again compared to the sequences available on the GenBank at the NCBI website.

3.8. BLAST

BLAST on the NCBI website indicated that N3 had a 99 % maximum identification with a *Synechococcus* sp. TAG 16S ribosomal RNA gene partial sequence with the accession number AF448066.1. FC2 results showed a 98 % maximum identification with a *Sellaphora pupula* clone RBG1 18S rRNA gene partial sequence with accession number EF151962.1. FC1 showed a 99 % maximum identification with a *Chlorella sorokiniana* 18S rRNA gene, strain Prag A14 with accession number X74001.1. D2 showed a 99 % maximum identification with *Scenedesmus abundans* gene for 18S small subunit rRNA with accession number X73995.1. For the control *Chlorella vulgaris* UTEX 259, there was a maximum identification of 99 % with *Chlorella vulgaris* genes for 18S rRNA, ITS1, 5.8S rRNA, and ITS2 with accession number AB162910.1 (Table 3.1). Both runs for sequences for each organism, which were obtained from independent PCR runs, were found to be consistent showing only minimal nucleotide differences between them.

3.9. Algae Inoculum

Cell counts from inoculums for both batches are depicted in Table 3.2. Some differences were noticed between batches with *Synechococcus* sp. having a higher cell count (1.12×10^8) in Batch A as compared to Batch B (4.82×10^7).

Table 3.1. Codes for isolated microorganisms with tentative identification based on morphology, similarity between amplified sequences and nearest recorded sequence in NCBI using BLAST and accession numbers for NCBI sequences.

Microorganism	Similarity with the nearest recorded sequence	Accession number
N3 (cyanobacteria)	99 % with <i>Synechococcus</i> sp. TAG 16S	AF448066.1
FC2 (diatom)	98 % with <i>Sellaphora pupula</i> clone RBG1	EF151962.1
FC1 (green algae)	99 % with <i>Chlorella sorokiniana</i> strain Prag A14	X74001.1
D2 (green algae)	99 % with <i>Scenedesmus abundans</i>	X73995.1
<i>C. vulgaris</i> *	99 % with <i>Chlorella vulgaris</i>	AB162910.1

*= *Chlorella vulgaris* UTEX 259 (control).

Table 3.2. Algae inoculums for Batch A and Batch B in cells ml⁻¹

Strain	Density	
	Batch A	Batch B
<i>Synechococcus</i> sp.	1.12 X 10 ⁸	4.82 X 10 ⁷
<i>Sellaphora pupula</i>	2.75 X 10 ⁴	1.50 X 10 ⁴
<i>Chlorella sorokiniana</i>	4.80 X 10 ⁶	6.20 X 10 ⁶
<i>Scenedesmus abundans</i>	1.06 X 10 ⁷	1.06 X 10 ⁷
<i>Chlorella vulgaris</i> UTEX 259	3 X 10 ⁶	8.1 X 10 ⁶

3.10. Biomass for Lipid Extraction

The dry weights of each microorganism used for lipid extraction studies are presented in Table 3.3. Compared to control (0.037 % CO₂), most strains resulted in an increase in total dry biomass when grown using 5 % CO₂, with the exception of *Synechococcus* sp., which mass decreased from 435 to 168 mg of dry biomass.

3.11. Total Extracted Lipid and Chlorophyll Content

Extracted lipids on dry weight biomass for each strain are summarized in Table 3.4. A decrease in the total lipid portion was observed on three out of the five evaluated strains when compared to the control batch (Batch A), with the exception of *Scenedesmus abundans* which resulted in an insignificant increment and *C. vulgaris* UTEX 259 with almost a two-fold increment.

Fatty acid profiles for identified strains are summarized in Tables 3.5. The total amount of fatty acids decreased for three of the strains when using CO₂. The most significant difference in the amount of fatty acids was observed with *Sellaphora pupula* when grown using air (54.8 mg g⁻¹ dry biomass) as compared to using CO₂ (24.4 mg g⁻¹ dry biomass). The amount of fatty acids for *Synechococcus* sp. and *Chlorella sorokiniana* decreased slightly from 20.63 mg g⁻¹ dry biomass and 29.82 mg g⁻¹ dry biomass to 17.62 mg g⁻¹ dry biomass and 23.99 mg g⁻¹ dry biomass, respectively. However, for *Scenedesmus abundans* an increment was seen from 14.14 mg g⁻¹ dry biomass to 31.63 mg g⁻¹ dry biomass. *Chlorella vulgaris* UTEX 259 presented the most significant increment from 15.14 mg g⁻¹ dry biomass to 47.83 mg g⁻¹ dry biomass, with more than half of that total belonging to the polyunsaturated fatty acid (PUFA) fraction. Recoveries for internal (methyl nervonate) and external (methyl tricosanoate) standards were 88% and 82% on average, respectively. The total chlorophyll concentration was negligible averaging 0.007 mg g⁻¹ of oil.

Table 3.3. Total dry biomass for lipid extraction.

Strain	mg dry biomass	
	Batch A *	Batch B**
<i>Synechococcus</i> sp.	435	168
<i>Sellaphora pupula</i>	62	100
<i>Chlorella sorokiniana</i>	158	847
<i>Scenedesmus abundans</i>	210	682
<i>Chlorella vulgaris</i> UTEX 259	294	575

* = Strains grown for 15 days with mixing using air (0.037 % CO₂). **= Strains grown for 10 days with mixing using 5 % CO enriched air.

Table 3.4. Total lipid extracted from biomass (g/100 g dry biomass).

Strain	Batch A*	Batch B**
<i>Synechococcus</i> sp.	9.17	6.9
<i>Sellaphora pupula</i>	19.52	12.12
<i>Chlorella sorokiniana</i>	11.56	10.95
<i>Scenedesmus abundans</i>	11.61	11.8
<i>Chlorella vulgaris</i> UTEX 259	7.36	13.65

* = Strains grown for 15 days with mixing using air (0.037 % CO₂). **= Strains grown for 10 days with mixing using 5 % CO₂ enriched air.

Table 3.5. Fatty acid profiles for evaluated strains grown in air (0.037 % CO₂) or 5 % CO₂ in mg g⁻¹ dry biomass. Results are the average means of duplicates along with standard deviations.

Fatty Acid	<i>Synechococcus</i> sp.		<i>Sellaphora pupula</i>		<i>Chlorella sorokiniana</i>		<i>Scenedesmus abundans</i>		<i>Chlorella vulgaris</i> UTEX 259	
	Air (0.037 % CO ₂)	5 % CO ₂	Air (0.037 % CO ₂)	5 % CO ₂	Air (0.037 % CO ₂)	5 % CO ₂	Air (0.037 % CO ₂)	5 % CO ₂	Air (0.037 % CO ₂)	5 % CO ₂
Saturated Fatty Acids (SFA's)										
C14:0	5.85 ± 0.03	5.24 ± 0.64	3.73 ± 0.14	1.33 ± 0.05	1.01 ± 0.008	1.22 ± 0.03	0.78 ± 0.04	2.24 ± 0.01	0.56 ± 0.06	2.29 ± 0.05
C16:0	3.57 ± 0.03	3.13 ± 0.16	8.99 ± 0.57	6.02 ± 0.39	7.76 ± 0.96	6.90 ± 2.24	3.15 ± 0.11	6.56 ± 0.84	3.20 ± 0.42	10.35 ± 0.77
C18:0	0.50 ± 0.05	0.85 ± 0.02	2.85 ± 0.13	1.26 ± 0.005	0.93 ± 0.01	1	2.09 ± 0.03	ND	0.79 ± 0.06	ND
Total SFAs	9.92	9.22	15.57	8.61	9.7	9.12	6.02	8.8	4.55	12.64
Monounsaturated Fatty Acids (MUFA's)										
C16:1	10.23 ± 0.004	6.24 ± 0.56	16.07 ± 1.50	9.24 ± 0.78	1.54 ± 0.01	2.03 ± 0.82	1.22 ± 0.03	4.13 ± 0.56	0.41 ± 0.09	5.41 ± 0.53
C18:1	0.48 ± 0.07	0.89 ± 0.03	3.06 ± 0.06	1.40 ± 0.007	1.42 ± 0.14	1.52 ± 0.21	1.14 ± 0.02	3.73 ± 0.47	0.71 ± 0.06	2.02 ± 0.45
Total MUFAs	10.71	7.12	19.13	10.64	2.96	3.55	2.36	7.86	1.12	7.43
Polyunsaturated Fatty Acids (PUFA's)										
C18:2	ND	0.34 ± 0.02	1.43 ± 0.03	0.73 ± 0.22	6.91 ± 0.16	5.87 ± 1.18	2.41 ± 0.07	2.20 ± 0.19	1.68 ± 0.24	3.59 ± 0.56
C18:3	ND	0.94	1.62 ± 0.05	1.87 ± 0.0001	10.25 ± 0.19	6.65 ± 0.21	3.35 ± 0.12	12.77 ± 1.81	7.79 ± 1.31	24.17 ± 4.78
C20:4	ND	ND	17.08 ± 0.06	2.55 ± 0.27	ND	ND	ND	ND	ND	ND
Total PUFAs	0	1.28	20.13	5.15	17.16	12.52	5.76	14.97	9.47	27.76
Fatty Acids*	20.63	17.63	54.8	24.4	29.82	23.99	14.14	31.63	15.14	47.83

ND= None detected; *= total fatty acids.

4. DISCUSSION

First described in the late 1980's, PCR is one of the most widely used methods in molecular biology (Saiki et al., 1988). Several decades later, attempts for PCR optimization continued to be made to meet its specific objectives (Dieffenbach and Dveksler, 2003). Templates known for being difficult to sequence include those with high guanine-cytosine (G/C) content, high adenine-thymine (A/T), as well as sequences with marked secondary structure or large regions of homopolymer (Stirling, 2003). In regards to sequences with high G/C content, several approaches have been taken to solve this problem and perhaps the most successful method for improving results is the inclusion of certain organic additives in the reaction mixture, such as DMSO, betaine, polyethylene glycol, glycerol and formamide (Chakrabarti and Schutt, 2001). The effect DMSO exerts in the PCR amplification of some GC-rich sequences is a largely studied one (Pomp and Medrano, 1991; Sun et al., 1993; Sidhu et al., 1996).

In this study, the DNA treatment containing neat DMSO and 3 mM MgCl₂ was determined to be the best PCR condition for FC1. Kang et al. (2005) conducted a study to find the best PCR conditions for the uniform amplification of random sequence templates, which can be disrupted by high GC content. They determined that the effect of additives on PCR was more outstanding in the presence of 5 % DMSO, which increased the ratio of full length to short products by 120 % (Kang et al., 2005). For D2 and *Chlorella vulgaris* UTEX 259, the best condition was determined to be the one that had glycerol at 50 % and MgCl₂ at 3 mM. Nagai et al, (1998) conducted a study to evaluate the effect of various additives, among them glycerol, on PCR. Their results indicated that the addition of either one in the reaction mixture allowed the specific amplification when an enterohemorrhagic *E. coli* DNA fragment was used as template. The improvement of glycerol on PCR efficiency and specificity can possibly be explained by (1)

glycerol's enhancement of the hydrophobic interactions between protein domains (Back et al., 1979); (2) lowering of the strand-separation temperature (Wang et al., 1993); and (3) a raise on proteins' thermal transition temperature (Back et al., 1979; Gekko and Timasheff, 1981).

Magnesium Chloride (MgCl_2) is a necessary co-factor for all Type II enzymes which include restriction endonucleases and polymerases. Mg^{2+} binds to the enzyme inducing a shift which puts the subunits in a proper configuration. It is assumed that a significant reduction in MgCl_2 concentration may prevent a sufficient number of enzyme molecules from being in the correct conformation for good amplification to happen (IDT, 2005). However, it is also known that excessive MgCl_2 can be equally harmful many times, with a substantial increase in secondary products produced by non-specific priming being the usual response to excess Mg^{2+} (IDT, 2005).

In the present study, three different concentrations of MgCl_2 were tested (1.5, 3, 4.5 mM). Three mM of MgCl_2 was optimum for the amplification of FC1, D2 and *C. vulgaris*, 1.5 mM for N3 and 4.5 mM for FC2. A study performed using genomic DNA from selected strains of *Vibrio vulnificus* to obtain DNA fingerprint profiles with arbitrarily primed polymerase chain reaction (AP-PCR) demonstrated that when using a MgCl_2 concentration of 2.5 mM and PCR cycling parameters in combination with 1 μg of purified genomic DNA, 1.04 mM of R-PSE420 oligonucleotide primer and thermal cycling protocols with stepwise increments in the annealing temperatures, reproducible DNA fingerprints free of primer artifacts were generated (Vickery et al., 1998). The aforementioned MgCl_2 concentration is close to the medium (3 mM) concentration used in the present study which was optimum for the amplification of three out of the five organisms subjected to algal identification.

The size of the PCR products obtained in this study were ~1.6 kb for N3, using a set of 16S rRNA primers, and ~1.8 kb for FC2, FC1 and *Chlorella vulgaris* UTEX 259. The PCR

product for D2; however, had a size of 2.2 kb, which were observed on both, the PCR product gel and on the restriction enzyme digestion gel. This was confirmed by the fact that the final sequence assembled for D2 had a total of 2, 200 bp, suggesting that this particular gene on D2 was different compared to the other isolates amplified by the 18S rRNA primers. Therefore, this organism could be considered a different strain within its genus since the query coverage for D2 was only 81 % (results not shown) and it was the lowest among the sample strains (query coverage for the other microorganisms was between 98-100 %).

The NCBI website identified matches for all four sample strains. N3 was identified as *Synechococcus* sp. TAG 16S ribosomal RNA gene, partial sequence, FC2 as *Sellaphora pupula* clone RBG1 18S rRNA gene, partial sequence, FC1 as *Chlorella sorokiniana* 18S rRNA gene, strain Prag A14, and D2 as *Scenedesmus abundans* gene for 18S small subunit rRNA.

PNNL Biofuels Scientific Focus Area (BSFA) have conducted research in cyanobacteria in an attempt to develop a better understanding of metabolic subsystems and regulatory networks involved in solar energy conversion to biofuel products (Bryant et al., 2011). A system approach to understanding issues such as metabolic modules, physiological constraints and maximum rates of carbon processing is being undertaken by exploiting the fast-growing cyanobacterium, *Synechococcus* sp. PCC 7002, which has the added benefit of a well-developed genetic system (Bryant et al., 2011). Cyanobacteria are said to be attractive candidates for biofuel producing microbial systems since they possess the favorable characteristics of both prokaryotes and plants. Cyanobacteria are photosynthetic organisms able to absorb solar energy and fix carbon dioxide, whereas other biofuel-producing microbes such as *E. coli*, *Zymomonas mobilis*, *Saccharomyces cerevisiae* are not capable of doing both (Lu, 2010).

Studies on oleaginous diatoms indicate that the average lipid content can be 22.7 % dry cell weight when grown under normal conditions, whereas under stress conditions, a total lipid content of 44.6 % dry cell weight was seen (Hu et al., 2008). *Sellaphora pupula* (Kutzing) Mereschkowsky (formerly *Navicula pupula* Kutzing) is a common, freshwater diatom species and a complex organism containing many pseudo- and semi-cryptic species (Mann, 1984, 1989, 1999; Mann and Droop, 1996; Mann and Kocioleck, 1990; Behnke et al., 2004; Evans et al., 2007, 2008). A member of this genera, namely *Navicula* sp., has been found to possess a fatty acid profile where 16:0, 16:1 and 20:5n3 make up more than 75 % of the total lipid content (Dunstan et al., 1994), and other studies have determined the presence of C₂₅ triene III, considered probably the most abundant and widely occurring highly branched isoprenoid (HIB) alkene observed in marine diatoms (Belt et al., 2001).

Chlorella species can be encountered in all water habitats, fresh and marine sources (Iwamoto, 2004). It contains essential amino acids, protein, minerals, vitamins, chlorophyll, and bioactive substances (Borowitzka, 1988; Schubert, 1988). Bio-energy generation from *Chlorella* is a relatively new aspect in renewable energy research (Phukan et al., 2011). Illman et al, (2000) studied calorific values of *Chlorella* strains (*Chlorella protothecoides*, *vulgaris*, *emersonii*, *sorokiniana*, and marine strain, *minutissima*) using medium with low nitrogen content and pointed out that *Chlorella* strains may be good prospects for diesel replacements. Chader et al. (2011) analyzed the major fatty acids in *Chlorella sorokiniana* using Gas Chromatography and by growing it on different types of media. It was observed that the strain was mainly composed of a mixture of unsaturated fatty acids, such as oleic (18:1), linoleic (18:2) and linolenic acid (18:3). Saturated fatty acids, palmitic (16:0) and stearic (18:0), were also present to a small extent. In a study by Knothe (2008), palmitic, stearic, oleic, and linoleic acids were recognized as the most

common fatty acids present in the biodiesel. In the three media tested (TAP, BG-11 and media +N), C18:1 and C18:2 were commonly dominant. Chader et al. (2011) reported that the most important fatty acids were C16:0, C18:0, C18:1, C18:2 and C18:3 regardless of the medium utilized. Scragg et al. (2003) used an emulsion consisting of transesterified rape seed oil, a surfactant and slurry of *Chlorella vulgaris* in an unmodified single cylinder diesel engine with good results; whereas, Xu et al. (2006) was able to produce high quality biodiesel using heterotrophically grown *Chlorella protothecoides*.

The genus *Scenedesmus*, belonging to the green algae, have become the equivalent of laboratory rats in many fields in limnology or fresh water science (Wiltshire et al., 2000). If one algal species had to be chosen for biodiesel production, the one with the most suitable fatty acid profile, specifically in terms of linolenic and polyunsaturated fatty acids, would be *Scenedesmus obliquus* (Gouveia and Oliveira, 2009). Mandal and Mallick, (2009), studied lipid accumulation in *Scenedesmus obliquus* when grown under nitrogen deficiency and observed that lipid was 43 % of dry cell weight compared to 12.7 % (dry cell weight) in the control; whereas, the lipid content increased up to 30 % when phosphorus deficiency and thiosulphate supplementation were used. When cultured for 8 days, the application of response surface methodology in combination with central composite rotary design (CCRD) resulted in a lipid yield of 61.3 % (against 58.3 % obtained experimentally) at 0.04, 0.03, and 1.0 g L⁻¹ of nitrate, phosphate, and sodium thiosulphate, respectively .

The four strains (*Synechococcus* sp., *Sellaphora pupula*, *Chlorella sorokiniana*, and *Scenedesmus abundans*) isolated from Louisiana's fresh and brackish water bodies represent potential candidates for the production of biodiesel. In regards to the algal biomass obtained, an increase in final dry biomass was observed with most strains when aeration with 5 % CO₂ was

used for culturing, with the exception of the cyanobacteria *Synechococcus* sp. *Synechococcus* sp. strain exhibited the lowest pH of all strains showing susceptibility to dissolved levels of CO₂. Wang et al. (2011) observed that significant inhibitory effects on the growth of two cyanobacterial species, *Microcystis aeruginosa* and *Anabaena spiroides*, were seen when three pH conditions (5.5, 6.0 and 6.5) were reached using aeration with CO₂. *Chlorella sorokiniana*, on the contrary, was the strain with the best response to CO₂ aeration resulting in a six-fold increase in dry biomass. A two-fold increase was observed for *Sellaphora pupula* and *Chlorella vulgaris* UTEX 259 and above a three-fold for *Scenedesmus abundans*. Huertas et al. (2000) reported that cultures of *Nannochloropsis gaditana* grown in the absence of dissolved inorganic carbon (DIC) and with aeration of less than 0.0001% (v:v) CO₂ considered to be low DIC conditions, presented a reduction in final cell biomass of approximately 56 % when compared to the biomass obtained in cultures grown under control conditions which equaled 2 mM DIC in the medium and 0.03% (v:v) CO₂. No growth was observed in *Nannochloris maculata* cultured under low DIC conditions. However, when a concentration of 1% (v:v) CO₂ in air (high DIC conditions) was used, only the growth of *N. maculata* was enhanced, concluding that the response to DIC was species dependent. It was also observed that the uptake of nitrate and phosphate by *N. maculata* was linked to the inorganic carbon level and determined that no nutrient absorption had been observed in the low DIC-culture. Growth, however, at the highest inorganic carbon concentration caused an acceleration of the uptake of these nutrients.

A decrease in the amount of total fatty acids extracted from *Synechococcus* sp., *Sellaphora pupula* and *Chlorella sorokiniana* was observed when the growth media was aerated with 5 % CO₂. The amount of total lipids decreased from 20.63, 54.8 and 29.82 g kg⁻¹ to 17.62, 24.4 and 23.99 g kg⁻¹, respectively. Similar observations have been reported by Sato (1989), when

supplementing 1 % and 3 % CO₂ to *Chlamydomonas reinhardtii* cells grown photoautotrophically. Total polar lipids decreased from 60.3 nmol/10⁷ using 0.04 % CO₂, to 44.2 nmol/10⁷ and 45.7 nmol/10⁷ when the media was supplemented with 1 and 3 % CO₂, respectively. The opposite was observed with *Scenedesmus abundans* and *Chlorella vulgaris* UTEX 259 where total lipids extracted increased from 14.14 g kg⁻¹ to 31.63 g kg⁻¹ and from 15.14 g kg⁻¹ to 47.83 g kg⁻¹, respectively. Tang et al. (2011) reported similar trends with *Scenedesmus obliquus* and *Chlorella pyrenoidosa* grown in media supplemented with 0.03 %, 5 %, 10 %, 20 %, 30 %, and 50 % CO₂. The total lipid content for *Scenedesmus obliquus* increased from 15.15 wt. % at 0.03 % CO₂ to 16.45 wt. % at 5 % CO₂, whereas for *Chlorella pyrenoidosa* the total lipid content decreased from 20.9 wt. % at 0.03 % CO₂ to 20.65 wt. % at 5 % CO₂. These results demonstrate that even within the green algae classification various responses could be observed.

Changes were exhibited in the distribution of saturated (SFAs), monounsaturated (MUFAs) and polyunsaturated (PUFAs) fatty acids when 5 % CO₂ was used. The trend observed for *Synechococcus* sp., *Sellaphora pupula* and *Chlorella sorokiniana* was a decrease on SFAs, MUFAs and PUFAs with the exception of a slight increase (~0.5 g kg⁻¹) in MUFAs for *Chlorella sorokiniana*. An increase in all three categories of fatty acids was observed for *Scenedesmus abundans* and *Chlorella vulgaris* UTEX 259. Similar observations on strains of *Scenedesmus obliquus* and *Chlorella pyrenoidosa* with an increase in total fatty acids when providing 5 % CO₂ compared to 0.03 % were reported by Tang et al. (2011). Tsuzuki et al. (1990) observed an increase in the composition of saturated fatty acids over that of unsaturated fatty acids for *Chlorella vulgaris* after 1h under CO₂ concentrations ranging from 0.036 % to 2 %.

The main fatty acids in *Synechococcus* sp. grown in media supplemented with air were C16:1, C14:0 and C16:0. These findings agree with those reported by Pratoomyot et al. (2005)

and Patil et al. (2007). For *Sellaphora pupula*, the main fatty acids were C20:4, C16:1 and C16:0. Dunstan et al. (1994) described 16:1n7, 16:0 and 20:5n3 as the main fatty acids found on *Navicula* sp. The main fatty acids detected in *Chlorella sorokiniana* were C18:3, C16:0, C17:0 and C18:2. Hu et al. (2008) reported C16:0, C18:2 and C18:3 as being the major fatty acids in a *Chlorella sorokiniana* strain. Significant amounts of 16:3 and 16:2n7 were also detected, which may possibly correspond to the C17:0 content identified for this strain in the present study since neither 16:3 nor 16:2n7 were part of the F.A.M.E. mix used for the characterization. Chader et al. (2011) reported a similar profile on a locally isolated *Chlorella sorokiniana* strain. The strain when grown using TAP media contained C18:3, C18:2, C18:1 and C16:0 as the main fatty acids. The main fatty acids for *Scenedesmus abundans* were C18:3, C16:0, and C18:2. These findings were in agreement with those reported by Pratoomyot et al. (2005). Isik et al. (1999) reported C16:0, C18:0 +1 and C18:3 as the main fatty acid components in *Scenedesmus abundans*. The main fatty acids detected in *Chlorella vulgaris* UTEX 259 were C18:3, C16:0 and C18:2 and are in agreement with the results reported by Isik et al. (1999).

The primary emission in flue gas is CO₂, which may be present at concentrations ranging from 3 % to 25 % (Packer, 2009). Microalgal fixation of CO₂ by photosynthesis is considered the most favorable potential method for CO₂ sequestration from flue gas through the incorporation of CO₂ into a biomass carbon source such as lipids (Lee and Lee, 2003; Doucha et al., 2005; Wang et al., 2008; Brune et al., 2009; Yoo et al., 2010; Ho et al., 2011). The use of biomass generated by capturing CO₂ from industrial processes through microalgae, followed by its utilization for transportation needs, can aid on CO₂ sequestration and it would also help to diminish overall carbon emissions (Chiu et al., 2011). Chiu et al. (2011) reported a *Chlorella* strain that when treated with intermittent flue gas aeration in a double-set photobioreactor system, had an average

efficiency of CO₂ removal from the flue gas that reached 60 %, and NO and SO₂ removal efficiency that was maintained at approximately 70 % and 50 %, respectively. A typical composition of emissions from flue gases from two different power plants is given in Tables 4.1 and 4.2.

Table 4.1. Flue gas composition of Wyodak PC power plant (Robertson, 2007).

Flue gas component	Concentration (by volume)
CO ₂	11.8 %
N ₂	67 %
H ₂ O	8 %
O ₂	12 %
SO ₂	180 ppm
NO _x	150 ppm
CO	300 ppm

Table 4.2. Typical untreated flue gas composition from a power plant burning low sulfur eastern bituminous coal. (Granite and Pennline, 2002).

Flue gas component	Concentration (by volume)
CO ₂	15-16 %
O ₂	3-4 %
H ₂ O	5-7 %
SO ₂	800 ppm
SO ₃	10 ppm
NO _x	500 ppm
CO	20 ppm
HCl	100 ppm
Total Hg	1 ppb
N ₂	Balance
Hydrocarbons	10 ppm

The fatty acid profiles in this study changed on each of the strains when grown in 5 % CO₂ enriched media. For *Synechococcus* sp., C16:1, C14:0 and C16:0 remained the major fatty acids with the appearance of C18:2 and C18:3 which were absent in the batch grown in the presence of air only. This can be attributed to the desaturation effect due to higher concentration of CO₂ as described by Tang et al. (2011). According to Knothe (2008), the most common fatty acid esters in biodiesel are C16:0, C18:0, C18:1, C18:2 and C18:3. This is true for biodiesel feedstocks such as soybean, sunflower, rapeseed, palm and peanut oils (Knothe, 2008).

The fatty acid profile of *Sellaphora pupula* changed considerably when grown using 5 % CO₂. In the case of the PUFA C20:4, it was reduced from 17.08 g kg⁻¹ to 2.55 g kg⁻¹ and a smaller yet not significant decrease was also observed for C16:1 and C16:0. A high content of PUFAs is usually not a desirable feature in a feedstock for biodiesel production since this can result in biodiesel with low oxidative stability, low viscosity and low cetane number, being the latter a dimensionless descriptor of the ignition quality of a diesel fuel (Knothe, 2008). *Sellaphora pupula* demonstrated the greatest lipid percentage yield (19.52 %) when grown with air fed into the media, and, just like in *Synechococcus* sp., additional testing to determine an ideal CO₂ level could result in better biomass yields. For *Chlorella sorokiniana*, C18:3, C16:0 and C16:2 remained the main fatty acids after CO₂ aeration. *Chlorella sorokiniana* had the best growth response yielding the most biomass when treated with 5 % CO₂. For *Scenedesmus abundans*, C18:3 and C16:0 remained the main fatty acids, but the MUFAs C16:1 and C18:1 displaced the PUFA C18:2 when grown using 5 % CO₂. This can be considered a positive shift since *Scenedesmus abundans* displayed the highest content of C18:1. Oleic acid has been suggested in the past as a compound for improving biodiesel fuel properties due to its low melting point (-20°C) (Knothe, 2008). Oleic acid possesses better oxidative stability than C18:2 and C18:3 and it

exceeds the minimum cetane number standards given by the EN 14214 and ASTM D6751 (Knothe, 2008). In addition, out of the four isolated strains, *Scenedesmus abundans* produced the most total fatty acids with 31.63 g kg⁻¹. The main fatty acids detected for *Chlorella vulgaris* UTEX 259 were C18:3, C16:0 and C16:1 in the presence of 5 % CO₂, compared to C18:3, C16:0 and C18:2 without CO₂ aeration. It can be highlighted here that a substantial increment in C18:3 from 7.79 g kg⁻¹ to 24.17 g kg⁻¹ was observed. However, since this was mostly PUFAs, *Chlorella vulgaris* UTEX 259, under these circumstances, would not have the fatty acid profile most suitable for biodiesel production; but, it did produce the most total fatty acids at 47.83 g kg⁻¹

Oxidative stability, poor low-temperature properties and a slight increase in NO_x exhaust emissions remain as the major challenges with biodiesel (Knothe, 2008). The challenge when addressing these problems is that solving one of them usually negatively affects another one and this observed pattern is related to the dependence of fuel properties on the fatty acid profile (Knothe, 2008). Cha et al. (2011) suggested that the use of a binary blending of microalgae system to attain an “ideal mix” of fatty acids could be pursued. In their study, they proposed a mixture of C16:1, C18:1 and C14:0 fatty acids in a ratio of 5:4:1 for biodiesel of very low oxidative potential but that could still offer both, a favorable cold filter plugging point (CFPP) rating and cetane number. Following this suggestion, not only could individual strains isolated in the present study be tested at a larger scale, but could also be combined to obtain a mixture that can offer an even better fatty acid profile and biomass productivity for biodiesel production when paired with the use of CO₂.

5. CONCLUSIONS

Four algae, two of them obtained from different brackish sources and the other two obtained from a fresh water source within the state of Louisiana's Southeast region, were tentatively identified as *Synechococcus* sp., *Sellaphora pupula*, *Chlorella sorokiniana*, and *Scenedesmus abundans*. The use of other taxonomical approaches is recommended to complement these findings.

The fatty acid profiles of the above mentioned organisms were altered when CO₂ aeration was incorporated into the growth media. *Synechococcus* sp.'s profile contained the major fatty acids that traditional biodiesel feedstocks possess. *Sellaphora pupula* had the best lipid percentage yield and the highest content of C16:1, which is considered to be the best fatty acid to improve biodiesel's characteristics after C18:1. *Scenedesmus abundans* displayed the highest content of C18:1. Oleic acid has been suggested in the past as a compound for enrichment in biodiesel fuels for improving biodiesel properties. *Scenedesmus abundans* can be considered as the second best candidate of all the strains analyzed for biodiesel feedstock. However, the lower C18:3 content observed in *Chlorella sorokiniana* allowed for a balanced fatty acid profile in terms of cetane number, oxidative stability, viscosity and low temperature conditions. In addition to the aforementioned properties, *Chlorella sorokiniana* had the greatest productivity when grown using CO₂, making it the best candidate for the production of biodiesel. The opposite can be said for the control *Chlorella vulgaris* UTEX 259 which contained 50 % of total fatty acid as C18:3 making it a less desirable candidate for biodiesel production in terms of oxidative stability.

An interesting approach could be the establishment of a binary production system between *Chlorella sorokiniana*, with a high biomass production, and *Scenedesmus abundans*, with a better fatty acid profile, to produce oil with superior quality and characteristics for the production of

biodiesel. Additionally, further testing should be made to determine if the isolated strains can tolerate higher CO₂ concentrations and evaluate their effect on biomass and fatty acid profile.

REFERENCES

- Abbasi, A., Amiri, S. (2008). Emulsifying behavior of an exopolysaccharide produced by *Enterobacter cloacae*. *African J Biotechnol.* 7: 1574-1576.
- Aita, G. M., Kim, M. (2010). Pretreatment Technologies for the Conversion of Lignocellulosic Materials to Bioethanol. In: Sustainability of the Sugar and Sugar-Ethanol Industries, American Chemical Society, pp. 117-145.
- Akkerman, I., Janssen, M., Rocha, J., Wijffels, R. H. (2002). Photobiological hydrogen production: photochemical efficiency and bioreactor design. *Int J Hydrogen Energy.* 27: pp. 1195-1208.
- Armbrust, V., Galindo, M. (2001). Rapid Evolution of a Sexual Reproduction Gene in Centric Diatoms of the Genus *Thalassiosira*. *Appl Environ Microb.* 67 (8): 3501-3513.
- Back, J. F., Oakenfull, D., Smith, M. B. (1979). Increased thermal stability of proteins in the presence of sugars and polyols. *Biochemistry.* 18: 5191-5196.
- Banerjee, A., Sharma, R., Chisti, Y., Banerjee, U. C. (2002). *Botryococcus braunii*: a renewable source of hydrocarbons and other chemicals. *Crit Rev Biotechnol.* 22: 245-79.
- Barbosa, M., Jansen, M., Ham, N., Tramper, J., Wijffels, R. H. (2001). Microalgae Cultivation in air-lift reactors. *Biotechnol Bioeng.* 82 (2): 170-179.
- Bashan, Y., Levanony, H. (1990). Current status of *Azospirillum* inoculation technology: *Azospirillum* as a challenge for agriculture. *Can J Microbiol.* 36: 591-608.
- Bashan, Y., Holguin, G. (1997). *Azospirillum*-plant relationships: environmental and physiological advances. *Can J Microbiol.* 43: 103-21.
- Bashan, Y., Holguin, G., de-Bashan, L. E. (2004). *Azospirillum* plant relationships: physiological, molecular, agricultural, and environmental advances. *Can J Microbiol.* 50: 521-577.
- Becker, E. W. (1994a). Culture Media. In: *Microalgae: Biotechnology and microbiology.* (Ed Becker, E. W.), pp. 9-41. Cambridge University Press. Cambridge, UK.
- Becker, E. W. (1994b). Chemical Composition: Lipids. In: *Microalgae: Biotechnology and microbiology.* (Ed Becker, E. W.), pp. 179-183. Cambridge University Press. Cambridge, UK.
- Behnke, A., Friedl, T., Chepurinov, V. A., Mann, D. G. (2004). Reproductive compatibility and rDNA sequence analyses in the *Sellaphora pupula* species complex (Bacillariophyta). *J Phycol.* 40: 193-208.

- Belt, S., Masse, G., Allarda, G., Robert, J. M., Rowland, S. (2001). Identification of a C₂₅ highly branched isoprenoid triene in the freshwater diatom *Navicula sclesvicensis*. *Org Geochem.* 32 (9): 1169-1172.
- Benemann, J. R., Weissman, J. C. (1984). Chemicals from Microalgae. In: *Bioconversion systems.* (Ed. Wise, D.), pp. 62-64. CRC Press, Inc. 2000 Corporate Blvd., N.W., Boca Raton, Florida, USA.
- Blanchard, M. M., Taillon-Miller, P., Nowotny, P. (1993). PCR buffer optimization with uniform temperature regimen to facilitate automation. *Genome Res.* 2: 234-240.
- Borowitzka, M. A. (1988). Vitamins and fine chemicals from micro-algae. In: *Micro-algal biotechnology.* (Ed. Borowitzka, L.J., Borowitzka M. A.), pp. 153-196. Cambridge University Press, UK.
- Brennan, L., Owende, P. (2010). Biofuels from Microalgae: A review of technologies for production, processing and extractions of biofuels and co-products. *Renew Sust Energ Rev.* 14: 557-577.
- Brune, D. E., Lundquist, T. J., Benemann, J. R. (2009). Microalgal biomass for greenhouse gas reductions: potential for replacement of fossil fuels and animal feeds. *J Environ Eng. ASCE* 135: 1136-1144.
- Bryant, D. A., Konopka. A. E., Pinchuk, G. E., Stolyar, S. M., Posewitz, M. C., Beliaev, P. I. (2011). *Synechococcus* sp. PCC 7002: A robust cyanobacterial platform for systems biology and biofuels engineering. Abstract 71 in *Systems Biology for DOE Energy Missions Joint Meeting 2011*, Crystal City, Virginia.
- Buchheim, M. A., Turmel, M., Zimmer E. A., Chapman, R. L. (1990). Phylogeny of *Chlamydomonas* (Chlorophyta) based on cladistic analysis of nuclear 18S rRNA sequence data. *J Phycol.* 26: 689-699.
- Buchheim, M. A., Lemieux, C., Otis, C., Gutell, R. R., Chapman, R. L., Turmel, M. (1996). Phylogeny of the Chlamydomonales (Chlorophyceae): a comparison of ribosomal RNA gene sequences from the nucleus and the chloroplast. *Mol Phyl Evol.* 5: 391-402.
- Buchheim, M. A., Buchheim, J. A., Chapman, R. L. (1997). Phylogeny of *Chloromonas* (Chlorophyceae): a study of 18S rRNA gene sequences. *J Phycol.* 33: 286-293.
- Camacho, R. F., Garcia, C. F., Fernandez, S. J. M., Chisti, Y., Molina, G. E. (2003). A mechanistic model of photosynthesis in microalgae. *Biotechnol Bioeng.* 81 (4): 459-473.
- Campbell, C. J. (1997). *The coming oil crisis.* Multi-science Publishing Company and Petro-Consultants S.A, Essex, England.

- Castro, A. S., Tavano Garcia, V. M. (2005). Growth and biochemical composition of the diatom *Chaetoceros cf. wighamii brightwell* under different temperature, salinity and carbon dioxide levels. I: protein, carbohydrate and lipids. *Aquaculture*. 246: 405-412.
- Cha, T. S., Chen, J. W., Goh, E. G., Aziz, A., Loh, S. H. (2011). Differential regulation of fatty acid biosynthesis in two *Chlorella* in response to nitrate treatments and the potential of binary blending of microalgae oils for biodiesel application. *Bioresour Technol*. 102 (22): 10633-10640.
- Chader, S., Mahmat, B., Chetehouna, K., Mignolet, E. (2011). Biodiesel production using *Chlorella sorokiniana* a green microalga. *REN*. 14: 543-552.
- Chakrabarti, R., Schutt, C. E. (2001). The enhancement of PCR amplification by low molecular weight amides. *Nucleic Acids Res*. 29: 2377-2381.
- Chien, A., Edgar, D. B., Trela, J. M. (1976). Deoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*. *J Bact*. 127 (3): 1550-1557.
- Chisti, Y. (2007). Biodiesel from microalgae. *Biotechnol Adv*. 25 (3): 294-306.
- Chisti, Y. (2008). Biodiesel from microalgae beats bioethanol. *Trends Biotechnol*. 26 (3): 126-131.
- Chiu, S. Y., Kao, C. Y., Huang T. T., Lin C. J., Ong S. C., Chen C. D., Chang J. S., Lin, C. S. (2011). Microalgal biomass production and on-site bioremediation carbon dioxide, nitrogen oxide and sulfur dioxide from flue gas using *Chlorella* sp. cultures. *Bioresour Technol*. 102: 9135-9142.
- Chu, W. L., Phang, S. M., Goh, S. H. (1996). Environmental effects on growth and biochemical compositions of *Nitzschia inconspicua* Grunow. *J Appl Phycol*. 8 (4-5): 389-396.
- Cline, J., Braman J. C., Hogrefe H. H. (1996). PCR fidelity of *Pfu* DNA polymerase and other thermostable DNA polymerases. *Nucleic Acids Res*. 24 (18): 3546-3551.
- Cobb, B. D., Clarkson, J. M. (1994). A simple procedure for optimizing the polymerase chain reaction (PCR) using modified Taguchi methods. *Nucleic Acids Res*. 22: 3801-3805.
- de-Bashan, L. E., Bashan, Y., Moreno, M., Levsky, V. K., Bustillos J. J. (2002). Increased pigment and lipid content, lipid variety, and cell and population size of the microalgae *Chlorella* spp. when co-immobilized in alginate beads with microalgae-growth-promoting bacterium *Azospirillum brasilenses*. *Can J Microbiol*. 48: 528-541.
- Dieffenbach, C. W., Dveksler, G. S. (2003). *PCR Primer: A laboratory manual*. Cold Spring Harbor Laboratory Press. New York, USA.

- Dote, Y., Sawayama, S., Inoue, S., Minowa, T., Yokoyama, S. (1994). Recovery of liquid fuel from hydrocarbon-rich microalgae by thermochemical liquefaction. *Fuel*. 73 (12): 1855-1857.
- Doucha, J., Straka, F., Livansky, K. (2005). Utilization of flue gas for cultivation of microalgae *Chlorella* sp. in an outdoor open thin-layer photobioreactor. *J Appl Phycol*. 17: 403-412.
- Dubinsky, Z., Matsukawa, R., Karube, I. (1995). Photobiological aspects of algal mass Culture. *J Mar Biotechnol*. 2: 61-65.
- Dunahay, T. G., Jarvis, E. E., Dais, S. S., Roessler, P. G. (1996). Manipulation of microalgal lipid production using genetic engineering. *Appl Biochem Biotechnol*. 57-58 (1): 223- 231.
- Dunstan, G., Volkman, J., Barrett, S., Leroi, J. M., Jeffrey, S. W. (1994). Essential polyunsaturated fatty acids from 14 species of diatom (Bacillariophyceae). *Phytochemistry*. 35 (1): 155-161.
- EIA (2006). International carbon dioxide emissions from the consumption of energy. Available from: <http://www.eia.doe.gov/pub/international/iealf/tableh1co2.xls>.
- Erwin, P., Thacker, R. (2008). Cryptic diversity of the symbiotic cyanobacterium *Synechococcus spongiarum* among sponge hosts. *Mol Ecol*. 17: 2937-2947.
- Evans, K. M., Wortley, A. H., Mann, D. G. (2007). An assessment of potential diatom “barcode” genes (cox1, rbcL, 18S and ITS rDNA) and their effectiveness in determining relationships in *Sellaphora* (Bacillariophyta). *Protist*. 158: 349-364.
- Evans, K. M., Wortley, A. H., Simpson, G. E., Chepurinov, V. A., Mann, D. G. (2008). A molecular systematic approach to explore diversity within the *Sellaphora pupula* species complex (Bacillariophyta). *J Phycol*. 44: 215-231.
- FAO (2007). Sustainable bioenergy: a framework for decision makers. United Nations Energy.
- FAO (2008). The state of food and agriculture 2008. New York: Food and Agriculture Organization.
- Favilli, F., Pastorelli, R., Gori, A. (1993). Response of sugar beet to *Azospirillum* bacterization in field experiments. *Agric Mediterr*. 123: 281-285.
- Fedorov, A. S., Kosourov, S., Ghirardi, M. L., Seibert, M. (2005). Continuous H₂ photoproduction by *Chlamydomonas reinhardtii* using a novel two-stage, sulfate-limited chemostat system. *Appl Biochem Biotechnol*. 121-124: 403-412.
- Folch, J., Lees, M., Sloane Stanley G. H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem*. 226 (1): 497-509.

- Frackman, S., Kobs, G., Simpson, D., Storts, D. (1998). Betaine and DMSO: Enhancing agents for PCR . *Promega Notes*. 65: 27.
- Galhano, V., Figueiredo, D., Alves, A., Correia, A., Pereira, M., Gomes-Laranjo, J., Peixoto, F. (2011). Morphological, biochemical and molecular characterization of *Anabaena*, *Aphanizomenon* and *Nostoc* strains (Cyanobacteria, Nostocales) isolated from Portuguese freshwater habitats. *Hydrobiologia*. 663 (1): 187-203.
- Gavrilescu, M., Chisti, Y. (2005). Biotechnology—a sustainable alternative for chemical industry. *Biotechnol Adv*. 23: 471-499.
- Gekko, K., Timasheff, S. N. (1981). Thermodynamic and kinetic examination of protein stabilization by glycerol. *Biochemistry*. 20: 4677-4686.
- Ghirardi, M. L., Zhang, J. P., Lee, J. W., Flynn, T., Seibert, M., Greenbaum E. (2000). Microalgae: a green source of renewable H₂ . *Trends Biotechnol*. 18: 506-511.
- Ginzburg, B. Z. (1993). Liquid fuel (oil) from halophilic algae: a renewable source of non-polluting energy. *Renew Energy*. 3 (2-3): 249-252.
- Gomez, A., Rodrigues, M., Montañes, C., Dopazo, C., Fueyo, N. (2011). The technical potential of first-generation biofuels obtained from energy crops in Spain. *Biomass Bioenerg*. 35 (5): 2143-2155.
- Gouveia, L., Oliveira, A. C. (2009). Microalgae as a raw material for biofuels production. *J Ind Microbiol Biot*. 36 (2): 269-274.
- Govedarica, M., Milosevic, N., Jarak, M., Vojvodic Vukovic, M. (1993). Effectiveness of *Azospirillum lipoferum* strains in carrot. *Zemljiste Biljka*. 42: 121-125. (In Serbian)
- Govedarica, M., Milosevic, N., Jarak, M. (1994). Efficient application of associative nitrogen fixing bacteria in vegetable production. *Savrem Poljopr*. 42: 303-308. (In Serbian)
- Granite, E. J., Pennline, H. W. (2002). Photochemical removal of mercury from flue gas. *Proceedings of the 11th international conference on Coal Science*, San Francisco, CA.
- Grobbelaar, J. (2004). Mineral Nutrition. In: *Handbook of microalgal culture* (ed. Richmond A.), pp. 104-106. Blackwell Publishing Company, 2121 State Avenue, Ames, Iowa, USA.
- Han, D., Fan, Y., Hu, Z. (2009). An evaluation of four phylogenetic markers in *Nostoc*: Implications for cyanobacterial phylogenetic studies at the intrageneric level. *Curr Microbiol*. 58: 170-176.
- Healy, F. P. (1973). Inorganic nutrient uptake and deficiency in algae. *CRC Crit Rev Microbiol*. 3: 69.

Ho, S. H., Chen C. Y., Lee, D. J., Chang, J. S. (2011). Perspectives in microalgae CO₂ emission mitigation systems- A review. *Biotechnol Adv.* 29: 189-198.

[http:// www.ncbi.nlm.nih.gov/blast/](http://www.ncbi.nlm.nih.gov/blast/)

Hu, Q. (2004). Environmental effects on cell composition. In: *Handbook of microalgal culture* (ed. Richmond A.), p. 84. Blackwell Publishing Company, 2121 State Avenue, Ames, Iowa, USA.

Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., Darzins, A. (2008). Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *Plant J.* 54: 621-639.

Huertas, E., Montero, O., Lubian, L. (2000). Effects of dissolved inorganic carbon availability on growth, nutrient uptake and chlorophyll fluorescence of two species of marine microalgae. *Aquacult Eng.* 22: 181-197.

IDT. (2005). Breaking PCR: A systematic investigation of Intentional Violations of a basic Polymerase Chain Reaction Amplification control. Last accessed on 12/12/11. Available at <http://www.idtdna.com/pages/docs/technical-reports/breaking-pcr.pdf>

EIA. (2006). *World Energy Outlook 2006*. Paris: International Energy Agency.

Illman, A. M., Scragg, A. H., Shales, S. W. (2000). Increase in *Chlorella* strains calorific values when grown in low nitrogen medium. *Enzyme Microb Technol.* 27: 631-635.

Isik, O., Sarihan, E., Kusvuran, E., Gul, O., Erbatur, O. (1999). Comparison of the fatty acid composition of the freshwater fish larva *Tilapia zillii*, the rotifer *Brachionus calyciflorus*, and the microalgae *Scenedesmus abundans*, *Monoraphidium minutum* and *Chlorella vulgaris* in the algae-rotifer-fish larvae food chains. *Aquaculture.* 174 (3): 299-311.

Iverson, S., Lang, S., Cooper, M. (2001). Comparison of Bligh & Dyer and Folch methods for total lipid determination in a broad range of marine tissue. *Lipids.* 36 (11): 1283-1287.

Iwamoto, H. (2004). Industrial Production of Microalgal cell-mass and secondary products-Major Industrial Species. In: *Handbook of microalgal culture* (ed. Richmond A.), p.225. Blackwell Publishing Company, 2121 State Avenue, Ames, Iowa, USA.

Janda, M., Abbott, S. (2007). 16s rRNA Gene Sequencing for Bacterial Identification in the Diagnostic Laboratory: Pluses, Perils, and Pitfalls. *J Clin Microbiol.* 45 (9): 2761-2764.

Janson, S., Matveyev, A., Bergman, B. (1998). The presence and expression of hetR in the non-heterocystous cyanobacterium *Synploca* PCC 8002. *FEMS Microbiol Lett.* 168: 173-179.

- Jeffrey, S. W., Welschmeyer, N. A. (1997). Spectrophotometric and fluorometric equations in common use in oceanography. In: *Phytoplankton Pigments in Oceanography: Guidelines to Modern Methods*. (eds. Jeffrey, S. W., Mantoura, R. F. C., Wright, S. W.), pp. 597-621. UNESCO Publ., Paris, France.
- Kaczmarska, I., Ehrman, J., Bates, S., Green, D., Leger, C., Harris, J. (2005). Diversity and distribution of epibiotic bacteria on *Pseudonitzschia multiseries* (Bacillariophyceae) in culture, and comparison with those on diatoms in native seawater. *Harmful Algae*. 4 (4): 725-741.
- Kaledin, A. S., Sliusarenko, A. G., Gorodetskii, S. I. (1980). Isolation and properties of DNA polymerase from extreme thermophilic bacteria *Thermus aquaticus* YT-1. *Biokhimiia*. 45 (4): 644-651.
- Kang, J., Lee, M. S., Gorenstein, D. (2005). The enhancement of PCR amplification of a random sequence DNA library by DMSO and betaine: Application to in vitro combinatorial selection of aptamers. *Biochem Biophys Methods*. 64 (2): 147-151.
- Kapdan, I. K., Kargi, F. (2006). Bio-hydrogen production from waste materials. *Enzyme Microb Technol*. 38: 569-582.
- Kaplan, D., Richmond A. E., Dubinsky, Z., Aaronson, A. (1986). Algal Nutrition. In: *Handbook of microalgal mass culture* (ed. A. Richmond), pp. 147-198. CRC Press, Boca Raton, FL, USA.
- Khosla, V. (2009). Where will biofuels and biomass feedstocks come from? Available from: <http://www.khoslaventures.com/presentations/>. p. 31.
- Knothe, G. (2008). "Designer Biodiesel: Optimizing Fatty Ester Composition to Improve Fuel Properties". *Energ Fuel*. 22 (2): 1358-1364.
- Kolmodin L. A., Fenton Williams J. (1997). PCR: Basic Principles and Routine Practice. In: *PCR Cloning Protocols, From Molecular Cloning to Genetic Engineering*. (Ed. White B.), p. 9. Humana Press Inc., Totowa, New Jersey, USA.
- Komarek, J. (2006). Cyanobacterial Taxonomy: Current problems and prospects for the integration of traditional and molecular approaches. *Algae*. 21: 349-375.
- Lawyer, F. C., Stoffel, S., Saiki, R. K., Chang, S. Y., Landre, P. A., Abramson, R. D. (1993). High-level expression, purification and enzymatic characterization of full-length *Thermus aquaticus* DNA polymerase and a truncated form deficient in 5' to 3' exonuclease activity. *PCR Method and Appl*. 2: 275-287.
- Lee, K., Lee, C. (2001). Effect of light/dark cycles on wastewater treatments by microalgae. *Biotechnol Bioproc Eng*. 6 (3): 194-199.

- Lee, J. S., Lee, J. P. (2003). Review of advances in biological CO₂ mitigation technology. *Biotechnol Bioproc Eng.* 8: 354-359.
- Lin, Q., Lin, J. (2011). Effects of nitrogen source and concentration on biomass and oil production of a *Scenedesmus rubescens* like microalga. *Bioresour Technol.* 102 (2): 1615-1621.
- Liu, Z., Wang, G., Zhou, B. (2007). Effect of iron on growth and lipid accumulation in *Chlorella vulgaris*. *Bioresour Technol.* 99: 4717-4722.
- Longley, M. J., Bennett, S. E., Mosbaugh, D. W. (1990). Characterization of the 5' to 3' exonuclease associated with *Thermus aquaticus* DNA polymerase. *Nucleic Acids Res.* 18: 7317-7322.
- Lu, X. (2010). A perspective: Photosynthetic production of fatty acid-based biofuels in genetically engineered cyanobacteria. *Biotechnol Adv.* 6: 742-746.
- Lyamichev, V., Brow, M. A., Dahlberg, J. E. (1993). Structure-specific endonucleolytic cleavage of nucleic acids by eubacterial DNA polymerases. *Science.* 260: 778-783.
- Macedo, I. C., Seabra, J. E. A., Silva, J. E. A. R. (2008). Green house gases emissions in the production and use of ethanol from sugarcane in Brazil: The 2005/2006 averages and a prediction for 2020. *Biomass Bioenerg.* 32 (7): 582-595.
- MacIntyre, H. L., Kana, T. M., Anning, T., Geider, R. J. (2002). Photoacclimation of photosynthesis irradiance response curves and photosynthetic pigments in microalgae and cyanobacteria. *J Phycol.* 38 (1): 17-38.
- Mandal, S., Mallick, N. (2009). Microalga *Scenedesmus obliquus* as a potential source for biodiesel production. *Appl Microbiol Biot.* 84 (2): 281-291.
- Mann, D. G. (1984). Observations on copulation in *Navicula pupula* and *Amphora ovalis* in relation to the nature of diatom species. *Ann Bot.* 54: 429-438.
- Mann, D. G. (1989). The species concept in diatoms: Evidence for morphologically distinct, sympatric gamodemes in four epipelagic species. *Plant Syst Evol.* 164: 215-237.
- Mann, D. G. (1999). The species concept in diatoms. *Phycologia.* 38: 437-495
- Mann, D. G., Droop, S. J. M. (1996). Biodiversity, biogeography and conservation of diatoms. *Hydrobiologia.* 336: 19-32.
- Mann, D. G. Kociolek, J. P. (1990). The species concept in diatoms: Report on a workshop. In: *Proceedings of the 10th International Diatom Symposium* (Ed. Simola, H.).

- Masojidek, J., Koblizek, M., Torzillo, G. (2004). Photosynthesis in microalgae. In: Handbook of microalgal culture (Ed. Richmond A.) pp. 20-39. Blackwell Publishing Company, 2121 State Avenue, Ames, Iowa, USA.
- Mata, T. M., Martins, A. A., Caetano, N. S. (2010). Microalgae for biodiesel production and other applications: a review. *Renew Sust Energ Rev.* 14: 217-232.
- Meier, R. L. (1955). Biological Cycles in the Transformation of Solar Energy into Useful Fuels. In: *Solar Energy Research*. (Eds. Daniels, F., Duffie, A.), p. 179. Univ. Wisconsin Press, Madison, Wisconsin, USA.
- Melis, A. (2002). Green alga hydrogen production: progress, challenges and prospects. *Int J Hydrogen Energy.* 27: 1217-1228.
- Meyer, A., Todt, C., Mikkelsen, N., Lieb, B. (2010). Fast evolving 18S rRNA sequences from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into mollusk substitution rate heterogeneity. *BMC Evol Biol.* 10: 70 (1-12)
- Miao, X., Wu, Q. (2006). Biodiesel production from heterotrophic microalgal oil. *Bioresour Technol.* 97 (6): 841-846.
- Milne, T. A., Evans, R. J., Nagle, N. (1990). Catalytic conversion of microalgae and vegetable oils to premium gasoline, with shape selective zeolites. *Biomass.* 21 (3): 219-232.
- Minowa, T., Yokoyama, S. Y., Kishimoto, M., Okakurat, T. (1995). Oil production from algal cells of *Dunaliella tertiolecta* by direct thermochemical liquefaction. *Fuel.* 74 (12): 1735-1738.
- Mishra, A., Jha, B. (2009). Isolation and Characterization of extracellular polymeric substances from microalgae *Dunaliella salina* under salt stress. *Bioresour Technol.* 100 (13): 3382-3386.
- Nagai, M., Yoshida, A., Sato, N. (1998). Additive effects of bovine serum albumin, dithiothreitol and glycerol on PCR. *IUBMB Life.* 44 (1): 157-163.
- Natrah, F., Yosoff, F. M., Shariff, M., Abas, F., Mariana, N. S. (2008). Screening of Malaysian indigenous microalgae for antioxidant properties and nutritional value. *J Appl Phycol.* 19 (6): 711-718.
- Neenan, B., Feinberg, D., Hill, A., Mcintosh, R., Terry, K. (1986). Fuels from microalgae: Technology status, potential, and research requirements. *Solar Energy Research Institute Report*.
- Okon, Y., Labandera-Gonzalez, C. A. (1994). Agronomic applications of *Azospirillum*: an evaluation of 20 years of worldwide field inoculation. *Soil Biol Biochem.* 26: 1591-1601.

- Packer, M. (2009). Algal capture of carbon dioxide; biomass generation as a tool for greenhouse gas mitigation with reference to New Zealand energy strategy and policy. *Energy Policy*. 37 (9): 3428-3437.
- Patil, V., Källqvist T., Olsen, E., Vogt, E., Gislerod, H. R. (2007). Fatty acid composition of 12 microalgae for possible use in aquaculture feed. *Aquacult Int*. 15: 1-9.
- Perez, S. (2006). Culturing Algae. Last accessed on 12/12/11. Protocol available at: <http://people.oregonstate.edu/~weisv/Protocols/Symbiodinium/Algal%20cultures.pdf>
- Phukan, M. M., Chutia, R. S., Konwar, B. K., Kataki, R. (2011). Microalgae *Chlorella* a potential bio-energy feedstock. *Appl Energ*. 88 (10): 3307-3312.
- Pienkos, P., Darzins, A. (2009). The promise and challenges of microalgal-derived biofuels. *Biofuel Bioprod Bior*. 3 (4): 431-440.
- Pohl, P. (1982). Lipids and fatty acids of microalgae. In: *CRC handbook of biosolar resources*. (Ed. Zaborsky, O. R.), pp. 383-404. 1. CRC Press Inc., Boca Raton, FL, USA.
- Pomp, D., Medrano, J. F. (1991). Organic solvents as facilitators of polymerase chain reaction. *BioTechniques*. 10: 58-59.
- Pratoomyot, J., Srivilas, P., Noiraksar, T. (2005). Fatty acids composition of 10 microalgal species. *Songklanakarin J Sci Technol*. 27 (6): 1179-1187.
- Raghavan, G., Haridevi, C. K., Gopinathan, C. P. (2008). Growth and proximate composition of the *Chaetoceros calcitrans* f. *pumilus* under different temperature, salinity and carbon dioxide levels. *Aquac Res*. 39: 1053-1058.
- Rana, R., Spada, V. (2007). Biodiesel production from ocean biomass. In: *Proceedings of the 15th European conference and exhibition*, Berlin.
- Rao A. R., Dayananda C., Sarada R., Shamala T., Ravishankar G. (2006). Effect of salinity on growth of green alga *Botryococcus braunii* and its constituents. *Bioresour Technol*. 98 (3): 560-564.
- Rasoul-Amini, S., Ghasemi, Y., Hossein, M, Mohagheghzadeh, A. (2009). PCR amplification of 18S rRNA, single cell protein production and fatty acid evaluation of some naturally isolated microalgae. *Food Chem*. 116 (1): 129-136.
- Renaud, S. M., Thunh L. V., Lambrinidis G., Parry D. L. (2002). Effect of temperature on growth, chemical composition and fatty acid composition of tropical Australian batch cultures. *Aquaculture*. 211: 195-214.

Richmond, A. (1986). Cell response to environmental factors. In: *Handbook of Microalgal Mass Culture*. (Ed. Richmond, A.), pp. 69–99 . CRC Press. Boca Raton, Florida, USA.

Robertson, E. P. (2007). Analysis of CO₂ separation from flue gas, pipeline transportation, and sequestration in coal. Idaho National Laboratory external report. No. INL/EXT-08-13816, Idaho Falls, Idaho.

Rodolfi, L., Bassi, N., Padovani, G., Bonini, G., Zitelli, G. C., Biondi, N., Tredici, M. R. (2007). Lipid production from microalgae: strain selection, induction of lipid synthesis and outdoor cultivation in pilot photobioreactors. In: Proceedings of the 15th European conference and exhibition, Berlin, Germany.

Roessler, P. (1987). Lipid accumulation in silicon deficient diatoms. In: *The Metabolism, Structure and Function of Plant Lipids*. (Ed. Stumpf, P. K., Mudd, J. B., and Ness, W. D.), pp-649-651. Plenum publishing Co., New York, USA.

Roessler, P. G., Brown, L. M., Dunahay, T. G., Heacox, D. A., Jarvis, E. E., Schneider, J. C. (1994). Genetic-engineering approaches for enhanced production of biodiesel fuel from microalgae. *ACS Symp Ser.* 566: 255-270.

Roux, K. H. (1995). Optimization and troubleshooting in PCR. In: *PCR Primer: A Laboratory Manual* (Eds. Dieffenbach, C. W., Dveksler, G. S.), pp. 53-62 . Cold Spring Harbor Press, Plainview, NY. USA.

Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A., Arnheim, N. (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science.* 230: 1350-1354.

Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science.* 239: 487-491.

Sanchez, S. M. P., Voltolina D. (2006). The growth rate, biomass production and composition of *Chaetoceros* sp. grown with different light sources. *Aquacult Eng.* 35 (2): 161-165.

Sato, N. (1989). Modulation of lipid and fatty acid content by carbon dioxide in *Chlamydomonas reinhardtii*. *Plant Sci.* 61: 17-21.

Sawayama, S., Inoue, S., Dote, Y., Yokoyama, S. Y. (1995). CO₂ fixation and oil production through microalga. *Energy Convers Manag.* 36: 729-731.

Schleifer, K. H. (2009). Classification of Bacteria and Archaea: past, present and future. *Syst Appl Microbiol.* 32: 533-542.

- Scott, S. A., Davey, M. P., Dennis, J. S., Horst, I., Howe, C. J., Lea-Smith, D. J., Smith A. G. (2010). Biodiesel from algae: challenges and prospects. *Curr Opin Biotech.* 21: 277-286.
- Schubert, L. E. (1988). The use of *Spirulina* and *Chlorella* as food resource for animals and humans. In: *Progressing physiological research.* (Eds. Round, F.E., Chapman, D.J.), p. 237. Biopress Ltd, Bristol, U.K.
- Schubert, C. (2006). Can biofuels finally take center stage? *Nat Biotechnol.* 24 (7): 777-784.
- Scragg, A. H., Morrison J., Shale, S. W. (2003). The use of a fuel containing *Chlorella vulgaris* in a diesel engine. *Enzyme Microb Technol.* 33: 884-889.
- Sheehan, J., Dunahay, T., Benemann, J., Roessler, P. (1998). A look back at the U.S. Department of Energy's Aquatic Species Program. National Renewable Energy Laboratory.
- Sidhu, M. K., Liao, M. J., Rashidbaigi, A. (1996). Dimethyl sulfoxide improves RNA amplification. *Biotechniques.* 21: 44-47.
- Singh, J., Gu, S. (2010). Commercialization potential of microalgae for biofuels production. *Renew Sust Energ Rev.* 14: 2596-2610.
- Solovchenko, A. E., Khozin-Goldberg, I., Didi-Cohen, S., Cohen, Z., Merzlyak, M. N. (2008). Effects of light intensity and nitrogen starvation on growth, total fatty acids and arachidonic acid in the green microalga *Parietochloris incise*. *J Appl Phycol.* 20: 245-251.
- Spolaore, P., Joannis-Casan, C., Duran, E., Isambert, A. (2006). Commercial applications of microalgae. *J Biosci Bioeng.* 101: 87-96.
- Status of ratification of Kyoto Protocol. Retrieved November 20, 2011. Available at: http://unfccc.int/kyoto_protocol/status_of_ratification/items/2613.php
- Stirling, D. (2003) Technical notes for sequencing difficult templates. In: *PCR Protocols* (Ed. Bartlett J.M.S., Stirling D.), pg. 401. Humana Press, Totowa, New Jersey.
- Su C. H., Fu C. C., Chang Y. C., Nair G. R., Ye J. L., Chu I. M., Wu W. T. (2007). Simultaneous estimation of Chlorophyll a and lipid contents in microalgae by three-color analysis. *Biotechnol Bioeng.* 99 (4): 1034-1039.
- Sun, Y., Hegamyer, G., Colburn N. H. (1993). PCR directed sequencing of a GC-rich region by inclusion of 10% DMSO: application to C-jun. *BioTechniques.* 15: 372-374.
- Taguchi, G. (1986). *Introduction to quality engineering: Designing quality in to products and processes.* Asian Productivity Organization, Tokyo, Japan.

- Tang, D., Han, W., Li, P., Miao, X., Zhong, J. (2011). CO₂ biofixation and fatty acid composition of *Scenedesmus obliquus* and *Chlorella pyrenoidosa* in response to different CO₂ levels. *Bioresour Technol.* 102 (3): 3071-3076.
- Tomaselli, L. (2004). The Microalgal cell. In: Handbook of microalgal culture (Ed. Richmond A.) pp. 11-17. Blackwell Publishing Company, 2121 State Avenue, Ames, Iowa, USA.
- Tornabene, T. G., Holzer, G., Lien, S., Burris, N. (1983). Lipid composition of the nitrogen starved green alga *Neochloris oleabundans*. *Enzyme Microb Technol.* 5: 435-440.
- Tsuzuki, M., Ohnuma, E., Sato, N., Takaku, T., Kawaguchi, A. (1990). Effects of CO₂ concentration during growth on fatty acid composition in microalgae. *Plant Physiol.* 93: 851-856.
- Tzovenis, I., De Pauw N., Sorgeloos, P. (1997). Effect of different light regimes on the docosahexaenoic acid (DHA) content of *Isochrysis* aff. *galbana* (clone- TISO). *Aquacult Int.* 5 (6): 489-507.
- Vazquez-Duhalt, R., Arredondo-Vega, B. O. (1991). Haloadaptation of the green alga *Botryococcus braunii* ('A' Race). *Phytochemistry.* 30 (9): 2919-2925.
- Vickery, M., Smith, A., De Paola, A., Jones, D., Steffan, R., Beja, A. (1998). Optimization of the arbitrarily-primed polymerase chain reaction (AP-PCR) for intra-species differentiation of *Vibrio vulnificus*. *J Microbiol Meth.* 33 (2): 181-189.
- Vieira Costa, J. A., Greque de Morais, M. (2011). The role of biochemical engineering in the production of biofuels from microalgae. *Bioresources.* 102 (1): 2-9.
- Vonshak, A., Maske, H. (1982). Algae: growth techniques and biomass production. In: Techniques in bioproductivity and photosynthesis. (Ed. Coombs, J., Hall, D. O.), p. 66. Pergamon Press Inc., Maxwell House, Fairview Park, Elmsford, New York, U.S.A.
- Wang, C., Altieri, F., Ferraro, A., Giartosio, A., Turano, C. (1993). The effect of polyols on the stability of duplex DNA. *Physiol Chem Phys Med NMR.* 25 (4): 273-280.
- Wang, B., Li, Y., Wu, N., Lan, C. (2008). CO₂ bio-mitigation using microalgae. *Appl Microbiol Biot.* 79 (5): 707-718.
- Wang, X., Hao, C., Zhang, F., Feng, C., Yang, Y. (2011). Inhibition of the growth of two blue-green algae species (*Microcystis aeruginosa* and *Anabaena spiroides*) by acidification treatments using carbon dioxide. *Bioresour Technol.* 102 (10): 5742-5748.
- Wiltshire, K. H., Lampert, W. (2000). Extraction of pigments and fatty acids from the green alga *Scenedesmus obliquus*. *Aquat Ecol.* 34: 119-126.

Worldwatch. (2007). Biofuels for transportation: global potential and implications for sustainable agriculture and energy in the 21st century. Washington, DC: Worldwatch Institute.

Wuebbles, D. J., Atul K. J. (2001). Concerns about climate and the role of fossil fuel use. Fuel Process Technol. 71: 99-119.

Xiong, W., Li, X., Xiang, J., Wu, Q. (2008). High density fermentation of microalga *Chlorella protothecoides* in bioreactor for microbiodiesel production. Appl Microbiol Biotechnol. 78: 29-36.

Xu, H., Miao, X., Wu, Q. (2006). High quality biodiesel production from a microalga *Chlorella protothecoides* by heterotrophic growth in fermenters. J Biotechnol. (126): 499-507.

Xu, Z., Harvey, K., Pavlina T., Dutot G., Zaloga G., Siddiqui, R. (2010). An improved method for determining medium- and long-chain FAMES using gas chromatography. Lipids. 45: 199-208.

Yoo, C., Jun, S. Y., Lee J. Y., Ahn, C. Y., Oh, H. M. (2010). Selection of microalgae for lipid production under high levels of carbon dioxide. Bioresour Technol. 101: S71-S74.

Zehr, J. P., McReynolds, L. A. (1989). Use of degenerate oligonucleotides for amplification of the *nifH* gene from the marine cyanobacterium *Trichodesmium thiebautii*. Appl Environ Microb. 55: 2522-2526.

Zhu, C. J., Lee Y. K., Chao, T. M. (1997). Effects of temperature and growth phase on lipid and biochemical composition of *Isochrysis galbana* TK1. J Appl Phycol. 9 (5): 451-457.

APPENDIX 1. F/2 MEDIA COMPOSITION

Component	Stock Solution	Quantity per liter of medium	Molar Concentration in Final Medium
NaNO ₃	75 g/L dH ₂ O	1 mL	8.82 x 10 ⁻⁴ M
NaH ₂ PO ₄ H ₂ O	5 g/L dH ₂ O	1 mL	3.62 x 10 ⁻⁵ M
Na ₂ SiO ₃ 9H ₂ O	30 g/L dH ₂ O	1 mL	1.06 x 10 ⁻⁴ M
trace metal solution	(recipe below)	1 mL	---
vitamin solution	(recipe below)	0.5 mL	---

F/2 Vitamin Solution

Component	Primary Stock Solution	Quantity per liter of medium	Molar Concentration in Final Medium
thiamine HCl (vit. B ₁)	---	200 mg	2.96 x 10 ⁻⁷ M
biotin (vit. H)	0.1 g/L dH ₂ O	10 mL	2.05 x 10 ⁻⁹ M
cyanocobalamin (vit. B ₁₂)	1.0 g/L dH ₂ O	1 mL	3.69 x 10 ⁻¹⁰ M

Trace metals solution

Component	Primary Stock Solution	Quantity per liter of medium	Molar Concentration in Final Medium
FeCl ₃ 6H ₂ O	---	3.15 g	1.17 x 10 ⁻⁵ M
Na ₂ EDTA 2H ₂ O	---	4.36 g	1.17 x 10 ⁻⁵ M
CuSO ₄ 5H ₂ O	9.8 g/L dH ₂ O	1 mL	3.93 x 10 ⁻⁸ M
Na ₂ MoO ₄ 2H ₂ O	6.3 g/L dH ₂ O	1 mL	2.60 x 10 ⁻⁸ M
ZnSO ₄ 7H ₂ O	22.0 g/L dH ₂ O	1 mL	7.65 x 10 ⁻⁸ M
CoCl ₂ 6H ₂ O	10.0 g/L dH ₂ O	1 mL	4.20 x 10 ⁻⁸ M
MnCl ₂ 4H ₂ O	180.0 g/L dH ₂ O	1 mL	9.10 x 10 ⁻⁷ M

APPENDIX 2. BLUE-GREEN 11 MEDIA COMPOSITION

Component	Quantity per liter of medium	Stock Solution Concentration	Final Concentration
NaNO ₃	10 mL	30 g/200 mL dH ₂ O	17.6 mM
K ₂ HPO ₄	10 mL	0.8 g/200 mL dH ₂ O	0.22 mM
MgSO ₄ ·7H ₂ O	10 mL	1.5 g/200 mL dH ₂ O	0.03 mM
CaCl ₂ ·2H ₂ O	10 mL	0.72 g/200 mL dH ₂ O	0.2 mM
Citric Acid·H ₂ O	10 mL	0.12 g/200 mL dH ₂ O	0.03 mM
Ammonium Ferric Citrate	10 mL	0.12 g/200 mL dH ₂ O	0.02 mM
Na ₂ EDTA·2H ₂ O	10 mL	0.02 g/200 mL dH ₂ O	0.002 mM
Na ₂ CO ₃	10 mL	0.4 g/200 mL dH ₂ O	0.18 mM
BG-11 Trace Metals Solution	1 mL		

BG-11 Trace Metals Solution

Component	Quantity per liter of medium	Stock Solution Concentration	Final Concentration
H ₃ BO ₃	2.86 g	----	46 μM
MnCl ₂ ·4H ₂ O	1.81 g	----	9 μM
ZnSO ₄ ·7H ₂ O	0.22 g	----	0.77 μM
Na ₂ MoO ₄ ·2H ₂ O	0.39 g	----	1.6 μM
CuSO ₄ ·5H ₂ O	0.079 g	----	0.3 μM
Co(NO ₃) ₂ ·6H ₂ O	49.4 mg	----	0.17 μM

APPENDIX 3. BOLD'S BASAL MEDIA COMPOSITION

Component	400 mL Stock Solution	1 Liter Stock Solution	quantity per liter of medium	Molar Concentration in Final Medium ^f
<u>Major Stock Solutions</u>				
NaNO ₃	10 g L ⁻¹ dH ₂ O	25.00 g L ⁻¹ dH ₂ O	10 mL	2.94 x 10 ⁻³ M
CaCl ₂ • 2H ₂ O	1 g L ⁻¹ dH ₂ O	2.50 g L ⁻¹ dH ₂ O	10 mL	1.70 x 10 ⁻⁴ M
MgSO ₄ • 7H ₂ O	3 g L ⁻¹ dH ₂ O	7.50 g L ⁻¹ dH ₂ O	10 mL	3.04 x 10 ⁻⁴ M
K ₂ HPO ₄	3 g L ⁻¹ dH ₂ O	7.50 g L ⁻¹ dH ₂ O	10 mL	4.31 x 10 ⁻⁴ M
KH ₂ PO ₄	7 g L ⁻¹ dH ₂ O	17.50 g L ⁻¹ dH ₂ O	10 mL	1.29 x 10 ⁻³ M
NaCl	1 g L ⁻¹ dH ₂ O	2.50 g L ⁻¹ dH ₂ O	10 mL	4.28 x 10 ⁻⁴ M
<u>Alkaline EDTA Stock Solution</u>			1 ml per liter of medium	
EDTA anhydrous		50 g L ⁻¹ dH ₂ O		4.28 x 10 ⁻⁴ M
KOH		31 g L ⁻¹ dH ₂ O		1.38 x 10 ⁻³ M

<u>Acidified Iron Stock Solution</u>			1 mL per liter of medium	Molar Concentration in Final Medium ^f
FeSO ₄ • 7H ₂ O		4.98 g L ⁻¹ dH ₂ O		4.48 x 10 ⁻⁵ M
H ₂ SO ₄		1.0 mL		
<u>Boron Stock Solution</u>			1 mL per liter of medium	
H ₃ BO ₃		11.42 g L ⁻¹ dH ₂ O		4.62 x 10 ⁻⁴ M
<u>Trace Metal Stock Solution</u>			1 mL per liter of medium	
ZnSO ₄ • 7H ₂ O		8.82 g L ⁻¹ dH ₂ O		7.67 x 10 ⁻⁵ M
MnCl ₂ • 4H ₂ O		1.44 g L ⁻¹ dH ₂ O		1.82 x 10 ⁻⁵ M
MoO ₃		0.71 g L ⁻¹ dH ₂ O		1.23 x 10 ⁻⁵ M
CuSO ₄ • 5H ₂ O		1.57 g L ⁻¹ dH ₂ O		1.57 x 10 ⁻⁵ M
Co(NO ₃) ₂ • 6H ₂ O		0.49 g L ⁻¹ dH ₂ O		4.21 x 10 ⁻⁶ M

APPENDIX 4. SAMPLES DILUTIONS FOR HEMACYTOMETER


Batch A

<u>Sample</u>	<u>Dilution factor</u>	<u>Method</u>
N3	1:5	B
FC2	1:1	A
FC1	1:3	B
D2	1:3	B
<i>Chlorella v.UTEX 259</i>	1:2	B


Batch B

<u>Sample</u>	<u>Dilution factor</u>	<u>Method</u>
N3	1:5	B
FC2	1:1	A
FC1	1:5	B
D2	1:3	B
<i>Chlorella v.UTEX 259</i>	1:3	B

APPENDIX 5. COUNTING PROCEDURES FOR HEMACYTOMETER READINGS



Method A (4 squares)



Method B (5 squares)

APPENDIX 6. FREEZE-DRYER PROGRAM

Freeze

Step #	1	2
T° (°C)	-40	-40
Minutes	0	240

Final Freeze

Final Freeze(°C)	-44
Extra Time	0
Prim Vac (mTorr)	200

Primary Dry

Step #	1	2	3	4	5	6
T° (°C)	-20	-20	0	0	20	20
Minutes	25	120	0	120	0	360
Vacuum (mTorr)	100	100	100	100	100	100

T° (°C)	0
Minutes	0
Vacuum (mTorr)	200
Final Setpoint (° C)	4

VITA

Reynaldo Moreno Uclés, son of Reynaldo Moreno Regalado and Carmen S. Uclés de Moreno, was born in San Pedro Sula, Honduras, in 1978. He graduated from Academia Americana High School in 1995. He attended Universidad de San Pedro Sula from 1997 to 2002, where he received a Bachelor of Science in Agricultural Engineering and Business Administration. In August 2005, he received a scholarship from Louisiana State University, Baton Rouge, to study music, and in 2008 he earned his Bachelor of Music. In August 2009, he began his Master's program in the Department of Food Science at Louisiana State University, Baton Rouge. In May 2012, he will receive his degree of master of science in food science.