



Inorganic arsenic in rice-based products for infants and young children



Antonio J. Signes-Pastor^{*}, Manus Carey, Andrew A. Meharg

Institute for Global Food Security, Queen's University Belfast, David Keir Building, Malone Road, Belfast BT9 5BN, Northern Ireland, United Kingdom

ARTICLE INFO

Article history:

Received 18 September 2014

Received in revised form 5 November 2014

Accepted 13 November 2014

Available online 24 November 2014

Keywords:

Arsenic speciation

Baby rice

Rice cereals

Rice crackers

Food regulations

ABSTRACT

Inorganic arsenic (As_i) is a chronic, non-threshold carcinogen. Rice and rice-based products can be the major source of As_i for many subpopulations. Baby rice, rice cereals and rice crackers are widely used to feed infants and young children. The As_i concentration in rice-based products may pose a health risk for infants and young children. As_i concentration was determined in rice-based products produced in the European Union and risk assessment associated with the consumption of these products by infants and young children, and compared to an identical US FDA survey. There are currently no European Union or United States of America regulations applicable to As_i in food. However, this study suggests that the samples evaluated may introduce significant concentration of As_i into infants' and young children's diets. Thus, there is an urgent need for regulatory limits on As_i in food, especially for baby rice-based products.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

1. Introduction

Arsenic (As) is ubiquitous in the environment and occurs in different forms. As_i is a non-threshold human carcinogen (IARC, 2004). Other than cancer, human exposure to As has been associated to diverse health problems (WHO, 2011a), which may be exacerbated with early-life exposure (Smith et al., 2006; Vahter, 2009). The main sources of human exposure to As are water and food (WHO, 2010). As_i in water is tightly regulated (Council Directive, 1998; WHO, 2011b), however, there is no European Union (EU) or United States of America (USA) standard for As_i in food products despite the fact that food sources dominate exposure, especially rice and rice-based products (EFSA, 2009; US FDA, 2014). Indeed, it has been demonstrated that consumption of rice and/or rice-based products increases the occurrence of As species in human urine (Cascio, Raab, Jenkins, Felman, & Meharg, 2011; Gilbert-Diamond et al., 2011; Wei, Zhu, & Hguyen, 2014). Furthermore, a cohort study in West Bengal, India, has reported elevated genotoxic effects, as measured by micronuclei in urothelial cells, associated with the staple consumption of cooked rice with $>200 \mu\text{g As/kg}$ (Banerjee et al., 2013). Recently, the JECFA proposed a maximum level of 0.2 mg/kg of As_i in polished rice (JECFA, 2014). The European Food Safety Authority (EFSA) has reviewed the diet of the European Union population and has recommend that dietary exposure to As_i should be reduced. Cereal and cereal-based products have been identified as contributors to daily As_i exposure in the general European population and young

children (<3 years of age) have been categorised as the most exposed to As_i (EFSA, 2009). The review was mainly based on data reported as total As and a number of assumptions to estimate the As_i exposure were made. It was highlighted that more speciation data for different food commodities are required to support a comprehensive dietary exposure assessment and to redefine risk assessment of As_i , especially to high exposure risk subpopulations. The JECFA also reviewed and stated that dietary exposure to As_i should be reduced. In addition, the Provisional Tolerable Weekly Intake (PTWI) was withdrawn since cancer of the lung and urinary bladder in addition to skin and a range of adverse effects were reported at As_i exposure lower than those reviewed at the time the PTWI was established (JECFA, 2010).

Rice accumulates significantly higher levels of As_i from soil and water than other crops due to anaerobic paddy soil culture, which renders As_i highly available for plant uptake, leading to ~ 10 -fold higher concentrations in grain compared to aerobically grown grains such as wheat or barley (Meharg & Zhao, 2012; William et al., 2007; Xu, McGrath, Meharg, & Zhao, 2008). Levels of As_i in rice are also of concern due to the fact that there is a high gut bio-availability of rice As_i (Juhasz et al., 2006; Signes-Pastor, Al-Rmalli, Jenkins, Carbonell-Barrachina, & Haris Parvez, 2012). Rice-based products also have high levels of As_i and show a positive correlation between rice content and their As concentration (Burlo, Ramirez-Gandolfo, Signes-Pastor, Haris, & Carbonell-Barrachina, 2012; Carbonell-Barrachina et al., 2012). Rice-based products are widely used during weaning and to feed young children due to its availability, bland taste, nutritional value and relatively low allergic potential (Da Sacco, Baldassarre, & Massotti, 2013; Meharg et al., 2008; Mennella, Ziegler, Briefel, & Novak, 2006).

^{*} Corresponding author.

E-mail address: a.signes-pastor@qub.ac.uk (A.J. Signes-Pastor).

Consumption of rice and rice-based foods is particularly high for infants and young children affected by celiac disease, which is a common condition that affects 1% of children in the EU and USA, while their only viable treatment is a gluten-free diet (Da Sacco et al., 2013; Munera-Picazo, Burló, & Carbonell-Barrachina, 2014; Newton & Singer, 2012). Furthermore, the fact that infants and young children have higher food consumption rates on body weight basis than adults, *circa* 3-fold, increases their exposure to As_i with respect to adults for any given item of food, which exacerbates toxicological issues for this subpopulation (EFSA, 2009; Meharg & Zhao, 2012).

Information on As speciation data of rice-based products consumed by infants and young children is needed to define risk assessment of As_i for one of the most vulnerable subpopulations, and to set regulations for As_i content in food to protect them. In this study, therefore, As speciation was measured in 29 commercial baby rice, 53 commercial rice cereals and 97 commercial rice crackers from the EU market, and compared to 85 commercial baby rice, 105 rice cereals and 199 rice crackers included in the US FDA survey on As_i in rice and rice-based products (US FDA, 2014), and the findings put in context of exposure risks to infants and young children.

2. Materials and methods

Samples of baby rice ($n=29$), rice cereals ($n=53$) and rice crackers ($n=97$) belonging to 21 different and most popular commercial brands or manufacturers were purchased from 36 food shops (15 local shops and 21 big supermarkets) in the United Kingdom (UK). Duplicate samples of the same product and brand were always purchased from different stores. The rice-based products sampled showed use by date between February 2014 and March 2016.

2.1. Sample preparation for As speciation

All samples of rice-based products were freeze-dried, and then powdered using a Retch PM100 rotary ball-mill using a zirconium oxide lined vessel and zirconium oxide grinding balls. The powdered samples were weighed accurately to a weight of 0.1 g into 50 ml polypropylene centrifuge tubes to which 10 ml of 1% conc. Aristar nitric acid was added and allowed to sit overnight. Batches of 30 samples approximately were prepared which also include a blank and rice CRM (NIST 1568b Rice flour) which has the As species As_i, dimethylarsinic acid (DMA) and monomethylarsonic acid (MMA) concentrations certified, then microwave digested in an CEM MARS 6 instrument for 30 min. at 95 °C using a 3 stage slow heating program: to 55 °C in 5 min. held for 10 min., to 75 °C in 5 min., held for 10 min. to 95 °C in 5 min., held for 30 min. A 1 ml aliquot was transferred to a 2 ml polypropylene vial and 10 µl of analytical grade hydrogen peroxide was added to convert any arsenite to arsenate to facilitate subsequent chromatographic detection.

2.2. Chemical analysis

To speciate As in rice-based products the diluted 1% nitric acid digested sample solutions were run on a Thermo Scientific IC5000 Ion Chromatography (IC) system, with an Thermo AS7, 2 × 250 mm column (and a Thermo AG7, 2 × 50 mm guard column) and a gradient mobile phase (A: 20 mM Ammonium Carbonate, B 200 mM Ammonium Carbonate – Starting at 100% A, changing to 100% B, in a linear gradient over 15 min.) interfaced with a Thermo ICAP Q ICP-MS that monitored m/z^+ 75, using He gas in collision cell mode. The resulting chromatogram was

compared with that for authentic standards; DMA, As_i, MMA, tetratmethyl arsonium (TETRA) and arsenobetaine (AB). As present under each chromatographic peak was calibrated using a DMA concentration series.

2.3. Statistical analyses

Data were subjected to general lineal model (GLM) and the Duncan multiple range test to determine significant differences among samples. The statistical analyses were performed using IBM SPSS Statistic version 21.0.

3. Results and discussion

The analysis of the As speciated CRM gave excellent recovery results (mean ± SE), based on $n=15$, with $105 \pm 3\%$ recovery for DMA, $93.4 \pm 5\%$ for MMA and $94.5 \pm 2\%$ recovery for As_i. The CRM had a certified concentration of 0.182, 0.012 and 0.092 mg/kg As for DMA, MMA and As_i, respectively. The limits of detection (LOD) for both DMA and As_i (calculated from a DMA calibration) was 0.003 mg/kg rice d.wt., $n=6$. All samples presented were above LOD for As_i and only a few samples were below LOD for DMA and MMA, and in this case 1/2 LOD was used in statistical analysis of the data.

3.1. Baby rice

The summation of As species concentration (Σ As) in 29 baby rice samples belonging to 5 different commercial brands ranged from 0.063 to 0.334 mg/kg (Table 1). The As_i percentage ranged from 51.4% to 84.6% of the Σ As species concentration. The baby rice brands M001, M003 and M005 had a significantly higher percentage of As_i (71.6%, 73.4% and 79.3%, respectively) than M002 and M004 (66.3% and 53.5%, respectively; $p < 0.001$). The As_i concentration ranged from 0.056 to 0.268 mg/kg. This shows that 14% of the baby rice samples evaluated would be above the JECFA proposed As_i maximum level for rice (0.200 mg/kg). The baby rice brand M005 had the highest As_i concentration (median of 0.190 mg/kg; $p < 0.001$). DMA ranged from 0.030 to 0.123 mg/kg (Table 1). The same trend of As speciation has been previously described in rice (Meharg et al., 2009; Torres-Escribano, Leal, Vélez, & Montoro, 2008). Baby rice samples labelled as produced under organic standard showed higher levels of As_i than non-organic samples (Fig. 1), which was associated with the inclusion of whole grain rice. In fact, organic baby rice samples including whole grain contain a higher concentration and percentage of As_i (median of 0.190 mg/kg and 79.3%) than those manufactured with milled rice (median of 0.121 mg/kg and 55.1%; $p < 0.001$) (Table 2). This is in agreement with previous studies that show that brown rice contains higher proportion of As_i than white rice, which is mainly concentrated at the surface of the whole grain, in the region corresponding to the pericarp and aleurone layer (Choi et al., 2014; Meharg et al., 2008). This is of particular concern as organic products are usually associated with a healthier and more nutritious option that is increasing production of organic baby food and may exacerbate As_i exposure and health risk for infants and young children (Da Sacco et al., 2013).

In September 2013 the US Food and Drug Administration (US FDA) released analytical results of As_i in approximately 1100 samples of rice and rice products from the US market, which were in addition to approximately 200 samples of rice and rice products initially tested, the results of which were released in September 2012 (USFDA, 2013). These results include 85 samples of baby rice (product subcategory: infant cereals and toddler cereals). The median and range of As_i in baby rice from the USA market is 0.114

Table 1
Concentration (mg/kg d.wt.) of As species in rice-based products (baby rice, rice cereals and rice crackers) ordered by food category and manufacturer.

Food category	Manufacture	n	As _i (mg/kg d.wt.)	DMA (mg/kg d.wt.)	MMA (mg/kg d.wt.)	Σ As species (mg/kg d.wt.)	As _i (%)
Baby rice	M001	4	0.093 ^{bc} (0.088–0.097) ^A	0.037 ^c (0.024–0.041)	0.001 (0.001–0.001)	0.131 ^b (0.112–0.137)	71.6 ^a (70.4–78.5)
	M002	7	0.066 ^c (0.056–0.092)	0.035 ^c (0.030–0.061)	0.001 (0.001–0.003)	0.099 ^b (0.063–0.090)	66.3 ^b (59.9–69.8)
	M003	4	0.118 ^{bc} (0.062–0.129)	0.041 ^c (0.023–0.046)	0.001 (0.001–0.001)	0.160 ^b (0.085–0.175)	73.4 ^a (73.2–74.3)
	M004	5	0.131 ^b (0.122–0.142)	0.112 ^a (0.107–0.120)	0.001 (0.001–0.003)	0.247 ^a (0.230–0.263)	53.5 ^c (51.4–55.1)
	M005	9	0.190 ^a (0.117–0.268)	0.057 ^b (0.038–0.123)	0.001 (0.001–0.004)	0.264 ^a (0.181–0.334)	79.3 ^a (57.2–84.6)
p-Value			<0.001	<0.001	0.342	<0.001	<0.001
Rice cereals	M006	5	0.068 ^c (0.066–0.123)	0.028 ^{ab} (0.024–0.037)	0.001 (0.001–0.001)	0.097 ^{bc} (0.920–0.160)	74.2 ^{ab} (70.6–77.8)
	M007	3	0.162 ^b (0.104–0.167)	0.037 ^{ab} (0.016–0.052)	0.001 (0.001–0.001)	0.204 ^b (0.122–0.214)	81.5 ^a (75.7–85.2)
	M008	6	0.074 ^c (0.044–0.097)	0.022 ^b (0.005–0.035)	0.001 (0.001–0.001)	0.098 ^c (0.049–0.132)	76.3 ^a (72.3–89.6)
	M009	2	0.234 ^a (0.146–0.323)	0.047 ^a (0.026–0.069)	0.002 (0.001–0.003)	0.284 ^a (0.173–0.396)	83.1 ^a (81.7–84.5)
	M010	30	0.075 ^c (0.008–0.188)	0.045 ^a (0.013–0.082)	0.001 (0.001–0.004)	0.120 ^{bc} (0.057–0.249)	62.7 ^b (14.2–81.6)
p-Value			<0.001	0.002	0.422	<0.001	0.001
Rice crackers	M006	2	0.132 ^{bc} (0.128–0.136)	0.021 (0.021–0.022)	0.001 (0.001–0.001)	0.154 ^{bc} (0.150–0.158)	85.7 (85.1–86.3)
	M012	3	0.140 ^b (0.099–0.196)	0.042 (0.035–0.049)	0.001 (0.001–0.003)	0.175 ^{ab} (0.141–0.249)	78.7 (70.0–80.2)
	M001	6	0.127 ^b (0.110–0.162)	0.023 (0.014–0.029)	0.001 (0.001–0.001)	0.150 ^{bc} (0.124–0.190)	85.2 (83.3–88.6)
	M013	4	0.099 ^{bcd} (0.032–0.160)	0.016 (0.006–0.047)	0.001 (0.001–0.001)	0.115 ^{bcd} (0.039–0.207)	83.2 (77.3–89.4)
	M014	9	0.125 ^{bc} (0.103–0.187)	0.028 (0.023–0.032)	0.001 (0.001–0.002)	0.157 ^{bc} (0.131–0.220)	80.7 (72.1–85.0)
	M015	3	0.095 ^{bcd} (0.082–0.106)	0.035 (0.026–0.038)	0.001 (0.001–0.001)	0.133 ^{bcd} (0.107–0.144)	73.6 (71.6–75.9)
	M009	19	0.101 ^{bcd} (0.035–0.184)	0.026 (0.011–0.172)	0.001 (0.001–0.006)	0.130 ^{bc} (0.046–0.328)	79.4 (44.8–85.1)
	M016	3	0.070 ^{cde} (0.063–0.088)	0.017 (0.015–0.020)	0.001 (0.001–0.001)	0.085 ^{cd} (0.081–0.109)	81.2 (78.2–82.1)
	M017	4	0.067 ^{bcd} (0.057–0.175)	0.020 (0.013–0.037)	0.001 (0.001–0.001)	0.088 ^{bcd} (0.076–0.215)	78.4 (71.1–82.8)
	M005	25	0.131 ^{bc} (0.068–0.188)	0.026 (0.004–0.056)	0.001 (0.001–0.001)	0.164 ^{bc} (0.074–0.244)	85.0 (73.6–95.0)
	M018	11	0.027 ^e (0.019–0.050)	0.012 (0.001–0.045)	0.001 (0.001–0.001)	0.034 ^d (0.019–0.095)	71.7 (52.3–100)
	M019	2	0.205 ^a (0.198–0.212)	0.040 (0.033–0.047)	0.001 (0.001–0.001)	0.245 ^a (0.245–0.246)	83.6 (80.6–86.6)
	M020	2	0.065 ^{de} (0.060–0.072)	0.012 (0.004–0.021)	0.001 (0.001–0.001)	0.078 ^{cd} (0.064–0.092)	85.4 (77.3–93.6)
	M011	2	0.130 ^{bc} (0.121–0.140)	0.022 (0.022–0.023)	0.001 (0.001–0.001)	0.152 ^{bc} (0.143–0.163)	85.2 (84.8–85.6)
	M021	2	0.109 ^{bcd} (0.091–0.127)	0.021 (0.021–0.022)	0.001 (0.001–0.001)	0.130 ^{bcd} (0.113–0.148)	83.4 (80.7–86.1)
p-Value			<0.001	0.230	0.700	<0.001	0.104

^A Median (max–min); values with the same letters were not significantly different at p -value < 0.05 for the variable studied (Duncan's multiple range test).




Fig. 1. Boxplot with lines representing median, 25th and 75th percentiles, whiskers 10th and 90th percentiles and dots outliers for As_i in baby rice, rice cereals and rice crackers samples. Boxplot with open bars and diagonal hashed bars show As_i concentration in the rice-based products (baby rice, rice cereals and rice crackers) produced under non-organic (Non-Org.) and organic standards (Org.), respectively.

(0.039–0.254) mg/kg. As_i represents a median of 65.5% of ΣAs species concentration. Our results are similar to these and no significant differences were found in the As_i, DMA, ΣAs species concentration or As_i percentage (Table 3). Our results are also similar to the study carried out by Meharg et al., 2008, who analysed 17 samples of baby rice bought in the city of Aberdeen, UK, and reported a median As_i content of 0.110 mg/kg. However, our results show a wider range of As_i (0.056–0.268 mg/kg) than this study (0.060–0.160 mg/kg).

The FAO/WHO Expert Committee on Food Additives has determined the lower limit of the benchmark dose confidence limit (BMDL) based on a 0.5% increase in the incidence of lung cancer

to be 3.0 µg/kg b.wt. per day (WHO, 2011a). The Contaminants in Food Chain Panel (CONTAM) of EFSA determined the BMDL 1% extra risk for lung, skin and bladder cancers and skin lesions to be in the range of 0.3–8 µg/kg b.wt. per day and recommended using this instead of a single reference point in the risk characterisation for As_i intake (EFSA, 2009). The EFSA (2009) report stated that As_i exposure from food and water ranges from 0.13 to 0.56 µg/kg b.wt. for average consumers in 19 European countries. This estimated dietary exposure is within the range of the proposed BMDL values, indicating that the risk of toxicity cannot be excluded, especially for children whose dietary exposure has been estimated to be from 2 to 3-fold (0.50–2.66 µg/kg b.wt. per day)

Table 2

Effect of production under organic or non-organic standards on the concentration (mg/kg d.wt.) of As species in rice-based products (baby rice, rice cereals and rice crackers).

Food Category	Production method	Type of rice	n	As _i (mg/kg)	DMA (mg/kg)	MMA (mg/kg)	Σ As species (mg/kg)	As _i (%)
Baby rice	Non-organic	Milled	11	0.082 ^b (0.056–0.128) ^A	0.038 (0.022–0.061)	0.001 (0.001–0.003)	0.126 ^c (0.084–0.175)	67.1 ^{ab} (59.9–74.3)
		Whole grain	0	–	–	–	–	–
	Organic	Milled	9	0.121 ^b (0.087–0.141)	0.107 (0.024–0.120)	0.001 (0.001–0.003)	0.230 ^b (0.111–0.262)	55.1 ^b (51.4–78.5)
		Whole grain	9	0.190 ^a (0.117–0.268)	0.057 (0.038–0.123)	0.001 (0.001–0.004)	0.264 ^a (0.180–0.334)	79.3 ^a (57.2–84.6)
		<i>p</i> -Value		<0.001	0.014	0.302	<0.001	<0.001
Rice cereals	Non-organic	Milled	48	0.070 ^b (0.008–0.188)	0.036 (0.005–0.081)	0.001 (0.001–0.004)	0.113 ^b (0.042–0.248)	69.6 (14.2–89.6)
		Whole grain	0	–	–	–	–	–
	Organic	Milled	0	–	–	–	–	–
		Whole grain	5	0.162 ^a (0.103–0.323)	0.037 (0.016–0.068)	0.001 (0.001–0.003)	0.204 ^a (0.121–0.395)	81.6 (75.7–85.2)
		<i>p</i> -Value		<0.001	0.828	0.442	<0.001	0.025
Rice crackers	Non-organic	Milled	23	0.071 ^b (0.021–0.159)	0.023 (0.004–0.046)	0.001 (0.001–0.002)	0.095 ^b (0.034–0.206)	79.8 (52.3–93.6)
		Whole grain	22	0.081 ^b (0.018–0.211)	0.018 (0.001–0.048)	0.001 (0.001–0.002)	0.098 ^b (0.018–0.245)	81.3 (66.0–100.0)
	Organic	Milled	0	–	–	–	–	–
		Whole grain	52	0.121 ^a (0.048–0.195)	0.026 (0.004–0.171)	0.001 (0.001–0.006)	0.150 ^a (0.06–0.328)	82.5 (44.8–95.0)
		<i>p</i> -Value		<0.001	0.040	0.576	<0.001	0.153

^A Median (max–min); values with the same letters were not significantly different at *p*-value < 0.05 for the variable studied (Duncan's multiple range test).

Table 3

Total As concentration (mg/kg) and concentration (mg/kg) of As species in rice-based products (baby rice, rice cereals and rice crackers) from USA (FDA survey). Concentration (mg/kg) of As species in rice-based products (baby rice, rice cereals and rice crackers) from EU (Present study).

Product category	Sampling place	n	As _i (mg/kg)	DMA (mg/kg)	MMA (mg/kg)	Σ As species (mg/kg)	Total As (mg/kg)	As _i (%)
Baby rice	US	85	0.114 (0.039–0.254) ^A	0.071 (0.015–0.204)	0.003a (0.001–0.012)	0.215 (0.550–0.341)	0.200 (0.060–0.373)	65.5 (22.0–82.9)
	EU	29	0.121 (0.056–0.268)	0.041 (0.023–0.123)	0.001 ^b (0.001–0.004)	0.175 (0.084–0.334)	–	69.8 (51.4–84.6)
	<i>p</i> -Value		0.235	0.085	0.002	0.676	–	0.028
Rice cereals	US	105	0.091 (0.023–0.283)	0.042 ^a (0.007–0.493)	0.001 (0.001–0.014)	0.135 (0.041–0.625)	0.132 (0.050–0.723)	63.0 (18.8–87.7)
	EU	53	0.075 (0.008–0.323)	0.037 ^b (0.005–0.081)	0.001 (0.001–0.004)	0.119 (0.042–0.396)	–	70.6 (14.2–89.6)
	<i>p</i> -Value		0.096	0.004	0.590	0.005	–	0.077
Rice crackers	US	199	0.079 (0.008–0.273)	0.034 ^a (0.001–0.477)	0.001 ^a (0.001–0.021)	0.116 (0.012–0.657)	0.121 (0.009–1.931)	65.1 ^b (18.4–97.6)
	EU	97	0.111 (0.018–0.211)	0.025 ^b (0.001–0.172)	0.001 ^b (0.001–0.006)	0.141 (0.019–0.328)	–	80.9 ^a (44.8–100.0)
	<i>p</i> -Value		0.011	<0.001	<0.001	0.224	–	<0.001

^A Median (max–min); values with the same letters were not significantly different at *p*-value < 0.05 for the variable studied.

that of adults due to their greater food consumption in relation to their body weight (EFSA, 2009).

The plots A and B, included in Fig. 2, show the As_i exposure for infants (4 months of age) consuming baby rice. The median of 0.70 µg As_i/serving (Table 4) and the 3th (5.35 kg), 15th (5.90 kg), 50th (6.70 kg), 85th (7.55 kg) and 97th (8.35 kg) percentiles of body weight (WHO, 2014) have been used for calculations in the plot A. The median of 6.70 kg b.wt. and the 3th (0.35 µg As_i/serving), 15th (0.41 µg As_i/serving), 50th (0.70 µg As_i/serving) 85th (1.40 µg As_i/serving) and 97th (1.77 µg As_i/serving) percentiles of As_i concentration per serving have been used for calculations in the plot B. When the median body weight and the median As_i concentration per serving are used for calculations, 3 serving lead to an exposure of 0.31 µg As_i/kg b.wt. per day. This could be 1.67 to 3.37-fold higher when 4 servings are consumed and the median As_i concentration per serving with the 3th percentile of body weight, or the median body weight with the 97th percentile of As_i concentration per serving is used for calculations, respectively. These values are within the BMDL₀₁ range identified by EFSA (2009), which highlights that the risk cannot be excluded for infants consuming baby rice.

3.2. Rice cereals

The ΣAs species concentration in 53 rice cereal samples belonging to 6 different commercial brands ranged from 0.042 to 0.396 mg/kg (Table 1). The As_i percentage ranged from 14.2% to 89.6% of the ΣAs species concentration. The rice cereal brands M006, M007, M008, M009 and M011 had a significantly higher

percentage of As_i (median of 74.2%, 81.5%, 76.6%, 83.1% and 76.6%, respectively) than the M010 (median of 62.7%; *p* < 0.001). The As_i concentration ranged from 0.008 to 0.323 mg/kg and showed a good correlation with the rice content, which means that most of the As_i is coming from the rice. Due to As_i concentration level 2% of the rice cereal samples evaluated would be above the JECFA proposed maximum level. In addition, the rice cereal brand M009 had a higher As_i concentration (median of 0.234 mg/kg; *p* < 0.001) than the other rice cereals samples analysed, which could reach up to 1.6 times the JECFA maximum level. The major organic As species for each baby rice was DMA, ranging from 0.005 to 0.082 mg/kg. Rice cereal samples labelled as produced under organic standards showed higher levels of As_i (median of 0.162 mg/kg) than non-organic samples (median of 0.070 mg/kg; *p* < 0.001) (Fig. 1). As pointed out in the section on baby rice this is associated with the common inclusion of whole grain rice to produce organic products (Table 2).

The US FDA dataset on As_i in rice and rice products from the US market includes 105 rice cereals (product subcategory: hot/ready-to-eat cereals). The median and range of As_i in rice cereals from the USA market is 0.091 (0.023–0.283) mg/kg. As_i represents a median of 63.0% of ΣAs species concentration. Our results are similar to those and no significant differences were found in the As_i concentration or percentage (Table 3). However, rice cereals from the USA market had higher levels of DMA (median of 0.042 mg/kg) and ΣAs species concentration (median of 0.135 mg/kg) than rice cereals from the EU market (0.037 mg/kg; *p* = 0.004 and 0.119 mg/kg; *p* = 0.005, respectively) (Table 3). This supports the fact that the geographical origin affects the As concentration and speciation in




Fig. 2. Estimation of As_i exposure for infants and young children from baby rice, rice cereals and rice crackers. (The A and B plots show the As_i exposure of infants consuming baby rice. The median of $0.70 \mu\text{g } As_i/\text{servicing}$ and the 3rd, 15th, 50th, 85th and 97th percentiles of body weight have been used for calculations in plot A. The median of 6.70 kg b.w.t. and the 3rd, 15th, 50th, 85th and 97th percentiles of $\mu\text{g } As_i/\text{servicing}$ have been used for calculations in plot B. The C, D, E and F plots show the As_i exposure of young children consuming rice cereals (C and D) and rice crackers (E and F). The median of $2.22 \mu\text{g } As_i/\text{servicing}$, the median of $3.58 \mu\text{g } As_i/\text{servicing}$ and the 3rd, 15th, 50th, 85th and 97th percentile of body weight have been used for calculations in the plots C and E, respectively. The median of 9.30 kg b.w.t. and the 3rd, 15th, 50th, 85th and 97th percentile of $\mu\text{g } As_i/\text{servicing}$ have been used for calculations in plot D and F).

Table 4

Estimation of the As_i intake ($\mu\text{g}/\text{servicing}$) from baby rice, rice cereals and rice crackers.

Food category	Sampling place	n	Moisture (%)	As_i (mg/kg w.wt.)	Recommended unit (g w.wt.)	Recommended serving (g w.wt.)	As_i intake per unit (mg)	As_i intake per serving (mg)
Baby rice	US	85	n.a	0.114 (0.039–0.254)	n.a	15.0 (15.0–15.0)	n.a	1.7 (0.6–3.8)
	EU	29	2.3 (0.8–5.0)	0.117 (0.056–0.262)	n.a	6.0 (6.0–12.0)	n.a	0.7 (0.3–2.2)
Rice cereals	US	105	n.a	0.091 (0.023–0.283)	n.a	30.0 (15.0–55.0)	n.a	2.6 (0.8–11.0)
	EU	53	1.1 (0.3–7.6)	0.075 (0.008–0.313)	n.a	30.0 (22.0–35.0)	n.a	2.2 (0.2–9.3)
Rice crackers	US	199	n.a	0.079 (0.008–0.273)	n.a	30.0 (7.0–30.0)	n.a	2.4 (0.2–8.2)
	EU	97	3.1 (0.3–5.8)	0.108 (0.018–0.201)	1.6 (1.4–19.2)	30.0 (8.0–50.0)	0.3 (0.0–2.0)	3.5 (0.5–9.1)

rice. Meharg et al. (2009) reported that the relationship between As_i content versus total As content differs among countries, with Bangladesh and India having the steepest slope in linear regression, and the US having the shallowest slope due to higher levels of DMA.

The plots C and D, included in Fig. 2, show the As_i exposure for young children (12 months of age) consuming rice cereals. The

median of $2.22 \mu\text{g } As_i/\text{servicing}$ (Table 4) and the 3rd (7.50 kg), 15th (8.20 kg), 50th (9.30 kg), 85th (10.50 kg) and 97th (12.10 kg) percentiles of body weight (WHO, 2014) have been used for calculations in the plot C. The median of 9.30 kg b.w.t. and the 3rd ($0.58 \mu\text{g } As_i/\text{servicing}$), 15th ($1.47 \mu\text{g } As_i/\text{servicing}$), 50th ($2.22 \mu\text{g } As_i/\text{servicing}$) 85th ($3.11 \mu\text{g } As_i/\text{servicing}$) and 97th ($5.10 \mu\text{g } As_i/\text{servicing}$) percentiles of As_i concentration per serving have been used for calculations in

the plot D. When the median body weight and the median As_i concentration per serving are used for calculations, 2 serving lead to an exposure of 0.47 µg As_i/kg b.wt. per day. This could be 2.48 to 4.58-fold higher when 4 servings are consumed and the median As_i concentration per serving with the 3th percentile of body weight, or the median body weight with the 97th percentile of As_i concentration per serving is used for calculations, respectively. These values are within the BMDL₀₁ range identified by EFSA (2009), which highlights that the risk cannot be excluded for infants consuming rice cereals.

3.3. Rice crackers

The ΣAs species concentration in 97 rice cracker samples belonging to 15 different commercial brands ranged from 0.019 to 0.246 mg/kg (Table 1). The As_i percentage ranged from 71.1% to 100% of the ΣAs species concentration. The As_i concentration ranged from 0.019 to 0.212 mg/kg and showed a good correlation with the rice content, which means that most of the As_i is coming from the rice. Due to the As_i concentration level 1% of the rice crackers samples evaluated would be above the JECFA proposed maximum level. The rice cracker brand M019 had the highest As_i concentration (median of 0.205 mg/kg; $p < 0.001$). The major organic As species for each rice cracker was DMA, ranging from 0.001 to 0.172 mg/kg. Rice crackers labelled as produced under organic standards showed higher levels of As_i (median of 0.121 mg/kg) than non-organic samples elaborated with milled rice (median of 0.071 mg/kg; $p < 0.001$) (Table 2 and Fig. 1).

The US FDA dataset on As_i in rice and rice products from the USA market includes 199 samples of rice crackers (product subcategory: rice cakes and savoury rice snacks). The median and range of As_i in rice crackers from the US market is 0.079 (0.008–0.273) mg/kg. As_i content represents a median of 65.1% of ΣAs species concentration. Our results are similar to those and no significant differences were found in the As_i or ΣAs species concentration (Table 3). However, rice crackers from the US market had higher levels of DMA (median of 0.034 mg/kg) than rice crackers from the EU market (median of 0.025 mg/kg; $p < 0.001$). Thus the As_i percentage in rice crackers from the EU market (median of 80.9%) was higher than the As_i percentage in rice crackers from the USA market (median of 65.1%; $p < 0.001$) (Table 3). This is in agreement with the geographical variation in As concentration in rice, which might be associated with breeding/cultivar selection and agronomic practices (Meharg & Zhao, 2012; Meharg et al., 2009). Rice processing is another factor that may affect the As concentration in the final product. Extensive rinsing of uncooked grain followed by cooking rice in a large excess of water and discarding the water on cessation of cooking may reduce the As_i content of food by up to 30% when As_i free water is used (Raab, Baskaran, Feldman, & Meharg, 2008; Sengupta et al., 2006; Signes, Mitra, Burlo, & Carbonell-Barrachina, 2008).

The plots E and F, included in Fig. 2, show the As_i exposure for young children (12 months of age) consuming rice crackers. The median of 3.58 µg As_i/serving (Table 4) and the 3th (7.50 kg), 15th (8.20 kg), 50th (9.30 kg), 85th (10.50 kg) and 97th (12.10 kg) percentiles of body weight (WHO, 2014) have been used for calculations in the plot E. The median of 9.30 kg b.wt. and the 3th (0.61 µg As_i/serving), 15th (0.90 µg As_i/serving), 50th (3.58 µg As_i/serving) 85th (6.48 µg As_i/serving) and 97th (8.01 µg As_i/serving) percentiles of As_i concentration per serving have been used for calculations in the plot F. When the median body weight and the median As_i concentration per serving are used for calculations, 1 serving lead to an exposure of 0.38 µg As_i/kg b.wt per day. This could be 4.95 to 8.93-fold higher when 4 servings are consumed and the median As_i concentration per serving with the 3th percentile of body weight, or the median body weight with the 97th percentile of As_i concen-

tration per serving is used for calculations, respectively. These values are within the BMDL₀₁ range identified by EFSA (2009), which highlights that the risk cannot be excluded for infants consuming rice crackers.

4. Conclusions

Baby rice, rice cereals and crackers are widely used during weaning and to feed young children due to its availability, bland taste, nutritional value and relatively low allergic potential. They are particularly used to feed infants and young children affected by celiac disease whose only viable treatment is a gluten-free diet. The commercial brand affected the As_i concentration in the baby rice, rice cereals and rice crackers evaluated, which may be associated to the amount of rice, rice origin and manufacturing process. In fact, rice-based products including whole grain rice showed the highest As_i concentrations and they were usually labelled as produced under organic standards. The As_i concentration found in several samples was higher than the JECFA maximum level. The As_i exposure for infants consuming baby rice and young children consuming rice cereals and rice crackers are close or within the range of BMDL₀₁ values identified by EFSA (2009) and therefore the risk cannot be excluded. In addition, this could be worse when these products represent a major part of the diet and As_i contribution is accumulated. Despite this scenario there are currently no EU or USA regulations applicable to As_i in food to protect the most vulnerable subpopulation. Thus, we conclude that there is an urgent need for regulatory limits on As_i in food, especially in rice and rice-based products.

Acknowledgement

This work was funded by a Marie Curie Intra-European Fellowship within the 7th European Community Framework Programme (PIEF-GA-2013-622096).

References

- Banerjee, M., Banerjee, N., Bhattacharjee, P., Mondal, D., Lythgoe, P. R., Martínez, M., et al. (2013). High arsenic in rice is associated with elevated genotoxic effects in humans. *Scientific Reports*, 3, 2195.
- Burlo, F., Ramirez-Gandolfo, A., Signes-Pastor, A. J., Haris, P. I., & Carbonell-Barrachina, A. A. (2012). Arsenic contents in Spanish infant rice, pureed infant foods, and rice. *Journal of Food Science*, 77, 15–19.
- Carbonell-Barrachina, A. A., Wu, X., Ramirez-Gandolfo, A., Norton, G. J., Burlo, F., Deacon, C., et al. (2012). Inorganic arsenic contents in rice-based infant foods from Spain, UK, China and USA. *Environmental Pollution*, 163, 77–83.
- Cascio, C., Raab, A., Jenkins, R. O., Felman, J., & Meharg, A. A. (2011). The impact of a rice based diet on urinary arsenic. *Journal of Environmental Monitoring*, 13, 257–265.
- Choi, S. H., Kim, J. S., Lee, Y. J., Jeon, J. S., Kim, J. W., Russo, R. E., et al. (2014). Analysis of arsenic in rice grains using ICP-MS and fs LA-ICP-MS. *Journal of Analytical Atomic Spectrometry*, 29, 1233–1237.
- Council Directive 98/83/EC. (1998). OJ. L330/32-L330/53. Available: https://www.fsai.ie/uploadedFiles/Legislation/Food_Legislation_Links/Water/EU_Directive_98_83_EC.pdf. [Online 20 July 2014].
- Da Sacco, L., Baldassarre, A., & Massotti, A. (2013). Diet's role in the toxicity of inorganic arsenic (iAs): A journey from soil to children's moth. *Journal of Geochemical Exploration*, 131, 45–51.
- EFSA (European Food Safety Authority) Panel on Contaminants in the Food Chain. (2009). Scientific opinion on arsenic in food. *EFSA Journal*, 7(10), 1351. <http://www.efsa.europa.eu/en/efsajournal/doc/1351.pdf> [Online 20 July 2014].
- Gilbert-Diamond, D., Cottingham, K. L., Gruber, J. F., Punshon, T., Sayarath, V., Gandolfi, A. J., et al. (2011). Arsenic exposure contributes to arsenic exposure in US women. *Proceedings of the National Academy of Science of the United States of America*, 108, 20656–20660.
- IARC (International Agency for Cancer Research). (2004). Some Drinking-Water Disinfectants and Contaminants, Including Arsenic. IARC Monograph on the Evaluation of Carcinogenic Risk to Humans Vol. 84. Available: <http://monographs.iarc.fr/ENG/Monographs/vol84/mono84.pdf>. [Online 20 July 2014].
- JECFA (Food and Agriculture Organization/ World Health Organization). (2010). Joint FAO/WHO food standards committee of the Codex Alimentarius. Available: http://www.who.int/foodsafety/chem/summary72_rev.pdf. [Online 20 July 2014].

- JECFA (Food and Agriculture Organization/ World Health Organization). (2014). Joint FAO/WHO food standards committee of the Codex Alimentarius. Available: http://www.codexalimentarius.org/input/download/report/776/REP12_CFe.pdf. [Online 20 July 2014].
- Juhász, A. L., Smith, E., Weber, J., Rees, M., Rofe, A., Kuchel, T., et al. (2006). In vivo assessment of arsenic bioavailability in rice and its significance for human health risk assessment. *Environmental Health Perspectives*, *114*, 1826–1831.
- Meharg, A. A., Lombi, E., Williams, P. N., Scheckel, K. G., Feldmann, J., Raab, A., et al. (2008). Speciation and localization of arsenic in white and brown rice grains. *Environmental Science and Technology*, *42*, 1051–1057.
- Meharg, A. A., Williams, P. N., Adomako, E., Lawgali, Y. Y., Deacon, C., Villada, A., et al. (2009). Geographical variation in total and inorganic arsenic content in polished (white) rice. *Environmental Science and Technology*, *43*, 1612–1617.
- Meharg, A. A., & Zhao, F. J. (2012). *Arsenic and rice*. Springer [ISBN 978-94-007-2946-9].
- Mennella, J. A., Ziegler, P., Briefel, R., & Novak, T. (2006). Feeding Infants and Toddlers Study: The types of foods fed to Hispanic infants and toddlers. *Journal of American Dietetic Association*, *106*, 96–106.
- Munera-Picazo, S., Burló, F., & Carbonell-Barrachina, A. (2014). Arsenic speciation in rice-based food for adults with celiac disease. *Food Additives and Contaminants*, *31*, 1358–1366.
- Newton, K. P., & Singer, S. A. (2012). Celiac disease in children and adolescents: Special considerations. *Seminars in Immunopathology*, *34*, 479–496.
- Raab, A., Baskaran, C., Feldman, J., & Meharg, A. A. (2008). Cooking rice in a high water to rice ratio reduces inorganic arsenic content. *Journal of Environmental Monitoring*, *11*, 41–44.
- Sengupta, M. K., Hossain, M. A., Mukherjee, A., Ahamed, S., Das, B., Pas, A., et al. (2006). Arsenic burden of cooked rice: Traditional and modern methods. *Food and Chemical Toxicology*, *44*, 1823–1829.
- Signes, A., Mitra, K., Burlo, F., & Carbonell-Barrachina, A. A. (2008). Effect of cooking method and rice type on arsenic concentration in cooked rice and the estimation of arsenic dietary intake in a rural village in West Bengal, India. *Food Additives and Contaminants*, *25*, 1345–1352.
- Signes-Pastor, A. J., Al-Rmalli, S. W., Jenkins, R. O., Carbonell-Barrachina, A. A., & Haris Parvez, I. H. (2012). Arsenic bioaccessibility in cooked rice as affected by arsenic in cooking water. *Journal of Food Science*, *77*, 201–2016.
- Smith, A. H., Marshall, G., Yuan, Y., Ferreccui, C., Liaw, J., von Ehrenstein, O., et al. (2006). Increased mortality from lung cancer and bronchiectasis in young adults after exposure to arsenic in utero and in early childhood. *Environmental Health Perspectives*, *114*, 1293–1296.
- Torres-Escribano, S., Leal, M., Vélez, D., & Montoro, R. (2008). Total and inorganic arsenic concentrations in rice sold in Spain, effect of cooking, and risk assessments. *Environmental Science and Technology*, *42*, 3867–3872.
- USFDA (U.S. Food and Drug Administration). (2013). Arsenic in rice: Full analytical results from rice/rice products. Available: <http://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm319916.htm>. [Online 23 July 2014].
- US FDA (U.S. Food and Drug Administration). (2014). Arsenic in rice and rice products. Available: <http://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm319948.htm>. [Online 22 July 2014].
- Vahter, M. (2009). Effects of arsenic on maternal and fetal health. *Annual Review of Nutrition*, *29*, 381–399.
- Wei, Y., Zhu, J., & Hguyen, A. (2014). Rice consumption and urinary concentrations of arsenic in US adults. *International Journal Environmental Health Research*, *24*, 459–470.
- WHO (World Health Organization). (2010). Exposure to arsenic: a major public health concern. Available: <http://www.who.int/ipcs/features/arsenic.pdf?ua=1>. [Online 22 July 2014].
- WHO (World Health Organization). (2011a). Safety evaluation of certain contaminants in food. WHO Food Additives Series No. 63. Available: http://whqlibdoc.who.int/publications/2011/9789241660631_eng.pdf. [Online 20 July 2014].
- WHO (World Health Organization). (2011b). Arsenic in drinking-water. Background document for preparation of WHO Guidelines for drinking-water quality. Available: http://www.who.int/water_sanitation_health/dwq/chemicals/arsenic.pdf. [Online 20 July 2014].
- WHO (World Health Organization). (2014). Child growth standard. Weight-for-age. Available: http://www.who.int/childgrowth/standards/weight_for_age/en/ [Online 20 July 2014].
- William, P. N., Villada, A., Deacon, C., Raab, A., Figuerola, J., Green, A., et al. (2007). Greatly enhanced arsenic shoot assimilation in rice lead to elevate grain levels compared to wheat and barley. *Environmental Science and Technology*, *41*, 6854–6859.
- Xu, X. Y., McGrath, S. P., Meharg, A. A., & Zhao, F. J. (2008). Growing rice aerobically markedly decreases arsenic accumulation. *Environmental Science and Technology*, *42*, 5574–5579.