




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Motivic periods and Grothendieck arithmetic invariants



F. Andreatta^a, L. Barbieri-Viale^{a,*}, A. Bertapelle^b,
appendix by B. Kahn^c

^a *Dipartimento di Matematica “F. Enriques”, Università degli Studi di Milano,
via C. Saldini, 50, Milano I-20133, Italy*

^b *Dipartimento di Matematica “T. Levi-Civita”, Università degli Studi di Padova,
via Trieste, 63, Padova I-35121, Italy*

^c *IMJ-PRG, Case 247, 4 place Jussieu, 75252 Paris Cedex 05, France*

ARTICLE INFO

Article history:

Received 9 January 2019

Received in revised form 27

September 2019

Accepted 9 October 2019

Available online 7 November 2019

Communicated by A. Asok

MSC:

14F42

14F40

19E15

14C30

14L15

Keywords:

Motives

Periods

Motivic and de Rham cohomology

ABSTRACT

We construct a period regulator for motivic cohomology of an algebraic scheme over a subfield of the complex numbers. For the field of algebraic numbers we formulate a period conjecture for motivic cohomology by saying that this period regulator is surjective. Showing that a suitable Betti–de Rham realization of 1-motives is fully faithful we can verify this period conjecture in several cases. The divisibility properties of motivic cohomology imply that our conjecture is a neat generalization of the classical Grothendieck period conjecture for algebraic cycles on smooth and proper schemes. These divisibility properties are treated in an appendix by B. Kahn (extending previous work of Bloch and Colliot-Thélène–Raskind).

© 2019 Elsevier Inc. All rights reserved.

* Corresponding author.

E-mail addresses: Fabrizio.Andreatta@unimi.it (F. Andreatta), Luca.Barbieri-Viale@unimi.it (L. Barbieri-Viale), Alessandra.Bertapelle@unipd.it (A. Bertapelle), Bruno.Kahn@imj-prg.fr (B. Kahn).

Contents

0. Introduction 2
 1. Periods: constructions and conjectures 5
 2. Periods of 1-motives: fullness of Betti-de Rham realizations 13
 3. Some evidence: description of some Grothendieck arithmetic invariants 29
 Appendix A. Divisibility properties of motivic cohomology (by B. Kahn) 42
 References 48

0. Introduction

Let X be a scheme which is separated and of finite type over a subfield K of the complex numbers. Consider the q -twisted singular cohomology $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ of the analytic space X_{an} associated to the base change of X to \mathbb{C} and the p th de Rham cohomology $H^p_{\text{dR}}(X)$, which is an algebraically defined K -vector space. We have the following natural \mathbb{C} -linear isomorphism

$$\varpi_X^{p,q} : H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)) \otimes_{\mathbb{Z}} \mathbb{C} \cong H^p_{\text{dR}}(X) \otimes_K \mathbb{C}$$

providing a comparison between these cohomology theories. As Grothendieck originally remarked, for X defined over the field of algebraic numbers $K = \overline{\mathbb{Q}}$ or a number field, the position of the whole $H^p_{\text{dR}}(X)$ with respect to $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ under $\varpi_X^{p,q}$ «yields an interesting arithmetic invariant, generalizing the “periods” of regular differential forms» (see [30, p. 101 & footnotes (9) and (10)], cf. [3, §7.5 & Chap. 23], [16], [17] and [33, Chap. 5 & 13]). For the comparison of several notions of “periods” and versions of the period conjecture we refer to Huber survey [32] (see also [17, §2.2.2]).

The main goal of this paper is to describe this arithmetic invariant, at least for $p = 1$ and all twists, notably, $q = 1$ and $q = 0$. In more details, we first reconstruct $\varpi_X^{p,q}$ (in Definition 1.2.4) by making use of Ayoub’s period isomorphism (see Lemma 1.2.2) in Voevodsky’s triangulated category $\text{DM}_{\text{ét}}^{\text{eff}}$ of motivic complexes for the étale topology. Denote by $H_{\varpi}^{p,q}(X)$ the named arithmetic invariant, *i.e.*, the subgroup of those cohomology classes in $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ which are landing in $H^p_{\text{dR}}(X)$ via $\varpi_X^{p,q}$. We then show the existence of a regulator map (see Corollary 1.2.6 and Definition 1.2.7)

$$r_{\varpi}^{p,q} : H^{p,q}(X) \rightarrow H_{\varpi}^{p,q}(X)$$

from étale motivic cohomology groups $H^{p,q}(X)$. Here we regard motivic cohomology canonically identified with $H_{\text{é}h}^p(X, \mathbb{Z}(q))$ where $\mathbb{Z}(q)$ is the Suslin-Voevodsky motivic complex (see [47, Def. 3.1]), as a complex of sheaves for the éh-topology (introduced in [12, §10.2]). We are mostly interested in the case of $q = 0, 1$ so that $\mathbb{Z}(0) \cong \mathbb{Z}[0]$ and $\mathbb{Z}(1) \cong \mathbb{G}_m[-1]$ by a theorem of Voevodsky (see [47, Thm. 4.1]).

Following Grothendieck’s idea, we conjecture that the period regulator $r_{\varpi}^{p,q}$ is surjective over $\overline{\mathbb{Q}}$ and we actually show some evidence. We easily see that $H_{\varpi}^{0,q}(X) = 0$

for $q \neq 0$ and $r_{\varpi}^{0,0}$ is an isomorphism: therefore, the first non-trivial case is for $p = 1$. Moreover, by making use of Suslin-Voevodsky rigidity theorem we can show that $r_{\varpi}^{p,q}$ is surjective on torsion (see Lemma 1.4.2). We can also show: if the vanishing $H^{p,q}(X) \otimes \mathbb{Q}/\mathbb{Z} = 0$ holds true then the surjectivity of $r_{\varpi}^{p,q}$ is equivalent to the vanishing $H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(q)) = 0$. The divisibility properties of motivic cohomology (see Appendix A) imply that our conjecture is a neat generalization of the classical period conjecture for algebraic cycles on smooth and proper schemes (see Proposition 1.4.4).

In order to study the case $p = 1$ we can make use of the description of H^1 via the Albanese 1-motive $L_1\text{Alb}(X)$. Recall the existence of the homological motivic Albanese complex $\text{LAlb}(X)$, a complex of 1-motives whose p th homology $L_p\text{Alb}(X)$ is a 1-motive with cotorsion (see [12, §8.2] for details). We can regard complexes of 1-motives as objects of $\text{DM}_{\text{ét}}^{\text{eff}}$ and by the adjunction properties of LAlb (proven in [12, Thm. 6.2.1]) we have a natural map

$$\mathbb{E}x\text{t}^p(\text{LAlb}(X), \mathbb{Z}(1)) \rightarrow H^{p,1}(X) \cong H_{\text{é}h}^{p-1}(X, \mathbb{G}_m)$$

which is an isomorphism, rationally, for all p (see the motivic Albanese map displayed in (3.2) and (3.3) below). We can also describe periods for 1-motives (see Definition 2.2.1) in such a way that we obtain suitable Betti-de Rham realizations in period categories (see Definitions 2.5.4 and 2.5.1): a key point is that these realizations are fully faithful over $\overline{\mathbb{Q}}$ (see Theorem 2.7.1). The main ingredient in the proof of fullness is a theorem due to Waldschmidt [54, Thm. 5.2.1] in transcendence theory, generalizing the classical Schneider-Lang theorem (see also [16, Thm. 4.2]). An alternative proof can be given using a theorem of Wüstholz [56] (see our second proof of Theorem 2.7.1). A version of Baker’s theorem and instances of Kontsevich period conjecture for 1-motives are further explored in a recent work of Huber and Wüstholz [34]. Note that Kontsevich’s period conjecture for 1-motives was formulated in [57] (see also [3, §23.3.3]).

Actually, we show that the regulator $r_{\varpi}^{p,1}$ can be revisited by making use of 1-motives (see Lemmas 3.2.1 to 3.2.3 and Proposition 3.2.7). As a byproduct, all this promptly applies to show the surjectivity of $r_{\varpi}^{1,1} : H_{\text{é}h}^0(X, \mathbb{G}_m) \rightarrow H_{\varpi}^{1,1}(X)$ via $\text{Ext}(\text{LAlb}(X), \mathbb{Z}(1))$ verifying the conjecture for $p = 1$ and $q = 1$ (see Theorem 3.2.4). In fact, we can use the motivic Picard complex $\text{RPic}(X)$ (see [12, §8.3]) so that $\text{Ext}^p(\mathbb{Z}(0), \text{RPic}(X)) = \text{Ext}^p(\text{LAlb}(X), \mathbb{Z}(1))$ by Cartier duality showing that $\text{Ext}(\text{LAlb}(X), \mathbb{Z}(1))$ is an extension of the finitely generated group $\text{Hom}(\mathbb{Z}, \text{R}^p\text{Pic}(X))$ by a divisible group; we thus conclude that $r_{\varpi}^{p,1}$ induces a map

$$\theta_{\varpi}^p : \text{Hom}(\mathbb{Z}, \text{R}^p\text{Pic}(X)) \rightarrow H_{\varpi}^{p,1}(X)$$

which in turn can be described making use of the mentioned Betti-de Rham realization. The surjectivity of $r_{\varpi}^{p,1}$ can be translated via θ_{ϖ}^p and the fullness of the Betti-de Rham realization. For $p = 1$, considering the 1-motive $\text{R}^1\text{Pic}(X) = [L_1^* \xrightarrow{u_1^*} G_1^*]$ which is the Cartier dual of $L_1\text{Alb}(X) = [L_1 \xrightarrow{u_1} G_1]$, we get a canonical isomorphism

$$\text{Ker } u_1^* \cong \text{Hom}(\mathbb{Z}, \text{R}^1\text{Pic}(X)) \cong H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}_{\text{an}}(1)) = H_{\varpi}^{1,1}(X).$$

In particular, we obtain that $H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}(1)) = 0$ if X is proper. This vanishing for smooth projective varieties was previously obtained by Bost-Charles [17, Thm. 4.2].

With some more efforts, making now use of the motivic complex $L\pi_0(X)$ along with its adjunction property (as stated in [12, §5.4]), we get a map

$$\mathbb{E}xt^p(L\pi_0(X), \mathbb{Z}(0)) \rightarrow H^{p,0}(X) \cong H_{\text{ét}}^p(X, \mathbb{Z}).$$

Analysing the composition of this map for $p = 1$ with $r_{\varpi}^{1,0}$ we see that

$$r_{\varpi}^{1,0} : H_{\text{ét}}^1(X, \mathbb{Z}) \cong H_{\varpi}^{1,0}(X) = H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}_{\text{an}})$$

is an isomorphism (see Theorem 3.3.1), which yields the case $p = 1$ and $q = 0$ of our conjecture. In particular, $H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}_{\text{an}}) = 0$ for X normal. This vanishing for smooth quasi-projective varieties was previously obtained by Bost-Charles [17, Thm. 4.1].

For $p = 1$ and $q \neq 0, 1$ we have that $H_{\varpi}^{1,q}(X) = 0$ (see Corollary 3.4.2) so that the period conjecture for motivic cohomology is trivially verified.

Remarkably, the description of the Grothendieck arithmetic invariants $H_{\varpi}^{p,q}(X)$ appears strongly related to the geometric properties encoded by motivic cohomology. These properties are almost hidden for smooth schemes, since the divisibility properties of motivic cohomology of X smooth yields that for $p \notin [q, 2q]$ the surjectivity of $r_{\varpi}^{p,q}$ is equivalent to the vanishing $H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(q)) = 0$. However, for X smooth with a smooth compactification \overline{X} and normal crossing boundary Y , we have that

$$\text{Ker} (\text{Div}_Y^0(\overline{X}) \xrightarrow{u_1^*} \text{Pic}_{\overline{X}/\mathbb{Q}}^0) \cong H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}_{\text{an}}(1))$$

where u_1^* is the canonical mapping sending a divisor D supported on Y to $\mathcal{O}_{\overline{X}}(D)$. In fact, here $\text{R}^1\text{Pic}(X)$ is Cartier dual of $\text{L}_1\text{Alb}(X) = [0 \rightarrow \mathcal{A}_{X/\mathbb{Q}}^0]$, the Serre-Albanese semi-abelian variety (see [12, Chap. 9]). Therefore, there exist smooth schemes X such that $H_{\varpi}^{1,1}(X)$ is non-zero and the vanishings in [17, Thm. 4.1 & 4.2] are particular instances of our descriptions.

With similar techniques one can make use of the Borel-Moore Albanese complex $\text{LAlb}^c(X)$ (see [12, Def. 8.7.1]) to describe the compactly supported variant $H_{c,\varpi}^{1,q}(X)$, for any twist q .

Finally, the cohomological Albanese complex $\text{LAlb}^*(X)$ (see [12, Def. 8.6.2]) shall be providing a description of $H_{\varpi}^{2d-j,q}(X)$ for $d = \dim(X)$, at least for $j = 0, 1$ and q an arbitrary twist. A homological version of period regulators is also feasible and will be discussed in a future work.

Acknowledgements

We would like to thank Y. André and J. Ayoub for some useful discussions on the matters treated in this paper. The first author was partially supported by the Cariplo project n. 2017-1570 and all Italian authors acknowledge the support of the *Ministero dell’Istruzione, dell’Università e della Ricerca* (MIUR) through the Research Project (PRIN 2010-11) “Arithmetic Algebraic Geometry and Number Theory”. We thank the referee for her/his many comments that helped us to considerably improve the exposition (starting with the suggestion to change the title to something reflecting more accurately the contents of the paper).

1. Periods: constructions and conjectures

Let DM_{τ}^{eff} be the effective (unbounded) triangulated category of Voevodsky motivic complexes of τ -sheaves over a field K of zero characteristic, *i.e.*, the full triangulated subcategory of $D(\text{Shv}_{\tau}^{\text{tr}}(\text{Sm}_K))$ given by \mathbb{A}^1 -local complexes (*e.g.* see [6, §4.1] and, for complexes bounded above, see also [47, Lect. 14]). We here generically denote by τ either the Nisnevich or étale Grothendieck topology on Sm_K , the category of smooth schemes which are of finite type over the field K . Let $\mathbb{Z}(q)$ for $q \geq 0$ be the Suslin-Voevodsky motivic complex regarded as a complex of étale sheaves with transfers. More precisely we consider a change of topology tensor functor

$$\alpha : DM_{\text{Nis}}^{\text{eff}} \rightarrow DM_{\text{ét}}^{\text{eff}}$$

and $\mathbb{Z}(q) = \alpha\mathbb{Z}_{\text{Nis}}(q)$ (see [12, Cor. 1.8.5 & Def. 1.8.6]) where $\mathbb{Z}_{\text{Nis}}(q)$ is the usual complex for the Nisnevich topology (see also [50, Def. 3.1]). We have the following canonical isomorphisms $\mathbb{Z}(0) \cong \mathbb{Z}[0]$, $\mathbb{Z}(1) \cong \mathbb{G}_m[-1]$ and $\mathbb{Z}(q) \otimes \mathbb{Z}(q') \cong \mathbb{Z}(q+q')$ for any $q, q' \geq 0$ (see [50, Lemma 3.2]). For any object $M \in DM_{\text{ét}}^{\text{eff}}$ we here denote $M(q) := M \otimes \mathbb{Z}(q)$. Recall that by inverting the Tate twist $M \rightsquigarrow M(1)$ we obtain DM_{τ} (where every compact object is isomorphic to $M(-n)$ for some $n \geq 0$ and M compact and effective). For $M \in DM_{\text{ét}}^{\text{eff}}$ we shall define its motivic cohomology as

$$H^{p,q}(M) := \text{Hom}_{DM_{\text{ét}}^{\text{eff}}}(M, \mathbb{Z}(q)[p]).$$

For any algebraic scheme X we have the Voevodsky étale motive $M(X) = \alpha C_{\bullet} \mathbb{Z}_{\text{tr}}(X) \in DM_{\text{ét}}^{\text{eff}}$ where C_{\bullet} is the Suslin complex and $\mathbb{Z}_{\text{tr}}(X)$ is the representable Nisnevich sheaf with transfers (see [47, Def. 2.8, 2.14 & Properties 14.5] and compare with [12, Lemma 1.8.7 & Sect. 8.1]). We then write $H^{p,q}(X) := H^{p,q}(M(X))$ and we refer to it as the étale motivic cohomology of X . We have an isomorphism

$$H^{p,q}(X) \cong H_{\text{ét}}^p(X, \mathbb{Z}(q))$$

where the last cohomology is, in general, computed by the éh-topology (see [12, §10.2] and cf. [6] and [50, Prop. 1.8 & Def. 3.1]). In particular, if X is smooth $H_{\text{é h}}^p(X, \mathbb{Z}(q)) \cong H_{\text{ét}}^p(X, \mathbb{Z}(q))$.

Note that we also have the triangulated category of motivic complexes without transfers $\text{DA}_{\text{ét}}^{\text{eff}}$ and if we are interested in rational coefficients we may forget transfers or keep the Nisnevich topology as we have equivalences

$$\text{DA}_{\text{ét}, \mathbb{Q}}^{\text{eff}} \cong \text{DM}_{\text{ét}, \mathbb{Q}}^{\text{eff}} \cong \text{DM}_{\text{Nis}, \mathbb{Q}}^{\text{eff}}$$

(see [6], [7, Cor. B.14] and [47, Thm. 14.30]). If we work with rational coefficients, we then have that motivic cohomology $H^{p,q}(X)_{\mathbb{Q}}$ is computed by the cdh-topology, *i.e.*, $H_{\text{é h}}^p(X, \mathbb{Q}(q)) \cong H_{\text{cdh}}^p(X, \mathbb{Q}(q))$, and $H_{\text{cdh}}^p(X, \mathbb{Q}(q)) \cong H_{\text{Zar}}^p(X, \mathbb{Q}(q))$ if X is smooth.

1.1. de Rham regulator

Denote by $\underline{\Omega}$ the object of $\text{DM}_{\text{ét}}^{\text{eff}}$ which represents de Rham cohomology. More precisely we here denote $\underline{\Omega} := \alpha \underline{\Omega}_{\text{Nis}}$ where $\underline{\Omega}_{\text{Nis}}$ is the corresponding object for the Nisnevich topology (see [45, §2.1] and cf. [7, §2.3] without transfers). This latter $\underline{\Omega}_{\text{Nis}}$ is given by the complex of presheaves with transfers that associates to $X \in \text{Sm}_K$ the global sections $\Gamma(X, \Omega_{X/K}^{\bullet})$ of the usual algebraic de Rham complex.

For $M \in \text{DM}_{\text{ét}}^{\text{eff}}$ we shall denote (cf. [44, §6] and [45, Def. 2.1.1 & Lemma 2.1.2])

$$H_{\text{dR}}^p(M) := \text{Hom}_{\text{DM}_{\text{ét}}^{\text{eff}}}(M, \underline{\Omega}[p]).$$

For any algebraic scheme X and $M = M(X)$ we here may also consider the sheafification of $\underline{\Omega}$ for the éh-topology. Actually, we set

$$H_{\text{dR}}^p(X) := H_{\text{dR}}^p(M(X)) \cong H_{\text{é h}}^p(X, \underline{\Omega})$$

(see [12, Prop. 10.2.3]). Remark that this definition is equivalent to the definition of the algebraic de Rham cohomology in [33, Chap. 3] via the h -topology (as one can easily see via blow-up induction [12, Lemma 10.3.1 b]) after [33, Prop. 3.2.4] and [33, Lemma 3.1.14]).

Note that for $q = 0$ we have a canonical map $r^0 : \mathbb{Z}(0) \rightarrow \underline{\Omega}$ yielding a map

$$H^{p,0}(X) \cong H_{\text{é h}}^p(X, \mathbb{Z}) \rightarrow H_{\text{dR}}^p(X) \cong H_{\text{é h}}^p(X, \underline{\Omega}).$$

For $q = 1$ we have $r^1 := d \log : \mathbb{Z}(1) \rightarrow \underline{\Omega}$ in $\text{DM}_{\text{ét}}^{\text{eff}}$ (see [45, Lemme 2.1.3] for the Nisnevich topology and apply α) yielding a map

$$H^{p,1}(X) \cong H_{\text{é h}}^{p-1}(X, \mathbb{G}_m) \rightarrow H_{\text{dR}}^p(X) \cong H_{\text{é h}}^p(X, \underline{\Omega}).$$

Following [45, (2.1.5)] an internal de Rham regulator r^q in $\text{DM}_{\text{ét}}^{\text{eff}}$ for $q \geq 2$ is then obtained as the composition of

$$r^q : \mathbb{Z}(q) \cong \mathbb{Z}(1)^{\otimes q} \xrightarrow{d \log^{\otimes q}} \underline{\Omega}^{\otimes q} \rightarrow \underline{\Omega}. \tag{1.1}$$

For $M \in \text{DM}_{\text{ét}}^{\text{eff}}$, composing a map $M \rightarrow \mathbb{Z}(q)[p]$ with $r^q[p]$ we get an external *de Rham regulator* map

$$r_{\text{dR}}^{p,q} : H^{p,q}(M) \rightarrow H_{\text{dR}}^p(M) \tag{1.2}$$

and in particular for $M = M(X)$ we get

$$r_{\text{dR}}^{p,q} : H^{p,q}(X) \cong H_{\text{éh}}^p(X, \mathbb{Z}(q)) \rightarrow H_{\text{dR}}^p(X) \cong H_{\text{éh}}^p(X, \underline{\Omega}).$$

Note that if X is smooth then $H_{\text{éh}}^p(X, \underline{\Omega}) \cong H_{\text{ét}}^p(X, \underline{\Omega}) \cong H_{\text{Zar}}^p(X, \Omega_X^\bullet)$ coincides with the classical algebraic de Rham cohomology (again, see [12, Prop. 10.2.3] and cf. [33, Prop. 3.2.4]) and we thus obtain $r_{\text{dR}}^{p,q} : H_{\text{ét}}^p(X, \mathbb{Z}(q)) \rightarrow H_{\text{Zar}}^p(X, \Omega_X^\bullet)$ in this case.

1.2. Periods

As soon as we have an embedding $\sigma : K \hookrightarrow \mathbb{C}$ we may consider a Betti realization (e.g. see [45, §3.3] or [5, Def. 2.1]) in the derived category of abelian groups $D(\mathbb{Z})$ as a triangulated functor

$$\beta_\sigma : \text{DM}_{\text{ét}}^{\text{eff}} \rightarrow D(\mathbb{Z}) \tag{1.3}$$

such that $\beta_\sigma(\mathbb{Z}(q)) \cong \mathbb{Z}_{\text{an}}(q) := (2\pi i)^q \mathbb{Z}[0]$. Actually, following Ayoub (see also [7, §2.1.2] and [8, §1.1.2]) if we consider the analogue of the Voevodsky motivic category $\text{DM}_{\text{an}}^{\text{eff}}$ obtained as the full subcategory of $D(\text{Shv}_{\text{an}}^{\text{tr}}(\text{An}_{\mathbb{C}}))$ given by \mathbb{A}_{an}^1 -local complexes, where we here replace smooth schemes Sm_K by the category $\text{An}_{\mathbb{C}}$ of complex analytic manifolds, we get an equivalence

$$\beta : \text{DM}_{\text{an}}^{\text{eff}} \xrightarrow{\sim} D(\mathbb{Z})$$

such that $M_{\text{an}}(X) \rightsquigarrow \text{Sing}_*(X)$ is sent to the singular chain complex of $X \in \text{An}_{\mathbb{C}}$. Moreover there is a natural triangulated functor

$$\sigma : \text{DM}_{\text{ét}}^{\text{eff}} \rightarrow \text{DM}_{\text{an}}^{\text{eff}}$$

such that $M(X) \rightsquigarrow M_{\text{an}}(X_{\text{an}})$ where the analytic space X_{an} is given by the \mathbb{C} -points of the base change $X_{\mathbb{C}}$ of any algebraic scheme X . We then set $\beta_\sigma := \beta \circ \sigma$. Thus it is clear that $\beta_\sigma(\mathbb{Z}[0]) = \beta_\sigma(M(\text{Spec}(K))) = \mathbb{Z}[0]$. Since a K -rational point of X yields $M(X) = \mathbb{Z} \oplus \widetilde{M}(X)$ we also see that $\beta_\sigma(\mathbb{Z}(1)[1]) \cong \beta_\sigma(\widetilde{M}(\mathbb{G}_m)) \cong \beta(\widetilde{M}_{\text{an}}(\mathbb{C}^*)) \cong \mathbb{Z}_{\text{an}}(1)[1]$ and then, $\beta_\sigma(\mathbb{Z}(q)) \cong \mathbb{Z}_{\text{an}}(q)$ in general, as it follows from the compatibility of β_σ with the tensor structures, i.e., we here use the fact that β_σ is unital and monoidal. For $M \in \text{DM}_{\text{ét}}^{\text{eff}}$, we denote

$$H_{\text{an}}^{p,q}(M) := \text{Hom}_{D(\mathbb{Z})}(\beta_\sigma M, \mathbb{Z}_{\text{an}}(q)[p])$$

and we have a *Betti regulator* map

$$r_{\text{an}}^{p,q} : H^{p,q}(M) \rightarrow H_{\text{an}}^{p,q}(M) \tag{1.4}$$

induced by β_σ . In particular, for $M = M(X)$, we obtain from Ayoub’s construction (see also [45, Prop. 4.2.7]):

Lemma 1.2.1. *For any algebraic K -scheme X and any field homomorphism $\sigma : K \hookrightarrow \mathbb{C}$ we have*

$$H_{\text{an}}^{p,q}(X) := \text{Hom}_{D(\mathbb{Z})}(\beta_\sigma M(X), \mathbb{Z}_{\text{an}}(q)[p]) \cong H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$$

and a *Betti regulator* map

$$r_{\text{an}}^{p,q} : H^{p,q}(X) \rightarrow H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)).$$

Recall that the functor β_σ admits a right adjoint $\beta^\sigma : D(\mathbb{Z}) \rightarrow \text{DM}_{\text{ét}}^{\text{eff}}$ (see [8, Def. 1.7]). Note that the Betti regulator (1.4) is just given by composition with the unit

$$r_\sigma^q : \mathbb{Z}(q) \rightarrow \beta^\sigma \beta_\sigma(\mathbb{Z}(q)) \tag{1.5}$$

of the adjunction. Actually, by making use of the classical Poincaré Lemma and Grothendieck comparison theorem ([30, Thm. 1’]) we get:

Lemma 1.2.2 (Ayoub). *There is a canonical quasi-isomorphism*

$$\varpi^q : \beta^\sigma \beta_\sigma(\mathbb{Z}(q)) \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\text{q.i.}} \underline{\Omega} \otimes_K \mathbb{C}$$

whose composition with r_σ^q in (1.5) is the regulator r^q in (1.1) after tensoring with \mathbb{C} .

Proof. See [7, Cor. 2.89 & Prop. 2.92] and also [9, §3.5]. \square

Remark 1.2.3. Note that applying β_σ to ϖ^q we obtain a quasi-isomorphism $\beta_\sigma(\varpi^q)$ such that

$$\begin{array}{ccc} \mathbb{C} \cong \beta_\sigma(\mathbb{Z}(q)) \otimes_{\mathbb{Z}} \mathbb{C} & \xrightarrow{\beta_\sigma(r_\sigma^q)} & \beta_\sigma \beta^\sigma \beta_\sigma(\mathbb{Z}(q)) \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\beta_\sigma(\varpi^q)} \beta_\sigma(\underline{\Omega}) \otimes_K \mathbb{C} \\ & \searrow \beta_\sigma(r_\sigma^q)_{\mathbb{C}} & \nearrow \beta_\sigma(r_\sigma^q)_{\mathbb{C}} \end{array}$$

where $\beta_\sigma(r_\sigma^q)_{\mathbb{C}}$ is a split injection but it is not a quasi-isomorphism (cf. [45, §4.1]).

For $M \in \text{DM}_{\text{ét}}^{\text{eff}}$, by composition with ϖ^q we get a period isomorphism

$$\varpi_M^{p,q} : H_{\text{an}}^{p,q}(M) \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\simeq} H_{\text{dR}}^p(M) \otimes_K \mathbb{C}.$$

Definition 1.2.4. For any scheme X we shall call *period isomorphism* the \mathbb{C} -isomorphism

$$\varpi_X^{p,q} : H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)) \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\simeq} H_{\text{dR}}^p(X) \otimes_K \mathbb{C}$$

obtained by setting $\varpi_X^{p,q} := \varpi_{M(X)}^{p,q}$ as above. We shall denote $\eta_X^{p,q} := (\varpi_X^{p,q})^{-1}$ the inverse of the period isomorphism.

We also get the following compatibility.

Proposition 1.2.5. For $M \in \text{DM}_{\text{ét}}^{\text{eff}}$ along with a fixed embedding $\sigma : K \hookrightarrow \mathbb{C}$ the inverse of the period isomorphism $\varpi_M^{p,q}$ above induces a commutative diagram

$$\begin{array}{ccccc} H^{p,q}(M) & \xrightarrow{r_{\text{an}}^{p,q}} & H_{\text{an}}^{p,q}(M) & & \\ \downarrow r_{\text{dR}}^{p,q} & & \searrow \iota_{\text{an}}^{p,q} & & \\ H_{\text{dR}}^p(M) & \longrightarrow & H_{\text{dR}}^p(M) \otimes_K \mathbb{C} & \xrightarrow{\simeq} & H_{\text{an}}^{p,q}(M) \otimes_{\mathbb{Z}} \mathbb{C} \\ & \searrow \iota_{\text{dR}}^{p,q} & & & \end{array}$$

where $\iota_{\text{dR}}^{p,q}$ and $\iota_{\text{an}}^{p,q}$ are the canonical mappings given by tensoring with \mathbb{C} .

Proof. This easily follows from Lemma 1.2.2. In fact, by construction, the claimed commutative diagram can be translated into the following commutative square:

$$\begin{array}{ccc} \text{Hom}_{\text{DM}^{\text{eff}}}(M, \mathbb{Z}(q)[p]) & \xrightarrow{\iota_{\text{an}}^{p,q} \circ r_{\text{an}}^{p,q}} & \text{Hom}_{\text{DM}^{\text{eff}}}(M, \beta^\sigma \beta_\sigma \mathbb{Z}(q)[p])_{\mathbb{C}} \quad \square \\ \downarrow r_{\text{dR}}^{p,q} := r^q[p] \circ - & \searrow \iota_{\text{dR}}^{p,q} & \downarrow \varpi^q[p] \circ - \\ \text{Hom}_{\text{DM}^{\text{eff}}}(M, \underline{\Omega}[p]) & \xrightarrow{- \otimes_K \mathbb{C}} & \text{Hom}_{\text{DM}^{\text{eff}}}(M, \underline{\Omega}[p])_{\mathbb{C}}. \end{array}$$

Corollary 1.2.6. Let X be an algebraic K -scheme along with a fixed embedding $\sigma : K \hookrightarrow \mathbb{C}$. The period isomorphism $\varpi_X^{p,q}$ above induces a commutative square

$$\begin{array}{ccc} H^{p,q}(X) & \xrightarrow{r_{\text{an}}^{p,q}} & H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)) \\ \downarrow r_{\text{dR}}^{p,q} & & \downarrow \iota_{\text{an}}^{p,q} \\ H_{\text{dR}}^p(X) & \xrightarrow{\iota_{\text{dR}}^{p,q}} & H^p(X_{\text{an}}, \mathbb{C}). \end{array}$$

Note that from Corollary 1.2.6 we get a refinement of the Betti regulator.

Definition 1.2.7. Define the *algebraic* singular cohomology classes as the elements of the subgroup $H_{\text{alg}}^{p,q}(X) := \text{Im } r_{\text{an}}^{p,q} \subseteq H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ given by the image of the motivic cohomology under the Betti regulator $r_{\text{an}}^{p,q}$.

Define the ϖ -*algebraic* singular cohomology classes by the subgroup

$$H_{\varpi}^{p,q}(X) := H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)) \subseteq H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$$

where \cap means that we take elements in $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ which are given by the inverse image (under $\iota_{\text{an}}^{p,q}$) of elements in $H_{\text{dR}}^p(X)$ regarded (under $\iota_{\text{dR}}^{p,q}$) inside $H^p(X_{\text{an}}, \mathbb{C})$ via the isomorphism $\varpi_X^{p,q}$ above.

The groups $H_{\varpi}^{p,q}(X)$ shall be called *period cohomology* groups and

$$r_{\varpi}^{p,q} : H^{p,q}(X) \rightarrow H_{\varpi}^{p,q}(X)$$

induced by $r_{\text{dR}}^{p,q}$ and $r_{\text{an}}^{p,q}$ shall be called the *period regulator*.

We get that:

Corollary 1.2.8. $H_{\text{alg}}^{p,q}(X) \subseteq H_{\varpi}^{p,q}(X)$.

For example, all torsion cohomology classes are ϖ -algebraic: we shall see in Lemma 1.4.2 that they are also algebraic.

In particular, if $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ is all algebraic, *i.e.*, the Betti regulator $r_{\text{an}}^{p,q}$ is surjective, then the canonical embedding $\iota_{\text{an}}^{p,q}$ of singular cohomology $H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(q))$ in the \mathbb{C} -vector space $H^p(X_{\text{an}}, \mathbb{C})$ factors through an embedding of $H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(q))$ into the K -vector space $H_{\text{dR}}^p(X)$. If $K = \overline{\mathbb{Q}}$ this rarely happens. For example, if $p = 0$ it happens only if $q = 0$ and in this case $r_{\varpi}^{0,q}$ is always surjective (as $H_{\varpi}^{0,q}(X) = 0$ for $q \neq 0$).

1.3. Period conjecture for motivic cohomology

Over $K = \overline{\mathbb{Q}}$ it seems reasonable to make the conjecture that all ϖ -algebraic classes are algebraic, *i.e.*, to conjecture that the period regulator $r_{\varpi}^{p,q}$ is surjective. In other words we may say that the period conjecture for motivic cohomology holds for X , in degree p and twist q , if

$$H_{\text{alg}}^{p,q}(X) = H_{\varpi}^{p,q}(X). \tag{1.6}$$

Over a number field we may expect that this holds rationally. If (1.6) holds we also have that $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ modulo torsion embeds into $H_{\text{dR}}^p(X)$ if and only if $H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))$ is all algebraic. Note that using Proposition 1.2.5 we can define $H_{\varpi}^{p,q}(M)$ providing a version of the period conjecture for any object $M \in \text{DM}_{\text{ét}}^{\text{eff}}$.

Proposition 1.3.1. *For any $q \geq 0$ the period conjecture (1.6) holds true for X , in degree p and twist r , if and only if it holds true for $M(X)(q)$, in degree p and twist $q + r$.*

Proof. By Voevodsky cancellation theorem [51] we have that twisting by q in motivic cohomology $H^{p,r}(X) \xrightarrow{\sim} H^{p,q+r}(M(X)(q))$ is an isomorphism of groups. If $M = M(X)(q)$ with $q \geq 0$ we also get $H_{\varpi}^{p,r}(X) \xrightarrow{\sim} H_{\varpi}^{p,q+r}(M(X)(q))$ canonically by twisting. In fact, we have a diagram induced by twisting

$$\begin{CD}
 H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(r)) \otimes \mathbb{C} @>{q_{\mathbb{C}}}>> H_{\text{an}}^{p,q+r}(M(X)(q)) \otimes_{\mathbb{Z}} \mathbb{C} \\
 @V{\omega_X^{p,r}}VV @VV{\varpi_{M(X)(q)}^{p,q+r}}V \\
 H_{\text{dR}}^p(X) \otimes_K \mathbb{C} @>{q_{\text{dR}}}>> H_{\text{dR}}^p(M(X)(q)) \otimes_K \mathbb{C}
 \end{CD}$$

where $q_{\mathbb{C}} := q \otimes \mathbb{C}$ is the \mathbb{C} -isomorphism given by the canonical integrally defined mapping $q: H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(r)) \xrightarrow{\sim} H_{\text{an}}^{p,q+r}(M(X)(q))$ which is sending a p -th cohomology class regarded as a map $\beta_{\sigma} M(X) = \text{Sing}_*(X_{\text{an}}) \rightarrow \mathbb{Z}_{\text{an}}(r)[p]$ in $D(\mathbb{Z})$ to the q -twist $\beta_{\sigma} M(X)(q) = \text{Sing}_*(X_{\text{an}})(q) \rightarrow \mathbb{Z}_{\text{an}}(q+r)[p]$. Similarly, the \mathbb{C} -isomorphism q_{dR} is induced by twisting, since $\underline{\Omega}(-q) \xrightarrow{q.i.} \underline{\Omega}$ is a canonical isomorphism in $\text{DM}_{\text{ét}}^{\text{eff}}$ and the claim follows. \square

For X smooth we have that $H^{p,q}(X) \cong H_{\text{ét}}^p(X, \mathbb{Z}(q))$ and with rational coefficients we have that $H_{\text{ét}}^p(X, \mathbb{Q}(q)) \cong CH^q(X, 2q-p)_{\mathbb{Q}}$. In particular, if X is smooth and $p = 2q$ we get that $r_{\varpi}^{2q,q}$ is the modern refinement of the classical cycle class map with rational coefficients

$$r_{\varpi}^{2q,q} = c\ell_{\varpi}^q : CH^q(X)_{\mathbb{Q}} \rightarrow H_{\varpi}^{2q,q}(X)_{\mathbb{Q}} \tag{1.7}$$

for codimension q cycles on X considered in [17]. In this case, the period conjecture (1.6) with rational coefficients coincides with the classical Grothendieck period conjecture for algebraic cycles: see [17, §1.1.3] and [17, Prop. 2.13-14] comparing it with the conjecture on torsors of periods.

Remark 1.3.2. For $K = \mathbb{C}$ we may also think to refine the Hodge conjecture as previously hinted by Beilinson, conjecturing the surjectivity of

$$r_{\text{Hodge}}^{p,q} : H^{p,q}(X)_{\mathbb{Q}} \rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^p(X)(q)).$$

However, such a generalization doesn't hold, in general, e.g. see [20].

1.4. Torsion cohomology classes are algebraic

Consider $\mathbb{Z}/n(q) := \mathbb{Z}(q) \otimes \mathbb{Z}/n$. By Suslin-Voevodsky rigidity we have a quasi-isomorphism of complexes of étale sheaves $\mu_n^{\otimes q} \rightarrow \mathbb{Z}/n(q)$ yielding $H_{\text{ét}}^p(X, \mathbb{Z}/n(q)) \cong$

$H_{\text{ét}}^p(X, \mu_n^{\otimes q}) \cong H_{\text{éh}}^p(X, \mu_n^{\otimes q})$. For a proof of this key result see [47, Thm. 10.2 & Prop. 10.7] for X smooth and make use of [12, Prop. 12.1.1] to get it in general.

Lemma 1.4.1. *For any algebraic scheme X over $K = \overline{K} \hookrightarrow \mathbb{C}$ we have $H_{\text{éh}}^p(X, \mathbb{Z}/n(q)) \cong H^p(X_{\text{an}}, \mathbb{Z}/n)$.*

Proof. As étale cohomology of $\mu_n^{\otimes q}$ is invariant under the extension $\sigma : K \hookrightarrow \mathbb{C}$ of algebraically closed fields we obtain the claimed comparison from the classical comparison result after choosing a root of unity. \square

We then have (cf. [49, Prop. 3.1]):

Lemma 1.4.2. *The regulator $r_{\varpi}^{p,q} |_{\text{tor}} : H^{p,q}(X)_{\text{tor}} \rightarrow H_{\varpi}^{p,q}(X)_{\text{tor}}$ is surjective on torsion and $r_{\varpi}^{p,q} \otimes \mathbb{Q}/\mathbb{Z} : H^{p,q}(X) \otimes \mathbb{Q}/\mathbb{Z} \hookrightarrow H_{\varpi}^{p,q}(X) \otimes \mathbb{Q}/\mathbb{Z}$ is injective.*

Proof. By construction, for any positive integer n , comparing the usual universal coefficient exact sequences, we have the following commutative diagram with exact rows

$$\begin{array}{ccccccc}
 0 \rightarrow & H_{\text{éh}}^p(X, \mathbb{Z}(q))/n & \longrightarrow & H_{\text{éh}}^p(X, \mathbb{Z}/n(q)) & \longrightarrow & {}_n H_{\text{éh}}^{p+1}(X, \mathbb{Z}(q)) & \rightarrow 0 \\
 & \downarrow r_{\text{an}}^{p,q}/n & & \parallel \text{1.4.1} & & \downarrow {}_n r_{\text{an}}^{p+1,q} & \\
 0 \rightarrow & H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}})/n & \longrightarrow & H^p(X_{\text{an}}, \mathbb{Z}/n) & \longrightarrow & {}_n H^{p+1}(X_{\text{an}}, \mathbb{Z}_{\text{an}}) & \rightarrow 0.
 \end{array}$$

Passing to the direct limit on n we easily get the claim. In fact, ${}_n H_{\varpi}^{p,q}(X) = {}_n H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}})$ and $r_{\text{an}}^{p,q}/n$ factors through $r_{\varpi}^{p,q}/n$. \square

Lemma 1.4.3. *We have that $r_{\varpi}^{p,q} \otimes \mathbb{Q}$ is surjective if and only if $r_{\varpi}^{p,q}$ is surjective; moreover, if this is the case $r_{\varpi}^{p,q} \otimes \mathbb{Q}/\mathbb{Z}$ is an isomorphism.*

Proof. This follows from a simple diagram chase. \square

In the situation that $H^{p,q}(X) \otimes \mathbb{Q}/\mathbb{Z} = 0$ the period conjecture for motivic cohomology (1.6) is then equivalent to

$$H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(q)) = 0. \tag{1.8}$$

In particular:

Proposition 1.4.4. *If X is smooth then (1.6) for $p \notin [q, 2q]$ is equivalent to (1.8). If X is smooth and proper then (1.6) is equivalent to the surjectivity of cl_{ϖ}^q in (1.7) for $p = 2q$ and to the vanishing (1.8) for $p \neq 2q$.*

Proof. In fact, by the Appendix A, Theorem A.1.3, we have that for $p \notin [q, 2q]$ the group $H^{p,q}(X)$ is an extension of torsion by divisible groups so that $H^{p,q}(X) \otimes \mathbb{Q}/\mathbb{Z} = 0$. If X is proper the latter vanishing holds true for all $p \neq 2q$. \square

Proposition 1.4.4 explains some weight properties related to the Grothendieck period conjecture, weight arguments which are also considered in [17].

Remark 1.4.5. For $K = \mathbb{C}$ we have that $r_{\beta}^{p,q} |_{\text{tor}}: H^{p,q}(X)_{\text{tor}} \rightarrow H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))_{\text{tor}}$ is surjective (as also remarked in [49] for X smooth projective): torsion motivic cohomology classes supply the defect of algebraic cycles providing the missing torsion algebraic cycles. In fact, from the well known Atiyah-Hirzebruch-Totaro counterexamples to the integral Hodge conjecture we know that $c\ell^p: CH^p(X) \rightarrow H^{2p}(X_{\text{an}}, \mathbb{Z}_{\text{an}}(p))$ cannot be surjective on torsion for $p \geq 2$ in general.

2. Periods of 1-motives: fullness of Betti-de Rham realizations

Let ${}^t\mathcal{M}_1(K)$ be the abelian category of 1-motives with torsion over K (see [12, App. C]). We shall drop the reference to K if it is clear from the context. We shall denote

$$M_K = [u_K: L_K \rightarrow G_K] \in {}^t\mathcal{M}_1(K)$$

a 1-motive with torsion with L_K in degree 0 and G_K in degree 1; for brevity, we shall write $M_K = L_K[0]$ if $G_K = 0$ and $M_K = G_K[-1]$ if $L_K = 0$ and we omit the reference to K if unnecessary. Let $M_{\text{tor}} := [L_{\text{tor}} \cap \text{Ker}(u) \rightarrow 0]$ be the torsion part of M_K , let $M_{\text{fr}} := [L/L_{\text{tor}} \rightarrow G/u(L_{\text{tor}})]$ be the free part of M_K , and let $M_{\text{tf}} := [L/L_{\text{tor}} \cap \text{Ker}(u) \rightarrow G]$ be the torsion free part of M_K . There are short exact sequences of complexes

$$0 \rightarrow M_{\text{tor}} \rightarrow M_K \rightarrow M_{\text{tf}} \rightarrow 0 \tag{2.1}$$

and

$$0 \rightarrow [F = F] \rightarrow M_{\text{tf}} \rightarrow M_{\text{fr}} \rightarrow 0, \tag{2.2}$$

where $F = L_{\text{tor}}/L_{\text{tor}} \cap \text{Ker}(u)$. Let M_{ab} denote the 1-motive with torsion $[L \rightarrow G/\mathbb{T}]$ where \mathbb{T} is the maximal subtorus of G . Recall (see [12, Prop. C.7.1]) that the canonical functor $\mathcal{M}_1 \rightarrow {}^t\mathcal{M}_1$ from Deligne 1-motives admits a left adjoint/left inverse given by $M \rightsquigarrow M_{\text{fr}}$.

Any 1-motive $M = [L \rightarrow G]$ is canonically endowed with an increasing filtration of sub-1-motives, the *weight filtration*, defined as follows:

$$W_i(M) = \begin{cases} M & \text{if } i \geq 0 \\ G[-1] & \text{if } i = -1 \\ \mathbb{T}[-1] & \text{if } i = -2 \\ 0 & \text{if } i \leq -3 \end{cases} \tag{2.3}$$

with \mathbf{T} the maximal subtorus of \mathbf{G} . We have that $D^b({}^t\mathcal{M}_1) \cong D^b(\mathcal{M}_1)$ (see [12, Thm. 1.11.1]) and that there is a canonical embedding (see [12, Def 2.7.1])

$$\text{Tot} : D^b(\mathcal{M}_1) \hookrightarrow \text{DM}_{\text{ét}}^{\text{eff}} \tag{2.4}$$

so that we can also regard 1-motives as motivic complexes of étale sheaves. The restriction of the Betti realization β_σ in (1.3) can be described explicitly for 1-motives via Deligne’s Hodge realization (see [12, Thm. 15.4.1]). Similarly, the restriction of the de Rham realization in [45] can be described via Deligne’s de Rham realization as follows.

2.1. de Rham realization

Let K be a field of characteristic zero and let $\mathbf{M}_K = [\mathbf{u}_K : \mathbf{L}_K \rightarrow \mathbf{G}_K] \in {}^t\mathcal{M}_1(K)$ be a 1-motive with torsion over K . Note that for $\mathbf{M}_K^\natural := [\mathbf{u}_K^\natural : \mathbf{L}_K \rightarrow \mathbf{G}_K^\natural]$ the universal \mathbb{G}_a -extension of \mathbf{M}_K we have

$$0 \rightarrow \mathbb{V}(\mathbf{M}) \rightarrow \mathbf{M}_K^\natural \xrightarrow{\rho^{\mathbf{M}}} \mathbf{M}_K \rightarrow 0$$

where $\mathbb{V}(\mathbf{M}) := \text{Ext}(\mathbf{M}_K, \mathbb{G}_a)^\vee$. The existence of universal extensions is well-known when \mathbf{L}_K is torsion-free; for the general case see [11, Proposition 2.2.1]. Recall (see [23, §10.1.7]) the following

Definition 2.1.1. The de Rham realization of \mathbf{M}_K is

$$T_{\text{dR}}(\mathbf{M}_K) := \text{Lie}(\mathbf{G}_K^\natural)$$

as a K -vector space.

Remark 2.1.2. Note that $\rho^{\mathbf{M}} = (id_{\mathbf{L}}, \rho_{\mathbf{G}})$ where $\rho_{\mathbf{G}} : \mathbf{G}_K^\natural \twoheadrightarrow \mathbf{G}_K$ is a quotient and $\text{Ker } \rho_{\mathbf{G}} = \mathbb{V}(\mathbf{M})$ so that \mathbf{G}_K is the semiabelian quotient of \mathbf{G}_K^\natural and \mathbf{u}_K^\natural is a canonical lifting of \mathbf{u}_K , i.e., $\mathbf{u}_K = \rho_{\mathbf{G}} \circ \mathbf{u}_K^\natural$. Further $\mathbb{V}(\mathbf{M}) \subseteq T_{\text{dR}}(\mathbf{M}_K)$ is also the kernel of the morphism

$$d\rho_{\mathbf{G}} : \text{Lie}(\mathbf{G}_K^\natural) \rightarrow \text{Lie}(\mathbf{G}_K)$$

induced by $\rho_{\mathbf{G}}$, so that $T_{\text{dR}}(\mathbf{M}_K)$ together with the K -subspace $\mathbb{V}(\mathbf{M})$ can be regarded as a filtered K -vector space. This datum is called the *Hodge filtration* of $T_{\text{dR}}(\mathbf{M}_K)$.

The algebraic K -group G_K^{\natural} fits in the following diagram [14, (2.15)]

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathbb{V}(A) & \longrightarrow & A^{\natural} \times_A G & \longrightarrow & G \longrightarrow 0 \\
 & & \downarrow i & & \downarrow & & \parallel \\
 0 & \longrightarrow & \mathbb{V}(M) & \longrightarrow & G^{\natural} & \xrightarrow{\rho_G} & G \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \mathbb{V}(L) & \xlongequal{\quad} & L \otimes G_a & &
 \end{array}$$

where we have omitted subscripts K and written $\mathbb{V}(A)$ for $\mathbb{V}(A[-1]) = \mathbb{V}(G[-1])$.

Lemma 2.1.3. For $K \subset K'$ we have a natural isomorphism

$$(M_K^{\natural})_{K'} \cong (M_{K'})^{\natural}.$$

2.2. Base change to \mathbb{C} and periods

Consider K a subfield of \mathbb{C} and let $M_{\mathbb{C}} = [u_{\mathbb{C}} : L_{\mathbb{C}} \rightarrow G_{\mathbb{C}}]$ be the base change of M_K to \mathbb{C} . Let $T_{\mathbb{Z}}(M_{\mathbb{C}})$ be the finitely generated abelian group in the usual Deligne-Hodge realization of $M_{\mathbb{C}}$ (see [23, 10.1.3] and [13, §1]) given by the pull-back

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_1(G_{\mathbb{C}}) & \longrightarrow & \text{Lie}(G_{\mathbb{C}}) & \xrightarrow{\text{exp}} & G_{\mathbb{C}} \longrightarrow 0 \\
 & & \parallel & & \uparrow \tilde{u}_{\mathbb{C}} & & \uparrow u_{\mathbb{C}} \\
 0 & \longrightarrow & T_{\mathbb{Z}}(G_{\mathbb{C}}) & \longrightarrow & T_{\mathbb{Z}}(M_{\mathbb{C}}) & \xrightarrow{\widehat{\text{exp}}} & L_{\mathbb{C}} \longrightarrow 0
 \end{array}$$

where for brevity $T_{\mathbb{Z}}(G_{\mathbb{C}})$ denotes $T_{\mathbb{Z}}(G_{\mathbb{C}}[-1])$ which by definition is $H_1(G_{\mathbb{C}})$. After base change to \mathbb{C} and Lemma 2.1.3 we then get $(M_K^{\natural})_{\mathbb{C}} \cong (M_{\mathbb{C}})^{\natural}$ hence an isomorphism

$$\iota : T_{\text{dR}}(M_{\mathbb{C}}) \xrightarrow{\cong} T_{\text{dR}}(M_K) \otimes_K \mathbb{C} \tag{2.5}$$

and a commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_1(G_{\mathbb{C}}) & \longrightarrow & \text{Lie}(G_{\mathbb{C}}) & \xrightarrow{\text{exp}} & G_{\mathbb{C}} \longrightarrow 0 \\
 & & \parallel & & \uparrow d\rho_G & & \uparrow \rho_G \\
 0 & \longrightarrow & H_1(G_{\mathbb{C}}^{\natural}) & \longrightarrow & \text{Lie}(G_{\mathbb{C}}^{\natural}) & \xrightarrow{\text{exp}} & G_{\mathbb{C}}^{\natural} \longrightarrow 0 \\
 & & \parallel & & \uparrow \tilde{u}_{\mathbb{C}} & & \uparrow u_{\mathbb{C}} \\
 0 & \longrightarrow & T_{\mathbb{Z}}(G_{\mathbb{C}}) & \longrightarrow & T_{\mathbb{Z}}(M_{\mathbb{C}}) & \xrightarrow{\widehat{\text{exp}}} & L_{\mathbb{C}} \longrightarrow 0
 \end{array} \tag{2.6}$$

where the dotted arrow exists by definition of $T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}})$ and the fact that the upper right-hand square is cartesian. Hence also the lower right-hand square is cartesian and the sequence on the bottom is equivalently obtained by pull-back of the upper sequence via $u_{\mathbb{C}}$ or of the sequence in the middle via $u_{\mathbb{C}}^{\natural}$. Further $T_{\mathbb{Z}}(\mathbf{G}_{\mathbb{C}})$ is identified with the kernel of both exponential maps and the dotted arrow gives a homomorphism $T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}) \rightarrow T_{\text{dR}}(\mathbf{M}_{\mathbb{C}}) := \text{Lie}(\mathbf{G}_{\mathbb{C}}^{\natural})$. Finally note that the weight filtration in (2.3) gives a filtration on $T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}})$: the immersion $\mathbf{T}_{\mathbb{C}} \rightarrow \mathbf{G}_{\mathbb{C}}$ gives an inclusion $T_{\mathbb{Z}}(\mathbf{T}_{\mathbb{C}}) = H_1(\mathbf{T}_{\mathbb{C}}) \subseteq H_1(\mathbf{G}_{\mathbb{C}}) = T_{\mathbb{Z}}(\mathbf{G}_{\mathbb{C}})$ while $T_{\mathbb{Z}}(\mathbf{G}_{\mathbb{C}}) \subseteq T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}})$ comes from the previous diagram.

Definition 2.2.1. The homomorphism of *periods* is the unique homomorphism

$$\varpi_{\mathbf{M},\mathbb{Z}} : T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}) \rightarrow T_{\text{dR}}(\mathbf{M}_K) \otimes_K \mathbb{C}$$

that yields $d\rho_{\mathbf{G}} \circ \varpi_{\mathbf{M},\mathbb{Z}} = \tilde{u}_{\mathbb{C}}$ and $\exp \circ \varpi_{\mathbf{M},\mathbb{Z}} = u_{\mathbb{C}}^{\natural} \circ \widetilde{\text{exp}}$ under the identification given by the isomorphism ι in (2.5).

Note that $\tilde{u}_{\mathbb{C}}$ is the pull-back of $u_{\mathbb{C}}$ along \exp and for $x \in L_{\mathbb{C}}$ we may pick $\widetilde{\log}(x) \in T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}})$, i.e., such that $\widetilde{\text{exp}}(\widetilde{\log}(x)) = x$. We then get

$$u_{\mathbb{C}}(x) = \exp(\tilde{u}_{\mathbb{C}}(\widetilde{\log}(x))) = \exp(d\rho_{\mathbf{G}}(\varpi_{\mathbf{M},\mathbb{Z}}(\widetilde{\log}(x)))) \tag{2.7}$$

Theorem 2.2.2. *The induced \mathbb{C} -linear mapping*

$$\varpi_{\mathbf{M},\mathbb{C}} : T_{\mathbb{C}}(\mathbf{M}_{\mathbb{C}}) := T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}) \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow T_{\text{dR}}(\mathbf{M}_K) \otimes_K \mathbb{C}$$

is an isomorphism.

Proof. Making use of the identification in (2.5) we are left to see that it holds true for $K = \mathbb{C}$. The case of L without torsion is treated by Deligne [23, 10.1.8]. Actually, an easy proof can be given by dévissage to the case of lattices, tori and abelian varieties. For the general case note that $\varpi_{\mathbf{M},\mathbb{C}} = \varpi_{\mathbf{M}_{\text{tf}},\mathbb{C}}$ by (2.1). Indeed $T_{\text{dR}}(\mathbf{M}_{\text{tor}}) = 0$ and the kernel of the canonical morphism $T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}) \rightarrow T_{\mathbb{Z}}(\mathbf{M}_{\text{tf},\mathbb{C}})$ is torsion. Further by (2.2) the map $T_{\mathbb{Z}}(\mathbf{M}_{\text{tf},\mathbb{C}}) \rightarrow T_{\mathbb{Z}}(\mathbf{M}_{\text{fr},\mathbb{C}})$ is an isomorphism and we have an exact sequence

$$0 \rightarrow [F = F] \rightarrow \mathbf{M}_{\text{tf}}^{\natural} \rightarrow \mathbf{M}_{\text{fr}}^{\natural} \rightarrow 0$$

so that the canonical morphism $T_{\text{dR}}(\mathbf{M}_{\text{tf}}) \rightarrow T_{\text{dR}}(\mathbf{M}_{\text{fr}})$ is an isomorphism too. Hence $\varpi_{\mathbf{M}_{\text{tf}},\mathbb{C}} = \varpi_{\mathbf{M}_{\text{fr}},\mathbb{C}}$. We conclude that $\varpi_{\mathbf{M},\mathbb{C}} = \varpi_{\mathbf{M}_{\text{fr}},\mathbb{C}}$ and the latter is an isomorphism since \mathbf{M}_{fr} is a Deligne 1-motive. \square

Examples 2.2.3. • If $\mathbf{M}_K = [0 \rightarrow \mathbf{G}_m]$, then $T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}) = \mathbb{Z}$ and the first and second rows in (2.6) are given by $0 \rightarrow \mathbb{Z} \xrightarrow{2\pi i} \mathbb{C} \xrightarrow{\text{exp}} \mathbb{C}^* \rightarrow 0$. Hence $\varpi_{\mathbf{M},\mathbb{Z}} = \tilde{u}_{\mathbb{C}} : \mathbb{Z} \rightarrow \mathbb{C}, x \mapsto 2\pi ix$ and $\varpi_{\mathbf{M},\mathbb{C}} : \mathbb{C} \rightarrow \mathbb{C}, z \mapsto 2\pi iz$.

- If $M_K = [L_K \rightarrow 0]$, then $T_{\mathbb{Z}}(M_{\mathbb{C}}) = L_{\mathbb{C}}$ and $T_{\text{dR}}(M_{\mathbb{C}}) = L_{\mathbb{C}} \otimes_{\mathbb{Z}} \mathbb{C}$ and the map $\varpi_{M, \mathbb{Z}}$ is the homomorphism $L_{\mathbb{C}} \rightarrow L_{\mathbb{C}} \otimes_{\mathbb{Z}} \mathbb{C}, x \mapsto x \otimes 1$ and $\varpi_{M, \mathbb{C}}$ is the identity map.

- If $M_K = [\mathbf{u}: \mathbb{Z} \rightarrow \mathbb{G}_m]$ with $\mathbf{u}(1) = a \in K^*$, then $\mathbb{G}^{\mathbb{Z}} = \mathbb{G}_m \times \mathbb{G}_a$. Once fixed a complex logarithm $\log a$ of a we can construct an isomorphism $\mathbb{Z} \times \mathbb{Z} \rightarrow T_{\mathbb{Z}}(M_{\mathbb{C}}) = \{(z, y) \in \mathbb{C} \times \mathbb{Z} \mid \exp(z) = a^y\}$ that maps the pair (k, y) to $(y \log a + 2\pi i k, y)$. The homomorphism of periods becomes then the map $\varpi_{M, \mathbb{Z}}: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{C} \times \mathbb{C}$ that sends (k, y) to $(y \log a + 2\pi i k, y)$.

Note that over \mathbb{C} the Hodge filtration $\mathbb{V}(M) \subseteq T_{\text{dR}}(M_K)$ of Remark 2.1.2 is obtained from the Hodge filtration of $T_{\mathbb{C}}(M_{\mathbb{C}})$ via $\varpi_{M, \mathbb{C}}$.

2.3. Periods and transcendence

The proof of Theorem 2.7.1, which is the main outcome of the second section of this paper, makes use of deep results of transcendence theory that we recall below. First consider [43, Theorem 2].

Theorem 2.3.1. *Let A_K be an abelian variety of dimension d over $K = \overline{\mathbb{Q}}$. Let $\Theta: \mathbb{C}^d \rightarrow A_{\text{an}}$ be the homomorphism given by the theta functions, inducing an isomorphism of the complex torus onto A_{an} . Assume that the derivations $\partial/\partial z_i, (i = 1, \dots, d)$ are defined over K . If $\alpha = (\alpha_i) \in \mathbb{C}^d$ is a complex vector $\neq 0$ such that all α_i lie in K , then $\Theta(\alpha)$ is transcendental over K . In particular, the periods are transcendental.*

It can be generalized to semiabelian varieties as follows.

Theorem 2.3.2. *Let G_K be a semiabelian variety over $K = \overline{\mathbb{Q}}$. If $0 \neq x \in \text{Lie}(G_K)$, then $\exp(x) \in G_{\mathbb{C}}(\mathbb{C})$ is transcendental over K . In particular, $\text{Lie}(G_K) \cap \text{Ker}(\exp) = \{0\}$ in $\text{Lie}(G_{\mathbb{C}})$.*

Proof. The assertion is known if $G_K = \mathbb{G}_{m, K}^d$ due to fundamental work of Hermite and Lindemann on the transcendence of e^{β} for β a non zero algebraic number. The case G_K an abelian variety is Theorem 2.3.1. The general case follows then by dévissage. \square

This type of results has been further generalised by Waldschmidt ([54, Thm. 5.2.1]):

Theorem 2.3.3. *Let G be a commutative connected algebraic group over $K = \overline{\mathbb{Q}}$. Let $\varphi: \mathbb{C}^n \rightarrow G_{\text{an}}$ be an analytic map such that the induced morphism on Lie algebras arises from a homomorphism of K -vector spaces $K^n \rightarrow \text{Lie } G$. Let $\Gamma \subset \mathbb{C}^n$ be a subgroup containing n elements which are \mathbb{C} -linearly independent and such that $\varphi(\Gamma) \subset G(K)$. Then the algebraic dimension of φ , i.e. the dimension of the Zariski closure of the image of φ , is $\leq n$.*

This implies the following result [17, Thm. 3.1], which is one of the technical inputs for the proof of our Theorem 2.7.1

Theorem 2.3.4. *Let G_K, H_K be two connected commutative algebraic groups over $K = \overline{\mathbb{Q}}$. If the group $H_1(G_{\mathbb{C}})$ generates $\text{Lie}(G_{\mathbb{C}})$ as a complex vector space, then the map*

$$\text{Lie}: \text{Hom}_{K\text{-gr}}(G_K, H_K) \rightarrow \{\psi \in \text{Hom}_K(\text{Lie } G_K, \text{Lie } H_K) \mid \psi_{\mathbb{C}}(H_1(G_{\mathbb{C}})) \subseteq H_1(H_{\mathbb{C}})\}$$

is an isomorphism of \mathbb{Z} -modules.

The condition on the group $H_1(G_{\mathbb{C}})$ is satisfied whenever $G_{\mathbb{C}}$ is a semiabelian variety (or its universal vectorial extension) since $T_{\mathbb{Z}}(G_{\mathbb{C}}) = H_1(G_{\mathbb{C}})$ generates $T_{\text{dR}}(G_{\mathbb{C}}) = \text{Lie}(G_{\mathbb{C}}^{\text{sa}})$ and hence $\text{Lie}(G_{\mathbb{C}})$. Note that Theorem 2.2.2 is a generalization of this fact. For completeness we also cite the following result [56, Thm 1], which is a consequence of the celebrated analytic subgroup theorem of Wüstholz. It implies the result of Waldschmidt is a special case. We will use it to give an alternative proof of Theorem 2.7.1:

Theorem 2.3.5. *Let W be a commutative connected algebraic group over $\overline{\mathbb{Q}}$. Let S be a subset of $\exp^{-1}(W(\overline{\mathbb{Q}}))$ and let $V \subset \text{Lie } W$ be the smallest $\overline{\mathbb{Q}}$ -vector subspace whose \mathbb{C} -span contains S . Then, there exists a connected algebraic subgroup $Z \subset W$ such that $\text{Lie } Z = V$.*

2.4. Period categories

For a fixed $\sigma : K \hookrightarrow \mathbb{C}$ we consider a homological category for Betti-de Rham realizations as follows. Let $\text{Mod}_{\mathbb{Z},K}$ be the following category: (i) objects are triples $(H_{\mathbb{Z}}, H_K, \omega)$ where $H_{\mathbb{Z}}$ is a finitely generated abelian group, H_K is a finite dimensional K -vector space, and $\omega : H_{\mathbb{Z}} \rightarrow H_K \otimes_K \mathbb{C}$ is a homomorphism of groups; (ii) morphisms $\varphi : (H_{\mathbb{Z}}, H_K, \omega) \rightarrow (H'_{\mathbb{Z}}, H'_K, \omega')$ are pairs $\varphi := (\varphi_{\mathbb{Z}}, \varphi_K)$ where $\varphi_{\mathbb{Z}} : H_{\mathbb{Z}} \rightarrow H'_{\mathbb{Z}}$ is a group homomorphism, $\varphi_K : H_K \rightarrow H'_K$ is a K -linear homomorphism and φ is compatible with ω and ω' , i.e., the following square

$$\begin{CD} H_{\mathbb{Z}} @>\omega>> H_K \otimes_K \mathbb{C} \\ @V\varphi_{\mathbb{Z}}VV @VV\varphi_K \otimes 1_{\mathbb{C}}V \\ H'_{\mathbb{Z}} @>\omega'>> H'_K \otimes_K \mathbb{C} \end{CD} \tag{2.8}$$

commutes. For $H = (H_{\mathbb{Z}}, H_K, \omega)$ in $\text{Mod}_{\mathbb{Z},K}$ let

$$\omega_{\mathbb{C}} : H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow H_K \otimes_K \mathbb{C} \tag{2.9}$$

be the induced \mathbb{C} -linear mapping and denote $\text{Mod}_{\mathbb{Z},K}^{\cong}$ the full subcategory of $\text{Mod}_{\mathbb{Z},K}$ given by those objects such that $\omega_{\mathbb{C}}$ is a \mathbb{C} -isomorphism.

There is a \mathbb{Q} -linear variant $\text{Mod}_{\mathbb{Q},K}$ of this category where objects are $(H_{\mathbb{Q}}, H_K, \omega)$ as above but $H_{\mathbb{Q}}$ is a finite dimensional \mathbb{Q} -vector space. Note that $\text{Mod}_{\mathbb{Q},K} \cong \text{Mod}_{\mathbb{Z},K} \otimes \mathbb{Q}$ is the category $\text{Mod}_{\mathbb{Z},K}$ modulo torsion objects (see [12, B.3] for this notion).

Definition 2.4.1. We shall call $\text{Mod}_{\mathbb{Z},K}^{\cong}$ (resp. $\text{Mod}_{\mathbb{Q},K}^{\cong}$) the category (resp. \mathbb{Q} -linear category) of *homological periods*.

Let $\text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ (resp. $\text{Mod}_{\mathbb{Z},K}^{\cong, \text{tor}}$) be the full subcategory of $\text{Mod}_{\mathbb{Z},K}^{\cong}$ given by those objects H such that $H_{\mathbb{Z}}$ is free (resp. is torsion). For any $r \in \mathbb{Z}$ we shall denote

$$\mathbb{Z}(r) := (\mathbb{Z}, K, (2\pi i)^r) \in \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}.$$

For $H = (H_{\mathbb{Z}}, H_K, \omega)$ and $H' = (H'_{\mathbb{Z}}, H'_K, \omega')$ we can define

$$H \otimes H' := (H_{\mathbb{Z}} \otimes_{\mathbb{Z}} H'_{\mathbb{Z}}, H_K \otimes_K H'_K, \omega \otimes \omega') \tag{2.10}$$

and set $H(r) := H \otimes \mathbb{Z}(r)$ the Tate twist. For $H \in \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$, say that $H = (H_{\mathbb{Z}}, H_K, \omega)$ with $H_{\mathbb{Z}}$ free, we have duals $H^{\vee} \in \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ given by

$$(H_{\mathbb{Z}}, H_K, \omega)^{\vee} := (H_{\mathbb{Z}}^{\vee}, H_K^{\vee}, \omega^{\vee}) \tag{2.11}$$

where $H_{\mathbb{Z}}^{\vee} = \text{Hom}(H_{\mathbb{Z}}, \mathbb{Z})$ is the dual abelian group, $H_K^{\vee} = \text{Hom}(H_K, K)$ is the dual K -vector space, and

$$\omega^{\vee} : H_{\mathbb{Z}}^{\vee} \rightarrow H_K^{\vee} \otimes_K \mathbb{C}$$

is the composition of the canonical mapping $H_{\mathbb{Z}}^{\vee} \rightarrow H_{\mathbb{Z}}^{\vee} \otimes_{\mathbb{Z}} \mathbb{C}$ with the \mathbb{C} -isomorphism $H_{\mathbb{Z}}^{\vee} \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\cong} H_K^{\vee} \otimes_K \mathbb{C}$ given by the inverse of the \mathbb{C} -dual of $\omega_{\mathbb{C}}$ in (2.9), i.e., $\omega^{\vee}(f) = (f \otimes_{\mathbb{Z}} \text{id}_{\mathbb{C}}) \circ \omega_{\mathbb{C}}^{-1}$ for any $f : H_{\mathbb{Z}} \rightarrow \mathbb{Z}$, up to the canonical isomorphism $H_K^{\vee} \otimes_K \mathbb{C} \simeq (H_K \otimes_K \mathbb{C})^{\vee}$. We clearly get that $(H^{\vee})^{\vee} = H$ and $(\)^{\vee} : \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}} \rightarrow \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ is a dualizing functor. Note that $\mathbb{Z}(r)^{\vee} = \mathbb{Z}(-r)$ so that $H(r)^{\vee} = H^{\vee}(-r)$ for $r \in \mathbb{Z}$.

Similar constructions can be done for the \mathbb{Q} -linear variant $\text{Mod}_{\mathbb{Q},K}^{\cong}$. Note that $\text{Mod}_{\mathbb{Q},K}^{\cong}$ (resp. $\text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$) admits an internal Hom defined via the internal Hom of the category of finite dimensional \mathbb{Q} -vector spaces (resp. lattices). Furthermore these categories do have an identity object: $\mathbb{1} = \mathbb{Z}(0) \in \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ and $\mathbb{1} = \mathbb{Q}(0) \in \text{Mod}_{\mathbb{Q},K}^{\cong}$, respectively. For any object H of $\text{Mod}_{\mathbb{Q},K}^{\cong}$ we have $H^{\vee} = \underline{\text{Hom}}(H, \mathbb{1})$ and $\text{End}(\mathbb{1}) = \mathbb{Q}$. Hence all objects of $\text{Mod}_{\mathbb{Q},K}^{\cong}$ are reflexive. Similarly, for $\text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$.

Lemma 2.4.2. *The categories $\text{Mod}_{\mathbb{Z},K}$ and $\text{Mod}_{\mathbb{Z},K}^{\cong}$ are abelian tensor categories. The category $\text{Mod}_{\mathbb{Q},K}^{\cong}$ is a neutral Tannakian category with fibre functor the forgetful functor to \mathbb{Q} -vector spaces.*

Note that there is a cohomological version of $\text{Mod}_{\mathbb{Z},K}^{\cong}$ and $\text{Mod}_{\mathbb{Q},K}^{\cong}$, which is called the de Rham–Betti category in the existing literature (cf. [3, 7.5]).

Definition 2.4.3. Let $\text{Mod}_{K,\mathbb{Z}}^{\cong}$ be the category whose objects are triples $(H_K, H_{\mathbb{Z}}, \eta)$ where H_K is a finite dimensional K -vector space, $H_{\mathbb{Z}}$ is a finitely generated abelian group and

$$\eta : H_K \otimes_K \mathbb{C} \xrightarrow{\cong} H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C}$$

is an isomorphism of \mathbb{C} -vector spaces. We shall call $\text{Mod}_{K,\mathbb{Z}}^{\cong}$ and its \mathbb{Q} -linear variant $\text{Mod}_{K,\mathbb{Q}}^{\cong}$ the categories of *cohomological periods*.

The category $\text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}$ is denoted \mathcal{C}_{dRB} in [17, §2.1] and in [16, §5.3]. The \mathbb{Q} -linear variant $\text{Mod}_{K,\mathbb{Q}}^{\cong}$ is denoted $(K, \mathbb{Q})\text{-Vect}$ in [33, Chap. 5]. For these categories we have an analogue of Lemma 2.4.2; in particular, a dualizing functor exists.

Lemma 2.4.4. *There is canonical equivalence given by the functor*

$$\varsigma : \text{Mod}_{\mathbb{Z},K}^{\cong} \rightarrow \text{Mod}_{K,\mathbb{Z}}^{\cong} \quad \varsigma(H_{\mathbb{Z}}, H_K, \omega) := (H_K, H_{\mathbb{Z}}, \omega_{\mathbb{C}}^{-1})$$

which induces an equivalence between the tensor subcategories $\text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ and $\text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}$.

We set

$$\mathbb{Z}(r) := \varsigma(\mathbb{Z}(r)) \in \text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}$$

Note that, for $H \in \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ we may consider $H^\circ \in \text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}$ setting

$$(H_{\mathbb{Z}}, H_K, \omega)^\circ := (H_K^\vee, H_{\mathbb{Z}}^\vee, \omega^\circ) = \varsigma(H^\vee) = \varsigma(H)^\vee \tag{2.12}$$

where $\omega^\circ : H_K^\vee \otimes_K \mathbb{C} \xrightarrow{\cong} H_{\mathbb{Z}}^\vee \otimes_{\mathbb{Z}} \mathbb{C}$ is just given by the \mathbb{C} -dual of $\omega_{\mathbb{C}}$ in (2.9). We then have $\mathbb{Z}(r)^\circ = \mathbb{Z}(-r) \in \text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}$ so that $H(r)^\circ = H^\circ(-r)$ for all $r \in \mathbb{Z}$.

The functor $(\)^\circ$ is an anti-equivalence and there is an induced equivalence $\text{Mod}_{\mathbb{Q},K}^{\cong} \cong (\text{Mod}_{K,\mathbb{Q}}^{\cong})^{\text{op}}$ of neutral Tannakian categories.

2.5. Betti–de Rham realization and Cartier duality

Now recall the period mapping $\varpi_{M,\mathbb{Z}} : T_{\mathbb{Z}}(M_{\mathbb{C}}) \rightarrow T_{\text{dR}}(M_K) \otimes_K \mathbb{C}$ provided by Definition 2.2.1. According to Theorem 2.2.2 we have that $\varpi_{M,\mathbb{C}}$ is a \mathbb{C} -isomorphism.

Definition 2.5.1. For K a subfield of \mathbb{C} , $M_K \in {}^t\mathcal{M}_1(K)$ and $\varpi_{M,\mathbb{Z}}$ we set

$$T_{\text{BdR}}(M_K) := (T_{\mathbb{Z}}(M_{\mathbb{C}}), T_{\text{dR}}(M_K), \varpi_{M,\mathbb{Z}}) \in \text{Mod}_{\mathbb{Z},K}^{\cong}$$

and the \mathbb{Q} -linear variant

$$T_{\text{BdR}}^{\mathbb{Q}}(\mathbf{M}_K) := (T_{\mathbb{Q}}(\mathbf{M}_{\mathbb{C}}), T_{\text{dR}}(\mathbf{M}_K), \varpi_{\mathbf{M}, \mathbb{Q}}) \in \text{Mod}_{\mathbb{Q}, K}^{\cong}$$

where $T_{\mathbb{Q}}(\mathbf{M}_{\mathbb{C}}) := T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}) \otimes_{\mathbb{Z}} \mathbb{Q}$. Call these realizations the *Betti-de Rham* realizations.

Since the period mapping $\varpi_{\mathbf{M}, \mathbb{Z}}$ in $T_{\text{BdR}}(\mathbf{M}_K)$ is covariantly functorial, by the constructions in (2.6) and (2.5), the Betti-de Rham realization yields a functor

$$T_{\text{BdR}} : {}^t\mathcal{M}_1(K) \rightarrow \text{Mod}_{\mathbb{Z}, K}^{\cong} \tag{2.13}$$

in the homological category $\text{Mod}_{\mathbb{Z}, K}^{\cong}$. Similarly, with rational coefficients, we get a functor from 1-motives up to isogenies $\mathcal{M}_1^{\mathbb{Q}}(K) \cong {}^t\mathcal{M}_1^{\mathbb{Q}}(K)$ to $\text{Mod}_{\mathbb{Q}, K}^{\cong}$. By Examples 2.2.3 we have $T_{\text{BdR}}(\mathbb{Z}[0]) = \mathbb{Z}(0)$ and $T_{\text{BdR}}(\mathbb{G}_m[-1]) = \mathbb{Z}(1)$.

Definition 2.5.2. For $H = (H_{\mathbb{Z}}, H_K, \omega) \in \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ define the *Cartier dual*

$$H^* := (H_{\mathbb{Z}}^{\vee}, H_K^{\vee}, 2\pi i \omega^{\vee}) = H^{\vee}(1) = H(-1)^{\vee} = \underline{\text{Hom}}(H, \mathbb{Z}(1)) \in \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$$

Note that this construction is reflexive.

Theorem 2.5.3. For $\mathbf{M}_K \in {}^t\mathcal{M}_1(K)$ free with Cartier dual \mathbf{M}_K^* we have that

$$T_{\text{BdR}}(\mathbf{M}_K)^* \cong T_{\text{BdR}}(\mathbf{M}_K^*)$$

Proof. It suffices to prove that the Poincaré biextension of \mathbf{M}_K provides a natural morphism $T(\mathbf{M}_K) \otimes T(\mathbf{M}_K^*) \rightarrow \mathbb{Z}(1)$ which induces the usual dualities $\langle \cdot, \cdot \rangle_{\mathbb{Z}}$ and $\langle \cdot, \cdot \rangle_{\text{dR}}$ on $T_{\mathbb{Z}}$'s and $\langle \cdot, \cdot \rangle_{\text{dR}}$ on T_{dR} 's constructed in [23, §10.2.3 & §10.2.7]. This is proved in [23, Prop. 10.2.8]. \square

Note that we also have a de Rham–Betti contravariant realization in the cohomological category $\text{Mod}_{K, \mathbb{Z}}^{\cong}$. Recall from [12, §1.13] that we also have the category of 1-motives with cotorsion ${}^t\mathcal{M}_1$. Cartier duality

$$(\cdot)^* : {}^t\mathcal{M}_1 \xrightarrow{\cong} {}^t\mathcal{M}_1 \tag{2.14}$$

is an anti-equivalence of abelian categories.

Definition 2.5.4. For $\mathbf{M} \in {}^t\mathcal{M}_1$ denote

$$T_{\text{dRB}}(\mathbf{M}) := \varsigma(T_{\text{BdR}}(\mathbf{M}^*)) = (T_{\text{dR}}(\mathbf{M}^*), T_{\mathbb{Z}}(\mathbf{M}^*), \eta_{\mathbf{M}^*}) \in \text{Mod}_{K, \mathbb{Z}}^{\cong}$$

where $\eta_{\mathbf{M}^*} := \varpi_{\mathbf{M}^*, \mathbb{C}}^{-1}$ is the inverse of the \mathbb{C} -linear period isomorphism $\varpi_{\mathbf{M}^*, \mathbb{C}}$ of the Cartier dual $\mathbf{M}^* \in {}^t\mathcal{M}_1$ (see Theorem 2.2.2). Call this realization (and its \mathbb{Q} -linear variant) the *de Rham–Betti* realization.

With this definition we get a functor

$$T_{\text{dRB}} : {}_t\mathcal{M}_1^{\text{op}} \rightarrow \text{Mod}_{K, \mathbb{Z}}^{\cong} \tag{2.15}$$

Now we have $T_{\text{dRB}}(\mathbb{Z}[0]) = \mathbb{Z}(1)$ and $T_{\text{dRB}}(\mathbb{G}_m[-1]) = \mathbb{Z}(0)$. With the notation adopted in (2.12), we also have

$$T_{\text{BdR}}(\mathbf{M})^\circ(1) = (T_{\text{dR}}(\mathbf{M})^\vee, T_{\mathbb{Z}}(\mathbf{M})^\vee, (2\pi i)^{-1}\varpi_{\mathbf{M}}^\circ).$$

Lemma 2.5.5. *We have a natural isomorphism of functors $T_{\text{dRB}}(\) \cong T_{\text{BdR}}(\)^\circ(1)$.*

Proof. For $\mathbf{M} \in \mathcal{M}_1$ and its Cartier dual \mathbf{M}^* we have that $T_{\text{BdR}}(\mathbf{M})^* \cong T_{\text{BdR}}(\mathbf{M}^*) \in \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ by Theorem 2.5.3. Thus the period isomorphism of the Cartier dual $\varpi_{\mathbf{M}^*, \mathbb{Z}} = 2\pi i \varpi_{\mathbf{M}, \mathbb{Z}}^\vee$ and its \mathbb{C} -inverse $\varpi_{\mathbf{M}^*, \mathbb{C}}^{-1} = (2\pi i)^{-1} \varpi_{\mathbf{M}}^\circ$. \square

2.6. Weight and Hodge filtrations

Consider the category $\text{FMod}_{\mathbb{Z}, K}$ given by objects in $\text{Mod}_{\mathbb{Z}, K}$ endowed with finite and exhaustive filtrations and morphisms that respect the filtrations.

More precisely, an object of $\text{FMod}_{\mathbb{Z}, K}$ is an abelian group $H_{\mathbb{Z}}$ endowed with a (weight) filtration $W_{\bullet}H_{\mathbb{Z}}$ and a K -vector space H_K endowed with two filtrations $W_{\bullet}H_K, F_{\bullet}H_K$, along with the corresponding compatibilities of the ω 's on weight filtrations.

Let $\mathbf{M}_K = [\mathbf{L}_K \rightarrow \mathbf{G}_K]$ be a 1-motive over K and, as usual, let \mathbf{T}_K denote the maximal subtorus of \mathbf{G}_K . Since the Betti-de Rham realization (2.13) is functorial and compatible with the canonical weight filtration (2.3) on $T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}})$ and $T_{\text{dR}}(\mathbf{M}_K)$ is filtered by $\mathbb{V}(\mathbf{M})$, the Hodge filtration as in Remark 2.1.2, we also get a realization functor

$${}^F T_{\text{BdR}} : {}_t\mathcal{M}_1(K) \rightarrow \text{FMod}_{\mathbb{Z}, K}. \tag{2.16}$$

We have

$$(T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}), T_{\text{dR}}(\mathbf{M}_K), \varpi_{\mathbf{M}, \mathbb{Z}}) \supseteq (T_{\mathbb{Z}}(\mathbf{G}_{\mathbb{C}}), T_{\text{dR}}(\mathbf{G}_K), \varpi_{\mathbf{G}, \mathbb{Z}}) \supseteq (T_{\mathbb{Z}}(\mathbf{T}_{\mathbb{C}}), T_{\text{dR}}(\mathbf{T}_K), \varpi_{\mathbf{T}, \mathbb{Z}}).$$

Note that:

Lemma 2.6.1. *Let $K = \overline{\mathbb{Q}}$. Let \mathbf{M}_K and \mathbf{N}_K be two free 1-motives over K . Then any morphism $\varphi : T_{\text{BdR}}(\mathbf{M}_K) \rightarrow T_{\text{BdR}}(\mathbf{N}_K)$ in $\text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ preserves the weight filtrations.*

Proof. Let $\varphi = (\varphi_{\mathbb{Z}}, \varphi_K) : (T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}}), T_{\text{dR}}(\mathbf{M}_K), \varpi_{\mathbf{M}, \mathbb{Z}}) \rightarrow (T_{\mathbb{Z}}(\mathbf{N}_{\mathbb{C}}), T_{\text{dR}}(\mathbf{N}_K), \varpi_{\mathbf{N}, \mathbb{Z}})$ for \mathbf{M}_K and \mathbf{N}_K one of the following pure 1-motives: $[\mathbb{Z}_K \rightarrow 0]$, $[0 \rightarrow \mathbb{G}_{m, K}]$ and $[0 \rightarrow \mathbf{A}_K]$, where \mathbf{A}_K is an abelian variety. We show that $\varphi = 0$ for different weights, in all cases. As K is algebraically closed this implies that $\varphi = 0$ for all pure 1-motives of different weights and this easily yields the claimed compatibility.

For $M_K = [\mathbb{Z}_K \rightarrow 0]$ and $N_K = [0 \rightarrow \mathbb{G}_{m,K}]$ (respectively $M_K = [0 \rightarrow \mathbb{G}_{m,K}]$ and $N_K = [\mathbb{Z}_K \rightarrow 0]$) we have $T_{\text{BdR}}([\mathbb{Z}_K \rightarrow 0]) = \mathbb{Z}(0)$, $T_{\text{BdR}}([0 \rightarrow \mathbb{G}_{m,K}]) = \mathbb{Z}(1)$ and $\varphi = 0$ as $\varphi_K: K \rightarrow K$ is given by the multiplication by an algebraic number but the compatibility (2.8) forces such algebraic number to be $n2\pi i$ (respectively $n/2\pi i$) for some $n \in \mathbb{Z}$.

Similarly, for $M_K = [\mathbb{Z}_K \rightarrow 0]$ and $N_K = [0 \rightarrow A_K]$ we have that $\varpi_{\mathbb{Z},\mathbb{N}} \circ \varphi_{\mathbb{Z}}(1) = \varphi_K(1)$ if, and only if, $\varphi = 0$. Indeed, the preceding equality implies that $d\rho_A \circ \varpi_{\mathbb{Z},\mathbb{N}} \circ \varphi_{\mathbb{Z}}(1) = d\rho_A \circ \varphi_K(1)$. Now, the right-hand term is in $\text{Lie}(A_K)$ while by Remark 2.3.2 the left-hand term would give a transcendental point of $\text{Lie}(A_{\mathbb{C}})$ if $\varphi_{\mathbb{Z}}(1) \neq 0$.

Dually, for $M_K = [0 \rightarrow A_K]$ and $N_K = [0 \rightarrow \mathbb{G}_{m,K}]$ by making use of Theorem 2.5.3 we then get $\varphi^* = 0$ thus $\varphi = 0$.

Finally, for $M_K = [0 \rightarrow \mathbb{G}_{m,K}]$ and $N_K = [0 \rightarrow A_K]$ we can apply Theorem 2.3.4 to the pair $(\varphi_K, \varphi_{\mathbb{Z}})$ so that, dually, making use of Theorem 2.5.3, the same holds for $M_K = [0 \rightarrow A_K]$ and $N_K = [\mathbb{Z}_K \rightarrow 0]$. \square

Let $M_K = [u_K: L_K \rightarrow G_K]$ and $N_K = [v_K: F_K \rightarrow H_K]$ be free and let $\varphi: T_{\text{BdR}}(M_K) \rightarrow T_{\text{BdR}}(N_K)$ be a morphism in $\text{Mod}_{\mathbb{Z},K}^{\cong}$. Then we have a K -linear mapping $\varphi_K: T_{\text{dR}}(M_K) \rightarrow T_{\text{dR}}(N_K)$ and a homomorphism $\varphi_{\mathbb{Z}}: T_{\mathbb{Z}}(M_{\mathbb{C}}) \rightarrow T_{\mathbb{Z}}(N_{\mathbb{C}})$ which is compatible with the weight filtrations, by Lemma 2.6.1. Moreover, $\varphi_{\mathbb{Z}}$ and φ_K are compatible with the ϖ 's as in (2.8). We have that $\varphi_{\mathbb{Z}}$ restricts to a homomorphism

$$W_{-1}\varphi_{\mathbb{Z}}: W_{-1}T_{\mathbb{Z}}(M_{\mathbb{C}}) := T_{\mathbb{Z}}(G_{\mathbb{C}}) \cong H_1(G_{\mathbb{C}}^{\natural}) \rightarrow W_{-1}T_{\mathbb{Z}}(N_{\mathbb{C}}) := T_{\mathbb{Z}}(H_{\mathbb{C}}) \cong H_1(H_{\mathbb{C}}^{\natural}) \tag{2.17}$$

and we get an induced map on gr_0^W as follows

$$\varphi_{\mathbb{Z},0}: \text{gr}_0^W T_{\mathbb{Z}}(M_{\mathbb{C}}) = T_{\mathbb{Z}}(M_{\mathbb{C}})/T_{\mathbb{Z}}(G_{\mathbb{C}}) = L_{\mathbb{C}} \rightarrow \text{gr}_0^W T_{\mathbb{Z}}(N_{\mathbb{C}}) = T_{\mathbb{Z}}(N_{\mathbb{C}})/T_{\mathbb{Z}}(H_{\mathbb{C}}) = F_{\mathbb{C}}. \tag{2.18}$$

Note that $\varphi_{\mathbb{Z},0}$ is indeed defined over $\overline{\mathbb{Q}}$.

Lemma 2.6.2. *Let $K = \overline{\mathbb{Q}}$. Let M_K and N_K be two free 1-motives over K . Then any morphism $\varphi: T_{\text{BdR}}(M_K) \rightarrow T_{\text{BdR}}(N_K)$ in $\text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ preserves the Hodge filtrations.*

Proof. Let $M_K = [u_K: L_K \rightarrow G_K]$ and $N_K = [v_K: F_K \rightarrow H_K]$ be free and let $\varphi: T_{\text{BdR}}(M_K) \rightarrow T_{\text{BdR}}(N_K)$ be a morphism in $\text{Mod}_{\mathbb{Z},K}^{\cong}$. We have to show that $\varphi_K(\mathbb{V}(M)) \subseteq \mathbb{V}(N)$ where $\mathbb{V}(M)$ is the additive part of G_K^{\natural} and $\mathbb{V}(N)$ is that of H_K^{\natural} ; see Remark 2.1.2. Recall the commutative diagram

$$\begin{CD}
 T_{\mathbb{Z}}(\mathbb{M}_{\mathbb{C}}) @>\varphi_{\mathbb{Z}}>> T_{\mathbb{Z}}(\mathbb{N}_{\mathbb{C}}) \\
 @V\varpi_{\mathbb{M},\mathbb{Z}}VV @VV\varpi_{\mathbb{N},\mathbb{Z}}V \\
 T_{\text{dR}}(\mathbb{M}_{\mathbb{C}}) @>\varphi_K \otimes \text{id}_{\mathbb{C}}>> T_{\text{dR}}(\mathbb{N}_{\mathbb{C}}).
 \end{CD} \tag{2.19}$$

By definition of $\varpi_{\mathbb{M},\mathbb{Z}}$ and (2.17) there is then a commutative diagram

$$\begin{CD}
 H_1(\mathbb{G}_{\mathbb{C}}^{\natural}) @>W_{-1}\varphi_{\mathbb{Z}}>> H_1(\mathbb{H}_{\mathbb{C}}^{\natural}) \\
 @VVV @VVV \\
 \text{Lie}(\mathbb{G}_{\mathbb{C}}^{\natural}) @>\varphi_K \otimes \text{id}_{\mathbb{C}}>> \text{Lie}(\mathbb{H}_{\mathbb{C}}^{\natural})
 \end{CD}$$

where the vertical arrows are those in the horizontal sequence in the middle of diagram (2.6) for \mathbb{M} and \mathbb{N} respectively. Hence there exists an analytic morphism $h_{\mathbb{C}}: \mathbb{G}_{\mathbb{C}}^{\natural} \rightarrow \mathbb{H}_{\mathbb{C}}^{\natural}$ with $dh_{\mathbb{C}} = \varphi_K \otimes \text{id}_{\mathbb{C}}$. It is sufficient to prove that $h_{\mathbb{C}}$ is algebraic and defined over K to conclude by the structure theorem of algebraic K -groups that $h_K(\mathbb{V}(\mathbb{M})) \subseteq \mathbb{V}(\mathbb{N})$ and hence that $\varphi_K = dh_K$ preserves the Hodge filtrations.

If $L_K = 0$, by Lemma 2.6.1, φ factors through $W_{-1}T_{\text{BdR}}(\mathbb{N}_K) = T_{\text{BdR}}(\mathbb{H}_K)$. Hence we may assume $F_K = 0$ as well. It follows then from Theorem 2.3.4 applied to \mathbb{G}^{\natural} and \mathbb{H}^{\natural} that the above morphism $h_{\mathbb{C}}$ is indeed algebraic and defined over K . Hence φ_K preserves the Hodge filtrations.

Now let $L_K \neq 0$ and set $L_K^{\natural} = L_K \otimes_{\mathbb{G}_{a,K}} \mathbb{G}_{a,K}$. Since φ preserves the weights by Lemma 2.6.1 we get $W_{-1}\varphi: T_{\text{BdR}}(\mathbb{G}_K) \rightarrow T_{\text{BdR}}(\mathbb{H}_K)$. By the previous step $W_{-1}\varphi_K(\mathbb{V}(\mathbb{G})) \subseteq \mathbb{V}(\mathbb{H}) \subseteq \mathbb{V}(\mathbb{N})$. We thus obtain the following commutative diagram

$$\begin{CD}
 T_{\mathbb{Z}}(\mathbb{M}_{\mathbb{C}}) @>\varpi_{\mathbb{M},\mathbb{Z}}>> T_{\text{dR}}(\mathbb{M})/\mathbb{V}(\mathbb{G}) \otimes \mathbb{C} @>\text{exp}>> \mathbb{G}_{\mathbb{C}}^{\natural}/\mathbb{V}(\mathbb{G})_{\mathbb{C}} \simeq \mathbb{G}_{\mathbb{C}} \oplus L_{\mathbb{C}}^{\natural} \\
 @V\varphi_{\mathbb{Z}}VV @V\gamma VV @VV\delta V \\
 T_{\mathbb{Z}}(\mathbb{N}_{\mathbb{C}}) @>\varpi_{\mathbb{N},\mathbb{Z}}>> T_{\text{dR}}(\mathbb{N})/\mathbb{V}(\mathbb{N}) \otimes \mathbb{C} @>\text{exp}>> \mathbb{H}_{\mathbb{C}}^{\natural}/\mathbb{V}(\mathbb{N})_{\mathbb{C}} \simeq \mathbb{H}_{\mathbb{C}}
 \end{CD}$$

where the mapping γ is induced by φ_K , we have the canonical identification of $\mathbb{G}_K^{\natural}/\mathbb{V}(\mathbb{G}) = \mathbb{G}_K \oplus L_K^{\natural}$ and $\delta = g_{\mathbb{C}} + \beta$ with $g_{\mathbb{C}} = g_K \otimes \text{id}_{\mathbb{C}}$ and $g_K: \mathbb{G}_K \rightarrow \mathbb{H}_K$ induced by $W_{-1}\varphi_K$. We are left to show that $\beta: L_K^{\natural} \rightarrow \mathbb{H}_{\mathbb{C}}$ is zero. Since the composition of the upper arrows in the previous diagram maps $T_{\mathbb{Z}}(\mathbb{G}_{\mathbb{C}})$ to $0 \oplus 0$, we obtain a commutative square

$$\begin{CD}
 L_{\mathbb{C}} @>(u,1)>> \mathbb{G}_{\mathbb{C}} \oplus L_{\mathbb{C}}^{\natural} \\
 @V\varphi_{\mathbb{Z},0}VV @VV\delta V \\
 F_{\mathbb{C}} @>v>> \mathbb{H}_{\mathbb{C}}
 \end{CD}$$

where $\varphi_{\mathbb{Z},0}$ is the induced map as in (2.18). In particular, for $x \in L_K(K)$ we have $\beta(x \otimes 1) = v(\varphi_{\mathbb{Z},0}(x)) - g_K(u(x)) = \gamma - dg_K \otimes \text{id}_{\mathbb{C}}$ is in $H_K(K)$. On the other hand $\beta(x \otimes 1) = \exp d\beta(x \otimes 1)$. Since $d\beta = d\delta - dg_K \otimes \text{id}_{\mathbb{C}}$ we have that $d\beta(x \otimes 1)$ belongs to $\text{Lie}(H_K)$ regarded as a K -linear subspace of $\text{Lie}(H_{\mathbb{C}})$. By Remark 2.3.2 we get that $\beta(x \otimes 1) = 0$ and therefore that $\beta = 0$. \square

2.7. Full faithfulness

We are now ready to show that our previous Lemmas 2.6.1 and 2.6.2 yield the full faithfulness of Betti–de Rham and de Rham–Betti realizations.

Theorem 2.7.1. *The functors T_{BdR} in (2.13) and T_{dRB} in (2.15) restricted to $\mathcal{M}_1(K)$ are fully faithful over $K = \overline{\mathbb{Q}}$.*

Proof. Clearly, the functor T_{BdR} (resp. T_{dRB}) is faithful (cf. [4, proof of Lemma 3.3.2]) and we are left to show the fullness. Making use of Lemma 2.5.5 we are left to check the fullness for T_{BdR} . Let $M_K = [u_K: L_K \rightarrow G_K]$ and $N_K = [v_K: F_K \rightarrow H_K]$ be free and let $\varphi: T_{\text{BdR}}(M_K) \rightarrow T_{\text{BdR}}(N_K)$ be a morphism in $\text{Mod}_{\overline{\mathbb{Z}},K}^{\cong}$.

For 0-motives, i.e., if $G_K = H_K = 0$, we have $L_K \cong \mathbb{Z}_K^r$ and $F_K \cong \mathbb{Z}_K^s$, $\varphi_{\mathbb{Z}}: T_{\mathbb{Z}}(M_{\mathbb{C}}) \cong \mathbb{Z}^r \rightarrow T_{\mathbb{Z}}(N_{\mathbb{C}}) \cong \mathbb{Z}_K^s$ provides a morphism $f: M_K \rightarrow N_K$ such that $T_{\text{BdR}}(f) = \varphi$.

If the weight -1 parts are non-zero, by Lemma 2.6.1 $\varphi_{\mathbb{Z}}$ restricts to a homomorphism $W_{-1}\varphi_{\mathbb{Z}}$ as in (2.17) and it yields a morphism $\varphi_{\mathbb{Z},0}$ as in (2.18) i.e., $\varphi_{\mathbb{Z},0}$ is the map induced by $\varphi_{\mathbb{Z}}$ on gr_0^W . If we set $f_{\mathbb{C}} := \varphi_{\mathbb{Z},0}$ the homomorphism $f_{\mathbb{C}}: L_{\mathbb{C}} \rightarrow F_{\mathbb{C}}$ trivially descends to a homomorphism $f_K: L_K \rightarrow F_K$ over $K = \overline{\mathbb{Q}}$.

Let’s now consider $\varphi_{\mathbb{C}} := \varphi_K \otimes_K \text{id}_{\mathbb{C}}$ and translate (2.17) and (2.19), as in the proof of Lemma 2.6.2, in the following commutative diagram with exact rows


$$\begin{CD}
 0 @>>> H_1(G_{\mathbb{C}}^{\natural}) @>>> \text{Lie}(G_{\mathbb{C}}^{\natural}) @>{\exp}>> G_{\mathbb{C}}^{\natural} @>>> 0 \\
 @. @VVW_{-1}\varphi_{\mathbb{Z}}V @VV\varphi_{\mathbb{C}}V @VV\psi V @. \\
 0 @>>> H_1(H_{\mathbb{C}}^{\natural}) @>>> \text{Lie}(H_{\mathbb{C}}^{\natural}) @>{\exp}>> H_{\mathbb{C}}^{\natural} @>>> 0
 \end{CD}
 \tag{2.20}$$

yielding a morphism of analytic groups $\psi: G_{\mathbb{C}}^{\natural} \rightarrow H_{\mathbb{C}}^{\natural}$ on the quotients via the exponential mapping \exp , as indicated above. Now, since by Lemma 2.6.2, we have $\varphi_K(\mathbb{V}(M)) \subseteq \mathbb{V}(N)$, $\psi(\mathbb{V}(M_{\mathbb{C}})) \subseteq \mathbb{V}(N_{\mathbb{C}})$, the diagram (2.20) induces a commutative diagram

$$\begin{CD}
 0 @>>> H_1(G_{\mathbb{C}}) @>>> \text{Lie}(G_{\mathbb{C}}) @>{\exp}>> G_{\mathbb{C}} @>>> 0 \\
 @. @VVW_{-1}\varphi_{\mathbb{Z}}V @VV\varphi'_{\mathbb{C}}V @VV\psi' V @. \\
 0 @>>> H_1(H_{\mathbb{C}}) @>>> \text{Lie}(H_{\mathbb{C}}) @>{\exp}>> H_{\mathbb{C}} @>>> 0.
 \end{CD}$$

As $\varphi'_\mathbb{C}$ is the base change of the K -linear map $\text{Lie}(\mathbf{G}_K) \rightarrow \text{Lie}(\mathbf{H}_K)$ induced by φ_K , it follows from Theorem 2.3.4 that $\psi' = g_\mathbb{C}$ is the base change of the morphism $g_K : \mathbf{G}_K \rightarrow \mathbf{H}_K$ over $K = \overline{\mathbb{Q}}$ induced by $W_{-1}\varphi_K$ (see the proof of Lemma 2.6.2).

We are left to check that $h := (f_K, g_K)$ gives a morphism $h : \mathbf{M}_K \rightarrow \mathbf{N}_K$, i.e., that $g_K \circ u_K = v_K \circ f_K$, and to see that $T_{\text{BdR}}(h) = \varphi$. To show that h is a morphism of 1-motives we may work after base change to \mathbb{C} and, using (2.6), it suffices to prove that $\psi \circ u_\mathbb{C}^h = v_\mathbb{C}^h \circ f_\mathbb{C}$. Consider the following diagram



All squares are commutative. Indeed, $\text{exp} \circ \varpi_{M,\mathbb{Z}} = u_{\mathbb{C}}^h \circ \widetilde{\text{exp}}$ and $\text{exp} \circ \varpi_{N,\mathbb{Z}} = v_{\mathbb{C}}^h \circ \widetilde{\text{exp}}$ by (2.6), $f_{\mathbb{C}} \circ \widetilde{\text{exp}} = \widetilde{\text{exp}} \circ \varphi_{\mathbb{Z}}$ by definition of $f_{\mathbb{C}}$, $\varphi_{\mathbb{C}} \circ \varpi_{M,\mathbb{Z}} = \varpi_{N,\mathbb{Z}} \circ \varphi_{\mathbb{Z}}$ by the compatibility of $\varphi_{\mathbb{Z}}$ with φ_K as in (2.8), and finally $\psi \circ \text{exp} = \text{exp} \circ \varphi_{\mathbb{C}}$ by (2.20). One concludes by the surjectivity of the map $\widetilde{\text{exp}}$ that also $\psi \circ u_{\mathbb{C}}^h = v_{\mathbb{C}}^h \circ f_{\mathbb{C}}$.

Now consider the morphism $\alpha := T_{\text{BdR}}(h) - \varphi : T_{\text{BdR}}(\mathbf{M}_K) \rightarrow T_{\text{BdR}}(\mathbf{N}_K)$ in $\text{Mod}_{\mathbb{Z},K}^{\cong}$. By construction $W_{-1}\varphi_{\mathbb{Z}} = W_{-1}T_{\mathbb{Z}}(h)$ so that α is vanishing on $T_{\text{BdR}}(\mathbf{G}_K)$. Moreover we have that $\text{gr}_0^W T_{\mathbb{Z}}(h) = \varphi_{\mathbb{Z},0}$ so that α induces a morphism in $\text{Mod}_{\mathbb{Z},K}^{\cong}$ from $T_{\text{BdR}}(\mathbf{L}_K)$ to $T_{\text{BdR}}(\mathbf{H}_K)$ which is trivial by Lemma 2.6.1. \square

Remark 2.7.2. In the proof of Theorem 2.7.1, in order to show that $(f_K, g_K) : \mathbf{M}_K \rightarrow \mathbf{N}_K$ is a morphism we are left to check that $g_K \circ u_K = v_K \circ f_K$. Remark that this also follows from two key facts: (i) the pullbacks $\tilde{u}_{\mathbb{C}}$ and $\tilde{v}_{\mathbb{C}}$ of $u_{\mathbb{C}}$ and $v_{\mathbb{C}}$ factor through the period mappings ϖ 's and (ii) the mappings $\varphi_{\mathbb{Z}}$ and φ_K are compatible with the ϖ 's.

In fact, according to the above notation, for $x \in \mathbf{L}_{\mathbb{C}}$ pick $\widetilde{\log}(x) \in T_{\mathbb{Z}}(\mathbf{M}_{\mathbb{C}})$ and note that $\varphi_{\mathbb{Z}}(\widetilde{\log}(x)) = \widetilde{\log}(f(x))$. Making use of (2.7) we obtain

$$g_{\mathbb{C}}(u_{\mathbb{C}}(x)) = g_{\mathbb{C}} \text{exp} d\rho_{\mathbf{G}} \varpi_{M,\mathbb{Z}}(\widetilde{\log}(x)) = \text{exp} d\rho_{\mathbf{H}} d g_{\mathbb{C}}^h \varpi_{M,\mathbb{Z}}(\widetilde{\log}(x))$$

by the functoriality of exp . Now $\psi = \varphi_K \otimes_K 1_{\mathbb{C}}$ and we are assuming the compatibility $(\varphi_K \otimes_K 1_{\mathbb{C}}) \circ \varpi_{M,\mathbb{Z}} = \varpi_{N,\mathbb{Z}} \circ \varphi_{\mathbb{Z}}$ so that

$$g_{\mathbb{C}}(u_{\mathbb{C}}(x)) = \text{exp} d\rho_{\mathbf{H}} \varpi_{N,\mathbb{Z}} \varphi_{\mathbb{Z}}(\widetilde{\log}(x)) = \text{exp} d\rho_{\mathbf{H}} \varpi_{N,\mathbb{Z}}(\widetilde{\log}(f(x))) = v_{\mathbb{C}}(f(x))$$

using (2.7) again, as claimed.

We notice that an alternative proof of Theorem 2.7.1 can be given using Wüstholz’s analytic subgroup Theorem 2.3.5 as follows:

Alternative Proof of Theorem 2.7.1. Let $M_K = [u_K : L_K \rightarrow G_K]$ and $N_K = [v_K : F_K \rightarrow H_K]$ be free 1-motives and let $\varphi = (\varphi_Z, \varphi_K) : T_{\text{BdR}}(M_K) \rightarrow T_{\text{BdR}}(N_K)$ be a morphism in $\text{Mod}_{\mathbb{Z}, K}^{\cong}$. Let $W = G_K^{\natural} \times H_K^{\natural}$ and note that we have commutative squares

$$\begin{CD}
 T := T_{\mathbb{Z}}(M_{\mathbb{C}}) @>(id, \varphi_{\mathbb{Z}})>> T_{\mathbb{Z}}(M_{\mathbb{C}}) \times T_{\mathbb{Z}}(N_{\mathbb{C}}) @>(\varpi_{M, \mathbb{Z}} \times \varpi_{N, \mathbb{Z}})>> \text{Lie } W_{\mathbb{C}} = \text{Lie } G_{\mathbb{C}}^{\natural} \oplus \text{Lie } H_{\mathbb{C}}^{\natural} \\
 @VVV @VVV @VV \text{exp} V \\
 L_{\mathbb{C}} @>(id, \varphi_{\mathbb{Z}, 0})>> L_{\mathbb{C}} \times F_{\mathbb{C}} @>u^{\natural} \times v^{\natural}>> W_{\mathbb{C}} := G_{\mathbb{C}}^{\natural} \times H_{\mathbb{C}}^{\natural}
 \end{CD}$$

where the horizontal arrows are injective. Let S denote the image of T in $\text{Lie } W_{\mathbb{C}}$; it is contained in $\text{exp}^{-1}(W(\overline{\mathbb{Q}}))$ since the image of $L_{\mathbb{C}} \times F_{\mathbb{C}}$ via $u^{\natural} \times v^{\natural}$ is contained in $W(\overline{\mathbb{Q}})$. Let V denote the image of $\text{Lie } G_K^{\natural}$ in $\text{Lie } W$ via the map $\text{id} \oplus \varphi_K$. By the compatibility of φ_Z and φ_K over \mathbb{C} via the homomorphisms of periods, $V_{\mathbb{C}}$ coincides with the \mathbb{C} -span of S . It then follows from Theorem 2.3.5 that there exists an algebraic subgroup $Z \subset W$ whose Lie algebra is V . Now, the composition of the inclusion $Z \rightarrow W$ with the projection $W \rightarrow G_K^{\natural}$ is an isogeny, since it is an isomorphism on Lie algebras. In fact, it is an isomorphism; indeed the injective map $T \rightarrow V_{\mathbb{C}} = \text{Lie } Z_{\mathbb{C}} \subset \text{Lie } W_{\mathbb{C}}$ maps $H_1(G_{\mathbb{C}}^{\natural}) \subset T$ into $H_1(Z_{\mathbb{C}}) \subset H_1(W_{\mathbb{C}})$ and hence the isomorphism $\text{Lie } Z_{\mathbb{C}} \xrightarrow{\sim} \text{Lie } G_{\mathbb{C}}^{\natural}$ restricts to an isomorphism $H_1(Z_{\mathbb{C}}) \xrightarrow{\sim} H_1(G_{\mathbb{C}}^{\natural})$. Let $\gamma : G_K^{\natural} \rightarrow H_K^{\natural}$ be the homomorphism of algebraic K -groups defined by composing the inverse of the isomorphism $Z \rightarrow G_K^{\natural}$ with the inclusion $Z \rightarrow W$ and the second projection $W \rightarrow H_K^{\natural}$. By construction $\text{Lie } \gamma = \varphi_K$.

In order to see that $f := (\varphi_{Z, 0}, \gamma)$ is a morphism of 1-motives with $T_{\text{BdR}}(f) = \varphi$ it suffices to check that $\gamma_{\mathbb{C}} \circ u_{\mathbb{C}}^{\natural} = v_{\mathbb{C}}^{\natural} \circ \varphi_{Z, 0}$ as morphisms $L_{\mathbb{C}} \rightarrow H_{\mathbb{C}}^{\natural}$. The latter fact is equivalent to the equality $(\text{id}_{G^{\natural}}, \gamma) \circ u^{\natural} = (u^{\natural}, v^{\natural} \circ \varphi_{Z, 0})$ as morphisms $L_{\mathbb{C}} \rightarrow W_{\mathbb{C}}$, and, by the above diagram, this is satisfied whenever $(\text{id}_{\text{Lie } G^{\natural}}, \text{Lie } \gamma) \circ \varpi_{M, \mathbb{Z}} = (\varpi_{M, \mathbb{Z}}, \varpi_{N, \mathbb{Z}} \circ \varphi_{\mathbb{Z}}) : T \rightarrow \text{Lie } W_{\mathbb{C}}$. Then we conclude by the commutativity of diagram (2.8) since $\text{Lie } \gamma_{\mathbb{C}} = \varphi_K \otimes \text{id}_{\mathbb{C}}$. \square

2.8. Descent to number fields

Let K'/K be a field extension with $K' \subseteq \overline{\mathbb{Q}}$. Note the following commutative diagram of functors

$$\begin{CD}
 \mathcal{M}_1(K) @>T_{\text{BdR}}>> \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}} \\
 @VVV @VVV \\
 \mathcal{M}_1(K') @>T_{\text{BdR}}>> \text{Mod}_{\mathbb{Z}, K'}^{\cong, \text{fr}}
 \end{CD} \tag{2.21}$$

where the functor on the left is the usual base-change and the vertical functor on the right maps $(H_{\mathbb{Z}}, H_K, \omega)$ to $(H_{\mathbb{Z}}, H_K \otimes_K K', \omega)$ using the canonical isomorphism $(H_K \otimes_K K') \otimes_{K'} \mathbb{C} \simeq H_K \otimes_K \mathbb{C}$.

Proposition 2.8.1. *Let K be a subfield of $\overline{\mathbb{Q}}$. The functor $T_{\text{BdR}}: \mathcal{M}_1(K) \rightarrow \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ is fully faithful.*

Proof. The functor T_{BdR} is fully faithful over $\overline{\mathbb{Q}}$ by Theorem 2.7.1; hence it is faithful over K , since the left-hand vertical functor in (2.21) is faithful.

Assume now $M_K = [u: L_K \rightarrow G_K], N_K = [v: F_K \rightarrow H_K]$ are 1-motives over K and let $(\varphi_{\mathbb{Z}}, \varphi_K): T_{\text{BdR}}(M_K) \rightarrow T_{\text{BdR}}(N_K)$ be a morphism in $\text{Mod}_{\mathbb{Z}, K}$. By Theorem 2.7.1 there exists a morphism $\psi: M_{\overline{\mathbb{Q}}} \rightarrow N_{\overline{\mathbb{Q}}}$ such that $T_{\text{BdR}}(\psi) = (\varphi_{\mathbb{Z}}, \varphi_K \otimes_K \text{id}_{\overline{\mathbb{Q}}})$. Note that there exists a subfield $K' \subset \overline{\mathbb{Q}}$ with K'/K finite Galois and $\psi = (f, g)$ is defined over K' . We may further assume that $L_{K'}, F_{K'}$ are constant free. Hence $\text{gr}_0^W \varphi_{\mathbb{Z}}$ descends over K' and we have a commutative square

$$\begin{CD}
 L_{K'} @>f=\text{gr}_0^W(\varphi_{\mathbb{Z}})>> F_{K'} \\
 @VVV @VVV \\
 L_{K'} \otimes G_{a, K'} @>\text{gr}_0^W(\varphi_{K'})>> F_{K'} \otimes G_{a, K'}
 \end{CD} \tag{2.22}$$

where the vertical morphisms map x to $x \otimes 1$ (and descend the homomorphism of periods for $\text{gr}_0^W(M)$ and $\text{gr}_0^W(N)$ respectively). By diagram (2.22) f descends over K since $\text{gr}_0^W(\varphi_{K'}) = \text{gr}_0^W(\varphi_K) \otimes \text{id}_{K'}$ and the vertical morphisms are injective on points. In order to check that ψ descends over K , we may then reduce to the case $L_K = F_K = 0$. By Cartier duality, we may further reduce to the case where $L_K = F_K = 0$ and $G_K = A_K, H_K = B_K$ are abelian varieties.

For any $\tau \in \text{Gal}(K'/K)$ let τ also denote the corresponding K -automorphism of $\text{Spec } K'$. Further let $\tau^*A_{K'}$ denote the base change of $A_{K'}$ along τ and let $\tau_{A_{K'}}: A_{K'} \rightarrow A_{K'}$ be equal to $\text{id}_{A_{K'}} \otimes \tau$. It is not a morphism of K' -schemes in general. Finally let $\iota_{A, \tau}: A_{K'} \rightarrow \tau^*A_{K'}$ be the canonical morphism of K' -schemes such that $\tau_{A_{K'}}$ is the composition of $\iota_{A, \tau}$ with the projection $\tau^*A_{K'} \rightarrow A_{K'}$.

In order to prove that the morphism $\psi: A_{K'} \rightarrow B_{K'}$ is defined over K we have to check that for any $\tau \in \text{Gal}(K'/K)$ it is $\tau_{B_{K'}} \circ \psi = \psi \circ \tau_{A_{K'}}$. In fact it is sufficient to check that $\iota_{B, \tau} \circ \psi = (\tau^*\psi) \circ \iota_{A, \tau}$ as morphisms of K' -schemes $A_{K'} \rightarrow \tau^*B_{K'}$, where $\tau^*\psi: \tau^*A_{K'} \rightarrow \tau^*B_{K'}$ is the obvious base change of ψ . By faithfulness of T_{BdR} , it is sufficient to check that

$$T_{\text{BdR}}(\iota_{B, \tau}) \circ T_{\text{BdR}}(\psi) = T_{\text{BdR}}(\tau^*\psi) \circ T_{\text{BdR}}(\iota_{A, \tau}). \tag{2.23}$$

Note that since A_K is a K -form of $A_{K'}$, we may identify $A_{K'}$ with $\tau^*A_{K'}$ so that $\iota_{A, \tau}$ becomes the identity map. Further $T_{\text{dR}}(\tau^*\psi) = \tau^*(\varphi_K \otimes \text{id}_{K'})$ may be identified with

$\varphi_K \otimes \text{id}_{K'}$, and $T_{\mathbb{Z}}(\tau^*\psi)$ with $\varphi_{\mathbb{Z}}$. We conclude that $T_{\text{BdR}}(\tau^*\psi)$ may be identified with $(\varphi_{\mathbb{Z}}, \varphi_K \otimes \text{id}_{K'})$ and hence (2.23) is clear. \square

3. Some evidence: description of some Grothendieck arithmetic invariants

Throughout this section we assume that $K = \overline{\mathbb{Q}}$ and by scheme we mean a separated scheme of finite type over K . In order to show the period conjecture for motivic cohomology (1.6) we are left to deal with rational coefficients. However, we prefer to keep the arguments integral when possible. In general, for any algebraic scheme X over K , by making use of the period isomorphism $\varpi_X^{p,q}$ and its inverse $\eta_X^{p,q}$ in Definition 1.2.4 we set

$$H_{\text{BdR}}^{p,q}(X) := (H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)), H_{\text{dR}}^p(X), \varpi_X^{p,q}) \in \text{Mod}_{\mathbb{Z}, K}^{\cong}$$

and

$$H_{\text{dRB}}^{p,q}(X) := (H_{\text{dR}}^p(X), H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q)), \eta_X^{p,q}) \in \text{Mod}_{K, \mathbb{Z}}^{\cong}.$$

Note that $\varsigma(H_{\text{BdR}}^{p,q}(X)) = H_{\text{dRB}}^{p,q}(X)$. We have that $\varpi_X^{p,q} = (2\pi i)^q \varpi_X^{p,0}$ and $\eta_X^{p,q} = (2\pi i)^{-q} \eta_X^{p,0}$ where $\eta_X^{p,0} : H_{\text{dR}}^p(X) \otimes_K \mathbb{C} \xrightarrow{\cong} H^p(X_{\text{an}}, \mathbb{C})$ is the usual de Rham–Betti comparison isomorphism (up to a sign cf. [33, Def. 5.3.1] and [45, Lemma 4.1.1 & Prop. 4.1.2] for the Nisnevich topology). In particular we have that $H_{\text{dRB}}^{p,q}(X) = H_{\text{dRB}}^{p,0}(X)(q)$.

3.1. Period cohomology revisited

For $H \in \text{Mod}_{\mathbb{Z}, K}$ we set

$$H_{\varpi} := \text{Hom}(\mathbb{Z}(0), H)$$

where the Hom-group is taken in $\text{Mod}_{\mathbb{Z}, K}$. This yields a functor

$$(\)_{\varpi} : \text{Mod}_{\mathbb{Z}, K} \rightarrow \text{Mod}_{\mathbb{Z}}$$

to the category of finitely generated abelian groups. Similarly, let $H_{\varpi} := \text{Hom}(\mathbb{Z}(0), H)$ for $H \in \text{Mod}_{K, \mathbb{Z}}^{\cong}$ where now the Hom-group is taken in $\text{Mod}_{K, \mathbb{Z}}^{\cong}$. By Lemma 2.4.4, for $H \in \text{Mod}_{\mathbb{Z}, K}^{\cong}$ we clearly have that

$$H_{\varpi} = \varsigma(H)_{\varpi}.$$

Moreover, for $H \in \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ we have $H^* = H(-1)^{\vee} \in \text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ so that

$$H_{\varpi}^* = \text{Hom}(\mathbb{Z}(0), H(-1)^{\vee}) = \text{Hom}(H, \mathbb{Z}(1)).$$

Note that for $H \in \text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}$ we also have $H^\circ(1) = H(-1)^\circ \in \text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}$ (see (2.12)) and we shall denote

$$H^\varpi := \text{Hom}_{\text{Mod}_{K,\mathbb{Z}}^{\cong, \text{fr}}}(\mathbb{Z}(0), H(-1)^\circ) = \text{Hom}_{\text{Mod}_{\mathbb{Z},K}^{\cong, \text{fr}}}(H, \mathbb{Z}(1)) = H_\varpi^*.$$

With rational coefficients, for $H \in \text{Mod}_{\mathbb{Z},K} \otimes \mathbb{Q}$ and the corresponding $H^\mathbb{Q} \in \text{Mod}_{\mathbb{Q},K}$ we then have $H_\varpi \otimes_{\mathbb{Z}} \mathbb{Q} \cong H_\varpi^\mathbb{Q} := \text{Hom}(\mathbb{Q}(0), H^\mathbb{Q})$ and, similarly, $H^\varpi \otimes_{\mathbb{Z}} \mathbb{Q} \cong H_\varpi^\varpi$. We have (cf. [17, Def. 2.1] and [16, (5.15)]):

Lemma 3.1.1. *For $H = (H_{\mathbb{Z}}, H_K, \omega) \in \text{Mod}_{\mathbb{Z},K}$ we have $H_\varpi \cong H_{\mathbb{Z}} \cap H_K$ where \cap is the inverse image of H_K under $\omega : H_{\mathbb{Z}} \rightarrow H_K \otimes_K \mathbb{C}$. Moreover, for $H = (H_K, H_{\mathbb{Z}}, \eta) \in \text{Mod}_{K,\mathbb{Z}}^{\cong}$, we have that $H_\varpi \cong H_K \cap H_{\mathbb{Z}}$ where \cap is the inverse image of H_K under the composition of $H_{\mathbb{Z}} \rightarrow H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\eta^{-1}} H_K \otimes_K \mathbb{C}$.*

Proof. The identifications are given by mapping $\varphi \in \text{Hom}(\mathbb{Z}(0), H)$ to $\varphi(1) \in H_{\mathbb{Z}} \cap H_K$. \square

Similarly, for $H = (H_K, H_{\mathbb{Q}}, \omega) \in \text{Mod}_{K,\mathbb{Q}}^{\cong}$ we have that $H_\varpi^\mathbb{Q} \cong H_K \cap H_{\mathbb{Q}}$. We then clearly obtain:

Corollary 3.1.2. *For $H = H_{\text{dRB}}^{p,q}(X)$ we have that $H_\varpi \cong H_\varpi^{p,q}(X)$ coincides with the period cohomology of Definition 1.2.7.*

Moreover, composing the functor $H \rightsquigarrow H_\varpi$ with the Betti–de Rham realization of 1-motives T_{BdR} in (2.13) we obtain a functor

$$T_\varpi := ()_\varpi \circ T_{\text{BdR}} : {}^t\mathcal{M}_1(K) \rightarrow \text{Mod}_{\mathbb{Z}}. \tag{3.1}$$

For a 1-motive $M \in {}^t\mathcal{M}_1(K)$ we also have $T_{\text{dRB}}(M) \in \text{Mod}_{K,\mathbb{Z}}^{\cong}$. Composing $H \rightsquigarrow H_\varpi$ with the de Rham–Betti realization T_{dRB} in (2.15) now yields a functor

$$T^\varpi := ()_\varpi \circ T_{\text{dRB}} : {}^t\mathcal{M}_1(K)^{op} \rightarrow \text{Mod}_{\mathbb{Z}}.$$

We also note that Lemma 2.5.5 yields:

Corollary 3.1.3. *For $M \in \mathcal{M}_1(K)$ we have that $T^\varpi(M) := T_{\text{dRB}}(M)_\varpi \cong T_{\text{BdR}}(M)^\varpi$.*

Working with rational coefficients we have $T_{\text{BdR}}^\mathbb{Q} := T_{\text{BdR}} \otimes \mathbb{Q}$ (resp. $T_{\text{dRB}}^\mathbb{Q} := T_{\text{dRB}} \otimes \mathbb{Q}$) and we then get a functor $T_\varpi^\mathbb{Q}$ (resp. a contravariant functor T_ϖ^ϖ) from the category of 1-motives up to isogenies $\mathcal{M}_1^\mathbb{Q} := \mathcal{M}_1 \otimes \mathbb{Q} \cong {}^t\mathcal{M}_1 \otimes \mathbb{Q} \cong {}^t\mathcal{M}_1 \otimes \mathbb{Q}$ to the category of finite dimensional \mathbb{Q} -vector spaces. Moreover, applying our Theorem 2.7.1 we have:

Corollary 3.1.4. For $M = [u : L \rightarrow G] \in \mathcal{M}_1(\overline{\mathbb{Q}})$ with Cartier dual $M^* = [u^* : L^* \rightarrow G^*] \in \mathcal{M}_1(\overline{\mathbb{Q}})$ we have that

$$T_{\varpi}(M) \cong T_{\mathbb{Z}}(M_{\mathbb{C}}) \cap T_{\text{dR}}(M_K) \cong \text{Ker } u$$

and

$$T^{\varpi}(M) \cong T_{\text{dR}}(M_K^*) \cap T_{\mathbb{Z}}(M_{\mathbb{C}}^*) \cong \text{Ker } u^*.$$

Proof. Note that $\mathbb{Z}(0) = T_{\text{BdR}}(\mathbb{Z}[0])$ and for $T_{\text{BdR}}(M) = (T_{\mathbb{Z}}(M_{\mathbb{C}}), T_{\text{dR}}(M_K), \varpi_{M, \mathbb{Z}})$ we have

$$\text{Ker } u \cong \text{Hom}_{\mathcal{M}_1(K)}(\mathbb{Z}[0], M) \xrightarrow{T_{\text{BdR}}} \text{Hom}(T_{\text{BdR}}(\mathbb{Z}[0]), T_{\text{BdR}}(M)) = T_{\varpi}(M)$$

which is an isomorphism over $K = \overline{\mathbb{Q}}$ as proven in Theorem 2.7.1. We just apply Lemma 3.1.1. Moreover, $T_{\text{dRB}}(M) = (T_{\text{dR}}(M_K^*), T_{\mathbb{Z}}(M_{\mathbb{C}}^*), \eta_{M^*})$, $\mathbb{Z}(0) = T_{\text{dRB}}(\mathbb{G}_m[-1])$ and we have an isomorphism

$$\text{Hom}_{\mathcal{M}_1(K)}(M, \mathbb{G}_m[-1]) \xrightarrow{T_{\text{dRB}}} \text{Hom}(T_{\text{dRB}}(\mathbb{G}_m[-1]), T_{\text{dRB}}(M)) = T^{\varpi}(M)$$

and $\text{Hom}_{\mathcal{M}_1(K)}(M, \mathbb{G}_m[-1]) \cong \text{Hom}_{\mathcal{M}_1(K)}(\mathbb{Z}[0], M^*) = \text{Ker } u^*$ showing the claim. \square

3.2. Period conjecture for $q = 1$

Recall that $\mathbb{Z}(1) \in \text{DM}_{\text{ét}}^{\text{eff}}$ is canonically identified with $\text{Tot}([0 \rightarrow \mathbb{G}_m]) = \mathbb{G}_m[-1]$ (see [12, Lemma 1.8.7]). We then have

$$H^{p,1}(X) \cong H_{\text{ét}}^{p-1}(X, \mathbb{G}_m)$$

for all $p \in \mathbb{Z}$. Recall the motivic Albanese triangulated functor

$$\text{LAlb} : \text{DM}_{\text{gm}}^{\text{eff}} \rightarrow D^b(\mathcal{M}_1)$$

where $\text{DM}_{\text{gm}}^{\text{eff}} \subset \text{DM}_{\text{Nis}}^{\text{eff}}$ is the subcategory of compact objects, i.e., the category of geometric motives, which has been constructed in [12, Def. 5.2.1] (see also [10, Thm. 2.4.1]). This functor is integrally defined. Rationally, LAlb yields a left adjoint to the inclusion functor given by Tot in (2.4) (see [12, Thm. 6.2.1]).

Applying LAlb to the motive of any algebraic scheme X we get $\text{LAlb}(X) \in D^b(\mathcal{M}_1)$, a complex of 1-motives whose p -th homology $L_p\text{Alb}(X) \in {}_t\mathcal{M}_1$ is a 1-motive (with cotorsion, see [12, Def. 8.2.1]). Dually, we have $\text{RPic}(X) \in D^b({}^t\mathcal{M}_1)$ (see [12, §8.3]). Taking the Cartier dual of $L_p\text{Alb}(X)$ we get $R^p\text{Pic}(X) \in {}^t\mathcal{M}_1$ and conversely via (2.14). Now, the motivic Albanese map

$$M(X) \rightarrow \text{Tot } \text{LAlb}(X)$$

in $DM_{\text{ét}}^{\text{eff}}$ (see [12, §8.2.7]) yields an integrally defined map

$$\text{Hom}_{D^b({}^t\mathcal{M}_1)}(\text{LAlb}(X), [0 \rightarrow \mathbb{G}_m][p]) \rightarrow \text{Hom}_{DM_{\text{ét}}^{\text{eff}}}(M(X), \mathbb{Z}(1)[p]) \cong H_{\text{éh}}^{p-1}(X, \mathbb{G}_m). \tag{3.2}$$

Rationally (by adjunction), this map becomes a \mathbb{Q} -linear isomorphism

$$H_{\text{éh}}^{p-1}(X, \mathbb{G}_m)_{\mathbb{Q}} \cong \text{Hom}_{DM_{\text{ét}, \mathbb{Q}}^{\text{eff}}}(M(X), \mathbb{Z}(1)[p]) \xleftarrow{\cong} \text{Hom}_{D^b(\mathcal{M}_1^{\mathbb{Q}})}(\text{LAlb}(X), \mathbb{G}_m[-1][p]). \tag{3.3}$$

Using (2.14) we set

$$\text{Ext}^p(\mathbb{Z}, \text{RPic}(X)) := \text{Hom}_{D^b({}^t\mathcal{M}_1)}(\mathbb{Z}, \text{RPic}(X)[p]) \cong \text{Hom}_{D^b({}^t\mathcal{M}_1)}(\text{LAlb}(X), \mathbb{G}_m[-1][p])$$

for all $p \in \mathbb{Z}$ and we also have (cf. [12, Lemma 10.5.1]):

Lemma 3.2.1. *For any X over $K = \overline{\mathbb{Q}}$ and $p \in \mathbb{Z}$ there is an extension*

$$0 \rightarrow \text{Ext}(\mathbb{Z}, R^{p-1}\text{Pic}(X)) \rightarrow \mathbb{E}\text{xt}^p(\mathbb{Z}, \text{RPic}(X)) \xrightarrow{\pi} \text{Hom}(\mathbb{Z}, R^p\text{Pic}(X)) \rightarrow 0$$

where the Hom and Ext are here taken in the category ${}^t\mathcal{M}_1$ of 1-motives with torsion. The composition of (3.2) with the period regulator $r_{\overline{\omega}}^{p,1} : H_{\text{éh}}^{p-1}(X, \mathbb{G}_m) \rightarrow H_{\overline{\omega}}^{p,1}(X)$ induces a mapping

$$\theta_{\overline{\omega}}^p : \text{Hom}(\mathbb{Z}, R^p\text{Pic}(X)) \rightarrow H_{\overline{\omega}}^{p,1}(X).$$

Proof. In fact, the canonical spectral sequence

$$E_2^{p,q} = \text{Ext}^p(\mathbb{Z}, R^q\text{Pic}(X)) \Rightarrow \mathbb{E}\text{xt}^{p+q}(\mathbb{Z}, \text{RPic}(X))$$

yields the claimed extension since the abelian category of 1-motives with torsion ${}^t\mathcal{M}_1(K)$ is of homological dimension 1 over the algebraically closed field $K = \overline{\mathbb{Q}}$. Moreover, for any 1-motive $M = R^p\text{Pic}(X) \in {}^t\mathcal{M}_1(K)$ the group $\text{Ext}(\mathbb{Z}, M)$ is divisible and the group $\text{Hom}(\mathbb{Z}, M)$ is finitely generated (as it follows easily by making use of [12, §C.8]). The horizontal mapping in the following commutative diagram

$$\begin{array}{ccccc} \text{Ext}(\mathbb{Z}, R^{p-1}\text{Pic}(X)) & & \xrightarrow{\text{zero}} & & \\ \downarrow & & \searrow & & \\ \mathbb{E}\text{xt}^p(\mathbb{Z}, \text{RPic}(X)) & \longrightarrow & H_{\text{éh}}^{p-1}(X, \mathbb{G}_m) & \xrightarrow{r_{\overline{\omega}}^{p,1}} & H_{\overline{\omega}}^{p,1}(X) \\ \downarrow \pi & & & & \nearrow \\ \text{Hom}(\mathbb{Z}, R^p\text{Pic}(X)) & \cdots & & \xrightarrow{\theta_{\overline{\omega}}^p} & \end{array}$$

obtained by the composition of (3.2) with the period regulator $r_{\varpi}^{p,1}$, is therefore sending $\text{Ext}(\mathbb{Z}, R^{p-1}\text{Pic}(X))$ to zero, since $H_{\varpi}^{p,1}(X)$ is finitely generated. We then get the induced mapping θ_{ϖ}^p as indicated in the diagram. \square

Also for the Betti realization, there is an integrally defined group homomorphism

$$\theta_{\mathbb{Z}}^p : T_{\mathbb{Z}}(R^p\text{Pic}(X)_{\mathbb{C}})_{\text{fr}} \rightarrow H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(1))_{\text{fr}}$$

induced via Cartier duality, by applying the Betti realization β_{σ} in (1.3) to the motivic Albanese (3.2) in a canonical way. This is justified after the natural identification of Deligne’s $T_{\mathbb{Z}}$ with the Betti realization β_{σ} on 1-motives (see [12, Thm. 15.4.1] and [53] for an explicit construction of the natural isomorphism $T_{\mathbb{Z}} \cong \beta_{\sigma} \text{Tot}$). Rationally, it yields an injection

$$\theta_{\mathbb{Q}}^p : T_{\mathbb{Q}}(R^p\text{Pic}(X)_{\mathbb{C}}) \cong H_{(1)}^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(1)) \subset H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(1))$$

where the notation $H_{(1)}^p$ is taken to indicate the largest 1-motivic part of $H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(1))$ (more precisely, this is given by the underlying \mathbb{Q} -vector space associated to the mixed Hodge structure, see [12, Cor. 15.3.1]).

For the de Rham realization, similarly, we have a K -linear mapping

$$\theta_{\text{dR}}^p : T_{\text{dR}}(R^p\text{Pic}(X)) \rightarrow H_{\text{dR}}^p(X).$$

Actually, for $M = L_p\text{Alb}(X)$ and $M^* = R^p\text{Pic}(X)$, we have η_{M^*} the \mathbb{C} -inverse of the period isomorphism $\varpi_{M^*, \mathbb{C}}$ in Theorem 2.2.2 and $\eta_X^{p,1}$ which is the inverse of the period isomorphism in Definition 1.2.4. Together with $\theta_{\mathbb{Z}}^p$ and θ_{dR}^p , we obtain a diagram

$$\begin{array}{ccccc} T_{\mathbb{Z}}(R^p\text{Pic}(X)_{\mathbb{C}})_{\mathbb{C}} & \xrightarrow{\theta_{\mathbb{Z}}^p \otimes \mathbb{C}} & H^p(X_{\text{an}}, \mathbb{Z}_{\text{an}}(1))_{\mathbb{C}} & \xrightarrow{2\pi i} & H^p(X_{\text{an}}, \mathbb{C}) \\ \eta_{M^*} \uparrow & & \eta_X^{p,1} \uparrow & \nearrow \eta_X^{p,0} & \\ T_{\text{dR}}(R^p\text{Pic}(X))_{\mathbb{C}} & \xrightarrow{\theta_{\text{dR}}^p \otimes \mathbb{C}} & H_{\text{dR}}^p(X) \otimes_K \mathbb{C} & & \end{array}$$

We have that this diagram commutes, in fact:

Lemma 3.2.2. *Let X be over the field $K = \overline{\mathbb{Q}}$ and $p \in \mathbb{Z}$. There is a morphism*

$$\theta_{\text{dRB}}^p := (\theta_{\text{dR}}^p, \theta_{\mathbb{Z}}^p) : T_{\text{dRB}}(L_p\text{Alb}(X))_{\text{fr}} \rightarrow H_{\text{dRB}}^{p,1}(X)_{\text{fr}}$$

in the category $\text{Mod}_{K, \mathbb{Z}}^{\cong, \text{fr}}$. Rationally $\theta_{\text{dRB}}^p \otimes \mathbb{Q}$ becomes injective. Moreover, θ_{dRB}^0 and θ_{dRB}^1 are integrally defined isomorphisms.

Proof. This is a consequence of [12, Cor. 16.3.2]. For $p = 0, 1$ it is straightforward that they are isomorphisms. \square

Lemma 3.2.3. *The map θ_{ϖ}^p defined in Lemma 3.2.1 factors through the de Rham-Betti realization via the Cartier duals (2.14), i.e., we have the following factorization*

$$\text{Hom}(\mathbb{Z}, R^p\text{Pic}(X)) \xlongequal{\theta_{\varpi}^p} \text{Hom}(\mathbb{L}_p\text{Alb}(X), \mathbb{G}_m[-1]) \xrightarrow{T_{\text{dRB}}} T^{\varpi}(\mathbb{L}_p\text{Alb}(X)) \xrightarrow{\iota} H_{\varpi}^{p,1}(X)$$

such that ι is given by θ_{dRB}^p in Lemma 3.2.2, using Corollary 3.1.2, as follows

$$T^{\varpi}(\mathbb{L}_p\text{Alb}(X)) = \text{Hom}(\mathbb{Z}(0), T_{\text{dRB}}(\mathbb{L}_p\text{Alb}(X))) \rightarrow \text{Hom}(\mathbb{Z}(0), H_{\text{dRB}}^{p,1}(X)) \cong H_{\varpi}^{p,1}(X)$$

and the latter Hom is here taken in $\text{Mod}_{K, \mathbb{Z}}^{\cong}$.

Proof. By construction θ_{ϖ}^p is induced by $r_{\varpi}^{p,1}$ on a quotient via the motivic Albanese (3.2) applying Betti and de Rham realizations so that the claimed factorization is clear. \square

Thus, showing the period conjecture (1.6) for $q = 1$ is equivalent to seeing that θ_{ϖ}^p is surjective, rationally. Recall (see [12, Prop. 10.4.2]) that for any X of dimension $d = \dim(X)$ the 1-motive $\mathbb{L}_{d+1}\text{Alb}(X)$ is a group of multiplicative type and

$$\mathbb{L}_p\text{Alb}(X) = \begin{cases} 0 & \text{if } p < 0 \\ [\mathbb{Z}[\pi_0(X)] \rightarrow 0] & \text{if } p = 0 \\ [L_1 \xrightarrow{u_1} G_1] & \text{if } p = 1 \\ 0 & \text{if } p > \max(2, d + 1) \end{cases}$$

where G_1 is connected, so that $\mathbb{L}_p\text{Alb}(X) \in \mathcal{M}_1$ is free for $p = 0, 1$ (see [12, Prop. 12.6.3 c)). Thus $R^0\text{Pic}(X) = [\mathbb{Z}[\pi_0(X)] \rightarrow 0]^* = [0 \rightarrow \mathbb{Z}[\pi_0(X)]^{\vee} \otimes \mathbb{G}_m]$ is a torus and we have that $\text{Ext}(\mathbb{Z}, R^0\text{Pic}(X)) = \text{Hom}_K(\mathbb{Z}, R^0\text{Pic}(X)) = K^* \otimes_{\mathbb{Z}} \mathbb{Z}[\pi_0(X)]^{\vee}$ (see [12, Prop. C.8.3 (b)]).

Theorem 3.2.4. *For any X over $K = \overline{\mathbb{Q}}$ we have that (1.6) holds true for $p = q = 1$, i.e., the period regulator $r_{\varpi}^{1,1} : H_{\text{eh}}^0(X, \mathbb{G}_m) \twoheadrightarrow H_{\varpi}^{1,1}(X)$ is surjective. Moreover, considering the 1-motive $R^1\text{Pic}(X) = [L_1^* \xrightarrow{u_1^*} G_1^*]$ which is the Cartier dual of $\mathbb{L}_1\text{Alb}(X)$ we have a canonical isomorphism*

$$\text{Ker } u_1^* \cong H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}_{\text{an}}(1)) = H_{\varpi}^{1,1}(X).$$

In particular, if X is proper $H_{\text{eh}}^0(X, \mathbb{G}_m) \cong K^* \otimes_{\mathbb{Z}} \mathbb{Z}[\pi_0(X)]^{\vee}$ and $H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}(1)) = 0$.

Proof. In fact, $R^1\text{Pic}(X)$ is free and therefore $\text{Hom}(\mathbb{L}_1\text{Alb}(X), \mathbb{G}_m[-1]) \cong \text{Hom}(\mathbb{Z}, R^1\text{Pic}(X)) \cong \text{Ker } u_1^*$. Thus the extension in Lemma 3.2.1 is

$$0 \rightarrow K^* \otimes_{\mathbb{Z}} \mathbb{Z}[\pi_0(X)]^\vee \rightarrow \mathbb{E}xt^1(\mathbb{Z}, \mathbb{R}Pic(X)) \rightarrow \text{Ker } u_1^* \rightarrow 0.$$

Moreover $\theta_{\varpi}^1 : \text{Hom}(\mathbb{Z}, \mathbb{R}^1Pic(X)) \cong \text{Ker } u_1^* \xrightarrow{\cong} H_{\varpi}^{1,1}(X)$ is an isomorphism, which in turn implies that $r_{\varpi}^{1,1}$ is a surjection. Actually, see Lemma 3.2.3, θ_{ϖ}^1 factors as follows

$$\text{Hom}(\mathbb{Z}, \mathbb{R}^1Pic(X)) \stackrel{(a)}{\cong} T^\varpi(L_1Alb(X)) \stackrel{(b)}{\cong} \text{Hom}(\mathbb{Z}(0), H_{dRB}^{1,1}(X)) \stackrel{(c)}{\cong} H_{\varpi}^{1,1}(X)$$

where: (a) is the isomorphism obtained applying Corollary 3.1.4 to $M = L_1Alb(X)$; (b) is the $\text{Hom}(\mathbb{Z}(0), -)$ of the isomorphism $\theta_{dRB}^1 : T_{dRB}(L_1Alb(X)) \cong H_{dRB}^{1,1}(X)$ given by $p = 1$ in Lemma 3.2.2; (c) is the isomorphism in Corollary 3.1.2. If X is proper then $L_1^* = 0$, i.e., $L_1Alb(X) = [L_1 \xrightarrow{u_1} G_1]$ with G_1 an abelian variety (see [12, Cor. 12.6.6]) in such a way that $\mathbb{R}^1Pic(X) = [0 \rightarrow G_1^*]$, and $H_{\text{eh}}^0(X, \mathbb{G}_m) \cong \mathbb{G}_m(\pi_0(X))$ (see [12, Lemma 12.4.1]). \square

Remark 3.2.5. We may actually compute $\mathbb{R}^1Pic(X)$ by using descent. For example, if X is normal let \overline{X} be a normal compactification of X , $p : X_\bullet \rightarrow X$ a smooth hypercovering and \overline{X}_\bullet a smooth compactification with normal crossing boundary Y_\bullet such that $\overline{p} : \overline{X}_\bullet \rightarrow \overline{X}$ is a hypercovering. Then $\overline{p}^* : Pic_{\overline{X}/K}^0 \xrightarrow{\cong} Pic_{\overline{X}_\bullet/K}^0$ is an abelian variety and

$$\mathbb{R}^1Pic(X) = [\text{Div}_{Y_\bullet}^0(\overline{X}_\bullet) \xrightarrow{u_1^*} Pic_{\overline{X}/K}^0]$$

where $\text{Div}_{Y_\bullet}^0(\overline{X}_\bullet) := \text{Ker}(\text{Div}_{Y_0}^0(\overline{X}_0) \rightarrow \text{Div}_{Y_1}^0(\overline{X}_1))$ (see [12, Prop. 12.7.2]).

For X smooth we have that (see [12, Cor. 9.2.3])

$$L_pAlb(X) = \begin{cases} [\mathbb{Z}[\pi_0(X)] \rightarrow 0] & \text{if } p = 0 \\ [0 \rightarrow \mathcal{A}_{X/K}^0] & \text{if } p = 1 \\ [0 \rightarrow \text{NS}_{X/K}^*] & \text{if } p = 2 \\ 0 & \text{otherwise,} \end{cases}$$

where $\mathcal{A}_{X/K}^0$ is the Serre-Albanese semi-abelian variety and $\text{NS}_{X/K}^*$ denotes the group of multiplicative type dual to the Néron-Severi group $\text{NS}_{X/K}$. In this case, we then have

$$\mathbb{R}^pPic(X) = \begin{cases} [0 \rightarrow \mathbb{Z}[\pi_0(X)]^*] & \text{if } p = 0 \\ [\text{Div}_{Y_\bullet}^0(\overline{X}_\bullet) \xrightarrow{u_1^*} Pic_{\overline{X}/K}^0] & \text{if } p = 1 \\ [\text{NS}_{X/K} \rightarrow 0] & \text{if } p = 2 \\ 0 & \text{otherwise,} \end{cases}$$

for a smooth compactification \overline{X} with normal crossing boundary Y . Note that, reducing to the smooth case by blow-up induction we can see that the map (3.2) is an isomorphism for $p = 0, 1$ (cf. [12, Lemma 12.6.4 b)). We deduce the following:

Corollary 3.2.6. *For any scheme X over $K = \overline{\mathbb{Q}}$ we have a short exact sequence*

$$0 \rightarrow K^* \otimes_{\mathbb{Z}} \mathbb{Z}[\pi_0(X)]^{\vee} \rightarrow H_{\text{é h}}^0(X, \mathbb{G}_m) \xrightarrow{r_{\varpi}^{1,1}} H_{\varpi}^{1,1}(X) \rightarrow 0.$$

In general, we also have:

Proposition 3.2.7. *For $K = \overline{\mathbb{Q}}$ the period regulator $r_{\varpi}^{p,1}$ admits a factorization*

$$\begin{aligned} H^{p,1}(X)_{\mathbb{Q}} &\cong H_{\text{é h}}^{p-1}(X, \mathbb{G}_m)_{\mathbb{Q}} \twoheadrightarrow T_{\mathbb{Q}}^{\varpi}(\text{L}_p \text{Alb}(X)) \hookrightarrow H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(1)) \\ &= H_{\varpi}^{p,1}(X)_{\mathbb{Q}} \end{aligned}$$

where the projection is given by Lemma 3.2.1 via $T_{\text{dRB}}^{\mathbb{Q}}$ and the inclusion is given by $\theta_{\text{dRB}}^p \otimes \mathbb{Q}$ in Lemma 3.2.2. Therefore, the conjecture (1.6) is equivalent to $T_{\mathbb{Q}}^{\varpi}(\text{L}_p \text{Alb}(X)) \cong H_{\varpi}^{p,1}(X)_{\mathbb{Q}}$.

Proof. In fact, using the adjunction (3.3), the Cartier dual π^* of π in Lemma 3.2.1, the factorization of Lemma 3.2.3 and Theorem 2.7.1 we have the following commutative diagram

$$\begin{array}{ccc} \text{Hom}_{D^b(\mathcal{M}_1^{\mathbb{Q}})}(\text{LAlb}(X), \mathbb{G}_m[-1][p]) & \xrightarrow{\cong} & H_{\text{é h}}^{p-1}(X, \mathbb{G}_m)_{\mathbb{Q}} \xrightarrow{r_{\varpi, \mathbb{Q}}^{p,1}} H_{\varpi}^{p,1}(X)_{\mathbb{Q}} \quad \square \\ \pi^* \otimes \mathbb{Q} \downarrow & \nearrow & \\ \text{Hom}(\text{L}_p \text{Alb}(X), \mathbb{G}_m[-1])_{\mathbb{Q}} & \xrightarrow[\text{T}_{\text{dRB}}^{\mathbb{Q}}]{\cong} & T_{\mathbb{Q}}^{\varpi}(\text{L}_p \text{Alb}(X)). \end{array}$$

For X smooth we further have that

$$H^{p,1}(X) \cong H_{\text{é h}}^{p-1}(X, \mathbb{G}_m) \cong H_{\text{ét}}^{p-1}(X, \mathbb{G}_m)$$

and this latter is vanishing after tensoring with \mathbb{Q} for all $p \neq 1, 2$ (see [29, Prop. 1.4]). Accordingly, the period conjecture (1.6) for X smooth and $p \neq 1, 2$ is in fact equivalent to (1.8), i.e.,

$$H_{\varpi}^{p,1}(X) = H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}(1)) = 0 \quad p \neq 1, 2. \tag{3.4}$$

For $p = 2$ and X smooth we have that $H^{2,1}(X) \cong \text{Pic}(X)$, $r_{\varpi}^{2,1} = c\ell$ is induced by the usual cycle class map and $T_{\mathbb{Q}}^{\varpi}(\text{L}_2 \text{Alb}(X)) = \text{NS}(X)_{\mathbb{Q}}$.

We here recover the results of Bost-Charles (see [16, Thm. 5.1] and [17, Cor. 3.9-3.10]) as follows. We refer to [12, Chap. 4] for the notion of biextension of 1-motives. The following is a generalization of [17, Thm. 3.8 2)] and of the discussion of the sign issue in [17, §3.4]:

Lemma 3.2.8. For $N, M \in \mathcal{M}_1(\overline{\mathbb{Q}})$ we have that

$$\text{Biext}(N, M; \mathbb{G}_m) \cong (T_{\text{BdR}}(N)^\vee \otimes T_{\text{BdR}}(M)^\vee \otimes \mathbb{Z}(1))_\varpi$$

and, when $N = M$, the subgroup of symmetric biextensions corresponds to alternating elements.

Proof. Recall that $\text{Biext}(-, M; \mathbb{G}_m)$ is representable by the Cartier dual M^* for $M \in \mathcal{M}_1(K)$ (see [12, Prop. 4.1.1]). Thus $\text{Biext}(N, M; \mathbb{G}_m) = \text{Hom}(N, M^*) \cong \text{Hom}(T_{\text{BdR}}(N), T_{\text{BdR}}(M)^*)$ where we here use Theorem 2.5.3 and Theorem 2.7.1. Now $T_{\text{BdR}}(M)^* = T_{\text{BdR}}(M)^\vee(1)$ in such a way that $\text{Hom}(T_{\text{BdR}}(N), T_{\text{BdR}}(M)^*) = \text{Hom}(\mathbb{Z}(0), T_{\text{BdR}}(N)^\vee \otimes T_{\text{BdR}}(M)^\vee \otimes \mathbb{Z}(1))$ making use of the tensor structure of the category $\text{Mod}_{\mathbb{Z}, K}^{\cong, \text{fr}}$ by Lemma 2.4.2.

Assume $N = M$. Since $\text{Biext}(M, M; \mathbb{G}_m) \cong \text{Hom}(T_{\text{BdR}}(M), T_{\text{BdR}}(M)^*)$, any biextension \mathcal{P} corresponds to a pairing $T_{\text{BdR}}(M) \otimes T_{\text{BdR}}(M) \rightarrow \mathbb{Z}(1)$ which induces the pairing [23, 10.2.3] on Deligne-Hodge realizations and the pairing [23, 10.2.7] on de Rham realizations; if \mathcal{P} is symmetric, the pairing is alternating by [23, 10.2.5 & 10.2.8]. \square

Corollary 3.2.9. For X over $K = \overline{\mathbb{Q}}$ we have that

$$\text{Biext}(\text{L}_1\text{Alb}(X), \text{L}_1\text{Alb}(X); \mathbb{G}_m)^{\text{sym}} \cong (H_{\text{dRB}}^{1,0}(X) \otimes H_{\text{dRB}}^{1,0}(X) \otimes \mathbb{Z}(1))_\varpi^{\text{alt}}.$$

Proof. Applying Lemma 3.2.8 to the free 1-motive $\text{L}_1\text{Alb}(X)$ we obtain the claimed formula. In fact, recall that $T_{\mathbb{Z}}(\text{L}_1\text{Alb}(X)) \cong H_1(X_{\text{an}}, \mathbb{Z}_{\text{an}})_{\text{fr}}$ and observe that $H_{\text{dRB}}^{1,0}(X)$ is identified with $T_{\text{BdR}}(\text{L}_1\text{Alb}(X))^\vee$ up to inverting the period isomorphism by the same argument of Lemma 3.2.2. \square

This implies that the period conjecture for $p = 2$ holds true in several cases, e.g. for abelian varieties, as previously proved by Bost (see [16, Thm. 5.1]).

3.3. The case $q = 0$

Consider the case of $\mathbb{Z}(0)$ which is canonically identified with $\text{Tot}([\mathbb{Z} \rightarrow 0]) = \mathbb{Z}[0]$. Note that $H^{p,0}(X) \cong H_{\text{eh}}^p(X, \mathbb{Z})$. Let $\mathcal{M}_0(K) \subset \mathcal{M}_1(K)$ be the full subcategory of 0-motives or Artin motives over K . Recall that the motivic π_0 (see [12, §5.4] and [10, Cor. 2.3.4]) is a triangulated functor

$$L\pi_0 : \text{DM}_{\text{gm}}^{\text{eff}} \rightarrow D^b(\mathcal{M}_0)$$

whence $L\pi_0(X) \in D^b(\mathcal{M}_0)$, a complex in the derived category of Artin motives, associated to the motive of X . We have that $M(X) \rightarrow \text{Tot } L\pi_0(X) \in \text{DM}_{\text{et}}^{\text{eff}}$ (see (2.4) for Tot) induces

$$\mathrm{Hom}_{D^b(\mathcal{M}_0)}(L\pi_0(X), \mathbb{Z}[p]) \rightarrow \mathrm{Hom}_{\mathrm{DM}_{\text{ét}}^{\text{eff}}}(M(X), \mathbb{Z}(0)[p]) \cong H_{\text{éh}}^p(X, \mathbb{Z}).$$

This map is an isomorphism, integrally, for $p = 0, 1$ (cf. [12, Lemma 12.6.4 b]) and it becomes, by adjunction, a \mathbb{Q} -linear isomorphism, for all p . Recall that for any $M \in \mathrm{DM}_{\text{gm}}^{\text{eff}}$ we have (see [12, Prop. 8.2.3])

$$\mathrm{LAlb}(M(q)) \cong \begin{cases} L\pi_0(M)(1) & \text{if } q = 1 \\ 0 & \text{for } q \geq 2 \end{cases}$$

where an Artin motive twisted by one is a 1-motive of weight -2 , i.e., the twist by one functor $(-)(1) : D^b(\mathcal{M}_0) \rightarrow D^b(\mathcal{M}_1)$ is induced by $L \rightsquigarrow [0 \rightarrow L \otimes \mathbb{G}_m]$. Note that as soon as $K = \overline{\mathbb{Q}}$ Artin motives are of homological dimension 0 and we have that

$$\mathrm{Hom}_{D^b(\mathcal{M}_0)}(L\pi_0(X), \mathbb{Z}[p]) = \mathrm{Hom}_{\mathcal{M}_0}(L_p\pi_0(X), \mathbb{Z}).$$

Moreover, we have that

$$H_{\text{éh}}^p(X, \mathbb{Z}) \cong \mathrm{Hom}_{\mathrm{DM}_{\text{ét}}^{\text{eff}}}(M(X)(1), \mathbb{Z}(1)[p])$$

by Voevodsky’s cancellation theorem [51].

Theorem 3.3.1. *For any X over $K = \overline{\mathbb{Q}}$ we have that (1.6) holds true for $p = 1$ and $q = 0$. Moreover, we have*

$$H_{\text{ét}}^1(X, \mathbb{Z}) \cong H_{\text{dR}}^1(X) \cap H^1(X_{\text{an}}, \mathbb{Z}_{\text{an}}) \cong H_{\varpi}^{1,0}(X)$$

which is vanishing if X is normal.

Proof. Making use of Proposition 1.3.1 we are left to show the period conjecture for $M(X)(1)$ in degree 1 and twist 1. We have that

$$\mathrm{Hom}_{\mathrm{DM}_{\text{ét}}^{\text{eff}}}(M(X)(1), \mathbb{Z}(1)[1]) \cong \mathrm{Hom}_{D(\mathcal{M}_1)}(\mathrm{LAlb}(M(X)(1)), \mathbb{Z}(1)[1]).$$

We have $L_0\mathrm{Alb}(M(X)(1)) \cong L_0\pi_0(X)(1) \cong [0 \rightarrow \mathbb{Z}[\pi_0(X)] \otimes \mathbb{G}_m]$ in such a way that

$$\mathrm{Ext}_{\mathcal{M}_1}(\mathrm{L}_0\mathrm{Alb}(M(X)(1)), \mathbb{G}_m[-1]) = 0$$

and (cf. (3.3) for $M(X)(1)$) we obtain

$$H_{\text{éh}}^1(X, \mathbb{Z}) \cong \mathrm{Hom}_{\mathcal{M}_1}(\mathrm{L}_1\mathrm{Alb}(M(X)(1)), \mathbb{G}_m[-1]).$$

Now $T^\varpi(\mathrm{L}_1\mathrm{Alb}(M(X)(1))) \cong H_{\varpi}^{1,1}(M(X)(1)) \cong H_{\varpi}^{1,0}(X)$ by Lemma 3.2.2 twisted by (-1) and the same argument in the proof of Theorem 3.2.4 applies here. Finally, recall that $H_{\text{ét}}^1(X, \mathbb{Z}) \cong H_{\text{éh}}^1(X, \mathbb{Z})$ for any scheme X and $H_{\text{éh}}^1(X, \mathbb{Z}) = 0$ if X is normal (see [12, Lemma 12.3.2 & Prop. 12.3.4]). \square

Remark 3.3.2. For X not normal (e.g. for the nodal curve) the group $H_{\text{ét}}^1(X, \mathbb{Z})$ can be non-zero. Moreover, for any X we have a geometric interpretation $H_{\text{ét}}^1(X, \mathbb{Z}) \cong L\text{Pic}(X) \hookrightarrow \text{Pic}(X[t, t^{-1}])$ by a theorem of Weibel [55, Thm. 7.6]. Note that this $L\text{Pic}(X)$ is also a sub-quotient of the negative K -theory group $K_{-1}(X)$ (see [55, Thm. 8.5]).

For X smooth we have a quasi-isomorphism $L\pi_0(X) \cong \mathbb{Z}[\pi_0(X)][0]$ (see [12, Prop. 5.4.1]) which means that $H^{p,0}(X)_{\mathbb{Q}} = 0$ for $p \neq 0$. This yields (as it also does Proposition 1.4.4 for X smooth) that the period conjecture (1.6) is equivalent to

$$H_{\text{dR}}^p(X) \cap H^p(X_{\text{an}}, \mathbb{Q}_{\text{an}}) = 0 \quad p \neq 0. \tag{3.5}$$

Remark 3.3.3. The period conjecture (1.6) for $q = 0$ and X smooth is also equivalent to the surjectivity of $f_{\varpi}^p : H_{\varpi}^{p,0}(\pi_0(X))_{\mathbb{Q}} \rightarrow H_{\varpi}^{p,0}(X)_{\mathbb{Q}}$ induced by the canonical morphism $f : X \rightarrow \pi_0(X)$, for all $p \geq 0$. In fact, the morphism f induces a map $M(X) \rightarrow M(\pi_0(X))$ and a commutative square by functoriality

$$\begin{array}{ccc} H^{p,0}(X)_{\mathbb{Q}} & \xrightarrow{r_{\varpi}^{p,0}} & H_{\varpi}^{p,0}(X)_{\mathbb{Q}} \\ f^p \parallel & & \uparrow f_{\varpi}^p \\ H^{p,0}(\pi_0(X))_{\mathbb{Q}} & \xrightarrow{\cong} & H_{\varpi}^{p,0}(\pi_0(X))_{\mathbb{Q}} \end{array}$$

where $f^p : \text{Hom}_{\text{DM}_{\text{ét}}^{\text{eff}}}(M(\pi_0(X)), \mathbb{Z}[p])_{\mathbb{Q}} \rightarrow \text{Hom}_{\text{DM}_{\text{ét}}^{\text{eff}}}(M(X), \mathbb{Z}[p])_{\mathbb{Q}}$ is an isomorphism for X smooth; since $\dim \pi_0(X) = 0$ then $r_{\varpi}^{p,0}$ is clearly an isomorphism for $\pi_0(X)$. For $p = 0$ the group $H^0(X_{\text{an}}, \mathbb{Z}_{\text{an}}(0))$ has rank equal to the rank of $\mathbb{Z}[\pi_0(X)]$ and f_{ϖ}^0 is an isomorphism; for $p \neq 0$ the surjectivity of f_{ϖ}^p is equivalent to the vanishing of all groups.

3.4. Arbitrary twists

We now apply Waldschmidt’s Theorem 2.3.3 to arbitrary twists.

Proposition 3.4.1. For $M = [L \rightarrow G]$ a free 1-motive over $K = \overline{\mathbb{Q}}$ and $q \in \mathbb{Z}$ an integer we have that

- 1) the group $\text{Hom}(\mathbb{Z}(q), T_{\text{BdR}}(M))$ of homomorphisms in $\text{Mod}_{\mathbb{Z}, K}^{\cong}$ or $\text{Mod}_{\mathbb{Q}, K}^{\cong}$ is trivial for $q \neq 0, 1$;
- 2) the group $\text{Hom}(\mathbb{Z}(q), T_{\text{dRB}}(M))$ of homomorphisms in $\text{Mod}_{K, \mathbb{Z}}^{\cong}$ or $\text{Mod}_{K, \mathbb{Q}}^{\cong}$ is trivial for $q \neq 0, 1$.

Proof. 1) We work in $\text{Mod}_{\mathbb{Z}, K}^{\cong}$ and leave the other case to the reader. We suppose first that $L = 0$. Consider a non trivial $\varphi \in \text{Hom}_{\text{BdR}}(\mathbb{Z}(q), T_{\text{BdR}}(M))$ and the subgroup $\Gamma = T_{\mathbb{Z}}(\mathbb{Z}(q)) = \mathbb{Z} \subset T_{\text{dR}}(\mathbb{Z}(q))_{\mathbb{C}} = \mathbb{C}$. Via the non trivial map $\varphi_K \otimes \mathbb{C} : T_{\text{dR}}(\mathbb{Z}(q))_{\mathbb{C}} \rightarrow$

$T_{\text{dR}}(\mathbb{G})_{\mathbb{C}} = \text{Lie}(\mathbb{G}_{\mathbb{C}}^{\natural})$ we can identify Γ with a subgroup of $\text{Lie}(\mathbb{G}_{\mathbb{C}}^{\natural})$. This subgroup is contained in $V_{\mathbb{C}}$ with $V \subset \text{Lie}(\mathbb{G}^{\natural})$ defined by the image $\varphi_K(T_{\text{dR}}(\mathbb{Z}(q)))$. Via the exponential map $\text{Lie}(\mathbb{G}_{\mathbb{C}}^{\natural}) \rightarrow \mathbb{G}^{\natural}(\mathbb{C})$ the image of Γ is $0 \in \mathbb{G}^{\natural}(K)$ as φ is a map in the category $\text{Mod}_{\mathbb{Z}, K}^{\cong}$ (respectively $\text{Mod}_{\mathbb{Q}, K}^{\cong}$). We deduce from Waldschmidt’s Theorem 2.3.3 that $V \subset \text{Lie}(\mathbb{G}^{\natural})$ is the Lie algebra of a 1-dimensional algebraic subgroup H of \mathbb{G}^{\natural} . There are only two possibilities $H = \mathbb{G}_a$ and $H = \mathbb{G}_m$. In both cases the period morphism for $\mathbb{Z}(q)$ identifies Γ with the subgroup $(2\pi i)^q \mathbb{Z} \subset \text{Lie}(H_{\mathbb{C}})$ that goes to 0 via $\exp_{H_{\mathbb{C}}}$. For $H = \mathbb{G}_a$ the map $\exp_{H_{\mathbb{C}}}$ is the identity, leading to a contradiction. For $H = \mathbb{G}_m$ the kernel of $\exp_{H_{\mathbb{C}}}$ is $(2\pi i)\mathbb{Z}$ forcing $q = 1$.

Secondly we suppose that $\mathbb{G} = 0$. Consider a non trivial $\varphi \in \text{Hom}_{\text{BdR}}(\mathbb{Z}(q), T_{\text{BdR}}(\mathbb{M}))$. Recall that $T_{\text{dR}}(\mathbb{M}) = \mathbb{L} \otimes K$ and the period map is induced by the inclusion $\mathbb{L} \subset \mathbb{L} \otimes K$. Let $e = \varphi_K(1) \in \mathbb{L} \otimes K$. It is a non-zero element. Using that $T_{\mathbb{Z}}(\mathbb{Z}(q))$ is identified via the period morphism for $\mathbb{Z}(q)$ with $(2\pi i)^q \mathbb{Z}$, we deduce that $\varphi_{\mathbb{Z}}(1) = (2\pi i)^q \cdot e$ should lie in $\mathbb{L} \subset \mathbb{L} \otimes K$. As π is transcendental, this forces $q = 0$.

For general $\mathbb{M} = [\mathbb{L} \rightarrow \mathbb{G}]$ we reduce to \mathbb{G} and \mathbb{L} to conclude the statement.

2) We prove the statement for $\text{Mod}_{K, \mathbb{Z}}^{\cong}$ using Lemma 2.4.4. The analogue for $\text{Mod}_{K, \mathbb{Q}}^{\cong}$ follows similarly. Given a 1-motive \mathbb{M} and its Cartier dual \mathbb{M}^* we have a natural identification

$$\varsigma : \text{Hom}(\mathbb{Z}(q), T_{\text{BdR}}(\mathbb{M}^*)) \cong \text{Hom}(\varsigma(\mathbb{Z}(q)), \varsigma(T_{\text{BdR}}(\mathbb{M}^*))) = \text{Hom}(\mathbb{Z}(q), T_{\text{dRB}}(\mathbb{M})).$$

The statement follows then from 1). \square

Denote $H_{\text{dRB},(1)}^{p,q}(X)_{\text{fr}} \subset H_{\text{dRB}}^{p,q}(X)_{\text{fr}}$ the image of $T_{\text{dRB}}(\text{L}_p \text{Alb}(X))_{\text{fr}}(q-1)$ under $\theta_{\text{dRB}}^{p,q}(q-1)$ of Lemma 3.2.2 twisted by $q-1$. We have:

Corollary 3.4.2. *We get that $H_{\varpi,(1)}^{p,q}(X)_{\text{fr}} = 0$ if $q \neq 0, 1$. For $p = 1$ we have $H_{\varpi,(1)}^{1,q}(X) = H_{\varpi}^{1,q}(X)$ and*

$$H_{\varpi}^{1,q}(X) = \begin{cases} H_{\text{ét}}^1(X, \mathbb{Z}) & \text{(see Theorem 3.3.1) if } q = 0 \\ \text{Ker } u_1^* & \text{(see Theorem 3.2.4) if } q = 1 \\ 0 & q \neq 0, 1. \end{cases}$$

Proof. We apply Proposition 3.4.1 2) to $\mathbb{M} = \text{L}_p \text{Alb}(X)$ to deduce that

$$H_{\varpi,(1)}^{p,1-q}(X)_{\text{fr}} = \text{Hom}_{\text{dRB}}(\mathbb{Z}(0), H_{\text{dRB},(1)}^{p,1}(X)(-q)) = \text{Hom}_{\text{dRB}}(\mathbb{Z}(q), T_{\text{dRB}}(\text{L}_p \text{Alb}(X))_{\text{fr}}) = 0$$

if $q \neq 0, 1$. \square

Thus, for the period conjecture in degree $p = 1$, the previous computations for the twists $q = 0, 1$ are the only relevant.

3.5. Higher odd degrees

Next, let X be a smooth and projective variety over $K = \overline{\mathbb{Q}}$. Denote $J^{2k+1}(X)$ the intermediate Jacobian: as a real analytic manifold, it is defined as the quotient of the image $H_{\mathbb{Z}}^{2k+1}(X)$ of $H^{2k+1}(X_{\text{an}}, \mathbb{Z}_{\text{an}}(k))$ in $H^{2k+1}(X_{\text{an}}, \mathbb{R}(k))$. This defines a full lattice of $H^{2k+1}(X_{\text{an}}, \mathbb{R}(k))$ so that $J^{2k+1}(X)$ is compact. It has also a natural complex analytic structure induced by the identification

$$J^{2k+1}(X) := H^{2k+1}(X_{\text{an}}, \mathbb{C}) / (F^{k+1}H^{2k+1}(X_{\text{an}}, \mathbb{C}) + (\varpi_X^{2k+1,k})^{-1}(H_{\mathbb{Z}}^{2k+1}(X))).$$

Thus $J^{2k+1}(X)$ is a complex torus.

For integers n define $N^n H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k)) \subset H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$, the n -th step of the geometric coniveau filtration, as the kernel of

$$H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k)) \longrightarrow \bigoplus_{Z \subset X} H^{2k+1}(X_{\text{an}} \setminus Z_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$$

for $Z \subset X$ varying among the codimension $\geq n$ closed subschemes.

Lemma 3.5.1. *Assume that $H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$ has geometric coniveau k , i.e., that we have $N^k H^{2k+1}(X_{\text{an}}, \mathbb{C}) = H^{2k+1}(X_{\text{an}}, \mathbb{C})$. Then $J^{2k+1}(X)$ is an abelian variety, which descends to an abelian variety $J^{2k+1}(X)_K$ over K with*

$$T_{\text{dRB}}(J^{2k+1}(X)_K) = (H_{\text{dR}}^{2k+1}(X), H_{\mathbb{Z}}^{2k+1}(X), \eta_X^{2k+1,k}).$$

Proof. Under the assumption, $H_{\mathbb{Z}}^{2k+1}(X)$ is a polarized Hodge structure of type $(1, 0)$ and $(0, 1)$ so that $J^{2k+1}(X)$ is polarizable and, hence, an abelian variety. The second statement follows from [1, Thm. A] where it is proven that there exists an abelian variety J over K and a correspondence $\Gamma \in \text{CH}^h(J \times_K X)$ over K , for $h = k + \dim J^{2k+1}(X)$, inducing an isomorphism $\Gamma_*: H^1(J_{\text{an}}, \mathbb{Q}_{\text{an}}) \cong H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$ (and hence in de Rham cohomology, compatibly with the period morphisms). Then set $J^{2k+1}(X)_K := J$. \square

The period conjecture (1.6) in odd degrees for X predicts that $H_{\varpi}^{2k+1,q}(X) = H_{\text{dR}}^{2k+1}(X) \cap H^{2k+1}(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))_{\text{fr}} = 0$ for every $k \in \mathbb{N}$ and every $q \in \mathbb{Z}$.

Proposition 3.5.2. *The period conjecture (1.6) in degree $p = 2k + 1$ and any twist q for X smooth and projective holds true if $H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$ has geometric coniveau k .*

Proof. Thanks to Lemma 3.5.1 we have that

$$\begin{aligned} H_{\varpi}^{2k+1,q}(X) &= \text{Hom}(\mathbb{Z}(0), T_{\text{dRB}}(J^{2k+1}(X)_K)(q - k)) \\ &= \text{Hom}(\mathbb{Z}(k - q), T_{\text{dRB}}(J^{2k+1}(X)_K)). \end{aligned}$$

This is trivial for $k - q \neq 0, 1$ by Proposition 3.4.1. Now use Theorem 2.7.1. For $k - q = 0$ we get that this coincides with the homomorphisms of 1-motives from $[\mathbb{Z} \rightarrow 0]$ to $[0 \rightarrow J^{2k+1}(X)_K]$, which is 0. For $k - q = 1$ this coincides with the homomorphisms of 1-motives from $[0 \rightarrow \mathbb{G}_m]$ to $[0 \rightarrow J^{2k+1}(X)_K]$, which is also 0. \square

Remark 3.5.3. Lemma 3.5.1 is proven more generally in [1] for the Hodge structure $N^k H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k)) \subset H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$ defined by the k -th step of the coniveau filtration. Namely, if X is defined over a number field $L \subset K$, there is an abelian variety $J_a^{2k+1}(X)$ over L with $T_{\text{dR}}(J_a^{2k+1}(X)) = (N^k H_{\text{dR}}^{2k+1}(X), N^k H_{\mathbb{Z}}^{2k+1}(X), \eta_X^{2k+1, k})$. The proof of Proposition 3.5.2 using $J_a^{2k+1}(X)$ gives the following weak version of the period conjecture:

$$N^k H_{\text{dR}}^{2k+1}(X) \cap N^k H^{2k+1}(X_{\text{an}}, \mathbb{Z}_{\text{an}}(q))_{\text{fr}} = 0 \text{ for every } k \in \mathbb{N} \text{ and every } q \in \mathbb{Z}. \tag{3.6}$$

The assumption in Lemma 3.5.1 amounts to saying that $N^k H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$ is equal to $H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$. This equality holds, for example, for $k = 1$ for uniruled smooth projective threefolds; see [2].

The assumption implies, and under the generalized Hodge conjecture is equivalent to, the fact that the Hodge structure $H^{2k+1}(X_{\text{an}}, \mathbb{Q})$ has Hodge coniveau k , i.e., $H^{2k+1}(X_{\text{an}}, \mathbb{C})$ is the sum of the $(k + 1, k)$ and $(k, k + 1)$ pieces of the Hodge decomposition. Under this weaker condition on the Hodge coniveau one can still prove that $H^{2k+1}(X_{\text{an}}, \mathbb{Q}_{\text{an}}(k))$ is the Hodge structure associated to the abelian variety $J_{\text{alg}}^{2k+1}(X)$ over \mathbb{C} , called the algebraic intermediate Jacobian in $J^{2k+1}(X)$. Unfortunately one lacks the descent to K . See the discussion in [2].

Appendix A. Divisibility properties of motivic cohomology (by B. Kahn)

In this appendix, some results of Colliot-Thélène and Raskind on the \mathcal{K}_2 -cohomology of smooth projective varieties over a separably closed field k are extended to the étale motivic cohomology of smooth, not necessarily projective, varieties over k . Some consequences are drawn, such as the degeneration of the Bloch-Lichtenbaum spectral sequence for any field containing k .

Recall that in [19], Colliot-Thélène and Raskind study the structure of the \mathcal{K}_2 -cohomology groups of a smooth projective variety X over a separably closed field. Following arguments of Bloch [15], their proofs use the Weil conjecture proven by Deligne [22] and the Merkurjev-Suslin theorem [48]. These results and proofs can be reformulated in terms of motivic cohomology, since

$$H_{\text{Zar}}^i(X, \mathcal{K}_2) \simeq H^{i+2}(X, \mathbb{Z}(2))$$

or even in terms of étale motivic cohomology, since

$$H^j(X, \mathbb{Z}(2)) \xrightarrow{\sim} H_{\text{ét}}^j(X, \mathbb{Z}(2)) \text{ for } j \leq 3$$

as follows again from the Merkurjev-Suslin theorem.

If we work in terms of étale motivic cohomology, the recourse to the latter theorem is irrelevant and only the results of [22] are needed; in this form, the results of [19] and their proofs readily extend to étale motivic cohomology of higher weights, as in [41, Prop. 4.17] and [42, Prop. 1] (see also [49, Prop. 1.3]).

Here we generalise these results to the étale motivic cohomology of smooth varieties over a separably closed field: see Theorem A.1.3. This could be reduced by a dévissage to the smooth projective case, using de Jong’s alteration theorem in the style of [40], but it is simpler to reason directly by using cohomology with compact supports, and Weil II [24] rather than Weil I [22]. (I thank H el ene Esnault and Eckart Viehweg for suggesting to use this approach). This descends somewhat to the case where the base field k is not separably closed, yielding information on the Hochschild-Serre filtration on  etale motivic cohomology (Theorem A.2.1). The rest of the appendix is concerned with implications on motivic cohomology of a field K containing a separably closed field: the main result, which uses the norm residue isomorphism theorem of Voevodsky, Rost et al. ([52], see also [31]) is that $H^i(K, \mathbb{Z}(n))$ is divisible for $i \neq n$ (Theorem A.3.1). As an immediate consequence, the “Bloch-Lichtenbaum” spectral sequence of K from motivic cohomology to algebraic K -theory degenerates (Theorem A.4.1). We also show that the cokernel of the cup-product map

$$H^{i-1}(K, \mathbb{Z}(n-1)) \otimes K^* \rightarrow H^i(K, \mathbb{Z}(n))$$

is uniquely divisible for $i < n$ (Theorem A.5.1).

Throughout, motivic cohomology is understood in the sense of Suslin and Voevodsky (hypercohomology of the Suslin-Voevodsky complexes [50]).

A.1. A weight and coniveau argument

Let X be a separated scheme of finite type over a finitely generated field k .

Proposition A.1.1. *Let $n \in \mathbb{Z}$, k_s a separable closure of k and $G = \text{Gal}(k_s/k)$. Let $\overline{X} = X \otimes_k k_s$. Then $H_c^j(\overline{X}, \mathbb{Z}_l(m))^G$ and $H_c^j(\overline{X}, \mathbb{Z}_l(m))_G$ are finite for $j \notin [2m, m+d]$ and any prime number l invertible in k , where $d = \dim X$.*

Proof. Suppose first that $k = \mathbb{F}_q$ is finite. By [21, Cor. 5.5.3 p. 394], the eigenvalues of Frobenius acting on $H_c^j(\overline{X}, \mathbb{Q}_l)$ are algebraic integers which are divisible by q^{j-d} if $j \geq d$. This yields the necessary bound $m \geq j - d$ for an eigenvalue 1. On the other hand, by [24], these eigenvalues have archimedean absolute values $\leq q^{j/2}$: this gives the necessary bound $m \leq j/2$ for an eigenvalue 1. The conclusion follows.

In general, we may choose a regular model S of k , of finite type over $\text{Spec } \mathbb{Z}$, such that X extends to a compactifiable separated morphism of finite type $f : \mathcal{X} \rightarrow S$. By [35, lemma 2.2.2 p. 274 and 2.2.3 p. 277], $R^j f_* \mathbb{Z}_l$ is a constructible \mathbb{Z}_l -sheaf on S and

its formation commutes with any base change. Shrinking S , we may assume that it is locally constant and that l is invertible on S . For a closed point $s \in S$, this gives an isomorphism

$$H_c^j(\overline{X}, \mathbb{Z}_l) \simeq H_c^j(\overline{X}_s, \mathbb{Z}_l)$$

compatible with Galois action, and the result follows from the first case. \square

Corollary A.1.2. *If X is smooth in Proposition A.1.1, then $H^i(\overline{X}, \mathbb{Z}_l(n))^{(G)}$ is finite for $i \notin [n, 2n]$, where the superscript (G) denotes the subset of elements invariant under some open subgroup of G . If X is smooth projective, then $H^i(\overline{X}, \mathbb{Z}_l(n))^{(G)}$ is finite for $i \neq 2n$ and are 0 for almost all l .*

Proof. By Poincaré duality and Proposition A.1.1, $H^i(\overline{X}, \mathbb{Z}_l(n))^{(G)}$ is finite for $i \notin [n, 2n]$; the claim follows since $H^i(\overline{X}, \mathbb{Z}_l(n))$ is a finitely generated \mathbb{Z}_l -module. In the projective case, the Weil conjecture [22] actually gives the finiteness of $H^i(\overline{X}, \mathbb{Z}_l(n))^{(G)}$, hence of $H^i(\overline{X}, \mathbb{Z}_l(n))^{(G)}$, for all $i \neq 2n$. But Gabber’s theorem [25] says that $H^i(\overline{X}, \mathbb{Z}_l(n))$ is torsion-free for almost all l , hence the conclusion. \square

Theorem A.1.3. *Let X be a smooth variety over a separably closed field k of exponential characteristic p . Then, for $i \notin [n, 2n]$, the group $H_{\text{ét}}^i(X, \mathbb{Z}(n))[1/p]$ is an extension of a direct sum T of finite l -groups by a divisible group. If X is projective, this is true for all $i \neq 2n$, and T is finite. If $p > 1$, $H_{\text{ét}}^i(X, \mathbb{Z}(n))$ is uniquely p -divisible for $i < n$. In particular, $H_{\text{ét}}^i(X, \mathbb{Z}(n)) \otimes \mathbb{Q}/\mathbb{Z} = 0$ for $i < n$. For $i \leq 1$, $H_{\text{ét}}^i(X, \mathbb{Z}(n))$ is divisible. The sequence*

$$0 \rightarrow H_{\text{ét}}^{i-1}(X, \mathbb{Q}/\mathbb{Z}(n)) \rightarrow H_{\text{ét}}^i(X, \mathbb{Z}(n)) \rightarrow H_{\text{ét}}^i(X, \mathbb{Z}(n)) \otimes \mathbb{Q} \rightarrow 0$$

is exact for $i < n$.

Proof. Away from p , it is identical to [42, proof of Prop. 1] (which is the projective case) in view of Corollary A.1.2. The unique p -divisibility of $H_{\text{ét}}^i(X, \mathbb{Z}(n))$ for $i < n$ follows from [26, Th. 8.4] and requires no separably closed hypothesis on k . \square

Corollary A.1.4. *Let K be a field containing a separably closed field k . Then, for $i < n$, the sequence*

$$0 \rightarrow H_{\text{ét}}^{i-1}(K, \mathbb{Q}/\mathbb{Z}(n)) \rightarrow H_{\text{ét}}^i(K, \mathbb{Z}(n)) \rightarrow H_{\text{ét}}^i(K, \mathbb{Z}(n)) \otimes \mathbb{Q} \rightarrow 0$$

is exact and the left group has no p -torsion if $p = \text{char } K$.

Proof. We may assume K/k finitely generated. By Theorem A.1.3, this is true for any smooth model of K over k , and we pass to the limit (see [37, Prop. 2.1 b]), or rather its proof, for the commutation of étale motivic cohomology with limits). \square

Remarks A.1.5. 1) At least away from p , the range of “bad” i ’s in Corollary A.1.2 and Theorem A.1.3 is $[n, 2n]$ in general but shrinks to $2n$ when X is projective. If we remove a smooth closed subset, this range becomes $[2n - 1, 2n]$. As the proof of Proposition A.1.1 shows, it depends on the length of the weight filtration on $H^*(\overline{X}, \mathbb{Q}_l)$. If $X = Y - D$, where Y is smooth projective and D is a simple normal crossing divisor with r irreducible components, the range is $[2n - r, 2n]$. It would be interesting to understand the optimal range in general, purely in terms of the geometry of X .

2) Using Proposition A.1.1 or more precisely its proof, one may recover the l -local version of [40, Th. 3] without a recourse to de Jong’s alteration theorem. I don’t see how to get the global finiteness of [40] with the present method, because one does not know whether the torsion of $H_c^j(\overline{X}, \mathbb{Z}_l)$ vanishes for l large when X is not smooth projective.

3) Using a cycle class map to Borel-Moore l -adic homology, one could use Proposition A.1.1 to extend Theorem A.1.3 to higher Chow groups of arbitrary separated k -schemes of finite type. Such a cycle class map was constructed in [36, §1.3]. Note that Borel-Moore l -adic cohomology is dual to l -adic cohomology with compact supports, so the bounds for finiteness are obtained from those of Proposition A.1.1 by changing signs.

A.2. Descent

Theorem A.2.1. *Let X be a smooth variety over a field k ; write k_s for a separable closure of k , X_s for $X \otimes_k k_s$ and G for $\text{Gal}(k_s/k)$. For a complex of sheaves C over $X_{\text{ét}}$, write $F^r H_{\text{ét}}^i(X, C)$ for the filtration on $H_{\text{ét}}^i(X, C)$ induced by the Hochschild-Serre spectral sequence*

$$E_2^{r,s}(C) = H^r(G, H_{\text{ét}}^s(X_s, C)) \Rightarrow H_{\text{ét}}^{r+s}(X, C).$$

Then, for $i < n$, the homomorphism

$$F^r H_{\text{ét}}^{i-1}(X, \mathbb{Q}/\mathbb{Z}(n)) \rightarrow F^r H_{\text{ét}}^i(X, \mathbb{Z}(n))$$

induced by the Bockstein homomorphism β is bijective for $r \geq 3$ and surjective for $r = 1, 2$.

Proof. By the functoriality of $E_m^{r,s}(C)$ with respect to morphisms of complexes, we have a morphism of spectral sequences

$$\delta_m^{r,s} : E_m^{r,s-1}(\mathbb{Q}/\mathbb{Z}(n)) \rightarrow E_m^{r,s}(\mathbb{Z}(n))$$

converging to the Bockstein homomorphisms. By Theorem A.1.3, $\delta_2^{r,i-r}$ is bijective for $r \geq 2$ and surjective for $r = 1$. It follows that, for $m \geq 3$, $\delta_m^{r,i-r}$ is bijective for $r \geq 3$ and surjective for $r = 1, 2$. The conclusion follows. \square

Remarks A.2.2. 1) Of course, $F^r H_{\text{ét}}^i(X, \mathbb{Z}(n))$ is torsion for $r > 0$ by a transfer argument, hence is contained in $\beta H^{i-1}(X, \mathbb{Q}/\mathbb{Z}(n))$. The information of Theorem A.2.1 is that it equals $\beta F^r H^{i-1}(X, \mathbb{Q}/\mathbb{Z}(n))$.

2) For $i \geq n$, we have a similar conclusion for higher values of r , with the same proof: this is left to the reader.

A.3. *Getting the norm residue isomorphism theorem into play*

Recall that for any field K and any $i \leq n$, we have an isomorphism

$$H^i(K, \mathbb{Z}(n)) \xrightarrow{\sim} H_{\text{ét}}^i(K, \mathbb{Z}(n)). \tag{A.1}$$

Indeed, this is seen after localising at l for all prime numbers l . For $l \neq \text{char } K$, this follows from [50,27] and the norm residue isomorphism theorem [52], while for $l = \text{char } K$ it follows from [26]. Finally, $H^i(K, \mathbb{Z}(n)) = 0$ for $i > n$. This yields:

Theorem A.3.1. *Let K be as in Corollary A.1.4. Then, for $i \neq n$, the group in (A.1) is divisible.*

Proof. Again it suffices to prove this statement after tensoring with $\mathbb{Z}_{(l)}$ for all prime numbers l . This is an immediate consequence of Corollary A.1.4 since, by [52], one has an isomorphism for $l \neq \text{char } k$

$$H_{\text{ét}}^{i-1}(K, \mathbb{Q}_l/\mathbb{Z}_l(n)) \simeq K_{i-1}^M(K) \otimes \mathbb{Q}_l/\mathbb{Z}_l(n - i + 1)$$

and the right hand side is divisible. \square

A.4. *Application: degeneration of the Bloch-Lichtenbaum spectral sequence*

Theorem A.4.1. *Let K be as in Corollary A.1.4. Then the Bloch-Lichtenbaum spectral sequence [46, (1.8)]*

$$E_2^{p,q} = H^{p-q}(K, \mathbb{Z}(-q)) \Rightarrow K_{-p-q}(K)$$

degenerates. For any $n > 0$, the map $K_n^M(K) \rightarrow K_n(K)$ is injective with divisible cokernel.

Proof. By the Adams operations, the differentials are torsion [46, Th. 11.7]. By Theorem A.3.1, they vanish on the divisible groups $E_2^{p,q}$ for $p < 0$. But $H^i(K, \mathbb{Z}(n)) = 0$ for $i > n$, so $E_2^{p,q} = 0$ for $p > 0$. The last statement follows from the degeneration plus Theorem A.3.1. \square

Remarks A.4.2. 1) Again by the Adams operations, the filtration on $K_n(K)$ induced by the Bloch-Lichtenbaum spectral sequence splits after inverting $(n - 1)!$ for any field K . On the other hand, we constructed in [39] a canonical splitting of the corresponding spectral sequence with finite coefficients, including the abutment; the hypothesis that K contains a separably closed field is not required there. This implies in particular that the map

$$K_n^M(K)/l^\nu \rightarrow K_n(K)/l^\nu$$

is split injective [39, Th. 1 (c)], hence bijective if K contains a separably closed subfield by Theorem A.4.1. Could it be that the mod l^ν splittings of [39] also exist integrally?

2) As in [39], Theorems A.3.1 and A.4.1 extend to regular semi-local rings of geometric origin containing a separably closed field; the point is that, for such rings R , the groups $H_{\text{ét}}^{i-1}(R, \mathbb{Q}_l/\mathbb{Z}_l(n))$ are divisible by the universal exactness of the Gersten complexes ([28], [18, Th. 6.2.1]).

A.5. The map $H^{i-1}(K, \mathbb{Z}(n - 1)) \otimes K^* \rightarrow H^i(K, \mathbb{Z}(n))$

Theorem A.5.1. *Let K be as in Corollary A.1.4. Then, for $i < n$,*

(i) *The cokernel of the cup-product map*

$$H^1(K, \mathbb{Z}(n - i + 1)) \otimes H^{i-1}(K, \mathbb{Z}(i - 1)) \xrightarrow{\gamma^{i,n}} H^i(K, \mathbb{Z}(n))$$

is uniquely divisible.

(ii) *The cokernel of the cup-product map*

$$H^{i-1}(K, \mathbb{Z}(n - 1)) \otimes K^* \xrightarrow{\delta^{i,n}} H^i(K, \mathbb{Z}(n))$$

is uniquely divisible.

Proof. By (A.1), we may use the étale version of these groups.

(i) Since $H_{\text{ét}}^i(K, \mathbb{Z}(n))$ is divisible by Theorem A.3.1, so is $\text{Coker } \gamma^{i,n}$. Let $\nu \geq 1$ and l prime $\neq \text{char } K$. The diagram

$$\begin{array}{ccc} H_{\text{ét}}^1(K, \mathbb{Z}(n - i + 1)) \otimes H_{\text{ét}}^{i-1}(K, \mathbb{Z}(i - 1)) & \xrightarrow{\gamma^{i,n}} & H_{\text{ét}}^i(K, \mathbb{Z}(n)) \\ \beta \otimes 1 \uparrow & & \beta \uparrow \\ H_{\text{ét}}^0(K, \mathbb{Z}/l^\nu(n - i + 1)) \otimes H_{\text{ét}}^{i-1}(K, \mathbb{Z}(i - 1)) & \xrightarrow{\cup} & H_{\text{ét}}^{i-1}(K, \mathbb{Z}/l^\nu(n)) \end{array}$$

commutes, where β denotes Bockstein. The bottom horizontal map is surjective (even bijective) by the norm residue isomorphism theorem (resp. by [26]). By Theorem A.3.1

again, $H_{\text{ét}}^1(K, \mathbb{Z}(n-i+1))$ is l -divisible, hence so is $H_{\text{ét}}^1(K, \mathbb{Z}(n-i+1)) \otimes H_{\text{ét}}^{i-1}(K, \mathbb{Z}(i-1))$, and Coker $\gamma^{i,n}$ is also l -torsion free by an easy diagram chase.

(ii) Consider the commutative diagram

$$\begin{CD}
 H_{\text{ét}}^1(K, \mathbb{Z}(n-i+1)) \otimes H_{\text{ét}}^{i-2}(K, \mathbb{Z}(i-2)) \otimes K^* @>\gamma^{i-1,n-1} \otimes 1>> H_{\text{ét}}^{i-1}(K, \mathbb{Z}(n-1)) \otimes K^* \\
 @V1 \otimes \cup VV @VV\delta^{i,n} V \\
 H_{\text{ét}}^1(K, \mathbb{Z}(n-i+1)) \otimes H_{\text{ét}}^{i-1}(K, \mathbb{Z}(i-1)) @>\gamma^{i,n}>> H_{\text{ét}}^i(K, \mathbb{Z}(n)).
 \end{CD}$$

Since the left vertical map is surjective, we see that Coker $\delta^{i,n}$ is the quotient of Coker $\gamma^{i,n}$ by the image of the divisible group $H_{\text{ét}}^{i-1}(K, \mathbb{Z}(n-1)) \otimes K^*$ (Theorem A.3.1), hence the claim follows from (i). \square

It would be very interesting to describe $\text{Ker } \delta^{i,n}$, but this seems out of range.

References

- [1] J.D. Achter, S. Casalaina-Martin, C. Vial, Distinguished models of intermediate jacobians, *J. Inst. Math. Jussieu* (2018) 1–28, <https://doi.org/10.1017/S1474748018000245>. Published online by Cambridge University Press: 08 June 2018.
- [2] J.D. Achter, S. Casalaina-Martin, C. Vial, On descending cohomology geometrically, *Compos. Math.* 153 (7) (2017) 1446–1478.
- [3] Y. André, Une introduction aux motifs (motifs purs, motifs mixtes, périodes), *Panoramas et Synthèses*, vol. 17, Société Mathématique de France, Paris, 2004.
- [4] F. Andreatta, L. Barbieri-Viale, A. Bertapelle, Ogus realization of 1-motives, *J. Algebra* 487 (2017) 294–316.
- [5] J. Ayoub, Note sur les opérations de Grothendieck et la réalisation de Betti, *J. Inst. Math. Jussieu* 9 (2) (2010) 225–263.
- [6] J. Ayoub, A guide to (étale) motivic sheaves, in: *Proceedings of the International Congress of Mathematicians—Seoul 2014*, Vol. II, Kyung Moon Sa, Seoul, 2014, pp. 1101–1124.
- [7] J. Ayoub, L’algèbre de Hopf et le groupe de Galois motiviques d’un corps de caractéristique nulle, I, *J. Reine Angew. Math.* 693 (2014) 1–149.
- [8] J. Ayoub, L’algèbre de Hopf et le groupe de Galois motiviques d’un corps de caractéristique nulle, II, *J. Reine Angew. Math.* 693 (2014) 151–226.
- [9] J. Ayoub, Une version relative de la conjecture des périodes de Kontsevich-Zagier, *Ann. of Math.* (2) 181 (3) (2015) 905–992.
- [10] J. Ayoub, L. Barbieri-Viale, 1-motivic sheaves and the Albanese functor, *J. Pure Appl. Algebra* 213 (5) (2009) 809–839.
- [11] L. Barbieri-Viale, A. Bertapelle, Sharp de Rham realization, *Adv. Math.* 222 (4) (2009) 1308–1338.
- [12] L. Barbieri-Viale, B. Kahn, On the derived category of 1-motives, *Astérisque* 381 (2016), xi+254.
- [13] L. Barbieri-Viale, A. Rosenschon, M. Saito, Deligne’s conjecture on 1-motives, *Ann. of Math.* (2) 158 (2) (2003) 593–633.
- [14] A. Bertapelle, Deligne’s duality for de Rham realizations of 1-motives, *Math. Nachr.* 282 (12) (2009) 1637–1655.
- [15] S. Bloch, Torsion algebraic cycles and a theorem of Roitman, *Compos. Math.* 39 (1) (1979) 107–127.
- [16] J.-B. Bost, Algebraization, transcendence, and D -group schemes, *Notre Dame J. Form. Log.* 54 (3–4) (2013) 377–434.
- [17] J.-B. Bost, F. Charles, Some remarks concerning the Grothendieck period conjecture, *J. Reine Angew. Math.* 714 (2016) 175–208.
- [18] J.-L. Colliot-Thélène, R.T. Hoobler, B. Kahn, The Bloch-Ogus-Gabber theorem, in: *Algebraic K-Theory*, Toronto, ON, 1996, in: *Fields Inst. Commun.*, vol. 16, Amer. Math. Soc., Providence, RI, 1997, pp. 31–94.

- [19] J.-L. Colliot-Thélène, W. Raskind, K_2 -cohomology and the second Chow group, *Math. Ann.* 270 (2) (1985) 165–199.
- [20] R. de Jeu, J.D. Lewis, Beilinson’s Hodge conjecture for smooth varieties, *J. K-Theory* 11 (2) (2013) 243–282. With an appendix by Masanori Asakura.
- [21] P. Deligne, Théorèmes d’intégralité, in: *Groupes de monodromie en géométrie algébrique (SGA7)*, Vol. II, in: *Lecture Notes in Mathematics*, vol. 340, Springer-Verlag, Berlin-Heidelberg, 1973, pp. 363–400. Appendix to N. Katz *Le niveau de la cohomologie des intersections complètes*, exposé XXI.
- [22] P. Deligne, La conjecture de Weil. I, *Publ. Math. Inst. Hautes Etudes Sci.* 43 (1974) 273–307.
- [23] P. Deligne, Théorie de Hodge. III, *Publ. Math. Inst. Hautes Etudes Sci.* 44 (1974) 5–77.
- [24] P. Deligne, La conjecture de Weil. II, *Publ. Math. Inst. Hautes Etudes Sci.* 52 (1980) 137–252.
- [25] O. Gabber, Sur la torsion dans la cohomologie l -adique d’une variété, *C. R. Acad. Sci. Paris Sér. I Math.* 297 (3) (1983) 179–182.
- [26] T. Geisser, M. Levine, The K -theory of fields in characteristic p , *Invent. Math.* 139 (3) (2000) 459–493.
- [27] T. Geisser, M. Levine, The Bloch-Kato conjecture and a theorem of Suslin-Voevodsky, *J. Reine Angew. Math.* 530 (2001) 55–103.
- [28] D.R. Grayson, Universal exactness in algebraic K -theory, *J. Pure Appl. Algebra* 36 (2) (1985) 139–141.
- [29] A. Grothendieck, Le groupe de Brauer. II. Théorie cohomologique, in: *Séminaire Bourbaki: années 1964/1965, 1965/1966, exposés 277-312*, vol. 9, Société mathématique de France, 1966, pp. 287–307, Exposé 297.
- [30] A. Grothendieck, On the de Rham cohomology of algebraic varieties, *Publ. Math. Inst. Hautes Etudes Sci.* 29 (1966) 95–103.
- [31] C. Haesemeyer, C.A. Weibel, *The Norm Residue Theorem in Motivic Cohomology*, *Annals of Mathematics Studies*, vol. 200, Princeton University Press, Princeton, NJ, 2019.
- [32] A. Huber, Galois theory of periods, arXiv:1811.06268 [math.NT], 2018.
- [33] A. Huber, S. Müller-Stach, *Periods and Nori Motives*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A, Series of Modern Surveys in Mathematics*, vol. 65, Springer, Cham, 2017. With contributions by Benjamin Friedrich and Jonas von Wangenheim.
- [34] A. Huber, G. Wüstholz, Transcendence and linear relations of 1-periods, arXiv:1805.10104v4 [math.NT], 2019.
- [35] J.-P. Jouanolou, Cohomologie l -adique, in: *Cohomologie l -adique et Fonctions L* , *Séminaire de Géométrie Algébrique du Bois-Marie 1965–1966 (SGA 5)*, in: *Lecture Notes in Mathematics*, vol. 589, Springer-Verlag, Berlin-New York, 1977, pp. 251–281, Exp. VI.
- [36] B. Kahn, The Geisser-Levine method revisited, Preliminary version to [38], K -theory preprint archives 438, <http://www.math.uiuc.edu/K-theory/0438/>.
- [37] B. Kahn, La conjecture de Milnor (d’après V. Voevodsky), in: *Séminaire Bourbaki: volume 1996/1997, exposés 820-834*, in: *Astérisque*, vol. 245, Société mathématique de France, 1997, pp. 379–418, Exposé 834.
- [38] B. Kahn, The Geisser-Levine method revisited and algebraic cycles over a finite field, *Math. Ann.* 324 (3) (2002) 581–617.
- [39] B. Kahn, K -theory of semi-local rings with finite coefficients and étale cohomology, *K-Theory* 25 (2) (2002) 99–138.
- [40] B. Kahn, Some finiteness results for étale cohomology, *J. Number Theory* 99 (1) (2003) 57–73.
- [41] B. Kahn, Classes de cycles motiviques étales, *Algebra Number Theory* 6 (7) (2012) 1369–1407.
- [42] B. Kahn, The Brauer group and indecomposable $(2, 1)$ -cycles, *Compos. Math.* 152 (5) (2016) 1041–1051.
- [43] S. Lang, Transcendental points on group varieties, *Topology* 1 (1962) 313–318.
- [44] F. Lecomte, N. Wach, Le complexe motivique de Rham, *Manuscripta Math.* 129 (1) (2009) 75–90.
- [45] F. Lecomte, N. Wach, Réalisation de Hodge des motifs de Voevodsky, *Manuscripta Math.* 141 (3–4) (2013) 663–697.
- [46] M. Levine, K -theory and motivic cohomology of schemes, preprint, <https://faculty.math.illinois.edu/K-theory/0336/mot.pdf>, 1999.
- [47] C. Mazza, V. Voevodsky, C. Weibel, *Lecture Notes on Motivic Cohomology*, *Clay Mathematics Monographs*, vol. 2, American Mathematical Society/Clay Mathematics Institute, Providence, RI/Cambridge, MA, 2006.
- [48] A.S. Merkurjev, A.A. Suslin, K -cohomology of Severi-Brauer varieties and the norm residue homomorphism, *Izv. Akad. Nauk SSSR Ser. Mat.* 46 (5) (1982) 1011–1046, 1135–1136.

- [49] A. Rosenschon, V. Srinivas, Étale motivic cohomology and algebraic cycles, *J. Inst. Math. Jussieu* 15 (3) (2016) 511–537.
- [50] A. Suslin, V. Voevodsky, Bloch-Kato conjecture and motivic cohomology with finite coefficients, in: *The Arithmetic and Geometry of Algebraic Cycles, Banff, AB, 1998*, in: NATO Sci. Ser. C Math. Phys. Sci., vol. 548, Kluwer Acad. Publ., Dordrecht, 2000, pp. 117–189.
- [51] V. Voevodsky, Cancellation theorem, *Doc. Math. Extra vol. Andrei A. Suslin sixtieth birthday* (2010) 671–685.
- [52] V. Voevodsky, On motivic cohomology with \mathbf{Z}/l -coefficients, *Ann. of Math. (2)* 174 (1) (2011) 401–438.
- [53] V. Vologodsky, Hodge realizations of 1-motives and the derived Albanese, *J. K-Theory* 10 (2) (2012) 371–412.
- [54] M. Waldschmidt, *Nombres transcendants et groupes algébriques*, Astérisque, vol. 69, Société Mathématique de France, Paris, 1979.
- [55] C.A. Weibel, Pic is a contracted functor, *Invent. Math.* 103 (1) (1991) 351–377.
- [56] G. Wüstholz, Some remarks on a conjecture of Waldschmidt, in: *Diophantine Approximations and Transcendental Numbers, Luminy, 1982*, in: *Progr. Math.*, vol. 31, Birkhäuser Boston, Boston, MA, 1983, pp. 329–336.
- [57] G. Wüstholz, Leibniz’ conjecture, periods & motives, in: *Colloquium De Giorgi 2009*, in: *Colloquia*, vol. 3, Ed. Norm, Pisa, 2012, pp. 33–42.