



Research  
Tissue Engineering—Review

## Noncoding RNAs and Their Potential Therapeutic Applications in Tissue Engineering

Shiying Li, Tianmei Qian, Xinghui Wang, Jie Liu, Xiaosong Gu\*

Key Laboratory of Neuroregeneration of Jiangsu and the Ministry of Education, Co-Innovation Center of Neuroregeneration, Nantong University, Nantong, Jiangsu 226001, China

### ARTICLE INFO

#### Article history:

Received 3 November 2016

Revised 19 December 2016

Accepted 11 January 2017

Available online 16 February 2017

#### Keywords:

Tissue engineering

Noncoding RNAs

MicroRNAs

Nerve

Skin

Liver

Vascular system

Muscle

### ABSTRACT

Tissue engineering is a relatively new but rapidly developing field in the medical sciences. Noncoding RNAs (ncRNAs) are functional RNA molecules without a protein-coding function; they can regulate cellular behavior and change the biological milieu of the tissue. The application of ncRNAs in tissue engineering is starting to attract increasing attention as a means of resolving a large number of unmet healthcare needs, although ncRNA-based approaches have not yet entered clinical practice. In-depth research on the regulation and delivery of ncRNAs may improve their application in tissue engineering. The aim of this review is: to outline essential ncRNAs that are related to tissue engineering for the repair and regeneration of nerve, skin, liver, vascular system, and muscle tissue; to discuss their regulation and delivery; and to anticipate their potential therapeutic applications.

© 2017 THE AUTHORS. Published by Elsevier LTD on behalf of the Chinese Academy of Engineering and Higher Education Press Limited Company. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Tissue engineering is a growing area in biomedical research that holds great promise for a range of potential applications in regenerative medicine. It applies the principles of engineering and life sciences in order to develop biological substitutes to repair diseased and injured tissues and organs and restore their functions. The essential characteristic of tissue engineering is the use—whether alone or combined—of living cells, biocompatible materials, biochemical factors (e.g., growth factors, GFs), and physical factors (e.g., cyclic mechanical loading) to create a biomimetic tissue-like structure [1]. The living cells can be derived from donor tissue, albeit with a limited supply; stem or progenitor cells can be used as an alternative cell source [1]. For tissue engineering applications, the cellular microenvironment must allow seed cells to enact their roles, as they do in native tissue, thus ensuring the effective regulation of cell behavior.

Noncoding RNAs (ncRNAs) are a large cluster of RNAs that have

multiple functions in diverse cellular processes, although they do not encode proteins. According to their biological functions, ncRNAs can be divided into infrastructural and regulatory types. Infrastructural RNAs include ribosomal RNAs (rRNAs), transfer RNAs (tRNAs), small nucleolar RNAs (snoRNAs), small nuclear RNAs (snRNAs), guide RNAs (gRNAs), and telomerase RNAs. Regulatory ncRNAs can be classified into microRNAs (miRNAs), small interfering RNAs (siRNAs), long noncoding RNAs (lncRNAs), Piwi-interacting RNAs (piRNAs), promoter-associated RNAs (PARs), and enhancer RNAs (eRNAs) [2–4].

ncRNAs are considered to be a class of molecular targets that may play an important role in tissue engineering. Approaches for ncRNA-based tissue regeneration therapy include altering endogenous cellular activity using ncRNAs, influencing the behavior of resident stem/progenitor cells or cells incorporated into tissue engineered constructs, or modulating the fate of both implanted and endogenous cells with selected ncRNAs. miRNAs, siRNAs, and lncRNAs are the main regulatory ncRNAs that have current

\* Corresponding author.

E-mail address: [nervegu@ntu.edu.cn](mailto:nervegu@ntu.edu.cn)

potential applications. miRNAs are a class of small ncRNAs that have attracted considerable interest; they can influence a wide range of cell functions, including the control of proliferation, migration, differentiation, apoptosis, and other processes, by down-regulating or up-regulating the expression of their target genes [5–7]. lncRNAs are a major class of eukaryotic transcripts that regulate gene expression, possibly by chromatin remodeling, alternative splicing modulation, interacting with proteins to affect protein activity and localization, or serving as a structural component [8,9]. Moreover, lncRNAs may compete for miRNA binding, thus affecting the regulation and function of miRNA target genes. lncRNAs influence almost every step in the life cycle of genes; their best-studied function occurs in the epigenetic regulation of allelic expression [10,11].

## 2. The applications of ncRNAs in tissue engineering

The applications of ncRNAs in tissue engineering have received considerable attention. In the following discussion, we outline a variety of ncRNAs that have been used for neural tissue engineering, liver tissue engineering, skin tissue engineering, muscle tissue engineering, and vascular tissue engineering.

### 2.1. Neural tissue engineering

The nervous system comprises two major components: the central nervous system (CNS) and the peripheral nervous system (PNS). In clinical practice, injuries to the nervous system are commonly encountered. Neural tissue engineering holds great promise for the treatment of diseased or injured nerves, which have a limited capacity to spontaneously regenerate. The poor regenerative capacity of nerve tissues results from the existence of a hostile microenvironment formed by a complex series of events after nerve diseases or injuries. Therefore, an important issue in neural tissue engineering is to manipulate and neutralize the local microenvironment, thus making it more permissive for regeneration.

In neural tissue engineering, the supporting cells that are implanted into the injured nerve may produce GFs or extracellular matrix (ECM) molecules to facilitate nerve regeneration. Neuronal cells and neuroglial cells are the main cell types for neural tissue engineering. Neural stem cells have also been widely used in neural tissue engineering due to their capacity to self-renew and terminally differentiate into mature neural cell types. Therefore, the regulation and potential application of ncRNAs for neural tissue engineering mainly involves neural stem cells, neuronal cells, and neuroglial cells (Table 1) [12–200].

#### 2.1.1. Neural stem cells

**Neural stem/progenitor cells (NSPCs):** The ability to control the self-renewal and differentiation of transplanted NSPCs is critical for the successful application of neural tissue engineering. miR-25, miR-124/124a, miR-200, and miR-106b-25 clusters can promote the neuronal differentiation of NSPCs, and miR-9 and let-7d can promote the neuronal and astrocytic differentiation of neural stem cells (NSCs) [16,17]. miR-34a can obviously increase the numbers of NeuN<sup>+</sup> cells, and can enhance neuronal maturation and the neurite elongation of NSPC-derived neurons. In addition, it is necessary to ensure the subsequent maturation of differentiated cells for proliferation and functionalities. miR-25, miR-137, miR-184, and miR-195 can enhance NSPC proliferation [12–15], which helps provide sufficient cells to restore tissue structure and functionality. miR-137, miR-184, and miR-195 also increase the number of neurons and astrocytes from NSPC differentiation [13–15].

**Mesenchymal stem cells (MSCs):** MSCs, also called bone-marrow stromal cells, are pluripotent stem cells that come from the stromal compartment of the bone marrow. MSCs are increasingly applied in cell-based therapies for various diseases because they are easily obtained from the bone marrow and can be expanded on a large scale by *in vitro* culture. miR-9 and miR-124 can promote neuronal differentiation of MSCs toward mature functional neurons, while miR-128 negatively regulates the differentiation of MSCs into neuron-like cells [16,21,22].

**Table 1**  
ncRNAs with potential applications for tissue engineering.

Cell type and function	ncRNAs
<b>Nerve</b>	
<i>Neural stem/progenitor cells</i>	
Promote proliferation	miR-25 [12]; miR-137 [13]; miR-184 [14]; miR-195 [15]
Induce differentiation	miR-9, siRNA-TLX [16]; let-7d [17]; miR-137 [13]; miR-184 [14]; miR-195 [15]; miR-34a [18]; lncRNA-BDNF-AS, siRNA-BDNF-AS [19]
<i>Mesenchymal stem cells</i>	
Induce differentiation	miR-9 [20]; miR-124 [21]
Reduce differentiation	miR-128 [22]
<i>Neuronal cells</i>	
Inhibit cell death	miR-223 [23]; miR-181c [24]; miR-592 [25]; miR-424 [26]; miR-23a-3p [27]; miR-23a/b, miR-27a/b, siRNA-Apaf-1 [28]
Promote cell death	miR-134 [29]; miR-200c [30]; miR-30a/b [31–33]; miR-124 [34]; miR-711 [35]
Regulate degeneration and apoptosis	miR-20a [36]; miR-29b [37]; miR-146a, siRNA-miR146a [38]
Promote neurite outgrowth	miR-7 [39]; miR-21 [40]; miR-222, siRNA-PTEN [41]; miR-8 [42]; miR-431 [43]; miR-145 [44]; lncRNA-uc.217 [45]; miR-138, siRNA-SIRT1 [46]
<i>Microglial cells</i>	
Inhibit inflammation	let-7c [47]; miR-124, siRNA-C/EBP- $\alpha$ [48]
Promote pro-inflammation	miR-155 [49]
Inhibit activation	let-7c-5p [50]
<i>Astrocytes</i>	
Promote proliferation	miR-17-5p [51]
Inhibit inflammation	miR-146a [52]
Promote activation and differentiation	miR-181 [24]
Inhibit proliferation and migration	lncRNA-SCIR1 [53]
<i>Schwann cells</i>	
Inhibit proliferation and migration	miR-182 [54]; let-7 [55]; miR-1 [56]
Promote proliferation and migration	miR-221, miR-222 [57]

**Table 1** (continued)

Cell type and function	ncRNAs
Inhibit migration	miR-9 [58]
Promote migration	miR-132 [59]
Regulate dedifferentiation and proliferation	miR-34a [60]
Regulate myelination	miR-140 [60]; miR-29a [61]
Regulate fibrinolysis	miR-340 [62]
<b>Liver</b>	
<i>Stem/progenitor cells</i>	
Reduce differentiation	let-7b [63]; let-7f [64]
Induce differentiation	miR-1246, miR-1290, miR-148a, miR-30a, miR-30a, miR-424 [65]
Induce/reduce differentiation	miR-122, siRNA-FoxA1 [66–68]
Reduce differentiation and engraftment	miR-199a-5p, siRNA-SMARCA4, siRNA-MST1 [69]
<i>Hepatocytes</i>	
Promote proliferation	miR-21 [70–72]; miR-211 [73]; lncRNA-URHC [74]
Inhibit proliferation	miR-26a [75]; miR-33 [76]; miR-127, siRNA-Bcl6 [77]; miR-378 [70]; lncRNA-H19 [78]
Regulate cholesterol metabolism	lncRNA-HC [79]
Promote migration	lncRNA-HOTAIR [80]
Inhibit apoptosis and inflammation	lncRNA-TUG1 [81]
<i>Cholangiocytes</i>	
Inhibit cell function	miR-506 [82]
<b>Skin</b>	
<i>Epithelial stem cells</i>	
Inhibit proliferation, induce differentiation	miR-203 [83–86]
Induce differentiation	miR-27b, miR-224 [87]; miR-574-3p, miR-31 [88]
Inhibit proliferation	miR-34, siRNA-P63 [89]; miR-720 [90]; miR-210, siRNA-E <sub>2</sub> F <sub>3</sub> [91]
Improve proliferation, reduce differentiation	miR-125b, siRNA-FGFR <sub>2</sub> [92,93]
Inhibit proliferation, induce differentiation	miR-24, siRNA-PAK4 [94]
<i>Keratinocytes</i>	
Improve migration	miR-205 [95–97]
Inhibit migration, improve proliferation	miR-483-3p [98]
Inhibit migration	miR-198 [99]
Improve migration, inhibit migration	miR-21 [100,101]
Improve migration and proliferation	miR-31, siRNA-EMP-1, siRNA-TGF- $\beta$ [102]
Regulate apoptosis	lncRNA-p21 [103]
<i>Fibroblasts</i>	
Inhibit proliferation	let-7, miR-125 [104]
Improve proliferation	miR-29 [104]; miR-21 [105]; miR-22 [106]; lncRNA-H19 [107]
Induce senescence	miR-152, miR-181a [108]; miR-141 [109]; miR-143 [110]; miR-519a [111]
Improve migration	miR-21 [112]
Induce epithelial-mesenchymal transition	miR-34 [113]; let-7 [114]
Reduce epithelial-mesenchymal transition	miR-200 [113]
Reduce transdifferentiation	miR-146a [115,116]; miR-7 [117]
Decrease extracellular matrix deposition	miR-29 [118,119]; miR-150 [120]; miR-19a [121]
Increase extracellular matrix deposition	miR-92a [122]
Increase mechano-transduction	miR-21 [123–125]
Control collagen stabilization	lncRNA-TSIX, siRNA-TSIX [126]
<i>Melanocytes</i>	
Inhibit apoptosis	miR-17 [127,128]
<b>Muscle</b>	
<i>Myoblasts</i>	
Induce differentiation	lncRNA-MD1 [129]; miR-1 [130]; lncRNA-MyoD [131]; lncRNA-Dum [132]; lncRNA-MUNC [133]; lncRNA-YY1 [134]; miR-29 [135]; miR-181 [136]; lncRNA-H19 [137]; miR-322, miR-503 [138]
Promote proliferation	miR-133 [130]
Promote proliferation and reduce differentiation	lncRNA-sirt1 AS [139]; lncRNA-Malat1 [31,140]; lncRNA-31 [141]
Reduce differentiation	miR-23a, siRNA-Myh [77]
Promote proliferation and migration	miR-486 [142]
<i>Skeletal muscle satellite cells</i>	
Induce differentiation	miR-206 [143]; miR-214, siRNA-Ezh2 [144]; miR-27b [145]
Inhibit proliferation and induce differentiation	miR-1, miR-206 [146]
<i>Skeletal muscle stem cells</i>	
Inhibit differentiation	miR-669a/q [147]
Induce cell-cycle arrest	miR-195, miR-497 [148]
<i>Cardiac progenitor cells</i>	
Control the balance between differentiation and proliferation	miR-1 [149]
Inhibit proliferation and induce differentiation	miR-133a [150]; miR-1, miR-499 [151]
Enhance cardiac remodeling and reduce survival	miR-208a [152,153]; miR-28b [154]
Inhibit apoptosis	miR-138 [155]

**Table 1** (continued)

Cell type and function	ncRNAs
<i>Cardiomyocytes</i>	
Inhibit apoptosis	miR-21 [156]; miR-214 [157]; miR-24 [52]; lncRNA-MHRT [158]
Regulate differentiation and remodeling	miR-21, miR-129, miR-212 [159]
Promote hypertrophy	miR-22 [160]
Promote muscle growth	miR-486 [142]
Induce proliferation	miR-199a, miR-590 [161]; miR-17-92 [162]
Inhibit hypertrophy	lncRNA-H19 [163]
Promote apoptosis	lncRNA-NRF [164]
<i>Cardiac fibroblasts</i>	
Inhibit proliferation	miR-101 [165]
Promote proliferation	lncRNA-H19 [166]
Induce reprogramming to cardiomyocytes	miR-1, miR-133, miR-208, miR-499 [167]
<b>Vascular</b>	
<i>Endothelial cells</i>	
Inhibit proliferation	miR-34a [168]; miR-19a [169]; miR-200c, siRNA-ZEB1 [170]
Promote proliferation and/or migration	miR-126-5p [171]; miR-210 [172]; miR-424 [173]; lncRNA-H19 [81]
Induce senescence and reduce angiogenesis	miR-34a [168]; miR-217 [174]; miR-17-92 [175]; miR-503 [176]; siRNA-ROBO4 [177]
Inhibit migration and angiogenesis	miR-101, siRNA-EZH2 [178]
Promote angiogenesis	miR-17-5p, miR-18a, miR-31, miR-155 [179]; miR-210 [172]; miR-424 [173]; lncRNA-H19 [81]; miR-126 [180]
Promote apoptosis and senescence	miR-200c, siRNA-ZEB1 [170]; PINC [181]
Inhibit proliferation, migration, and apoptosis	miR-503 [176]; miR-155, siRNA-RhoA, siRNA-MYLK [182]
Regulate inflammation	miR-92a, siRNA-KLF4 [183]; miR-663 [184]; miR-10a [185]; miR-712, miR-502, siRNA-TIMP3, siRNA-RECK [186]
<i>Smooth muscle cells</i>	
Induce differentiation and inhibit proliferation	miR-143, miR-145 [187]; lncRNA-MYOSLID [166]
Inhibit proliferation, migration, and apoptosis	miR-503 [176]
Promote migration	miR-712, miR-502, siRNA-TIMP3, siRNA-RECK [186]; miR-24, siRNA-Trb3 [188]
Promote proliferation	miR-24, siRNA-Trb3 [188]; miR-221, miR-222, siRNA-Kip1, siRNA-Kip2 [189]; miR-34a [190]
Promote proliferation and inhibit apoptosis	miR-21 [191]
Inhibit proliferation and promote apoptosis	lncRNA-HIF1A-AS1 [33]; lncRNA-p21 [192]
Reduce elastin levels	miR-29a [193]
Regulate phenotype	siRNA-Jagged1 [194]
<i>Fibroblasts</i>	
Reduce elastin levels	miR-29a [193]
<i>Stem cells</i>	
Induce differentiation	miR-145 [195]; miR-200c, miR-150, siRNA-ZEB1 [196]; miR-1 [197]; miR-10a [198]
Reduce differentiation	siRNA-NOX4, siRNA-TGF- $\beta$ [199]
<i>Endothelial progenitor cells</i>	
Inhibit survival and migration	miR-15a, miR-16 [200]

### 2.1.2. Neuronal cells

After nerve injury, neuronal death is one of the events that influences recovery; therefore, protecting neurons from cell death is important. miR-223, miR-181c, miR-592, miR-424, miR-23a-3p, miR-23a/b, and miR-27a/b can protect neurons from cell death after ischemic brain injury [23–28], while miR-134, miR-200c, miR-30a/b, miR-124, and miR-711 promote neuronal cell death [29–32,34,35,201]. In spinal cord injury, miR-20a causes motor neuron degeneration by targeting *Ngn1* [36], and miR-29b regulates neuronal apoptosis by reducing the expression of Bad, Bim, Puma, and Noxa [37]. Following peripheral nerve injury, overexpression of miR-21 and miR-222 reduces apoptosis and enhances the viability of cultured dorsal root ganglion (DRG) neurons [40,41]. miR-146a mediates apoptosis in DRG neurons under hyperglycemic conditions [38].

The outgrowth of neurites/axons from lesioned neurons is the essence of peripheral nerve regeneration. miR-21 and miR-222 promote neurite outgrowth by targeting *Sprouty2* and *PTEN*, respectively [40,41]. miR-8, miR-431, miR-145, and miR-138 have been shown to play regulatory roles in neurite outgrowth [42–44,46]. In addition, lncRNA-uc.217 regulates neurite outgrowth in DRG neurons following peripheral nerve injury [45].

### 2.1.3. Neuroglial cells

**Microglial cells:** Microglial cells are important cell types in the CNS. *let-7c* suppresses the activation of microglial cells against ischemic damage [47]. miR-124 can decrease the inflammatory response toward nerve injury in order to prevent secondary injury in microglial cells [48]. miR-155 regulates the M1/M2 phenotype

ratio, and further regulates the microglia-mediated neurotoxic response and enhanced axonal regeneration [49].

**Astrocytes:** Astrocytes, which are specialized glial cells, carry out supportive, metabolic, and homeostatic functions in the CNS. The injury of myelinated axons results in axonal degeneration and the accumulation of myelin debris, which contains a variety of axonal growth inhibitors. Removal of these inhibitors via microglia and astrocytes can facilitate axon regeneration. Astrocytes also affect the immune response by mediating different signaling pathways. miR-17-5p promotes the proliferation of astrocytes by targeting the cell-cycle inhibitors p21 and RB1 [51]. miR-181 affects inflammatory cytokine secretion of astrocytes, and modulates astrocyte activation and differentiation [24]. miR-146a carries out an anti-inflammatory role by regulating the release of cytokines from astrocytes, suggesting that miR-146a treatment has the potential to prevent secondary injuries and promote tissue repair [52].

**Schwann cells (SCs):** SCs are the main glial cells in the PNS and play an essential role in peripheral nerve regeneration. SCs can also produce a high level of different GFs such as nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), and so forth [202]. In addition, SCs possess the capacity of phagocytosis to clear away myelin debris. These characteristics make SCs the most widely used support cells in neural tissue engineering for peripheral nerve regeneration. SCs are able to dedifferentiate back to an immature-like state following axonal damage. After dedifferentiation, SCs increase their cell number by proliferation in order for the repair process to begin. The migration of SCs to injured sites is also necessary in order for them to carry out their

functions. Therefore, enhancing the efficiency of SC proliferation and migration during this preparation phase may promote nerve regeneration. miR-182 inhibits the proliferation and migration of SCs by targeting FGF9 and NTM, respectively, at an early stage following sciatic nerve injury [54], while miR-221 and miR-222 promote the proliferation and migration of SCs by targeting *LASS2* [57]. miR-9 is an important functional regulator of SC migration by targeting *CTHRC1*, which in turn regulates Rac1 GTPase [58]. Overexpression of miR-132 facilitates SC migration in order to regulate peripheral nerve regeneration [59]. miR-34a can regulate SC dedifferentiation and proliferation following peripheral nerve injury by targeting Notch1 and cyclin D1 [60].

NGF, the first-discovered member of the neurotrophin family, contributes to neuronal survival and axon growth of the PNS, and ensures the functional integrity of neurons in the CNS. Many experimental studies have determined the beneficial effects of NGF on nerve regeneration. However, clinical applications of NGF are still limited by several constraints, including its deleterious side effects and the complexity of its delivery. let-7 miRNAs have been found to significantly regulate cell proliferation and migration of primary SCs by targeting *NGF* and suppressing its protein translation. Inhibition of let-7 miRNAs increases NGF secretion from primary cultured SCs, and enhances axonal outgrowth from a co-culture of primary SCs and DRG neurons. The inhibitory effect of let-7 miRNAs on SC apoptosis may also serve as an early stress response to nerve injury [55]. In addition, NGF expression that is inhibited by let-7 miRNA can regulate miR-221/222 expression in order to affect the SC phenotype, suggesting that a cascade of let-7 miRNA, through NGF, to miR-221/222 may represent a bypass for the let-7 regulation of SC phenotype modulation. Another neurotrophin, BDNF, is regulated by miR-1, and miR-1 regulates the proliferation and migration of SCs [56].

SCs are responsible for synthesizing myelin sheath in the PNS. miR-140 can modulate axonal myelination in co-cultures of DRG neurons and SCs by targeting the transcription factor *Egr2*, a master regulator of myelination [60]. miR-29a may regulate the myelination of SCs by targeting *PMP22*, a dose-sensitive, disease-associated protein primarily expressed in myelinating SCs [61].

After peripheral nerve injury, the degenerative debris and inflammatory alterations at the injury site may block the elongation of regenerating axons from reaching target organs. miR-340 regulates fibrinolysis, and also influences debris removal and axonal regrowth during sciatic nerve regeneration by targeting tPA, a serine protease with the capability of degrading matrix molecules and cell adhesions [62].

## 2.2. Liver tissue engineering

The liver is one of the largest organs in the human body. Acute or acute-on-chronic failure of the liver results in a life-threatening situation. Survival rates have improved substantially in recent years through advances in critical-care management and the use of liver transplantation. Unfortunately, the number of available liver grafts does not meet the continuously growing need [203,204]. The use of bioengineered livers, instead of procuring organs from brain-dead donors or removing parts of the liver from living donors, is the most promising approach for liver support [205].

Tissue-engineered solutions are under development to temporarily or definitively support or replace a diseased liver. Studies have shown that differentially expressed ncRNAs in the liver are associated with several physiological and pathological processes [206–210]. Some reports have suggested that ncRNAs could improve hepatocyte proliferation or induce the differentiation of stem cells into hepatocyte-like cells (Table 1). miR-21 and miR-

378 promote DNA synthesis in hepatocytes after a partial (2/3) hepatectomy by inhibiting *Btg2* and ornithine decarboxylase, respectively [70]. miR-21 can regulate liver regeneration by influencing the progression through G1 and into the S phase of the cell cycle, by targeting cyclin D1 [71]. The inhibition of miR-33 can obviously increase liver regeneration [76]. miR-26a and miR-127 also regulate hepatocyte proliferation [75,77,211].

The differentiation of stem cells or other types of progenitor cells to hepatocyte-like cells is the second major field in which ncRNAs are already in use, with potential applications for liver tissue engineering. The overexpression of miR-122 can improve hepatic differentiation [66,212], and miR-122 expression gradually increased during the maturation of mouse embryonic stem cells toward hepatocytes [68]. let-7 can regulate the secretion of hepatic-specific factors in human adipose-tissue-derived MSCs [63]. Using a set of miRNAs instead of a single miRNA is another approach to optimize hepatic differentiation. Overexpression of seven miRNAs (miR-1246, miR-1290, miR-148a, miR-30a, miR-424, miR-542-5p, and miR-122) induced human MSC (hMSC) conversion into functionally mature hepatocytes, while a single miRNA could not initiate hepatic differentiation [65].

Compared with the hepatocyte system, less-extensive attention has been paid to the biliary tree. Tissue engineering approaches mimicking biliary function would require a more complex microarchitecture of scaffolds that enable cell-cell interactions, allowing for biliary metabolism and transport [213,214]. Although little is known about the role of miRNAs in regulating cholangiocyte proliferation and function [215], several studies implicated specific miRNAs in the pathogenesis of cholangiocarcinoma. For example, the overexpression of miR-31 in cholangiocarcinoma cells altered RAS/MAPK signaling [216], and miR-138 can regulate cholangiocarcinoma proliferation, cell-cycle control, and migration, possibly by directly targeting *RhoC* [217]. miR-506 regulates the expression of anion exchanger 2 (AE2), and the suppression of miR-506 leads to improved AE2 function in primary biliary cirrhosis cholangiocytes [82]. In addition, the regulation of miR-125b/let-7a expression in cholangiocytes could be a therapeutic approach for biliary diseases [218].

## 2.3. Skin tissue engineering

The demand for clinical intervention for skin loss has increased in recent years. Tissue engineering represents a feasible approach to obtain replacement skin. A key point in the development of engineered skin is controlling cellular behavior. As a new and exciting field of RNA interference, ncRNAs have emerged to overcome the barriers of engineered skin design. The regulation of cell behavior by ncRNA modulation provides a realistic and precise method of affecting cell behavior in bioengineered skin equivalents. Generally speaking, miRNAs may be modulated by overexpression or silencing in skin tissue engineering [219]. The marriage of ncRNA with skin tissue engineering offers the promise of creating safer, more effective skin tissue engineering for critically important clinical conditions [220].

Epithelial stem cells are skin-specific, making them the ideal choice for skin tissue engineering. miRNA can govern the transition of epithelial stem cells from proliferative pools to differentiated keratinocytes [221]. miR-203 plays an important role in maintaining “stemness” in the skin and in other stratified epithelial tissues [222–224]. In addition, miR-203, miR-720, and miR-574-3p can regulate the initiation of epithelial stratification and the maintenance of basal keratinocyte proliferation by directly targeting p63 [89,90].

Keratinocytes, which differentiate from epithelial stem cells, are the key cells that provide a barrier function to the skin. miR-



205 can regulate the migration of keratinocytes [95,96], and miR-198 and miR-21 were found to be associated with chronic wound healing by regulating the migration of keratinocytes [99]. miR-31 promoted the migration and proliferation of keratinocytes [102], and miR-483-3p can affect the growth arrest of keratinocytes during the final steps of re-epithelialization [225].

Fibroblasts are the primary cells of the dermis. let-7 plays an important function in dermal fibroblast proliferation in serum-starved quiescent fibroblasts [104]. miR-22 promotes the proliferation of fibroblasts by regulating several cell-cycle genes [106]. Faraonio et al. [226] described how 24 miRNAs influence senescence-dependent changes in human diploid fibroblasts. Of these, miR-210, miR-376a, miR-486-5p, miR-494, and miR-542-5p can enhance DNA damage and promote senescence. miR-21 was found to regulate fibroblast migration, which is critical to the success of skin tissue engineering [100,101].

It has also been reported that melanocytes can be incorporated into epidermal and dermoepidermal bioengineered skin equivalents [227]. Several studies have shown that some miRNAs could affect the development and progression of melanoma [228–230]. miR-17 has the ability to partially rescue the apoptosis of melanocytes after Dicer ablation by targeting Bim [127]. miR-137, miR-182, and miR-340 can regulate microphthalmia-associated transcription factor in melanoma cells [231–233].

#### 2.4. Muscle tissue engineering

In the human body, muscles develop strength and body work, and are responsible for constant blood pumping (i.e., the cardiac muscle) and posture and movements (i.e., the skeletal muscles). Muscle injury and degeneration account for significant fractions of the global adult mortality and disease burden [234].

Regenerative treatments are not always available for the clinical care of striated muscle disorders; therefore, muscle tissue engineering and stem-cell-based treatments are being promisingly explored. The aim of tissue engineering is the functional recovery of damaged striated muscles by combining biocompatible scaffolds with bioactive molecules and/or cells [235,236]. ncRNAs are emerging as key players in regulating the phenotype of seed cells and the adaptability of both exogenous and resident stem cells [129]. Thus, the roles of ncRNAs (mainly miRNAs) are widely reported in the development of skeletal and cardiac muscle, and in regulating the regenerative potential of muscle progenitors (Table 1).

Studies have shown that muscle-related miRNAs, such as miR-1, miR-133, and miR-206, play a critical role in modulating muscle formation and regeneration [237]. miR-1 promotes differentiation of cultured myoblasts by targeting HDAC4, while miR-133 stimulates myoblast proliferation, mostly through the repression of Srf expression [130]. miR-206 improves skeletal muscle regeneration in Duchenne muscular dystrophy [143] and slows the progression of amyotrophic lateral sclerosis [238]. In addition, lncRNA-MD1 serves as a competing inhibitor to titrate miR-133 and miR-135 away from their targets, MAML1 and MEF2C, further influencing myoblast differentiation [129].

At present, the seed cells of tissue engineering for muscle repair are often stem cells; these provide the degenerating muscle tissue with progenitors to reconstitute genetically suitable myocytes and restore functionality. Sato et al. [148] showed that the transplantation of skeletal muscle stem cells (MuSCs), which were treated with miR-195 and miR-497, improved the efficiency of muscle regeneration through target genes involved in the cell-cycle progression in dystrophin-deficient mice.

miRNAs are found to play an important role in cardiac tissue, where they stimulate cardiomyocyte proliferation in neonatal

mice, rat hearts, and adult mice following myocardial infarction [161]. miRNAs can carry out entwined spatiotemporal roles in the expansion and terminal differentiation of cardiac progenitor cells (CPCs). For example, miR-1 inhibits the expansion of cardiac progenitors by targeting *Hand2*, while miR-1-lacking murine embryos die at embryonic day (E) 10.5 because of severe cardiac malformations [149]. The regulatory role of miRNAs has also been reported in the aberrant switch of fetal programs in response to cardiac stress; miR-21, miR-129, and miR-212 led to hypertrophy and reactivation of a fetal cardiac gene program in rat neonatal cardiomyocytes [159]. As in skeletal myogenesis, miR-22 and miR-133a can affect hypertrophic remodeling by regulating key epigenetic regulators in cardiomyocytes [239].

Overall, as a tissue engineering strategy, the combination of MuSCs, postnatal cardiomyocytes or cardiac stem cells treated with selected ncRNAs, and biomimetic scaffolds should be able to improve their respective engraftment in the damaged tissue, which may make it possible to increase tissue regeneration ability in skeletal muscle disorders and heart diseases, respectively [240].

#### 2.5. Vascular tissue engineering

The vascular system forms an extensive network throughout the body, mediating gas exchange, nutrient transport, and waste removal, as well as delivering cells and mediators in the immune response [241,242]. Vascular tissue engineering is based on the use of scaffolds that can be combined with seed cells such as stem cells, along with other cellular and molecular products, to build vascular conduits, which can be used to restore, maintain, or improve vascular tissue function [241]. In vascular tissue engineering, ncRNAs can enhance the quantity or quality of cells available for cell-based therapeutic angiogenesis, promote stem cell differentiation to vascular cells to be seeded in the scaffolds, improve the function of cells acting at different levels in the vascular scaffold, and correct antiangiogenic molecular defects [243,244].

Endothelial cells (ECs) and smooth muscle cells (SMCs) are essential components of blood vessels; thus, ECs and SMCs are regarded as the main supporting cells in vascular tissue engineering. For this reason, ncRNA application in vascular tissue engineering mainly focuses on the regulatory roles of ncRNAs toward these two cells. ECs line the inner layer of the entire vascular system and ensure vascular homeostasis. They have an essential role during developmental and post-natal angiogenesis [241]. miR-34a and miR-217 promote endothelial senescence, and their inhibitor reduces senescence and increases angiogenesis [168,174]. miR-424, miR-17-5p, miR-18a, miR-31, and miR-155 promote vascular integrity and angiogenesis [173,179]. miR-210 and miR-126-5p promote the proliferation of ECs [171,172]. In addition, ncRNAs can be applied alone or in combination with GFs in order to improve endothelial coverage and endothelial function in vascular tissue engineering.

SMCs make up the middle layer of the vascular wall, and perform the physiological functions of contracting and relaxing vessels and regulating blood pressure and blood flow distribution. They also play important roles in the vascular remodeling processes that follow injury [245–247]. Following vascular injury, SMCs dedifferentiate to promote vessel repair. Healthy SMCs should return to their contractile phenotype once the injury is resolved [246]. While miR-221, miR-222, and miR-24 promote SMC proliferation, miR-143 and miR-145 stimulate differentiation.

miRNAs also regulate the process of stem cell differentiation to vascular cell, and the functional capacities of vascular progenitor cells. Some studies focused on techniques to promote vascular differentiation from stem cells; for example, miR-1, miR-10, and miR-

145 can regulate stem cell differentiation to vascular lineages [195].

To sum up, ncRNAs carry out important roles in the cells associated with tissue engineering; therefore, they may have potential therapeutic applications in tissue engineering as regulators of the function and phenotype of some seed cells. Table 1 provides more information about regulation relationships.

### 3. Application methods

It is now widely recognized that an effective and safe delivery system is key to ncRNA application in tissue engineering. In brief, an ncRNA delivery system should ensure low cytotoxicity and high transfection efficiency, and should also allow controlled release of ncRNAs during the lengthy process of tissue regeneration [248]. Here, we focus on some delivery methods of ncRNAs that have proven useful for tissue engineering applications, according to available works in the literature (Fig. 1).

#### 3.1. Viral transduction

Due to the high transfection efficiency, viral transduction is a good choice for delivering ncRNAs. Several viruses such as retroviruses, lentiviruses, adenoviruses, and adeno-associated viruses (AAVs) can be used in tissue engineering; retroviruses and lentiviruses are widely used.

Retroviruses can be used effectively in various types of dividing cells, including NSPCs. However, the drawback of retroviruses is their inability to infect quiescent cells [249]. In contrast, lentiviruses can transduce non-dividing cells [250–253], which makes these viruses popular for use in the transfection of cells for tissue engineering. The drawbacks of using retroviruses and lentiviruses are that these viruses are unstable when kept in storage and are unable to achieve high viral titers. Adenoviruses can efficiently achieve high viral titers in dividing and non-dividing cells; however, they have significant immunogenicity and toxicity [254,255].

AAVs are a safe and efficient method for the delivery of ncRNAs because of their small size. AAVs are stable in the nucleus, and can maintain high levels of gene expression for months or years [254].


Of these methods, viral transduction provides high transfection efficiency and sustained expression of ncRNAs in transfected cells due to the integration of the viral genome into the host genome. However, genomic integration can lead to uncontrolled insertional mutagenesis, so the use of viral-mediated ncRNA delivery in translational therapeutic applications is limited [256–258].

#### 3.2. Non-viral transduction

In comparison with viral transduction, non-viral delivery methods have unique advantages, including low immunogenicity and mutagenesis, and the capacity of delivering a large quantity of therapeutic agents [256]. Hence, researchers often prefer non-viral approaches to viral transduction.

The liposome is widely used in methods of ncRNA delivery. Its advantages are its biocompatibility, reproducibility, and ease of large-scale production [259]. A number of conventional reagents are currently available, including Lipofectamine<sup>®</sup>, siPORT<sup>™</sup>, HiPerFect, Oligofectamine<sup>™</sup>, MaxSuppressor<sup>™</sup>, DharmaFECT<sup>®</sup>, SilentFect<sup>™</sup>, and NeuroPorter<sup>™</sup>. Despite their variation in structure, these reagents share some common features [256]. In particular, they contain positively charged groups, which interact with the negatively charged sugar-phosphate backbone of RNA molecules. This interaction helps the contact between the RNA/reagent complex and the cell membrane, thereby promoting subsequent cellular uptake [260]. However, lipid-based delivery systems cause toxicity and nonspecific uptake *in vivo*. Moreover, this toxicity, which is accompanied by gene changes, could in turn hinder the desired outcome of the application [261].

Numerous polymers such as poly(lactide-co-glycolide) (PLGA) and polyethylenimine (PEI) are also commonly used as ncRNA



**Fig. 1.** Schematic illustration of ncRNA delivery. (a) Viral transduction of ncRNAs into cells through virus infection using different viral vectors; (b) non-viral transduction through: liposomes; polymers including polyethylene glycol (PEG), poly(lactide-co-glycolide) (PLGA), and polyethylenimine (PEI); chemical modification; or nanoparticles; by which ncRNAs are taken into cells, mainly by cell endocytosis with the help of the above carriers; (c) scaffold-mediated delivery, in which ncRNAs are released into cells from a scaffold loaded with non-viral ncRNA vectors, followed by matrix degradation to enable cell ingrowth.

carriers for gene therapy [262]. PLGA has been widely used for drug delivery because it is safe, biocompatible, and biodegradable. In addition, it can be further optimized to control pharmacodynamics via the surface modification of PLGA nanoparticles [248]. PLGA-based nanoparticles are a potential approach for efficient ncRNA delivery. PEI, another widely used material for ncRNA delivery, is water-soluble and positively charged. The positively charged PEI can encapsulate negatively charged ncRNAs by electrostatic interaction. After endocytosis of these nanocomplexes, the strong buffer effect of the complex results in endosome swelling, and subsequently causes endosome destabilization and the release of ncRNA-encapsulated nanoparticles into the cytosol [262–265].

Several chemical modifications can enhance the stability and affinity of ncRNAs, and thus improve systemic delivery efficacy by increasing the degradation resistance of nucleases in cells and tissues. For example, the 2'-O-methyl, 2'-O-methoxyethyl, or 2'-O-fluorol ncRNAs produced by 2'-OH group modification have enhanced stability and higher binding affinity [266]. Cholesterol was also conjugated into these chemical modifications in order to improve the cellular uptake of ncRNAs; many papers reported that cholesterol-conjugated ncRNAs can enter cells directly via intravenous injection or tissue injection.

### 3.3. Scaffold-mediated delivery

Scaffold-mediated delivery is another common delivery method in tissue engineering and regenerative medicine. For cell engraftment, proliferation, differentiation, and migration at the injured site, biomimetic scaffolds provide a proper microenvironment for tissue repair and regeneration [240]. In addition, topographical features of scaffolds can ensure the sustained delivery of genes and perform additional physical signals to regulate cellular behavior and gene-uptake efficiencies [256]. Moreover, scaffold-mediated delivery can improve local therapy, thus directly enhancing the dose in the target tissue relative to an off-target site. Scaffold architecture also affects cell phenotype. For example, fiber scaffolds—particularly those with aligned fibers—can regulate the maturation of SCs [256,267]. A number of studies have demonstrated the delivery of ncRNAs from tissue engineering scaffolds, thus introducing a novel method for the regulation of gene expression from the delivery platform [220].

## 4. Concluding remarks and future perspectives

### 4.1. Potential risks of ncRNA-based therapy

ncRNA therapeutics, a new concept that differs from conventional chemical drug design, has emerged in recent years in treatment involving tissue engineering and regenerative medicine. However, numerous challenges exist for the therapeutic application of ncRNAs in tissue engineering and regenerative medicine. For example, although the fact that miRNAs target multiple target genes can be an advantage, it also causes ambiguity regarding the scope of the exact genes that are regulated by miRNAs. For clinical evaluation, the functional phenotype and regulatory mechanisms of an miRNA need to be well elucidated and validated. Because of their regulation of a wide variety of cellular events for tissue regeneration, the diversified effects of miRNAs need to be carefully controlled [268]. For example, miR-221 and miR-222 not only affect the proliferation and migration of SCs by regulating *LASS2*, but also regulate the aggressive growth of human glioblastomas by targeting p27<sup>Kip1</sup> [57,269].

For the application of lncRNAs, the challenges are even greater. lncRNAs can regulate a series of cellular processes including

proliferation, differentiation, migration, survival, and apoptosis through diverse mechanisms; however, identification of the functioning mechanisms is limited [270]. In addition, lncRNAs are tissue-specific protein-coding genes, resulting in additional challenges when targeting ncRNAs to a specific tissue or to subcellular compartments.

For ncRNAs with well-defined molecular regulation mechanisms, the delivery system is the greatest challenge. The pathways underlying the delivery process are not well elucidated; therefore, the established guidelines may not always lead to the expected biological phenomenon—a problem that needs to be solved.

### 4.2. Future perspectives

Tissue engineering generates biological substitutes to replace compromised tissues or organs mainly by means of scaffold-based implants, through which seed cells are often introduced. The concomitant introduction of ncRNAs for phenotype modulation of the seed cells is also suggested. To date, siRNAs have been used in diverse scaffold-based tissue engineering strategies; however, the application of other ncRNAs, especially miRNAs, is just beginning.

Although some interesting studies have illustrated the application of ncRNAs in tissue engineering, a better comprehension of the effects and specific targets of ncRNAs in different types of tissues remains to be achieved in view of ncRNA-based therapeutics. In addition, the development of a delivery system to protect ncRNAs against degradation, and to enable them to reach the target tissue/organ, is a major hurdle to overcome. To fulfill these goals, a close collaboration among specialists from different research areas, including medical science, biology, and engineering, is required.

In conclusion, research on ncRNA applications in tissue engineering is still in its infancy. However, ncRNA-based therapy is developing rapidly and has provided a new horizon for tissue engineering strategies in transporting ncRNA safely into seed cells. Given an improved understanding of ncRNA biology and ncRNA delivery, we believe that the utility of ncRNAs in tissue engineering and regenerative medicine will be dramatically improved in the near future.

## Acknowledgements

This work was supported by the National Basic Research Program of China (973 Program, 2014CB542202), the National Hi-Tech Research and Development Program of China (863 Program, 2012AA020502), the National Natural Science Foundation of China (81130080 and 31300879), and the Key University Science Research Project of Jiangsu Province (16KJA310005). It was also a project funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

## Compliance with ethics guidelines

Shiyang Li, Tianmei Qian, Xinghui Wang, Jie Liu, and Xiaosong Gu declare that they have no conflict of interest or financial conflicts to disclose.

## References

- [1] Berthiaume F, Maguire TJ, Yarmush ML. Tissue engineering and regenerative medicine: history, progress, and challenges. *Annu Rev Chem Biomol Eng* 2011;2:403–30.
- [2] Zhou S, Ding F, Gu X. Non-coding RNAs as emerging regulators of neural injury responses and regeneration. *Neurosci Bull* 2016;32(3):253–64.
- [3] Ponting CP, Oliver PL, Reik W. Evolution and functions of long noncoding RNAs. *Cell* 2009;136(4):629–41.
- [4] Kaikkonen MU, Lam MT, Glass CK. Non-coding RNAs as regulators of gene



- expression and epigenetics. *Cardiovasc Res* 2011;90(3):430–40.
- [5] Ivey KN, Srivastava D. microRNAs as regulators of differentiation and cell fate decisions. *Cell Stem Cell* 2010;7(1):36–41.
- [6] Khoshgoo N, Kholdebarin R, Iwaszow BM, Keijzer R. microRNAs and lung development. *Pediatr Pulmonol* 2013;48(4):317–23.
- [7] Small EM, Olson EN. Pervasive roles of microRNAs in cardiovascular biology. *Nature* 2011;469(7330):336–42.
- [8] Amodio N, Di Martino MT, Neri A, Tagliaferri P, Tassone P. Non-coding RNA: a novel opportunity for the personalized treatment of multiple myeloma. *Expert Opin Biol Ther* 2013;13 Suppl 1:S125–37.
- [9] Sana J, Faltejskova P, Svoboda M, Slaby O. Novel classes of non-coding RNAs and cancer. *J Transl Med* 2012;10:103.
- [10] Lin M, Pedrosa E, Shah A, Hrabovsky A, Maqbool S, Zheng D, et al. RNA-Seq of human neurons derived from iPSCs reveals candidate long non-coding RNAs involved in neurogenesis and neuropsychiatric disorders. *PLoS One* 2011;6(9):e23356.
- [11] Ng SY, Johnson R, Stanton LW. Human long non-coding RNAs promote pluripotency and neuronal differentiation by association with chromatin modifiers and transcription factors. *EMBO J* 2012;31(3):522–33.
- [12] Brett JO, Renault VM, Rafalski VA, Webb AE, Brunet A. The microRNA cluster miR-106b–25 regulates adult neural stem/progenitor cell proliferation and neuronal differentiation. *Aging (Albany, NY)* 2011;3(2):108–24.
- [13] Szulwach KE, Li X, Smrt RD, Li Y, Luo Y, Lin L, et al. Cross talk between microRNA and epigenetic regulation in adult neurogenesis. *J Cell Biol* 2010;189(1):127–41.
- [14] Liu C, Teng Z, Santistevan NJ, Szulwach KE, Guo W, Jin P, et al. Epigenetic regulation of miR-184 by MBD1 governs neural stem cell proliferation and differentiation. *Cell Stem Cell* 2010;6(5):433–44.
- [15] Liu C, Teng Z, McQuate AL, Jobe EM, Christ CC, von Hoyningen-Huene SJ, et al. An epigenetic feedback regulatory loop involving microRNA-195 and MBD1 governs neural stem cell differentiation. *PLoS One* 2013;8(1):e51436.
- [16] Zhao C, Sun G, Li S, Shi Y. A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol* 2009;16(4):365–71.
- [17] Zhao C, Sun G, Ye P, Li S, Shi Y. microRNA let-7d regulates the TLX/microRNA-9 cascade to control neural cell fate and neurogenesis. *Sci Rep* 2013;3:1329.
- [18] Aranha MM, Santos DM, Solá S, Steer CJ, Rodrigues CM. miR-34a regulates mouse neural stem cell differentiation. *PLoS One* 2011;6(8):e21396.
- [19] Zheng X, Lin C, Li Y, Ye J, Zhou J, Guo P. Long noncoding RNA BDNF-AS regulates ketamine-induced neurotoxicity in neural stem cell derived neurons. *Biomed Pharmacother* 2016;82:722–8.
- [20] Han R, Kan Q, Sun Y, Wang S, Zhang G, Peng T, et al. miR-9 promotes the neural differentiation of mouse bone marrow mesenchymal stem cells via targeting zinc finger protein 521. *Neurosci Lett* 2012;515(2):147–52.
- [21] Zou D, Chen Y, Han Y, Lv C, Tu G. Overexpression of microRNA-124 promotes the neuronal differentiation of bone marrow-derived mesenchymal stem cells. *Neural Regen Res* 2014;9(12):1241–48.
- [22] Wu R, Tang Y, Zang W, Wang Y, Li M, Du Y, et al. microRNA-128 regulates the differentiation of rat bone mesenchymal stem cells into neuron-like cells by Wnt signaling. *Mol Cell Biochem* 2014;387(1–2):151–8.
- [23] Harraz MM, Eacker SM, Wang X, Dawson TM, Dawson VL. microRNA-223 is neuroprotective by targeting glutamate receptors. *Proc Natl Acad Sci USA* 2012;109(46):18962–7.
- [24] Hutchison ER, Kawamoto EM, Taub DD, Lal A, Abdelmohsen K, Zhang Y, et al. Evidence for miR-181 involvement in neuroinflammatory responses of astrocytes. *Glia* 2013;61(7):1018–28.
- [25] Irmady K, Jackman KA, Padov VA, Shahani N, Martin LA, Cerchetti L, et al. miR-592 regulates the induction and cell death-promoting activity of p75<sup>NTR</sup> in neuronal ischemic injury. *J Neurosci* 2014;34(9):3419–28.
- [26] Liu P, Zhao H, Wang R, Wang P, Tao Z, Gao L, et al. microRNA-424 protects against focal cerebral ischemia and reperfusion injury in mice by suppressing oxidative stress. *Stroke* 2015;46(2):513–9.
- [27] Zhao H, Tao Z, Wang R, Liu P, Yan F, Li J, et al. microRNA-23a-3p attenuates oxidative stress injury in a mouse model of focal cerebral ischemia-reperfusion. *Brain Res* 2014;1592:65–72.
- [28] Chen Q, Xu J, Li L, Li H, Mao S, Zhang F, et al. microRNA-23a/b and microRNA-27a/b suppress A $\alpha$ 1 protein and alleviate hypoxia-induced neuronal apoptosis. *Cell Death Dis* 2014;5:e1132.
- [29] Chi W, Meng F, Li Y, Wang Q, Wang G, Han S, et al. Downregulation of miRNA-134 protects neural cells against ischemic injury in N2A cells and mouse brain with ischemic stroke by targeting HSPA12B. *Neuroscience* 2014;277:111–22.
- [30] Stary CM, Xu L, Sun X, Ouyang YB, White RE, Leong J, et al. microRNA-200c contributes to injury from transient focal cerebral ischemia by targeting Reelin. *Stroke* 2015;46(2):551–6.
- [31] Wang P, Zhang N, Liang J, Li J, Han S, Li J. micro-RNA-30a regulates ischemia-induced cell death by targeting heat shock protein HSPA5 in primary cultured cortical neurons and mouse brain after stroke. *J Neurosci Res* 2015;93(11):1756–68.
- [32] Wang P, Liang J, Li Y, Li J, Yang X, Zhang X, et al. Down-regulation of miRNA-30a alleviates cerebral ischemic injury through enhancing Beclin 1-mediated autophagy. *Neurochem Res* 2014;39(7):1279–91.
- [33] Wang S, Zhang X, Yuan Y, Tan M, Zhang L, Xue X, et al. BRG1 expression is increased in thoracic aortic aneurysms and regulates proliferation and apoptosis of vascular smooth muscle cells through the long non-coding RNA HIF1A-AS1 *in vitro*. *Eur J Cardiothorac Surg* 2015;47(3):439–46.
- [34] Zhu F, Liu J, Li J, Xiao F, Zhang Z, Zhang L. microRNA-124 (miR-124) regulates Ku70 expression and is correlated with neuronal death induced by ischemia/reperfusion. *J Mol Neurosci* 2014;52(1):148–55.
- [35] Sabirzhanov B, Stoica BA, Zhao Z, Loane DJ, Wu J, Dorsey SG, et al. miR-711 upregulation induces neuronal cell death after traumatic brain injury. *Cell Death Differ* 2016;23(4):654–68.
- [36] Jee MK, Jung JS, Im YB, Jung SJ, Kang S. Silencing of miR20a is crucial for Ngn1-mediated neuroprotection in injured spinal cord. *Hum Gene Ther* 2012;23(5):508–20.
- [37] Liu X, Zheng X, Zhang R, Guo Y, Wang J. Combinatorial effects of miR-20a and miR-29b on neuronal apoptosis induced by spinal cord injury. *Int J Clin Exp Pathol* 2015;8(4):3811–8.
- [38] Wang L, Chopp M, Szalad A, Zhang Y, Wang X, Zhang R, et al. The role of miR-146a in dorsal root ganglia neurons of experimental diabetic peripheral neuropathy. *Neuroscience* 2014;259:155–63.
- [39] Liu J, Githinji J, McLaughlin B, Wilczek K, Nolte J. Role of miRNAs in neuronal differentiation from human embryonic stem cell-derived neural stem cells. *Stem Cell Rev* 2012;8(4):1129–37.
- [40] Strickland IT, Richards L, Holmes FE, Wynick D, Uney JB, Wong LF. Axotomy-induced miR-21 promotes axon growth in adult dorsal root ganglion neurons. *PLoS One* 2011;6(8):e23423.
- [41] Zhou S, Shen D, Wang Y, Gong L, Tang X, Yu B, et al. microRNA-222 targeting PTEN promotes neurite outgrowth from adult dorsal root ganglion neurons following sciatic nerve transection. *PLoS One* 2012;7(9):e44768. Erratum in: *PLoS One* 2012;7(9).
- [42] Lu CS, Zhai B, Mauss A, Landgraf M, Gygi S, Van Vactor D. microRNA-8 promotes robust motor axon targeting by coordinate regulation of cell adhesion molecules during synapse development. *Philos Trans R Soc Lond B Biol Sci* 2014;369(1652):20130517.
- [43] Wu D, Murashov AK. microRNA-431 regulates axon regeneration in mature sensory neurons by targeting the Wnt antagonist *Kremen1*. *Front Mol Neurosci* 2013;6:35.
- [44] Zhang H, Zheng S, Zhao J, Zhao W, Zheng L, Zhao D, et al. microRNAs 144, 145, and 214 are down-regulated in primary neurons responding to sciatic nerve transection. *Brain Res* 2011;1383:62–70.
- [45] Yao C, Wang J, Zhang H, Zhou S, Qian T, Ding F, et al. Long non-coding RNA uc.217 regulates neurite outgrowth in dorsal root ganglion neurons following peripheral nerve injury. *Eur J Neurosci* 2015;42(1):1718–25.
- [46] Liu C, Wang R, Saijilafu, Jiao Z, Zhang B, Zhou F. microRNA-138 and SIRT1 form a mutual negative feedback loop to regulate mammalian axon regeneration. *Genes Dev* 2013;27(13):1473–83.
- [47] Banerjee S, Xie N, Cui H, Tan Z, Yang S, Icyuz M, et al. microRNA let-7c regulates macrophage polarization. *J Immunol* 2013;190(12):6542–9.
- [48] Ponomarev ED, Veremeyko T, Barteneva N, Krichevsky AM, Weiner HL. microRNA-124 promotes microglia quiescence and suppresses EAE by deactivating macrophages via the C/EBP- $\alpha$ -PU.1 pathway. *Nat Med* 2011;17(1):64–70.
- [49] Cardoso AL, Guedes JR, Pereira de Almeida L, Pedrosa de Lima MC. miR-155 modulates microglia-mediated immune response by down-regulating SOCS-1 and promoting cytokine and nitric oxide production. *Immunology* 2012;135(1):73–88.
- [50] Ni J, Wang X, Chen S, Liu H, Wang Y, Xu X, et al. microRNA let-7c-5p protects against cerebral ischemia injury via mechanisms involving the inhibition of microglia activation. *Brain Behav Immun* 2015;49:75–85.
- [51] Hong P, Jiang M, Li H. Functional requirement of *Dicer1* and miR-17-5p in reactive astrocyte proliferation after spinal cord injury in the mouse. *Glia* 2014;62(12):2044–60.
- [52] Iyer A, Zurolo E, Prabowo A, Fluiter K, Spliet WG, van Rijen PC, et al. microRNA-146a: a key regulator of astrocyte-mediated inflammatory response. *PLoS One* 2012;7(9):e44789.
- [53] Wang J, Hu B, Cao F, Sun S, Zhang Y, Zhu Q. Down regulation of lncSCIR1 after spinal cord contusion injury in rat. *Brain Res* 2015;1624:314–20.
- [54] Yu B, Qian T, Wang Y, Zhou S, Ding G, Ding F, et al. miR-182 inhibits Schwann cell proliferation and migration by targeting FGF9 and NTM, respectively at an early stage following sciatic nerve injury. *Nucleic Acids Res* 2012;40(20):10356–65.
- [55] Li S, Wang X, Gu Y, Chen C, Wang Y, Liu J, et al. let-7 microRNAs regenerate peripheral nerve regeneration by targeting nerve growth factor. *Mol Ther* 2015;23(3):423–33. Erratum in: *Mol Ther* 2015;23(4):790.
- [56] Yi S, Yuan Y, Chen Q, Wang X, Gong L, Liu J, et al. Regulation of Schwann cell proliferation and migration by miR-1 targeting brain-derived neurotrophic factor after peripheral nerve injury. *Sci Rep* 2016;6:29121.
- [57] Yu B, Zhou S, Wang Y, Qian T, Ding G, Ding F, et al. miR-221 and miR-222 promote Schwann cell proliferation and migration by targeting LASS2 after sciatic nerve injury. *J Cell Sci* 2012;125(11):2675–83.
- [58] Zhou S, Gao R, Hu W, Qian T, Wang N, Ding G, et al. miR-9 inhibits Schwann cell migration by targeting *Cthrc1* following sciatic nerve injury. *J Cell Sci* 2014;127(5):967–76.
- [59] Yao C, Shi X, Zhang Z, Zhou S, Qian T, Wang Y, et al. Hypoxia-induced up-regulation of miR-132 promotes Schwann cell migration after sciatic nerve injury by targeting PRKAG3. *Mol Neurobiol* 2016;53(8):5129–39.
- [60] Viader A, Chang LW, Fahrner T, Nagarajan R, Milbrandt J. microRNAs modulate Schwann cell response to nerve injury by reinforcing transcriptional silencing of dedifferentiation-related genes. *J Neurosci* 2011;31(48):17358–

- 69.
- [61] Verrier JD, Lau P, Hudson L, Murashov AK, Renne R, Notterpek L. Peripheral myelin protein 22 is regulated post-transcriptionally by miRNA-29a. *Glia* 2009;57(12):1265–79.
- [62] Li S, Zhang R, Yuan Y, Yi S, Chen Q, Gong L, et al. miR-340 regulates fibrinolysis and axon regrowth following sciatic nerve injury. *Mol Neurobiol*. Epub 2016 Jun 25.
- [63] Alizadeh E, Akbarzadeh A, Eslaminejad MB, Barzegar A, Hashemzadeh S, Nejati-Koshki K, et al. Up regulation of liver-enriched transcription factors HNF4a and HNF6 and liver-specific microRNA (miR-122) by inhibition of let-7b in mesenchymal stem cells. *Chem Biol Drug Des* 2015;85(3):268–79.
- [64] Davoodian N, Lotfi AS, Soleimani M, Mola SJ, Arjmand S. let-7f microRNA negatively regulates hepatic differentiation of human adipose tissue-derived stem cells. *J Physiol Biochem* 2014;70(3):781–9.
- [65] Cui L, Shi Y, Zhou X, Wang X, Wang J, Lan Y, et al. A set of microRNAs mediate direct conversion of human umbilical cord lining-derived mesenchymal stem cells into hepatocytes. *Cell Death Dis* 2013;4:e918.
- [66] Doddapaneni R, Chawla YK, Das A, Kalra JK, Ghosh S, Chakraborti A. Overexpression of microRNA-122 enhances *in vitro* hepatic differentiation of fetal liver-derived stem/progenitor cells. *J Cell Biochem* 2013;114(7):1575–83.
- [67] Davoodian N, Lotfi AS, Soleimani M, Mowla SJ. microRNA-122 overexpression promotes hepatic differentiation of human adipose tissue-derived stem cells. *J Cell Biochem* 2014;115(9):1582–93.
- [68] Deng X, Qiu R, Wu Y, Li Z, Xie P, Zhang J, et al. Overexpression of miR-122 promotes the hepatic differentiation and maturation of mouse ESCs through a miR-122/FoxA1/HNF4a-positive feedback loop. *Liver Int* 2014; 34(2): 281–95.
- [69] Möbus S, Yang D, Yuan Q, Lüdtkke TH, Balakrishnan A, Sgodda M, et al. microRNA-199a-5p inhibition enhances the liver repopulation ability of human embryonic stem cell-derived hepatic cells. *J Hepatol* 2015;62(1):101–10.
- [70] Song G, Sharma AD, Roll GR, Ng R, Lee AY, Belloch RH, et al. microRNAs control hepatocyte proliferation during liver regeneration. *Hepatology* 2010;51(5):1735–43.
- [71] Ng R, Song G, Roll GR, Frandsen NM, Willenbring H. A microRNA-21 surge facilitates rapid cyclin D1 translation and cell cycle progression in mouse liver regeneration. *J Clin Invest* 2012;122(3):1097–108.
- [72] Bai Y, Yu Z, Luo L, Yi J, Xia Q, Zeng Y. microRNA-21 accelerates hepatocyte proliferation *in vitro* via PI3K/Akt signaling by targeting PTEN. *Biochem Biophys Res Commun* 2014;443(3):802–7.
- [73] Yuan Q, Loya K, Rani B, Möbus S, Balakrishnan A, Laml J, et al. microRNA-221 overexpression accelerates hepatocyte proliferation during liver regeneration. *Hepatology* 2013;57(1):299–310.
- [74] Xu W, Zhang J, Dang Z, Li X, Zhou T, Liu J, et al. Long non-coding RNA URHC regulates cell proliferation and apoptosis via ZAK through the ERK/MAPK signaling pathway in hepatocellular carcinoma. *Int J Biol Sci* 2014;10(7):664–76.
- [75] Zhou J, Ju W, Wang D, Wu L, Zhu X, Guo Z, et al. Down-regulation of microRNA-26a promotes mouse hepatocyte proliferation during liver regeneration. *PLoS One* 2012;7(4):e33577.
- [76] Cirera-Salinas D, Pauta M, Allen RM, Salerno AG, Ramírez CM, Chamorro-Jorganes A, et al. miR-33 regulates cell proliferation and cell cycle progression. *Cell Cycle* 2012;11(5):922–33.
- [77] Pan C, Chen H, Wang L, Yang S, Fu H, Zheng Y, et al. Down-regulation of miR-127 facilitates hepatocyte proliferation during rat liver regeneration. *PLoS One* 2012;7(6):e39151.
- [78] Wang S, Wu X, Liu Y, Yuan J, Yang F, Huang J, et al. Long noncoding RNA H19 inhibits the proliferation of fetal liver cells and the Wnt signaling pathway. *FEBS Lett* 2016;590(4):559–70.
- [79] Lan X, Yan J, Ren J, Zhong B, Li J, Li Y, et al. A novel long noncoding RNA lnc-HC binds hnRNP2B1 to regulate expressions of Cyp7a1 and Abca1 in hepatocytic cholesterol metabolism. *Hepatology* 2016;64(1):58–72.
- [80] Ding C, Cheng S, Yang Z, Lv Z, Xiao H, Du C, et al. Long non-coding RNA HOTAIR promotes cell migration and invasion via down-regulation of RNA binding motif protein 38 in hepatocellular carcinoma cells. *Int J Mol Sci* 2014;15(3):4060–76.
- [81] Su S, Liu J, He K, Zhang M, Feng C, Peng F, et al. Overexpression of the long noncoding RNA TUG1 protects against cold-induced injury of mouse livers by inhibiting apoptosis and inflammation. *FEBS J* 2016;283(7):1261–74.
- [82] Banales JM, Sáez E, Úriz M, Sarvide S, Urribarri AD, Splinter P, et al. Up-regulation of microRNA 506 leads to decreased Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> anion exchanger 2 expression in biliary epithelium of patients with primary biliary cirrhosis. *Hepatology* 2012;56(2):687–97.
- [83] Yi R, Poy MN, Stoffel M, Fuchs E. A skin microRNA promotes differentiation by repressing 'stemness'. *Nature* 2008;452(7184):225–9.
- [84] Lena AM, Shalom-Feuerstein R, Rivetti di Val Cervo P, Aberdam D, Knight RA, Melino G, et al. miR-203 represses 'stemness' by repressing ΔNp63. *Cell Death Differ* 2008;15(7):1187–95.
- [85] Koster MI, Kim S, Mills AA, DeMayo FJ, Roop DR. p63 is the molecular switch for initiation of an epithelial stratification program. *Genes Dev* 2004;18(2):126–31.
- [86] Sonkoly E, Wei T, Pavez Loriè E, Suzuki H, Kato M, Törmä H, et al. Protein kinase C-dependent upregulation of miR-203 induces the differentiation of human keratinocytes. *J Invest Dermatol* 2010;130(1):124–34.
- [87] Hildebrand J, Rütze M, Walz N, Gallinat S, Wenck H, Deppert W, et al. A comprehensive analysis of microRNA expression during human keratinocyte differentiation *in vitro* and *in vivo*. *J Invest Dermatol* 2011;131(1):20–9.
- [88] Peng H, Kaplan N, Hamanaka RB, Katsnelson J, Blatt H, Yang W, et al. microRNA-31/factor-inhibiting hypoxia-inducible factor 1 nexus regulates keratinocyte differentiation. *Proc Natl Acad Sci USA* 2012;109(35):14030–4.
- [89] Antonini D, Russo MT, De Rosa L, Gorrese M, Del Vecchio L, Missero C. Transcriptional repression of miR-34 family contributes to p63-mediated cell cycle progression in epidermal cells. *J Invest Dermatol* 2010;130(5):1249–57.
- [90] Chikh A, Matin RN, Senatore V, Hufbauer M, Lavery D, Raimondi C, et al. iASP/p63 autoregulatory feedback loop is required for the homeostasis of stratified epithelia. *EMBO J* 2011;30(20):4261–73.
- [91] Biswas S, Roy S, Banerjee J, Hussain SRA, Khanna S, Meenakshisundaram G, et al. Hypoxia inducible microRNA 210 attenuates keratinocyte proliferation and impairs closure in a murine model of ischemic wounds. *Proc Natl Acad Sci USA* 2010;107(15):6976–81.
- [92] Zhang L, Stokes N, Polak L, Fuchs E. Specific microRNAs are preferentially expressed by skin stem cells to balance self-renewal and early lineage commitment. *Cell Stem Cell* 2011;8(3):294–308.
- [93] Xu N, Brodin P, Wei T, Meisgen F, Eidsmo L, Nagy N, et al. miR-125b, a microRNA downregulated in psoriasis, modulates keratinocyte proliferation by targeting FGFR2. *J Invest Dermatol* 2011;131(7):1521–9.
- [94] Amelio I, Lena AM, Viticchiè G, Shalom-Feuerstein R, Terrinoni A, Dinsdale D, et al. miR-24 triggers epidermal differentiation by controlling actin adhesion and cell migration. *J Cell Biol* 2012;199(2):347–63.
- [95] Yi R, O'Carroll D, Pasolli HA, Zhang Z, Dietrich FS, Tarakhovskiy A, et al. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat Genet* 2006;38(3):356–62.
- [96] Yu J, Peng H, Ruan Q, Fatima A, Getsios S, Lavker RM. microRNA-205 promotes keratinocyte migration via the lipid phosphatase SHIP2. *FASEB J* 2010;24(10):3950–9.
- [97] Yu J, Ryan DG, Getsios S, Oliveira-Fernandes M, Fatima A, Lavker RM. microRNA-184 antagonizes microRNA-205 to maintain SHIP2 levels in epithelia. *Proc Natl Acad Sci USA* 2008;105(49):19300–5.
- [98] Bertero T, Gastaldi C, Bourget-Ponzio I, Imbert V, Loubat A, Selva E, et al. miR-483-3p controls proliferation in wounded epidermal cells. *FASEB J* 2011;25(9):3092–105.
- [99] Sundaram GM, Common JE, Gopal FE, Srikanta S, Lakshman K, Lunny DP, et al. 'See-saw' expression of microRNA-198 and FSTL1 from a single transcript in wound healing. *Nature* 2013;495(7439):103–6.
- [100] Pastar I, Khan AA, Stojadinovic O, Lebrun EA, Medina MC, Brem H, et al. Induction of specific microRNAs inhibits cutaneous wound healing. *J Biol Chem* 2012;287(35):29324–35.
- [101] Yang X, Wang J, Guo S, Fan K, Li J, Wang Y, et al. miR-21 promotes keratinocyte migration and re-epithelialization during wound healing. *Int J Biol Sci* 2011;7(5):685–90.
- [102] Li D, Li X, Wang A, Meisgen F, Pivarcsi A, Sonkoly E, et al. microRNA-31 promotes skin wound healing by enhancing keratinocyte proliferation and migration. *J Invest Dermatol* 2015;135(6):1676–85.
- [103] Hall JR, Messenger ZJ, Tam HW, Phillips SL, Recio L, Smart RC. Long non-coding RNA lincRNA-p21 is the major mediator of UVB-induced and p53-dependent apoptosis in keratinocytes. *Cell Death Dis* 2015;6(3):e1700.
- [104] Suh EJ, Remillard MY, Legesse-Miller A, Johnson EL, Lemons JM, Chapman TR, et al. A microRNA network regulates proliferative timing and extracellular matrix synthesis during cellular quiescence in fibroblasts. *Genome Biol* 2012;13(12):R121.
- [105] Zhu H, Li C, Bai W, Su L, Liu J, Li Y, et al. microRNA-21 regulates hTERT via PTEN in hypertrophic scar fibroblasts. *PLoS One* 2014;9(5):e97114.
- [106] Polioudakis D, Bhinge AA, Killian PJ, Lee BK, Abell NS, Iyer VR. A Myc-microRNA network promotes exit from quiescence by suppressing the interferon response and cell-cycle arrest genes. *Nucleic Acids Res* 2013;41(4):2239–54.
- [107] Zhang J, Liu C, Wan Y, Peng L, Li W, Qiu J. Long non-coding RNA H19 promotes the proliferation of fibroblasts in keloid scarring. *Oncol Lett* 2016;12(4):2835–9.
- [108] Mancini M, Saintigny G, Mahé C, Annicchiarico-Petruzzelli M, Melino G, Candi E. microRNA-152 and -181a participate in human dermal fibroblasts senescence acting on cell adhesion and remodeling of the extra-cellular matrix. *Aging (Albany, NY)* 2012;4(11):843–53.
- [109] Dimri M, Carroll JD, Cho JH, Dimri GP. microRNA-141 regulates BMI1 expression and induces senescence in human diploid fibroblasts. *Cell Cycle* 2013;12(22):3537–46.
- [110] Bonifacio LN, Jarstfer MB. miRNA profile associated with replicative senescence, extended cell culture, and ectopic telomerase expression in human foreskin fibroblasts. *PLoS One* 2010;5(9):e12519.
- [111] van Kouwenhove M, Kedde M, Agami R. microRNA regulation by RNA-binding proteins and its implications for cancer. *Nat Rev Cancer* 2011;11(9):644–56.
- [112] Madhyastha R, Madhyastha H, Nakajima Y, Omura S, Maruyama M. microRNA signature in diabetic wound healing: promotive role of miR-21 in fibroblast migration. *Int Wound J* 2012;9(4):355–61.
- [113] Lu M, Jolly MK, Levine H, Onuchic JN, Ben-Jacob E. microRNA-based regulation of epithelial-hybrid-mesenchymal fate determination. *Proc Natl Acad Sci USA* 2013;110(45):18144–9.
- [114] Huleihel L, Ben-Yehudah A, Milosevic J, Yu G, Pandit K, Sakamoto K, et al. let-7d microRNA affects mesenchymal phenotypic properties of lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol* 2014;306(6):L534–42.
- [115] Liu Z, Lu C, Cui L, Hu Y, Yu Q, Jiang Y, et al. microRNA-146a modulates TGF-

- $\beta$ 1-induced phenotypic differentiation in human dermal fibroblasts by targeting SMAD4. *Arch Dermatol Res* 2012;304(3):195–202.
- [116] Lan HY. Diverse roles of TGF- $\beta$ /SMADs in renal fibrosis and inflammation. *Int J Biol Sci* 2011;7(7):1056–67.
- [117] Midgley AC, Bowen T, Phillips AO, Steadman R. microRNA-7 inhibition rescues age-associated loss of epidermal growth factor receptor and hyaluronan-dependent differentiation in fibroblasts. *Aging Cell* 2014;13(2):235–44.
- [118] Cheng J, Wang Y, Wang D, Wu Y. Identification of collagen 1 as a post-transcriptional target of miR-29b in skin fibroblasts: therapeutic implication for scar reduction. *Am J Med Sci* 2013;346(2):98–103.
- [119] Maurer B, Stanczyk J, Jüngel A, Akhmetshina A, Trenkmann M, Brock M, et al. microRNA-29, a key regulator of collagen expression in systemic sclerosis. *Arthritis Rheum* 2010;62(6):1733–43.
- [120] Honda N, Jinnin M, Kira-Etoh T, Makino K, Kajihara I, Makino T, et al. miR-150 down-regulation contributes to the constitutive type I collagen overexpression in scleroderma dermal fibroblasts via the induction of integrin  $\beta$ 3. *Am J Pathol* 2013;182(1):206–16.
- [121] Ciechomska M, O'Reilly S, Suwara M, Bogunia-Kubik K, van Laar JM. miR-29a reduces TIMP-1 production by dermal fibroblasts via targeting TGF- $\beta$  activated kinase 1 binding protein 1, implications for systemic sclerosis. *PLoS One* 2014;9(12):e115596.
- [122] Sing T, Jinnin M, Yamane K, Honda N, Makino K, Kajihara I, et al. microRNA-92a expression in the sera and dermal fibroblasts increases in patients with scleroderma. *Rheumatology (Oxford)* 2012;51(9):1550–6.
- [123] Weber M, Baker MB, Moore JP, Searles CD. miR-21 is induced in endothelial cells by shear stress and modulates apoptosis and eNOS activity. *Biochem Biophys Res Commun* 2010;393(4):643–8.
- [124] Wang T, Feng Y, Sun H, Zhang L, Hao L, Shi C, et al. miR-21 regulates skin wound healing by targeting multiple aspects of the healing process. *Am J Pathol* 2012;181(6):1911–20.
- [125] Zanotti S, Gibertini S, Savadori P, Curcio M, Mantegazza R, Morandi L, et al. P20.4 antithetic role of miR-21 and miR-29 in the progression of fibrosis in Duchenne muscular dystrophy. *Neuromuscul Disord* 2013;23(9–10):839–40.
- [126] Wang Z, Jinnin M, Nakamura K, Harada M, Kudo H, Nakayama W, et al. Long non-coding RNA TSIX is upregulated in scleroderma dermal fibroblasts and controls collagen mRNA stabilization. *Exp Dermatol* 2016;25(2):131–6.
- [127] Levy C, Khaled M, Robinson KC, Veguilla RA, Chen PH, Yokoyama S, et al. Lineage-specific transcriptional regulation of DICER by MITF in melanocytes. *Cell* 2010;141(6):994–1005.
- [128] Bouillet P, Cory S, Zhang L, Strasser A, Adams JM. Degenerative disorders caused by Bcl-2 deficiency prevented by loss of its BH3-only antagonist Bim. *Dev Cell* 2001;1(5):645–53.
- [129] Cesana M, Cacchiarelli D, Legnini I, Santini T, Sthandier O, Chinappi M, et al. A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA. *Cell* 2011;147(2):358–69.
- [130] Chen J, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, et al. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet* 2006;38(2):228–33.
- [131] Gong C, Li Z, Ramanujan K, Clay I, Zhang Y, Lemire-Brachat S, et al. A long non-coding RNA, lncMyoD, regulates skeletal muscle differentiation by blocking IMP2-mediated mRNA translation. *Dev Cell* 2015;34(2):181–91.
- [132] Wang L, Zhao Y, Bao X, Zhu X, Kwok YK, Sun K, et al. lncRNA Dum interacts with DNMTs to regulate Dppa2 expression during myogenic differentiation and muscle regeneration. *Cell Res* 2015;25(3):335–50.
- [133] Mueller AC, Cichewicz MA, Dey BK, Layer R, Reon BJ, Gagan JR, et al. MUNC, a long noncoding RNA that facilitates the function of MyoD in skeletal myogenesis. *Mol Cell Biol* 2015;35(3):498–513.
- [134] Zhou L, Sun K, Zhao Y, Zhang S, Wang X, Li Y, et al. *Linc-YY1* promotes myogenic differentiation and muscle regeneration through an interaction with the transcription factor YY1. *Nat Commun* 2015;6:10026.
- [135] Wang H, Garzon R, Sun H, Ladner KJ, Singh R, Dahlman J, et al. NF- $\kappa$ B-YY1-miR-29 regulatory circuitry in skeletal myogenesis and rhabdomyosarcoma. *Cancer Cell* 2008;14(5):369–81.
- [136] Naguibneva I, Ameyar-Zazoua M, Poleskaya A, Ait-Si-Ali S, Groisman R, Souidi M, et al. The microRNA miR-181 targets the homeobox protein Hox-A11 during mammalian myoblast differentiation. *Nat Cell Biol* 2006;8(3):278–84.
- [137] Dey BK, Pfeifer K, Dutta A. The H19 long noncoding RNA gives rise to microRNAs miR-675-3p and miR-675-5p to promote skeletal muscle differentiation and regeneration. *Genes Dev* 2014;28(5):491–501.
- [138] Sarkar S, Dey BK, Dutta A. miR-322/424 and -503 are induced during muscle differentiation and promote cell cycle quiescence and differentiation by down-regulation of Cdc25A. *Mol Biol Cell* 2010;21(13):2138–49.
- [139] Wang G, Wang Y, Xiong Y, Chen X, Ma M, Cai R, et al. *Sirt1* AS lncRNA interacts with its mRNA to inhibit muscle formation by attenuating function of miR-34a. *Sci Rep* 2016;6:21865.
- [140] Watts R, Johnsen VL, Shearer J, Hittel DS. Myostatin-induced inhibition of the long noncoding RNA Malat1 is associated with decreased myogenesis. *Am J Physiol Cell Physiol* 2013;304(10):C995–1001.
- [141] Ballarino M, Cazzella V, D'Andrea D, Grassi L, Bisceglie L, Cipriano A, et al. Novel long noncoding RNAs (lncRNAs) in myogenesis: a miR-31 overlapping lncRNA transcript controls myoblast differentiation. *Mol Cell Biol* 2015;35(4):728–36.
- [142] Alexander MS, Casar JC, Motohashi N, Myers JA, Eisenberg I, Gonzalez RT, et al. Regulation of DMD pathology by an *ankyrin*-encoded miRNA. *Skelet Muscle* 2011;1(1):27.
- [143] Liu N, Williams AH, Maxeiner JM, Bezprozvannaya S, Shelton JM, Richardson JA, et al. microRNA-206 promotes skeletal muscle regeneration and delays progression of Duchenne muscular dystrophy in mice. *J Clin Invest* 2012;122(6):2054–65.
- [144] Juan AH, Kumar RM, Marx JG, Young RA, Sartorelli V. miR-214-dependent regulation of the polycomb protein Ezh2 in skeletal muscle and embryonic stem cells. *Mol Cell* 2009;36(1):61–74.
- [145] Crist CG, Montarras D, Pallafacchina G, Rocancourt D, Cumano A, Conway SJ, et al. Muscle stem cell behavior is modified by microRNA-27 regulation of Pax3 expression. *Proc Natl Acad Sci USA* 2009;106(32):13383–7.
- [146] Chen J, Tao Y, Li J, Deng Z, Yan Z, Xiao X, et al. microRNA-1 and microRNA-206 regulate skeletal muscle satellite cell proliferation and differentiation by repressing Pax7. *J Cell Biol* 2010;190(5):867–79.
- [147] Crippa S, Cassano M, Messina G, Galli D, Galvez BG, Curk T, et al. miR669a and miR669q prevent skeletal muscle differentiation in postnatal cardiac progenitors. *J Cell Biol* 2011;193(7):1197–212.
- [148] Sato T, Yamamoto T, Sehara-Fujisawa A. miR-195/497 induce postnatal quiescence of skeletal muscle stem cells. *Nat Commun* 2014;5:4597.
- [149] Zhao Y, Samal E, Srivastava D. Serum response factor regulates a muscle-specific microRNA that targets *Hand2* during cardiogenesis. *Nature* 2005;436(7048):214–20.
- [150] Liu N, Bezprozvannaya S, Williams AH, Qi X, Richardson JA, Bassel-Duby R, et al. microRNA-133a regulates cardiomyocyte proliferation and suppresses smooth muscle gene expression in the heart. *Genes Dev* 2008;22(23):3242–54.
- [151] Sluijter JP, van Mil A, van Vliet P, Metz CH, Liu J, Doevendans PA, et al. microRNA-1 and -499 regulate differentiation and proliferation in human-derived cardiomyocyte progenitor cells. *Arterioscler Thromb Vasc Biol* 2010;30(4):859–68.
- [152] Montgomery RL, Hullinger TG, Semus HM, Dickinson BA, Seto AG, Lynch JM, et al. Therapeutic inhibition of miR-208a improves cardiac function and survival during heart failure. *Circulation* 2011;124(14):1537–47.
- [153] Callis TE, Pandya K, Seok HY, Tang RH, Tatsuguchi M, Huang ZP, et al. microRNA-208a is a regulator of cardiac hypertrophy and conduction in mice. *J Clin Invest* 2009;119(9):2772–86.
- [154] van Rooij E, Quiat D, Johnson BA, Sutherland LB, Qi X, Richardson JA, et al. A family of microRNAs encoded by myosin genes governs myosin expression and muscle performance. *Dev Cell* 2009;17(5):662–73.
- [155] He S, Liu P, Jian Z, Li J, Zhu Y, Feng Z, et al. miR-138 protects cardiomyocytes from hypoxia-induced apoptosis via MLK3/JNK/c-jun pathway. *Biochem Biophys Res Commun* 2013;441(4):763–9.
- [156] Dong S, Cheng Y, Yang J, Li J, Liu X, Wang X, et al. microRNA expression signature and the role of microRNA-21 in the early phase of acute myocardial infarction. *J Biol Chem* 2009;284(43):29514–25.
- [157] Aurora AB, Mahmoud AI, Luo X, Johnson BA, van Rooij E, Matsuzaki S, et al. microRNA-214 protects the mouse heart from ischemic injury by controlling Ca<sup>2+</sup> overload and cell death. *J Clin Invest* 2012;122(4):1222–32.
- [158] Zhang J, Gao C, Meng M, Tang H. Long noncoding RNA MHRT protects cardiomyocytes against H<sub>2</sub>O<sub>2</sub>-induced apoptosis. *Biomol Ther (Seoul)* 2016;24(1):19–24.
- [159] Thum T, Galuppo P, Wolf C, Fiedler J, Kneitz S, van Laake LW, et al. microRNAs in the human heart: a clue to fetal gene reprogramming in heart failure. *Circulation* 2007;116(3):258–67.
- [160] Gurha P, Wang T, Larimore AH, Sassi Y, Abreu-Goodger C, Ramirez MO, et al. microRNA-22 promotes heart failure through coordinate suppression of PPAR/ERR-nuclear hormone receptor transcription. *PLoS One* 2013;8(9):e75882.
- [161] Eulalio A, Mano M, Dal Ferro M, Zentilin L, Sinagra G, Zacchigna S, et al. Functional screening identifies miRNAs inducing cardiac regeneration. *Nature* 2012;492(7429):376–81.
- [162] Chen J, Huang Z, Seok HY, Ding J, Kataoka M, Zhang Z, et al. miR-17-92 cluster is required for and sufficient to induce cardiomyocyte proliferation in postnatal and adult hearts. *Circ Res* 2013;112(12):1557–66.
- [163] Liu L, An X, Li Z, Song Y, Li L, Zuo S, et al. The H19 long noncoding RNA is a novel negative regulator of cardiomyocyte hypertrophy. *Cardiovasc Res* 2016;111(1):56–65.
- [164] Wang K, Liu F, Liu C, An T, Zhang J, Zhou LY, et al. The long noncoding RNA NRF regulates programmed necrosis and myocardial injury during ischemia and reperfusion by targeting miR-873. *Cell Death Differ* 2016;23(8):1394–405.
- [165] Pan Z, Sun X, Shan H, Wang N, Wang J, Ren J, et al. microRNA-101 inhibited postinfarct cardiac fibrosis and improved left ventricular compliance via the FBJ osteosarcoma oncogene/transforming growth factor- $\beta$ 1 pathway. *Circulation* 2012;126(7):840–50.
- [166] Zhao J, Zhang W, Lin M, Wu W, Jiang P, Tou E, et al. MYOSLID is a novel serum response factor-dependent long noncoding RNA that amplifies the vascular smooth muscle differentiation program. *Arterioscler Thromb Vasc Biol* 2016;36(10):2088–99.
- [167] Jayawardena TM, Egemnazarov B, Finch EA, Zhang L, Payne JA, Pandya K, et al. microRNA-mediated *in vitro* and *in vivo* direct reprogramming of cardiac fibroblasts to cardiomyocytes. *Circ Res* 2012;110(11):1465–73.
- [168] Ito T, Yagi S, Yamakuchi M. microRNA-34a regulation of endothelial senescence. *Biochem Biophys Res Commun* 2010;398(4):735–40.



- [169] Qin X, Wang X, Wang Y, Tang Z, Cui Q, Xi J, et al. microRNA-19a mediates the suppressive effect of laminar flow on cyclin D1 expression in human umbilical vein endothelial cells. *Proc Natl Acad Sci USA* 2010;107(7):3240–4.
- [170] Magenta A, Cencioni C, Fasanaro P, Zaccagnini G, Greco S, Sarra-Ferraris G, et al. miR-200c is upregulated by oxidative stress and induces endothelial cell apoptosis and senescence via ZEB1 inhibition. *Cell Death Differ* 2011;18(10):1628–39.
- [171] Schober A, Nazari-Jahantigh M, Wei Y, Bidzhekov K, Gremse F, Grommes J, et al. microRNA-126-5p promotes endothelial proliferation and limits atherosclerosis by suppressing Dlk1. *Nat Med* 2014;20(4):368–76.
- [172] Fasanaro P, D'Alessandra Y, Di Stefano V, Melchionna R, Romani S, Pompilio G, et al. microRNA-210 modulates endothelial cell response to hypoxia and inhibits the receptor tyrosine kinase ligand Ephrin-A3. *J Biol Chem* 2008;283(23):15878–83.
- [173] Ghosh G, Subramanian IV, Adhikari N, Zhang X, Joshi HP, Basi D, et al. Hypoxia-induced microRNA-424 expression in human endothelial cells regulates HIF- $\alpha$  isoforms and promotes angiogenesis. *J Clin Invest* 2010;120(11):4141–54.
- [174] Menghini R, Casagrande V, Cardellini M, Martelli E, Terrinoni A, Amati F, et al. microRNA 217 modulates endothelial cell senescence via silent information regulator 1. *Circulation* 2009;120(15):1524–32.
- [175] Doebele C, Bonauer A, Fischer A, Scholz A, Reiss Y, Urbich C, et al. Members of the microRNA-17-92 cluster exhibit a cell-intrinsic antiangiogenic function in endothelial cells. *Blood* 2010;115(23):4944–50.
- [176] Caporali A, Meloni M, Völlenklee C, Bonci D, Sala-Newby GB, Addis R, et al. Deregulation of microRNA-503 contributes to diabetes mellitus-induced impairment of endothelial function and reparative angiogenesis after limb ischemia. *Circulation* 2011;123(3):282–91.
- [177] Paul JD, Coulombe KL, Toth PT, Zhang Y, Marsboom G, Bindokas VP, et al. SLIT3-ROBO4 activation promotes vascular network formation in human engineered tissue and angiogenesis *in vivo*. *J Mol Cell Cardiol* 2013;64:124–31.
- [178] Smits M, Mir SE, Nilsson RJ, van der Stoop PM, Niers JM, Marquez VE, et al. Down-regulation of miR-101 in endothelial cells promotes blood vessel formation through reduced repression of EZH2. *PLoS One* 2011;6(1):e16282.
- [179] Suárez Y, Fernández-Hernando C, Yu J, Gerber SA, Harrison KD, Pober JS, et al. Dicer-dependent endothelial microRNAs are necessary for postnatal angiogenesis. *Proc Natl Acad Sci USA* 2008;105(37):14082–7.
- [180] Nicoli S, Standley C, Walker P, Hurlstone A, Fogarty KE, Lawson ND. microRNA-mediated integration of haemodynamics and Vegf signalling during angiogenesis. *Nature* 2010;464(7292):1196–200.
- [181] Jiang C, Fang X, Jiang Y, Shen F, Hu Z, Li X, et al. TNF- $\alpha$  induces vascular endothelial cells apoptosis through overexpressing pregnancy induced noncoding RNA in Kawasaki disease model. *Int J Biochem Cell Biol* 2016;72:118–24.
- [182] Weber M, Kim S, Patterson N, Rooney K, Searles CD. miRNA-155 targets myosin light chain kinase and modulates actin cytoskeleton organization in endothelial cells. *Am J Physiol Heart Circ Physiol* 2014;306(8):H1192–203.
- [183] Fang Y, Davies PF. Site-specific microRNA-92a regulation of Kruppel-like factors 4 and 2 in atherosusceptible endothelium. *Arterioscler Thromb Vasc Biol* 2012;32(4):979–87.
- [184] Ni CW, Qiu H, Jo H. microRNA-663 upregulated by oscillatory shear stress plays a role in inflammatory response of endothelial cells. *Am J Physiol Heart Circ Physiol* 2011;300(5):H1762–9.
- [185] Fang Y, Shi C, Manduchi E, Civelek M, Davies PF. microRNA-10a regulation of proinflammatory phenotype in athero-susceptible endothelium *in vivo* and *in vitro*. *Proc Natl Acad Sci USA* 2010;107(30):13450–5.
- [186] Kim CW, Kumar S, Son DJ, Jang IH, Griendling KK, Jo H. Prevention of abdominal aortic aneurysm by anti-microRNA-712 or anti-microRNA-205 in angiotensin II-infused mice. *Arterioscler Thromb Vasc Biol* 2014;34(7):1412–21.
- [187] Cordes KR, Sheehy NT, White MP, Berry EC, Morton SU, Muth AN, et al. miR-145 and miR-143 regulate smooth muscle cell fate and plasticity. *Nature* 2009;460(7256):705–10.
- [188] Chan MC, Hilyard AC, Wu C, Davis BN, Hill NS, Lal A, et al. Molecular basis for antagonism between PDGF and the TGF $\beta$  family of signalling pathways by control of miR-24 expression. *EMBO J* 2010;29(3):559–73.
- [189] Liu X, Cheng Y, Zhang S, Lin Y, Yang J, Zhang C. A necessary role of miR-221 and miR-222 in vascular smooth muscle cell proliferation and neointimal hyperplasia. *Circ Res* 2009;104(4):476–87.
- [190] Badi I, Burbia I, Ruggeri C, Zeni F, Bertolotti M, Scopece A, et al. microRNA-34a induces vascular smooth muscle cells senescence by SIRT1 downregulation and promotes the expression of age-associated pro-inflammatory secretory factors. *J Gerontol A Biol Sci Med Sci* 2015;70(11):1304–11.
- [191] Ji R, Cheng Y, Yue J, Yang J, Liu X, Chen H, et al. microRNA expression signature and antisense-mediated depletion reveal an essential role of microRNA in vascular neointimal lesion formation. *Circ Res* 2007;100(11):1579–88.
- [192] Wu G, Cai J, Han Y, Chen J, Huang ZP, Chen C, et al. lincRNA-p21 regulates neointima formation, vascular smooth muscle cell proliferation, apoptosis, and atherosclerosis by enhancing p53 activity. *Circulation* 2014;130(17):1452–65.
- [193] Zhang P, Huang A, Ferruzzi J, Mechem RP, Starcher BC, Tellides G, et al. Inhibition of microRNA-29 enhances elastin levels in cells haploinsufficient for elastin and in bioengineered vessels—brief report. *Arterioscler Thromb Vasc Biol* 2012;32(3):756–9.
- [194] Bhattacharyya A, Lin S, Sandig M, Mequanint K. Regulation of vascular smooth muscle cell phenotype in three-dimensional coculture system by Jagged1-selective Notch3 signaling. *Tissue Eng Part A* 2014;20(7–8):1175–87.
- [195] Yamaguchi S, Yamahara K, Homma K, Suzuki S, Fujii S, Morizane R, et al. The role of microRNA-145 in human embryonic stem cell differentiation into vascular cells. *Atherosclerosis* 2011;219(2):468–74.
- [196] Luo Z, Wen G, Wang G, Pu X, Ye S, Xu Q, et al. microRNA-200c and -150 play an important role in endothelial cell differentiation and vasculogenesis by targeting transcription repressor ZEB1. *Stem Cells* 2013;31(9):1749–62.
- [197] Xie C, Huang H, Sun X, Guo Y, Hamblin M, Ritchie RP, et al. microRNA-1 regulates smooth muscle cell differentiation by repressing Kruppel-like factor 4. *Stem Cells Dev* 2011;20(2):205–10.
- [198] Huang H, Xie C, Sun X, Ritchie RP, Zhang J, Chen Y. miR-10a contributes to retinoid acid-induced smooth muscle cell differentiation. *J Biol Chem* 2010;285(13):9383–9.
- [199] Xiao Q, Luo Z, Pepe AE, Margariti A, Zeng L, Xu Q. Embryonic stem cell differentiation into smooth muscle cells is mediated by Nox4-produced H<sub>2</sub>O<sub>2</sub>. *Am J Physiol Cell Physiol* 2009;296(4):C711–23.
- [200] Spinetti G, Fortunato O, Caporali A, Shantikumar S, Marchetti M, Meloni M, et al. microRNA-15a and microRNA-16 impair human circulating proangiogenic cell functions and are increased in the proangiogenic cells and serum of patients with critical limb ischemia. *Circ Res* 2013;112(2):335–46.
- [201] Han F, Huo Y, Huang C, Chen C, Ye J. microRNA-30b promotes axon outgrowth of retinal ganglion cells by inhibiting Semaphorin3A expression. *Brain Res* 2015;1611:65–73.
- [202] Gu X, Ding F, Yang Y, Liu J. Construction of tissue engineered nerve grafts and their application in peripheral nerve regeneration. *Prog Neurobiol* 2011;93(2):204–30.
- [203] Nebrig M, Neuhaus P, Pascher A. Advances in the management of the explanted donor liver. *Nat Rev Gastroenterol Hepatol* 2014;11(8):489–96.
- [204] Dutkowski P, Linecker M, DeOliveira ML, Müllhaupt B, Clavien PA. Challenges to liver transplantation and strategies to improve outcomes. *Gastroenterology* 2015;148(2):307–23.
- [205] Uygun BE, Yarmush ML. Engineered liver for transplantation. *Curr Opin Biotechnol* 2013;24(5):893–9.
- [206] McDaniel K, Herrera L, Zhou T, Francis H, Han Y, Levine P, et al. The functional role of microRNAs in alcoholic liver injury. *J Cell Mol Med* 2014;18(2):197–207.
- [207] Padgett KA, Lan RY, Leung PC, Lleo A, Dawson K, Pfeiff J, et al. Primary biliary cirrhosis is associated with altered hepatic microRNA expression. *J Autoimmun* 2009;32(3–4):246–53.
- [208] Szabo G, Bala S. MicroRNAs in liver disease. *Nat Rev Gastroenterol Hepatol* 2013;10(9):542–52.
- [209] Janssen HL, Reesink HW, Lawitz EJ, Zeuzem S, Rodriguez-Torres M, Patel K, et al. Treatment of HCV infection by targeting microRNA. *N Engl J Med* 2013;368(18):1685–94.
- [210] Finch ML, Marquardt JU, Yeoh GC, Callus BA. Regulation of microRNAs and their role in liver development, regeneration and disease. *Int J Biochem Cell Biol* 2014;54:288–303.
- [211] Chen X, Murad M, Cui Y, Yao L, Venugopal SK, Dawson K, et al. miRNA regulation of liver growth after 50% partial hepatectomy and small size grafts in rats. *Transplantation* 2011;91(3):293–9.
- [212] Davoodian N, Lotfi AS, Soleimani M, Mowla SJ. microRNA-122 overexpression promotes hepatic differentiation of human adipose tissue-derived stem cells. *J Cell Biochem* 2014;115(9):1582–93.
- [213] Kasuya J, Tanishita K. Microporous membrane-based liver tissue engineering for the reconstruction of three-dimensional functional liver tissues *in vitro*. *Biomater* 2012;2(4):290–5.
- [214] Katsuda T, Kojima N, Ochiya T, Sakai Y. Biliary epithelial cells play an essential role in the reconstruction of hepatic tissue with a functional bile ductular network. *Tissue Eng Part A* 2013;19(21–22):2402–11.
- [215] O'Hara SP, Gradilone SA, Masyuk TV, Tabibian JH, LaRusso NF. microRNAs in cholangiopathies. *Curr Pathobiol Rep* 2014;2(3):133–42.
- [216] Hu C, Huang F, Deng G, Nie W, Huang W, Zeng X. miR-31 promotes oncogenesis in intrahepatic cholangiocarcinoma cells via the direct suppression of RASA1. *Exp Ther Med* 2013;6(5):1265–70.
- [217] Wang Q, Tang H, Yin S, Dong C. Downregulation of microRNA-138 enhances the proliferation, migration and invasion of cholangiocarcinoma cells through the upregulation of RhoC/p-ERK/MMP-2/MMP-9. *Oncol Rep* 2013;29(5):2046–52.
- [218] Glaser S, Meng F, Han Y, Onori P, Chow BK, Francis H, et al. Secretin stimulates biliary cell proliferation by regulating expression of microRNA 125b and microRNA let7a in mice. *Gastroenterology* 2014;146(7):1795–808.
- [219] van Rooij E, Purcell AL, Levin AA. Developing microRNA therapeutics. *Circ Res* 2012;110(3):496–507.
- [220] Miller KJ, Brown DA, Ibrahim MM, Ramchal TD, Levinson H. microRNAs in skin tissue engineering. *Adv Drug Deliv Rev* 2015;88:16–36.
- [221] Yi R, Fuchs E. microRNA-mediated control in the skin. *Cell Death Differ* 2010;17(2):229–35.
- [222] Mills AA, Zheng B, Wang X, Vogel H, Roop DR, Bradley A. p63 is a p53 homologue required for limb and epidermal morphogenesis. *Nature* 1999;398(6729):708–13.
- [223] Yang A, Schweitzer R, Sun D, Kaghad M, Walker N, Bronson RT, et al. p63 is essential for regenerative proliferation in limb, craniofacial and epithelial



- development. *Nature* 1999;398(6729):714–8.
- [224] Senoo M, Pinto F, Crum CP, McKeon F. p63 is essential for the proliferative potential of stem cells in stratified epithelia. *Cell* 2007;129(3):523–36.
- [225] Bertero T, Gastaldi C, Bourget-Ponzio I, Mari B, Meneguzzi G, Barbry P, et al. CDC25A targeting by miR-483-3p decreases CCND-CDK4/6 assembly and contributes to cell cycle arrest. *Cell Death Differ* 2013;20(6):800–11.
- [226] Faraonio R, Salerno P, Passaro F, Sedia C, Iaccio A, Bellelli R, et al. A set of miRNAs participates in the cellular senescence program in human diploid fibroblasts. *Cell Death Differ* 2012;19(4):713–21.
- [227] Rehder J, Souto LRM, Issa CMBM, Puzzi MB. Model of human epidermis reconstructed *in vitro* with keratinocytes and melanocytes on dead de-epidermized human dermis. *Sao Paulo Med J* 2004;122(1):22–5.
- [228] Müller DW, Bosserhoff AK. Integrin  $\beta_3$  expression is regulated by let-7a miRNA in malignant melanoma. *Oncogene* 2008;27(52):6698–706.
- [229] Schultz J, Lorenz P, Gross G, Ibrahim S, Kunz M. microRNA let-7b targets important cell cycle molecules in malignant melanoma cells and interferes with anchorage-independent growth. *Cell Res* 2008;18(5):549–57.
- [230] Müller DW, Rehli M, Bosserhoff AK. miRNA expression profiling in melanocytes and melanoma cell lines reveals miRNAs associated with formation and progression of malignant melanoma. *J Invest Dermatol* 2009;129(7):1740–51.
- [231] Segura MF, Hanniford D, Menendez S, Reavie L, Zou X, Alvarez-Diaz S, et al. Aberrant miR-182 expression promotes melanoma metastasis by repressing FOXO3 and microphthalmia-associated transcription factor. *Proc Natl Acad Sci USA* 2009;106(6):1814–9.
- [232] Bemis LT, Chen R, Amato CM, Classen EH, Robinson SE, Coffey DG, et al. microRNA-137 targets microphthalmia-associated transcription factor in melanoma cell lines. *Cancer Res* 2008;68(5):1362–8.
- [233] Goswami S, Tarapore RS, Teslaa JJ, Grinblat Y, Setaluri V, Spiegelman VS. microRNA-340-mediated degradation of microphthalmia-associated transcription factor mRNA is inhibited by the coding region determinant-binding protein. *J Biol Chem* 2010;285(27):20532–40. Retraction in: Goswami S, Tarapore RS, Teslaa JJ, Grinblat Y, Setaluri V, Spiegelman VS. *J Biol Chem* 2014;289(17):11859.
- [234] Quattrocchi M, Sampaolesi M. The mesmiRizing complexity of microRNAs for striated muscle tissue engineering. *Adv Drug Deliv Rev* 2015;88:37–52.
- [235] Cezar CA, Mooney DJ. Biomaterial-based delivery for skeletal muscle repair. *Adv Drug Deliv Rev* 2015;84:188–97.
- [236] Pascual-Gil S, Garbayo E, Díaz-Herráez P, Prosper F, Blanco-Prieto MJ. Heart regeneration after myocardial infarction using synthetic biomaterials. *J Control Release* 2015;203:23–38.
- [237] Luo W, Nie Q, Zhang X. microRNAs involved in skeletal muscle differentiation. *J Genet Genomics* 2013;40(3):107–16.
- [238] Williams AH, Valdez G, Moresi V, Qi X, McAnally J, Elliott JL, et al. microRNA-206 delays ALS progression and promotes regeneration of neuromuscular synapses in mice. *Science* 2009;326(5959):1549–54.
- [239] Chaturvedi P, Tyagi SC. Epigenetic mechanisms underlying cardiac degeneration and regeneration. *Int J Cardiol* 2014;173(1):1–11.
- [240] Gori M, Trombetta M, Santini D, Rainer A. Tissue engineering and microRNAs: future perspectives in regenerative medicine. *Expert Opin Biol Ther* 2015;15(11):1601–22.
- [241] Caputo M, Saif J, Rajakaruna C, Brooks M, Angelini GD, Emanuelli C. microRNAs in vascular tissue engineering and post-ischemic neovascularization. *Adv Drug Deliv Rev* 2015;88:78–91.
- [242] Pober JS, Sessa WC. Evolving functions of endothelial cells in inflammation. *Nat Rev Immunol* 2007;7(10):803–15.
- [243] Yang W, Yang D, Na S, Sandusky G, Zhang Q, Zhao G. Dicer is required for embryonic angiogenesis during mouse development. *J Biol Chem* 2005;280(10):9330–5.
- [244] Kane NM, Howard L, Descamps B, Meloni M, McClure J, Lu R, et al. Role of microRNAs 99b, 181a, and 181b in the differentiation of human embryonic stem cells to vascular endothelial cells. *Stem Cells* 2012;30(4):643–54.
- [245] Zhang L, Zhou Y, Zhu J, Xu Q. An updated view on stem cell differentiation into smooth muscle cells. *Vascul Pharmacol* 2012;56(5–6):280–7.
- [246] Owens GK, Kumar MS, Wamhoff BR. Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 2004;84(3):767–801.
- [247] Lacolley P, Regnault V, Nicoletti A, Li Z, Michel JB. The vascular smooth muscle cell in arterial pathology: a cell that can take on multiple roles. *Cardiovasc Res* 2012;95(2):194–204.
- [248] Peng B, Chen Y, Leong KW. microRNA delivery for regenerative medicine. *Adv Drug Deliv Rev* 2015;88:108–22.
- [249] Dufait I, Liechtenstein T, Lanna A, Bricogne C, Laranga R, Padella A, et al. Retroviral and lentiviral vectors for the induction of immunological tolerance. *Scientifica (Cairo)* 2012;2012:694137.
- [250] McBride JL, Behrstock SP, Chen EY, Jakel RJ, Siegel I, Svendsen CN, et al. Human neural stem cell transplants improve motor function in a rat model of Huntington's disease. *J Comp Neurol* 2004;475(2):211–9.
- [251] Englund U, Fricker-Gates RA, Lundberg C, Björklund A, Victorin K. Transplantation of human neural progenitor cells into the neonatal rat brain: extensive migration and differentiation with long-distance axonal projections. *Exp Neurol* 2002;173(1):1–21.
- [252] Dugas JC, Cuellar TL, Scholze A, Ason B, Ibrahim A, Emery B, et al. Dicer1 and miR-219 are required for normal oligodendrocyte differentiation and myelination. *Neuron* 2010;65(5):597–611.
- [253] McTigue DM, Wei P, Stokes BT. Proliferation of NG2-positive cells and altered oligodendrocyte numbers in the contused rat spinal cord. *J Neurosci* 2001;21(10):3392–400.
- [254] Beavers KR, Nelson CE, Duvall CL. miRNA inhibition in tissue engineering and regenerative medicine. *Adv Drug Deliv Rev* 2015;88:123–37.
- [255] Lin E, Nemunaitis J. Oncolytic viral therapies. *Cancer Gene Ther* 2004;11(10):643–64.
- [256] Nguyen LH, Diao HJ, Chew SY. microRNAs and their potential therapeutic applications in neural tissue engineering. *Adv Drug Deliv Rev* 2015;88:53–66.
- [257] Yau WW, Rujitanaroj PO, Lam L, Chew SY. Directing stem cell fate by controlled RNA interference. *Biomaterials* 2012;33(9):2608–28.
- [258] Muthiah M, Park IK, Cho CS. Nanoparticle-mediated delivery of therapeutic genes: focus on miRNA therapeutics. *Expert Opin Drug Deliv* 2013;10(9):1259–73.
- [259] Torrecilla J, Rodríguez-Gascón A, Solinís MÁ, del Pozo-Rodríguez A. Lipid nanoparticles as carriers for RNAi against viral infections: current status and future perspectives. *Biomed Res Int* 2014;2014:161794.
- [260] Dalby B, Cates S, Harris A, Ohki EC, Tilkins ML, Price PJ, et al. Advanced transfection with Lipofectamine 2000 reagent: primary neurons, siRNA, and high-throughput applications. *Methods* 2004;33(2):95–103.
- [261] Nguyen LT, Atobe K, Barichello JM, Ishida T, Kiwada H. Complex formation with plasmid DNA increases the cytotoxicity of cationic liposomes. *Biol Pharm Bull* 2007;30(4):751–7.
- [262] Chen Y, Gao D, Huang L. *In vivo* delivery of miRNAs for cancer therapy: challenges and strategies. *Adv Drug Deliv Rev* 2015;81:128–41.
- [263] Boussif O, Lezoualc'h F, Zanta MA, Mergny MD, Scherman D, Demeneix B, et al. A versatile vector for gene and oligonucleotide transfer into cells in culture and *in vivo*: polyethylenimine. *Proc Natl Acad Sci USA* 1995;92(16):7297–301.
- [264] Park TG, Jeong JH, Kim SW. Current status of polymeric gene delivery systems. *Adv Drug Deliv Rev* 2006;58(4):467–86.
- [265] Akinc A, Thomas M, Klibanov AM, Langer R. Exploring polyethylenimine-mediated DNA transfection and the proton sponge hypothesis. *J Gene Med* 2005;7(5):657–63.
- [266] Davis S, Lollo B, Freier S, Esau C. Improved targeting of miRNA with anti-sense oligonucleotides. *Nucleic Acids Res* 2006;34(8):2294–304.
- [267] Chew SY, Mi R, Hoke A, Leong KW. The effect of the alignment of electrospun fibrous scaffolds on Schwann cell maturation. *Biomaterials* 2008;29(6):653–61.
- [268] Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor suppressors. *Dev Biol* 2007;302(1):1–12.
- [269] Gillies JK, Lorimer IA. Regulation of p27<sup>Kip1</sup> by miRNA 221/222 in glioblastoma. *Cell Cycle* 2007;6(16):2005–9.
- [270] Chen Q, Wei C, Wang Z, Sun M. Long non-coding RNAs in anti-cancer drug resistance. *Oncotarget*. Epub 2016 Oct 4.