Contents lists available at ScienceDirect

Discrete Mathematics

journal homepage: www.elsevier.com/locate/disc

A classification of cubic *s*-regular graphs of order 16*p*

Ju-Mok Oh

Combinatorial and Computational Mathematics Center, Pohang University of Science and Technology, Pohang, 790-784, Republic of Korea

ARTICLE INFO

Article history: Received 25 November 2007 Received in revised form 10 August 2008 Accepted 1 September 2008 Available online 21 September 2008

Keywords: Cubic s-regular graph Covering graph The Möbius–Kantor graph

1. Introduction

Throughout this paper, graphs are finite, simple, undirected and connected. For a graph *X*, let *V*(*X*), *E*(*X*) and Aut(*X*) denote the vertex set, the edge set and the full automorphism group of *X*, respectively. The *arc set A*(*X*) of a graph *X* is defined to be the set { $(u, v), (v, u) | \{u, v\} \in E(X)$ }. For a vertex $v \in V(X)$, by N(v) we denote the set of vertices adjacent to *v*. A graph \widetilde{X} is called a *covering* of *X* with a projection $p : \widetilde{X} \to X$ if *p* is a surjection from $V(\widetilde{X})$ to V(x) such that $\mathbf{p}|_{N(\widetilde{v})} : N(\widetilde{v}) \to N(v)$ is a bijection for any vertex $v \in V(X)$ and $\widetilde{v} \in \mathbf{p}^{-1}(v)$. The graph *X* is usually referred to as the *base graph* and \widetilde{X} as the *covering graph*. The *fibre* of an arc or a vertex is its preimage under *p*. The group CT(\mathbf{p}) of all automorphisms of \widetilde{X} which fix each of the fibres setwise is called the *covering transformation group*.

An *s*-arc in a graph X is an ordered (s + 1)-tuple (v_0, v_1, \ldots, v_s) of vertices of X such that v_{i-1} is adjacent to v_i for $1 \le i \le s$, and $v_{i-1} \ne v_{i+1}$ for $1 \le i < s$. A graph X is said to be *s*-arc-transitive if Aut(X) is transitive on the set of *s*-arcs in X. In particular, 0-arc-transitive means *vertex-transitive*, and 1-arc-transitive means *arc-transitive* or *symmetric*. A subgroup of the automorphism group of a graph X is said to be *s*-regular if it acts regularly on the set of *s*-arcs of X. In particular, if the subgroup is the full automorphism group Aut(X) of X then X is said to be *s*-regular. Thus, if a graph X is *s*-regular then Aut(X) is transitive on the set of *s*-arcs and the only automorphism fixing an *s*-arc is the identity automorphism of X.

A covering $\mathbf{p} : \widetilde{X} \to X$ is said to be *regular* (or *N*-covering) if there is a semiregular subgroup *N* of the automorphism group Aut(\widetilde{X}) such that the graph *X* is isomorphic to the quotient graph \widetilde{X}/N , say by an isomorphism *h*, and the quotient map $\widetilde{X} \to \widetilde{X}/N$ is the composition $h\mathbf{p}$ of *h* and \mathbf{p} . If the covering graph \widetilde{X} is connected, then *N* is the covering transformation group. An automorphism of a covering graph \widetilde{X} is said to be *fibre-preserving* if it maps a fibre to a fibre, while a covering transformation maps a fibre onto itself. An automorphism $\alpha \in Aut(X)$ lifts along \mathbf{p} if there exists an automorphism $\widetilde{\alpha} \in Aut(\widetilde{X})$ such that $\alpha \mathbf{p} = \mathbf{p}\widetilde{\alpha}$. In this case we also say that \mathbf{p} is α -admissible. A subgroup $G \leq Aut(X)$ lifts along \mathbf{p} if each $\alpha \in G$ lifts. The set of all lifts *G* forms a group $\widetilde{G} \leq Aut(\widetilde{X})$, called the *lift* of *G*. A regular covering projection \mathbf{p} is *arc-transitive* if some arc-transitive subgroup of Aut(X) lifts along \mathbf{p} .

Two coverings \widehat{X} and $\widehat{X'}$ with projections \mathbf{p} and $\mathbf{p'}$, respectively, are said to be *isomorphic* if there exist an automorphism $\alpha \in \operatorname{Aut}(X)$ and an isomorphism $\widetilde{\alpha} : \widetilde{X} \to \widetilde{X'}$ such that $\alpha \mathbf{p} = \mathbf{p'}\widetilde{\alpha}$. In particular, if α is the identity automorphism of X, then we say that \widetilde{X} and $\widetilde{X'}$ are *equivalent*.

ELSEVIER



ABSTRACT

A graph is *s*-*regular* if its automorphism group acts regularly on the set of its *s*-arcs. In this paper, we classify all cubic *s*-regular graphs of order 16*p* for every $s \ge 1$ and every prime *p*. As a result, a new infinite family of cubic 1-regular graphs with girth 10 is constructed. © 2008 Elsevier B.V. All rights reserved.

E-mail address: jumokoh@postech.ac.kr.

⁰⁰¹²⁻³⁶⁵X/ $\$ - see front matter $\$ 2008 Elsevier B.V. All rights reserved. doi:10.1016/j.disc.2008.09.001

Table 1
Cubic symmetric graphs of order 16p with $p \leq 47$

Graph	Order	s-regular	Girth	Diameter	Bipartite?
F ₃₂	16 · 2	2	6	5	Yes
F ₄₈	16 · 3	2	8	6	Yes
F ₈₀	16 · 5	3	10	8	Yes
F _{112A}	16 · 7	1	10	7	Yes
F _{112B}	16 · 7	2	8	7	Yes
F _{112C}	16 · 7	3	8	10	Yes
F ₂₀₈	16 · 13	1	10	9	Yes
F ₃₀₄	16 · 19	1	10	11	Yes
F496	16 · 31	1	10	15	Yes
F ₅₉₂	16 · 37	1	10	15	Yes
F ₆₈₈	16 · 43	1	10	17	Yes

Let *X* be a connected graph and *N* be a finite group, called the *voltage group*. Assign to each arc of *X* a *voltage* $\xi(u, v) \in N$ such that $\xi(v, u) = \xi(u, v)^{-1}$. This function ξ is called an (*ordinary*) *voltage assignment* of *X*. Let $Cov(X, \xi)$ be the *derived graph* with vertex set $V \times N$ and adjacency relation defined by $(u, a) \sim (v, a\xi(u, v))$ whenever $u \sim v$ in *X*. Then the first coordinate projection is a regular covering $\mathbf{p}_{\xi} : Cov(X, \xi) \to X$ where the group *N*, viewed as $CT(\mathbf{p}_{\xi})$, acts via left multiplication on itself. Given a spanning tree *T* of the graph *X*, a voltage assignment ξ is called *T-reduced* if the voltages on the tree arcs are the identity. Gross and Tucker [14] showed that every regular covering projection $\mathbf{p} : \widehat{X} \to X$ is equivalent to $\mathbf{p}_{\xi} : Cov(X, \xi) \to X$ for some *T*-reduced voltage assignment $\xi : X \to N$ with respect to an arbitrary fixed spanning tree *T* of *X*.

Tutte [25,26] showed that every finite cubic symmetric graph is *s*-regular for some $s \ge 1$, and this *s* is at most five. It follows that every cubic symmetric graph has an order of the form 2mp for a positive integer *m* and a prime number *p*. In order to know all cubic symmetric graphs, we need to classify the cubic *s*-regular graphs of order 2mp for a fixed positive integer *m* and each prime *p*. Conder and Dobcsányi [3,4] classified the cubic *s*-regular graphs up to order 2048 with the help of the "Low index normal subgroups" routine in MAGMA system [1]. Cheng and Oxley [2] classified the cubic *s*-regular graphs of order $2p^2$, $2p^3$, 4p, $4p^2$, 6p, $6p^2$, 8p, $8p^2$, 10p, $10p^2$ and 14p were classified in [7–12,21].

In this paper, we classify all cubic s-regular cubic graphs with order 16*p* for each $s \ge 1$ and each prime *p*. As a result, a new infinite family of cubic 1-regular graphs with girth 10 is constructed.

2. The cubic symmetric graphs of order 16p

We will use the following well-known results in group theory.

Proposition 2.1 ([15, Chapter IV, Theorem 2.6]). Let G be a finite group and P a Sylow p-subgroup of G. Let $N_G(P)$ be the normalizer of P in G and $C_G(P)$ the centralizer of P in G. If $N_G(P) = C_G(P)$, then G has a normal subgroup N such that $G/N \cong P$.

Proposition 2.2. (1) [22, Theorem 8.5.3] Let p and q be primes and let a and b be non-negative integers. Then every group of order $p^a q^b$ is solvable.

(2) [13, Feit-Thompson Theorem] Every finite group of odd order is solvable.

Let *X* be a graph and let *N* be a subgroup of Aut(*X*). Denote by \underline{X} the quotient graph corresponding to the orbits of *N*, that is the graph having the orbits of *N* as vertices with two orbits adjacent in \underline{X} whenever there is an edge between those orbits in *X*.

Proposition 2.3 ([16, Theorem 9]). Let X be a connected symmetric graph of a prime valency and let G be an s-arc-transitive subgroup of Aut(X) for some $s \ge 1$. If a normal subgroup N of G has more than two orbits, then it is semiregular and G/N is an s-arc-transitive subgroup of Aut(X) where X is the quotient graph of X corresponding to the orbits of N.

By [3,4] we have the following.

Lemma 2.4. Let p be a prime. Let X be a cubic symmetric graph of order 16p. If $p \le 47$, then X is isomorphic to one of the graphs in Table 1.

Assume that a connected graph X and a subgroup $G \leq \operatorname{Aut}(X)$ are given. Choose a spanning tree T of X and a set of arcs $\{x_1, \ldots, x_r\} \subseteq A(X)$ containing exactly one arc from each edge in $E(X \setminus T)$. Let \mathcal{B}_T be the corresponding basis of the first homology group $H_1(X, \mathbb{Z}_p)$ determined by $\{x_1, \ldots, x_r\}$. Further, denote by $G^{\#_h} = \{\alpha^{\#_h} \mid \alpha \in G\} \leq \operatorname{GL}(H_1(X, \mathbb{Z}_p))$ the induced action of G on $H_1(X, \mathbb{Z}_p)$, and let $M_G \leq \mathbb{Z}_p^{r \times r}$ be the matrix representation of $G^{\#_h}$ with respect to the basis \mathcal{B}_T . By M_G^t we denote the dual group consisting of all transposes of matrices in M_G .

The following proposition is a special case of [18, Proposition 6.3, Corollary 6.5] (also see [6,23]).



Fig. 1. The Möbius-Kantor graph.

Proposition 2.5. Let *T* be a spanning tree of a connected graph *X* and let the set $\{x_1, x_2, ..., x_r\} \subseteq A(X)$ contain exactly one arc from each cotree edge. Let $\xi : A(X) \rightarrow \mathbb{Z}_p$ be a voltage assignment on *X* which is trivial on *T*, and let $Z(\xi) = [\xi(x_1), \xi(x_2), ..., \xi(x_r)]^t \in \mathbb{Z}_p^{r \times 1}$. Then the following holds.

- (a) A group $G \leq \text{Aut}(X)$ lifts along $p_{\xi} : \text{Cov}(X, \xi) \to X$ if and only if the induced subspace $\langle Z(\xi) \rangle$ is an M_G^t -invariant 1-dimensional subspace.
- (b) If $\xi' : A(X) \to \mathbb{Z}_p$ is another voltage assignment satisfying (a), then $Cov(X, \xi')$ is equivalent to $Cov(X, \xi)$ if and only if $\langle Z(\xi) \rangle = \langle Z(\xi') \rangle$, as subspaces. Moreover, $Cov(X, \xi')$ is isomorphic to $Cov(X, \xi)$ if and only if there exists an automorphism $\alpha \in Aut(X)$ such that the matrix M_{α}^t maps $\langle Z(\xi') \rangle$ onto $\langle Z(\xi) \rangle$.

The Möbius–Kantor graph F_{16} is illustrated in Fig. 1. It is known that F_{16} is a unique cubic symmetric graph of order 16, which is 2-regular (see [3,4]). We choose

 $\begin{aligned} \alpha &:= (2, 8, 9)(3, 16, 14)(4, 13, 6)(7, 12, 10), \\ \beta &:= (1, 2)(3, 8)(4, 7)(5, 6)(9, 10)(11, 16)(12, 15)(13, 14), \\ \gamma &:= (1, 2)(3, 9)(4, 14)(5, 6)(7, 13)(8, 10)(11, 12)(15, 16) \end{aligned}$

as automorphisms of F_{16} . Then Aut $(F_{16}) = \langle \alpha, \beta, \gamma \rangle$ and Aut (F_{16}) has two proper arc-transitive subgroups $H := \langle \alpha, \beta \rangle$ and $K := \langle \alpha, \gamma \rangle$. This can be checked by GAP [24].

Thus, in order to determine all arc-transitive \mathbb{Z}_p -covering projections of F_{16} , it suffices to find those which are H- or K-admissible. By Proposition 2.5, this is equivalent to finding all invariant 1-dimensional subspaces of the representations M_H^t or M_K^t .

We choose a spanning tree T of F_{16} consisting of the edges

- $\{\{1, 9\}, \{2, 10\}, \{3, 11\}, \{4, 12\}, \{5, 13\}, \{6, 14\}, \{7, 15\}, \{8, 16\}, \{1, 9\}, \{2, 10\}, \{3, 11\}, \{4, 12\}, \{5, 13\}, \{6, 14\}, \{7, 15\}, \{8, 16\}, \{1, 12\}, \{1, 12\}, \{2, 12\}, \{3, 12\}, \{4, 12\}, \{5, 13\}, \{4, 12\}, \{5, 13\}, \{6, 14\}, \{7, 15\}, \{8, 16\}, \{1, 12\}, \{1, 12\}, \{2, 12\}, \{3, 12\}, \{4, 12\}, \{5, 13\}, \{6, 14\}, \{7, 15\}, \{8, 16\}, \{1, 12\}, \{1, 12\}, \{1, 12\}, \{2, 12\}, \{2, 12\}, \{3, 12\}, \{4$
- $\{9, 12\}, \{9, 14\}, \{10, 13\}, \{10, 15\}, \{11, 14\}, \{12, 15\}, \{13, 16\}\}.$

We orient the cotree arcs by setting

 $x_1 = (1, 2),$ $x_2 = (2, 3),$ $x_3 = (3, 4),$ $x_4 = (4, 5),$ $x_5 = (5, 6),$ $x_6 = (6, 7),$ $x_7 = (7, 8),$ $x_8 = (8, 1),$ $x_9 = (11, 16).$

Let $\mathcal{B} = \{C_{x_i} \mid 1 \le i \le 9\}$ be the standard ordered basis of $H_1(F_{16}, \mathbb{Z}_p)$ associated with the spanning tree *T* and the arcs x_i (i = 1, ..., 9). Let $p \ge 5$ be a prime number such that $p = 1 \mod(6)$ and let ζ be a solution of the equation $x^2 + x + 1 = 0$ in \mathbb{Z}_p . We define a *T*-reduced voltage assignment $\xi : \{x_i \mid 1 \le i \le 9\} \rightarrow \mathbb{Z}_p$ by setting

$$x_1 \mapsto \zeta, \quad x_2 \mapsto 1-\zeta, \quad x_3 \mapsto \zeta, \quad x_4 \mapsto -\zeta - 1, \quad x_5 \mapsto \zeta + 2,$$

 $x_6 \mapsto -\zeta - 1, \quad x_7 \mapsto \zeta, \quad x_8 \mapsto 1-\zeta, \quad x_9 \mapsto -2.$

We remark that the voltage assignment ξ is derived from the M_{H}^{t} -invariant 1-dimensional subspace $\langle k_{1} \rangle$ (see Section 3). Let $CF_{16p}(p \geq 5)$ be the derived graph from the voltage assignment ξ .

Malnič et al. [17] classified *semisymmetric elementary abelian* covers of F_{16} . One might derive the following theorem from [17]. But, we give its (simpler) proof in the next section.

Theorem 2.6. Let $p \ge 5$ be a prime. Let \widetilde{X} be an arc-transitive \mathbb{Z}_p -cover of the Möbius–Kantor graph F_{16} . Then \widetilde{X} is isomorphic to the 1-regular graph CF_{16p} of girth 10 where $p = 1 \mod(6)$.

Remark. Marušič et al. [19,20] gave the relation between *half-transitive* group action with vertex stabilizer \mathbb{Z}_2 and 1-regular group action with cyclic vertex stabilizer, which give us infinitely many finite half-transitive graphs of valency 4.

The following is the main result in this paper.

Theorem 2.7. Let p be a prime and let X be a connected cubic symmetric graph of order 16p. Then X is 1-, 2- or 3-regular. Furthermore,

(1) *X* is 1-regular if and only if *X* is isomorphic to the graph $CF_{16p}(p \ge 7)$, where $p = 1 \mod(6)$.

(2) X is 2-regular if and only if X is isomorphic to one of the three graphs F_{32} , F_{48} and F_{112B} .

(3) X is 3-regular if and only if X is isomorphic to one of the two graphs F_{80} and F_{112C} .

Proof. Let *X* be a cubic symmetric graph of order 16*p*. By [3,4] we may assume p > 47. Let A = Aut(X) and let *P* be a Sylow *p*-subgroup of *A*. If *P* is normal in *A*, by Proposition 2.3 *X* is a regular covering of the graph F_{16} with the covering transformation group \mathbb{Z}_p and the normality of *P* implies that the fibre-preserving group is arc-transitive. By Theorem 2.6, *X* is isomorphic to CF_{16p} . Thus, it suffices to show that *P* is normal in *A*.

Let $N_A(P)$ be the normalizer of P in A. By Sylow's theorem, the number of Sylow p-subgroups of A is $np + 1 = |A : N_A(P)|$. Since X is at most 5-regular, |A| is a divisor of $48 \cdot 16p$. Thus np + 1 is a divisor of $48 \cdot 16$. Suppose to the contrary that P is not normal in A. Since $np + 1 \ge 54$ and $np + 1 | 2^8 \cdot 3$, we have (n, p) = (13, 59), (1, 127), (1, 191) or (1, 383). If $N_A(P) = P$ then $C_A(P) = P$, where $C_A(P)$ is the centralizer of P in A. By Proposition 2.1, A has a normal subgroup N such that $A/N \cong P$, and by Proposition 2.3, the quotient graph corresponding to the orbits of N has odd order and valency 3, a contradiction. Thus one may assume $(n, p) \neq (13, 59)$. Since $|A : N_A(P)| = 2^7, 2^6 \cdot 3$ or $2^7 \cdot 3$, |A| has a divisor $2^7 \cdot 3 \cdot p$ where p = 127, 191or 383, implying that X is at least 3-arc-transitive. Let M be a minimal normal subgroup of A and X the quotient graph of Xcorresponding to the orbits of M.

If M is elementary abelian then by Proposition 2.3 \underline{X} is 3-arc-transitive with order 2⁴, 2p, 4p or 8p, which is impossible by the result in [3,4], [8, Theorem 5.2] and [11, Theorem 5.1]. Thus, one may assume that $M = T_1 \times T_2 \times \cdots \times T_t$, where T_i ($1 \le i \le t$) are isomorphic non-abelian simple groups. By Proposition 2.2, $|T_i|$ has at least three prime factors. Notice that |A| is a divisor of $2^8 \cdot 3 \cdot p$ where p = 127, 191 or 383. Then t = 1 and M is a non-abelian simple group. Thus M has order $2^\ell \cdot 3 \cdot p$ for some $1 \le \ell \le 8$. However, there is no simple group with such orders (see [5]).

3. The proof of Theorem 2.6

Let $p \ge 5$ be a prime. It is known that a polynomial $x^2 + x + 1 = 0$ has a solution in \mathbb{Z}_p if and only if -3 is a square root in \mathbb{Z}_p , which is if and only if $p = 1 \pmod{6}$.

Let *R*, *T* and *S* be the *transposes* of the matrices which represent the linear transformations $\alpha^{\#_h}$, $\beta^{\#_h}$ and $\gamma^{\#_h}$ relative to \mathcal{B} , respectively. Then

In order to find (R, T)- or (R, S)-invariant 1-dimensional subspaces in \mathbb{Z}_p , it is useful to consider R, T and S as matrices over the splitting field $\mathbb{Z}_p(\zeta)$ where ζ is a solution of the polynomial $x^2 + x + 1 = 0$. The respective characteristic and minimal polynomials of R, T and S are

$$\begin{aligned} \Delta_R(x) &= (x-1)(x-\zeta)^4(x-\zeta^2)^4, \qquad m_R(x) = (x-1)(x-\zeta)(x-\zeta^2), \\ \Delta_T(x) &= (x-1)^3(x+1)^6, \qquad m_T(x) = (x-1)(x+1), \\ \Delta_S(x) &= (x-1)^4(x+1)^5, \qquad m_S(x) = (x-1)(x+1). \end{aligned}$$

By a straightforward calculation, we have

$$\begin{split} & \operatorname{Ker}(R-I) = \langle u_1 \rangle \,, \qquad \operatorname{Ker}(R-\zeta I) = \langle u_2, u_3, u_4, u_5 \rangle \,, \qquad \operatorname{Ker}(R-\zeta^2 I) = \langle u_6, u_7, u_8, u_9 \rangle \,, \\ & \operatorname{Ker}(T-I) = \langle v_1, v_2, v_3 \rangle \,, \qquad \operatorname{Ker}(T+I) = \langle v_4, v_5, v_6, v_7, v_8, v_9 \rangle \,, \\ & \operatorname{Ker}(S-I) = \langle w_1, w_2, w_3, w_4 \rangle \,, \qquad \operatorname{Ker}(S+I) = \langle w_5, w_6, w_7, w_8, w_9 \rangle \end{split}$$

where

Solving homogeneous linear equations over the splitting field $\mathbb{Z}_p(\zeta)$, one can see that

 $\operatorname{Ker}(R-I) \cap \operatorname{Ker}(T \pm I) = \operatorname{Ker}(R-I) \cap \operatorname{Ker}(S \pm I) = 0,$ $\operatorname{Ker}(R - \zeta I) \cap \operatorname{Ker}(S \pm I) = \operatorname{Ker}(R - \zeta^2 I) \cap \operatorname{Ker}(S \pm I) = 0,$ $\operatorname{Ker}(R-\zeta I)\cap\operatorname{Ker}(T-I)=\operatorname{Ker}(R-\zeta^2 I)\cap\operatorname{Ker}(T-I)=0,$ $\operatorname{Ker}(R-\zeta I)\cap\operatorname{Ker}(T+I)=\langle k_1\rangle,$ $\operatorname{Ker}(R - \zeta^2 I) \cap \operatorname{Ker}(T + I) = \langle k_2 \rangle$

where

$$k_{1} := \begin{bmatrix} \zeta \\ 1 - \zeta \\ \zeta \\ -\zeta - 1 \\ \zeta + 2 \\ -\zeta - 1 \\ \zeta \\ 1 - \zeta \\ -2 \end{bmatrix} \text{ and } k_{2} := \begin{bmatrix} 1 \\ \zeta - 1 \\ 1 \\ -\zeta - 1 \\ 2\zeta + 1 \\ -\zeta - 1 \\ 1 \\ \zeta - 1 \\ -2\zeta \end{bmatrix}.$$

Hence, there exist only two $\langle R, T \rangle$ -invariant 1-dimensional subspaces $\langle k_1 \rangle$ and $\langle k_2 \rangle$. Furthermore, since $Sk_1 = \zeta k_2$, two spaces $\langle k_1 \rangle$ and $\langle k_2 \rangle$ induce isomorphic covering projections whose maximal lifting group is H. By considering the induced subgraph

 $\langle N_0(1,0) \cup N_1(1,0) \cup N_2(1,0) \cup N_3(1,0) \cup N_4(1,0) \cup N_5(1,0) \rangle$

of CF_{16p} , one can see that the girth of CF_{16p} is 10. This completes the proof that any arc-transitive \mathbb{Z}_p -covering ($p \ge 5$) graph of F_{16} is isomorphic to the graph CF_{16p} with girth 10.

By Lemma 2.4, the graph CF₁₆₋₇ is 1-regular and isomorphic to F_{112A} because the girth of CF₁₆₋₇ is 10. Thus, one can assume p > 11. Let **p** : $CF_{16n} \rightarrow F_{16}$ be the associated covering projection from the voltage assignment ξ and $A := Aut(CF_{16n})$. Suppose to the contrary that CF_{16p} is s-regular for some $s \ge 2$. By Tutte [25,26], $s \le 5$ and so $|A| \mid 16 \cdot p \cdot 48$. Thus $L := CT(\mathbf{p})$ is a Sylow *p*-subgroup of *A*. Let *B* be the 1-regular subgroup of Aut(CF_{16p}) lifted by $\langle \alpha, \beta \rangle$. Then $|B| = 16 \cdot 3 \cdot p$. The normality of L in B implies that $B < N_4(L)$, where $N_4(L)$ is the normalizer of L in A. Since \widetilde{X} is at most 5-regular, $|A| : N_4(L) | = 16$. By Sylow's theorem, the number of Sylow *p*-subgroups of *A* is np + 1 and $np + 1 = |A| : N_A(L)|$. Since $p \ge 11$, we have np + 1 = 1. Thus L is normal in A. By Proposition 2.3, A/L is an s-regular subgroup of Aut(F_{16}). This is impossible because otherwise s-regular subgroup A/L(s > 2) of Aut(F_{16}) lifts. This completes the proof of Theorem 2.6.

As continuation of this work, we have classified the cubic s-regular graphs of order 18p and 20p for every $s \ge 1$ and every prime p.

Acknowledgement

This paper was supported by Korea Research Foundation Grant (KRF-2007-313-C00011).

References

- W. Bosma, J. Cannon, Handbook of Magma Function, Sydney University Press, Sydney, 1994.

- 4
- Y. Cheng, J. Oxley, On weakly symmetric graphs of order twice a prime, J. Combin. Theory Ser. B 42 (1987) 196–211. M.D.E. Conder, Trivalent (cubic) symmetric graphs on up to 2048 vertices, 2006. http://www.math.auckland.ac.nz~conder/symmcubic2048list.txt. M.D.E. Conder, P. Dobcsányi, Trivalent symmetric graphs on up to 768 vertices, J. Combin. Math. Combin. Comput. 40 (2002) 41–63. J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, R.A. Wilson, An ATLAS of Finite Groups, Oxford University Press, Oxford, 1985. S.F. Du, J.H. Kwak, M.Y. Xu, Linear criteria for lifting automorphisms of the elementary abelian regular coverings, Linear Algebra Appl. 373 (2003) 101–119.
 Y.Q. Feng, J.H. Kwak, Classifying cubic symmetric graphs of order 10p or 10p², Sci. China Ser. A 49 (2006) 300–319.
 Y.Q. Feng, J.H. Kwak, Cubic symmetric graphs of order a small number times a prime or a prime square, J. Combin. Theory Ser. B 97 (2007) 627–646.
 Y.Q. Feng, J.H. Kwak, Cubic symmetric graphs of order twice an odd prime-power, J. Aust. Math. Soc. 81 (2006) 153–164.
 Y.Q. Feng, J.H. Kwak, One-regular cubic graphs of order a small number times a prime or a prime square, J. Aust. Math. Soc. 76 (2004) 345–356.
 Y.Q. Feng, J.H. Kwak, M.Y. Xu, Cubic s-regular graphs of order 2p³, J. Graph Theory 52 (2006) 341–352.
 W. Feit, J.G. Thompson, Solvability of groups of odd order, Pacific J. Math. 13 (1936) 775–1029.
 J.L. Gross, T.W. Tucker, Generating all graph coverings by permutation voltage assignments, Discrete Math. 18 (1977) 273–283.
 B. Huppert, Endliche Gruppen I, Springer, Berlin, 1979.
 P. Lorimer, Vertex-transitive graphs of prime valency 1 Graph Theory 8 (1984) 55–68. 101 - 119

- 12
- 13
- 14
- 16] P. Lorimer, Vertex-transitive graphs: Symmetric graphs of prime valency, J. Graph Theory 8 (1984) 55-68.
- A. Malnič, D. Marušič, Š. Miklavič, P. Potočnik, Semisymmetric elementary abelian covers of the Möbius-Kantor graph, Discrete Math. 307 (2007) 2156-2175
- A. Malnič, D. Marušič, P. Potočnik, Elementary abelian covers of graphs, J. Algebraic Combin. 20 (2004) 71–97. D. Marušič, R. Nedela, Maps and half-transitive graphs of valency 4, European J. Combin. 19 (1998) 345–354.
- [19] D. Marušić, K. Nedela, Maps and nan-transitive graphs of valency 4, European J. Combin. 19 (1998) 545–554.
 [20] D. Marušić, M.Y. Xu, A ¹/₂-transitive graphs of valency 4 with a nonsolvable group of automorphisms, J. Graph Theory 25 (1994) 133–138.
 [21] J.M. Oh, A classification of cubic s-regular graphs of order 14p, Discrete Math., in press (doi:10.1016/j.disc.2008.06.025).
 [22] D.J. Robinson, A Course in the Theory of Groups, Springer-Verlag, Berlin, 1979.
 [23] J. Siráň, Coverings of graphs and maps, orthogonality, and eigenvectors, J. Algebraic Combin. 14 (2001) 57–72.
 [24] The GAP group, GAP Groups, algorithms, and programming, Version 4.4, 2007. http://www.gap-system.org.
 [25] W.T. Tutte, A family of cubical graphs, Proc. Cambridge Philos. Soc. 43 (1947) 459–474.
 [26] W.T. Tutte, On the symmetry of cubic graphs, Canad. J. Math. 11 (1959) 621–624.