

A class of II_1 factors with many non conjugate Cartan subalgebras

An Speelman¹, Stefaan Vaes*

KU Leuven, Department of Mathematics, Belgium

Received 18 November 2011; accepted 2 July 2012

Available online 14 August 2012

Communicated by Alain Connes

Abstract

We construct a class of II_1 factors M that admit unclassifiably many Cartan subalgebras in the sense that the equivalence relation of being conjugate by an automorphism of M is complete analytic, in particular non Borel. We also construct a II_1 factor that admits uncountably many non isomorphic group measure space decompositions, all involving the same group G . So G is a group that admits uncountably many non stably orbit equivalent actions whose crossed product II_1 factors are all isomorphic.

© 2012 Elsevier Inc. All rights reserved.

Keywords: II_1 factor; Cartan subalgebra; Deformation/rigidity theory

1. Introduction

Free ergodic probability measure preserving (p.m.p.) actions $\Gamma \curvearrowright (X, \mu)$ of countable groups give rise to II_1 factors $L^\infty(X) \rtimes \Gamma$ by Murray and von Neumann's group measure space construction. In [34] it was shown that the isomorphism class of $L^\infty(X) \rtimes \Gamma$ only depends on the orbit equivalence relation on (X, μ) given by $x \sim y$ iff $x \in \Gamma \cdot y$. More precisely, two free ergodic p.m.p. actions $\Gamma \curvearrowright X$ and $\Lambda \curvearrowright Y$ are orbit equivalent iff there exists an isomorphism of $L^\infty(X) \rtimes \Gamma$ onto $L^\infty(Y) \rtimes \Lambda$ sending $L^\infty(X)$ onto $L^\infty(Y)$.

* Corresponding author.

E-mail addresses: an.speelman@wis.kuleuven.be (A. Speelman), stefaan.vaes@wis.kuleuven.be (S. Vaes).

¹ Research Assistant of the Research Foundation – Flanders (FWO).

If $\Gamma \curvearrowright (X, \mu)$ is essentially free and $M := L^\infty(X) \rtimes \Gamma$, then $A := L^\infty(X)$ is a maximal abelian subalgebra of M that is regular: the group of unitaries $u \in \mathcal{U}(M)$ that normalize A in the sense that $uAu^* = A$, spans a dense $*$ -subalgebra of M . A regular maximal abelian subalgebra of a II_1 factor is called a *Cartan subalgebra*.

In order to classify classes of group measure space II_1 factors $M := L^\infty(X) \rtimes \Gamma$ in terms of the group action $\Gamma \curvearrowright (X, \mu)$, one must solve two different problems. First one should understand whether the Cartan subalgebra $L^\infty(X) \subset M$ is, in some sense, unique. If this is the case, any other group measure space decomposition $M = L^\infty(Y) \rtimes \Lambda$ must come from an orbit equivalence between $\Gamma \curvearrowright X$ and $\Lambda \curvearrowright Y$. The second problem is then to classify the corresponding class of orbit equivalence relations $\mathcal{R}(\Gamma \curvearrowright X)$ in terms of the group action. Both problems are notoriously hard, but a huge progress has been made over the last 10 years thanks to Sorin Popa's deformation/rigidity theory (see [27,11,35] for surveys). In particular, many uniqueness results for Cartan subalgebras and group measure space Cartan subalgebras have been established recently in [18,19,21,32,10,1,13,36,15,2].

In this paper we concentrate on “negative” solutions to the first problem by exhibiting a class of II_1 factors with many Cartan subalgebras. First recall that two Cartan subalgebras $A \subset M$ and $B \subset M$ of a II_1 factor are called *unitarily conjugate* if there exists a unitary $u \in \mathcal{U}(M)$ such that $B = uAu^*$. The Cartan subalgebras are called *conjugate by an automorphism* if there exists $\alpha \in \text{Aut } M$ such that $B = \alpha(A)$. In [3] it is shown that the hyperfinite II_1 factor R has a unique Cartan subalgebra up to conjugacy by an automorphism of R . By [9, Theorem 7] (see [20, Section 4] for details) the hyperfinite II_1 factor however admits uncountably many Cartan subalgebras that are not unitarily conjugate. In [4] Connes and Jones constructed the first example of a II_1 factor that admits two Cartan subalgebras that are not conjugate by an automorphism. More explicit examples of this phenomenon were found in [19, Section 7] and [32, Example 5.8].

In [29, Section 6.1] Popa discovered a free ergodic p.m.p. action $\Gamma \curvearrowright (X, \mu)$ such that the orbit equivalence relation $\mathcal{R}(\Gamma \curvearrowright X)$ has trivial fundamental group, while the II_1 factor $M := L^\infty(X) \rtimes \Gamma$ has fundamental group \mathbb{R}_+ . Associated with this is a one parameter family A_t of Cartan subalgebras of M that are not conjugate by an automorphism of M . Indeed, put $A := L^\infty(X)$ and for every $t \in (0, 1]$, choose a projection $p_t \in A$ of trace t and a $*$ -isomorphism $\pi_t : p_t M p_t \rightarrow M$. Put $A_t := \pi_t(A p_t)$. Then $A_t \subset M$, $t \in (0, 1]$, are Cartan subalgebras that are not conjugate by an automorphism. Note that earlier (see [23, Corollary of Theorem 3] and [22, Corollary 4.7.2]) Popa found free ergodic p.m.p. actions $\Gamma \curvearrowright (X, \mu)$ such that the equivalence relation $\mathcal{R}(\Gamma \curvearrowright X)$ has countable fundamental group, while the II_1 factor $L^\infty(X) \rtimes \Gamma$ has fundamental group \mathbb{R}_+ . By the same construction, there are uncountably many t such that the Cartan subalgebras A_t are not conjugate by an automorphism. In all these cases, although the Cartan subalgebras $A_t \subset M$ are not conjugate by an automorphism of M , they are by construction conjugate by a stable automorphism of M . More precisely, we say that two Cartan subalgebras $A \subset M$ and $B \subset M$ are *conjugate by a stable automorphism of M* if there exist non zero projections $p \in A$, $q \in B$ and a $*$ -isomorphism $\alpha : pMp \rightarrow qMq$ satisfying $\alpha(Ap) = Bq$.

In this paper we provide the first class of II_1 factors M that admit uncountably many Cartan subalgebras that are not conjugate by a stable automorphism of M . Actually we obtain II_1 factors M such that the equivalence relation “being conjugate by an automorphism” on the set of Cartan subalgebras of M is complete analytic, in particular non Borel. In this precise sense the decision problem whether two Cartan subalgebras of these II_1 factors M are conjugate by an automorphism, is as hard as it can possibly get (see Remark 14). Using Popa's conjugacy criterion for Cartan subalgebras [24, Theorem A.1], we show that for any II_1 factor with separable predual,

the set of Cartan subalgebras is a standard Borel space and the equivalence relation “being unitarily conjugate” is always Borel. We provide examples of II_1 factors where this equivalence relation is not concretely classifiable.

The “many” Cartan subalgebras in the II_1 factors M in the previous paragraph are not of group measure space type. In the final section of the paper we give the first example of a II_1 factor that admits uncountably many Cartan subalgebras that are not conjugate by a (stable) automorphism and that all arise from a group measure space construction. They actually all arise as the crossed product with a single group G . This means that we construct a group G that admits uncountably many free ergodic p.m.p. actions $G \curvearrowright (X_i, \mu_i)$ that are non stably orbit equivalent, such that the II_1 factors $L^\infty(X_i) \rtimes G$ are all isomorphic. This group G should be opposed to the classes of groups considered in [32,10,1,13,36] for which isomorphism of the group measure space II_1 factors always implies orbit equivalence of the group actions.

Terminology

A Cartan subalgebra A of a II_1 factor M is a maximal abelian von Neumann subalgebra whose normalizer

$$\mathcal{N}_M(A) := \{u \in \mathcal{U}(M) \mid uAu^* = A\}$$

generates M as a von Neumann algebra. As mentioned above, we say that two Cartan subalgebras $A \subset M$ and $B \subset M$ are *unitarily conjugate* if there exists a unitary $u \in \mathcal{U}(M)$ such that $uAu^* = B$. We say that they are *conjugate by an automorphism of M* if there exists $\alpha \in \text{Aut} M$ such that $\alpha(A) = B$. Finally A and B are said to be *conjugate by a stable automorphism of M* if there exist non zero projections $p \in A, q \in B$ and a $*$ -isomorphism $\alpha : pMp \rightarrow qMq$ satisfying $\alpha(p) = Bq$.

By [9] every II_1 equivalence relation \mathcal{R} on (X, μ) gives rise to a II_1 factor $L\mathcal{R}$ that contains $L^\infty(X)$ as a Cartan subalgebra. Conversely, if $L^\infty(X) = A \subset M$ is an arbitrary Cartan subalgebra of a II_1 factor M , we get a natural II_1 equivalence relation \mathcal{R} on (X, μ) and a measurable 2-cocycle on \mathcal{R} with values in \mathbb{T} such that $A \subset M$ is canonically isomorphic with $L^\infty(X) \subset L_\Omega \mathcal{R}$.

If \mathcal{R}_i are II_1 equivalence relations on the standard probability spaces (X_i, μ_i) , we say that \mathcal{R}_1 and \mathcal{R}_2 are *orbit equivalent* if there exists a measure space isomorphism $\Delta : X_1 \rightarrow X_2$ such that $(x, y) \in \mathcal{R}_1$ iff $(\Delta(x), \Delta(y)) \in \mathcal{R}_2$ almost everywhere. More generally, \mathcal{R}_1 and \mathcal{R}_2 are called *stably orbit equivalent* if there exist non negligible measurable subsets $\mathcal{U}_i \subset X_i$ such that the restricted equivalence relations $\mathcal{R}_i|_{\mathcal{U}_i}$ are orbit equivalent.

Throughout this paper compact groups are supposed to be second countable. Two subgroups $K_1, K_2 < K$ of the same group K are called *commensurate* if $K_1 \cap K_2$ has finite index in both K_1 and K_2 .

Statement of the main results

Let Γ be a countable group and K a compact abelian group with countable dense subgroup $\Lambda < K$. Consider the II_1 factor

$$M := L^\infty(K^\Gamma) \rtimes (\Gamma \times \Lambda)$$

where $\Gamma \curvearrowright K^\Gamma$ is the Bernoulli action given by $(g \cdot x)_h = x_{hg}$ and $\Lambda \curvearrowright K^\Gamma$ acts by diagonal translation: $(\lambda \cdot x)_h = \lambda x_h$.

Whenever $K_1 < K$ is a closed subgroup such that $\Lambda_1 := \Lambda \cap K_1$ is dense in K_1 , denote

$$\mathcal{C}(K_1) := L^\infty(K^\Gamma/K_1) \rtimes \Lambda_1 = L^\infty(K^\Gamma/K_1) \overline{\otimes} L\Lambda_1.$$

Note that $\mathcal{C}(K_1)$ is an abelian von Neumann subalgebra of M . In Lemma 6 we prove that $\mathcal{C}(K_1) \subset M$ is always a Cartan subalgebra. As we will see there, the density of $\Lambda \cap K_1$ in K_1 is equivalent with $\mathcal{C}(K_1)$ being maximal abelian in M .

The following is our main result, providing criteria for when the Cartan subalgebras $\mathcal{C}(K_1)$ are unitarily conjugate, conjugate by a (stable) automorphism or inducing (stably) orbit equivalent relations. As we shall see later, concrete choices of $\Lambda \subset K$ then lead to examples of II_1 factors where the equivalence relation of “being conjugate by an automorphism” is complete analytic and where the equivalence relation of “being unitarily conjugate” is not concretely classifiable.

Theorem 1. *Let Γ be an icc property (T) group with $[\Gamma, \Gamma] = \Gamma$. Let K be a compact abelian group with countable dense subgroup Λ . As above, write $M = L^\infty(K^\Gamma) \rtimes (\Gamma \times \Lambda)$. Take closed subgroups $K_i < K$, $i = 1, 2$, such that $\Lambda \cap K_i$ is dense in K_i .*

1. *The following statements are equivalent.*
 - *The Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are unitarily conjugate.*
 - *The subgroups $K_1, K_2 < K$ are commensurate.*
 - *$\Lambda K_1 = \Lambda K_2$.*
2. *The following statements are equivalent.*
 - *The Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are conjugate by an automorphism of M .*
 - *The Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are stably conjugate by an automorphism.*
 - *There exists a continuous automorphism $\delta \in \text{Aut} K$ such that $\delta(\Lambda) = \Lambda$ and $\delta(\Lambda K_1) = \Lambda K_2$.*
3. *The following statements are equivalent.*
 - *The II_1 equivalence relations associated with $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are orbit equivalent.*
 - *The II_1 equivalence relations associated with $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are stably orbit equivalent.*
 - *There exists a continuous automorphism $\delta \in \text{Aut} K$ such that $\delta(\Lambda K_1) = \Lambda K_2$.*

In Section 4 we shall see that for any II_1 factor M with separable predual, the set $\text{Cartan}(M)$ of its Cartan subalgebras is a standard Borel space and that moreover the equivalence relation of “being unitarily conjugate” is Borel, while the equivalence relation of “being conjugate by an automorphism” is analytic (see Proposition 12). As the following examples show, the first can be non concretely classifiable, while the second can be complete analytic, hence non Borel.

Whenever $\Lambda \subset K$ is a compact group with countable dense subgroup, we call $K_1 \subset K$ an *admissible* subgroup if K_1 is closed and $K_1 \cap \Lambda$ is dense in K_1 .

Theorem 2. *Let Γ be any icc property (T) group with $[\Gamma, \Gamma] = \Gamma$, e.g. $\Gamma = \text{SL}(3, \mathbb{Z})$.*

- *Whenever \mathcal{P} is a set of distinct prime numbers, denote $K := \prod_{p \in \mathcal{P}} \mathbb{Z}/p\mathbb{Z}$ and consider the dense subgroup $\Lambda := \bigoplus_{p \in \mathcal{P}} \mathbb{Z}/p\mathbb{Z}$. For every subset $\mathcal{P}_1 \subset \mathcal{P}$, the subgroup of K given by $K(\mathcal{P}_1) := \prod_{p \in \mathcal{P}_1} \mathbb{Z}/p\mathbb{Z}$ is admissible and gives rise to a Cartan subalgebra $\mathcal{C}(\mathcal{P}_1)$ of $M := L^\infty(K^\Gamma) \rtimes (\Gamma \times \Lambda)$. Two such Cartan subalgebras $\mathcal{C}(\mathcal{P}_1)$ and $\mathcal{C}(\mathcal{P}_2)$ or their associated equivalence relations are conjugate in any of the possible senses if and only if the symmetric difference $\mathcal{P}_1 \Delta \mathcal{P}_2$ is finite.*
In particular the equivalence relations on $\text{Cartan}(M)$ given by “being unitarily conjugate” or “being conjugate by a (stable) automorphism” are not concretely classifiable.
- *Denote by $\Lambda := \mathbb{Q}^{(\infty)}$ the countably infinite direct sum of copies of the additive group \mathbb{Q} . Then Λ admits a second countable compactification K such that for the associated II_1 factor $M := L^\infty(K^\Gamma) \rtimes (\Gamma \times \Lambda)$ the equivalence relation of “being conjugate by an automorphism of M ” is a complete analytic equivalence relation on $\text{Cartan}(M)$.*

In [Theorem 1](#) the second set of statements is, at least formally, stronger than the third set of statements. [Example 3](#) shows that this difference really occurs. So in certain cases the Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ give rise to isomorphic equivalence relations, but are nevertheless non conjugate by an automorphism of M . The reason for this is the following: the Cartan inclusions $\mathcal{C}(K_i) \subset M$ come with 2-cocycles Ω_i ; although the associated equivalence relations are isomorphic, this isomorphism does not map Ω_1 onto Ω_2 .

We finally observe that if $K_1 < K$ is an infinite subgroup, the Cartan subalgebra $\mathcal{C}(K_1) \subset M$ is never of group measure space type. The induced equivalence relation is generated by a free action of a countable group (see [Lemma 6](#)), but the 2-cocycle is non trivial (see [Remark 7](#)).

Example 3. Choose elements $\alpha, \beta, \gamma \in \mathbb{T}$ that are rationally independent (i.e. generate a copy of $\mathbb{Z}^3 \subset \mathbb{T}$). Consider $K = \mathbb{T}^3$ and define the countable dense subgroup $\Lambda < K$ generated by $(\alpha, 1, \alpha)$, $(1, \alpha, \beta)$ and $(1, 1, \gamma)$. Put $K_1 := \mathbb{T} \times \{1\} \times \mathbb{T}$ and $K_2 := \{1\} \times \mathbb{T} \times \mathbb{T}$. Then, $\Lambda \cap K_1$ is generated by $(\alpha, 1, \alpha)$, $(1, 1, \gamma)$ and hence dense in K_1 . Similarly $\Lambda \cap K_2$ is dense in K_2 .

We have $\Lambda K_1 = \mathbb{T} \times \alpha^{\mathbb{Z}} \times \mathbb{T}$ and $\Lambda K_2 = \alpha^{\mathbb{Z}} \times \mathbb{T} \times \mathbb{T}$. Hence the automorphism $\delta(x, y, z) = (y, x, z)$ of \mathbb{T}^3 maps ΛK_1 onto ΛK_2 . It is however easy to check that there exists no automorphism δ of K such that $\delta(\Lambda K_1) = \Lambda K_2$ and $\delta(\Lambda) = \Lambda$.

So, by [Theorem 1](#), the Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ of M give rise to orbit equivalent relations, but are not conjugate by a (stable) automorphism of M .

In the final section of the paper we give an example of a group G that admits uncountably many non stably orbit equivalent actions that all give rise to the same II_1 factor. The precise statement goes as follows and will be an immediate consequence of the more concrete, but involved, [Proposition 15](#).

Theorem 4. *There exists a group G that admits uncountably many free ergodic p.m.p. actions $G \curvearrowright (X_i, \mu_i)$ that are non stably orbit equivalent but with all the II_1 factors $L^\infty(X_i) \rtimes G$ being isomorphic with the same II_1 factor M . So this II_1 factor M admits uncountably many group measure space Cartan subalgebras that are non conjugate by a (stable) automorphism of M .*

2. Proof of the main [Theorem 1](#)

We first prove the following elementary lemma that establishes the essential freeness of certain actions on compact groups.

Lemma 5. *Let Y be a compact group and $\Lambda < Y$ a countable subgroup. Assume that a countable group Γ acts on Y by continuous group automorphisms $(\alpha_g)_{g \in \Gamma}$ preserving Λ . Assume that for $g \neq e$ the subgroup $\{y \in Y \mid \alpha_g(y) = y\}$ has infinite index in Y . Equip Y with its Haar measure.*

Then the action of the semidirect product $\Lambda \rtimes \Gamma$ on Y given by $(\lambda, g) \cdot y = \lambda \alpha_g(y)$ is essentially free.

Proof. Take $(\lambda, g) \neq e$ and consider the set of y with $(\lambda, g) \cdot y = y$. If this set is non empty, it is a coset of $\{y \in Y \mid \alpha_g(y) = y\}$. If $g \neq e$, the latter is a closed subgroup of Y with infinite index and hence has measure zero. So we may assume that $g = e$ and $\lambda \neq e$, which is the trivial case. \square

Take a countable group Γ and a compact abelian group K with dense countable subgroup $\Lambda < K$. Put $X := K^\Gamma$ and consider the action $\Gamma \times \Lambda \curvearrowright X$ given by $((g, s) \cdot x)_h = s x_{hg}$ for all $g, h \in \Gamma, s \in \Lambda, x \in X$. Write $M = L^\infty(X) \rtimes (\Gamma \times \Lambda)$.

Assume that $K_1 < K$ is a closed subgroup. We view $L^\infty(X/K_1)$ as the subalgebra of K_1 -invariant functions in $L^\infty(X)$. We put $\Lambda_1 := \Lambda \cap K_1$ and define $\mathcal{C}(K_1)$ as the von Neumann subalgebra of M generated by $L^\infty(X/K_1)$ and $L(\Lambda_1)$. So,

$$\mathcal{C}(K_1) = L^\infty(X/K_1) \rtimes \Lambda_1 \subset M.$$

Note that we can identify $\mathcal{C}(K_1) = L^\infty(X/K_1) \overline{\otimes} L\Lambda_1 = L^\infty(X/K_1 \times \widehat{\Lambda_1})$.

Lemma 6. *With the above notations, $\mathcal{C}(K_1)' \cap M = \mathcal{C}(\overline{\Lambda_1})$. So $\mathcal{C}(K_1) \subset M$ is maximal abelian iff $\Lambda \cap K_1$ is dense in K_1 .*

In that case, $\mathcal{C}(K_1)$ is a Cartan subalgebra of M and the induced equivalence relation on $X/K_1 \times \widehat{\Lambda_1}$ is given by the orbits of the product action of $(\Gamma \times \Lambda K_1/K_1) \times \widehat{K_1}$ on $X/K_1 \times \widehat{\Lambda_1}$.

For arbitrary closed subgroups $K_1 < K$, the intersection $\Lambda \cap K_1$ need not be dense in K_1 . But putting $K_2 = \overline{\Lambda \cap K_1}$, we always have that $\Lambda \cap K_2$ is dense in K_2 .

Proof. By writing the Fourier decomposition of an element in $M = L^\infty(X) \rtimes (\Gamma \times \Lambda)$, one easily checks that

$$(L\Lambda_1)' \cap M = L^\infty(X/\overline{\Lambda_1}) \rtimes (\Gamma \times \Lambda). \quad (1)$$

View $X = K^\Gamma$ as a compact group and view K_1 as a closed subgroup of X sitting in X diagonally. Put $Y = X/K_1$ and view Λ/Λ_1 as a subgroup of Y sitting in Y diagonally. By Lemma 5 the action $\Gamma \times \Lambda/\Lambda_1 \curvearrowright Y$ is essentially free. From this we conclude that

$$L^\infty(X/K_1)' \cap M = L^\infty(X) \rtimes \Lambda_1.$$

In combination with (1) we get that $\mathcal{C}(K_1)' \cap M = \mathcal{C}(\overline{\Lambda_1})$. This proves the first part of the lemma.

Assume now that $\Lambda \cap K_1$ is dense in K_1 . We know that $\mathcal{C}(K_1) \subset M$ is maximal abelian. For every $\omega \in \widehat{K}$ we define the unitary $U_\omega \in L^\infty(X)$ given by $U_\omega(x) = \omega(x_e)$. One checks that $\mathcal{C}(K_1)$ is normalized by all unitaries in $\mathcal{C}(K_1)$, $\{U_\omega \mid \omega \in \widehat{K}\}$, $\{u_g \mid g \in \Gamma\}$ and $\{u_s \mid s \in \Lambda\}$. These unitaries generate M so that $\mathcal{C}(K_1)$ is indeed a Cartan subalgebra of M .

It remains to analyze which automorphisms of $\mathcal{C}(K_1) = L^\infty(X/K_1 \times \widehat{\Lambda_1})$ are induced by the above normalizing unitaries. The automorphism $\text{Ad}U_\omega$ only acts on $\widehat{\Lambda_1}$ by first restricting ω to a character on Λ_1 and then translating by this character on $\widehat{\Lambda_1}$. The automorphism $\text{Ad}u_g$, $g \in \Gamma$, only acts on X/K_1 by the quotient of the Bernoulli shift. Finally the automorphism $\text{Ad}u_s$, $s \in \Lambda$, also acts only on X/K_1 by first projecting s onto $\Lambda/\Lambda_1 = \Lambda K_1/K_1$ and then diagonally translating with this element. The resulting orbit equivalence relation indeed corresponds to the direct product of $\mathcal{R}(\Gamma \times \Lambda K_1/K_1 \curvearrowright X/K_1)$ and $\mathcal{R}(\widehat{K_1} \curvearrowright \widehat{\Lambda_1})$. \square

Remark 7. Lemma 6 says that the equivalence relation induced by $\mathcal{C}(K_1) \subset M$ is the direct product of the orbit relation $\mathcal{R}((\Gamma \times \Lambda K_1/K_1) \curvearrowright X/K_1)$ and the hyperfinite II_1 equivalence relation. Nevertheless, if Γ is an icc property (T) group, the II_1 factor M is not McDuff, i.e. cannot be written as a tensor product of some II_1 factor with the hyperfinite II_1 factor. Indeed, by property (T), all central sequences in M lie asymptotically in the relative commutant $(L\Gamma)' \cap M = L\Lambda$. Since $L\Lambda$ is abelian, it follows that M is not McDuff. So whenever Γ is an icc property (T) group, it follows that the 2-cocycle associated with the Cartan subalgebra $\mathcal{C}(K_1) \subset M$ is non trivial.

To prove Theorem 1, we first have to classify the orbit equivalence relations of the product actions $(\Gamma \times \Lambda K_1/K_1) \times \widehat{K_1}$ on $X/K_1 \times \widehat{\Lambda_1}$, when K_1 runs through closed subgroups of K

with $\Lambda \cap K_1$ dense in K_1 . These orbit equivalence relations are the direct product of the orbit equivalence relation of $(\Gamma \times \Lambda K_1/K_1) \curvearrowright X/K_1$ and the unique hyperfinite II_1 equivalence relation. In [31, Theorem 4.1], the actions $(\Gamma \times \Lambda K_1/K_1) \curvearrowright X/K_1$ were classified up to stable orbit equivalence. In Lemma 9, we redo the proof in the context of a direct product with a hyperfinite equivalence relation. The main difficulty is to make sure that the orbit equivalence “does not mix up too much” the first and second factors in the direct product. The orbit equivalence will however not simply split as a direct product of orbit equivalences. The main reason for this is that the II_1 factor $N := L^\infty(X/K_1) \rtimes (\Gamma \times \Lambda K_1/K_1)$ has property Gamma. Therefore we cannot apply [28, Theorem 5.1], which says the following: if N is a II_1 factor without property Gamma and R is the hyperfinite II_1 factor, then every automorphism of $N \bar{\otimes} R$ essentially splits as a tensor product of two automorphisms.

To state and prove Lemma 9, we use the following point of view on stable orbit equivalences. Assume that $\mathcal{G}_i \curvearrowright Z_i, i = 1, 2$, are free ergodic p.m.p. actions of countable groups \mathcal{G}_i . By definition, a stable orbit equivalence between $\mathcal{G}_1 \curvearrowright Z_1$ and $\mathcal{G}_2 \curvearrowright Z_2$ is a measure space isomorphism $\Delta : \mathcal{U}_1 \rightarrow \mathcal{U}_2$ between non negligible subsets $\mathcal{U}_i \subset Z_i$ satisfying

$$\Delta(\mathcal{G}_1 \cdot x \cap \mathcal{U}_1) = \mathcal{G}_2 \cdot \Delta(x) \cap \mathcal{U}_2$$

for a.e. $x \in \mathcal{U}_1$. By ergodicity of $\mathcal{G}_1 \curvearrowright Z_1$, we can choose a measurable map $\Delta_0 : Z_1 \rightarrow \mathcal{U}_1$ satisfying $\Delta_0(x) \in \mathcal{G}_1 \cdot x$ for a.e. $x \in Z_1$. Denote $\Psi := \Delta \circ \Delta_0$. By construction Ψ is a local isomorphism from Z_1 to Z_2 , meaning that $\Psi : Z_1 \rightarrow Z_2$ is a Borel map such that Z_1 can be partitioned into a sequence of non negligible subsets $\mathcal{W} \subset Z_1$ such that the restriction of Ψ to any of these subsets \mathcal{W} is a measure space isomorphism of \mathcal{W} onto some non negligible subset of Z_2 . Also by construction Ψ is orbit preserving, meaning that for a.e. $x, y \in Z_1$ we have that $x \in \mathcal{G}_1 \cdot y$ iff $\Psi(x) \in \mathcal{G}_2 \cdot \Psi(y)$.

Using the inverse $\Delta^{-1} : \mathcal{U}_2 \rightarrow \mathcal{U}_1$ we analogously find an orbit preserving local isomorphism $\bar{\Psi} : Z_2 \rightarrow Z_1$. By construction $\bar{\Psi}(\Psi(x)) \in \mathcal{G}_1 \cdot x$ for a.e. $x \in Z_1$ and $\Psi(\bar{\Psi}(y)) \in \mathcal{G}_2 \cdot y$ for a.e. $y \in Z_2$. We call $\bar{\Psi}$ a *generalized inverse* of Ψ .

A measurable map $\Psi : Z_1 \rightarrow Z_2$ between standard measure spaces is called a *factor map* iff $\Psi^{-1}(\mathcal{V})$ has measure zero whenever $\mathcal{V} \subset Z_2$ has measure zero. If $\mathcal{G}_i \curvearrowright Z_i, i = 1, 2$, are free ergodic p.m.p. actions of the countable groups \mathcal{G}_i and if $\Psi : Z_1 \rightarrow Z_2, \bar{\Psi} : Z_2 \rightarrow Z_1$ are factor maps satisfying

$$\begin{aligned} \Psi(\mathcal{G}_1 \cdot x) &\subset \mathcal{G}_2 \cdot \Psi(x) \quad \text{and} \quad \bar{\Psi}(\Psi(x)) \in \mathcal{G}_1 \cdot x \quad \text{for a.e. } x \in Z_1, \\ \bar{\Psi}(\mathcal{G}_2 \cdot y) &\subset \mathcal{G}_1 \cdot \bar{\Psi}(y) \quad \text{and} \quad \Psi(\bar{\Psi}(y)) \in \mathcal{G}_2 \cdot y \quad \text{for a.e. } y \in Z_2, \end{aligned}$$

then Ψ and $\bar{\Psi}$ are local isomorphisms, stable orbit equivalences and each other's generalized inverse.

If $\Psi, \Psi' : Z_1 \rightarrow Z_2$ are factor maps that send a.e. orbits into orbits, we say that Ψ and Ψ' are *similar* if $\Psi'(x) \in \mathcal{G}_2 \cdot \Psi(x)$ for a.e. $x \in Z_1$.

If $\Gamma \curvearrowright (X, \mu)$ and $\Lambda \curvearrowright (Y, \eta)$ are p.m.p. actions of second countable locally compact groups and if $\delta : \Gamma \rightarrow \Lambda$ is an isomorphism, we call $\Delta : X \rightarrow Y$ a δ -conjugacy if Δ is a measure space isomorphism satisfying $\Delta(g \cdot x) = \delta(g) \cdot \Delta(x)$ a.e.

We start off by the following well known and elementary lemma. For the convenience of the reader, we give a proof.

Lemma 8. *Let $\mathcal{G}_i \curvearrowright Z_i$ be free ergodic p.m.p. actions of countable groups. Let $\Psi : Z_1 \rightarrow Z_2$ be a stable orbit equivalence.*

- If $\Lambda < \mathcal{G}_1$ is a subgroup and $\Psi(g \cdot x) = \Psi(x)$ for all $g \in \Lambda$ and a.e. $x \in Z_1$, then Λ is a finite group.
- If $\delta : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ is an isomorphism of \mathcal{G}_1 onto \mathcal{G}_2 and $\Psi(g \cdot x) = \delta(g) \cdot \Psi(x)$ for all $g \in \mathcal{G}_1$ and a.e. $x \in Z_1$, then Ψ is a measure space isomorphism.

Proof. To prove the first point, take a non negligible subset $\mathcal{U} \subset Z_1$ such that $\Psi|_{\mathcal{U}}$ is a measure space isomorphism of \mathcal{U} onto $\mathcal{V} \subset Z_2$. Since $\mathcal{G}_1 \curvearrowright Z_1$ is essentially free, it follows that $g \cdot \mathcal{U} \cap \mathcal{U}$ is negligible for all $g \in \Lambda - \{e\}$. Hence the sets $(g \cdot \mathcal{U})_{g \in \Lambda}$ are essentially disjoint. Since Z_1 has finite measure and \mathcal{U} is non negligible, it follows that Λ is finite.

We now prove the second point. Since Ψ is a local isomorphism, the image $\Psi(Z_1)$ is a measurable, $\delta(\mathcal{G}_1)$ -invariant, non negligible subset of Z_2 . Since $\delta(\mathcal{G}_1) = \mathcal{G}_2$ and $\mathcal{G}_2 \curvearrowright Z_2$ is ergodic, it follows that Ψ is essentially surjective. Assume that Ψ is not a measure space isomorphism. Since Ψ is a local isomorphism, we find non negligible disjoint subsets $\mathcal{U}_1, \mathcal{U}_2 \subset Z_1$ such that $\Psi|_{\mathcal{U}_i}$, $i = 1, 2$, are measure space isomorphisms onto the same subset $\mathcal{V} \subset Z_2$. Since Ψ is a stable orbit equivalence, $(\Psi|_{\mathcal{U}_2})^{-1}(\Psi(x)) \in \mathcal{G}_1 \cdot x$ for a.e. $x \in \mathcal{U}_1$. Making $\mathcal{U}_1, \mathcal{U}_2$ smaller, we may assume that there exists a $g \in \mathcal{G}_1$ such that $(\Psi|_{\mathcal{U}_2})^{-1}(\Psi(x)) = g \cdot x$ for a.e. $x \in \mathcal{U}_1$. Since \mathcal{U}_1 and \mathcal{U}_2 are disjoint, it follows that $g \neq e$. But it also follows that $\delta(g) \cdot \Psi(x) = \Psi(x)$ for a.e. $x \in \mathcal{U}_1$. Since $\mathcal{G}_2 \curvearrowright Z_2$ is essentially free, we arrive at the contradiction that $\delta(g) = e$. \square

Lemma 9. Let Γ be an icc property (T) group with $[\Gamma, \Gamma] = \Gamma$. As above take a compact abelian group K with dense countable subgroup $\Lambda < K$. Put $X := K^\Gamma$.

Let $K_i < K$, $i = 1, 2$, be closed subgroups. Denote $G_i := \Gamma \times \Lambda K_i$ with its natural action on X given by

$$G_i \curvearrowright K^\Gamma : ((g, k) \cdot x)_h = k x_{hg} \quad \text{for all } g, h \in \Gamma, k \in \Lambda K_i, x \in K^\Gamma.$$

Let $H_i \curvearrowright Y_i$ be free ergodic p.m.p. actions of the countable abelian groups H_i . Assume that

$$\Delta : X/K_1 \times Y_1 \rightarrow X/K_2 \times Y_2$$

is a stable orbit equivalence between the product actions $G_i/K_i \times H_i \curvearrowright X/K_i \times Y_i$. Then there exists

- a compact subgroup $K'_1 < \Lambda K_1$ that is commensurate with K_1 , with corresponding canonical stable orbit equivalence $\Delta_1 : X/K'_1 \rightarrow X/K_1$ between the actions $G_1/K'_1 \curvearrowright X/K'_1$ and $G_1/K_1 \curvearrowright X/K_1$;
- an automorphism $\delta_1 \in \text{Aut} \Gamma$ and an isomorphism $\delta_2 : \Lambda K_1/K'_1 \rightarrow \Lambda K_2/K_2$;
- a stable orbit equivalence $\Psi : Y_1 \rightarrow Y_2$ between the actions $H_i \curvearrowright Y_i$;
- a measurable family $(\Theta_y)_{y \in Y_1}$ of $(\delta_1 \times \delta_2)$ -conjugacies between the actions $G_1/K'_1 \curvearrowright X/K'_1$ and $G_2/K_2 \curvearrowright X/K_2$;
- a δ_1 -conjugacy Θ_1 of $\Gamma \curvearrowright X/K$;

such that the stable orbit equivalence $\Delta \circ (\Delta_1 \times \text{id})$ is similar with the stable orbit equivalence

$$\Theta : X/K'_1 \times Y_1 \rightarrow X/K_2 \times Y_2 : (x, y) \mapsto (\Theta_y(x), \Psi(y))$$

and such that $\Theta_y(x K'_1)K = \Theta_1(xK)$ a.e.

Note that we turn G_i into a locally compact group by requiring that $K_i < G_i$ is a compact open subgroup.

Proof. Denote

$$\Theta : X \times Y_1 \rightarrow X/K_2 \times Y_2 : \Theta(x, y) = \Delta(xK_1, y).$$

Associated with the orbit map Θ are two measurable families of cocycles:

$$\begin{aligned} \omega^y : G_1 \times X &\rightarrow G_2/K_2 \times H_2 \quad \text{determined by } \Theta(g \cdot x, y) = \omega^y(g, x) \cdot \Theta(x, y), \\ \mu^x : H_1 \times Y_1 &\rightarrow G_2/K_2 \times H_2 \quad \text{determined by } \Theta(x, h \cdot y) = \mu^x(h, y) \cdot \Theta(x, y). \end{aligned}$$

By a combination of Popa's cocycle superrigidity theorem for $\Gamma \curvearrowright K^\Gamma$ (see [26, Theorem 0.1]) and the elementary [33, Lemma 5.5], we can replace Θ by a similar factor map and find measurable families of continuous group homomorphisms

$$\delta_y : G_1 \rightarrow G_2/K_2 \quad \text{and} \quad \eta_y : G_1 \rightarrow H_2$$

such that

$$\omega^y(g, x) = (\delta_y(g), \eta_y(g)) \quad \text{and} \quad \Theta(g \cdot x, y) = (\delta_y(g), \eta_y(g)) \cdot \Theta(x, y) \quad \text{a.e.}$$

Since $[\Gamma, \Gamma] = \Gamma$ and H_2 is abelian, it follows that $\eta_y(g) = e$ for all $g \in \Gamma$ and a.e. $y \in Y_1$. Writing $\Theta(x, y) = (\Theta_1(x, y), \Theta_2(x, y))$, it follows that $\Theta_2(g \cdot x, y) = \Theta_2(x, y)$ for all $g \in \Gamma$ and a.e. $(x, y) \in X \times Y_1$. By ergodicity of $\Gamma \curvearrowright X$, it follows that $\Theta_2(x, y) = \Psi(y)$ for some factor map $\Psi : Y_1 \rightarrow Y_2$. But then

$$\eta_y(g) \cdot \Psi(y) = \eta_y(g) \cdot \Theta_2(x, y) = \Theta_2(g \cdot x, y) = \Psi(y)$$

for all $g \in G_1$ and a.e. $(x, y) \in X \times Y_1$. It follows that $\eta_y(g) = e$ for all $g \in G_1$ and a.e. $y \in Y_1$.

Denote $\mu^x(h, y) = (\mu_1^x(h, y), \mu_2^x(h, y))$. Because

$$\mu_2^x(h, y) \cdot \Psi(y) = \mu_2^x(h, y) \cdot \Theta_2(x, y) = \Theta_2(x, h \cdot y) = \Psi(h \cdot y)$$

for all $h \in H_1$ and a.e. $(x, y) \in X \times Y_1$, it follows that $\mu_2^x(h, y)$ is essentially independent of the x -variable. So we write $\mu_2(h, y)$ instead of $\mu_2^x(h, y)$.

Next we observe that $\Theta_1(g \cdot x, h \cdot y)$ can be computed in two ways, leading to the equality

$$\delta_{h \cdot y}(g) \mu_1^x(h, y) \cdot \Theta_1(x, y) = \mu_1^{g \cdot x}(h, y) \delta_y(g) \cdot \Theta_1(x, y) \quad \text{a.e.}$$

So, $\mu_1^{g \cdot x}(h, y) = \delta_{h \cdot y}(g) \mu_1^x(h, y) \delta_y(g)^{-1}$ for all $g \in G_1, h \in H_1$ and a.e. $(x, y) \in X \times Y_1$.

It follows that for all $h \in H_1$ and a.e. $y \in Y_1$, the non negligible set $\{(x, x') \in X \times X \mid \mu_1^x(h, y) = \mu_1^{x'}(h, y)\}$ is invariant under the diagonal action of Γ on $X \times X$. Since $\Gamma \curvearrowright X$ is weakly mixing, we conclude that $\mu_1^x(h, y)$ is essentially independent of the x -variable. We write $\mu_1(h, y)$ instead of $\mu_1^x(h, y)$. The above formula becomes

$$\delta_{h \cdot y}(g) = \mu_1(h, y) \delta_y(g) \mu_1(h, y)^{-1}. \quad (2)$$

It follows that the map $y \mapsto \text{Ker} \delta_y$ is H_1 -invariant and hence, by ergodicity of $H_1 \curvearrowright Y_1$, there is a unique closed subgroup $K'_1 < G_1$ such that $\text{Ker} \delta_y = K'_1$ for a.e. $y \in Y_1$. Since G_2/K_2 is a countable group, we know that its compact subgroup $\delta_y(K_1)$ must be finite. Hence, $K'_1 \cap K_1 < K_1$ has finite index. We claim that also $K'_1 \cap K_1 < K'_1$ has finite index.

To prove this claim, first observe that the formula $\Theta(g \cdot x, y) = \delta_y(g) \cdot \Theta(x, y)$ implies that we can view Θ as a map from $X/K'_1 \times Y_1$ to $X/K_2 \times Y_2$. In particular Θ can be viewed as a map from $X/(K_1 \cap K'_1) \times Y_1$ to $X/K_2 \times Y_2$. Since $X/(K_1 \cap K'_1)$ is a finite covering of X/K_1 and since the original Δ was a local isomorphism, also

$$\Theta : X/(K_1 \cap K'_1) \times Y_1 \rightarrow X/K_2 \times Y_2$$

is a local isomorphism and hence a stable orbit equivalence. Since moreover $\Theta(g \cdot x, y) = \Theta(x, y)$ for all $g \in K'_1/(K_1 \cap K'_1)$, it follows from [Lemma 8](#) that $K'_1/(K_1 \cap K'_1)$ is a finite group. This proves the claim, meaning that K'_1 and K_1 are commensurate subgroups of G_1 . In particular, K'_1 is compact. Denoting by $p_\Gamma : G_1 \rightarrow \Gamma$ the projection homomorphism onto the first factor, we get that $p_\Gamma(K'_1)$ is a finite normal subgroup of Γ . Since Γ is icc, it follows that $p_\Gamma(K'_1) = \{e\}$ and thus $K'_1 < \Lambda K_1$.

By construction the map

$$\Theta : X/K'_1 \times Y_1 \rightarrow X/K_2 \times Y_2$$

is a stable orbit equivalence with the following properties:

- Θ is similar with $\Delta \circ (\Delta_1 \times \text{id})$;
- Θ is of the form $\Theta(x, y) = (\Theta_y(x), \Psi(y))$;
- Θ satisfies $\Theta_y(g \cdot x) = \delta_y(g) \cdot \Theta_y(x)$ a.e. where $\delta_y : G_1/K'_1 \rightarrow G_2/K_2$ is a measurable family of injective group homomorphisms.

It remains to prove that the δ_y are group isomorphisms that do not depend on y , Ψ is a stable orbit equivalence and a.e. Θ_y is a measure space isomorphism of X/K'_1 onto X/K_2 .

Applying the same reasoning to the inverse of the stable orbit equivalence $\Delta \circ (\Delta_1 \times \text{id})$, we find a closed subgroup $K'_2 < \Lambda K_2$ that is commensurate with K_2 , a measurable family of injective group homomorphisms $\bar{\delta}_y : G_2/K'_2 \rightarrow G_1/K'_1$ and a map

$$\bar{\Theta} : X/K'_2 \times Y_2 \rightarrow X/K'_1 \times Y_1$$

of the form $\bar{\Theta}(x, y) = (\bar{\Theta}_y(x), \bar{\Psi}(y))$ satisfying the following two properties:

$$\bar{\Theta}_y(g \cdot x) = \bar{\delta}_y(g) \cdot \bar{\Theta}_y(x) \quad \text{and}$$

$$\Theta(\bar{\Theta}(xK'_2, y)) \in (G_2/K_2 \times H_2) \cdot (xK_2, y) \quad \text{a.e.} \quad (3)$$

The second of these formulas yields a measurable map $\varphi : X \times Y_2 \rightarrow G_2/K_2$ such that

$$\Theta_{\bar{\Psi}(y)}(\bar{\Theta}_y(xK'_2)) = \varphi(x, y) \cdot xK_2 \quad \text{a.e.}$$

Computing $\Theta_{\bar{\Psi}(y)}(\bar{\Theta}_y(gK'_2 \cdot xK'_2))$ it follows that

$$\delta_{\bar{\Psi}(y)}(\bar{\delta}_y(gK'_2)) \varphi(x, y) = \varphi(g \cdot x, y) gK_2 \quad \text{for all } g \in G_2 \text{ and a.e. } (x, y) \in X \times Y_2.$$

So, for almost every $y \in Y_2$, the non negligible set $\{(x, x') \in X \times X \mid \varphi(x, y) = \varphi(x', y)\}$ is invariant under the diagonal action of Γ on $X \times X$. Since $\Gamma \curvearrowright X$ is weakly mixing, it follows that $\varphi(x, y)$ is essentially independent of the x -variable. We write $\varphi(y)$ instead of $\varphi(x, y)$ and find that

$$(\delta_{\bar{\Psi}(y)} \circ \bar{\delta}_y)(gK'_2) = \varphi(y) gK_2 \varphi(y)^{-1}.$$

It follows that $K'_2 = K_2$ and that $\delta_{\bar{\Psi}(y)} \circ \bar{\delta}_y$ is an inner automorphism of G_2/K_2 . It follows in particular that $\delta_{\bar{\Psi}(y)}$ is surjective, so that δ_y is an isomorphism of G_1/K'_1 onto G_2/K_2 for all y in a non negligible subset of Y_1 . Because of (2) and the ergodicity of $H_1 \curvearrowright Y_1$, it follows that δ_y is an isomorphism for a.e. $y \in Y_1$.

It also follows from (3) that $\bar{\Psi}(\bar{\Psi}(y)) \in H_2 \cdot y$ for a.e. $y \in Y_2$ and similarly with $\bar{\Psi} \circ \Psi$. Hence Ψ and $\bar{\Psi}$ are local isomorphisms and hence a stable orbit equivalence between $H_1 \curvearrowright Y_1$ and $H_2 \curvearrowright Y_2$. Since $\Theta(x, y) = (\Theta_y(x), \Psi(y))$ and since Θ is a local isomorphism, it then follows

that a.e. Θ_y is a local isomorphism. But Θ_y is also a δ_y -conjugation w.r.t. the isomorphism δ_y . Lemma 8 implies that Θ_y is a measure space isomorphism for a.e. $y \in Y_1$.

We know that $[\Gamma, \Gamma] = \Gamma$, Γ has trivial center and the groups $\Lambda K_i/K'_i$ are abelian. So it follows that δ_y is the direct product of an automorphism δ_y^1 of Γ and an isomorphism δ_y^2 of $\Lambda K_1/K'_1$ onto $\Lambda K_2/K_2$. Formula (2) and the fact that $\Lambda K_2/K_2$ is the center of G_2/K_2 imply that $\delta_{h \cdot y}^2 = \delta_y^2$ a.e. By ergodicity of $H_1 \curvearrowright Y_1$, it follows that δ_y^2 is essentially independent of y . We write δ_2 instead of δ_y^2 . Next denote by $\mu_\Gamma(h, y)$ the projection of $\mu_1(h, y) \in \Gamma \times \Lambda K_2/K_2$ onto Γ . Formula (2) says that

$$\delta_{h \cdot y}^1 = \text{Ad} \mu_\Gamma(h, y) \circ \delta_y^1.$$

Since Γ has property (T), Γ is finitely generated and $\text{Aut} \Gamma$ is a countable group. In particular, $\text{Out} \Gamma = \text{Aut} \Gamma / \text{Inn} \Gamma$ is a countable group. Then the map $y \mapsto (\delta_y^1 \text{ mod } \text{Inn} \Gamma)$ is an H_1 -invariant measurable map from Y_1 to $\text{Out} \Gamma$ and hence is essentially constant. So we find a measurable map $\varphi : Y_1 \rightarrow \Gamma$ and an automorphism $\delta_1 \in \text{Aut} \Gamma$ such that $\delta_y^1 = \text{Ad} \varphi(y) \circ \delta_1$ a.e. Replacing $\Theta_y(x)$ by $\varphi(y)^{-1} \cdot \Theta_y(x)$ we may assume that δ_y is a.e. equal to $\delta_1 \times \delta_2$. After this replacement, also $\mu_\Gamma(h, y) = e$ a.e. meaning that $\mu_1(h, y) \in \Lambda K_2/K_2$.

Since Θ_y is a δ_2 -conjugacy and $\Lambda K_1, \Lambda K_2$ are dense subgroups of K , then Θ_y induce δ_1 -conjugacies ρ_y of the action $\Gamma \curvearrowright X/K$ given by $\rho_y(xK) = \Theta_y(xK'_1)K$. Since $\mu_1(h, y) \in \Lambda K_2/K_2$, it follows that $\rho_{h \cdot y}(x) = \rho_y(x)$ a.e. Hence ρ_y is a.e. equal to a single δ_1 -conjugacy Θ_1 of the action $\Gamma \curvearrowright X/K$. \square

Lemma 10. *Let K be a compact abelian group with countable dense subgroup Λ . Let $K_1, K_2 < K$ be closed subgroups and denote $\Lambda_i := \Lambda \cap K_i$.*

1. *If $K_2 \subset \Lambda K_1$ and if $\Lambda K_1/K_2$ is countable, then K_1 and K_2 are commensurate.*
2. *Assume that K_1 and K_2 are commensurate and that $K_2 \subset \Lambda K_1$. If Λ_1 is dense in K_1 , also Λ_2 is dense in K_2 .*
3. *Assume that Λ_i is dense in K_i for $i = 1, 2$. Then the following statements are equivalent.*
 - (a) *Λ_1 and Λ_2 are commensurate.*
 - (b) *K_1 and K_2 are commensurate.*
 - (c) *$\Lambda K_1 = \Lambda K_2$.*

Proof. 1. Since $\Lambda K_1/K_2$ is countable, the image of K_1 in K/K_2 is a countable compact group, hence a finite group. This means that $K_1 \cap K_2 < K_1$ has finite index. Since $K_2 \subset \Lambda K_1$, also the image of K_2 in K/K_1 is countable, hence finite. This means that $K_1 \cap K_2 < K_2$ has finite index. So, K_1 and K_2 are commensurate.

2. Assume that K_1 and K_2 are commensurate, $K_2 \subset \Lambda K_1$ and Λ_1 is dense in K_1 . Since $K_1 \cap K_2 < K_1$ has finite index, $K_1 \cap K_2$ is an open subgroup of K_1 . Because Λ_1 is dense in K_1 , we get that $\Lambda_1 \cap (K_1 \cap K_2)$ is dense in $K_1 \cap K_2$. So, $\Lambda_1 \cap \Lambda_2$ is dense in $K_1 \cap K_2$. Also the image of Λ_1 in $K_1/(K_1 \cap K_2)$ is dense. But $K_1/(K_1 \cap K_2)$ is finite and it follows that $K_1 = \Lambda_1(K_1 \cap K_2)$. Hence also $\Lambda K_1 = \Lambda(K_1 \cap K_2)$. Since by assumption $K_2 \subset \Lambda K_1$, we get that $K_2 \subset \Lambda(K_1 \cap K_2)$, i.e. $K_2 = \Lambda_2(K_1 \cap K_2)$. We already saw that $\Lambda_1 \cap \Lambda_2$ is dense in $K_1 \cap K_2$ and it follows that Λ_2 is dense in K_2 .

3. (a) \Rightarrow (b). The image of Λ_i in $K_i/(K_1 \cap K_2)$ is dense, but also finite because $\Lambda_1 \cap \Lambda_2$ has finite index in Λ_i . So, $K_i/(K_1 \cap K_2)$ is finite for $i = 1, 2$, meaning that K_1 and K_2 are commensurate.

(b) \Rightarrow (c). Since Λ_i is dense in K_i and $K_1 \cap K_2 < K_i$ has finite index, we get as in the proof of 2 that $K_i = \Lambda_i(K_1 \cap K_2)$. Hence, $\Lambda K_i = \Lambda(K_1 \cap K_2)$ for $i = 1, 2$. In particular, $\Lambda K_1 = \Lambda K_2$.

(c) \Rightarrow (b). We have $K_2 \subset \Lambda K_1$ and we have that $\Lambda K_1/K_2 = \Lambda K_2/K_2$ is countable. So, statement 1 implies that K_1 and K_2 are commensurate.

(b) \Rightarrow (a). Since $K_i/(K_1 \cap K_2)$ is finite, also the image of Λ_i in $K_i/(K_1 \cap K_2)$ is finite, meaning that $\Lambda_i/(\Lambda_1 \cap \Lambda_2)$ is a finite group. So, Λ_1 and Λ_2 are commensurate. \square

Proof of Theorem 1

Proof. *Proof of The Equivalence of the Statements in 1.* Put $\Lambda_i = \Lambda \cap K_i$. Using Lemma 10, it suffices to prove that $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are unitarily conjugate if and only if Λ_1 and Λ_2 are commensurate. First assume that Λ_1 and Λ_2 are commensurate. By Lemma 10, also K_1 and K_2 are commensurate. So, $K_i < K_1 K_2$ has finite index. It follows that $L^\infty(K^\Gamma/K_1 K_2) \bar{\otimes} L(\Lambda_1 \cap \Lambda_2)$ is a subalgebra of finite index in both $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$. By [24, Theorem A.1], the Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are unitarily conjugate.

Conversely, assume that Λ_1 and Λ_2 are not commensurate. Assume for instance that $\Lambda_1 \cap \Lambda_2 < \Lambda_1$ has infinite index. We find a sequence $\lambda_n \in \Lambda_1$ such that for every finite subset $\mathcal{F} \subset \Lambda_1$ the element λ_n lies outside $\mathcal{F} \Lambda_2$ eventually. Denote by $u_n := u_{\lambda_n}$ the unitaries in $\mathcal{C}(K_1)$ that correspond to the group elements λ_n . Denoting by $E_{\mathcal{C}(K_2)}$ the trace preserving conditional expectation, one deduces that

$$\|E_{\mathcal{C}(K_2)}(au_nb)\|_2 \rightarrow 0 \quad \text{for all } a, b \in M.$$

So, by [24, Theorem A.1], the Cartan subalgebras $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are not unitarily conjugate.

Proof of the equivalence of the statements in 3. Denote $X = K^\Gamma$. Assume first that there exists a continuous automorphism $\delta \in \text{Aut} K$ satisfying $\delta(\Lambda K_1) = \Lambda K_2$. Then $K'_1 := \delta^{-1}(K_2)$ is a closed subgroup of K with the properties that $K'_1 \subset \Lambda K_1$ and that $\Lambda K_1/K'_1$ is countable. By Lemma 10.1, we get that K'_1 and K_1 are commensurate. Then Lemma 10.2 implies that $\Lambda \cap K'_1$ is dense in K'_1 . By the equivalence of the statements in 1, we know that the Cartan subalgebras $\mathcal{C}(K'_1)$ and $\mathcal{C}(K_1)$ are unitarily conjugate. In particular, they give rise to isomorphic equivalence relations. So we may replace K_1 by K'_1 and assume that $\delta(K_1) = K_2$. By “applying everywhere” δ , the actions $(\Gamma \times \Lambda K_i/K_i) \curvearrowright X/K_i$ are conjugate, hence orbit equivalent. Also their respective direct products with the hyperfinite equivalence relation are orbit equivalent. By Lemma 6, the equivalence relations induced by $\mathcal{C}(K_i)$, $i = 1, 2$, are orbit equivalent.

Conversely, assume that the equivalence relations induced by $\mathcal{C}(K_i)$, $i = 1, 2$, are stably orbit equivalent. Put $G_i := \Gamma \times \Lambda K_i$. We combine Lemmas 6 and 9, and we replace K_1 by a commensurate group that is contained in ΛK_1 . By Lemma 10.2, the intersection $\Lambda \cap K_1$ remains dense in K_1 . We have found a measure space isomorphism $\Delta : X/K_1 \rightarrow X/K_2$, an automorphism $\delta_1 \in \text{Aut} \Gamma$ and an isomorphism $\delta_2 : \Lambda K_1/K_1 \rightarrow \Lambda K_2/K_2$ such that Δ is a $(\delta_1 \times \delta_2)$ -conjugacy. In particular, Δ is a δ_1 -conjugacy of the Γ -action. By [31, Lemma 5.2] there exists an isomorphism $\alpha : K_1 \rightarrow K_2$ and a $(\delta_1 \times \alpha)$ -conjugacy $\bar{\Delta} : X \rightarrow X$ satisfying $\bar{\Delta}(x)K_2 = \Delta(xK_1)$ for a.e. $x \in X$. Since Δ is a δ_2 -conjugacy, it follows that $\bar{\Delta}(g \cdot x) \in \Lambda K_2 \cdot \Delta(x)$ for all $g \in \Lambda K_1$ and a.e. $x \in X$. We find a measurable map $\omega : \Lambda K_1 \times X \rightarrow \Lambda K_2$ such that $\bar{\Delta}(g \cdot x) = \omega(g, x) \cdot \bar{\Delta}(x)$ a.e. Since $\bar{\Delta}$ is a δ_1 -conjugacy, it follows that $\omega(g, \gamma \cdot x) = \omega(g, x)$ for all $\gamma \in \Gamma$, $g \in \Lambda K_1$ and a.e. $x \in X$. By ergodicity of $\Gamma \curvearrowright X$, the map $\omega(g, x)$ is essentially independent of the x -variable. So we can extend α to a homomorphism $\alpha : \Lambda K_1 \rightarrow \Lambda K_2$. By symmetry (i.e. considering $\bar{\Delta}^{-1}$) it follows that α is an isomorphism of ΛK_1 onto ΛK_2 . Viewing K as a closed subgroup of the group $\text{Aut}(X)$ of measure preserving automorphisms of X (up to equality almost everywhere), it follows that $\bar{\Delta} \Lambda K_1 \bar{\Delta}^{-1} = \Lambda K_2$. Taking closures we also have $\bar{\Delta} K \bar{\Delta}^{-1} = K$. This means that α can be

extended to a continuous automorphism of K . This concludes the proof of the equivalence of the statements in 3.

Proof of the equivalence of the statements in 2. Denote $X = K^\Gamma$. If $\delta \in \text{Aut}K$ is a continuous automorphism satisfying $\delta(\Lambda) = \Lambda$ and $\delta(\Lambda K_1) = \Lambda K_2$, we may replace as above K_1 by the commensurate $\delta^{-1}(K_2)$ and assume that $\delta(K_1) = K_2$. Then also $\delta(\Lambda_1) = \Lambda_2$. So, “applying everywhere” δ yields an automorphism $\alpha \in \text{Aut}M$ satisfying $\alpha(\mathcal{C}(K_1)) = \mathcal{C}(K_2)$.

Conversely, assume that $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ are stably conjugate by an automorphism. Interchanging the roles of K_1 and K_2 if necessary, we find a projection $p \in \mathcal{C}(K_2)$ and an isomorphism $\alpha : M \rightarrow pMp$ satisfying $\alpha(\mathcal{C}(K_1)) = \mathcal{C}(K_2)p$. This α induces a stable orbit equivalence between the equivalence relations associated with $\mathcal{C}(K_1)$ and $\mathcal{C}(K_2)$ respectively. If one of the K_i is finite, the equivalence of the statements in 3 implies that the other one is finite as well and that $\Lambda = \Lambda K_1 = \Lambda K_2$. So we can exclude this trivial case.

We claim that there exist automorphisms $\delta_1 \in \text{Aut}\Gamma$, $\delta \in \text{Aut}K$ and a $(\delta_1 \times \delta)$ -conjugacy $\Delta \in \text{Aut}(X)$ of the action $\Gamma \times K \curvearrowright X$ such that $\delta(K_1) = K_2$ and such that after a unitary conjugacy of α and p , we have

$$p \in L\Lambda_2 \quad \text{and} \quad \alpha(F) = (F \circ \Delta^{-1})p \quad \text{for all } F \in L^\infty(X/K).$$

To prove this claim, denote $Y_i = \widehat{\Lambda_i}$ and $H_i = \widehat{K_i}$. Also denote $G_i = \Gamma \times \Lambda K_i$. [Lemma 6](#) describes the equivalence relations associated with $\mathcal{C}(K_i)$. The isomorphism $\alpha : M \rightarrow pMp$ therefore induces a stable orbit equivalence $\Pi : X/K_1 \times Y_1 \rightarrow X/K_2 \times Y_2$ between the product actions $G_i/K_i \times H_i \curvearrowright X/K_i \times Y_i$. We apply [Lemma 9](#) to Π . We find in particular a compact subgroup $K'_1 < \Lambda K_1$ that is commensurate with K_1 . Denote $\Lambda'_1 := \Lambda \cap K'_1$, $H'_1 := \widehat{K'_1}$ and $Y'_1 := \widehat{\Lambda'_1}$. By [Lemma 10.2](#), Λ'_1 is dense in K'_1 . By the equivalence of the statements in 1, we know that the Cartan subalgebras $\mathcal{C}(K'_1)$ and $\mathcal{C}(K_1)$ are unitarily conjugate. The corresponding orbit equivalence between the product actions $G_1/K'_1 \times H'_1 \curvearrowright X/K'_1 \times Y'_1$ and $G_1/K_1 \times H_1 \curvearrowright X/K_1 \times Y_1$ is similar with $\Delta_1 \times \Delta_2$ where Δ_1, Δ_2 are the natural stable orbit equivalences (whose compression constants are each other's inverse). So replacing $\mathcal{C}(K_1)$ by $\mathcal{C}(K'_1)$ amounts to replacing Π by $\Pi \circ (\Delta_1 \times \Delta_2)$. [Lemma 9](#) says that $\Pi \circ (\Delta_1 \times \text{id})$ is similar to a stable orbit equivalence of a very special form. Then $\Pi \circ (\Delta_1 \times \Delta_2)$ is still of this very special form. So after replacing $\mathcal{C}(K_1)$ by $\mathcal{C}(K'_1)$ we find that Π is similar to a stable orbit equivalence $\Theta : X/K_1 \times Y_1 \rightarrow X/K_2 \times Y_2$ satisfying the following properties.

- Θ is of the form $\Theta(x, y) = (\Theta_y(x), \Psi(y))$ where $\Psi : Y_1 \rightarrow Y_2$ is a stable orbit equivalence between the actions $H_i \curvearrowright Y_i$ and where $(\Theta_y)_{y \in Y_1}$ is a measurable family of measure preserving conjugacies between $G_1/K_1 \curvearrowright X/K_1$ and $G_2/K_2 \curvearrowright X/K_2$.
- There exist automorphisms $\delta_1 \in \text{Aut}\Gamma$, $\delta \in \text{Aut}K$ and a $(\delta_1 \times \delta)$ -conjugacy $\Delta \in \text{Aut}(X)$ such that $\delta(K_1) = K_2$ and $\Theta_y(x)K = \Delta(xK)$ for a.e. $(x, y) \in X/K_1 \times Y_1$.

To deduce the precise form of Δ in the last item, we proceed as in the proof above of the equivalence of the statements in 3. The compression constant of Ψ is equal to the compression constant of Π and hence is equal to $\tau(p)$. So replacing Ψ by a similar stable orbit equivalence, we may assume that Ψ is a measure space isomorphism (scaling the measure by the factor $\tau(p)$) of Y_1 onto a measurable subset $\mathcal{U} \subset Y_2$ of measure $\tau(p)$. Keeping $\Theta_y(x)$ unchanged, the map Θ then becomes a measure space isomorphism of $X/K_1 \times Y_1$ onto $X/K_2 \times \mathcal{U}$. Define the projection $p \in L\Lambda_2 = L^\infty(Y_2)$ by $p = \chi_{\mathcal{U}}$. Since Π and Θ are similar stable orbit equivalences and Θ is moreover a measure space isomorphism, we can unitarily conjugate α so that $\alpha|_{\mathcal{C}(K_1)} = \Theta_*$. This proves the claim above.

It remains to prove that $\delta(\Lambda) = \Lambda$.

Since every automorphism $\delta_1 \in \text{Aut} \Gamma$ defines a natural automorphism of M that globally preserves all the Cartan subalgebras $\mathcal{C}(K_1)$, we may assume that $\delta_1 = \text{id}$. For $g \in \Gamma \times \Lambda$, we denote by $u_g \in M = L^\infty(X) \rtimes (\Gamma \times \Lambda)$ the canonical unitaries. Since the relative commutant of $L^\infty(X/K)$ inside M equals $L^\infty(X) \rtimes \Lambda$, it follows that

$$\alpha(u_g) = \omega_g u_g p \quad \text{for all } g \in \Gamma, \text{ where } \omega_g \in \mathcal{U}(p(L^\infty(X) \rtimes \Lambda)p).$$

Denote by $(\sigma_g)_{g \in \Gamma}$ the action of Γ on $L^\infty(X) \rtimes \Lambda$, implemented by $\text{Ad} u_g$ and corresponding to the Bernoulli action on $L^\infty(X)$ and the trivial action on $L\Lambda$. Since p is invariant under $(\sigma_g)_{g \in \Gamma}$ we can restrict σ_g to $p(L^\infty(X) \rtimes \Lambda)p$. Note that

$$\omega_{gh} = \omega_g \sigma_g(\omega_h) \quad \text{for all } g, h \in \Gamma.$$

Denote $N^\infty := B(\ell^2(\mathbb{N})) \overline{\otimes} N$ whenever N is a von Neumann algebra. By Theorem 11 there exists a projection $q \in (L\Lambda)^\infty$ with $(\text{Tr} \otimes \tau)(q) = \tau(p)$, a partial isometry $v \in B(\mathbb{C}, \ell^2(\mathbb{N})) \overline{\otimes} (L^\infty(X) \rtimes \Lambda)$ and a group homomorphism $\gamma : \Gamma \rightarrow \mathcal{U}(q(L\Lambda)^\infty q)$ satisfying

$$v^* v = p, \quad v v^* = q \quad \text{and} \quad \omega_g = v^* \gamma_g \sigma_g(v) \quad \text{for all } g \in \Gamma.$$

We replace α by $\text{Ad} v \circ \alpha$. Then $\alpha : M \rightarrow q M^\infty q$. Note that $L^\infty(X/K)$ commutes with v so that

$$\alpha(F) = (F \circ \Delta^{-1})q \quad \text{for all } F \in L^\infty(X/K) \quad \text{and} \quad \alpha(u_g) = \gamma_g u_g = u_g \gamma_g$$

$$\text{for all } g \in \Gamma.$$

Denote $\Delta_* \in \text{Aut}(L^\infty(X))$ given by $\Delta_*(F) = F \circ \Delta^{-1}$ and note that $\Delta_*(L^\infty(X/K)) = L^\infty(X/K)$. Also $\alpha(F) = \Delta_*(F)q$ whenever $F \in L^\infty(X/K)$.

Recall that $X = K^\Gamma$. Whenever $\omega \in \widehat{K}$, denote by $U_\omega \in \mathcal{U}(L^\infty(X))$ the unitary given by $U_\omega(x) = \omega(x_\ell)$. Fix $\omega \in \widehat{K}$ and $g \in \Gamma$. Observe that $U_\omega u_g U_\omega^* u_g^*$ belongs to $L^\infty(X/K)$. Hence, since Δ is an id-conjugacy for $\Gamma \curvearrowright X$, we have

$$\alpha(U_\omega u_g U_\omega^* u_g^*) = \Delta_*(U_\omega \sigma_g(U_\omega^*))q = \Delta_*(U_\omega) \sigma_g(\Delta_*(U_\omega^*))q$$

$$= \Delta_*(U_\omega) u_g \Delta_*(U_\omega)^* u_g^* q.$$

On the other hand

$$\alpha(U_\omega u_g U_\omega^* u_g^*) = \alpha(U_\omega) u_g \gamma_g \alpha(U_\omega)^* \gamma_g^* u_g^*.$$

Combining both formulas we conclude that

$$u_g^* \Delta_*(U_\omega)^* \alpha(U_\omega) u_g = \Delta_*(U_\omega)^* \gamma_g \alpha(U_\omega) \gamma_g^* \quad \text{for all } \omega \in \widehat{K}, g \in \Gamma. \quad (4)$$

We claim that for all $\omega \in \widehat{K}$, $\Delta_*(U_\omega)^* \alpha(U_\omega) \in (L\Lambda)^\infty$. To prove this claim, fix $\omega \in \widehat{K}$ and put $a := \Delta_*(U_\omega)^* \alpha(U_\omega)$. Define for every finite subset $\mathcal{F} \subset \Gamma$, the von Neumann subalgebra $N_{\mathcal{F}} \subset (L^\infty(X) \rtimes \Lambda)^\infty$ given by

$$N_{\mathcal{F}} := (L^\infty(K^{\mathcal{F}}) \rtimes \Lambda)^\infty.$$

Denote by $E_{\mathcal{F}}$ the trace preserving conditional expectation of $(L^\infty(X) \rtimes \Lambda)^\infty$ onto $N_{\mathcal{F}}$. Note that $(L\Lambda)^\infty \subset N_{\mathcal{F}}$ for all $\mathcal{F} \subset \Gamma$. Choose $\varepsilon > 0$ and denote by $\|\cdot\|_2$ the 2-norm on $(L^\infty(X) \rtimes \Lambda)^\infty$ given by the semi-finite trace $\text{Tr} \otimes \tau$. Since $\alpha(U_\omega)$ commutes with $L^\infty(X/K)q$, it follows that $\alpha(U_\omega) \in q(L^\infty(X) \rtimes \Lambda)^\infty q$. So we can take a large enough finite subset $\mathcal{F} \subset \Gamma$ such that

$$\|\alpha(U_\omega) - E_{\mathcal{F}}(\alpha(U_\omega))\|_2 < \varepsilon \quad \text{and} \quad \|q \Delta_*(U_\omega) - E_{\mathcal{F}}(q \Delta_*(U_\omega))\|_2 < \varepsilon.$$

It follows that for all $g \in \Gamma$, the element $\Delta_*(U_\omega)^* \gamma_g \alpha(U_\omega) \gamma_g^*$ lies at distance at most 2ε from $N_{\mathcal{F}}$. By (4), the element a then lies at distance at most 2ε from $N_{\mathcal{F}g^{-1}}$. Since a also lies at

distance at most 2ε from $N_{\mathcal{F}}$, we conclude that a lies at distance at most 4ε from $N_{\mathcal{F} \cap \mathcal{F}g^{-1}}$ for all $g \in \Gamma$. We can choose g such that $\mathcal{F} \cap \mathcal{F}g^{-1} = \emptyset$ and conclude that a lies at distance at most 4ε from $(L\Lambda)^\infty$. Since $\varepsilon > 0$ is arbitrary, the claim follows.

Put $V_\omega := \Delta_*(U_\omega)^* \alpha(U_\omega)$ and $q_\omega := V_\omega V_\omega^*$. Denote by $z \in L\Lambda$ the central support of q in $(L\Lambda)^\infty$. Define

$$q_1 = \bigvee_{\omega \in \widehat{K}} V_\omega V_\omega^*.$$

Since $V_\omega \in (L\Lambda)^\infty$ and $V_\omega^* V_\omega = q$, it follows that $q_1 \in (L\Lambda)^\infty$ and $q_1 \leq z$. Since

$$V_\omega V_\omega^* = \Delta_*(U_\omega)^* q \Delta_*(U_\omega)$$

it also follows that q_1 commutes with all the unitaries $\Delta_*(U_\omega)$, $\omega \in \widehat{K}$. Being an element of $(L\Lambda)^\infty$, the projection q_1 certainly commutes with $L^\infty(X/K) = \Delta_*(L^\infty(X/K))$. Hence, q_1 commutes with the whole of $\Delta_*(L^\infty(X)) = L^\infty(X)$. So, $q_1 \in B(\ell^2(\mathbb{N})) \otimes 1$. Since $q_1 \leq z$, it follows that $z = 1$. Because $(\text{Tr} \otimes \tau)(q) = \tau(p) \leq 1$, we must have $p = 1$ and find an element $w \in B(\ell^2(\mathbb{N}), \mathbb{C}) \overline{\otimes} L\Lambda$ satisfying $ww^* = 1$ and $w^*w = q$. Replacing α by $\text{Ad}w \circ \alpha$, we have found that α is an automorphism of M satisfying

$$\begin{aligned} \alpha(F) &= \Delta_*(F) \quad \text{for all } F \in L^\infty(X/K), & \alpha(U_\omega) &= \Delta_*(U_\omega) V_\omega \quad \text{for all } \omega \in \widehat{K}, \text{ and} \\ \alpha(u_g) &= \gamma_g u_g \quad \text{for all } g \in \Gamma. \end{aligned}$$

Here $V_\omega \in L\Lambda$ are unitaries and $g \mapsto \gamma_g$ is a homomorphism from Γ into the abelian group $\mathcal{U}(L\Lambda)$. Since $[\Gamma, \Gamma] = \Gamma$, we actually have that $\gamma_g = 1$ for all $g \in \Gamma$. In particular $\alpha(L\Gamma) = L\Gamma$. Taking the relative commutant, it follows that $\alpha(L\Lambda) = L\Lambda$.

Since $(\text{Ad}U_\omega)(u_s) = \omega(s) u_s$ for all $s \in \Lambda$, $\omega \in \widehat{K}$, we also have that $(\text{Ad}\alpha(U_\omega))(\alpha(u_s)) = \omega(s) \alpha(u_s)$ for all $s \in \Lambda$, $\omega \in \widehat{K}$. Since $\alpha(U_\omega) = \Delta_*(U_\omega) V_\omega$ and V_ω belongs to the abelian algebra $L\Lambda$, we conclude that $(\text{Ad}\Delta_*(U_\omega))(\alpha(u_s)) = \omega(s) \alpha(u_s)$ for all $s \in \Lambda$, $\omega \in \widehat{K}$. On the other hand, since Δ is a δ -conjugation for the action $K \curvearrowright X$, we also have that $(\text{Ad}\Delta_*(U_\omega))(u_k) = \omega(\delta^{-1}(k)) u_k$ for all $k \in \Lambda$ and $\omega \in \widehat{K}$. Writing the Fourier decomposition

$$\alpha(u_s) = \sum_{k \in \Lambda} \lambda_k^s u_k \quad \text{with } \lambda_k^s \in \mathbb{C},$$

we conclude that

$$\omega(s) \lambda_k^s = \omega(\delta^{-1}(k)) \lambda_k^s \quad \text{for all } s, k \in \Lambda, \omega \in \widehat{K}.$$

So, for every $s \in \Lambda$ there is precisely one $k \in \Lambda$ with $\lambda_k^s \neq 0$ and this element $k \in \Lambda$ moreover satisfies $\omega(s) = \omega(\delta^{-1}(k))$ for all $\omega \in \widehat{K}$. So, $k = \delta(s)$. In particular $\delta(s) \in \Lambda$ for all $s \in \Lambda$. So, $\delta(\Lambda) \subset \Lambda$. Since we can make a similar reasoning on $\alpha^{-1}(u_s)$, $s \in \Lambda$, it follows that $\delta(\Lambda) = \Lambda$. \square

3. A non commutative cocycle superrigidity theorem

We prove the following twisted version of Popa's cocycle superrigidity theorem [26, Theorem 0.1] for Bernoulli actions of property (T) groups.

Theorem 11. *Let K be a compact group with countable subgroup $\Lambda < K$. Let Γ be a property (T) group. Put $X = K^\Gamma$ and denote by $\Lambda \curvearrowright X$ the action by diagonal translation. Put $N = L^\infty(X) \rtimes \Lambda$. Denote by $(\sigma_g)_{g \in \Gamma}$ the action of Γ on N such that σ_g is the Bernoulli shift on $L^\infty(X)$ and the identity on $L\Lambda$. Let $p \in L\Lambda$ be a non zero projection.*

- Assume that $q \in \mathcal{B}(\ell^2(\mathbb{N})) \overline{\otimes} \mathcal{L}A$ is a projection, $\gamma : \Gamma \rightarrow \mathcal{U}(q(\mathcal{B}(\ell^2(\mathbb{N})) \overline{\otimes} \mathcal{L}A)q)$ is a group homomorphism and $v \in \mathcal{B}(\mathbb{C}, \ell^2(\mathbb{N})) \overline{\otimes} N$ is a partial isometry satisfying $v^*v = p$ and $vv^* = q$. Then the formula

$$\omega_g := v^* \gamma_g \sigma_g(v)$$

defines a 1-cocycle for the action $(\sigma_g)_{g \in \Gamma}$ on pNp , i.e. a family of unitaries satisfying $\omega_{gh} = \omega_g \sigma_g(\omega_h)$ for all $g, h \in \Gamma$;

- conversely, every 1-cocycle for the action $(\sigma_g)_{g \in \Gamma}$ on pNp is of the above form with γ being uniquely determined up to unitary conjugacy in $\mathcal{B}(\ell^2(\mathbb{N})) \overline{\otimes} \mathcal{L}A$.

Note that if A is icc, we can take $q = p$, i.e. every 1-cocycle for the action $(\sigma_g)_{g \in \Gamma}$ on pNp is cohomologous with a homomorphism from Γ to $\mathcal{U}(p\mathcal{L}(A)p)$. If A is not icc, this is no longer true since q and $e_{11} \otimes p$ need not be equivalent projections in $\mathcal{B}(\ell^2(\mathbb{N})) \overline{\otimes} \mathcal{L}A$, although they are equivalent in $\mathcal{B}(\ell^2(\mathbb{N})) \overline{\otimes} N$.

Proof. It is clear that the formulas in the theorem define 1-cocycles. Conversely, let $p \in \mathcal{L}A$ be a projection and assume that the unitaries $\omega_g \in pNp$ define a 1-cocycle for the action $(\sigma_g)_{g \in \Gamma}$ of Γ on pNp . Consider the diagonal translation action $\Lambda \curvearrowright X \times K$ and put $\mathcal{N} := L^\infty(X \times K) \rtimes \Lambda$. We embed $N \subset \mathcal{N}$ by identifying the element $Fu_\lambda \in N$ with the element $(F \otimes 1)u_\lambda \in \mathcal{N}$ whenever $F \in L^\infty(X)$, $\lambda \in \Lambda$. Also $(\sigma_g)_{g \in \Gamma}$ extends naturally to a group of automorphisms of \mathcal{N} with $\sigma_g(1 \otimes F) = 1 \otimes F$ for all $F \in L^\infty(K)$. Define $P = L^\infty(K) \rtimes \Lambda$ and view P as a subalgebra of \mathcal{N} by identifying the element $Fu_\lambda \in P$ with the element $(1 \otimes F)u_\lambda \in \mathcal{N}$ whenever $F \in L^\infty(K)$, $\lambda \in \Lambda$. Note that we obtained a commuting square

$$\begin{array}{ccc} N & \subset & \mathcal{N} \\ \cup & & \cup \\ \mathcal{L}A & \subset & P. \end{array}$$

Define $\Delta : X \times K \rightarrow X \times K : \Delta(x, k) = (k \cdot x, k)$ and denote by Δ_* the corresponding automorphism of $L^\infty(X \times K)$ given by $\Delta_*(F) = F \circ \Delta^{-1}$. One checks easily that the formula

$$\begin{aligned} \Psi : L^\infty(X) \overline{\otimes} P &\rightarrow \mathcal{N} : \Psi(F \otimes Gu_\lambda) = \Delta_*(F \otimes G)u_\lambda \\ &\text{for all } F \in L^\infty(X), G \in L^\infty(K), \lambda \in \Lambda \end{aligned}$$

defines a $*$ -isomorphism satisfying $\Psi \circ (\sigma_g \otimes \text{id}) = \sigma_g \circ \Psi$ for all $g \in \Gamma$. Put $\mu_g := \Psi^{-1}(\omega_g)$. It follows that $(\mu_g)_{g \in \Gamma}$ is a 1-cocycle for the action $\Gamma \curvearrowright X$ with values in the Polish group $\mathcal{U}(pPp)$. By Popa's cocycle superrigidity theorem [26, Theorem 0.1] and directly applying Ψ again, we find a unitary $w \in \mathcal{U}(p\mathcal{N}p)$ and a group homomorphism $\rho : \Gamma \rightarrow \mathcal{U}(pPp)$ such that

$$\omega_g = w^* \rho_g \sigma_g(w) \quad \text{for all } g \in \Gamma.$$

Consider the basic construction for the inclusion $N \subset \mathcal{N}$ denoted by $\mathcal{N}_1 := \langle \mathcal{N}, e_N \rangle$. Put $T := we_N w^*$. Since $\sigma_g(w) = \rho_g^* w \omega_g$, it follows that $\sigma_g(T) = \rho_g^* T \rho_g$. Also note that $T \in L^2(\mathcal{N}_1)$, $T = pT$ and $\text{Tr}(T) = \tau(p)$.

Since we are dealing with a commuting square, we can identify the basic construction $P_1 := \langle P, e_{\mathcal{L}A} \rangle$ for the inclusion $\mathcal{L}A \subset P$ with the von Neumann subalgebra of \mathcal{N}_1 generated by P and e_N . In this way, one checks that there is a unique unitary

$$\begin{aligned} W : L^2(\mathcal{N}_1) &\rightarrow L^2(X) \otimes L^2(P_1) \quad \text{satisfying } W(ab) = a \otimes b \\ &\text{for all } a \in L^\infty(X), b \in L^2(P_1). \end{aligned}$$

The unitary W satisfies $W(\rho_g \sigma_g(\xi) \rho_g^*) = (\sigma_g \otimes \text{Ad} \rho_g)(W(\xi))$ for all $g \in \Gamma$, $\xi \in L^2(\mathcal{N}_1)$. The projection $T \in L^2(\mathcal{N}_1)$ satisfies $T = \rho_g \sigma_g(T) \rho_g^*$. Since the unitary representation $(\sigma_g)_{g \in \Gamma}$ on $L^2(X) \ominus \mathbb{C}1$ is a multiple of the regular representation, it follows that $W(T) \in 1 \otimes L^2(P_1)$. This means that $T \in P_1$ and $\rho_g T = T \rho_g$ for all $g \in \Gamma$.

So we can view T as the orthogonal projection of $L^2(P)$ onto a right $L\Lambda$ submodule of dimension $\tau(p)$ that is globally invariant under left multiplication by ρ_g , $g \in \Gamma$. Since $pT = T$, the image of T is contained in $pL^2(P)$. Take projections $q_n \in L\Lambda$ with $\sum_n \tau(q_n) = \tau(p)$ and a right $L\Lambda$ -linear isometry

$$\Theta : \bigoplus_{n \in \mathbb{N}} q_n L^2(L\Lambda) \rightarrow L^2(P)$$

onto the image of T . Denote by $w_n \in L^2(P)$ the image under Θ of q_n sitting in position n . Note that

$$w_n = p w_n, \quad E_{L\Lambda}(w_n^* w_m) = \delta_{n,m} q_n \quad \text{and} \quad T = \sum_{n \in \mathbb{N}} w_n e_{L\Lambda} w_n^*$$

where in the last formula we view T as an element of P_1 . Identifying P_1 as above with the von Neumann subalgebra of \mathcal{N}_1 generated by P and e_N , it follows that

$$T = \sum_{n \in \mathbb{N}} w_n e_N w_n^*.$$

Since $T = w e_N w^*$, we get that $e_N = \sum_n w^* w_n e_N w_n^* w$. We conclude that $w^* w_n \in L^2(N)$ for all n . So $w_n^* w_m \in L^1(N)$ for all n, m . Since also $w_n^* w_m \in L^1(P)$, we have $w_n^* w_m \in L^1(L\Lambda)$. But then,

$$\delta_{n,m} q_n = E_{L\Lambda}(w_n^* w_m) = w_n^* w_m.$$

So, the elements w_n are partial isometries in P with mutually orthogonal left supports lying under p and with right supports equal to q_n . Since $\sum_n \tau(q_n) = \tau(p)$, we conclude that the formula

$$W := \sum_n e_{1,n} \otimes w_n$$

defines an element $W \in B(\ell^2(\mathbb{N}), \mathbb{C}) \overline{\otimes} P$ satisfying $WW^* = p$, $W^*W = q$ where $q \in (L\Lambda)^\infty = B(\ell^2(\mathbb{N})) \overline{\otimes} L\Lambda$ is the projection given by $q := \sum_n e_{nn} \otimes q_n$. We also have that $v := W^*w$ belongs to $B(\mathbb{C}1, \ell^2(\mathbb{N})) \overline{\otimes} N$ and satisfies $v^*v = p$, $vv^* = q$.

Finally, let $\gamma : \Gamma \rightarrow \mathcal{U}(q(L\Lambda)^\infty q)$ be the unique group homomorphism satisfying

$$\Theta(\gamma_g \xi) = \rho_g \Theta(\xi) \quad \text{for all } g \in \Gamma, \xi \in \bigoplus_n q_n L^2(L\Lambda).$$

By construction $W\gamma_g = \rho_g W$ for all $g \in \Gamma$. Since $\omega_g = w^* \rho_g \sigma_g(w)$ and since W is invariant under $(\sigma_g)_{g \in \Gamma}$, we conclude that $\omega_g = v^* \gamma_g \sigma_g(v)$.

We finally prove that γ is unique up to unitary conjugacy. So assume that we also have q_1, v_1 and $\varphi : \Gamma \rightarrow \mathcal{U}(q_1(L\Lambda)^\infty q_1)$ satisfying $\omega_g = v_1^* \varphi_g \sigma_g(v_1)$. It follows that the element $v_1 v^* \in q_1 N^\infty q$ satisfies $\sigma_g(v_1 v^*) = \varphi_g^* v_1 v^* \gamma_g$ for all $g \in \Gamma$. As above this forces $v_1 v^*$ to belong to $q_1(L\Lambda)^\infty q$, providing the required unitary conjugacy between γ and φ . \square

4. The classification of Cartan subalgebras up to automorphisms can be complete analytic; proof of Theorem 2

Let M be a II_1 factor with separable predual. By [7, Theorem 3 and its corollaries] the set of von Neumann subalgebras $A \subset M$ carries the standard Effros–Borel structure. In this II_1 setting it can be described as the smallest σ -algebra such that for all $x, y \in M$ the map

$$A \mapsto \tau(E_A(x)y) \quad (5)$$

is measurable. Here τ denotes the unique tracial state on M and E_A is the unique trace preserving conditional expectation of M onto A .

Proposition 12. *Let M be a II_1 factor with separable predual. Equip the set of its von Neumann subalgebras with the Effros–Borel structure, as explained above.*

- *The set $\text{Cartan}(M)$ of Cartan subalgebras of M is a Borel set and as such a standard Borel space.*
- *The equivalence relation of “being unitarily conjugate” is a Borel equivalence relation on $\text{Cartan}(M)$.*
- *The equivalence relation of “being conjugate by an automorphism of M ” is an analytic equivalence relation on $\text{Cartan}(M)$.*

Proof. Denote by $\text{vNalg}(M)$ the standard Borel space of von Neumann subalgebras of M . By [7, Theorem 3 and its corollaries] the map $A \mapsto A' \cap M$ is a Borel map on $\text{vNalg}(M)$. Hence the set

$$\text{masa}(M) = \{A \in \text{vNalg}(M) \mid A' \cap M = A\}$$

of maximal abelian subalgebras of M is a Borel subset of $\text{vNalg}(M)$.

Consider the Polish group $\mathcal{U}(M)$ of unitaries in M equipped with the $\|\cdot\|_2$ distance. Denote by $\text{subgr}(\mathcal{U}(M))$ the set of closed subgroups of $\mathcal{U}(M)$ equipped with the Effros–Borel structure. This Borel structure can be described as the smallest σ -algebra such that for all $u \in \mathcal{U}(M)$ the map

$$\text{subgr}(\mathcal{U}(M)) \rightarrow \mathbb{R} : \mathcal{G} \mapsto d(u, \mathcal{G})$$

is measurable. Here and in what follows, $d(u, \mathcal{G})$ denotes the $\|\cdot\|_2$ distance of u to \mathcal{G} . We claim that the map

$$\text{masa}(M) \rightarrow \text{subgr}(\mathcal{U}(M)) : A \mapsto \mathcal{N}_M(A) := \{u \in \mathcal{U}(M) \mid uAu^* = A\}$$

is Borel. So we have to prove that for all $u \in \mathcal{U}(M)$, the map $A \mapsto d(u, \mathcal{N}_M(A))$ is Borel. But [30, Theorem 6.2 and Lemma 6.3] provides us with the following inequalities for every maximal abelian subalgebra $A \subset M$ and all $u \in \mathcal{U}(M)$.

$$\frac{1}{4} \|E_A - E_{uAu^*}\|_{\infty,2} \leq d(u, \mathcal{N}_M(A)) \leq 90 \|E_A - E_{uAu^*}\|_{\infty,2}.$$

So in order to prove that $A \mapsto \mathcal{N}_M(A)$ is Borel, it suffices to prove that for all $u \in \mathcal{U}(M)$, the map $A \mapsto \|E_A - E_{uAu^*}\|_{\infty,2}$ is Borel. Choosing sequences (x_n) and (y_n) in M such that (x_n) is dense in the unit ball of M while (y_n) is dense in the unit ball of $L^2(M)$, we have that

$$\|E_A - E_{uAu^*}\|_{\infty,2} = \sup_{n,m} |\tau(E_A(x_n)y_m) - \tau(E_A(u^*x_nu)y_m)|.$$

Since the maps in (5) are Borel, also $A \mapsto \|E_A - E_{uAu^*}\|_{\infty,2}$ is Borel. This proves the claim that $A \mapsto \mathcal{N}_M(A)$ is a Borel map from $\text{masa}(M)$ to $\text{subgr}(\mathcal{U}(M))$.

Repeating the proof of [7, Theorem 3] the map $\text{subgr}(\mathcal{U}(M)) \rightarrow \text{vNalg}(M) : \mathcal{G} \mapsto \mathcal{G}''$ is Borel. Together with the previous paragraph it follows that the map $\text{masa}(M) \rightarrow \text{vNalg}(M) : A \mapsto \mathcal{N}_M(A)''$ is Borel. In particular,

$$\text{Cartan}(M) = \{A \in \text{masa}(M) \mid \mathcal{N}_M(A)'' = M\}$$

is a Borel subset of $\text{masa}(M)$.

Consider the equivalence relation of “being unitarily conjugate” on the standard Borel space $\text{Cartan}(M)$. Let $(u_n)_{n \in \mathbb{N}}$ be a $\|\cdot\|_2$ dense sequence in $\mathcal{U}(M)$. We claim that $A, B \in \text{Cartan}(M)$ are unitarily conjugate if and only if

$$\inf_n \|E_{u_n A u_n^*} - E_B\|_{\infty,2} < \frac{1}{2}. \quad (6)$$

Once this claim is proven, we observe as above that for every $n \in \mathbb{N}$ the map $(A, B) \mapsto \|E_{u_n A u_n^*} - E_B\|_{\infty,2}$ is Borel. It then follows that the unitarily conjugate Cartan subalgebras (A, B) form a Borel subset of $\text{Cartan}(M) \times \text{Cartan}(M)$.

It remains to prove the claim. If A and B are unitarily conjugate, i.e. $uAu^* = B$, it suffices to take n such that $\|u - u_n\|_2$ is small enough and (6) follows. Conversely, if (6) holds, take n such that $\|E_{u_n A u_n^*} - E_B\|_{\infty,2} < 1/2$. In particular, for every $v \in \mathcal{U}(u_n A u_n^*)$ we have that $\|E_B(v)\|_2 > 1/2$. Popa’s conjugacy criterion [24, Theorem A.1] implies that $u_n A u_n^*$ and B are unitarily conjugate. Hence also A and B are unitarily conjugate.

Finally, repeating the proof of [8, Lemma 2.1] the map $\text{Aut}(M) \times \text{Cartan}(M) \rightarrow \text{Cartan}(M) : (\alpha, A) \mapsto \alpha(A)$ is seen to be Borel. Since the equivalence relation of “being conjugate by an automorphism of M ” is precisely the orbit equivalence relation of this Borel action, it is an analytic equivalence relation. \square

In [6] it was shown that the isomorphism relation for torsion free abelian groups is complete analytic. This is done through the construction of a Borel reduction from the complete analytic problem of checking whether a tree has an infinite path. In this context a *tree* is a set of finite sequences of natural numbers which is closed under taking initial segments. An *infinite path* in a tree T is an infinite sequence of natural numbers $x_1 x_2 \dots$ such that $x_1 \dots x_n \in T$ for every $n \in \mathbb{N}$. A construction by Hjorth [12] associates to every tree T a subgroup $G(T)$ of the countably infinite direct sum $\Lambda := \mathbb{Q}^{(\infty)}$ such that the isomorphism class of the torsion free abelian group $G(T)$ remembers whether or not T admits an infinite path. We recall this construction in the following lemma and provide a second countable compactification $\Lambda \subset K$ such that $\overline{G(T)} \cap \Lambda = G(T)$ for every tree T .

Recall that $\mathbb{Q}/\mathbb{Z} = \bigoplus_{p \in \mathcal{P}} \mathbb{Z}(p^\infty)$ where \mathcal{P} denotes the set of prime numbers and $\mathbb{Z}(p^\infty) = \mathbb{Z}[p^{-1}]/\mathbb{Z}$. Denote by ρ_p the corresponding homomorphism $\rho_p : \mathbb{Q} \rightarrow \mathbb{Z}(p^\infty)$ uniquely characterized by $\rho_p(p^{-k}) = p^{-k} + \mathbb{Z}$ and $\rho_p(q^{-k}) = 0$ for all primes $q \neq p$ and all $k \in \mathbb{N}$.

Denote by $\mathbb{N}^{<\mathbb{N}}$ the set of finite sequences of natural numbers. For every $\sigma = (x_1 \dots x_n)$ denote by $|\sigma| = n$ the length of the sequence. Observe that the empty sequence $()$ belongs to $\mathbb{N}^{<\mathbb{N}}$ and that it is the unique element of length 0. Recall that a tree T is a subset $T \subset \mathbb{N}^{<\mathbb{N}}$ that is closed under taking initial segments: if $n \geq 1$ and $(x_1 \dots x_n) \in T$, then $(x_1 \dots x_k) \in T$ for all $k \leq n$. Whenever $\sigma \in \mathbb{N}^{<\mathbb{N}}$ with $|\sigma| \geq 1$ we denote by σ^- the sequence obtained by removing the last element of σ . Whenever $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $b \in \mathbb{N}$ we denote by σb the sequence obtained by appending b at the end of σ .

Lemma 13. Choose for every prime number p a second countable compactification L_p of $\mathbb{Z}(p^\infty)$. Choose a second countable compactification $K_{\mathbb{Q}}$ of \mathbb{Q} such that the homomorphisms $\rho_p : \mathbb{Q} \rightarrow L_p$ extend continuously to $K_{\mathbb{Q}}$ for all $p \in \mathcal{P}$.

Consider the countable group Λ given as the direct sum $\Lambda := \mathbb{Q}^{(\mathbb{N}^{<\mathbb{N}})}$. For every $\sigma \in \mathbb{N}^{<\mathbb{N}}$ define the homomorphisms

$$\pi_\sigma : \Lambda \rightarrow \mathbb{Q} : x \mapsto x_\sigma \quad \text{and} \quad \gamma_\sigma : \Lambda \rightarrow \mathbb{Q} : x \mapsto x_\sigma + \sum_{b \in \mathbb{N}} x_{\sigma b}.$$

Define K as the smallest compactification of Λ such that the homomorphisms $\pi_\sigma : \Lambda \rightarrow K_{\mathbb{Q}}$ and $\gamma_\sigma : \Lambda \rightarrow K_{\mathbb{Q}}$ extend continuously to K for all $\sigma \in \mathbb{N}^{<\mathbb{N}}$.

For every $\sigma \in \mathbb{N}^{<\mathbb{N}}$ we denote by δ_σ the obvious basis element of $\Lambda = \mathbb{Q}^{(\mathbb{N}^{<\mathbb{N}})}$. Let $p_0 < p_1 < \dots$ be an enumeration of the prime numbers. Following [12] we define for every tree T the subgroup $G(T)$ of Λ generated by

$$\{ p_{2|\sigma|}^{-k} \delta_\sigma \mid \sigma \in T, k \in \mathbb{N} \} \cup \{ p_{2|\sigma|-1}^{-k} (\delta_\sigma - \delta_{\sigma^-}) \mid \sigma \in T, |\sigma| > 0, k \in \mathbb{N} \}.$$

Denote by $\overline{G(T)}$ the closure of $G(T)$ inside K .

For every tree T we have $\overline{G(T)} \cap \Lambda = G(T)$.

Proof. The definition of $G(T)$ and the continuity of the homomorphisms ρ_p , π_σ and γ_σ imply the following properties.

1. For all $\sigma \in \mathbb{N}^{<\mathbb{N}}$ with $\sigma \notin T$ and for all $g \in \overline{G(T)}$, we have $\pi_\sigma(g) = 0$.
2. For all $\sigma \in T$, all prime numbers $p \notin \{p_{2|\sigma|-1}, p_{2|\sigma|}, p_{2|\sigma|+1}\}$ and all $g \in \overline{G(T)}$, we have $\rho_p(\pi_\sigma(g)) = 0$.
3. For all $\sigma \in T$ and all $g \in \overline{G(T)}$, we have $\rho_{p_{2|\sigma|+1}}(\gamma_\sigma(g)) = 0$.

For all $g \in \Lambda$ we consider the finite set $S_g = \{(p, \sigma) \in \mathcal{P} \times \mathbb{N}^{<\mathbb{N}} \mid \rho_p(\pi_\sigma(g)) \neq 0\}$. Note that $S_g = \emptyset$ if and only if $\pi_\sigma(g) \in \mathbb{Z}$ for every $\sigma \in \mathbb{N}^{<\mathbb{N}}$.

Assume that $g \in \overline{G(T)} \cap \Lambda$ with $g \notin G(T)$. It follows that $S_g \neq \emptyset$. Indeed, otherwise we get that $\pi_\sigma(g) \in \mathbb{Z}$ for all $\sigma \in \mathbb{N}^{<\mathbb{N}}$. By property 1 above, we know that $\pi_\sigma(g) = 0$ for all $\sigma \notin T$. Hence g belongs to the group generated by $\{\delta_\sigma \mid \sigma \in T\}$. In particular, $g \in G(T)$.

Assume that $\overline{G(T)} \cap \Lambda$ is strictly larger than $G(T)$. Take $g \in (\overline{G(T)} \cap \Lambda) \setminus G(T)$ minimizing $|S_g|$. By the previous paragraph $S_g \neq \emptyset$. So we can choose $(p, \sigma) \in S_g$ maximizing $|\sigma|$. By property 1 above we have $\sigma \in T$. Property 2 implies that $p \in \{p_{2|\sigma|-1}, p_{2|\sigma|}, p_{2|\sigma|+1}\}$. We prove that each of these three possibilities leads to a contradiction.

First assume that $|\sigma| > 0$ and $p = p_{2|\sigma|-1}$. By property 3 above we have $\rho_p(\gamma_{\sigma^-}(g)) = 0$. Using property 1 it follows that

$$\rho_p(g_{\sigma^-}) + \sum_{b \in \mathbb{N}, \sigma^- b \in T} \rho_p(g_{\sigma^- b}) = 0.$$

So we can take h in the group generated by $\{p^{-k}(\delta_{\sigma^- b} - \delta_{\sigma^-}) \mid b \in \mathbb{N}, \sigma^- b \in T, k \in \mathbb{N}\}$ such that $\rho_p(h_{\sigma^-}) = \rho_p(g_{\sigma^-})$ and $\rho_p(h_{\sigma^- b}) = \rho_p(g_{\sigma^- b})$ for all $b \in \mathbb{N}$. Note that $h \in G(T)$ and hence $g - h \in (\overline{G(T)} \cap \Lambda) \setminus G(T)$. By construction

$$S_{g-h} = S_g \setminus (\{(p, \sigma^-)\} \cup \{(p, \sigma^- b) \mid b \in \mathbb{N}\}).$$

So $|S_{g-h}| < |S_g|$ contradicting the minimality of $|S_g|$.

Next assume that $p = p_{2|\sigma|}$. Choose h in the group generated by $\{p^{-k} \delta_\sigma \mid k \in \mathbb{N}\}$ such that $\rho_p(h_\sigma) = \rho_p(g_\sigma)$. Note that $h \in G(T)$ and hence $g - h \in (\overline{G(T)} \cap \Lambda) \setminus G(T)$. Again $|S_{g-h}| < |S_g|$ contradicting the minimality of $|S_g|$.

Finally assume that $p = p_{2|\sigma|+1}$. By property 3 above we have that $\rho_p(\gamma_\sigma(g)) = 0$. Hence

$$\rho_p(\pi_\sigma(g)) + \sum_{b \in \mathbb{N}} \rho_p(\pi_{\sigma b}(g)) = 0.$$

As $\rho_p(\pi_\sigma(g)) \neq 0$, it follows that there exists a $b \in \mathbb{N}$ such that $\rho_p(\pi_{\sigma b}(g)) \neq 0$. This contradicts the maximality of $|\sigma|$.

In all three cases we obtained a contradiction. This ends the proof of the lemma. \square

We are now ready to prove [Theorem 2](#).

Proof of Theorem 2. The first part of [Theorem 2](#) follows immediately from [Theorem 1](#) and the following observations.

- The subgroups $K(\mathcal{P}_1)$ and $K(\mathcal{P}_2)$ are commensurate if and only if the symmetric difference $\mathcal{P}_1 \Delta \mathcal{P}_2$ is finite.
- Considering the order of the elements in K , it follows that every automorphism $\delta \in \text{Aut}(K)$ globally preserves each direct summand $\mathbb{Z}/p\mathbb{Z}$. In particular $\delta(\Lambda) = \Lambda$ and $\delta(\Lambda K(\mathcal{P}_1)) = \Lambda K(\mathcal{P}_1)$ for every subset $\mathcal{P}_1 \subset \mathcal{P}$.

It remains to prove the second part of [Theorem 2](#). Put $\Lambda = \mathbb{Q}^{(\mathbb{N}^{<\mathbb{N}})}$ and denote by K the compactification of Λ given by [Lemma 13](#). Consider the II_1 factor $M := L^\infty(K^\Gamma) \rtimes (\Gamma \times \Lambda)$. For every tree $T \subset \mathbb{N}^{<\mathbb{N}}$ denote by $G(T) \subset \Lambda$ the subgroup defined in [Lemma 13](#). Denote by $\mathcal{C}(T) := \mathcal{C}(\overline{G(T)})$ the associated Cartan subalgebra.

We denote the set of trees by Trees . Viewing Trees as a subset of the power set of $\mathbb{N}^{<\mathbb{N}}$, it is a standard Borel space. It is straightforward though tedious to check that the map

$$\text{Trees} \rightarrow \text{Cartan}(M) : T \mapsto \mathcal{C}(T)$$

is Borel.

A classical result (see e.g. [17, Theorem 27.1]) says that the subset of Trees consisting of the trees that admit an infinite path, is complete analytic. In combination with [6, Lemma 3.1] we get a complete analytic set $X \subset \mathbb{R}$ and Borel functions $F_1, F_2 : \mathbb{R} \rightarrow \text{Trees}$ such that the following statements hold.

- If $x \in X$, then $F_1(x) \cong F_2(x)$.
- If $x \notin X$, then $F_1(x)$ has an infinite path and $F_2(x)$ has no infinite path.

Moreover we may assume that for all $x \in \mathbb{R}$ the trees $F_1(x)$ and $F_2(x)$ consist of sequences of even numbers only (for instance by “multiplying by 2” the original F_1, F_2).

Define the Borel map

$$h : \mathbb{R} \rightarrow \text{Cartan}(M) \times \text{Cartan}(M) : h(x) = (\mathcal{C}(F_1(x)), \mathcal{C}(F_2(x))).$$

Denote by $\mathcal{R} \subset \text{Cartan}(M) \times \text{Cartan}(M)$ the equivalence relation given by the Cartan subalgebras (A, B) that are conjugate by an automorphism of M . In [Proposition 12](#) we have seen that \mathcal{R} is an analytic subset of $\text{Cartan}(M) \times \text{Cartan}(M)$. We prove below that $h^{-1}(\mathcal{R}) = X$. Since X is complete analytic, it then follows that \mathcal{R} is complete analytic.

The formula $h^{-1}(\mathcal{R}) = X$ follows, once we have shown the following two statements.

- If $S, T \subset \mathbb{N}^{<\mathbb{N}}$ are trees that consist of sequences of even numbers and if $S \cong T$, then the Cartan subalgebras $\mathcal{C}(S)$ and $\mathcal{C}(T)$ are conjugate by an automorphism of M .
- If $S, T \subset \mathbb{N}^{<\mathbb{N}}$ are trees such that S admits an infinite path and such that the Cartan subalgebras $\mathcal{C}(S)$ and $\mathcal{C}(T)$ are conjugate by an automorphism of M , then T also admits an infinite path.

To prove the first statement, let $\psi : S \rightarrow T$ be an isomorphism of trees, i.e. a bijection that preserves initial segments. Since S and T only contain sequences of even numbers, ψ can be extended, by induction on the length $|\sigma|$, to an isomorphism $\psi : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}^{<\mathbb{N}}$. Denote by $\alpha \in \text{Aut}(A)$ the corresponding automorphism given by $\alpha(\delta_\sigma) = \delta_{\psi(\sigma)}$ for all $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Using the notation of [Lemma 13](#) we have that $\pi_{\psi(\sigma)} \circ \alpha = \pi_\sigma$ and $\gamma_{\psi(\sigma)} \circ \alpha = \gamma_\sigma$ for all $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Hence α extends to a continuous automorphism of K . By construction $\alpha(G(S)) = G(T)$ and so $\alpha(\overline{G(S)}) = \overline{G(T)}$. It follows from [Theorem 1](#) that the Cartan subalgebras $\mathcal{C}(S)$ and $\mathcal{C}(T)$ are conjugate by an automorphism of M .

To prove the second statement, assume that S admits an infinite path and that the Cartan subalgebras $\mathcal{C}(S)$ and $\mathcal{C}(T)$ are conjugate by an automorphism of M . By [Theorem 1](#) we get a continuous automorphism α of K such that $\alpha(A) = A$ and $\alpha(\overline{AG(S)}) = \overline{AG(T)}$. It follows that $\alpha(\overline{G(S)})$ is commensurate with $\overline{G(T)}$ and that $\alpha(G(S) \cap A)$ is commensurate with $\overline{G(T)} \cap A$. We conclude that $\alpha(G(S))$ and $G(T)$ are commensurate.

Let $(x_1 x_2 \dots)$ be an infinite path in S . Define $g_0 = \delta_0$ and $g_i = \delta_{(x_1 \dots x_i)}$. By construction, g_i is a sequence of elements in $G(S)$ such that for all i we have that g_i is divisible by all the powers of p_{2i} and that $g_{i+1} - g_i$ is divisible by all the powers of p_{2i+1} . Denote by $N < \infty$ the index of $\alpha(G(S)) \cap G(T)$ in $G(T)$. Then $N\alpha(g_0), N\alpha(g_1), N\alpha(g_2), \dots$ is a sequence of elements in $G(T)$ satisfying the same divisibility conditions. It follows from [\[6, Lemma 2.2\]](#) that the tree T admits an infinite path. \square

Remark 14. [Theorem 2](#) provides examples of II_1 factors M such that the equivalence relation given by “being conjugate by an automorphism” on the set of Cartan subalgebras of M , is complete analytic. This is a mathematically rigorous way of saying that the decision problem whether two Cartan subalgebras of M are conjugate by an automorphism, is as hard as it can possibly be. The decision problem whether two Cartan subalgebras of M are unitarily conjugate is strictly easier. Indeed, Popa’s unitary conjugacy criterion in [\[24, Theorem A.1\]](#) is rather concrete and, as we have seen in [Proposition 12](#), shows that unitary conjugacy of Cartan subalgebras is always Borel. The fact that conjugacy by an automorphism can be complete analytic shows that there can never be a similar concrete criterion for conjugacy of Cartan subalgebras by an automorphism.

There are also other mathematically rigorous undecidability and non classifiability statements. It sounds plausible that there exist II_1 factors M for which the Cartan subalgebras cannot be classified by countable structures. Entirely new techniques would however be needed to construct such II_1 factors. Indeed, the Cartan subalgebras, up to any kind of conjugacy, given by the admissible subgroups associated with a pair $A \subset K$, are by definition classifiable by countable structures.

As far as we know, it is still an open problem whether or not the isomorphism relation on countable abelian groups is universal among the equivalence relations that are classifiable by countable structures. If this problem can be solved in the affirmative, it is very likely that a technique similar to [Theorem 2](#) will provide II_1 factors M for which the classification of

Cartan subalgebras up to conjugacy by an automorphism, is in the same sense universal among equivalence relations that are classifiable by countable structures.

5. Factors with many group measure space Cartan subalgebras

In this section we prove [Theorem 4](#). We actually construct a concrete group G and a concrete uncountable family $G \curvearrowright (X_{\mathcal{F}}, \mu_{\mathcal{F}})$, indexed by all possible subsets $\mathcal{F} \subset \mathbb{N}$, of free ergodic p.m.p. actions that are non stably orbit equivalent and that give rise to isomorphic crossed product Π_1 factors.

The basic idea is due to [\[29, Section 6.1\]](#) yielding the following concrete example of two group actions $H \curvearrowright (X_i, \mu_i)$, $i = 1, 2$, that are not orbit equivalent (although stably orbit equivalent) and yet have $L^\infty(X_1) \rtimes H \cong L^\infty(X_2) \rtimes H$. Let H_0 be a finite non commutative group. Define the infinite direct sum $H_1 := H_0^{(\mathbb{N})}$ with compactification $K := H_0^{\mathbb{N}}$. Consider the action $H_1 \curvearrowright K$ by left translation. Define $X_1 := K^{\text{SL}(3, \mathbb{Z})}$ and consider the action of $H := \text{SL}(3, \mathbb{Z}) \times H_1$ on X_1 where $\text{SL}(3, \mathbb{Z})$ acts by Bernoulli shift and H_1 acts diagonally. Take the first copy of H_0 in H_1 and put $X_2 := X_1/H_0$. We have a natural action of $H \cong H/H_0$ on X_2 . By construction $H \curvearrowright X_2$ is stably orbit equivalent (with compression constant $|H_0|^{-1}$) with $H \curvearrowright X_1$. Put $M = L^\infty(X_1) \rtimes H$. In [\[29, Section 6.1\]](#) it is shown that M has fundamental group \mathbb{R}_+ , while the orbit equivalence relation of $H \curvearrowright X_1$ has trivial fundamental group. Hence $L^\infty(X_1) \rtimes H \cong L^\infty(X_2) \rtimes H$ although $H \curvearrowright X_1$ and $H \curvearrowright X_2$ are not orbit equivalent.

We call two free ergodic p.m.p. actions W^* -equivalent if their associated group measure space Π_1 factors are isomorphic.

We now perform the following general construction. We start with two free ergodic p.m.p. actions $H \curvearrowright X_i$, $i = 1, 2$, that are not orbit equivalent but that are W^* -equivalent, and we construct a group G with uncountably many non stably orbit equivalent but W^* -equivalent actions.

Assume that Γ is a countable group and that $\Gamma_n < \Gamma$ is a sequence of subgroups of infinite index. Define the disjoint union $I := \bigsqcup_{n \in \mathbb{N}} \Gamma/\Gamma_n$ with the natural action $\Gamma \curvearrowright I$. To avoid trivialities, assume that $\Gamma \curvearrowright I$ is faithful, i.e. the intersection of all $g\Gamma_n g^{-1}$, $g \in \Gamma$, $n \in \mathbb{N}$, reduces to $\{e\}$. Define the generalized wreath product group

$$G := H \wr_I \Gamma = H^{(I)} \rtimes \Gamma.$$

Whenever $\mathcal{F} \subset \mathbb{N}$, define the space

$$X_{\mathcal{F}} := \prod_{n \in \mathcal{F}} X_1^{\Gamma/\Gamma_n} \times \prod_{n \notin \mathcal{F}} X_2^{\Gamma/\Gamma_n}.$$

Taking an infinite product of copies of the action $H \curvearrowright X_i$, we get an action $H^{(\Gamma/\Gamma_n)} \curvearrowright X_i^{\Gamma/\Gamma_n}$. Taking a further infinite product of these actions over $n \in \mathcal{F}$ (with $i = 1$) and $n \in \mathbb{N} - \mathcal{F}$ (with $i = 2$), we obtain an action of $H^{(I)}$ on $X_{\mathcal{F}}$. This action is compatible with the Bernoulli shift of Γ on $X_{\mathcal{F}}$. So we have constructed a free ergodic p.m.p. action

$$G \overset{\sigma_{\mathcal{F}}}{\curvearrowright} X_{\mathcal{F}}.$$

Denoting $M := L^\infty(X_1) \rtimes H \cong L^\infty(X_2) \rtimes H$, it is easily checked that

$$L^\infty(X_{\mathcal{F}}) \rtimes G \cong \left(\overline{\otimes}_{i \in I} M \right) \rtimes \Gamma \quad \text{where } \Gamma \text{ acts on } \overline{\otimes}_{i \in I} M \text{ by shifting the indices.}$$

Hence all the actions $G \curvearrowright X_{\mathcal{F}}$ are W^* -equivalent.

Theorem 4 is an immediate consequence of the following more concrete result, showing that under the right assumptions, the actions $\sigma_{\mathcal{F}}$ are mutually non stably orbit equivalent. We did not try to formulate the most general conditions under which such a result holds, but provide a concrete example of $\Gamma_n < \Gamma$ for which the proof is rather straightforward.

Proposition 15. *Consider the group actions $H \curvearrowright X_i, i = 1, 2$, defined above: $H \curvearrowright X_i$ are W^* -equivalent, non orbit equivalent and H is a residually finite group. Define the field*

$$K := \mathbb{Q}(\sqrt{n} \mid n \in \mathbb{N})$$

by adding all square roots of positive integers to \mathbb{Q} . For all $n \geq 2$ we put $K_n := \mathbb{Q}(\sqrt{n})$. Define the countable group $\Gamma := \text{PSL}(4, K)$ and the sequence of subgroups

$$\Gamma_n := \{aA \mid a \in K, A \text{ is an upper triangular matrix with entries in } K_n, \det(aA) = 1\}.$$

As above we put $I = \bigsqcup_{n \geq 2} \Gamma/\Gamma_n$ and $G = H \wr_I \Gamma$. For every subset $\mathcal{F} \subset \mathbb{N} - \{0, 1\}$, we are given the action $\sigma_{\mathcal{F}}$ of G on $X_{\mathcal{F}}$.

The actions $\sigma_{\mathcal{F}}$ and $\sigma_{\mathcal{F}'}$ are stably orbit equivalent iff $\mathcal{F} = \mathcal{F}'$.

We freely make use of the following properties of $\Gamma_n < \Gamma$.

- Γ is a simple group. In particular, Γ has no non trivial amenable quotients.
- The subgroups Γ_n are amenable.
- If $n \geq 2$ and $g \in \Gamma - \Gamma_n$, then $g\Gamma_n g^{-1} \cap \Gamma_n$ has infinite index in Γ_n , i.e. the quasi-normalizer of Γ_n inside Γ equals Γ_n . Equivalently: Γ_n acts with infinite orbits on $\Gamma/\Gamma_n - \{e\Gamma_n\}$.
- If $n \neq m$ and $\delta \in \text{Aut}\Gamma$, then $\delta(\Gamma_n) \cap \Gamma_m$ has infinite index in Γ_m . This follows because the automorphism group of Γ is generated by the inner automorphisms, the automorphisms given by automorphisms of the field K and the automorphism transpose-inverse, and because every field automorphism of K globally preserves every K_n .
- The previous two items imply that $(\text{Stab } i) \cdot j$ is infinite for all $i \neq j$.

We also use the following lemma. It is a very special case of general von Neumann algebra results (see [25, Theorem 4.1], [14, Theorem 0.1] and [16, Theorem 4.2]). Our lemma is also closely related to [5, Theorem 6.7]. We include a short and elementary proof for the convenience of the reader.

Lemma 16. *Let $\Gamma \curvearrowright I$ be any action of a countable group Γ by permutations of a countable set I . Let H be any countable group and put $G := H \wr_I \Gamma$. Assume that $\Lambda < G$ is a subgroup with the relative property (T). Then either a finite index subgroup of Λ is contained in $H \wr_I \text{Stab } i$ for some $i \in I$, or there exists a $g \in G$ such that $g\Lambda g^{-1} \subset \Gamma$.*

Proof. Define the unitary representation $\pi : G \rightarrow \mathcal{U}(\ell^2(H \times I))$ given by

$$(\pi(h)\xi)(k, i) = \xi(h_i^{-1}k, i) \quad \text{if } h \in H^{(I)}, \text{ and } (\pi(\gamma)\xi)(k, i) = \xi(k, \gamma^{-1} \cdot i) \text{ if } \gamma \in \Gamma.$$

Define the function

$$\eta : H \times I \rightarrow \mathbb{C} : \eta(k, i) = \begin{cases} 1 & \text{if } k = e, \\ 0 & \text{if } k \neq e. \end{cases}$$

The formal expression $b(g) := \pi(g)\eta - \eta$ provides a well defined 1-cocycle of G with values in $\ell^2(H \times I)$. Since $\Lambda < G$ has the relative property (T), the 1-cocycle b is inner on Λ . So we find $\xi \in \ell^2(H \times I)$ satisfying

$$b(\lambda) = \pi(\lambda)\xi - \xi \quad \text{for all } \lambda \in \Lambda.$$

Denote by $\delta_h : H \rightarrow \mathbb{C}$ the function that equals 1 in h and 0 elsewhere. We get that

$$\xi(k, i) = \xi(h_i^{-1}k, \gamma^{-1} \cdot i) - \delta_e(h_i^{-1}k) + \delta_e(k) \quad \text{for all } (h, \gamma) \in \Lambda, k \in H, i \in I.$$

For every $i \in I$, we write $\xi_i \in \ell^2(H)$ given by $\xi_i(k) = \xi(k, i)$. We also denote by $(\lambda_h)_{h \in H}$ the left regular representation of H on $\ell^2(H)$. The above formula becomes

$$\xi_i = \lambda_{h_i} \xi_{\gamma^{-1} \cdot i} - \delta_{h_i} + \delta_e \quad \text{for all } (h, \gamma) \in \Lambda, i \in I. \quad (7)$$

Since $\xi \in \ell^2(H \times I)$, it follows that $\|\xi_i\| \rightarrow 0$ as $i \rightarrow \infty$. Assume that finite index subgroups of Λ are never contained in $H \wr_I \text{Stab } i$. We then find a sequence $(h_n, \gamma_n) \in \Lambda$ such that for all $i \in I$ we have that $\gamma_n^{-1} \cdot i \rightarrow \infty$ as $n \rightarrow \infty$. Denote by $V \subset \ell^2(H)$ the subset of vectors $V := \{\delta_e - \delta_h \mid h \in H\}$. Note that $V \subset \ell^2(\mathbb{N})$ is closed. We apply (7) to $(h_n, \gamma_n) \in \Lambda$ and let $n \rightarrow \infty$. Since $\|\xi_{\gamma_n^{-1} \cdot i}\| \rightarrow 0$, we conclude that $\xi_i \in V$ for every $i \in I$. Denote by $b_i \in H$ the unique element such that $\xi_i = \delta_e - \delta_{b_i}$. Since $\|\xi_i\| \rightarrow 0$ as $i \rightarrow \infty$, we conclude that $b_i = e$ for all but finitely many $i \in I$. So $b := (b_i)_{i \in I}$ is a well defined element of $H^{(I)}$. Formula (7) becomes

$$b_i^{-1} h_i b_{\gamma^{-1} \cdot i} = e \quad \text{for all } (h, \gamma) \in \Lambda, i \in I,$$

which precisely means that $b^{-1} \Lambda b \subset \Gamma$. \square

We can now prove [Proposition 15](#). We use the terminology and conventions concerning stable orbit equivalences that we introduced after [Remark 7](#).

Proof. Let $\mathcal{F}, \mathcal{F}' \subset \mathbb{N} - \{0, 1\}$ be subsets. Assume that $\Delta : X_{\mathcal{F}} \rightarrow X_{\mathcal{F}'}$ is a stable orbit equivalence with corresponding Zimmer 1-cocycle

$$\omega : G \times X_{\mathcal{F}} \rightarrow G, \quad \Delta(g \cdot x) = \omega(g, x) \cdot \Delta(x) \quad \text{for all } g \in G \text{ and a.e. } x \in X_{\mathcal{F}}.$$

Since both X_1 and X_2 are standard non atomic probability spaces, the restriction of the action $G \curvearrowright^{\sigma_{\mathcal{F}}} X_{\mathcal{F}}$ to Γ is the generalized Bernoulli action $\Gamma \curvearrowright [0, 1]^I$. Note that its further restriction to the property (T) group $\Lambda := \text{PSL}(4, \mathbb{Z}) < \Gamma$ is weakly mixing, since $\Lambda \cdot i$ is infinite for every $i \in I$. By Popa's cocycle superrigidity theorem [26, Theorem 0.1] we can replace Δ by a similar stable orbit equivalence and assume that $\omega(\lambda, x) = \delta(\lambda)$ for all $\lambda \in \Lambda$ and a.e. $x \in X_{\mathcal{F}}$, where $\delta : \Lambda \rightarrow G$ is a group homomorphism.

For all $i, j \in \{1, 2, 3, 4\}$ denote by e_{ij} the matrix having 1 in position ij and 0 elsewhere. Define, for $i \neq j$ and $k \in K$, the elementary matrix $E_{ij}(k) = 1 + ke_{ij}$. The elementary subgroups $E_{ij}(K)$ generate Γ . Given distinct $i, j \in \{1, 2, 3, 4\}$, denote by i', j' the remaining elements of $\{1, 2, 3, 4\}$ and denote by Λ_{ij} the subgroup of Λ generated by $E_{i'j'}(\mathbb{Z})$ and $E_{jj'}(\mathbb{Z})$. Note that Λ_{ij} is non amenable and that Λ_{ij} commutes with $E_{ij}(K)$. Since $\Lambda_{ij} \curvearrowright X_{\mathcal{F}}$ is weakly mixing and ω is a homomorphism on Λ_{ij} , [26, Proposition 3.6] implies that ω is a homomorphism on $E_{ij}(K)$. Since this holds for all $i \neq j$, we conclude that ω is a homomorphism on the whole of Γ . Denote this homomorphism by $\delta : \Gamma \rightarrow G$.

Since Γ is simple, δ is injective. Recall that $G = H \wr_I \Gamma$. We shall prove that there exists a $g \in G$ such that $g\delta(\Gamma)g^{-1} \subset \Gamma$. Denote by Γ_{12} the copy of $\text{PSL}(2, K)$ inside Γ generated by $E_{12}(K)$ and $E_{21}(K)$. We first claim that there is no $i \in I$ such that $\delta(\Gamma_{12}) \subset H \wr_I \text{Stab } i$. Assume the contrary. Denote by $\pi : G \rightarrow \Gamma$ the natural quotient map. Since Γ_{12} is simple and non amenable, $\pi(\delta(\Gamma_{12})) \subset \text{Stab } i$ must be the trivial group. So, $\delta(\Gamma_{12}) \subset H^{(I)}$. Since H is residually finite and δ is injective, this is absurd. This proves the claim. Since Γ_{12} has no non

trivial finite index subgroups, we only retain the existence of $\gamma_n \in \Gamma_{12}$ such that for all $i \in I$ we have that $\pi(\delta(\gamma_n)) \cdot i \rightarrow \infty$ as $n \rightarrow \infty$.

We next claim that there is no finite index subgroup $\Lambda_0 < \Lambda$ and $i \in I$ with $\delta(\Lambda_0) \subset H \wr_I \text{Stab } i$. To prove this claim, assume the contrary. Since $\text{Stab } i$ is amenable and Λ_0 has property (T), it follows that $\pi(\delta(\Lambda_0))$ is finite. So making Λ_0 smaller but still of finite index, we get that $\delta(\Lambda_0) \subset H^{(I)}$. Since Λ_0 is finitely generated, we can take $I_1 \subset I$ finite such that $\delta(\Lambda_0) \subset H^{(I_1)}$. Recall the subgroup $\Lambda_{12} < \Lambda$ that commutes with the subgroup $\Gamma_{12} < \Gamma$ that we considered in the previous paragraph. Also recall that we have found elements $\gamma_n \in \Gamma_{12}$ such that for all $i \in I$ we have that $\pi(\delta(\gamma_n)) \cdot i \rightarrow \infty$ as $n \rightarrow \infty$. Since $\delta(\Lambda_0 \cap \Lambda_{12}) \subset H^{(I_1)}$, we have

$$\delta(\gamma_n) \delta(\Lambda_0 \cap \Lambda_{12}) \delta(\gamma_n)^{-1} \in H^{(\pi(\delta(\gamma_n)) \cdot I_1)}.$$

Since γ_n and Λ_{12} commute, the left hand side equals $\delta(\Lambda_0 \cap \Lambda_{12})$. Letting $n \rightarrow \infty$, the sets $\pi(\delta(\gamma_n)) \cdot I_1$ and I_1 are eventually disjoint. We conclude that $\delta(\Lambda_0 \cap \Lambda_{12}) = \{e\}$. Since $\Lambda_0 < \Lambda$ has finite index, the group $\Lambda_0 \cap \Lambda_{12}$ is infinite, contradicting the injectivity of δ .

Combining the claim that we proved in the previous paragraph with Lemma 16, we can conjugate δ , replace Δ by a similar stable orbit equivalence and assume that $\delta(\Lambda) \subset \Gamma$. For all i, j we know that $\delta(\Lambda_{ij})$ is a non amenable subgroup of Γ . Therefore $\delta(\Lambda_{ij}) \cdot k$ is infinite for all $k \in I$ and the centralizer of $\delta(\Lambda_{ij})$ inside G is a subgroup of Γ . Therefore $\delta(E_{ij}(K)) \subset \Gamma$ for all i, j . Hence, $\delta(\Gamma) \subset \Gamma$.

We can perform a similar reasoning on the generalized inverse stable orbit equivalence $\bar{\Delta} : X_{\mathcal{F}'} \rightarrow X_{\mathcal{F}}$ and may assume that $\bar{\Delta}(g \cdot x) = \bar{\delta}(g) \cdot \bar{\Delta}(x)$ for all $g \in \Gamma$ and a.e. $x \in X_{\mathcal{F}'}$, where $\bar{\delta} : \Gamma \rightarrow \Gamma$ is an injective group homomorphism. By definition $\bar{\Delta}(\Delta(x)) \in G \cdot x$ for a.e. $x \in X_{\mathcal{F}}$. So we find a measurable function $\varphi : X_{\mathcal{F}} \rightarrow G$ such that $\bar{\Delta}(\Delta(x)) = \varphi(x) \cdot x$ for a.e. $x \in X_{\mathcal{F}}$. It follows that

$$\varphi(g \cdot x) = \bar{\delta}(\delta(g)) \varphi(x) g^{-1} \quad \text{for all } g \in \Gamma \text{ and a.e. } x \in X_{\mathcal{F}}.$$

So the non negligible subset $\{(x, x') \in X_{\mathcal{F}} \times X_{\mathcal{F}} \mid \varphi(x) = \varphi(x')\}$ is invariant under the diagonal Γ -action. Since $\Gamma \curvearrowright X_{\mathcal{F}}$ is weakly mixing, we conclude that φ is essentially constant. This constant value necessarily belongs to Γ and we conclude that $\bar{\delta} \circ \delta$ is an inner automorphism of Γ . A similar reasoning holds for $\delta \circ \bar{\delta}$. In particular $\Delta(\Gamma \cdot x) = \Gamma \cdot \Delta(x)$ for a.e. $x \in X_{\mathcal{F}}$. Lemma 8 implies that Δ is a measure space isomorphism of $X_{\mathcal{F}}$ onto $X_{\mathcal{F}'}$ and that Δ is a δ -conjugacy of the respective Γ -actions. In particular, the compression constant of Δ equals 1.

Denote

$$X_{\mathcal{F}}^i := \begin{cases} X_1 & \text{if } i \in \Gamma/\Gamma_n \text{ and } n \in \mathcal{F}, \\ X_2 & \text{if } i \in \Gamma/\Gamma_n \text{ and } n \notin \mathcal{F}. \end{cases}$$

By construction, $X_{\mathcal{F}} = \prod_{i \in I} X_{\mathcal{F}}^i$ and similarly for $X_{\mathcal{F}'}$.

By [32, Lemma 6.15] the δ -conjugacy Δ necessarily decomposes as a product of measure space isomorphisms. More precisely, we find a permutation η of the set I and measure space isomorphisms $\Delta_i : X_{\mathcal{F}}^i \rightarrow X_{\mathcal{F}'}^{\eta(i)}$ such that

$$\eta(g \cdot i) = \delta(g) \cdot \eta(i) \quad \text{and} \quad \Delta(x)_{\eta(i)} = \Delta_i(x_i) \quad \text{for all } g \in \Gamma \text{ and a.e. } x \in X_{\mathcal{F}}.$$

Since η preserves the orbits of $\Gamma \curvearrowright I$, it defines a permutation $\bar{\eta}$ of \mathbb{N} such that $\eta(i) \in \Gamma/\Gamma_{\bar{\eta}(n)}$ iff $i \in \Gamma/\Gamma_n$. Also Δ_i only depends on the orbit of i , yielding $\Delta_n : X_{\mathcal{F}}^i \rightarrow X_{\mathcal{F}'}^{\eta(i)}$ whenever $i \in \Gamma/\Gamma_n$.

Note that $\delta(\text{Stab } i) = \text{Stab } \eta(i)$. For all $n \neq m$ and all $\delta \in \text{Aut } \Gamma$ we have that $\delta(\Gamma_n) \neq \Gamma_m$. Hence $\bar{\eta}$ is the identity. Denote by H^i the copy of H in $H^{(I)}$ sitting in position i . Fix $n \in \mathbb{N}$ and $i \in \Gamma/\Gamma_n$. Put $j := \eta(i)$ and note that $j \in \Gamma/\Gamma_n$. Since the quasi-normalizer of Γ_n inside Γ equals Γ_n and since Γ_n acts with infinite orbits on Γ/Γ_m for all $m \neq n$, it follows that the action

$$\text{Stab } i \curvearrowright \prod_{j, j \neq i} X_{\mathcal{F}}^j$$

is weakly mixing. Since the 1-cocycle ω is a homomorphism on $\text{Stab } i$ and H^i commutes with $\text{Stab } i$, it follows from [26, Proposition 3.6] that for all $h \in H^i$, the map $x \mapsto \omega(h, x)$ only depends on the coordinate x_i and that $\omega(h, x)$ commutes with $\delta(\text{Stab } i) = \text{Stab } j$. Since $(\text{Stab } j) \cdot k$ is infinite for all $k \neq j$ and since $\text{Stab } j < \Gamma$ has a trivial centralizer in Γ , it follows that $\omega(h, x) \in H^j$. A similar reasoning holds for the inverse 1-cocycle and we conclude that $\Delta_n : X_{\mathcal{F}}^i \rightarrow X_{\mathcal{F}'}^j$ is an orbit equivalence between $H \curvearrowright X_{\mathcal{F}}^i$ and $H \curvearrowright X_{\mathcal{F}'}^j$. Since both $i, j \in \Gamma/\Gamma_n$ and since the actions $H \curvearrowright X_1$ and $H \curvearrowright X_2$ are not orbit equivalent, it follows that either n belongs to both \mathcal{F} and \mathcal{F}' or that n belongs to both $\mathbb{N} - \mathcal{F}$ and $\mathbb{N} - \mathcal{F}'$. This holds for all n and we conclude that $\mathcal{F} = \mathcal{F}'$. \square

Acknowledgments

The second author was partially supported by ERC Starting Grant VNALG-200749, Research Programme G.0639.11 of the Research Foundation – Flanders (FWO) and KU Leuven BOF research grant OT/08/032.

References

- [1] I. Chifan, J. Peterson, Some unique group-measure space decomposition results. Preprint [arXiv:1010.5194](#).
- [2] I. Chifan, T. Sinclair, On the structural theory of II_1 factors of negatively curved groups. Preprint [arXiv:1103.4299](#).
- [3] A. Connes, J. Feldman, B. Weiss, An amenable equivalence relation is generated by a single transformation, *Ergodic Theory Dynam. Systems* 1 (1981) 431–450.
- [4] A. Connes, V.F.R. Jones, A II_1 factor with two non-conjugate Cartan subalgebras, *Bull. Amer. Math. Soc.* 6 (1982) 211–212.
- [5] Y. Cornuier, Y. Stalder, A. Valette, Proper actions of wreath products and generalizations, *Trans. Amer. Math. Soc.* 364 (2012) 3159–3184.
- [6] R. Downey, A. Montalbán, The isomorphism problem for torsion-free abelian groups is analytic complete, *J. Algebra* 320 (2008) 2291–2300.
- [7] E.G. Effros, The Borel space of von Neumann algebras on a separable Hilbert space, *Pacific J. Math.* 15 (1965) 1153–1164.
- [8] E.G. Effros, Global structure in von Neumann algebras, *Trans. Amer. Math. Soc.* 121 (1966) 434–454.
- [9] J. Feldman, C.C. Moore, Ergodic equivalence relations, cohomology, and Von Neumann algebras, I, II, *Trans. Amer. Math. Soc.* 234 (1977) 289–324, 325–359.
- [10] P. Fima, S. Vaes, HNN extensions and unique group measure space decomposition of II_1 factors, *Trans. Amer. Math. Soc.* 364 (2012) 2601–2617.
- [11] D. Gaboriau, Orbit equivalence and measured group theory, in: *Proceedings of the International Congress of Mathematicians (Hyderabad, India, 2010)*, Vol. III, Hindustan Book Agency, 2010, pp. 1501–1527.
- [12] G. Hjorth, The isomorphism relation on torsion-free abelian groups, *Fund. Math.* 175 (2002) 241–257.
- [13] C. Houdayer, S. Popa, S. Vaes, A class of groups for which every action is W^* -superrigid. *Groups Geom. Dyn.* (in press) [arXiv:1010.5077](#).
- [14] A. Ioana, Rigidity results for wreath product II_1 factors, *J. Funct. Anal.* 252 (2007) 763–791.
- [15] A. Ioana, Uniqueness of the group measure space decomposition for Popa’s \mathcal{HT} factors. *Geom. Funct. Anal.* (in press) [arXiv:1104.2913](#).
- [16] A. Ioana, S. Popa, S. Vaes, A class of superrigid group von Neumann algebras. Preprint [arXiv:1007.1412](#).

- [17] A.S. Kechris, *Classical Descriptive Set Theory*, in: Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.
- [18] N. Ozawa, S. Popa, On a class of II_1 factors with at most one Cartan subalgebra, *Ann. of Math.* 172 (2010) 713–749.
- [19] N. Ozawa, S. Popa, On a class of II_1 factors with at most one Cartan subalgebra, II, *Amer. J. Math.* 132 (2010) 841–866.
- [20] J.A. Packer, Point spectrum of ergodic abelian group actions and the corresponding group-measure factors, *Pacific J. Math.* 119 (1985) 381–405.
- [21] J. Peterson, Examples of group actions which are virtually W^* -superrigid. Preprint [arXiv:1002.1745](https://arxiv.org/abs/1002.1745).
- [22] S. Popa, Correspondences, INCREST Preprint, vol. 56, 1986. Available at: www.math.ucla.edu/~popa/preprints.html.
- [23] S. Popa, Some rigidity results in type II_1 factors, *C. R. Acad. Sci. Paris Sér. I Math.* 311 (1990) 535–538.
- [24] S. Popa, On a class of type II_1 factors with Betti numbers invariants, *Ann. of Math.* 163 (2006) 809–899.
- [25] S. Popa, Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups, I, *Invent. Math.* 165 (2006) 369–408.
- [26] S. Popa, Cocycle and orbit equivalence superrigidity for malleable actions of w -rigid groups, *Invent. Math.* 170 (2007) 243–295.
- [27] S. Popa, Deformation and rigidity for group actions and von Neumann algebras, in: *Proceedings of the International Congress of Mathematicians (Madrid, 2006)*, Vol. I, European Mathematical Society Publishing House, 2007, pp. 445–477.
- [28] S. Popa, On Ozawa’s property for free group factors, *Int. Math. Res. Not.* (2007) Article ID rnm036.
- [29] S. Popa, On the superrigidity of malleable actions with spectral gap, *J. Amer. Math. Soc.* 21 (2008) 981–1000.
- [30] S. Popa, A.M. Sinclair, R.R. Smith, Perturbations of subalgebras of type II_1 factors, *J. Funct. Anal.* 213 (2004) 346–379.
- [31] S. Popa, S. Vaes, Strong rigidity of generalized Bernoulli actions and computations of their symmetry groups, *Adv. Math.* 217 (2008) 833–872.
- [32] S. Popa, S. Vaes, Group measure space decomposition of II_1 factors and W^* -superrigidity, *Invent. Math.* 182 (2010) 371–417.
- [33] S. Popa, S. Vaes, Cocycle and orbit superrigidity for lattices in $\text{SL}(n, \mathbb{R})$ acting on homogeneous spaces, in: B. Farb, D. Fisher (Eds.), *Geometry, Rigidity and Group Actions*, The University of Chicago Press, 2011, pp. 419–451.
- [34] I.M. Singer, Automorphisms of finite factors, *Amer. J. Math.* 77 (1955) 117–133.
- [35] S. Vaes, Rigidity for von Neumann algebras and their invariants, in: *Proceedings of the International Congress of Mathematicians (Hyderabad, 2010)*, Vol. III, Hindustan Book Agency, 2010, pp. 1624–1650.
- [36] S. Vaes, One-cohomology and the uniqueness of the group measure space decomposition of a II_1 factor. *Math. Ann.* (in press) [arXiv:1012.5377](https://arxiv.org/abs/1012.5377).